

Weyl's Law and Connes' Trace Theorem for Noncommutative Two Tori

FARZAD FATHIZADEH AND MASOUD KHALKHALI

Abstract

We prove the analogue of Weyl's law for a noncommutative Riemannian manifold, namely the noncommutative two torus \mathbb{T}_θ^2 equipped with a general translation invariant conformal structure and a Weyl conformal factor. This is achieved by studying the asymptotic distribution of the eigenvalues of the perturbed Laplacian on \mathbb{T}_θ^2 . We also prove the analogue of Connes' trace theorem by showing that the Dixmier trace and a noncommutative residue coincide on pseudodifferential operators of order -2 on \mathbb{T}_θ^2 .

Mathematics Subject Classification (2010). 46L87, 58B34, 35S05.

Keywords. Weyl's law, heat kernel, Connes' trace theorem, noncommutative residue, noncommutative tori.

1 Introduction

A celebrated theorem of Weyl states that one can compute the volume of a compact Riemannian manifold from the asymptotic growth of the eigenvalues of its Laplacian. This result, known as Weyl's law, has been the starting point of numerous developments and conjectures in mathematics and physics. Some of these results are now presented in the context of spectral geometry, inverse problems and quantum mechanics. It is pointed out in [19] that a subtle asymptotic identity seems to be the only intrinsic link between a classical observable h and its quantization H . That is, the eigenvalue counting function for H defined by $N(\Lambda) = \#\{\lambda \leq \Lambda\}$, and the phase space volume $\text{Vol}(h \leq \Lambda)$ satisfy the asymptotic identity

$$N(\Lambda) \sim \gamma \text{Vol}(h \leq \Lambda) \quad \text{as} \quad \Lambda \rightarrow \infty,$$

where γ is a constant that does not depend on h or H . Here h is a real-valued function on a symplectic phase space and H is a selfadjoint operator on a Hilbert space.

Like similar results in spectral geometry which encode geometric information in terms of spectral data, Weyl's law has played an important role in the development of metric aspects of noncommutative geometry [6, 7, 8, 9, 11, 3, 10].

A noncommutative geometric space is described by a spectral triple $(\mathcal{A}, \mathcal{H}, D)$. Here \mathcal{A} is an algebra acting by bounded operators on a Hilbert space \mathcal{H} , and D is an unbounded selfadjoint operator acting in \mathcal{H} such that the commutators $[D, a]$ are bounded for all $a \in \mathcal{A}$ [8, 11, 9, 10]. Under suitable summability and regularity conditions, the geometric invariants such as volume and curvature of a noncommutative space $(\mathcal{A}, \mathcal{H}, D)$ with metric dimension d are obtained by considering small time asymptotic expansions of the form

$$\text{Trace}(ae^{-tD^2}) \sim \sum_{n=0}^{\infty} a_n(a, D^2)t^{\frac{n-d}{2}}, \quad a \in \mathcal{A}.$$

There is an equivalent formulation of such geometric invariants in terms of spectral zeta functions.

The main results of this paper are Theorem 4.2 and Theorem 5.1 in which we prove the analogue of Weyl's law and Connes' trace theorem [7] for the noncommutative two torus \mathbb{T}_θ^2 . In Section 2 we recall the construction of a spectral triple on \mathbb{T}_θ^2 via a complex number τ in the upper half-plane and a positive invertible element $k \in C^\infty(\mathbb{T}_\theta^2)$. The complex parameter τ defines a translation invariant conformal structure on \mathbb{T}_θ^2 and the element k plays the role of a Weyl conformal factor [13, 15]. In Section 3 we recall Connes' pseudodifferential calculus [5] for the canonical dynamical system defining the noncommutative two torus. Using this pseudodifferential calculus and heat kernel techniques [18, 13, 15], in Section 4 we find the first coefficient in the small time heat kernel expansion of the perturbed Laplacian on \mathbb{T}_θ^2 . The volume of \mathbb{T}_θ^2 with respect to the perturbed volume form manifests itself in this coefficient, and we use a Tauberian theorem to find its relation with the asymptotic distribution of the eigenvalues of the Laplacian. This establishes the analogue of Weyl's law for \mathbb{T}_θ^2 . In Section 5 we prove the analogue of Connes' trace theorem [7] for \mathbb{T}_θ^2 . That is, we show that the Dixmier trace [14] and a noncommutative residue defined in [17] coincide on pseudodifferential operators of order -2 on \mathbb{T}_θ^2 .

Recently there has been much progress in understanding the local differential geometry of the noncommutative two torus equipped with a general metric. In [15], the Gauss-Bonnet theorem of [13] is extended to full generality following the pioneering work [4] in the subject. The much more delicate question of computing the scalar curvature of this noncommutative manifold is now also fully settled in [12, 16]. In [12], among other results, one can also find a closed formula for the Ray-Singer analytic torsion in terms of the Dirichlet quadratic form, and the corresponding evolution equation for the metric is shown to give the appropriate analogue of Ricci curvature. A different approach to the question of Ricci curvature is proposed in [2].

The first author would like to thank IHES for kind support and excellent environment during his visit in Summer 2011 where part of this work was carried out. We would like to thank the referees for carefully reading the paper and making several useful suggestions.

2 Preliminaries

Let Σ be a closed, oriented, 2-dimensional smooth manifold equipped with a Riemannian metric g . The spectrum of the Laplacian $\Delta_g = d^*d$, where d is the de Rham differential operator acting on smooth functions on Σ , encodes important geometric and topological information about (Σ, g) . In fact there is an asymptotic expansion for the trace of the heat kernel $e^{-t\Delta_g}$ which is of the form

$$\text{Trace}(e^{-t\Delta_g}) \sim t^{-1} \sum_{n=0}^{\infty} a_{2n}(\Delta_g)t^n \quad (t \rightarrow 0), \quad (1)$$

with $a_0(\Delta_g) = \frac{1}{4\pi}\text{Vol}(\Sigma)$ and $a_2(\Delta_g) = \frac{1}{24\pi}K(\Sigma)$. Here $\text{Vol}(\Sigma)$ and $K(\Sigma)$ respectively denote the volume and total curvature of the Riemannian manifold.

On the other hand $\text{Trace}(e^{-t\Delta_g})$ depends only on the spectrum of the Laplacian since

$$\text{Trace}(e^{-t\Delta_g}) = \sum e^{-t\lambda_j}, \quad (2)$$

where the summation is over all eigenvalues λ_j of Δ_g counted with multiplicity. Comparing relations (1), (2), and using Karamata's Tauberian theorem [1] (page 91) the term $a_0 = \frac{1}{4\pi}\text{Vol}(\Sigma)$ in the above asymptotic expansion is seen to depend only on the asymptotic growth of the eigenvalues λ_j . More precisely, one obtains

$$\lim_{j \rightarrow \infty} \frac{\lambda_j}{j} = \frac{4\pi}{\text{Vol}(\Sigma)}.$$

This provides a proof for the celebrated Weyl's law which states that one can hear the volume of a Riemannian manifold from the asymptotic distribution of the eigenvalues of its Laplacian.

In the following subsections we recall the definition of the noncommutative two torus and the construction of a spectral triple on it. The spectral triple will encode the metric information corresponding to general translation invariant conformal structures on \mathbb{T}_θ^2 and a Weyl conformal factor by means of which one can vary inside the conformal class [4, 13, 15].

2.1 The irrational rotation algebra

Let θ be an irrational number. Recall that the irrational rotation C^* -algebra A_θ is, by definition, the universal unital C^* -algebra generated by two unitaries U, V satisfying

$$VU = e^{2\pi i\theta}UV.$$

One usually thinks of A_θ as the algebra of continuous functions on the noncommutative 2-torus \mathbb{T}_θ^2 . There is a continuous action of the ordinary torus \mathbb{T}^2 , $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$, on A_θ by C^* -algebra automorphisms $\{\alpha_s\}_{s \in \mathbb{R}^2}$ defined by

$$\alpha_s(U^m V^n) = e^{is \cdot (m, n)} U^m V^n, \quad m, n \in \mathbb{Z}.$$

The space of smooth elements for this action, denoted by A_θ^∞ , consists of those elements $a \in A_\theta$ for which the map $s \mapsto \alpha_s(a)$ is a smooth map from \mathbb{R}^2 to

A_θ . It is a dense subalgebra of A_θ which can be alternatively described as the algebra of elements of the form

$$\sum_{m,n \in \mathbb{Z}} a_{m,n} U^m V^n$$

with rapidly decreasing coefficients, *i.e.* for any non-negative integers k, q , we have

$$\sup_{m,n \in \mathbb{Z}} ((1 + |m|)^k (1 + |n|)^q |a_{m,n}|) < \infty.$$

There is a unique normalized trace \mathfrak{t} on A_θ whose restriction to the smooth elements is given by

$$\mathfrak{t}\left(\sum_{m,n \in \mathbb{Z}} a_{m,n} U^m V^n\right) = a_{0,0}.$$

This trace functional is positive and faithful, and is used to define an inner product on A_θ :

$$\langle a, b \rangle = \mathfrak{t}(b^* a), \quad a, b \in A_\theta. \quad (3)$$

Completing A_θ with respect to this inner product, we obtain a Hilbert space which will be denoted by \mathcal{H}_0 .

The infinitesimal generators of the above action of \mathbb{T}^2 on A_θ are the derivations $\delta_1, \delta_2 : A_\theta^\infty \rightarrow A_\theta^\infty$ defined by

$$\delta_1(U^m V^n) = m U^m V^n, \quad \delta_2(U^m V^n) = n U^m V^n, \quad m, n \in \mathbb{Z}.$$

In fact, δ_1, δ_2 are analogues of the differential operators $\frac{1}{i}\partial/\partial x, \frac{1}{i}\partial/\partial y$ acting on smooth functions on the ordinary two torus. We have $\delta_j(a^*) = -\delta_j(a)^*$ for $j = 1, 2$, and all $a \in A_\theta^\infty$. Moreover, since $\mathfrak{t} \circ \delta_j = 0$ for $j = 1, 2$, we have the analogue of integration by parts:

$$\mathfrak{t}(a \delta_j(b)) = -\mathfrak{t}(\delta_j(a) b), \quad a, b \in A_\theta^\infty.$$

As unbounded operators on \mathcal{H}_0 , the derivations δ_1, δ_2 are formally selfadjoint and have unique selfadjoint extensions.

2.2 Conformal structures on A_θ

In this subsection we recall the analogues of complex and conformal structures for noncommutative two tori. Our definitions are motivated by their commutative counterparts. To any complex number τ in the upper half-plane, $\Im(\tau) > 0$, we can associate a complex structure on the noncommutative two torus by defining the operators

$$\partial = \delta_1 + \bar{\tau} \delta_2, \quad \partial^* = \delta_1 + \tau \delta_2.$$

These derivations on A_θ^∞ are analogues of Dolbeault operators on Riemann surfaces. We note that ∂ is an unbounded densely defined operator on \mathcal{H}_0 and ∂^* is its formal adjoint.

It is explained in [8] (§VI. 2) that the information on the conformal structure represented by τ is encoded in the positive Hochschild 2-cocycle on A_θ^∞ given by

$$\psi(a, b, c) = -\mathfrak{t}(a\partial(b)\partial^*(c)), \quad a, b, c \in A_\theta^\infty.$$

Therefore, the analogue of the space of $(1, 0)$ -forms on the ordinary two torus is defined to be the Hilbert space completion of the space of finite sums $\sum a\partial b$, $a, b \in A_\theta^\infty$, with respect to the inner product defined by

$$\langle a\partial(b), a'\partial(b') \rangle = \mathfrak{t}((a')^*a\partial(b)\partial(b')^*), \quad a, b, a', b' \in A_\theta^\infty.$$

This Hilbert space will be denoted by $\mathcal{H}^{(1,0)}$.

In order to vary inside the conformal class represented by τ , we choose a selfadjoint element $h \in A_\theta^\infty$ (cf. [13]), and define a positive linear functional φ on A_θ by

$$\varphi(a) = \mathfrak{t}(ae^{-h}), \quad a \in A_\theta.$$

This linear functional is not a trace. However, it is a twisted trace and satisfies the KMS condition at $\beta = 1$ for the 1-parameter group $\{\sigma_t\}_{t \in \mathbb{R}}$ of inner automorphisms

$$\sigma_t = \Delta^{-it},$$

where the modular operator Δ is defined by (cf. [13])

$$\Delta(a) = e^{-h}ae^h, \quad a \in A_\theta.$$

We define a new inner product $\langle \cdot, \cdot \rangle_\varphi$ on A_θ by

$$\langle a, b \rangle_\varphi = \varphi(b^*a), \quad a, b \in A_\theta.$$

The Hilbert space obtained from completing A_θ with respect to this inner product will be denoted by \mathcal{H}_φ .

2.3 A spectral triple on A_θ

Following the above notations, let ∂_φ be the same operator as ∂ but viewed as an unbounded operator from \mathcal{H}_φ to $\mathcal{H}^{(1,0)}$. Consider the left action of A_θ on the Hilbert space

$$\mathcal{H} = \mathcal{H}_\varphi \oplus \mathcal{H}^{(1,0)}$$

defined by $a(b, c) = (ab, ac)$ for all $a \in A_\theta$, $b \in \mathcal{H}_\varphi$, and $c \in \mathcal{H}^{(1,0)}$. The operator

$$D = \begin{pmatrix} 0 & \partial_\varphi^* \\ \partial_\varphi & 0 \end{pmatrix} : \mathcal{H} \rightarrow \mathcal{H}$$

is an unbounded selfadjoint operator and one checks that this data defines a spectral triple which we denote by $(A_\theta, \mathcal{H}, D)$ [13]. This is an even spectral triple with the grading operator given by

$$\gamma = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} : \mathcal{H} \rightarrow \mathcal{H}.$$

The above spectral triple endows the noncommutative two torus with a metric structure in the sense of noncommutative geometry (cf. [8], §VI). More precisely it encodes the metric information represented by the complex structure and the Weyl conformal factor introduced in Subsection 2.2. Apart from this paper, this spectral triple played an important role in recent work on the Gauss-Bonnet theorem and scalar curvature for noncommutative two tori [13, 15, 12, 16].

The full Laplacian

$$D^2 = \begin{pmatrix} \partial_\varphi^* \partial_\varphi & 0 \\ 0 & \partial_\varphi \partial_\varphi^* \end{pmatrix}$$

consists of the Laplacian on \mathcal{H}_φ and the Laplacian on $\mathcal{H}^{(1,0)}$. Since these operators have the same spectrum away from 0, for the purpose of this paper, we only work with the first Laplacian

$$\Delta' := \partial_\varphi^* \partial_\varphi : \mathcal{H}_\varphi \rightarrow \mathcal{H}_\varphi.$$

In the flat case, when $h = 0$, one can easily see that this Laplacian reduces to the flat Laplacian

$$\Delta := \partial^* \partial = \delta_1^2 + 2\Re(\tau)\delta_1\delta_2 + |\tau|^2\delta_2^2.$$

In general, $\Delta' : \mathcal{H}_\varphi \rightarrow \mathcal{H}_\varphi$ is a complicated operator. However, it can be shown that it is anti-unitarily equivalent to the operator

$$k\Delta k : \mathcal{H}_0 \rightarrow \mathcal{H}_0,$$

where $k := e^{h/2}$ acts on \mathcal{H}_0 by left multiplication (cf. [13, 15]). In particular these two operators have the same spectrum. This observation plays a crucial role in this paper.

3 Pseudodifferential Operators on \mathbb{T}_θ^2

Connes' pseudodifferential calculus [5] may be used to compute spectral invariants of noncommutative tori as in [13, 12, 15, 16]. Our proof of Weyl's law and Connes' trace theorem for noncommutative tori is also based on this calculus.

A differential operator of order $n \geq 0$ on the noncommutative two torus is an operator of the form

$$\sum_{j_1, j_2} a_{j_1, j_2} \delta_1^{j_1} \delta_2^{j_2} : A_\theta^\infty \rightarrow A_\theta^\infty,$$

where $a_{j_1, j_2} \in A_\theta^\infty$ and j_1, j_2 are non-negative integers such that $j_1 + j_2 \leq n$. The elements a_{j_1, j_2} act by left multiplication. Using operator-valued symbols, the above notion of differential operators can be generalized to the notion of pseudodifferential operators. Here we briefly recall these symbols and the corresponding pseudodifferential calculus. We shall use the notation $\partial_1 = \frac{\partial}{\partial \xi_1}$, $\partial_2 = \frac{\partial}{\partial \xi_2}$ for partial differentiation with respect to the coordinates of \mathbb{R}^2 .

Definition 3.1. Let n be an integer. A smooth map $\rho : \mathbb{R}^2 \rightarrow A_\theta^\infty$ is said to be a symbol of order n if for all non-negative integers i_1, i_2, j_1, j_2 we have

$$\|\delta_1^{i_1} \delta_2^{i_2} \partial_1^{j_1} \partial_2^{j_2} \rho(\xi)\| \leq c(1 + |\xi|)^{n-j_1-j_2},$$

where c is a constant, and if there exists a smooth map $k : \mathbb{R}^2 \setminus \{0\} \rightarrow A_\theta^\infty$ such that

$$\lim_{\lambda \rightarrow \infty} \lambda^{-n} \rho(\lambda \xi_1, \lambda \xi_2) = k(\xi_1, \xi_2), \quad (\xi_1, \xi_2) \in \mathbb{R}^2 \setminus \{0\}.$$

Here $\|\cdot\|$ denotes the C^* -norm of A_θ . The space of symbols of order n is denoted by S_n .

To a symbol $\rho \in S_n$ one can associate an operator P_ρ on A_θ^∞ defined by

$$P_\rho(a) = (2\pi)^{-2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} e^{-is \cdot \xi} \rho(\xi) \alpha_s(a) ds d\xi, \quad a \in A_\theta^\infty. \quad (4)$$

The operator P_ρ is said to be a pseudodifferential operator of order n . For example, the differential operator $\sum_{j_1+j_2 \leq n} a_{j_1, j_2} \delta_1^{j_1} \delta_2^{j_2}$ is associated with the symbol $\sum_{j_1+j_2 \leq n} a_{j_1, j_2} \xi_1^{j_1} \xi_2^{j_2}$ via the above formula.

Definition 3.2. Two symbols $\rho, \rho' \in S_k$ are said to be equivalent if $\rho - \rho'$ is in S_n for all integers n . The equivalence of symbols will be denoted by $\rho \sim \rho'$.

The following Proposition in [5] shows that the space of pseudodifferential operators on \mathbb{T}_θ^2 is an algebra and gives an explicit formula for the symbol of the product of two pseudodifferential operators up to the above equivalence relation. It also shows that the formal adjoint of a pseudodifferential operator, with respect to the inner product defined on A_θ^∞ by (3), is again a pseudodifferential operator and gives a formula for its symbol.

Proposition 3.3. Let P and Q be pseudodifferential operators with the symbols ρ and ρ' respectively. Then the adjoint P^* and the product PQ are pseudodifferential operators with the following symbols

$$\begin{aligned} \sigma(P^*) &\sim \sum_{\ell_1, \ell_2 \geq 0} \frac{1}{\ell_1! \ell_2!} \partial_1^{\ell_1} \partial_2^{\ell_2} \delta_1^{\ell_1} \delta_2^{\ell_2} (\rho(\xi))^*, \\ \sigma(PQ) &\sim \sum_{\ell_1, \ell_2 \geq 0} \frac{1}{\ell_1! \ell_2!} \partial_1^{\ell_1} \partial_2^{\ell_2} (\rho(\xi)) \delta_1^{\ell_1} \delta_2^{\ell_2} (\rho'(\xi)). \end{aligned}$$

We also recall the notion of ellipticity for these pseudodifferential operators:

Definition 3.4. Let ρ be a symbol of order n . It is said to be elliptic if $\rho(\xi)$ is invertible for $\xi \neq 0$, and if there exists a constant c such that

$$\|\rho(\xi)^{-1}\| \leq c(1 + |\xi|)^{-n}$$

for sufficiently large $|\xi|$.

The flat Laplacian $\Delta = \delta_1^2 + 2\Re(\tau)\delta_1\delta_2 + |\tau|^2\delta_2^2$ is an example of an elliptic operator on \mathbb{T}_θ^2 .

4 Weyl's Law for \mathbb{T}_θ^2

The resolvent of the Laplacian of a closed Riemannian manifold can be approximated by pseudodifferential operators. Combining this with the Cauchy integral formula one can find a small time asymptotic expansion for the trace of the heat kernel of the Laplacian [18]. As stated in [13] (*cf.* also [15, 12, 16]), one can use Connes' pseudodifferential calculus to find similar asymptotic expansions in noncommutative settings. We recall the main idea of this technique and will use it to establish the analogue of Weyl's law for \mathbb{T}_θ^2 .

Let Δ' be the perturbed Laplacian on A_θ introduced in Subsection 2.3. Using the Cauchy integral formula, one has

$$e^{-t\Delta'} = \frac{1}{2\pi i} \int_C e^{-t\lambda} (\Delta' - \lambda)^{-1} d\lambda,$$

where C is a curve in the complex plane that goes around the non-negative real axis in such a way that

$$e^{-ts} = \frac{1}{2\pi i} \int_C e^{-t\lambda} (s - \lambda)^{-1} d\lambda, \quad s \geq 0.$$

Appealing to this formula, one can use Connes' pseudodifferential calculus to employ similar arguments to those in [18] and derive an asymptotic expansion of the form

$$\text{Trace}(e^{-t\Delta'}) \sim t^{-1} \sum_{n=0}^{\infty} B_{2n}(\Delta') t^n \quad (t \rightarrow 0).$$

That is, one can approximate $(\Delta' - \lambda)^{-1}$ by pseudodifferential operators B_λ whose symbols are of the form

$$b_0(\xi, \lambda) + b_1(\xi, \lambda) + b_2(\xi, \lambda) + \dots,$$

where for $j = 0, 1, 2, \dots$, $b_j(\xi, \lambda)$ is a symbol of order $-2 - j$.

Since $\text{Trace}(e^{-t\Delta'})$ depends only on the eigenvalues of Δ' , for the purpose of deriving the above asymptotic expansion one can work with the operator $k\Delta k$ which as shown in [13, 15] is anti-unitarily equivalent to Δ' . The symbol of $k\Delta k$ as a pseudodifferential operator [15] is equal to $a_2(\xi) + a_1(\xi) + a_0(\xi)$, where

$$a_2(\xi) = \xi_1^2 k^2 + |\tau|^2 \xi_2^2 k^2 + 2\Re(\tau) \xi_1 \xi_2 k^2,$$

$$a_1(\xi) = 2\xi_1 k \delta_1(k) + 2|\tau|^2 \xi_2 k \delta_2(k) + 2\Re(\tau) \xi_1 k \delta_2(k) + 2\Re(\tau) \xi_2 k \delta_1(k),$$

$$a_0(\xi) = k \delta_1^2(k) + |\tau|^2 k \delta_2^2(k) + 2\Re(\tau) k \delta_1 \delta_2(k).$$

Therefore, using the calculus of symbols described in Section 3, one has to solve the equation

$$(b_0 + b_1 + b_2 + \dots)((a_2 - \lambda) + a_1 + a_0) \sim 1.$$

Here, λ is treated as a symbol of order 2 and we let $a'_2 = a_2 - \lambda$, $a'_1 = a_1$, $a'_0 = a_0$. Then the above equation yields

$$\sum_{\substack{j, \ell_1, \ell_2 \geq 0, \\ k=0,1,2}} \frac{1}{\ell_1! \ell_2!} \partial_1^{\ell_1} \partial_2^{\ell_2} (b_j) \delta_1^{\ell_1} \delta_2^{\ell_2} (a'_k) \sim 1.$$

Comparing symbols of the same order on both sides, one concludes that

$$b_0 = a'_2{}^{-1} = (a_2 - \lambda)^{-1} = (\xi_1^2 k^2 + |\tau|^2 \xi_2^2 k^2 + 2\Re(\tau) \xi_1 \xi_2 k^2 - \lambda)^{-1},$$

and

$$b_n = - \sum_{\substack{2+j+\ell_1+\ell_2-k=n, \\ 0 \leq j < n, 0 \leq k \leq 2}} \frac{1}{\ell_1! \ell_2!} \partial_1^{\ell_1} \partial_2^{\ell_2} (b_j) \delta_1^{\ell_1} \delta_2^{\ell_2} (a_k) b_0, \quad n > 0.$$

Similar to [18] one can use these symbols to approximate $e^{-tk\Delta k}$ with suitable infinitely smoothing operators and derive the desired asymptotic expansion. We record this result, already stated in [13], in the following proposition.

Proposition 4.1. *Let Δ' be the perturbed Laplacian on A_θ as above. There is an asymptotic expansion*

$$\text{Trace}(e^{-t\Delta'}) \sim t^{-1} \sum_{n=0}^{\infty} B_{2n}(\Delta') t^n \quad (t \rightarrow 0), \quad (5)$$

where for each $n = 0, 1, 2, \dots$,

$$B_{2n}(\Delta') = \frac{1}{2\pi i} \int \int_C e^{-\lambda t} (b_{2n}(\xi, \lambda)) d\lambda d\xi.$$

The following theorem is the analogue of Weyl's law for the noncommutative two torus equipped with a general metric. In fact we show that the first term in the above asymptotic expansion is intimately related to a natural definition for the volume of $(\mathbb{T}_\theta^2, \tau, k)$, and by means of a Tauberian theorem the relation between the volume and the eigenvalue counting function of Δ' is derived. We recall that τ is a complex parameter in the upper half-plane associated to which is a conformal structure on \mathbb{T}_θ^2 , and $k = e^{h/2}$ is an invertible selfadjoint element of A_θ^∞ that plays the role of a Weyl conformal factor.

Theorem 4.2. *For any positive real number λ , let $N(\lambda)$ denote the number of eigenvalues of the perturbed Laplacian Δ' that are less than λ . Then as $\lambda \rightarrow \infty$, we have*

$$N(\lambda) \sim \frac{\pi}{\Im(\tau)} \varphi(1) \lambda. \quad (6)$$

Proof. Using Proposition 4.1 and passing to the coordinates [15]

$$\xi_1 = r \cos \theta - r \frac{\Re(\tau)}{\Im(\tau)} \sin \theta, \quad \xi_2 = \frac{r}{\Im(\tau)} \sin \theta,$$

where θ ranges from 0 to 2π and r goes from 0 to ∞ , we can compute the first coefficient in the asymptotic expansion (5) for the trace of the heat kernel of Δ' directly:

$$B_0(\Delta') = \frac{1}{2\pi i} \int \int_C e^{-\lambda} \mathfrak{t}(b_0(\xi, \lambda)) d\lambda d\xi = \frac{\pi}{\Im(\tau)} \mathfrak{t}(k^{-2}).$$

Note that

$$b_0 = a_2'^{-1} = (\xi_1^2 k^2 + |\tau|^2 \xi_2^2 k^2 + 2\Re(\tau)\xi_1\xi_2 k^2 - \lambda)^{-1}.$$

Now the asymptotic behavior of the eigenvalue counting function N can be determined as follows. It follows from (5) that

$$\lim_{t \rightarrow 0^+} t \sum e^{-t\lambda_j} = B_0(\Delta') = \frac{\pi}{\Im(\tau)} \mathfrak{t}(k^{-2}),$$

where the summation is over all eigenvalues λ_j of Δ' . Then it follows immediately from Karamata's Tauberian theorem [1] (page 91) that the eigenvalue counting function N satisfies

$$N(\lambda) \sim \frac{\pi}{\Im(\tau)\Gamma(2)} \mathfrak{t}(k^{-2})\lambda = \frac{\pi}{\Im(\tau)} \mathfrak{t}(k^{-2})\lambda \quad \text{as } \lambda \rightarrow \infty.$$

□

A corollary of this theorem is that $(1 + \Delta')^{-1}$ is in the domain of the Dixmier trace. Before stating the corollary we quickly review the Dixmier trace and the noncommutative integral, following [8].

We denote the ideal of compact operators on a Hilbert space \mathcal{H} by $\mathcal{K}(\mathcal{H})$. For any $T \in \mathcal{K}(\mathcal{H})$, let $\mu_n(T), n = 1, 2, \dots$, denote the sequence of eigenvalues of its absolute value $|T| = (T^*T)^{\frac{1}{2}}$ written in decreasing order with multiplicity:

$$\mu_1(T) \geq \mu_2(T) \geq \dots \geq 0.$$

The Dixmier trace is a trace functional on an ideal of compact operators $\mathcal{L}^{1,\infty}(\mathcal{H})$ defined as

$$\mathcal{L}^{1,\infty}(\mathcal{H}) = \left\{ T \in \mathcal{K}(\mathcal{H}); \quad \sum_{n=1}^N \mu_n(T) = O(\log N) \right\}.$$

This ideal of operators is equipped with a natural norm:

$$\|T\|_{1,\infty} := \sup_{N \geq 2} \frac{1}{\log N} \sum_{n=1}^N \mu_n(T), \quad T \in \mathcal{L}^{1,\infty}(\mathcal{H}).$$

It is clear that trace class operators are automatically in $\mathcal{L}^{1,\infty}(\mathcal{H})$. The Dixmier trace of an operator $T \in \mathcal{L}^{1,\infty}(\mathcal{H})$ measures the logarithmic divergence of its

ordinary trace. More precisely, for any positive operator $T \in \mathcal{L}^{1,\infty}(\mathcal{H})$ we are interested in the limiting behavior of the sequence

$$\frac{1}{\log N} \sum_{n=1}^N \mu_n(T), \quad N = 2, 3, \dots$$

While by our assumption this sequence is bounded, its usual limit may not exist and must be replaced by a suitable generalized limit. The limiting procedure is carried out by means of a state on a C^* -algebra. Recall that a state on a C^* -algebra is a non-zero positive linear functional on the algebra.

To define the Dixmier trace of a positive operator $T \in \mathcal{L}^{1,\infty}(\mathcal{H})$, consider the partial trace

$$\text{Trace}_N(T) = \sum_{n=1}^N \mu_n(T), \quad N = 1, 2, \dots,$$

and its piecewise affine interpolation denoted by $\text{Trace}_r(T)$ for $r \in [1, \infty)$. Then let

$$\tau_\Lambda(T) := \frac{1}{\log \Lambda} \int_e^\Lambda \frac{\text{Trace}_r(T)}{\log r} \frac{dr}{r}, \quad \Lambda \in [e, \infty),$$

be the Cesàro mean of the function $\text{Trace}_r(T)/\log r$ over the multiplicative group \mathbb{R}_+^* . Now choosing a normalized state $\omega : C_b[e, \infty) \rightarrow \mathbb{C}$ on the algebra of bounded continuous functions on $[e, \infty)$ such that $\omega(f) = 0$ for all f vanishing at ∞ , the Dixmier trace of $T \geq 0$ is defined as

$$\text{Tr}_\omega(T) = \omega(\tau_\Lambda(T)).$$

Then one can extend Tr_ω to all of $\mathcal{L}^{1,\infty}(\mathcal{H})$ by linearity.

The resulting linear functional Tr_ω is a positive trace on $\mathcal{L}^{(1,\infty)}(\mathcal{H})$ which in general depends on the limiting procedure ω . The operators $T \in \mathcal{L}^{1,\infty}(\mathcal{H})$ whose Dixmier trace $\text{Tr}_\omega(T)$ is independent of the choice of the state ω are called measurable and we will denote their Dixmier trace by $\int T$. If for a compact positive operator T we have

$$\mu_n(T) \sim \frac{c}{n} \quad \text{as } n \rightarrow \infty,$$

where c is a constant, then T is measurable and $\int T = c$. We use this fact to compute the Dixmier trace of $(1 + \Delta')^{-1}$ where Δ' is the perturbed Laplacian on the noncommutative two torus:

Corollary 4.3. *The operator $(1 + \Delta')^{-1}$ belongs to $\mathcal{L}^{1,\infty}(\mathcal{H}_\varphi)$. Moreover it is measurable and*

$$\int (1 + \Delta')^{-1} = \frac{\pi\varphi(1)}{\mathfrak{S}(\tau)}.$$

Proof. Assuming that the eigenvalues λ_j of Δ' are written in increasing order

$$\lambda_0 \leq \lambda_1 \leq \lambda_2 \leq \dots,$$

the asymptotic estimate (6) can be reformulated as

$$\lambda_j \sim \frac{\Im(\tau)}{\pi\varphi(1)} j \quad \text{as } j \rightarrow \infty.$$

Therefore

$$(1 + \lambda_j)^{-1} \sim \frac{\pi\varphi(1)}{\Im(\tau)} \left(\frac{\pi\varphi(1)}{\Im(\tau)} + j \right)^{-1} \quad \text{as } j \rightarrow \infty,$$

and it follows that

$$\int (1 + \Delta')^{-1} = \frac{\pi\varphi(1)}{\Im(\tau)}.$$

□

Considering the classical case explained at the beginning of Section 2, the above theorem suggests the following formula for the volume of the noncommutative two torus \mathbb{T}_θ^2 equipped with the metric associated to the conformal structure represented by τ , $\Im(\tau) > 0$, and the Weyl factor k :

$$\text{Vol}(\mathbb{T}_\theta^2) := \frac{4\pi^2}{\Im(\tau)} \varphi(1) = \frac{4\pi^2}{\Im(\tau)} \mathfrak{t}(k^{-2}).$$

5 Connes' Trace Theorem for \mathbb{T}_θ^2

Let M be a closed smooth manifold of dimension n . Wodzicki defined a trace functional on the algebra of pseudodifferential operators of arbitrary order on M , and proved that it was the only non-trivial trace [21]. This functional, denoted by Res , is called the noncommutative residue. The restriction of Res to pseudodifferential operators of order $-n$ was discovered independently by Guillemin and its properties were studied in [19]. In general, unlike the Dixmier trace, Res is not a positive linear functional. However its restriction to pseudodifferential operators of order $-n$ is positive. One of the main results proved in [7] is that if E is a smooth vector bundle on M then the Dixmier trace Tr_ω and Res coincide on pseudodifferential operators of order $-n$ acting on L^2 sections of E . In fact it is proved that such operators P are measurable operators in $\mathcal{L}^{1,\infty}(L^2(M, E))$ and

$$\int P = \frac{1}{n} \text{Res}(P).$$

This result is known as Connes' trace theorem.

The noncommutative residue of a classical pseudodifferential operator P acting on smooth sections of E is defined as

$$\text{Res}(P) = (2\pi)^{-n} \int_{S^*M} \text{tr}(\rho_{-n}(x, \xi)) dx d\xi,$$

where $S^*M \subset T^*M$ is the unit cosphere bundle on M and ρ_{-n} is the component of order $-n$ of the complete symbol of P . For detailed discussion of traces on classical pseudodifferential operators on manifolds and uniqueness results we refer the reader to [20] and references therein.

In [17], a noncommutative residue on the algebra of classical pseudodifferential operators on the noncommutative two torus is defined and it is proved that up to multiplication by a constant, it is the unique continuous trace on the algebra of symbols. A symbol ρ of order n on A_θ is said to be classical if there is an asymptotic expansion of the form

$$\rho(\xi) \sim \sum_{j=0}^{\infty} \rho_{n-j}(\xi) \quad \text{as} \quad \xi \rightarrow \infty,$$

where each $\rho_{n-j} : \mathbb{R}^2 \setminus \{0\} \rightarrow A_\theta^\infty$ is smooth and positively homogeneous of order $n-j$. It is shown that the homogeneous terms in such an asymptotic expansion are uniquely determined by ρ and the noncommutative residue of P_ρ is defined by

$$\text{res}(P_\rho) = \int_{\mathbb{S}^1} \mathfrak{t}(\rho_{-2}(\xi)) d\Omega,$$

where $d\Omega$ is the Lebesgue measure on the unit circle. This is the analogue of Wodzicki's noncommutative residue.

In the following theorem we prove the analogue of Connes' trace theorem [7] for the noncommutative two torus. That is, we show that the Dixmier trace and the noncommutative residue defined above coincide on classical pseudodifferential operators of order -2 on \mathbb{T}_θ^2 .

Theorem 5.1. *Let ρ be a classical pseudodifferential symbol of order -2 on the noncommutative two torus. Then $P_\rho \in \mathcal{L}^{1,\infty}(\mathcal{H}_0)$ and*

$$\text{Tr}_\omega(P_\rho) = \frac{1}{2} \text{res}(P_\rho).$$

Proof. We have $P_\rho = A(1 + \Delta_0)^{-1}$ where $\Delta_0 = \delta_1^2 + \delta_2^2$ is the Laplacian of the metric associated with $\tau = i$ and $k = 1$, and $A = P_\rho(1 + \Delta_0)$. Since A is a pseudodifferential operator of order 0, it is a bounded operator on \mathcal{H}_0 . It follows from Corollary 4.3 that P_ρ belongs to the ideal $\mathcal{L}^{1,\infty}(\mathcal{H}_0)$. Similarly, one can see that any pseudodifferential operator of order -3 is a trace-class operator. Since the Dixmier trace vanishes on trace-class operators, $\text{Tr}_\omega(P_\rho)$ only depends on the principal part of ρ . That is, if we write

$$\rho(\xi) \sim \rho'(\xi) + \sum_{j=1}^{\infty} \rho_{-2-j}(\xi) \quad \text{as} \quad \xi \rightarrow \infty,$$

where ρ' and ρ_{-2-j} are positively homogeneous of order -2 and $-2-j$, respectively, then

$$\text{Tr}_\omega(P_\rho) = \text{Tr}_\omega(P_{\rho'}). \quad (7)$$

Now we prove that

$$\mathrm{Tr}_\omega(P_{\rho'}) = \mathrm{Tr}_\omega(P_{\rho'_{0,0}}), \quad (8)$$

where

$$\rho'_{0,0} := \mathfrak{t} \circ \rho' : \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{C}.$$

First we make the following observations. The symbol of $P_{\rho'}P_U - P_UP_{\rho'}$ is equivalent to

$$\rho'U - U\rho' + (\partial_1\rho')U + \frac{1}{2!}(\partial_1^2\rho')U + \frac{1}{3!}(\partial_1^3\rho')U + \dots.$$

Since

$$(\partial_1\rho')U + \frac{1}{2!}(\partial_1^2\rho')U + \frac{1}{3!}(\partial_1^3\rho')U + \dots$$

is equivalent to a symbol of order -3 , the Dixmier trace vanishes on the corresponding operator. Therefore, since the Dixmier trace is a trace, we have

$$\mathrm{Tr}_\omega(P_{\rho'}U - P_UP_{\rho'}) = \mathrm{Tr}_\omega(P_{\rho'}P_U - P_UP_{\rho'}) = 0.$$

Hence

$$\mathrm{Tr}_\omega(P_{\rho'}U) = \mathrm{Tr}_\omega(P_UP_{\rho'}). \quad (9)$$

Using a similar argument for V instead of U , we have

$$\mathrm{Tr}_\omega(P_{\rho'}V) = \mathrm{Tr}_\omega(P_V\rho'). \quad (10)$$

Now, in order to prove (8), we consider the expansion of $\rho'(\xi)$ with rapidly decaying coefficients:

$$\rho'(\xi) = \sum_{m,n \in \mathbb{Z}} \rho'_{m,n}(\xi) U^m V^n.$$

Note that since ρ' is positively homogeneous of order -2 , so are the functions $\rho'_{m,n} : \mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{C}$ for all $m, n \in \mathbb{Z}$. Using (9) and (10) we see that for fixed $m, n \in \mathbb{Z}$, the pseudodifferential operator associated with $\rho'_{m,n} U^m V^n$ has the same Dixmier trace as the ones associated with the following symbols:

$$\rho'_{m,n} U^{m-1} V^n U = e^{2\pi i n \theta} \rho'_{m,n} U^m V^n, \quad V \rho'_{m,n} U^m V^{n-1} = e^{2\pi i m \theta} \rho'_{m,n} U^m V^n.$$

Since θ is irrational, it follows that

$$\mathrm{Tr}_\omega(P_{\rho'_{m,n} U^m V^n}) = 0 \quad \text{if} \quad (m, n) \neq (0, 0).$$

Therefore, in order to conclude (8), it suffices to show that if we define

$$q(\xi) = \rho'(\xi) - \sum_{|m|, |n| \leq M} \rho'_{m,n}(\xi) U^m V^n,$$

then $\mathrm{Tr}_\omega(P_q)$ goes to 0 as $M \rightarrow \infty$. This can be proved as follows. We write $P_q = B(1 + \Delta_0)^{-1}$ where B is a pseudodifferential operator of order 0, and we

denote its symbol by b . Since $B = P_q(1 + \Delta_0)$ we have $b \sim q(1 + \xi_1^2 + \xi_2^2)$. Therefore

$$|\mathrm{Tr}_\omega(P_q)| = |\mathrm{Tr}_\omega(P_{q(1+|\xi|^2)}(1 + \Delta_0)^{-1})| \leq \|P_{q(1+|\xi|^2)}\|_{\mathcal{L}(\mathcal{H}_0)} \|(1 + \Delta_0)^{-1}\|_{1,\infty}.$$

Since q is positively homogeneous of order -2 , as M goes to infinity, the A_θ -norm of $q(1+|\xi|^2)$ and its derivatives go to 0 uniformly in $|\xi| \geq 1$. So $\|P_{q(1+|\xi|^2)}\|_{\mathcal{L}(\mathcal{H}_0)}$ approaches 0 as $M \rightarrow \infty$. Hence

$$\lim_{M \rightarrow \infty} \mathrm{Tr}_\omega(P_q) = 0, \quad (11)$$

and the identity (8) is proved.

Combining (7) and (8), so far we have proved that

$$\mathrm{Tr}_\omega(P_\rho) = \mathrm{Tr}_\omega(P_{\rho'}) = \mathrm{Tr}_\omega(P_{\rho'_{0,0}}).$$

Note that $\rho'_{0,0} = \mathfrak{t} \circ \rho'$ is a complex-valued function that is viewed as a symbol of order -2 on \mathbb{T}_θ^2 . This map is determined by its values on \mathbb{S}^1 since it is positively homogeneous of order -2 . In order to analyze $\mathrm{Tr}_\omega(P_{\rho'_{0,0}})$ we define a linear functional μ on the space of smooth complex-valued functions on \mathbb{S}^1 as follows. Given a smooth map $f : \mathbb{S}^1 \rightarrow \mathbb{C} \subset A_\theta^\infty$, which will be viewed as an A_θ -valued map, first, we extend it to a homogeneous map of order -2 from $\mathbb{R}^2 \setminus \{0\}$ to A_θ^∞ , and denote the extension by \tilde{f} . Then we define $\mu(f)$ to be the Dixmier trace of the pseudodifferential operator corresponding to \tilde{f} :

$$\mu(f) := \mathrm{Tr}_\omega(P_{\tilde{f}}), \quad f \in C^\infty(\mathbb{S}^1).$$

Considering the positivity of the Dixmier trace and the formulas for product and adjoint of pseudodifferential operators explained in Proposition 3.3, it follows that μ is a positive linear functional. The argument used for proving identity (11) shows that μ extends by continuity to a positive linear functional on continuous functions on \mathbb{S}^1 . Therefore it follows from the Riesz representation theorem that μ is given by integration against a Borel measure on \mathbb{S}^1 .

Now we show that μ is rotation invariant. Since the range of any smooth map $f : \mathbb{S}^1 \rightarrow \mathbb{C} \subset A_\theta^\infty$ is in the center of A_θ^∞ , in the definition of $P_{\tilde{f}}$ given by (4) the noncommutativity of the algebra does not play a role. So we can think of $P_{\tilde{f}}$ as a pseudodifferential operator on the ordinary two torus. By a straightforward computation one can see that if $\sigma(x, \xi)$ is an ordinary pseudodifferential symbol on \mathbb{R}^2 and if T is a rotation of the plane then

$$P_{\sigma(Tx, T\xi)} = \mathcal{U}_T^{-1} P_{\sigma(x, \xi)} \mathcal{U}_T,$$

where \mathcal{U}_T is the unitary operator given by

$$\mathcal{U}_T(g) := g \circ T^{-1}, \quad g \in C_c^\infty(\mathbb{R}^2).$$

This fact and the trace property of the Dixmier trace imply that

$$\mathrm{Tr}_\omega(P_{\sigma(Tx, T\xi)}) = \mathrm{Tr}_\omega(\mathcal{U}_T^{-1} P_{\sigma(x, \xi)} \mathcal{U}_T) = \mathrm{Tr}_\omega(P_{\sigma(x, \xi)}).$$

In case $\sigma(x, \xi) = \tilde{f}(\xi)$, this implies that $\mu(f \circ T) = \mu(f)$.

Since the linear functional μ is invariant under rotations of the circle, it is given by integration against a constant multiple of the Lebesgue measure on \mathbb{S}^1 . Denoting this constant by c , and using (7) and (8), we can write:

$$\begin{aligned} \mathrm{Tr}_\omega(P_\rho) &= \mathrm{Tr}_\omega(P_{\rho'}) = \mathrm{Tr}_\omega(P_{\rho'_{0,0}}) = \mu\left(\rho'_{0,0} \Big|_{\mathbb{S}^1}\right) \\ &= c \int_{\mathbb{S}^1} \rho'_{0,0} d\Omega = c \mathrm{res}(P_\rho). \end{aligned}$$

Finally the constant c is fixed by $(1 + \Delta_0)^{-1}$ as follows. Using Corollary 4.3, we have $\mathrm{Tr}_\omega((1 + \Delta_0)^{-1}) = \pi$. On the other hand one can easily see that $\mathrm{res}((1 + \Delta_0)^{-1}) = 2\pi$, because the asymptotic expansion with homogeneous terms for its symbol starts with $1/(\xi_1^2 + \xi_2^2)$. Hence $c = \frac{1}{2}$. □

References

- [1] N. Berline, E. Getzler, M. Vergne, *Heat kernels and Dirac operators*, Springer-Verlag, Berlin, 2004.
- [2] T. A. Bhuyain and M. Marcolli, *The Ricci flow on noncommutative two-tori*, Letters in Mathematical Physics, DOI 10.1007/s11005-012-0550-0.
- [3] A. Chamseddine and A. Connes, *The Spectral action principle*, Comm. Math. Phys. Vol.186 (1997), 731-750.
- [4] P. B. Cohen and A. Connes, *Conformal geometry of the irrational rotation algebra*, Preprint MPI (92-93).
- [5] A. Connes, *C*-algèbres et géométrie différentielle*, C.R. Acad. Sc. Paris, t. 290, Série A, 599-604, 1980.
- [6] A. Connes, *Noncommutative differential geometry*, Inst. Hautes Études Sci. Publ. Math. No. **62**, 257-360, 1985.
- [7] A. Connes, *The action functional in noncommutative geometry*, Comm. Math. Phys. 117, no. 4, 673-683, 1988.
- [8] A. Connes, *Noncommutative geometry*, Academic Press, 1994.
- [9] A. Connes, *Gravity coupled with matter and the foundation of noncommutative geometry*, Comm. Math. Phys. 182 (1996), no. 1, 155-176.
- [10] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives*, American Mathematical Society Colloquium Publications, 55, 2008.
- [11] A. Connes and H. Moscovici, *The local index formula in noncommutative geometry*, Geom. Funct. Anal. 5, no. 2, 174-243, 1995.

- [12] A. Connes and H. Moscovici, *Modular curvature for noncommutative two-tori*, arXiv:1110.3500.
- [13] A. Connes and P. Tretkoff, *The Gauss-Bonnet theorem for the noncommutative two torus*, Noncommutative geometry, arithmetic, and related topics, 141–158, Johns Hopkins Univ. Press, 2011.
- [14] J. Dixmier, *Existence de traces non normales*, C. R. Acad. Sci. Paris Sér. A-B 262, A1107-A1108, 1966.
- [15] F. Fathizadeh and M. Khalkhali, *The Gauss-Bonnet theorem for noncommutative two tori with a general conformal structure*, To appear in the Journal of Noncommutative Geometry.
- [16] F. Fathizadeh and M. Khalkhali, *Scalar curvature for the noncommutative two torus*, arXiv:1110.3511.
- [17] F. Fathizadeh and M. W. Wong, *Noncommutative residues for pseudodifferential operators on the noncommutative two-torus*, Journal of Pseudo-Differential Operators and Applications, **2**(3) (2011), 289–302.
- [18] P. Gilkey, *Invariance theory, the heat equation, and the Atiyah-Singer index theorem*, Mathematics Lecture Series, 11. Publish or Perish, Inc., Wilmington, DE, 1984.
- [19] V. Guillemin, *A new proof of Weyl's formula on the asymptotic distribution of eigenvalues*, Adv. in Math. 55 (1985), no. 2, 131–160.
- [20] M. Lesch and C. Neira Jiménez, *Classification of traces and hypertraces on spaces of classical pseudodifferential operators*, To appear in the Journal of Noncommutative Geometry.
- [21] M. Wodzicki, *Noncommutative residue. I. Fundamentals*, in *K-theory, Arithmetic and Geometry* (Moscow, 1984–1986), Lecture Notes in Mathematics **1289**, Springer, 320-399, 1987.

Department of Mathematics and Statistics, York University, Toronto, Ontario, Canada, M3J 1P3
ffathiza@mathstat.yorku.ca

Department of Mathematics, The University of Western Ontario, London, Ontario, Canada, N6A 5B7
masoud@uwo.ca