

RATIONAL SMOOTHNESS, CELLULAR DECOMPOSITIONS AND GKM THEORY

RICHARD GONZALES

ABSTRACT. We introduce the notion of \mathbb{Q} -filtrable varieties: projective varieties with a torus action and a finite number of fixed points, such that the cells of the associated Bialynicki-Birula decomposition are all rationally smooth. Our main results develop GKM theory in this setting. We also supply a method for building nice combinatorial bases on the equivariant cohomology of any \mathbb{Q} -filtrable GKM variety. Applications to the theory of group embeddings are provided.

INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Let X be a smooth complex projective algebraic variety with a \mathbb{C}^* -action and finitely many fixed points x_1, \dots, x_m . The method of Bialynicki-Birula ([BB1]) gives rise to a decomposition of X into locally closed subvarieties:

$$W_i = \{x \in X \mid \lim_{t \rightarrow 0} tx = x_i\}.$$

Clearly, $X = \bigsqcup_i W_i$. The subvarieties W_i are called cells of the decomposition. Theorem 4.3 of [BB1] asserts that all cells are isomorphic to affine spaces, that is, $W_i \simeq \mathbb{C}^{n_i}$ for all i . From this, one concludes that X has no cohomology in odd degrees. This method for breaking down a projective variety into pieces, also known as *BB-decomposition*, allows to compute important topological invariants, e.g. Betti numbers. It is worth emphasizing that many of the ideas of [BB1] extend to the singular case. In fact, the BB-decomposition makes sense even if X is singular, though, this time, the cells need not be so well behaved.

Goresky, Kottwitz and MacPherson in their seminal paper [GKM], developed a theory, nowadays called GKM theory, that makes it possible to describe the equivariant cohomology of certain *T-skeletal* varieties: projective algebraic varieties upon which an algebraic torus T acts with a finite number of fixed points and invariant curves. Cohomology, in this article, is considered with rational coefficients. Let X be a T -skeletal variety and denote by X^T the fixed point set. The main purpose of GKM theory is to

2010 *Mathematics Subject Classification*. Primary 14F43, 14L30; Secondary 55N91, 14M15.

Key words and phrases. rational smoothness, algebraic torus actions, GKM theory, equivariant cohomology, algebraic monoids, group embeddings.

identify the image of the functorial map

$$i^* : H_T^*(X) \rightarrow H_T^*(X^T),$$

assuming X has no cohomology in odd degrees (*equivariantly formal*). GKM theory asserts that if X is a *GKM variety*, i.e. T -skeletal and equivariantly formal, then the equivariant cohomology ring $H_T^*(X)$ can be identified with certain ring of piecewise polynomial functions $PP_T^*(X)$ (Theorem 2.5).

Mostly, GKM theory has been applied to smooth projective T -skeletal varieties, because they all have trivial cohomology in odd degrees (BB-decomposition). Furthermore, the GKM data issued from the fixed points and invariant curves has been explicitly obtained for some interesting subclasses: flag varieties ([C], [Br1]), toric varieties ([Br2], [VV], [U]) and regular embeddings of reductive groups ([Br3], [U]). Additionally, GKM theory has been applied to Schubert varieties ([C], [Br1], [Br5]). The latter ones, even though singular, are GKM varieties and their BB-cells (relative to an appropriate action of \mathbb{C}^*) are exactly the Bruhat cells.

Let now X be a complex algebraic variety of dimension n , and $x \in X$. We say that X is *rationally smooth at x* , if there exists a neighborhood U of x (in the complex topology) such that, for all $y \in U$, we have

$$H^m(X, X - \{y\}) = (0) \text{ if } m \neq 2n, \text{ and } H^{2n}(X, X - \{y\}) = \mathbb{Q}.$$

If X is rationally smooth at every $x \in X$, then X is called *rationally smooth*. Such varieties satisfy Poincaré duality with rational coefficients [M]. See [Br4] for an up-to-date discussion of rationally smooth singularities on complex algebraic varieties with torus action.

Let G be a connected reductive group. Recall that a normal irreducible projective variety X is called an *embedding* of G , or a *group embedding*, if X is a $G \times G$ -variety containing an open orbit isomorphic to G . Let M be a reductive monoid with zero and unit group G . Then there exists a central one-parameter subgroup $\epsilon : \mathbb{C}^* \rightarrow G$, with image Z contained in the center of G , such that $\lim_{t \rightarrow 0} \epsilon(t) = 0$. Moreover, the quotient space

$$\mathbb{P}_\epsilon(M) := (M \setminus \{0\})/Z$$

is a normal projective embedding of the quotient group G/Z . Embeddings of the form $\mathbb{P}_\epsilon(M)$ are called *standard group embeddings*. It is known that all normal projective embeddings of a connected reductive group are standard ([AB]). Using methods from the theory of algebraic monoids, Renner ([R2], [R3]) investigated those standard embeddings that are rationally smooth.

The purpose of this article is to establish GKM theory in the setting of *\mathbb{Q} -filtrable varieties*: projective varieties with a torus action having finitely many fixed points, such that the cells of the associated BB-decomposition are all rationally smooth, i.e. rational cells. In general, \mathbb{Q} -filtrable varieties have singularities. As an application of our theory, we show that rationally

smooth standard embeddings are \mathbb{Q} -filtrable. Our results lay down the topological foundations for the study of rationally smooth standard embeddings via GKM theory.

This article is organized as follows. The first two sections briefly review GKM theory. In Section 3, we devote ourselves to the study of rational cells and state their main topological features (Thm. 3.9, Prop. 3.11, and Thm. 3.15). In Section 4 we introduce the notion \mathbb{Q} -filtrable varieties. Our main result in this section is given below.

Theorem 4.8. *Let X be a normal projective T -variety. Suppose that X is \mathbb{Q} -filtrable. Then*

- (a) *X admits a filtration into T -stable closed subvarieties X_i , $i = 0, \dots, m$, such that*

$$\emptyset = X_0 \subset X_1 \subset \dots \subset X_{m-1} \subset X_m = X.$$

- (b) *each cell $C_i = X_i \setminus X_{i-1}$ is a rational cell, for $i = 1, \dots, m$.*
 (c) *For each $i = 1, \dots, m$, the singular rational cohomology of X_i vanishes in odd degrees. In other words, each X_i is equivariantly formal.*
 (d) *If, in addition, the T -action on X is T -skeletal, then each X_i is a GKM-variety.*

It is worth noting that \mathbb{Q} -filtrable spaces need not be rationally smooth. For instance, Schubert varieties admit a decomposition into affine cells but they are not always rationally smooth. As a way of example, the Schubert variety of codimension one in the Grassmannian of 2-planes in \mathbb{C}^4 does not satisfy Poincaré duality, and hence is not rationally smooth. References [Ar] and [Br2] supply some criteria for rational smoothness of Schubert varieties. What is remarkable about \mathbb{Q} -filtrable varieties is that they are equivariantly formal.

In Section 6, after recalling Arabia's notion of equivariant Euler classes (Section 5), we construct free module generators on the equivariant cohomology of any \mathbb{Q} -filtrable GKM variety. Our findings extend the earlier works of Arabia ([Ar]) and Guillemin-Kogan ([GK]). The main result of Section 6 is the following.

Theorem 6.9. *Let X be a \mathbb{Q} -filtrable GKM-variety. Let $x_1 < x_2 < \dots < x_m$ be the order relation on X^T compatible with the filtration of X given in Theorem 4.8. Then there exist unique classes $\theta_i \in H_T^*(X)$, $i = 1, \dots, m$, with the following properties:*

- (i) $I_i(\theta_i) = 1$,
 (ii) $I_j(\theta_i) = 0$ for all $j \neq i$,
 (iii) the restriction of θ_i to $x_j \in X^T$ is zero for all $j < i$, and
 (iv) $\theta_i(x_i) = \text{Eu}_T(i, C_i)$.

Moreover, the θ_i 's generate $H_T^*(X)$ freely as a module over $H_T^*(pt)$.

Here $I_i : H_T^*(X) \rightarrow H_T^*(pt)$ is the H_T^* -linear map obtained by integrating, along X_i , the pullback of a class in X to X_i , and $\text{Eu}_T(i, C_i)$ stands for the equivariant Euler class.

Although the class of \mathbb{Q} -filtrable varieties includes smooth projective T -skeletal varieties and Schubert varieties, its crucial attribute is that it also includes a large and interesting family of singular group embeddings, namely, rationally smooth standard embeddings. Indeed, in the last section of this article, we show that the notion of \mathbb{Q} -filtrable variety is well suited to the study of group embeddings and, in doing so, we provide our theory with its major set of fundamental examples. Our main result in this direction can be stated as follows.

Theorem 7.3. *Let $X = \mathbb{P}_e(M)$ be a standard group embedding. If X is rationally smooth, then X is \mathbb{Q} -filtrable and so it has no cohomology in odd degrees.*

Acknowledgments. Some of these results were part of the author's doctoral dissertation under the supervision of Lex Renner. I would like to thank him and Michel Brion for their invaluable help.

1. EQUIVARIANT COHOMOLOGY AND LOCALIZATION

Throughout this article, we work with complex algebraic varieties. Cohomology is always considered with rational coefficients.

1.1. The Borel construction. Let $T = (\mathbb{C}^*)^r$ be an algebraic torus and let X be a T -variety, that is, a complex algebraic variety with an algebraic action of T . Let $ET \rightarrow BT$ be a universal principal bundle for T . The **equivariant cohomology** of X (with rational coefficients) is defined to be

$$H_T^*(X) := H^*(X_T),$$

where $X_T = (X \times ET)/T$ is the total space associated to the fibration

$$X \hookrightarrow X_T \xrightarrow{p_X} BT .$$

This construction was introduced by Borel [Bo]. Here, BT is simply connected, the map p_X is induced by the canonical projection $ET \times X \rightarrow ET$, and T acts diagonally on $ET \times X$. Notice that $H_T^*(X)$ is, via p_X^* , an algebra over $H_T^*(pt)$. To simplify notation, we sometimes write H_T^* instead of $H_T^*(pt)$.

It can be shown that $H_T^*(X)$ is independent of the choice of universal T -bundle. See [Bo] and [Q] for more details.

Example 1.1. Let $T = (\mathbb{C}^*)^r$ be an algebraic torus. In this case, $BT = (\mathbb{C}P^\infty)^r$, and consequently $H_T^*(pt) = H^*(BT) = \mathbb{Q}[x_1, \dots, x_r]$, where $\deg(x_i) = 2$. A more intrinsic description of $H_T^*(pt)$ is given as follows. Denote by $\Xi(T)$ the character group of T . Any $\chi \in \Xi(T)$ defines a one-dimensional complex

representation of T with space \mathbb{C}_χ . Here T acts on \mathbb{C}_χ via $t \cdot z := \chi(t)z$. Consider the associated complex line bundle

$$L(\chi) := (E_T \times_T \mathbb{C}_\chi \rightarrow BT)$$

and its first Chern class $c(\chi) \in H^2(BT)$. Let S be the symmetric algebra over \mathbb{Q} of the group $\Xi(T)$. Then S is a polynomial ring on r generators of degree 1, and the map $\chi \rightarrow c(\chi)$ extends to a ring isomorphism

$$c : S \rightarrow H_T^*(pt)$$

which doubles degrees: the *characteristic homomorphism* ([Bo]).

1.2. Localization Theorem for torus actions. Let $S \subset H_T^*$ be the multiplicative system $H_T^* \setminus \{0\}$. For a given T -variety X , denote by X^T the fixed point set. The following is a classical theorem due to Borel ([Bo]). See also [Hs], Theorem III.1.

Theorem 1.2. *Let X be a T -variety. Suppose $H_T^*(X)$ is a finite H_T^* -module. Then the localized restriction homomorphism*

$$S^{-1}H_T^*(X) \longrightarrow S^{-1}H_T^*(X^T) = H^*(X^T) \otimes_{\mathbb{Q}} (S^{-1}H_T^*)$$

is an isomorphism. □

2. GKM THEORY

GKM theory is a relatively recent tool that owes its name to the work of Goresky, Kottwitz and MacPherson [GKM]. This theory encompasses techniques that date back to the early works of Atiyah ([At], [AS]), Segal ([Seg]), Borel ([Bo]) and Chang-Skjelbred ([CS]).

2.1. Equivariant formality.

Definition 2.1. Suppose an algebraic torus T acts on a (possibly singular) space X . Let $p_X : X_T \rightarrow BT$ be the fibration associated to the Borel construction. We say that X is **equivariantly formal** if the Serre spectral sequence

$$E_2^{p,q} = H^p(BT; H^q(X)) \implies H_T^{p+q}(X)$$

for this fibration degenerates at E_2 .

The following theorem characterizes equivariant formality. For a proof, see [GKM], Theorem 1.6.2, or [Br5], Lemma 1.2.

Theorem 2.2. *Consider the following conditions for a T -variety X .*

- (a) X is equivariantly formal.
- (b) The edge homomorphism $H_T^*(X) \rightarrow H^*(X)$ is surjective; that is, the ordinary rational cohomology is given by extension of scalars,

$$H^*(X) \simeq H_T^*(X) \otimes_{H_T^*} \mathbb{Q}.$$

- (c) $H_T^*(X, \mathbb{Q})$ is a free $H_T^*(pt)$ -module.
- (d) The singular rational cohomology of X vanishes in odd degrees.

Then $(a) \Leftrightarrow (b) \Leftrightarrow (c) \Leftarrow (d)$.

If X^T is finite, then all these conditions are equivalent. \square

It follows that X is equivariantly formal if and only if there is an isomorphism of H_T^* -modules between $H_T^*(X)$ and $H^*(X) \otimes_{\mathbb{Q}} H_T^*$. Every smooth projective T -variety is equivariantly formal ([GKM], Theorem 14.1 (7)).

A joint application of Theorem 1.2 and Theorem 2.2 leads to the following.

Corollary 2.3. *Let X be a T -variety with a finite number of fixed points. Then X is equivariantly formal if and only if $H_T^*(X)$ is a free H_T^* -module of rank $|X^T|$, the number of fixed points.* \square

2.2. T -Skeletal Actions.

Definition 2.4. Let X be a projective T -variety. Let $\mu : T \times X \rightarrow X$ be the action map. We say that μ is a **T -skeletal action** if

- (1) X^T is finite, and
- (2) The number of one-dimensional orbits of T on X is finite.

In this context, X is called a **T -skeletal variety**. If a T -skeletal variety X is also equivariantly formal, then we say that X is a **GKM variety**.

Let X be a normal projective T -skeletal variety. Then X has an equivariant embedding into a projective space with a linear action of T ([Su], Theorem 1), and so the closure of any orbit of dimension one in X contains exactly two fixed points. Accordingly, it is possible to define a ring $PP_T^*(X)$ of **piecewise polynomial functions**. Indeed, let $R = \bigoplus_{x \in X^T} R_x$, where R_x is a copy of the polynomial algebra H_T^* . We then define $PP_T^*(X)$ as the subalgebra of R defined by

$$PP_T^*(X) = \{(f_1, \dots, f_n) \in \bigoplus_{x \in X^T} R_x \mid f_i \equiv f_j \text{ mod } (\chi_{i,j})\}$$

where x_i and x_j are the two *distinct* fixed points in the closure of the one-dimensional T -orbit $\mathcal{C}_{i,j}$, and $\chi_{i,j}$ is the character of T associated with $\mathcal{C}_{i,j}$. The character $\chi_{i,j}$ is uniquely determined up to sign (permuting the two fixed points changes $\chi_{i,j}$ to its opposite).

Theorem 2.5 ([CS], [At], [GKM]). *Let X be a normal projective T -skeletal variety. Suppose that X is a GKM variety. Then the restriction mapping*

$$H_T^*(X) \longrightarrow H_T^*(X^T) = \bigoplus_{x_i \in X^T} H_T^*$$

is injective, and its image is the subalgebra $PP_T^(X)$.* \square

Theorem 2.2 characterizes normal projective GKM-varieties among all T -skeletal varieties.

Theorem 2.6. *Let X be a normal projective variety with a T -skeletal action $\mu : T \times X \rightarrow X$. Then X is a GKM-variety if and only if X has no (rational) cohomology in odd degrees.* \square

We will show that the class of equivariantly formal spaces incorporates certain subclass of singular varieties, namely, \mathbb{Q} -filtrable varieties (Theorems 4.8 and 6.9). This subclass encompasses all rationally smooth standard embeddings of a reductive group (Theorem 7.3). As such, it is much larger than the subclass of smooth varieties.

3. RATIONAL CELLS

This section is devoted to the study of our most important topological tool: rational cells.

Definition 3.1. Let X be an algebraic variety with an action of a torus T and a fixed point x . We say that x is an **attractive fixed point** if there exists a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ and a neighborhood U of x , such that $\lim_{t \rightarrow 0} \lambda(t) \cdot y = x$ for all points y in U .

There is an important characterization of attractive fixed points. A proof of the following result can be found in [Br4], Proposition A2.

Proposition 3.2. *For a torus T acting on a variety X with a fixed point x , the following conditions are equivalent:*

- (i) *The weights of T in the Zariski tangent space $T_x(X)$ are contained in an open half space.*
- (ii) *There exists a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ such that, for all y in a neighborhood of x , we have $\lim_{t \rightarrow 0} \lambda(t)y = x$.*

If (ii) holds, then the set

$$X_x := \{y \in X \mid \lim_{t \rightarrow 0} \lambda(t)y = x\}$$

is the unique affine T -invariant open neighborhood of x in X . Moreover, X_x admits a closed T -equivariant embedding into $T_x X$. \square

Lemma 3.3. *Let X be an irreducible affine variety with a T -action and an attractive fixed point $x_0 \in X$. Then X is rationally smooth at x_0 if and only if X is rationally smooth everywhere.*

Proof. If X is rationally smooth everywhere, then it is rationally smooth at x_0 . For the converse, we use Proposition 3.2 (ii) and the affineness of X to guarantee the existence of a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ such that

$$X = \{y \in X \mid \lim_{t \rightarrow 0} \lambda(t)y = x_0\}.$$

In symbols, $x_0 \in \overline{\mathbb{C}^* \cdot y}$, for any $y \in X$. Now consider the complex topology on X . We claim that any non-empty open T -stable subset of X containing x_0 is all of X . In effect, let U be a T -stable neighborhood of x_0 . Then, for any $y \in X$, there exists $s_y \in \mathbb{C}^*$, such that $s_y \cdot y \in U$. Indeed, because x_0 is attractive, one can find a sequence $\{t_n\} \subset \mathbb{C}^*$ such that $t_n \cdot y$ converges to x_0 . That is, there exists N with the property that $t_N \cdot y$ belongs to U . Setting $s_y = t_N$ yields $s_y \cdot y \in U$. However, U is T -stable, and therefore it contains the entire orbit $\mathbb{C}^* \cdot y$. In short, $y \in U$ or, equivalently, $U = X$.

Hence, the non-empty open T -stable subset of rationally smooth points of X is, *a fortiori*, equal to X . \square

Definition 3.4. Let X be an irreducible affine variety with a T -action and an attractive fixed point $x_0 \in X$. If X is rationally smooth at x_0 (and thus everywhere), we refer to (X, x_0) as a **rational cell**.

It follows from Definition 3.4 and Proposition 3.2 that if (X, x_0) is a rational cell, then

$$X = \{y \in X \mid \lim_{t \rightarrow 0} \lambda(t)y = x_0\},$$

for a suitable one-parameter subgroup λ . Notably, $\{x_0\}$ is the unique closed T -orbit in X .

Example 3.5. Certainly \mathbb{C}^n is a rational cell with the usual \mathbb{C}^* -action by scalar multiplication. Here the origin is the unique attractive fixed point.

Example 3.6. Let $V = \{xy = z^2\} \subset \mathbb{C}^3$. The standard \mathbb{C}^* -action by scalar multiplication makes V a rational cell with $(0, 0, 0)$ as its attractive fixed point. This is clear once we observe that V is the quotient of \mathbb{C}^2 by the finite group with two elements, where the non-trivial element acts on $(s, t) \in \mathbb{C}^2$ via $(s, t) \mapsto (-s, -t)$. So Proposition A1 (iii) of [Br4] implies that V is rationally smooth.

Example 3.7. A normal variety is not necessarily rationally smooth. For instance, consider the hypersurface $H \subset \mathbb{C}^4$ defined by $\{xy = uv\}$. Because the singular locus of H , namely $\{(0, 0, 0, 0)\}$, has codimension three, it follows that H is normal ([Sha], p. 128, comments after Theorem II.5.1.3). Nevertheless, H is not rationally smooth at the origin. To see this, let $T = (\mathbb{C}^*)^2$ act on H via $(t, s) \cdot (x, y, u, v) = (tx, ts^2y, su, st^2v)$. Then H has the origin as its unique attractive fixed point. Moreover, H contains four T -invariant curves (the four coordinate axes) passing through $(0, 0, 0, 0)$. If H were rationally smooth at the origin, then, by a result of Brion (Theorem 3.15), the dimension of H would equal the number of its T -invariant curves. This is a contradiction, since H is only three dimensional.

Definition 3.8. Let Z be a rationally smooth complex projective variety. Let n be the (complex) dimension of Z . We say that Z is a **rational cohomology complex projective space** if there is a ring isomorphism

$$H^*(Z) \simeq \mathbb{Q}[t]/(t^{n+1}),$$

where $\deg(t) = 2$.

Let (X, x) be a rational cell. Then, by Proposition 3.2, X admits a closed T -equivariant embedding into $T_x X$. Set \dot{X} to be $X - \{x\}$. Choose an injective one-parameter subgroup $\lambda: \mathbb{C}^* \rightarrow T$ as in Definition 3.4. Then all weights of the \mathbb{C}^* -action on $T_x X$ via λ are positive. Thus, the quotient

$$\mathbb{P}(X) := \dot{X}/\mathbb{C}^*$$

exists and is a projective variety ([Br4]). Indeed, it is a closed subvariety of $\mathbb{P}(T_x X)$, a weighted projective space. The variety $\mathbb{P}(X)$ can be viewed as an algebraic version of the link of X at x .

The following result, except for parts (b) and (c), is due to Brion ([Br4]). The idea of the proof of part (b) is due to Renner.

Theorem 3.9. *Let (X, x_0) be a rational cell of dimension n . Then,*

(a) X is contractible.

(b) $X - \{x_0\}$ is homeomorphic to $\mathbb{S}(X) \times \mathbb{R}^+$, where $\mathbb{S}(X) := X - \{x_0\}/\mathbb{R}^+$ is a compact topological space.

(c) $X - \{x_0\}$ deformation retracts to $\mathbb{S}(X)$. In addition, X is rationally smooth at x_0 if and only if $X - \{x_0\}$, and thus $\mathbb{S}(X)$, is a rational cohomology sphere \mathbb{S}^{2n-1} .

(d) The space $\mathbb{P}(X) = X - \{x_0\}/\mathbb{C}^*$ is a rationally smooth complex projective variety of dimension $n - 1$. Furthermore, X is rationally smooth if and only if $\mathbb{P}(X)$ is a rational cohomology complex projective space $\mathbb{C}\mathbb{P}^{n-1}$.

Proof. For part (a) simply notice that the action of \mathbb{C}^* on X extends to a map $\mathbb{C} \times X \rightarrow X$ sending $0 \times X$ to x_0 and restricting to the identity $1 \times X \rightarrow X$. Since the proof of (d) can be found in [Br4], Lemma 1.3, it suffices to prove parts (b) and (c).

(b) From Proposition 3.2, we know that X admits a closed T -equivariant embedding into $T_{x_0} X \simeq \mathbb{C}^d$, which identifies x_0 with 0. Choosing a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ as in Definition 3.4 yields a \mathbb{C}^* -action on \mathbb{C}^d with only positive weights m_1, \dots, m_d . Specifically, $\lambda \in \mathbb{C}^*$ acts on \mathbb{C}^d via

$$\lambda \cdot (z_1, \dots, z_d) = (\lambda^{m_1} z_1, \dots, \lambda^{m_d} z_d).$$

Next, define an \mathbb{R}^+ -equivariant map $N : \mathbb{C}^d \rightarrow \mathbb{R}$ by

$$N(z_1, \dots, z_d) = \sqrt{\sum_{i=1}^d (z_i \bar{z}_i)^{1/m_i}}.$$

Clearly, for $\lambda \in \mathbb{C}$ and $z \in \mathbb{C}^d$, the definition favors $N(\lambda \cdot z) = |\lambda|N(z)$ (here $\lambda \cdot z$ means $(\lambda^{m_1} z_1, \dots, \lambda^{m_d} z_d)$).

Since \mathbb{R}^+ acts freely on $X - \{0\} \subseteq \mathbb{C}^d - \{0\}$, the quotient map

$$X - \{0\} \rightarrow \mathbb{S}(X)$$

is a principal \mathbb{R}^+ -fibration. We claim that this fibration is trivial, i.e.

$$X - \{0\} \simeq \mathbb{S}(X) \times \mathbb{R}^+.$$

To prove the claim, we just need to provide a global section s . In fact, we can do so canonically. Let $s : \mathbb{S}(X) \rightarrow X - \{0\}$ be the map defined by

$$s([x]) = \frac{1}{N(x)} \cdot x.$$

This map is well defined (given that we are using the \mathbb{C}^* -action mentioned above) and not only defines a global section, but also a homeomorphism between $\mathbb{S}(X)$ and $X \cap N^{-1}(1)$, where $N^{-1}(1)$ is the “unit” sphere. Thus, $\mathbb{S}(X)$ is compact.

(c) The first claim follows immediately from part (b). As for the second assertion, remember that X is contractible. Thus, the long exact sequence of the pair $(X, X - \{x_0\})$ splits into short exact sequences

$$0 \longrightarrow H^*(X - \{x_0\}) \longrightarrow H^{*+1}(X, X - \{x_0\}) \longrightarrow 0.$$

Therefore X is rationally smooth if and only if $X - \{x_0\}$ is a rational homology sphere of dimension $2n - 1$. \square

Corollary 3.10. *Keeping the same notation as in Theorem 3.9, the rational cell X is homeomorphic to the open cone over $\mathbb{S}(X)$. Moreover, $\mathbb{P}(X)$ is equivariantly formal.*

Proof. The first assertion follows from Theorem 3.9 (c). As for the second, by Theorem 3.9 again, $\mathbb{P}(X)$ is a rational cohomology complex projective space and thus has no cohomology in odd degrees. Theorem 2.2 concludes the proof. \square

Proposition 3.11. *Let (X, x_0) be a rational cell of dimension n . Denote by X^+ its one point compactification. Then X^+ is simply connected and has the rational homotopy type of \mathbb{S}^{2n} , the Euclidean $2n$ -sphere.*

Proof. First, observe that X^+ is path-connected. As a consequence of Theorem 3.9, we can write X^+ as a union of two open cones D_0 and D_∞ ; namely, $D_0 = S \times [0, 1]/S \times \{0\}$ and $D_\infty = S \times (\epsilon, \infty]/S \times \{\infty\}$, where S stands for $\mathbb{S}(X) = (X \setminus \{x_0\})/\mathbb{R}_+$, and ϵ is a positive number less than 1. Given that $X - \{x_0\}$ is path-connected, the intersection $D_0 \cap D_\infty = S \times (\epsilon, 1)$ is path-connected as well. In summary, X^+ can be written as the union of two contractible open subsets with path-connected intersection. Thus, by van Kampen’s theorem, X^+ itself is simply connected. To finish the proof, we need to show that X^+ is a rational cohomology $2n$ -sphere. This is a simple exercise, using the Mayer-Vietoris sequence of the cover $\{D_0, D_\infty\}$. \square

Lemma 3.12 (One-dimensional rational cells). *Let (X, x) be a rational cell of dimension one. Then*

- (1) X is a cone over a topological circle.
- (2) X is homeomorphic to \mathbb{C} .
- (3) If, additionally, X is normal, then X is isomorphic to \mathbb{C} as an algebraic variety.

Proof. Without loss of generality, we can assume that T acts faithfully on X . Thus, T is isomorphic to \mathbb{C}^* . Now assertions (1) and (2) can be proved as follows. Since X is one-dimensional, then the singular locus is an invariant discrete set. Nonetheless, x_0 is the unique attractive fixed point, and \mathbb{C}^* is connected, so the singular locus is either empty or consists of only one point,

namely, x_0 . As a result, $X \setminus \{x_0\}$ is smooth. Next notice that X has two \mathbb{C}^* -orbits: the attractive fixed point x_0 , and a dense open orbit of the form \mathbb{C}^* . Hence, X is homeomorphic to \mathbb{C} and it is a cone over the circle S^1 .

Finally, if we also assume that X is normal and one-dimensional, then *a fortiori* X is smooth ([Har]). This proves (3). \square

Lemma 3.13. *Let (X, x) be a rational cell. Suppose x is a smooth point. Then X is isomorphic to its tangent space at x .*

Proof. By Proposition 3.2, we know that X admits an equivariant *closed* embedding into $T_x X$. If x is a smooth point, then both X and $T_x X$ have the same dimension. For affine varieties this can only happen if $X = T_x X$. \square

We are now ready to state what we call the Equivariant Normalization Theorem for rational cells. It is due to Brion ([Br2], Proof of Theorem 18, implication (i) \Rightarrow (ii)) and Arabia ([Ar], Section 3.2.1).

Theorem 3.14. *Let (X, x) be a rational cell. Then there exists a T -module V and an equivariant finite surjective map $\pi : X \rightarrow V$ such that $\pi(x) = 0$ and $V^T = \{0\}$.* \square

We now specialize a result of Brion ([Br4], Section 1.4, Corollary 2) to rational cells.

Theorem 3.15. *Let (X, x) be a rational cell. Suppose that the number of closed irreducible T -stable curves on X is finite. Let $n(X, x)$ be this number. Then $n(X, x) = \dim(X)$.* \square

4. HOMOLOGY AND BETTI NUMBERS OF \mathbb{Q} -FILTRABLE SPACES

We define \mathbb{Q} -filtrable varieties, spaces that come equipped with a paving by rational cells, and show that they are equivariantly formal (Theorem 4.8).

4.1. The Bialynicki-Birula decomposition. Let X be a projective algebraic variety with a \mathbb{C}^* -action and a finite number of fixed points x_1, \dots, x_m . Consider the associated BB-decomposition $X = \bigsqcup_i W_i$, where each cell is defined as follows

$$W_i = \{x \in X \mid \lim_{t \rightarrow 0} t \cdot x = x_i\}.$$

In the present section, we show that rational cells are a good substitute for the notion of affine space in the topological study of singular varieties.

Remark 4.1. In general, the *BB*-decomposition of a projective variety is not a stratification; that is, it may happen that the closure of a cell is not the union of cells, even if we assume our variety to be smooth. For a justification of this claim, see [BB2].

Definition 4.2. Let X be a complex algebraic variety endowed with a \mathbb{C}^* -action and a finite number of fixed points. A **BB-decomposition** $\{W_i\}$ is said to be **filtrable** if there exists a finite increasing sequence $X_0 \subset X_1 \subset \dots \subset X_m$ of closed subvarieties of X such that:

- a) $X_0 = \emptyset$, $X_m = X$,
 b) For each $j = 1, \dots, m$, the “stratum” $X_j \setminus X_{j-1}$ is a cell of the decomposition $\{W_i\}$.

The following result is due to Bialynicki-Birula ([BB2]).

Theorem 4.3. *Let X be a normal projective variety with a torus action and a finite number of fixed points. Then the stable decomposition is filtrable. \square*

4.2. \mathbb{Q} -filtrable spaces. To begin with, let us introduce a few technical results.

Lemma 4.4. *Let X be a complex projective algebraic variety with a \mathbb{C}^* -action. Suppose X can be decomposed as the disjoint union*

$$X = Y \sqcup C,$$

where Y is a closed stable subvariety and C is an open rational cell containing a fixed point of X , say c_0 , as its unique attractive fixed point. Denote by n the (complex) dimension of C . Then

$$H^k(X, Y) = \begin{cases} 0 & \text{if } k \neq 2n \\ \mathbb{Q} & \text{if } k = 2n. \end{cases}$$

Furthermore, if Y has vanishing odd cohomology, then

$$H^k(X, \mathbb{Q}) = \begin{cases} H^k(Y, \mathbb{Q}) & \text{if } k \neq 2n \\ H^{2n}(Y, \mathbb{Q}) \oplus \mathbb{Q} & \text{if } k = 2n. \end{cases}$$

Proof. Let $H_c^*(-)$ denote cohomology with compact supports. It is well-known that $H^*(X) = H_c^*(X)$ and $H^*(Y) = H_c^*(Y)$, because X and Y are complex projective varieties. Moreover, by Corollary B.14 of [PS], one has

$$H^*(X, Y) \simeq H_c^*(X - Y) = H_c^*(C).$$

Given that C is a rational cell, and a cone over a rational cohomology sphere of dimension $2n - 1$ (Corollary 3.10), it follows easily that

$$H_c^*(C) = H^*(C, C - \{c_0\}) = \begin{cases} 0 & \text{if } k \neq 2n \\ \mathbb{Q} & \text{if } k = 2n. \end{cases}$$

So the first claim is proved.

As for the second assertion, consider the long exact sequence of the pair (X, Y) , namely,

$$\dots \rightarrow H^{*-1}(Y) \rightarrow H^*(X, Y) \rightarrow H^*(X) \rightarrow H^*(Y) \rightarrow H^{*+1}(X, Y) \rightarrow \dots$$

By our previous remarks, this long exact sequence can be rewritten as

$$\dots \rightarrow H^{*-1}(Y) \rightarrow H_c^*(C) \rightarrow H^*(X) \rightarrow H^*(Y) \rightarrow H_c^{*+1}(C) \rightarrow \dots$$

If Y has no cohomology in odd degrees, then the long exact sequence splits, yielding the identifications $H^i(X) = H^i(Y)$, when $i \neq 2n$, and

$$H^{2n}(X) = H^{2n}(Y) \oplus H_c^{2n}(C) = H^{2n}(Y) \oplus \mathbb{Q}.$$

The proof is now complete. \square

Corollary 4.5. *Keeping the notation of Lemma 4.4, attaching a complex n -dimensional rational cell produces no changes in cohomology, except in degrees $2n-1$ and $2n$. Furthermore, if Y has no cohomology in odd degrees, then X has no odd cohomology either, and there is a short exact sequence of the form*

$$0 \longrightarrow H_c^{2n}(C) \longrightarrow H^{2n}(X) \longrightarrow H^{2n}(Y) \rightarrow 0.$$

Proof. We simply observe that the long exact sequence of the pair (X, Y) gives

$$H^k(X) \simeq H^k(Y)$$

for $k \neq 2n-1, 2n$. Besides, we also obtain the exact sequence

$$0 \rightarrow H^{2n-1}(X) \rightarrow H^{2n-1}(Y) \rightarrow H_c^{2n}(C) = \mathbb{Q} \rightarrow H^{2n}(X) \rightarrow H^{2n}(Y) \rightarrow 0.$$

So in general $H^{2n-1}(X)$ injects into $H^{2n-1}(Y)$. In case we assume Y to have vanishing odd cohomology, we obtain X with vanishing odd cohomology as well, and a “lifting of generators” sequence:

$$0 \longrightarrow H_c^{2n}(C) \longrightarrow H^{2n}(X) \longrightarrow H^{2n}(Y) \rightarrow 0.$$

□

Corollary 4.6. *Let X be a normal complex projective variety endowed with a \mathbb{C}^* -action and a finite number of fixed points. Suppose that each BB -cell is a rational cell. Then X has vanishing odd cohomology over the rationals, and the dimension of its cohomology group in degree $2k$ equals the number of rational cells of complex dimension k . Furthermore, X is equivariantly formal and $\chi(X) = |X^T|$.*

Proof. Since the BB -decomposition on X is filtrable, the result follows from the previous lemma as we move up in the filtration by attaching one rational cell at the time. This process is systematic and preserves cohomology in lower and higher degrees at each step. □

Let T be an algebraic torus acting on a variety X . A one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ is called *generic* if $X^{\mathbb{C}^*} = X^T$, where \mathbb{C}^* acts on X via λ . Generic one-parameter subgroups always exist. Note that the BB -cells of X , obtained using λ , are T -invariant.

Our results in this section suggest the following definition.

Definition 4.7. Let X be a projective variety equipped with a T -action. We say that X is **\mathbb{Q} -filtrable** if

- (1) X is normal,
- (2) the fixed point set X^T is finite, and
- (3) there exists a generic one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ for which the associated BB -decomposition of X consists of *rational cells*.

Theorem 4.8. *Let X be a normal projective T -variety. Suppose that X is \mathbb{Q} -filtrable. Then*

(a) X admits a filtration into closed subvarieties X_i , $i = 0, \dots, m$, such that

$$\emptyset = X_0 \subset X_1 \subset \dots \subset X_{m-1} \subset X_m = X.$$

(b) each cell $C_i = X_i \setminus X_{i-1}$ is a rational cell, for $i = 1, \dots, m$.

(c) For each $i = 1, \dots, m$, the singular rational cohomology of X_i vanishes in odd degrees. In other words, each X_i is equivariantly formal.

(d) If, in addition, the T -action on X is T -skeletal, then each X_i is a GKM-variety.

Proof. Assertions (a) and (b) are a direct consequence of Definition 4.7 and Theorem 4.3. Applying Corollary 4.6 and Theorem 2.2 at each step of the filtration yields claim (c). For statement (d), we argue as follows. Notice that all the X_i 's have vanishing odd cohomology, as it is guaranteed by (c). Moreover, since the X_i 's are T -invariant and the T -action on X is T -skeletal, then each X_i contains only a finite number of fixed points and T -invariant curves. In consequence, Theorem 2.5 applied to each X_i gives (d). \square

Thus, we obtain the applicability of GKM-theory at each step of the filtration, even though the various X_i 's are not necessarily rationally smooth. This approach is more flexible than the general approach (by comparing singular cohomology with intersection cohomology), used, for instance, in Theorem 3.7 of [R3] or in [W]. Such flexibility will become apparent from our results in Section 6, where we supply a method for constructing free module generators on the equivariant cohomology of \mathbb{Q} -filtrable GKM-varieties (Theorem 6.9). This method is based on the notion of equivariant Euler classes.

5. EQUIVARIANT EULER CLASSES

We quickly review the theory of equivariant Euler classes. For a complete treatment of the subject, the reader is invited to consult [Hs] and [Ar].

Let X be a T -variety with an isolated fixed point x . To fix ideas, let us first assume that (X, x) is a rational cell of dimension n . Recall that $\mathbb{S}(X) = [X - \{x\}]/\mathbb{R}^+$ is a rational cohomology sphere \mathbb{S}^{2n-1} and that X is homeomorphic to the (open) cone over $\mathbb{S}(X)$ (Theorem 3.9 and Corollary 3.10). The Borel construction yields the fibration

$$\mathbb{S}(X) \hookrightarrow \mathbb{S}(X)_T \longrightarrow BT.$$

Observe that the E_2 -term of the corresponding Serre spectral sequence consists of only two lines, namely,

$$E_2^{p,q} = H^p(BT) \otimes H^q(\mathbb{S}(X)) \neq 0 \text{ only when } q = 0 \text{ and } q = 2n - 1.$$

Let $\text{Eu}_T(x, X) \in H^{2n}(BT)$ be the transgression of the generator λ_X of $H^{2n-1}(\mathbb{S}(X))$. We call $\text{Eu}_T(x, X)$ the **Equivariant Euler class of X at x** . It follows from [Hs], Theorem IV.6, that $\text{Eu}_T(x, X)$ splits into a product of linear polynomials, namely

$$\text{Eu}_T(x, X) = \omega_1^{k_1} \cdots \omega_s^{k_s},$$

where $\omega_i \in H^2(BT) \simeq \Xi(T) \otimes \mathbb{Q}$. Here $\Xi(T)$ stands for the character group of T , and the isomorphism is provided in Example 1.1.

Since X is a cone over $\mathbb{S}(X)$, then $H_c^*(X) \simeq H^*(X, X - \{x\}) \simeq \mathbb{Q}$, where $H_c^*(-)$ denotes cohomology with compact supports. Using the Serre spectral sequence, one notices that these isomorphisms are also valid in equivariant cohomology:

$$H_{T,c}^*(X) \simeq H_T^*(X, X - \{x\}) \simeq H_T^*.$$

Let \mathcal{T}_X be the canonical generator of $H_T^*(X, X - \{x\})$. This generator can be described by the commutative diagram

$$\begin{array}{ccc} H_T^*(X, X - \{x\}) & \xrightarrow{i^*} & H_T^*(X) \\ \Phi_X^* \uparrow & & \downarrow \text{res} \\ & \int_{[X]} & \\ H_T^*(x) & \xrightarrow{\times(\text{Eu}_T(x, X))} & H_T^*(x), \end{array}$$

where Φ_X^* is multiplication by \mathcal{T}_X . In other words, \mathcal{T}_X is the unique class in $H_T^*(X, X - \{x\})$ whose restriction to $H_T^*(pt)$ coincides with $\text{Eu}_T(x, X)$. It is customary in the literature to call \mathcal{T}_X the **Thom class** of X . Let us bear in mind that the map Φ_X^* raises degree by $2n$. Clearly, $H_T^*(X, X - \{x\}) \simeq H_c^*(X) \otimes H_T^*(pt)$ and so, $H_{T,c}^j(X) = 0$ for $j < 2n$. As for the integral appearing here, it is, by definition, the inverse of Φ_X^* .

Let \mathcal{Q}_T be the quotient field of H_T^* . If $\mu \in H_{T,c}^*(X)$, then

$$\text{Eu}_T(x, X) \wedge \int_{[X]} \mu = \mu_x,$$

where μ_x denotes restriction of the class μ to x . Hence, the identity

$$\frac{1}{\text{Eu}_T(x, X)} = \frac{1}{\mu_x} \int_{[X]} \mu,$$

holds in \mathcal{Q}_T , for every non-zero μ in $H_T^*(X, X - \{x\})$.

More generally, let X be a T -variety with an isolated fixed point x . Suppose that X is rationally smooth at x . As pointed out in [Ar], we can replace X by a conical neighborhood U_x of x and define $\text{Eu}_T(x, X) := \text{Eu}_T(x, U_x)$. For instance, if x is also an attractive fixed point, we can let U_x be a rational cell (Proposition 3.2).

In case the isolated fixed point $x \in X$ is not necessarily a rationally smooth point, Arabia ([Ar]) has shown that we can still define an Euler class $\text{Eu}_T(x, X)$. The key ingredient here is that, by Theorem 1.2, the map

$$i^* : H_T^*(X, X - \{x\}) \rightarrow H_T^*(x)$$

is an isomorphism modulo H_T^* -torsion. Therefore, the function that assigns to a non-torsion element $\mu \in H_T^*(X, X - \{x\})$ the fraction $\frac{1}{\mu_x} \int_X \mu \in \mathcal{Q}_T$ is *constant*.

Definition 5.1. Let X be a T -variety. Suppose that $x \in X^T$ is an isolated fixed point. The fraction

$$\frac{1}{\text{Eu}_T(x, X)} := \frac{1}{\mu_x} \int_X \mu \in \mathcal{Q}_T,$$

where μ is any non-torsion element of $H_T^*(X, X - \{x\})$, is called the *inverse of the equivariant Euler class of X at x* . When this fraction is non-zero, we denote its inverse by $\text{Eu}_T(x, X)$ and call it the *Equivariant Euler class of X at x* .

From Remark 2.3-2(b) of [Ar], it follows that if x is a rationally smooth point of X , then $\text{Eu}_T(x, X)$ is a non-zero polynomial, and splits into a product of linear factors.

Example 5.2. When $X = \mathbb{C}^n$, $x = 0$, and the algebraic torus T acts linearly on \mathbb{C}^n , one proves

$$\text{Eu}_T(0, \mathbb{C}^n) = (-1)^n \prod_{\alpha \in \mathcal{A}} \alpha,$$

where \mathcal{A} is the collection of weights. Furthermore, if the weights in \mathcal{A} are pairwise linearly independent, then the associated complex projective space $\mathbb{P}(\mathbb{C}_{\mathcal{A}}^n)$ has exactly n T -fixed points: the lines \mathbb{C}_{α_i} . One also verifies that

$$\text{Eu}_T([\mathbb{C}_{\alpha_i}], \mathbb{P}(\mathbb{C}_{\mathcal{A}}^n)) = \prod_{j \neq i} (\alpha_i - \alpha_j).$$

See [Ar], Remark 2.4.1-1.

Remark 5.3. The inverse of the equivariant Euler class coincides with the equivariant multiplicity at a nondegenerate fixed point ([Br1], Section 4).

Proposition 5.4 (Atiyah-Bott Localization formula, [Ar]). *Let X be a complex projective variety. Suppose that a torus T acts on X with only a finite number of fixed points. Then*

$$\int_X \mu = \sum_{x \in X^T} \frac{\mu|_x}{\text{Eu}_T(x, X)},$$

for any $\mu \in H_T^*(X)$. □

Let (X, x) be a rational cell. Then, by Proposition 3.2, X admits a closed T -equivariant embedding into its tangent space $T_x X$. Notice that there are only a finite number of codimension-one subtori T_1, \dots, T_m of T for which $X^{T_j} \neq X^T$, since each one of them is contained in the kernel of a weight of T in $T_x X$.

Theorem 5.5 ([Ar], [Br2]). *Let (X, x) be a rational cell of dimension n . Let $\pi : X \rightarrow \mathbb{C}^n$ be the equivariant finite surjective map from Theorem 3.14. Then*

(a) *The induced morphism in cohomology*

$$\pi^* : H_c^{2n}(\mathbb{C}^n) \longrightarrow H_c^{2n}(X)$$

is an isomorphism and satisfies $\int_Y \pi^*(\mu) = \deg(\pi) \int_{\mathbb{C}^n} \mu$, where $\deg(\pi)$ is the cardinality of a general fibre of π . This formula also holds in equivariant cohomology, in particular

$$\mathrm{Eu}_T(0, \mathbb{C}^n) = \deg(\pi) \cdot \mathrm{Eu}_T(x_0, X).$$

(b) $\mathrm{Eu}_T(X, x) = c \prod_{T_i} \mathrm{Eu}_T(X^{T_i}, x)$, where c is a positive rational number, and the product runs over the finite number of codimension-one subtori T_i of T for which $X^{T_i} \neq X^T$.

Proof. We refer the reader to [Ar], Proposition 3.2.1-1, for the proof of part (a). Finally, part (b) follows from Remark 5.3 and [Br2], Theorem 18 (iii). \square

Let (X, x) be a rational cell. At the beginning of this section it was shown that $\mathrm{Eu}_T(x, X)$ splits into a product of characters. The following result provides a geometric interpretation of this factorization.

Corollary 5.6. *Let (X, x) be a rational cell of dimension n . Suppose that X contains only a finite number of closed irreducible T -invariant curves \mathcal{C}_i , $i = 1, \dots, n$. Let χ_i be the character associated with the action of T on \mathcal{C}_i . Then*

$$\mathrm{Eu}_T(x, X) = c \cdot \chi_1 \cdots \chi_n,$$

where c is a positive rational number.

Proof. There is only a finite number of codimension-one subtori T_i such that $X^{T_i} \neq X^T$. Notice that T acts on each X^{T_i} through its quotient $T/T_i \simeq \mathbb{C}^*$. Because x is an attractive fixed point of X , we can assume, without loss of generality, that x is an attractive fixed point of each X^{T_i} , for the induced action of $\mathbb{C}^* \simeq T/T_i$. It follows from Corollary 2 of Sect. 1.4 of [Br4] that $X^{T_i} = \mathcal{C}_i$. Moreover, by Theorem 1.1 of [Br4], each X^{T_i} is rationally smooth at x . Hence, each X^{T_i} is a one-dimensional rational cell with attractive fixed point x (see Lemma 3.12 for a characterization of these cells). The result can now be deduced from Theorem 5.5 and Example 5.2. \square

6. LOCAL INDICES AND MODULE GENERATORS FOR THE EQUIVARIANT COHOMOLOGY OF \mathbb{Q} -FILTRABLE GKM VARIETIES.

We supply a method for building canonical free module generators on the equivariant cohomology of any \mathbb{Q} -filtrable GKM-variety. Our findings here extend the work of Guillemin-Kogan ([GK]), on the equivariant K -theory of orbifolds, to the equivariant cohomology of a much larger class of singular varieties.

Let X be a \mathbb{Q} -filtrable GKM-variety. In other words, X is a normal projective T -variety with only a finite number of fixed points and T -invariant

curves. Moreover, there exists a *BB*-decomposition of X as a disjoint union of rational cells, say $(C_1, x_1), \dots, (C_m, x_m)$, each one containing $x_i \in X^T$ as its unique attractive fixed point. This decomposition induces a filtration of X ,

$$\emptyset = X_0 \subset X_1 \subset X_2 \dots \subset X_m = X,$$

by closed invariant subvarieties X_i , so that each difference $X_i \setminus X_{i-1}$ equals C_i , for $i = 1, \dots, m$. The key observation here is provided by Theorem 4.8. It states that every X_i is equivariantly formal and is made up of rational cells. In consequence, GKM-theory can be applied to each X_i . We will refer to X_i as the *i-th filtered piece of X*, and m will be called the *length of the filtration*.

Denote by x_1, \dots, x_m the fixed points of X . The filtration induces a total ordering of the fixed points, namely,

$$x_1 < x_2 < \dots < x_m.$$

Let (C_i, x_i) be a rational cell of X . From the previous section, we know that

$$H_{T,c}^*(C_i) \simeq H_T^*(C_i, C_i - \{x_i\}) \simeq H_T^*(x_i),$$

where the second isomorphism is provided by the Thom class \mathcal{T}_i , a well-known element of $H_T^*(C_i, C_i - \{x_i\})$. When restricted to $H_T^*(x_i)$, the Thom class \mathcal{T}_i becomes a product of linear polynomials: the Euler class $\text{Eu}_T(c_i, C_i)$.

In section 4 we built non-equivariant short exact sequences of the form

$$0 \longrightarrow H_c^{2k}(C_i) \longrightarrow H^{2k}(X_i) \longrightarrow H^{2k}(X_{i-1}) \longrightarrow 0,$$

for every i . Since the spaces involved have zero cohomology in odd degrees, then these short exact sequences naturally generalize to the equivariant case, so we also have equivariant short exact sequences

$$0 \longrightarrow H_{T,c}^{2k}(C_i) \longrightarrow H_T^{2k}(X_i) \longrightarrow H_T^{2k}(X_{i-1}) \longrightarrow 0,$$

for each i . On the other hand, by equivariant formality, the singular equivariant cohomology of each X_i injects into $H_T^*(X_i^T) = \bigoplus_{j \leq i} H_T^*(x_j)$.

In summary, for each i , we have the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_{T,c}^*(C_{i+1}) & \longrightarrow & H_T^*(X_{i+1}) & \longrightarrow & H_T^*(X_i) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_T^*(x_{i+1}) & \longrightarrow & \bigoplus_{j \leq i+1} H_T^*(x_j) & \longrightarrow & \bigoplus_{j \leq i} H_T^*(x_j) \longrightarrow 0 \end{array}$$

where the vertical maps are all injective. Indeed, such maps correspond to the various restrictions to fixed point sets. We will use this diagram to build cohomology generators. The next two lemmas are inspired in Theorem 2.3 and Proposition 4.1 of [HHH], where Kac-Moody flag varieties are studied.

Lemma 6.1. *Let X be a \mathbb{Q} -filtrable variety. Then there exists a non-canonical isomorphism of H_T^* -modules*

$$H_T^*(X) \simeq \bigoplus_{x_i \in X^T} \mathrm{Eu}_T(C_i, x_i) H_T^*(pt),$$

which is compatible with restriction to the various i -th filtered pieces $X_i \subset X$.

Proof. We argue by induction on the length of the filtration. The case $m = 1$ is simple, because it corresponds to $X = \{x_1\}$, a singleton. Assuming that we have proved the assertion for m , let us prove the case $m + 1$. Substitute $i = m$ in the commutative diagram above. Then

$$H_T^*(X_{m+1}) = H_T^*(X) \simeq H_{T,c}^*(C_{m+1}) \oplus H_T^*(X_m).$$

By induction, $H_T^*(X_m) \simeq \prod_{i \leq m} \mathrm{Eu}_T(C_i, x_i) H_T^*(pt)$. So the claim for $m + 1$ follows directly from the isomorphism $H_{T,c}^*(C_{m+1}) \simeq \mathrm{Eu}_T(C_{m+1}, x_{m+1}) H_T^*(pt)$. \square

The isomorphism of the previous lemma is not canonical because the cellular decomposition of X depends on a particular choice of generic one-parameter subgroup and a compatible ordering of the fixed points.

Given a class $\mu \in H_T^*(X)$, denote by $\mu(x_i)$ its restriction to the fixed point x_i .

Lemma 6.2. *Let X be a projective T -variety. Assume that X is \mathbb{Q} -filtrable and let $x_1 < x_2 < \dots < x_m$ be the order relation on X^T compatible with the filtration of X . For each i , let $\varphi_i \in H_T^*(X)$ be a class such that*

$$\varphi_i(x_j) = 0 \text{ for } j < i,$$

and

$$\varphi_i(x_i) \text{ is a scalar multiple of } \mathrm{Eu}_T(i, C_i).$$

Then the classes $\{\varphi_i\}$ generate $H_T^*(X)$ freely as a module over $H_T^*(pt)$.

Proof. Since X is equivariantly formal, we know that $H_T^*(X)$ injects into $H_T^*(X^T)$ and is a free H_T^* -module of rank $m = |X^T|$. First, we show that the φ_i 's are linearly independent. Arguing by contradiction, suppose there is a linear combination

$$\sum_{i=0}^m f_i \varphi_i = 0,$$

with $f_i \in H_T^*$, not all of them zero. Let k be the minimum of the set $\{i \mid f_i \neq 0\}$. Then we have

$$f_k \varphi_k + f_{k+1} \varphi_{k+1} + \dots + f_m \varphi_m = 0$$

where $f_k \neq 0$. Let us restrict this linear combination to x_k . Then

$$f_k \varphi_k(x_k) + f_{k+1} \varphi_{k+1}(x_k) + \dots + f_m \varphi_m(x_k) = 0.$$

But $\varphi_\ell(x_k) = 0$ for all $\ell > k$. Thus we obtain

$$f_k \varphi(x_k) = 0.$$

However, $\varphi(x_k)$ is a non-zero multiple of the Euler class $\text{Eu}_T(x_k, C_k)$ and, as such, it is non-zero. We conclude that f_k must be zero. This is a contradiction.

To conclude the proof, we need to show that the φ_i 's generate $H_T^*(X)$ as a module. But this is a routine exercise, using induction on the length of the filtration of X (the base case being trivial). The commutative diagram of page 18 then disposes of the inductive step. \square

As for the existence of classes satisfying Lemma 6.2, we will show that they can always be constructed on GKM-varieties. First, we need two technical lemmas.

Lemma 6.3. *Let X be a normal projective T -variety with finitely many fixed points. Choose a generic one-parameter subgroup and write X as $X = C \sqcup Y$, where*

$$C = \{z \in X \mid \lim_{t \rightarrow 0} tz = x\}$$

is the stable cell of $x \in X^T$, and Y is closed and T -stable. Then any closed irreducible T -stable curve that passes through x is contained in the Zariski closure of C .

Proof. Let ℓ be a closed irreducible T -stable curve passing through x . Recall that ℓ is the closure of a one-dimensional orbit Tz . Moreover, $\ell = \overline{Tz}$ has two fixed points, namely, x and a fixed point $y_{i(\ell)}$ contained *necessarily* in Y . We claim that $z \in C$. For otherwise, $\lim_{t \rightarrow 0} tz = y_{i(\ell)}$, which implies that z belongs to the stable subvariety of $y_{i(\ell)}$. Since Y is T -invariant and closed, then $\ell = \overline{Tz} \subset Y$. That is, $x \in \partial\ell$ would belong to Y , which is absurd. Thus $z \in C$.

The fact that C is also T -stable gives the inclusion $Tz \subset C$. We conclude that $\ell = \overline{Tz} \subset \overline{C}$. \square

Lemma 6.4. *Let X be a normal projective variety on which a torus acts with a finite number of fixed points and one-dimensional orbits. Suppose X is equivariantly formal and there is a generic one-parameter subgroup such that X can be written as a disjoint union $X = C \sqcup Y$, where*

$$C = \{z \in X \mid \lim_{t \rightarrow 0} tz = x\}$$

is a rational cell with unique attractive fixed point $x \in X^T$, and Y is closed and T -stable. Then the cohomology class $\tau \in \bigoplus_{w \in X^T} H_T^(w)$, defined by*

$$\tau(x) = \text{Eu}_T(x, C) \quad \text{and} \quad \tau(y) = 0 \quad \text{for all } y \in Y^T,$$

belongs to the image of $H_T^(X)$ in $H_T^*(X^T)$.*

Proof. The hypotheses imply that X is a GKM-variety. As a result, the equivariant cohomology of X can be described by the GKM-relations of Theorem 2.5. So, to prove the lemma, it is enough to verify that τ satisfies such relations.

Because τ restricts to zero at every fixed point except x , we need only show that

$$\tau(x) = \tau(x) - \tau(y) = \text{Eu}_T(x, C)$$

is divisible by χ_i whenever the fixed points $x \in C$ and $y_i \in Y^T$ are joined by a T -curve ℓ_i in X , and T acts on ℓ_i through χ_i . Let p be the total number of ℓ_i 's.

By Lemma 6.3, the curve ℓ_i is contained in the Zariski closure \overline{C} of C . In fact, $\ell_i \setminus \{y_i\} \subset C$. Also, it follows from Theorem 3.15 that $p = \dim(C)$. Thus, using Corollary 5.6, we conclude that $\text{Eu}_T(x, C)$ is a non-zero multiple of the χ_i 's. In short, τ belongs to $H_T^*(X)$. \square

It is noticeable that, in the previous lemmas, no assumption on the irreducibility of X has been made. Surely we allow for some flexibility in this matter, since the various filtered pieces X_i of a \mathbb{Q} -filtrable space X need not be irreducible.

Theorem 6.5. *Let X be a \mathbb{Q} -filtrable GKM-variety. Then cohomology generators $\{\varphi_i\}$ of $H_T^*(X)$ with the properties described in Lemma 6.2 exist.*

Proof. We proceed by induction on m , the length of the filtration of X . If $m = 1$, then $X = \{x_1\}$ and the statement is clear, since we can just choose $\varphi_1 = 1$. Assuming we have proved the statement for varieties with a filtration of length m , let us prove the case when the length is $m + 1$. First, notice that $X_{m+1} = X$ and, by the inductive hypothesis, there are classes $\varphi_1, \dots, \varphi_m \in H_T^*(X_m)$ which satisfy the desired properties in $H_T^*(X_m)$. Using the commutative diagram of page 18, we can lift them to classes $\tilde{\varphi}_1, \dots, \tilde{\varphi}_m$ which still satisfy the required conditions, though this time they lie in $H_T^*(X_{m+1}) = H_T^*(X)$. In consequence, we just need to construct a class $\varphi_{m+1} \in H_T^*(X)$ with the sought-after qualities. So set $\varphi_{m+1}(x_{m+1}) = \text{Eu}_T(x_{m+1}, C_{m+1})$ and $\varphi_{m+1}(x_j) = 0$ for all $j \leq m$. By Lemma 6.4, this class surely belongs to $H_T^*(X)$. Thus the result also holds for varieties with a filtration of length $m + 1$. This provides the inductive step in view of Lemma 6.2. \square

Definition 6.6. Let X be a \mathbb{Q} -filtrable T -variety. Fix an ordering of the fixed points, say $x_1 < x_2 < \dots < x_m$. Given $\mu \in H_T^*(X)$, we define its **local index at x_i** , denoted $I_i(\mu)$, by the following formula:

$$I_i(\mu) = \int_{X_i} p_i^*(\mu),$$

where $p_i : X_i \rightarrow X$ denotes the inclusion of the i -th filtered piece into X . It follows from the definition that assigning local indices yields H_T^* -linear morphisms

$$I_i : H_T^*(X) \rightarrow H_T^*(pt).$$

Using the localization formula (Proposition 5.4), one can easily prove the following

Lemma 6.7. *The local index of μ at x_i satisfies*

$$I_i(\mu) = \sum_{j \leq i} \frac{\mu(x_j)}{\text{Eu}_T(x_j, X_i)},$$

where $\mu(x_j)$ denotes the restriction of μ to x_j . □

Corollary 6.8. *Let $x_i \in X^T$, be a fixed point. Suppose that $\mu \in H_T^*(X)$ is a cohomology class that satisfies $\mu(x_j) = 0$ for all $j < i$. Then*

$$\mu(x_i) = I_i(\mu)\text{Eu}_T(x_i, X_i).$$

□

Our most important result in this Section is the following generalization of the work of Guillemin and Kogan ([GK], Theorems 1.1 and 1.6) to \mathbb{Q} -filtrable GKM-varieties.

Theorem 6.9. *Let X be a \mathbb{Q} -filtrable GKM-variety. Let $x_1 < x_2 < \dots < x_m$ be the order relation on X^T compatible with the filtration of X . Then, for each $i = 1, \dots, m$, there exists a unique class $\theta_i \in H_T^*(X)$ with the following properties:*

- (i) $I_i(\theta_i) = 1$,
- (ii) $I_j(\theta_i) = 0$ for all $j \neq i$,
- (iii) the restriction of θ_i to $x_j \in X^T$ is zero for all $j < i$, and
- (iv) $\theta_i(x_i) = \text{Eu}_T(i, C_i)$.

Moreover, the θ_i 's generate $H_T^*(X)$ freely as a module over $H_T^*(pt)$.

Proof. By Theorem 6.5, choose a set of free generators $\{\varphi_i\}$ which satisfy the properties described in Lemma 6.2, together with the additional condition $\varphi_i(x_i) = \text{Eu}_T(i, C_i)$.

Given i , notice that $I_j(\varphi_i) = 0$, for all $j < i$, and $I_i(\varphi_i) = 1$. We will show that we can modify these φ_i 's accordingly to obtain the generators θ_i . In fact, given $i \in \{1, \dots, m\}$, the only obstruction to setting $\theta_i = \varphi_i$ is that $I_j(\varphi_i)$ can be non-zero for some $j > i$.

Let $i \in \{1, \dots, m\}$. If $I_j(\varphi_i) = 0$ for all $j > i$, then let $\theta_i = \varphi_i$. Otherwise, proceed as follows. Let k_0 be the minimum of all $k > i$ such that $I_k(\varphi_i) \neq 0$. Define $\Psi_i = \varphi_i - I_{k_0}(\varphi_i)\varphi_{k_0}$. Let us compute the local indices of Ψ_i . Clearly, if $j < i$, we have $I_j(\Psi_i) = 0$. Also, if $j = i$, then $I_i(\Psi_i) = 1$. It is worth noticing that Ψ_i restricts to 0 at each x_j with $j < i$. Now if j satisfies $i < j \leq k_0$, then $I_j(\Psi_i) = 0$. So, arguing by induction, we can provide a class $\widetilde{\Psi}_i$ such that $I_j(\widetilde{\Psi}_i) = 0$ for all $j \neq i$, and $I_i(\widetilde{\Psi}_i) = 1$. Thus, set $\theta_i = \widetilde{\Psi}_i$. Working on each i at a time, we conclude that there exist classes θ_i satisfying conditions (i)-(iv) of the Theorem.

Let us now prove uniqueness. Suppose there are classes $\{\theta_i\}$ and $\{\theta'_i\}$ satisfying all the properties of the theorem. Fix i and let $\tau = \theta_i - \theta'_i$. It is clear that τ is an element of $H_T^*(X)$ whose local index $I_j(\tau)$ is zero for all j . Suppose that τ is not zero. Then, since $H_T^*(X)$ injects into $H_T^*(X^T)$, there

should be a k such that $\tau(x_k) \neq 0$. Take the minimum of all k 's for which $\tau(x_k) \neq 0$. Denote this minimum by s . Then, by Corollary 6.8, one would have $\tau(x_s) = I_s(\tau)\text{Eu}_T(x_s, X_s) = 0$. This is absurd. Therefore $\tau = 0$. Since i can be chosen arbitrarily, we conclude that $\theta_i = \theta'_i$ for all i .

Finally, notice that properties (iii) and (iv) together with Lemma 6.2 imply that the θ_i 's freely generate $H_T^*(X)$. We are done. \square

7. RATIONAL CELLS AND STANDARD GROUP EMBEDDINGS

Previously, we have developed the theory of \mathbb{Q} -filtrable varieties. In this last section, we provide the theory with a large class of examples, namely, rationally smooth standard embeddings. We show that these varieties admit BB -decompositions into rational cells (Theorem 7.3). Thus, they are \mathbb{Q} -filtrable and satisfy Theorems 4.8 and 6.9.

First, let us set the stage. An affine algebraic monoid M is called **reductive** if it is irreducible, normal, and its unit group is a reductive algebraic group. See [R4] for many of the details. A reductive monoid is called **semisimple** if it has a zero element, and its unit group has a one-dimensional center.

Let M be a reductive monoid with zero. Denote by G its unit group and by T a maximal torus of G . Associated to M , there is a torus embedding $\overline{T} \subset M$ defined as follows

$$\overline{T} = \{x \in M \mid xt = tx, \text{ for all } t \in T\}.$$

Certainly, $T \subseteq \overline{T}$. Let $E(\overline{T})$ be the idempotent set of \overline{T} ; that is,

$$E(\overline{T}) = \{e \in \overline{T} \mid e^2 = e\}.$$

The **Renner monoid**, \mathcal{R} , is defined to be $\mathcal{R} := \overline{N_G(\overline{T})}/T$. It is a finite monoid whose group of units is W (the Weyl group) and contains $E(\overline{T})$ as idempotent set. In fact, any $x \in \mathcal{R}$ can be written as $x = fu$, where $f \in E(\overline{T})$ and $u \in W$. Recall that W is generated by reflections $\{s_\alpha\}_{\alpha \in \Phi}$, where Φ is the set of roots of G with respect to T .

Denote by \mathcal{R}_k the set of elements of rank k in \mathcal{R} , that is,

$$\mathcal{R}_k = \{x \in \mathcal{R} \mid \dim Tx = k\}.$$

Definition 7.1. Let M be a reductive monoid with unit group G and zero element $0 \in M$. There exists a central one-parameter subgroup $\epsilon : \mathbb{C}^* \rightarrow G$ with image Z contained in the center of G , that converges to 0 ([Br6], Lemma 1.1.1). Then \mathbb{C}^* acts attractively on M via ϵ , and hence the quotient

$$\mathbb{P}_\epsilon(M) = [M \setminus \{0\}]/\mathbb{C}^*$$

is a normal projective variety. Notice also that $G \times G$ acts on $\mathbb{P}_\epsilon(M)$ via

$$G \times G \times \mathbb{P}_\epsilon(M) \rightarrow \mathbb{P}_\epsilon(M), (g, h, [x]) \mapsto [gxh^{-1}].$$

Furthermore, $\mathbb{P}_\epsilon(M)$ is a normal projective embedding of the reductive group G/Z . In the sequel, $X = \mathbb{P}_\epsilon(M)$ will be called a **Standard Group Embedding**.

For an up-to-date description of these and other embeddings, see [AB].

Example 7.2. Let G_0 be a semisimple algebraic group over the complex numbers and let $\rho : G_0 \rightarrow \text{End}(V)$ be a representation of G_0 . Define Y_ρ to be the Zariski closure of $G = [\rho(G_0)]$ in $\mathbb{P}(\text{End}(V))$, the projective space associated with $\text{End}(V)$. Finally, let X_ρ be the normalization of Y_ρ . By definition, X_ρ is a standard group embedding of G . Notice that M_ρ , the Zariski closure of $\mathbb{C}^*\rho(G_0)$ in $\text{End}(V)$, is a semisimple monoid whose group of units is $\mathbb{C}^*\rho(G_0)$. Rationally smooth standard embeddings of the form X_ρ , with ρ irreducible, have been classified combinatorially in [R2].

We now come to the main result of this section. It states that rationally smooth standard embeddings are equivariantly formal for the induced $T \times T$ -action.

Theorem 7.3. *Let $X = \mathbb{P}_\epsilon(M)$ be a standard group embedding. If X is rationally smooth, then X is \mathbb{Q} -filtrable.*

Proof. Renner has shown that X comes equipped with the following BB-decomposition:

$$X = \bigsqcup_{r \in \mathcal{R}_1} C_r,$$

where $\mathcal{R}_1 = X^{T \times T}$. See Theorem 3.4 of [R1], and Theorem 4.3 of [R3] for more details. Our strategy is to show that if X is rationally smooth, then each cell C_r is rationally smooth.

With this purpose in mind, we call the reader's attention to the fact that, in the terminology of [R1], M is quasismooth (Definition 2.2 of [R1]) if and only if $M \setminus \{0\}$ is rationally smooth. The equivalence between these two notions follows from Theorem 2.1 of [R1] and Theorems 2.1, 2.3, 2.4 and 2.5 of [R2].

Next, by Lemma 4.6 and Theorem 4.7 of [R1], each C_r equals

$$U_1 \times C_r^* \times U_2,$$

where the U_i 's are affine spaces. Moreover, if we write $r \in \mathcal{R}_1$ as $r = ew$, with $e \in E_1(\overline{T})$ and $w \in W$, then $C_r^* = C_e^*w$. So it is enough to show that C_e^* is rationally smooth, for $e \in E_1(\overline{T})$.

By Theorem 5.1 of [R1], it follows that, if $X = \mathbb{P}_\epsilon(M)$ is rationally smooth, then

$$C_e^* = [f_e M(e)]/Z,$$

for some unique $f_e \in E(\overline{T})$, where $M(e) = M_e Z$ and M_e is rationally smooth (Theorem 2.5 of [R2]). Furthermore, the proof of Theorem 5.1 of [R1] also

implies that $[e]$ is the zero element of the rationally smooth, reductive, affine monoid $M(e)/Z$. Additionally,

$$C_e^* = \{x \in M(e)/Z \mid \lim_{s \rightarrow 0} sx = [e]\},$$

for some generic one-parameter subgroup. Using Lemma 7.4 below, one concludes that C_e^* is rationally smooth.

Finally, since X is normal, projective and admits a BB -decomposition into rational cells, we have compiled all the necessary data to conclude that X is \mathbb{Q} -filtrable. \square

Lemma 7.4. *Let M be a reductive monoid with zero. Suppose that zero 0 is a rationally smooth point of M . Let $f \in E(M)$, be an idempotent of M . Then $0 \in fM$ is a rationally smooth point of the closed subvariety fM .*

Proof. By Lemma 1.1.1 of [Br6], one can find a one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$, with image S , such that $\lambda(0) = f$. Notice that

$$fM = \{x \in M \mid \lambda(t)x = x, \forall t \in \mathbb{C}^*\}.$$

That is, fM is the fixed point set of the subtorus S of T . Thus, by Theorem 1.1 of [Br4], one concludes that 0 is also a rationally smooth point of fM . \square

We conclude this section providing a partial converse to Theorem 7.3.

Theorem 7.5. *Let $X = \mathbb{P}_e(M)$ be a standard embedding. Suppose that X contains a unique closed $G \times G$ -orbit. If X is \mathbb{Q} -filtrable, then X is rationally smooth.*

Proof. Since X contains a unique closed $G \times G$ -orbit, it follows from [R4], Chapter 7, that $W \times W$ acts transitively on \mathcal{R}_1 , the set of representatives of the $T \times T$ -fixed points of X . Because X is irreducible, there exists a unique cell, say C_σ , with $\sigma \in \mathcal{R}_1$, such that $X = \overline{C_\sigma}$. By assumption, C_σ is rationally smooth at σ , and so, X is rationally smooth at σ . We claim that X is rationally smooth at every $r \in \mathcal{R}_1 = X^{T \times T}$. Indeed, by the previous remarks, $r = w \cdot \sigma \cdot v$, for some $(w, v) \in W \times W$ and rational smoothness is a local property invariant under homeomorphisms. Now Lemma 7.6 below concludes the proof. \square

Lemma 7.6. *Let X be a projective T -variety with a finite number of fixed points x_1, \dots, x_m . Then X is rationally smooth at every $x \in X$ if and only if X is rationally smooth at every fixed point x_i .*

Proof. One direction is clear. For the converse, pick a generic one-parameter subgroup $\lambda : \mathbb{C}^* \rightarrow T$ such that $X^T = X^{\mathbb{C}^*}$. Let $x \in X$. Then, there exists $x_k \in X^T$ such that $x_k = \lim_{t \rightarrow 0} tx$ (BB -decomposition). Moreover, since X is rationally smooth at x_k , there exists a neighborhood V_k of x_k with the property that X is rationally smooth at every $y \in V_k$. By construction, there exists $s \in \mathbb{C}^*$ satisfying $sx \in V_k$. To see this, simply notice that we can find a sequence $\{s_n\} \subset \mathbb{C}^*$ for which $s_n \cdot x$ converges to x_k , i.e. there is N such that $s_n \cdot x$ belongs to V_k , for all $n \geq N$. Now setting $s = s_N$ yields

$s \cdot x \in V_k$. In other words, sx is a rationally smooth point of X . But the set of rationally smooth points is T -invariant. Hence, x is a rationally smooth point of X . Inasmuch as the point x was chosen arbitrarily, the argument is complete. \square

In the author's thesis it was shown that all standard embeddings are $T \times T$ -skeletal. Consequently, rationally smooth standard embeddings are also GKM-varieties. In a forthcoming paper ([G-2]) we find explicitly all the GKM-data (i.e. fixed points, invariant curves and associated characters) of any rationally smooth standard embedding $\mathbb{P}_\epsilon(M)$, and describe $H_{T \times T}^*(\mathbb{P}_\epsilon(M))$ as a complete combinatorial invariant of M . The results will appear elsewhere.

REFERENCES

- [AB] Alexeev, V.; Brion, M. *Stable reductive varieties II*. Projective case, Adv. Math. 184 (2004), 380-408.
- [Ar] Arabia, A. *Classes d'Euler équivariantes et points rationnellement lisses*. Annales de l'institut Fourier, tome 48, No. 3 (1998), p. 861-912.
- [AS] Atiyah, M.F.; Segal, G. B. *The index of elliptic operators: II*. The Annals of Mathematics, 2nd Ser., Vol. 87, No. 3. May, 1968. pp. 531-545.
- [At] Atiyah, M. F. *Elliptic Operators and Compact Groups*. Lecture Notes in Mathematics, 401. Springer-Verlag. 1974.
- [BB1] Bialynicki-Birula, A. *Some theorems on actions of algebraic groups*. The Annals of Mathematics, 2nd Ser., Vol 98, No. 3, Nov. 1973, pp. 480-497.
- [BB2] Bialynicki-Birula, A. *Some properties of the decompositions of algebraic varieties determined by actions of a torus*. Bull. Acad. Polon. Sci. Sr. Sci. Math. Astronom. Phys. 24 (1976), no. 9, 667674.
- [BDP] Bifet, E; De Concini, C.; Procesi, C. *Cohomology of Regular Embeddings*. Advances in Mathematics, 82, pp. 1-34 (1990).
- [Bo] Borel, A. *Seminar on transformation groups*. Annals of Math Studies, No. 46, Princeton University Press, Princeton, N.J. 1960.
- [Br1] Brion, M. *Equivariant Chow groups for torus actions*. Transformation groups, vol. 2, No. 3, 1997, pp. 225-267.
- [Br2] Brion, M. *Equivariant cohomology and equivariant intersection theory*. Notes by Alvaro Rittatore. NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 514, Representation theories and algebraic geometry (Montreal, PQ, 1997), 1-37, Kluwer Acad. Publ., Dordrecht, 1998.
- [Br3] Brion, M. *The behaviour at infinity of the Bruhat decomposition*. Comment. Math. Helv. 73, 1998, pp. 137-174.
- [Br4] Brion, M. *Rational smoothness and fixed points of torus actions*. Transformation Groups, Vol. 4, No. 2-3, 1999, pp. 127-156.
- [Br5] Brion, M. *Poincaré duality and equivariant cohomology*. Michigan Math. J. 48, 2000, pp. 77-92.
- [Br6] Brion, M. *Local structure of algebraic monoids*. Mosc. Math. J. 8 (2008), no. 4, 647666, 846.
- [BV] Brion, M., Vergne, M. *An equivariant Riemann-Roch theorem for complete, simplicial toric varieties*. J. Reine Angew. Math. 482 (1997), 6792.
- [C] Carrell, J. *The Bruhat graph of a Coxeter group, a conjecture of Deodhar, and rational smoothness of Schubert varieties*. Algebraic groups and their generalizations: classical methods (University Park, PA, 1991), 5361, Proc. Sympos. Pure Math., 56, Part 1, Amer. Math. Soc., Providence, RI, 1994.

- [CS] T. Chang, T. Skjelbred, *The topological Shur lemma and related results*, Annals of Mathematics, 2nd Ser., Vol. 100, No. 2, (Sept., 1974), pp. 307-321.
- [D] Danilov, V. *The geometry of toric varieties*. Uspekhi Mat. Nauk 33 (1978), no. 2(200), 85134, 247.
- [G-1] Gonzales, R. *GKM theory of rationally smooth group embeddings*. PhD. thesis, The University of Western Ontario, 2011.
- [G-2] Gonzales, R. *GKM theory of rationally smooth standard embeddings*. Preprint.
- [GK] Guillemin, V., Kogan, M. *Morse theory on Hamiltonian G -spaces and equivariant K -theory*. J. Differential Geometry, 66, 2004, pp. 345-375.
- [GKM] Goresky, M., Kottwitz, R., MacPherson, R. *Equivariant cohomology, Koszul duality, and the localization theorem*. Invent. math. 131, 25-83 (1998)
- [GZ] Guillemin, V., Zara, C. *1-skeleta, Betti numbers, and equivariant cohomology*. Duke Math. J. 107 (2001), 283-349.
- [Har] Hartshorne, R. *Algebraic Geometry*, GTM, 52, Springer-Verlag, 1978.
- [HHH] Harada, M.; Henriques, A.; Holm, T. S. *Computation of generalized equivariant cohomologies of Kac-Moody flag varieties*. Adv. Math. 197 (2005), no. 1, 198-221.
- [Hs] Hsiang, Wu Yi. *Cohomology Theory of Topological Transformation Groups*. Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 85. Springer-Verlag, New York-Heidelberg, 1975.
- [M] McCrory, C. *A Characterization of Homology Manifolds*. J. London Math. Soc. 16 (1977), 146-159.
- [PS] Peters, C. and Steenbrink, J. *Mixed Hodge Structures*. Springer Verlag, 2008.
- [Q] D. Quillen. *The Spectrum of an Equivariant Cohomology Ring: I, II*. The Annals of Mathematics, 2nd. Ser., Vol. 94, No. 3. Nov., 1971, pp. 549-572, pp.573-602.
- [R1] Renner, L. *The H -polynomial of a semisimple monoid*. J. Alg. 319 (2008), 360-376.
- [R2] Renner, L. *Rationally smooth algebraic monoids*. Semigroup Forum 78 (2009), 384-395.
- [R3] Renner, L. *The H -polynomial of an Irreducible Representation*. Journal of Algebra 332 (2011) 159186.
- [R4] Renner, L. *Linear Algebraic Monoids*. Encyclopedia of Mathematical Sciences, vol. 134. Invariant Theory and Algebraic Transformation Groups, V. Springer-Verlag, Berlin, 2005.
- [Seg] Segal, G. *Equivariant K -theory*. Publ. math. l'I.H.E.S., tome 34 (1968), p. 129-151.
- [Sha] Shafarevich, I. *Basic Algebraic Geometry 1*. 2nd rev. and expanded ed. Springer-Verlag, 1994.
- [Su] Sumihiro, H. *Equivariant completion*, J. Math. Kyoto University 14 (1974), 1-28.
- [U] Uma, V. *Equivariant K -theory of compactifications of algebraic groups*. Transform. Groups 12 (2007), No. 2, 371-406.
- [VV] Vezzosi, G., Vistoli, A. *Higher algebraic K -theory for actions of diagonalizable algebraic groups*. Invent. Math. 153 (2003), No. 1, 1-44.
- [W] Weber, A. *Formality of equivariant intersection cohomology of algebraic varieties*. Proc. Amer. Math. Soc. 131 (2003), no. 9, 2633-2638.

DEPARTMENT OF MATHEMATICS, GALATASARAY UNIVERSITY, CIRAGAN CAD. NO.36, BESIKTAS, 34357 ISTANBUL, TURKEY

E-mail address: rgonzalesv@gmail.com