

Solitary and periodic solutions of the generalized Kuramoto - Sivashinsky equation

Nikolai A. Kudryashov

Department of Applied Mathematics
 Moscow Engineering and Physics Institute
 (State university)
 31 Kashirskoe Shosse, 115409
 Moscow, Russian Federation

Abstract

The generalized Kuramoto - Sivashinsky in the case of the power nonlinearity with arbitrary degree is considered. New exact solutions of this equation are presented.

Keywords: Exact solution, nonlinear differential equation, Kuramoto - Sivashinsky equation

PACS: 02.30.Hq - Ordinary differential equations

1 Introduction

In this paper we present exact solutions of the generalized Kuramoto - Sivashinsky equation

$$u_t + u^m u_x + \alpha u_{xx} + \beta u_{xxx} + \gamma u_{xxxx} = 0 \quad (1.1)$$

Nonlinear evolution equation (1.1) at $m = 1$ has been studied by a number of authors from various viewpoints. This equation has drawn much attention not only because it is interesting as a simple one - dimensional nonlinear evolution equation including effects of instability, dissipation and dispersion but also it is important for description in engineering and scientific problems. Equation (1.1) in work [1] was used for explanation of the origin of persistent wave propagation through medium of reaction - diffusion type. In paper [2] equation (1.1) was obtained at $m = 1$ for description of the nonlinear evolution of the disturbed flame front. We can meet the application of equation (1.1) for studying of motion of a viscous incompressible flowing down an inclined plane [3-5]. Mathematical model for consideration dissipative waves in plasma physics by means of equation (1.1) was presented in [6]. Elementary particles as solutions of the Kuramoto - Sivashinsky equation were studied in [7]. Equation (1.1) at $m \neq 1$ also can be used for the description of physical applications. For a example this equation were derived for description of nonlinear long waves in viscous - elastic tube [8].

Exact solution of equation (2.2) in the form of the solitary wave at $m = 1$ and at $\beta = 0$ were obtained in [1, 9]. Solitary wave solutions of equation (2.2) at $m = 1$ in the case $\beta \neq 0$ were found in works [10–12] for cases:

$$\alpha = \frac{\beta^2}{16\gamma}, \quad \alpha = \frac{47\beta^2}{144\gamma}, \quad \alpha = \frac{73\beta^2}{256\gamma}. \quad (1.2)$$

Other forms of these solutions were presented in works [13–17]. Periodical solutions of equation (2.2) were found in [18, 19] at $\alpha = \frac{\beta^2}{16\gamma}$.

Exact solutions of equation (2.2) at $m = 2$ recently were obtained [8] for the four cases:

$$\alpha = \frac{2\beta^2}{25\gamma}, \quad \alpha = \frac{71\beta^2}{225\gamma}, \quad \alpha = \frac{121\beta^2}{55\gamma}, \quad \alpha = \frac{374\beta^2}{2025\gamma}. \quad (1.3)$$

The aim of this paper is to search for exact solutions of equation (1.1) in general case of arbitrary degree m . We present exact solutions of equation (1.1) at $m \neq -1$, $m \neq -3$ and $m \neq 0$. Some of these solutions are new.

2 Solitary waves of equation (1.1)

Using traveling wave

$$u(x, t) = w(z), \quad z = x - C_0 t \quad (2.1)$$

from equation (1.1) we have

$$w_{zzz} + \sigma w_{zz} + w_z - C_0 w + \frac{1}{m+1} w^{m+1} = 0, \quad m \neq -1 \quad (2.2)$$

Assuming in (2.2)

$$w(z) = v(z)^{\frac{1}{m}} \quad (2.3)$$

we have equation in the form

$$\begin{aligned} & \gamma m^3 v^2 v_{zzz} + \gamma m^2 v^2 v_{zzz} + 3\gamma m v v_z v_{zz} + \beta m^3 v^2 v_{zz} - 2\gamma m v_z^3 - \\ & - 3\gamma m^3 v v_z v_{zz} + \gamma v_z^3 - \gamma m^2 v_z^3 + 2\gamma m^3 v_z^3 + \beta m v v_z^2 - \\ & - \beta m^3 v v_z^2 + \beta m^2 v^2 v_{zz} + \alpha m^3 v^2 v_z + \alpha m^2 v^2 v_z + m^3 v^4 - \\ & - C_0 m^4 v^3 - C_0 m^3 v^3 = 0 \end{aligned} \quad (2.4)$$

Study of analytical properties of equation (2.4) allows us to determine that in the general case the meromorphic solutions of equation (2.4) can be found taking into account two cases:

$$\alpha = \frac{\beta^2 m}{\gamma (m+3)^2}, \quad \alpha = \frac{\beta^2 (2m^2 + 18m + 27)}{9\gamma (m+3)^2} \quad (2.5)$$

Consider the first case. The pole order of the solution of equation (2.4) is equal to three therefore we look for exact solutions of equation (2.4) in the form

$$v(z) = A_0 + A_1 Y(z) + A_2 Y(z)^2 + A_3 Y(z)^3, \quad (2.6)$$

where $Y(z)$ is solution of equation with the first order pole. Let us take simplest equation in the form [20, 21]

$$Y_z = -Y^2 + b. \quad (2.7)$$

Substituting (2.6) into (2.4) and taking into account equation (2.7) we have

$$b = \frac{\beta^2 m^2}{4 \gamma^2 (m+3)^2}. \quad (2.8)$$

We have also the following values of coefficients in (2.6)

$$A_3 = \frac{3 \gamma (2m+3)(m+3)(m+1)}{m^3},$$

$$A_2 = -\frac{3(2m+3)(m+1)\beta}{2m^2}, \quad A_1 = -\frac{3(2m+3)(m+1)\beta^2}{4(m+3)m\gamma}, \quad (2.9)$$

$$C_0 = \frac{2(m+2)\beta^3}{\gamma^2(m+3)^3}, \quad A_0 = \frac{3(2m+3)(m+1)\beta^3}{8\gamma^2(m+3)^2}.$$

We have solutions of equation (2.7) in the form

$$Y(z) = \frac{\beta m}{2\gamma(m+3)} \tanh \left\{ \frac{\beta m (z - z_0)}{2\gamma(m+3)} \right\}. \quad (2.10)$$

Solution of equation (2.2) takes the form

$$w(z) = \left(\frac{3(2m+3)(m+1)\beta^3}{8\gamma(m+3)} \right)^{\frac{1}{m}} \left(1 - \tanh \left\{ \frac{\beta m (z - z_0)}{2\gamma(m+3)} \right\} \right)^{\frac{2}{m}} \left(1 + \tanh \left\{ \frac{\beta m (z - z_0)}{2\gamma(m+3)} \right\} \right)^{\frac{1}{m}}, \quad m \neq 0, \quad m \neq -3, \quad m \neq -1. \quad (2.11)$$

This is new solution of equation (1.1). Solution (2.11) at $m = 1$ takes the form of the solitary wave that was obtained in [10]. In the case $m = 2$ this solution corresponds to exact solution of equation (1.1) published in [8].

Consider the second case. Solutions of equation (2.4) we look for taking into consideration the transformation (2.6) again. Function $Y(z)$ satisfies equation (2.8) as well. In this case we obtain the expression for the parameter b in the form

$$b = \frac{\beta^2 m^2}{36 \gamma^2 (m+3)^2}. \quad (2.12)$$

We also obtain coefficients A_3, A_2, A_1, C_0 and A_0 as following

$$A_3 = \frac{3\gamma(2m+3)(m+3)(m+1)}{m^3},$$

$$A_2 = -\frac{3(2m+3)(m+1)\beta}{2m^2}, \quad A_1 = \frac{(2m+3)(m+1)\beta^2}{4\gamma m(m+3)}, \quad (2.13)$$

$$C_0 = -\frac{(2m+3)\beta^3}{9(m+3)^2\gamma^2}, \quad A_0 = -\frac{\beta^3(2m+3)(m+1)}{72(m+3)^2\gamma^2}.$$

Using these coefficients we have solution of equation (2.2). It takes the form

$$w(z) = \left(\frac{\beta^3(2m+3)(m+1)}{72\gamma^2(m+3)^2} \right)^{\frac{1}{m}} \left(\tanh \left\{ \frac{\beta m(z-z_0)}{6\gamma(m+3)} \right\} - 1 \right)^{\frac{3}{m}}, \quad (2.14)$$

$$m \neq 0, \quad m \neq -1, \quad m \neq -3.$$

Other forms of this solution were found in works [22–24]. At $m = 1$ from (2.14) we have one of the solitary solutions of work [10]. In the case $m = 2$ we get solution [8].

3 Periodical waves of equation (1.1)

Let us find periodical solutions of equation (2.2) taking into consideration the formula [18]

$$w(z) = A_0 + A_1 R + A_2 R_z, \quad R \equiv R(z), \quad (3.1)$$

taking into account the relation for the parameter α

$$\alpha = \frac{\beta^2 m}{\gamma(m+3)^2}. \quad (3.2)$$

We assume that $R(z)$ is solution of equation for the Weierstrass function

$$R_{zz} + 6R^2 - aR - b = 0 \quad (3.3)$$

From equation (3.3) we have equation

$$R_z^2 + 4R^3 - aR^2 - 2bR + d = 0 \quad (3.4)$$

Substituting (3.1) into equation (2.3) and taking into account equations (3.3)

and (3.4) we have

$$b = \frac{m^4 \beta^4 - \gamma^4 a^2 (m+3)^2}{24 \gamma^4 (m+3)^4}, \quad A_2 = \frac{3 \gamma (2m+3)(m+3)(m+1)}{2m^3},$$

$$A_1 = \frac{3 \beta (2m+3)(m+1)}{2m^2}, \quad C_0 = \frac{2 \beta^3 (m+2)}{\gamma^2 (m+3)^3}, \quad (3.5)$$

$$A_0 = \frac{\beta (2m+3)(m+1) (\beta^2 m^2 - \gamma^2 (m+3)^2 a)}{8 \gamma^2 m^2 (m+3)^2},$$

$$d = \frac{(2 \beta^2 m^2 + \gamma^2 (m+3)^2 a) (\beta^2 m^2 - \gamma^2 (m+3)^2 a)^2}{432 \gamma^6 (m+3)^6} \quad (3.6)$$

Solution of equation (2.2) in this case takes the form

$$w(z) = \left(\frac{3(m+1)(2m+3)}{2m^2} \right)^{\frac{1}{m}}$$

$$\left(\beta R(z) + \frac{\gamma(m+3)}{m} R_z + \frac{\beta (\beta^2 m^2 - \gamma^2 (m+3)^2 a)}{12 \gamma^2 (m+3)^2} \right)^{\frac{1}{m}}, \quad (3.7)$$

where $R(z)$ satisfies equation

$$R_z^2 + 4R^3 - aR^2 - \frac{(m^4 \beta^4 - \gamma^4 a^2 (m+3)^4)}{12 \gamma^4 (m+3)^4} R -$$

$$- \frac{(2 \beta^2 m^2 + \gamma^2 (m+3)^2 a) (\beta^2 m^2 - \gamma^2 (m+3)^2 a)^2}{432 \gamma^6 (m+3)^6} = 0 \quad (3.8)$$

In general case solutions of equation (3.8) expressed via hyperbolic function because the algebraic equation

$$4R^3 - aR^2 - \frac{(m^4 \beta^4 - \gamma^4 a^2 (m+3)^4)}{12 \gamma^4 (m+3)^4} R -$$

$$\frac{(2 \beta^2 m^2 + \gamma^2 (m+3)^2 a) (\beta^2 m^2 - \gamma^2 (m+3)^2 a)^2}{432 \gamma^6 (m+3)^6} = 0 \quad (3.9)$$

has two equal roots.

However there are periodic solutions of equation (1.1) at $m = 1$ and $m = 3$. Parameter C_0 in these cases will be arbitrary constant. Other coefficients can be found from expressions (3.5). Let us present these solutions using formula (3.1).

In the case $m = 1$ we have $\alpha = \frac{\beta^2}{16\gamma}$. Solution of equation (1.1) takes the form [18]

$$y(z) = C_0 - \frac{5}{4}\beta a - \frac{1}{64}\frac{\beta^3}{\gamma^2} + 15\beta R + 60\gamma R_z \quad (3.10)$$

where $R(z)$ satisfies equation for the Weierstrass function

$$R_z^2 + 4R^3 - aR^2 - 2\left(\frac{\beta^4}{6144\gamma^4} - \frac{a^2}{24}\right)R + \frac{13\beta^6}{4423680\gamma^6} + \frac{a\beta^4}{36864\gamma^4} - \frac{C_0^2}{2160\gamma^2} - \frac{a^3}{432} = 0 \quad (3.11)$$

In the case $m = 3$ we have $\alpha = \frac{\beta^2}{12\gamma}$. Periodic solution of equation (1.1) in this case can be written in the form

$$y(z) = \left(3C_0 - \frac{1}{2}\beta a - \frac{1}{72}\frac{\beta^3}{\gamma^2} + 6\beta R + 12\gamma R_z\right)^{\frac{1}{3}} \quad (3.12)$$

where $R(z)$ satisfies equation for the Weierstrass function in the form

$$R_z^2 + 4R^3 - aR^2 - 2\left(\frac{C_0\beta}{8\gamma^2} - \frac{11\beta^4}{3456\gamma^4} - \frac{a^2}{24}\right)R - \frac{11\beta^4 a}{20736\gamma^4} + \frac{13\beta^6}{373248\gamma^6} + \frac{C_0\beta^3}{1728\gamma^4} + \frac{C_0\beta a}{48\gamma^2} - \frac{C_0^2}{16\gamma^2} - \frac{a^3}{432} = 0 \quad (3.13)$$

This solution of equation (1.1) is a new. Let us note that periodical solutions of equation (1.1) are appeared at $m = 1$ and at $m = 3$ when there is meromorphic solutions in the form of solitary waves of equation (1.1).

References

- [1] *Y. Kuramoto, T. Tsuzuki*, Persistent propagation of concentration waves in dissipative media far from thermal equilibrium, Progress of theoretical physics, vol. 55, No. 2, 1976, 356 - 369
- [2] *G.I. Sivashinsky*, Instabilities, pattern, formation, and turbulence in flames, Ann. Rev. Fluid Mech., vol. 15, 1983, 179 - 199
- [3] *D.J. Benney*, Long waves on liquid films, Journal of Mathematical Physics, 45 (1966) 150
- [4] *J. Topper, T. Kawahara*, Approximate equation for long nonlinear waves on a viscous fluid, Journal of the Physical society of Japan, vol. 44, No. 2, 1978, 663 - 666
- [5] *V.Y. Shkadov*, Solitary waves in layer of viscous fluid, Fluid Dynamics, No. 1, 1977, 63 - 66

- [6] *B.I. Cohen, J.A. Krommers, W.M. Tang, M.N. Rosenbluth*, Non-linear saturation of the dissipative trapped - ion mode by mode coupling, Nuclear fusion, vol.16(6), 1976, 971 - 992
- [7] *D. Michelson*, Elementary particles as solutions of the Sivashinsky equation, Physica D, 44, (1990) 502 - 556
- [8] *N.A. Kudryashov, D.I. Sinelschikov, I.L. Chernyavsky*, Nonlinear evolution equations for description of perturbation in tube, Nonlinear dynamics, vol. 4, No. 1, 2008, 69 - 86
- [9] *R. Conte, M. Musette*, Painlevé analysis and Backlund transformations in the Kuramoto - Sivashinsky equation, Journal of Physics A.: Math. Gen. 22, (1989) 169 - 177
- [10] *N.A. Kudryashov*, Exact soliton solutions of the generalized evolution equation of wave dynamics, Journal of Applied Mathematics and Mechanics, 52 (3) (1988) 360 - 365
- [11] *N.A. Kudryashov*, On types of nonlinear nonintegrable equations with exact solutions Phys Lett. A., **155**, (1991), 269 – 275
- [12] *N.A. Kudryashov, E.D. Zargaryan*, Solitary waves in active-dissipative dispersive media, J. Phys. A. Math. and Gen. **29**, 1996, 8067-8077
- [13] *N.G. Berloff, L.N. Howard*, Solitary and Periodic Solutions of nonlinear nonintegrable equations, Studies in Applied Mathematics, 99 (1997), 1 - 24
- [14] *S. Fu, S. Liu, S. Liu*, New exact solutions to the KdV - Burgers - Kuramoto equation, Chaos, Solitons and Fractals, 23 (2005) 609 - 616
- [15] *S. Zhang*, New exact solutions of the KdV - Burgers - Kuramoto equation, Phys Lett. A., **358**, (2006), 414-420
- [16] *S.A. Khuri*, Traveling wave solutions for nonlinear differential equations: a unified ansatz approach, Chaos, Solitons and Fractals, 32 (2007) 252 - 258
- [17] *J. Nickel*, Traveling wave solutions to the Kuramoto - Sivashinsky equation, Chaos, Solitons and Fractals, 33 (2007) 1376 - 1382
- [18] *N.A. Kudryashov*, Exact solutions of the generalized Kuramoto - Sivashinsky equation, Phys Lett. A., **147**, 1990, 287-291
- [19] *N.A. Kudryashov*, Exact soliton solutions of nonlinear wave equations arising in mechanics, Journal of Applied Mathematics and Mechanics, 54 (3) (1990) 450 - 453
- [20] *N.A. Kudryashov*, Exact solitary waves of the Fisher equation, Physics Letters A., v. 342, 2005, pp.99 - 106,
- [21] *N.A. Kudryashov*, Simplest equation method to look for exact solutions of nonlinear differential equations, Chaos, Solitons and Fractals, v.24, 2005, pp. 1217 - 1231

- [22] *Z.N. Zhu*, Exact Solutions to the n -dimensional generalized fifth order Kuramoto - Sivashinsky type equation, Chinese Journal of Physics, 84 (2) (1996), 85 -90
- [23] *N.A. Kudryashov, M.V. Demina*, Polygons of differential equation for finding exact solutions, Chaos, Solitons and Fractals, v.33, (2007), 1480 - 1496
- [24] *M. Qin, G. Fan*, An effective method for finding special solutions of nonlinear differential equations with variable coefficients, Physics Letters a, 372 (2008) 3240 - 3242