Quantum uncertainty relation saturated by the eigenstates of the harmonic oscillator

A. Mandilara and N. J. Cerf

Quantum Information and Communication, École Polytechnique de Bruxelles, CP 165/59, Université Libre de Bruxelles, 1050 Brussels, Belgium

⁻ 105/59, Universite Liore de Druzeites, 1050 Drusseis, Deigraf

We re-derive the Schrödinger-Robertson uncertainty principle for the position and momentum of a quantum particle in one dimension. Our derivation does not directly employ commutation relations, but works by reduction to an eigenvalue problem, which can then be further exploited to find a larger class of constrained uncertainty relations. We derive an uncertainty relation under the constraint of a fixed degree of Gaussianity, and prove that, remarkably, it is saturated by all eigenstates of the harmonic oscillator. This generalizes the well-known property that the (Gaussian) ground state of the harmonic oscillator saturates the uncertainty relation.

The Heisenberg uncertainty relation [1] captures the difference between classical and quantum states, and sets a limit on the precision of incompatible quantum measurements. It has been introduced in the early days of quantum mechanics, but its form has evolved with the understanding and formulation of quantum physics throughout the years. The first rigorous mathematical proof of Heisenberg's uncertainty relation for the canonical operators of position \hat{x} and momentum $\hat{p}([\hat{x}, \hat{p}] = i\hbar)$

$$(\langle \hat{x}^2 \rangle - \langle \hat{x} \rangle^2)(\langle \hat{p}^2 \rangle - \langle \hat{p} \rangle^2) \ge \hbar^2/4, \tag{1}$$

is due to Kennard [2] and Weyl [3], but only pure states were considered there. The full proof was later derived following different methodologies [4–6] (see [7] for more details), while the properties of the states saturating this inequality were also progressively unveiled.

The original uncertainty relation (1) only concerned the operators \hat{x} and \hat{p} , but it was generalized to any pair of Hermitian operators by Schrödinger [8] and Robertson [9], in the case of pure states. In the same works, the anticommutator of \hat{x} and \hat{p} was also included in Eq. (1), yielding a stronger uncertainty relation

$$(\langle \hat{x}^2 \rangle - \langle \hat{x} \rangle^2) (\langle \hat{p}^2 \rangle - \langle \hat{p} \rangle^2) - \frac{1}{4} (\langle \hat{x}\hat{p} + \hat{p}\hat{x} \rangle - 2 \langle \hat{x} \rangle \langle \hat{p} \rangle)^2 \ge \hbar^2/4,$$
 (2)

that bears their name. The first proof of the Schrödinger-Robertson (SR) uncertainty relation for position and momentum, Eq. (2), in the general case including mixed states is probably due to Moyal [5], and, for any pair of not-necessarily Hermitian operators, to Dodonov, Kurmyshev and Man'ko [10]. In this latter work, the states of minimum uncertainty or *minimizing states* (MS) for the SR inequality were identified as the Gaussian pure states. These states are accessible in practice, and widely used in the field of quantum optics, where they are known as coherent and squeezed states.

In this Letter, we re-derive inequality (2) and the corresponding minimizing states using the Lagrange multipliers method and tools from quantum optics. The exhibited method can be applied to define a much larger class of *bounded* uncertainty relations [11], which provide a stronger bound than Eq. (2) for states on which some

information is known, such as their purity [7] or von Neumann entropy [12]. In particular, we derive a stronger bound, which we name *Gaussianity-bounded* uncertainty relation, depending on the degree of Gaussianity of the state as measured by a parameter g that we introduce. We identify the corresponding set of minimizing states and find, among them, all the eigenstates of the harmonic oscillator. Non-Gaussian states have proven to be essential in the field of continuous-variable quantum information [13–17], but they remain very hard to classify. Identifying non-Gaussian states of minimum uncertainty among them may help to a better understanding of the structure of the state space in infinite dimensions.

Unconstrained SR relation. Our method is simply introduced as a way to find the MS of the SR uncertainty relation. Consider an arbitrary quantum state characterized by its density operator $\hat{\rho}$. Following the notations of quantum optics, its covariance matrix γ is defined as

$$\gamma_{ij} = \operatorname{Tr}(\{(\hat{r}_i - d_i), (\hat{r}_j - d_j)\}\hat{\rho})$$
(3)

where $\hat{\mathbf{r}} = (\hat{x}, \hat{p})^T$ is the vector of position and momentum observables, $\mathbf{d} = Tr(\hat{\mathbf{r}}\hat{\rho})$ is the displacement vector, and $\{\cdot, \cdot\}$ stands for the anticommutator. Hereafter, we use the variable α to denote the dimensionless *uncertainty* of the state, that is, the left-hand side of Eq. (2) divided by \hbar^2 . In terms of the covariance matrix, it is simply expressed as

$$\alpha = \det \gamma / (4\hbar^2). \tag{4}$$

The determinant of γ , and consequently the uncertainty, remains invariant under the action of the linear canonical group, the semidirect product of the special linear group Sp(2, R) with the translation group T(2). In quantum optics, these correspond to the Gaussian operations, combining displacements and symplectic transformations [18]. Therefore, without loss of generality, we may confine our search of MS among states with a covariance matrix in Williamson normal form (γ proportional to the identity) and $\mathbf{d} = \mathbf{0}$. In other words, we may search for states $\hat{\rho}$ that minimize α while satisfying the constraints

$$\operatorname{Tr}\left(\hat{\rho}\hat{x}\right) = \operatorname{Tr}\left(\hat{\rho}\hat{p}\right) = 0 \tag{5}$$

$$\operatorname{Tr}\left(\hat{\rho}\left(\hat{x}\hat{p}+\hat{p}\hat{x}\right)\right) = \operatorname{Tr}\left(\hat{\rho}\left(\hat{x}^{2}-\hat{p}^{2}\right)\right) = 0 \quad (6)$$

 $\mathbf{2}$

Under these conditions, Eq. (4) can be expressed as

$$\alpha = \operatorname{Tr}\left(\hat{\rho}\left(2\hat{n}+1\right)\right)/4,\tag{7}$$

where $\hat{n} = \hat{a}^{\dagger} \hat{a}$ the number operator.

Now, let us proceed with the optimization of α under the constraints of Eqs. (5)-(6). It is known that for every density matrix $\hat{\rho}$, a unique eigenbasis $\{|\Psi_n\rangle\}$ exists such that $\hat{\rho} = \sum c_n |\Psi_n\rangle \langle \Psi_n|$ with $0 \le c_n \le 1$ and $\sum c_n =$ 1. It is more convenient for our purposes to define the unnormalized vectors $|\psi_n\rangle = \sqrt{c_n} |\Psi_n\rangle$, and rewrite $\hat{\rho}$ as

$$\hat{\rho} = \sum_{n} |\psi_n\rangle \langle\psi_n| \tag{8}$$

while imposing the additional constraint

$$\operatorname{Tr}\left(\hat{\rho}\right) = 1.\tag{9}$$

Then, we choose an orthonormal basis $\{|i\rangle\}$ to decompose the vectors $|\psi_n\rangle = \sum \psi_n^i |i\rangle$ and accordingly re-express the uncertainty (7) as

$$\alpha = \sum_{n,i,j} \psi_n^{i*} \psi_n^j \left\langle i \right| \left(2\hat{n} + 1 \right) \left| j \right\rangle / 4 \tag{10}$$

and the constraints (5), (6), and (9) as

$$\sum_{n,i,j} \psi_n^{i*} \psi_n^j \left\langle i \right| \hat{x} \left| j \right\rangle = 0 \tag{11}$$

$$\sum_{n,i,j} \psi_n^{i*} \psi_n^j \left\langle i \right| \hat{p} \left| j \right\rangle = 0 \tag{12}$$

$$\sum_{n,i,j} \psi_n^{i*} \psi_n^j \left\langle i \right| \left(\hat{x} \hat{p} + \hat{p} \hat{x} \right) \left| j \right\rangle = 0 \tag{13}$$

$$\sum_{n,i,j} \psi_n^{i*} \psi_n^j \left\langle i \right| \left(\hat{x}^2 - \hat{p}^2 \right) \left| j \right\rangle = 0 \tag{14}$$

$$\sum_{n,i} \psi_n^i \psi_n^{i*} = 1. \tag{15}$$

We define the Lagrange multipliers λ_k' and consider the functional

$$\tilde{\alpha} = \alpha + \lambda_1' \operatorname{Tr} \left(\hat{\rho} \right) + \lambda_2' \operatorname{Tr} \left(\hat{\rho} \hat{x} \right) + \lambda_3' \operatorname{Tr} \left(\hat{\rho} \hat{p} \right) + \lambda_4' \operatorname{Tr} \left(\hat{\rho} \left(\hat{x} \hat{p} + \hat{p} \hat{x} \right) \right) + \lambda_5' \operatorname{Tr} \left(\hat{\rho} \left(\hat{x}^2 - \hat{p}^2 \right) \right)$$
(16)

which depends, through Eqs. (11)-(15), on the complex amplitudes ψ_n^i . Extremizing $\tilde{\alpha}$ yields conditions on these amplitudes for the eigenvectors $|\psi_n\rangle$ of the minimizing state $\hat{\rho}$, which read as

$$\left(\hat{n} + \lambda_1 \hat{\mathbf{I}} + \lambda_2 \hat{x} + \lambda_3 \hat{p} + \lambda_4 \left(\hat{x} \hat{p} + \hat{p} \hat{x} \right) \right. \\ \left. + \lambda_5 \left(\hat{x}^2 - \hat{p}^2 \right) \right) \left| \psi_n \right\rangle = 0.$$
 (17)

Introducing the Hermitian operator

$$\hat{H} = \hat{n} + \lambda_2 \hat{x} + \lambda_3 \hat{p} + \lambda_4 \left(\hat{x} \hat{p} + \hat{p} \hat{x} \right) + \lambda_5 \left(\hat{x}^2 - \hat{p}^2 \right), \quad (18)$$

we can rewrite Eq. (17) as $\hat{H} |\psi_n\rangle = -\lambda_1 |\psi_n\rangle$, leading to the following necessary condition: the eigenvectors $|\psi_n\rangle$ of the MS are degenerate eigenvectors of \hat{H} corresponding all to the same eigenvalue $-\lambda_1$.

In order to proceed with the identification of the MS, one should first identify the eigenvectors and eigenvalues of H. This is an easy task because one can always apply a squeezing and displacement operation that transforms H onto the Hamiltonian of the harmonic oscillator, $H_0 =$ $\hat{n} + 1/2$. More precisely, there exists a linear canonical transformation $U = \exp(\gamma a^{\dagger} - \gamma^* a) \exp(\beta a^{\dagger 2} - \beta^* a^2)$ such that $\hat{H} = U\hat{H}_0 U^{\dagger}$, up to a real constant. Obviously, this means that H has the same eigenvalues as H_0 , up to this constant, and that its eigenvectors are the accordingly transformed number states, $U|n\rangle$, remembeing that the number states $|n\rangle$ are the eigenstates of \hat{H}_0 . Since \hat{H}_0 does not possess any degeneracy in its spectrum, the same holds for H and therefore the only possibility is that the MS is a pure state of the type $\hat{\rho} = U |n\rangle \langle n| U^{\dagger}$. Finally, we need to find among these states those that satisfy the constraints (11)-(15). These are obviously the number states $|n\rangle$, and, among them, the one that minimizes the uncertainty Eq. (7) is the ground state of \hat{H}_0 , that is, the vacuum state $|0\rangle$. Of course, we recover the lower bound of the SR relation, $\hbar^2/4$, by plugging in $\hat{\rho} = |0\rangle \langle 0|$ into Eq. (7). Clearly, the orbit of states that are connected to $|0\rangle$ by linear canonical transformations (i.e., all Gaussian pure states) coincides with the well-known set of all MS for the SR uncertainty relation.

Gaussianity-bounded SR relation. Our method only provides necessary conditions on the extremal solution, since it relies on the Lagrange multipliers method, and concluding on a solution may be complicated when the eigenvectors of \hat{H} cannot be identified analytically. We now apply it to find an uncertainty relation under the constraint of a fixed degree of Gaussianity, which is a case where the eigenvalue problem of H is solvable analytically. Several measures of non-Gaussianity have been used in the literature [11, 12, 19–21], but here we instead suggest using a parameter q capturing the degree of Gaussianity, inspired from our work on non-Gaussian states with positive Wigner function [22]. It is more appropriate for our purposes and also has merits on its own, see [23]. Denoting as $\hat{\rho}_G$ the Gaussian state that has the same covariance matrix γ as $\hat{\rho}$, we define the degree of Gaussianity of $\hat{\rho}$ as

$$g = \operatorname{Tr}\left(\hat{\rho}\hat{\rho}_{G}\right) / \operatorname{Tr}\left(\hat{\rho}_{G}^{2}\right).$$
(19)

It is equal to one for all Gaussian states, while it can take values larger or lower than one for other states [23].

As we have seen, our method works by reduction to a constrained optimization problem, even for solving the unconstrained SR inequality. Thus, it can be simply adapted to find the MS with an extra constraint on the degree of Gaussianity g. As before, we can confine our search on states with $\mathbf{d} = \mathbf{0}$ and γ in the Williamson

normal form, since neither α nor g change under linear canonical transformations. For these states, the corresponding Gaussian state $\hat{\rho}_G$ can be expressed simply as

$$\hat{\rho}_G = e^{-\beta \hat{n}} / N \tag{20}$$

where $e^{-\beta} = \frac{4\alpha-1}{4\alpha+1}$ and $N = 2\alpha + 1/2$. In addition, Tr $(\hat{\rho}_G^2) = 1/(4\alpha)$. Instead of minimizing the uncertainty α for a fixed g, it is easier here to fix α and search for states of extremal g, or extremal Tr $(\hat{\rho}\hat{\rho}_G)$. The result of this optimization is equivalent to the original one, but in this way the presentation is more clear. The procedure for deriving the MS is completely analogous, but we extremize the degree of Gaussianity (19) using a constraint on the uncertainty (4), in addition to Eqs. (5)-(6) and (9). The Lagrangian is now

$$\tilde{g} = g + \lambda'_1 \operatorname{Tr}(\hat{\rho}) + \lambda'_2 \operatorname{Tr}(\hat{\rho}\hat{x}) + \lambda'_3 \operatorname{Tr}(\hat{\rho}\hat{p}) + \lambda'_4 \operatorname{Tr}(\hat{\rho}(\hat{x}\hat{p} + \hat{p}\hat{x})) + \lambda'_5 \operatorname{Tr}(\hat{\rho}(\hat{x}^2 - \hat{p}^2))$$
(21)
+ $\lambda'_6 \operatorname{Tr}(\hat{\rho}(2\hat{n} + 1))$ (22)

The extremization conditions that the amplitudes ψ_n^i of the unnormalized eigenvectors $|\psi_n\rangle$ of the solution state must satisfy can be summarized as

$$\left(e^{-\beta \hat{n}} + \lambda_1 \hat{\mathbf{I}} + \lambda_2 \hat{x} + \lambda_3 \hat{p} + \lambda_4 \left(\hat{x} \hat{p} + \hat{p} \hat{x} \right) \right.$$

$$+ \lambda_5 \left(\hat{x}^2 - \hat{p}^2 \right) + \lambda_6 \hat{n} \left. \right) \left| \psi_n \right\rangle = 0.$$

$$(23)$$

By defining a Hermitian operator

$$\hat{H}_{1} = e^{-\beta \hat{n}} + \lambda_{2} \hat{x} + \lambda_{3} \hat{p} + \lambda_{4} \left(\hat{x} \hat{p} + \hat{p} \hat{x} \right) + \lambda_{5} \left(\hat{x}^{2} - \hat{p}^{2} \right) + \lambda_{6} \hat{n}$$
(24)

we conclude that $|\psi_n\rangle$ should be degenerate eigenvectors of \hat{H}_1 . It can be shown that, without loss of generality, we can confine ourselves to states $|\psi_n\rangle = \psi_n |n\rangle$, with ψ_n being complex amplitudes [23]. For these states, the constraints (5)-(6) are satisfied, and \hat{H}_1 is replaced by

$$\hat{H}_2 = e^{-\beta \hat{n}} + \lambda_6 \hat{n}. \tag{25}$$

The eigenvectors of \hat{H}_2 are the number states $|n\rangle$, but, unlike for the harmonic oscillator, double degeneracies are possible if $\lambda_6 < 0$. As a result, we look for mixtures of two number states $\hat{\rho} = |\psi_i|^2 |i\rangle \langle i| + |\psi_j|^2 |j\rangle \langle j|$ satisfying the normalization constraint $|\psi_i|^2 + |\psi_j|^2 = 1$ and the constraint of a fixed uncertainty α ,

$$|\psi_i|^2 (2i+1) + |\psi_j|^2 (2j+1) = 4\alpha$$
 (26)

that achieve the minimum or maximum

$$g = |\psi_i|^2 \frac{8\alpha (4\alpha - 1)^i}{(4\alpha + 1)^{i+1}} + |\psi_j|^2 \frac{8\alpha (4\alpha - 1)^j}{(4\alpha + 1)^{j+1}}.$$
 (27)

By supervision, one can see that the minimum g (corresponding to positive eigenvalues of \hat{H}_2) is achieved by mixtures of two successive number states

$$\hat{\rho}_{\min} = r \left| n \right\rangle \left\langle n \right| + (1 - r) \left| n + 1 \right\rangle \left\langle n + 1 \right|, \qquad (28)$$



FIG. 1: Extremal degree of Gaussianity g for a fixed uncertainty α shown as a dashed line (blue in the online figure). It is achieved by states $\hat{\rho}_{\min}$ for $g \leq 1$, and $\hat{\rho}_{\max}$ for g > 1. The line connecting the number states $|n\rangle$ and $|n+1\rangle$ is realized by mixtures of them. The Gaussianity-bounded uncertainty relation corresponds to a part of this extremal line shown as a solid line (black in the online figure). The uncertainty α must have a value larger or equal to the solid line for a given value of g. Figure (b) is a magnified view of figure (a), where the discontinuity of the uncertainty relation becomes evident.

with the parameters n and $r \in [0, 1]$ depending on α . The number states $|n\rangle$ are naturally included in the set for r = 0. The maximum g (corresponding to negative eigenvalues of \hat{H}_2) is achieved by mixtures

$$\hat{\rho}_{\max} = r \left| 0 \right\rangle \left\langle 0 \right| + (1 - r) \left| n \right\rangle \left\langle n \right| \tag{29}$$

in the limit $n \to \infty$, $r \to 1$, while $4\alpha = r + (1-r)(2n+1)$ is kept finite.

In Figure 1, we plot as a dashed line the two extremal values of g for a fixed value of α as realized by the states in Eqs.(28)-(29). The MS (i.e., the states minimizing the uncertainty α for fixed degree of Gaussianity g) correspond only to some part of this line, which we represent as a solid line. For g > 1, the situation is simple and all states $\hat{\rho}_{\max}$ are MS. In contrast, for $\frac{2}{e} < g \leq 1$, the minimum α for fixed g displays discontinuities. For

$$\frac{\left(m+1\right)^{m+1}\left(3+2m\right)}{\left(2+m\right)^{2+m}} < g \le \frac{m^m \left(1+2m\right)}{\left(1+m\right)^{1+m}} \tag{30}$$

 α is minimized by the states $\hat{\rho}_{\min}$ with n = m. In par-

$$16\alpha \frac{(4\alpha - 1)^m}{(1 + 4\alpha)^{2+m}} (1 + m) = g$$
(31)

in order to derive the dependence of α on g. Beyond this general solution for the MS, we notice from Fig. 1 that for the largest interval $\frac{3}{4} < g \leq 1$ corresponding to m = 0, the uncertainty α is saturated by mixtures of $|0\rangle$ and $|1\rangle$. Thus, ignoring the remaining tiny interval $\frac{2}{e} < g \leq \frac{3}{4}$ (note $\frac{3}{4} - \frac{2}{e} = 0.014$), we can summarize the Gaussianity-bounded uncertainty relation as

$$\alpha \ge \begin{cases} \frac{2+2\sqrt{1-g}-g}{4g}, & \text{if } \frac{3}{4} < g \le 1\\ \frac{g}{8-4g}, & \text{if } g > 1. \end{cases}$$
(32)

Note that, in addition to the above states $\hat{\rho}_{\min}$ and $\hat{\rho}_{\max}$, which are phase invariant, the set of MS also comprises all states that can be transformed onto them by phase averaging, see [23]. For example, superpositions of successive number states are also MS since they can be transformed onto Eq.(28) by phase averaging. Furthermore, all states connected to $\hat{\rho}_{\min}$ and $\hat{\rho}_{\max}$ by linear canonical transformations are MS as well, since the uncertainty α and gaussianity g are invariants of the group.

Our method follows an "inverse" path to the common procedure where the lower bound on the uncertainty relation is derived first based on commutators, and then the MS are identified. Such an inverse procedure was put forward by Dodonov and Man'ko for the derivation of purity-bounded uncertainty relations [7, 11], but, as it appears here, our method is more generally applicable because it is based on the amplitudes of the eigenstates of the MS instead of its density matrix elements, see [23].

Conclusions. We have exhibited an alternative method to derive the uncertainty relation and minimizing states as an eigenvalue problem. This method is especially useful when constraints are included that account for some knowledge on the state. We have found an uncertainty relation that is bounded by the degree of Gaussianity gof the state. The non-Gaussian states with the lowest uncertainty α for a fixed g include (up to linear canonical transformations) mixtures of successive number states $|n\rangle$ and $|n+1\rangle$, as well as the unphysical mixture of vacuum and $|n\rangle$, with $n \to \infty$. Among these minimizing states, the number states play a prominant role as they are the only phase-invariant pure states. Thus, we have proven here that they are also *extremal* according to this new uncertainty relation, thereby extending to all (non-Gaussian) eigenstates of the harmonic oscillator the celebrated uncertainty-minimization property of its (Gaussian) ground state. Given the considerable attention that non-Gaussian states are attracting in continuous-variable quantum information theory, see e.g. [24], unveiling this

extremization property of all number states may hopefully contribute to further progress in the field.

AM gratefully acknowledges financial support from the F.R.S-FNRS. This work was also carried out with the financial support of the F.R.S-FNRS via project HIPER-COM and the support of the Belgian Federal program IUAP via project Photonics@be.

Appendix

Degree of Gaussianity

In our work, we have introduced the quantity

$$g = \operatorname{Tr}\left(\hat{\rho}\hat{\rho}_{G}\right) / \operatorname{Tr}\left(\hat{\rho}_{G}^{2}\right) \tag{A.1}$$

in order to characterize the degree of Gaussianity of a state $\hat{\rho}$. In Eq. (A.1), $\hat{\rho}_G$ is the reference Gaussian state of $\hat{\rho}$ in the sense that it possess the same covariance matrix γ as $\hat{\rho}$. Here, we exhibit the mathematical properties of g, elaborate on the physical intuition behind its definition, and discuss on possible experimental scenarios for its estimation. Let us start by listing some basic mathematical properties of g.

(i) The Gaussianity is invariant under Gaussian operations, i.e., symplectic transformations Sp(2, R) and translations T(2).

Proof. A Gaussian operation U_G acting on a state $\hat{\rho}$ can be always translated as a sequence of displacement, rotation, and squeezing of the Wigner function of $\hat{\rho}$ in the phase-space [25]. The Wigner function of the reference Gaussian state $\hat{\rho}_G$ experiences the same deformation in phase space as that of $\hat{\rho}$, and therefore one may conclude that $\hat{\rho}_G$ experiences the same Gaussian operation U_G as $\hat{\rho}$. Then, using the invariance of trace under cyclic permutations, we have

$$\operatorname{Tr}\left(U_{G}\hat{\rho}U_{G}^{\dagger}U_{G}\hat{\rho}_{G}U_{G}^{\dagger}\right) = \operatorname{Tr}\left(\hat{\rho}\hat{\rho}_{G}\right)$$
$$\operatorname{Tr}\left(U_{G}\hat{\rho}_{G}U_{G}^{\dagger}U_{G}\hat{\rho}_{G}U_{G}^{\dagger}\right) = \operatorname{Tr}\left(\hat{\rho}_{G}^{2}\right), \quad (A.2)$$

so we conclude that g is invariant under Gaussian operations.

(ii) The Gaussianity is a bounded quantity, that is, $2/e \leq g \leq 2$. For Gaussian states g = 1, while the converse is not necessarily true.

Proof. Having identified in this work the extremum values for g as a function of the uncertainty α , it is straightforward to derive the lower and upper bounds of the interval [2/e, 2], which correspond to the limit $\alpha \to \infty$. The state which realizes the lower bound (up to Gaussian transformations) is the number state $|n\rangle$ as $n \to \infty$, while the upper bound is saturated by the state

$$\hat{\rho}_{\max} = (1 - r) \left| 0 \right\rangle \left\langle 0 \right| + r \left| n \right\rangle \left\langle n \right| \tag{A.3}$$

with $r \to 0$ and $n \to \infty$. It is also straightforward to prove that if a state is Gaussian, then g = 1. On the

other hand, g is not constructed on the basis of a mathematical distance, unlike the measures of non-Gaussianity of refs. [19–21]. As a consequence, the inverse statement is not necessarily true and non-Gaussian states might exist possessing a degree of Gaussianity equal to 1.

(*iii*) The Gaussianity provides necessary criterion for strict positivity of the Wigner function of a state.

Proof. In a previous work [22], we have derived bounds on the trace overlap Tr $(\hat{\rho}\hat{\rho}_G)$ for states with strictly positive Wigner function. These bounds have been derived partially analytically and can be easily translated into bounds on the Gaussianity g. Based on the formulas derived in [22], we can prove that a state with strictly positive Wigner function and with an uncertainty $\alpha = \det \gamma/(4\hbar^2)$ is restricted to have a Gaussianity g that lies in the interval $[g_{\min}, g_{\max}]$, where

$$g_{\min} = 0.038\alpha + 0.62 + 0.1778/\alpha - 0.021/\alpha^{2}$$
$$g_{\max} = \sqrt{\frac{2}{1 + \frac{1}{16\alpha^{2}}}}.$$
(A.4)

Finally, it should be noted that the bounds on the Gaussianity g given by our Gaussianity-bounded uncertainty relation provide a necessary criterion for a quantum state to be physical. The classical bivariate distributions which achieve the minimum degree of Gaussianity among all positive-definite distributions (classical and quantum quasi-distributions) of the same covariance matrix have been identified in a previous work [26]. Interestingly, one can always find a classical distribution that possess a smaller g than what is allowed for all quantum states. This indicates that the positivity of the density matrix that is imposed to derive the Gaussianitybounded uncertainty relation is more restrictive than the positivity of a distribution that is imposed on the proof of [26].

To obtain more intuition about the quantity g, it is instructive to explicitly express it in terms of the moments of the Wigner quasi-probability distribution. Let us restrict ourselves to a state $\hat{\rho}$ with a covariance matrix $\gamma = 2\hbar \operatorname{diag}(\alpha, \alpha)$, since by linear canonical transformation all states can be reduced to this form and we have proven that g remains invariant along this transformation. In the Wigner representation, the corresponding reference Gaussian state $\hat{\rho}_G$ of such a state is phaseinvariant; hence, it can be expressed as

$$W_G(r) = \frac{1}{4\pi\alpha} e^{-r^2/4\alpha}, \qquad r = \sqrt{x^2 + p^2}.$$
 (A.5)

In contrast, the state $\hat{\rho}$ itself may possess an angulardependent Wigner function $W(r, \varphi)$. The trace overlap between $\hat{\rho}$ and $\hat{\rho}_G$ takes the following form in the Wigner representation

$$\operatorname{Tr}\left(\hat{\rho}\hat{\rho}_{G}\right) = 2\pi \iint W\left(r,\varphi\right)W_{G}\left(r\right)r\mathrm{d}r\mathrm{d}\varphi \qquad (A.6)$$

while we have $g = 4\alpha \operatorname{Tr}(\hat{\rho}\hat{\rho}_G)$.

Now let us average the phase of the Wigner function $W(r, \varphi)$ in order to construct a new state $\hat{\rho}_s$ with phase-invariant Wigner function $W_s(r)$,

$$W_{s}(r) = \frac{1}{2\pi} \int W(r,\varphi) \, d\varphi. \tag{A.7}$$

Employing Eq.(A.7), the trace overlap Eq. (A.6) can be re-written as Tr $(\hat{\rho}\hat{\rho}_G) = 4\pi^2 \int W_s(r) W_G(r) r dr$, and by expanding $W_G(r)$ in Taylor series we arrive to

$$g = 4\pi \sum_{n=0}^{\infty} \left(\frac{(-1)^n \langle r^{2n+1} \rangle}{n! (4\alpha)^n} \right).$$
 (A.8)

For a Gaussian bivariate distribution independent of the phase as in Eq.(A.5), a simple expression exists for the radial moments $\langle r^{2n+1} \rangle_G = (4\alpha)^n \Gamma(n+1)/2\pi$. Since by definition $\langle r^3 \rangle = \langle r^3 \rangle_G$, we conclude that the Gaussianity g accounts for the difference of the odd (≥ 3) radial moments of the Wigner function of the phase-averaged state $\hat{\rho}_s$ as compared to those of the reference Gaussian state.

Finally, we can use now the phase averaged state $\hat{\rho}_s$ introduced in Eq. (A.7) to prove that in our search for quantum states that extremize g while possessing a covariance matrix of the form $\gamma = 2\hbar \operatorname{diag}(\alpha, \alpha)$, we are allowed to restrict to mixtures of number states. By substitution of Eq. (A.7) into Eq. (A.6), it is evident that

$$\operatorname{Tr}\left(\hat{\rho}\hat{\rho}_{G}\right) = \operatorname{Tr}\left(\hat{\rho}_{s}\hat{\rho}_{G}\right). \tag{A.9}$$

In addition the reference Gaussian state (and therefore the covariance matrix) of $\hat{\rho}_s$ is the same as for $\hat{\rho}$, since phase averaging cannot affect the phase-independent Wigner function Eq. (A.5). From this, one may conclude that g is the same for $\hat{\rho}$ and $\hat{\rho}_s$ and therefore, invariant under the phase randomization procedure for states with covariance matrix of the form $\gamma = 2\hbar \operatorname{diag}(\alpha, \alpha)$. Therefore, we may with no loss of generality confine our search to states with phase-independent Wigner function, which can be expressed as mixtures of number states, i.e., $\hat{\rho} = \sum c_n |n\rangle \langle n|$.

Let us briefly address possible experimental means for a direct estimation of the Gaussianity q of a state $\hat{\rho}$. without going through a full state tomography procedure. The trace overlap between $\hat{\rho}$ and $\hat{\rho}_G$ can be estimated without the need for a full state tomography, using a modified heterodyne detection scheme where the state $\hat{\rho}$ is fed together with $\hat{\rho}_G$ (instead of the vacuum state $|0\rangle$) in the beam splitter preceding the quadrature measurements. However, this would first require performing homodyne measurements on $\hat{\rho}$ for the identification of its covariance and the reconstruction of the reference Gaussian state $\hat{\rho}_G$. This is probably not very practical as multiple copies of $\hat{\rho}$ would be need in order to characterize $\hat{\rho}_G$ before the modified heterodyne measurement could be applied. Other scenarios for measuring g without constructing $\hat{\rho}_G$ from $\hat{\rho}$ may exist, and it is a subject that deserves further investigation.

Generalization to non-linear constraints

The method we have developed here in order to re-derive the SR uncertainty relation and derive the Gaussianity-bounded uncertainty relation, is applicable not only in the case where all constraints are linear in the density matrix elements of $\hat{\rho}$, i.e., of the form Tr $(B\hat{\rho})$, but also in the case where we have non-linear constraints such as the purity Tr $(\hat{\rho}^2)$ or the von Neumann entropy $-\text{Tr}(\hat{\rho} \ln \hat{\rho})$. It is not difficult to show that in the presence of non-linear constraints, the necessary condition on the existence of degeneracies in the spectrum of the Hermitian operator constructed from the constraints is lifted. The necessary condition is, in this case, that every eigenvector $|\psi_n\rangle$ of the density matrix of the solution $\hat{\rho} = \sum |\psi_n\rangle \langle \psi_n|$ should be an eigenvector with positive

- [1] W. Heisenberg, Z. Phys. 43, 122 (1927).
- [2] E. H. Kennard, Z. Phys. 44, 326 (1927).
- [3] H. Weyl, "Theory of groups and quantum mechanics", New York: Dutton, pp. 77, 393-394 (1927).
- [4] L. I. Mandel'shtam and I. E. Tamm, "The uncertainty relation time-energy in nonrelativistic quantum mechanics", Izv. AN USSR, Seriya Fiziki., vol. 9, no.1/2, pp. 122-128 (1945).
- [5] J. E. Moyal, Proc. Cambridge Philos. Soc. 45, 99 (1949).
- [6] D. Stoler and S. Newman, Phys. Lett. A 38, 433 (1972).
- [7] V. V. Dodonov and V. I. Man'ko, in: Invariants and Evolution of Nonstationary Quantum Systems, Proc. Lebedev Physics Institute, Vol. 183, edited by M. A. Markov (Nova Science, Commack, NY, 1989), pp. 3-101.
- [8] E. Schrödinger, Ber. Kgl. Akad. Wiss., Berlin, 24, 296 (1930)
- [9] H. P. Robertson, Phys. Rev. 35, 667A (1930); H. P. Robertson, Phys. Rev. 46, 794 (1934)
- [10] V. V. Dodonov, E. V. Kurmyshev and V. I. Man'ko, Phys. Lett. A 79, 150 (1980).
- [11] V. V. Dodonov, J. Opt. B: Quantum Semiclass. Opt. 4, S98-S108 (2002).
- [12] M. J. Bastiaans, J. Opt. Soc. Am. A, 1243 (1986).
- [13] J. Eisert, S. Scheel, and M. B. Plenio, Phys. Rev. Lett. 89, 137903 (2002).
- [14] J. Fiurasek, Phys. Rev. Lett. 89, 137904 (2002).

eigenvalue of a Hermitian operator derived in a similar way as in the main paper

$$(H - \hat{\rho}) |\psi_n\rangle = 0. \tag{A.10}$$

Moreover, as it is dictated by the above condition Eq. (A.10), the eigenvalues are the mixing amplitudes of the eigenvectors

$$H |\psi_n\rangle = \langle \psi_n | \psi_n \rangle |\psi_n\rangle = c_n |\psi_n\rangle.$$
 (A.11)

In a further work [27], we explicitly show via an example how the method develops in this obviously more complicated case and we draw parallels with the method of derivation of purity bounded uncertainty relations [7],[11].

- [15] G. Giedke and J. I. Cirac, Phys. Rev. A 66, 032316 (2002).
- [16] J. Niset, J. Fiurasek, and N. J. Cerf, Phys. Rev. Lett. 102, 120501 (2009).
- [17] L. Magnin, F. Magniez, A. Leverrier, and N. J. Cerf, Phys. Rev. A 81, 010302(R) (2010).
- [18] E. C. G. Sudarshan, C. B. Chiu and G. Bhamathi, Phys. Rev. A 52, 43 (1995).
- [19] M. G. Genoni, M. G. A. Paris, and K. Banaszek, Phys. Rev. A 76, 042327 (2007).
- M. G. Genoni, M. G. A. Paris and K. Banaszek, Phys. Rev. A 78, 060303(R) (2008); M. G. Genoni and M. G. A. Paris, Phys. Rev. A 82, 052341 (2010).
- [21] J. Solomon Ivan, M. Sanjay Kumar, and R. Simon, Quantum Inf. Process, DOI 10.1007/s11128-011-0314-2 (2011).
- [22] A. Mandilara, E. Karpov and N. J. Cerf, J. Phys.: Conf. Ser. 254, 012011 (2010).
- [23] Appendix.
- [24] A. Ourjoumtsev, H. Jeong, R. Tualle-Brouri and P. Grangier, Nature 448, 784 (2007).
- [25] W.-M. Zhang, D. H. Feng and R. Gilmore, Rev. Mod. Phys. 62, 867 (1990).
- [26] A. Mandilara, E. Karpov and N. J. Cerf, Phys. Rev. A79, 062302 (2009).
- [27] A. Mandilara, E. Karpov and N. J. Cerf, in preparation.