

Proof of two conjectures of Z.-W. Sun on congruences for Franel numbers

Victor J. W. Guo

Department of Mathematics, East China Normal University,
Shanghai 200062, People's Republic of China

jwguo@math.ecnu.edu.cn, <http://math.ecnu.edu.cn/~jwguo>

Abstract. For all nonnegative integers n , the Franel numbers are defined as

$$f_n = \sum_{k=0}^n \binom{n}{k}^3.$$

We confirm two conjectures of Z.-W. Sun on congruences for Franel numbers:

$$\begin{aligned} \sum_{k=0}^{n-1} (3k+2)(-1)^k f_k &\equiv 0 \pmod{2n^2}, \\ \sum_{k=0}^{p-1} (3k+2)(-1)^k f_k &\equiv 2p^2(2^p-1)^2 \pmod{p^5}, \end{aligned}$$

where n is a positive integer and $p > 3$ is a prime.

Keywords: Franel numbers, binomial coefficients, multinomial coefficients, congruences

AMS Subject Classifications: 11A07, 11B65, 05A10, 05A19

1 Introduction

The numbers f_n are defined to be the sums of cubes of binomial coefficients:

$$f_n = \sum_{k=0}^n \binom{n}{k}^3.$$

In 1894, Franel [3, 4] obtained the following recurrence relation for f_n :

$$(n+1)^2 f_{n+1} = (7n^2 + 7n + 2)f_n + 8n^2 f_{n-1}, \quad n = 1, 2, \dots \quad (1.1)$$

Nowadays, the numbers f_n are usually called Franel numbers. The Franel numbers also appear in the first and second Strehl identities [12, 13] (see also Koepf [7, p. 55]):

$$\begin{aligned} \sum_{k=0}^n \binom{n}{k}^3 &= \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{n}, \\ \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 &= \sum_{k=0}^n \sum_{j=0}^k \binom{n}{k} \binom{n+k}{k} \binom{k}{j}^3. \end{aligned}$$

Applying the recurrence relation (1.1), Jarvis and Verrill [6] proved the following congruence for Franel numbers

$$f_n \equiv (-8)^n f_{p-1-n} \pmod{p},$$

where p is a prime and $0 \leq n \leq p-1$. Recently, Z.-W. Sun [15], among other things, proved several interesting congruences for Franel numbers, such as

$$\begin{aligned} \sum_{k=0}^{p-1} (-1)^k f_k &\equiv \left(\frac{p}{3}\right) \pmod{p^2}, \\ \sum_{k=0}^{p-1} (-1)^k k f_k &\equiv -\frac{2}{3} \left(\frac{p}{3}\right) \pmod{p^2}, \\ \sum_{k=0}^{p-1} (-1)^k k^2 f_k &\equiv \frac{10}{27} \left(\frac{p}{3}\right) \pmod{p^2}, \end{aligned}$$

where $p > 3$ is a prime and $\left(\frac{a}{3}\right)$ denotes the Legendre symbol. Sun [15] also proposed many amazing conjectures on congruences for f_n . The main purpose of this paper is to prove the following results, which were conjectured by Sun [15].

Theorem 1.1. *For any positive integer n , there holds*

$$\sum_{k=0}^{n-1} (3k+2)(-1)^k f_k \equiv 0 \pmod{2n^2}. \quad (1.2)$$

Theorem 1.2. *For any prime $p > 3$, there holds*

$$\sum_{k=0}^{p-1} (3k+2)(-1)^k f_k \equiv 2p^2(2^p-1)^2 \pmod{p^5}. \quad (1.3)$$

2 Proof of Theorem 1.1

We need the following identity due to MacMahon [8, p. 122] (see also Foata [2] and Riordan [10, p. 41]):

$$\sum_{k=0}^n \binom{n}{k}^3 x^k = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n+k}{3k} \binom{3k}{2k} \binom{2k}{k} x^k (1+x)^{n-2k}. \quad (2.1)$$

When $x = 1$, the above identity (2.1) gives a new expression for Franel numbers:

$$\sum_{k=0}^n \binom{n}{k}^3 = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n+k}{3k} \binom{3k}{2k} \binom{2k}{k} 2^{n-2k}. \quad (2.2)$$

Differentiating both sides of (2.1) twice with respect to x and then letting $x = 1$, we get

$$\sum_{k=0}^n \binom{n}{k}^3 k(k-1) = \sum_{k=0}^{\lfloor n/2 \rfloor} 2^{n-2k} \binom{n+k}{3k} \binom{3k}{2k} \binom{2k}{k} \frac{n(n-1)-2k}{4}. \quad (2.3)$$

Moreover, by induction, we can easily prove that

$$\sum_{\ell=2k}^{n-1} (-1)^\ell (3\ell+2) \binom{\ell+k}{3k} 2^{\ell-2k} = (n-2k)(-1)^{n-1} \binom{n+k}{3k} 2^{n-2k}. \quad (2.4)$$

Applying (2.2) and then exchanging the summation order, we have

$$\sum_{k=0}^{n-1} (3k+2)(-1)^k f_k = (-1)^{n-1} \sum_{k=0}^{n-1} 2^{n-2k} (n-2k) \binom{n+k}{3k} \binom{3k}{2k} \binom{2k}{k} \quad (2.5)$$

in view of (2.4). Noticing that

$$n-2k = 4 \frac{n(n-1)-2k}{4} - (n^2-2n),$$

by (2.2) and (2.3), we can write the right-hand side of (2.5) as

$$\begin{aligned} & (-1)^{n-1} 4 \sum_{k=0}^n \binom{n}{k}^3 k(k-1) + (-1)^n (n^2-2n) \sum_{k=0}^n \binom{n}{k}^3 \\ &= (-1)^{n-1} 4n^2 \sum_{k=0}^n \binom{n}{k} \binom{n-1}{k-1}^2 + (-1)^n n^2 \sum_{k=0}^n \binom{n}{k}^3, \end{aligned}$$

where we have used the following identities:

$$\begin{aligned} k \binom{n}{k} &= n \binom{n-1}{k-1}, \\ \sum_{k=0}^n k \binom{n}{k}^3 &= \sum_{k=0}^n (n-k) \binom{n}{k}^3 = \frac{n}{2} \sum_{k=0}^n \binom{n}{k}^3. \end{aligned}$$

Namely, we have proved that

$$\frac{1}{2n^2} \sum_{k=0}^{n-1} (3k+2)(-1)^k f_k = (-1)^{n-1} 2 \sum_{k=0}^n \binom{n}{k} \binom{n-1}{k-1}^2 + (-1)^n \frac{f_n}{2}. \quad (2.6)$$

The proof then follows from the fact

$$f_n = \sum_{k=0}^n \binom{n}{k}^3 \equiv \sum_{k=0}^n \binom{n}{k} = 2^n \equiv 0 \pmod{2}.$$

By (2.2), it is easily seen that, for $n \geq 1$,

$$f_n \equiv \begin{cases} 2, & \text{if } n \text{ is a power of 2,} \\ 0, & \text{otherwise,} \end{cases} \pmod{4}.$$

Thus, we may further refine Theorem 1.1 as follows:

Theorem 2.1. *For any positive integer n , there holds*

$$\sum_{k=0}^{n-1} (3k+2)(-1)^k f_k \equiv \begin{cases} 2n^2, & \text{if } n \text{ is a power of 2,} \\ 0, & \text{otherwise,} \end{cases} \pmod{4n^2}. \quad (2.7)$$

3 Proof of Theorem 1.2

The following lemma is due to Sun [15, Lemma 2.1].

Lemma 3.1 (Sun). *For any prime $p > 3$, there holds*

$$f_{p-1} \equiv 1 + 3(2^{p-1} - 1) + 3(2^{p-1} - 1)^2 \pmod{p^3}. \quad (3.1)$$

To prove Theorem 1.2, we also need the following variation of Lemma 3.1.

Lemma 3.2. *For any prime $p > 3$, there holds*

$$\sum_{k=1}^{p-1} \binom{p-1}{k} \binom{p-1}{k-1}^2 \equiv 2^{p-1} - 2^{2p-2} \pmod{p^3}. \quad (3.2)$$

Proof. It is easy to verify the following identity

$$\sum_{k=0}^p \binom{p-1}{k}^3 = \sum_{k=0}^p \binom{p}{k}^3 - \sum_{k=0}^p \binom{p-1}{k-1}^3 - 3 \sum_{k=0}^p \binom{p}{k} \binom{p-1}{k} \binom{p-1}{k-1}.$$

Hence, by Sun's congruence (3.1) and the known congruence

$$\sum_{k=0}^p \binom{p}{k}^3 \equiv 2 \pmod{p^3}, \quad (3.3)$$

we immediately get

$$\sum_{k=0}^p \binom{p}{k} \binom{p-1}{k} \binom{p-1}{k-1} \equiv 2^p - 2^{2p-1} \pmod{p^3}. \quad (3.4)$$

On the other hand, replacing k by $p-k$, we have

$$\sum_{k=1}^{p-1} \binom{p-1}{k} \binom{p-1}{k-1}^2 = \sum_{k=1}^{p-1} \binom{p-1}{k}^2 \binom{p-1}{k-1} = \frac{1}{2} \sum_{k=0}^p \binom{p}{k} \binom{p-1}{k} \binom{p-1}{k-1}. \quad (3.5)$$

Combining (3.4) and (3.5), complete the proof. $\square$

Proof of Theorem 1.2. By (2.6) and (3.3), we have

$$\begin{aligned} \frac{1}{2p^2} \sum_{k=0}^{p-1} (3k+2)(-1)^k f_k &= 2 \sum_{k=0}^n \binom{p}{k} \binom{p-1}{k-1}^2 - 1, \\ &= 2 \sum_{k=0}^n \binom{p-1}{k} \binom{p-1}{k-1}^2 + 2 \sum_{k=0}^n \binom{p-1}{k-1}^3 - 1. \end{aligned}$$

The proof then follows from (3.1) and (3.2). $\square$

4 Concluding remarks and open problems

For any nonnegative integers n and r , let

$$f_n^{(r)} = \sum_{k=0}^n \binom{n}{k}^r.$$

Then $f_n^{(3)} = f_n$ are the Franel numbers. Calkin [1, Proposition 3] proved the following congruence:

$$f_n^{(2r)} \equiv 0 \pmod{p},$$

where p is a prime such that $\frac{n}{m} < p < \frac{n+1}{m} + \frac{n+1-m}{m(2mr-1)}$ for some positive integer m . Guo and Zeng [5, Theorem 4.4] proved that

$$f_n^{(2r)} \equiv 0 \pmod{n+1}. \quad (4.1)$$

Sun [14, Conjecture 3.5] conjectured that

$$\sum_{k=0}^{n-1} (3k+2) f_k^{(4)} \equiv 0 \pmod{2n}. \quad (4.2)$$

It is easy to see that $f_n^{(0)} = n+1$, $f_n^{(1)} = 2^n$, $f_n^{(2)} = \binom{2n}{n}$ by the Chu-Vandermonde identity (see [7, p. 41]), and

$$\sum_{k=0}^{n-1} (3k+2) f_k^{(0)} = n^3 + n^2, \quad (4.3)$$

$$\sum_{k=0}^{n-1} (-1)^k (3k+2) f_k^{(1)} = (-1)^{n-1} 2^n n, \quad (4.4)$$

$$\sum_{k=0}^{n-1} (3k+2) f_k^{(2)} = n \binom{2n}{n}. \quad (4.5)$$

Motivated by the identities (4.3)–(4.5), the congruence (1.2), and Sun's conjecture (4.2), we would like to propose the following conjecture on congruences for $f_n^{(r)}$.

Conjecture 4.1. *Let $n \geq 1$ and $r \geq 0$ be two integers. Then*

$$\sum_{k=0}^{n-1} (-1)^{rk} (3k+2) f_k^{(r)} \equiv 0 \pmod{2n}. \quad (4.6)$$

By (4.1), if the congruence (4.6) holds, then we have

$$\sum_{k=0}^{n-1} (3k+2) f_k^{(2r)} \equiv 0 \pmod{n(n+1)}.$$

Moreover, for $n > 1$, the following congruence seems to be true:

$$\sum_{k=0}^{n-1} (-1)^k (3k+2) f_k^{(2r+1)} \equiv 0 \pmod{4n}.$$

Recall that the multinomial coefficients are given by

$$\binom{n}{k_1, \dots, k_m} = \frac{n!}{k_1! \cdots k_m!},$$

where $k_1, \dots, k_m \geq 0$ and $k_1 + \cdots + k_m = n$. Let

$$M_{m,n}^{(r)} = \sum_{k_1 + \cdots + k_m = n} \binom{n}{k_1, \dots, k_m}^r$$

be the sums of r th powers of multinomial coefficients. Then $M_{m,n}^{(0)} = \binom{m+n-1}{n}$, $M_{m,n}^{(1)} = m^r$, $M_{1,n}^{(r)} = 1$, $M_{2,n}^{(r)} = f_n^{(r)}$, and

$$M_{3,n}^{(2)} = \sum_{k=0}^n \sum_{j=0}^k \binom{n}{k}^2 \binom{k}{j}^2 = \sum_{k=0}^n \binom{n}{k}^2 \binom{2k}{k}.$$

Note that Osburn and Sahu [9] studied supercongruences for the numbers $\sum_{k=0}^n \binom{n}{k}^r \binom{2k}{k}^s$. The sequence $\{M_{3,n}^{(2)}\}_{n \geq 0}$ is the A002893 sequence of Sloane [11]. It also appears in Zagier [16, #8 of Table 1]. Z.-W. Sun [15] proved the following identity involving $M_{3,k}^{(2)}$:

$$\sum_{k=0}^{n-1} (4k+3) M_{3,k}^{(2)} = 3n^2 \sum_{k=0}^{n-1} \frac{1}{k+1} \binom{2k}{k} \binom{n-1}{k}^2. \quad (4.7)$$

It seems that Conjecture 4.1 can be further generalized as follows.

Conjecture 4.2. *Let $m, n \geq 1$ and $r \geq 0$ be integers. Then*

$$\sum_{k=0}^{n-1} (-1)^{rk} ((m+1)k+m) M_{m,k}^{(r)} \equiv 0 \pmod{mn}. \quad (4.8)$$

It is not hard to prove the following identities by induction.

$$\sum_{k=0}^{n-1} ((m+1)k+m) M_{m,k}^{(0)} = mn \binom{m+n-1}{m}, \quad (4.9)$$

$$\sum_{k=0}^{n-1} (-1)^k ((m+1)k+m) M_{m,k}^{(1)} = (-1)^{n-1} m^n n, \quad (4.10)$$

$$\sum_{k=0}^{n-1} (2k+1) M_{1,k}^{(2r)} = n^2, \quad (4.11)$$

$$\sum_{k=0}^{n-1} (-1)^k (2k+1) M_{1,k}^{(2r+1)} = (-1)^{n-1} n. \quad (4.12)$$

Combining the congruence (1.2), the identities (4.5), (4.7) and (4.9)–(4.12), we have the following result to support Conjecture 4.2.

Proposition 4.3. *The congruence (4.8) holds if (m, r) belongs to*

$$\{(m, 0), (m, 1), (1, r), (2, 2), (2, 3), (3, 2) : m, r \geq 1\}.$$

Acknowledgments. This work was partially supported by the Fundamental Research Funds for the Central Universities.

References

- [1] N.J. Calkin, Factors of sums of powers of binomial coefficients, *Acta Arith.* 86 (1998), 17–26.
- [2] D. Foata, Etude algébrique de certains problèmes d’analyse combinatoire et du calcul des probabilités, *Publ. Inst. Statist. Univ. Paris* 14 (1965), 81–241.
- [3] J. Franel, On a question of Laisant, *L’intermédiaire des mathématiciens* 1 (1894), 45–47.
- [4] J. Franel, On a question of J. Franel, *L’intermédiaire des mathématiciens* 2 (1895), 33–35.
- [5] V. J. W. Guo and J. Zeng, New congruences for sums involving Apéry numbers or central Delannoy numbers, preprint, arXiv: 1008.2894.
- [6] F. Jarvis and H. A. Verrill, Supercongruences for the Catalan-Larcombe-French numbers, *Ramanujan J.* 22 (2010), 171–186.
- [7] W. Koepf, *Hypergeometric Summation, an Algorithmic Approach to Summation and Special Function Identities*, Friedr. Vieweg & Sohn, Braunschweig, 1998.
- [8] P. A. MacMahon, *Combinatorial Analysis*, Vol. 1, Cambridge University Press, London, 1915.
- [9] R. Osburn and B. Sahu, Supercongruences for Apéry-like numbers, *Adv. Appl. Math.* 47 (2011), 631–638.
- [10] J. Riordan, *Combinatorial Identities*, J. Wiley, New York, 1979.

- [11] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, available at <http://www.research.att.com/~njas/sequences/>
- [12] V. Strehl, Binomial sums and identities, Maple Technical Newsletter 10 (1993), 37–49.
- [13] V. Strehl, Binomial identities — combinatorial and algorithmic aspects, Discrete Math. 136 (1994), 309–346.
- [14] Z.-W. Sun, Conjectures and results on $x^2 \bmod p^2$ with $4p = x^2 + dy^2$, In: Number Theory and the Related Topics, Eds., Y. Ouyang, C. Xing, F. Xu and P. Zhang, Higher Education Press & International Press, Beijing and Boston, 2012, to appear.
- [15] Z.-W. Sun, Congruences for Franel numbers, preprint, arXiv:1112.1034v6.
- [16] D. Zagier, Integral solutions of Apéry-like recurrence equations, In: J. Harnad, P. Winternitz, Eds., Group and Symmetries: From Neolithic Scots to John McKay, CRM Proceedings & Lecture Notes, 47, American Mathematical Society, Providence, RI, 2009, pp. 349–366.