

What if Superluminal Neutrinos Exist but not Higgs Bosons?

B.G. Sidharth

International Institute for Applicable Mathematics & Information Sciences
Hyderabad (India) & Udine (Italy)

B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500 063 (India)

(Based on the Bhagawantham Memorial Lecture 2011)

Abstract

As 2011 ended, two results stood out which seemed to go against twentieth century ideas. The first was the OPERA superluminal neutrino observation contradicting Special Relativity. The second was lack of a definitive appearance of the Higgs Boson. While both these hopefully will be decided by the end of 2012, we investigate a single mechanism that explains both these anomalies.

1 Introduction

One of the pillars of twentieth century physics has been Einstein's Special Theory of Relativity according to which the speed of light is the maximum limit in the universe.

Similarly another pillar of last century's physics has been the Standard Model of Particle Physics which has been in place from around 1970. The year 2011 saw doubts cast on both these well established theories. First came an announcement on 23rd September that neutrinos which were let off from CERN in Geneva, reached the GRAN SASSO Lab in Central Italy some sixty nano seconds too early, thus apparently breaching the speed of light barrier and the Special Theory of Relativity. This was a 6σ result. Even so, the experiment was then repeated, with smaller bunches of neutrinos to avoid possible

errors. Again the same result was obtained.

On the other hand, after several false rumours that the Higgs Boson, a missing but vital piece of the Standard Model had been detected at LHC in CERN, a Conference was held on 13th December to unveil the latest conclusions of the CMS and ATLAS teams. It was widely anticipated that the discovery of the Higgs Boson would be announced. Contrary to the build up of expectations, the announcement was bland. There were mere hints of the Higgs Boson, but no definite sighting. The definitive information that came out was that the Higgs Boson, if it exists, would have a mass of about 125GeV . Both these puzzles may be resolved by end 2012.

This prompts us to play the role of the devil's advocate and ask, what if superluminal neutrinos are a reality, but not the Higgs Boson? There are alternative theories that explain separately either of these two anomalies. But we look for a single explanation for both these apparently unrelated issues. This is to be found if we discard a common assumption – or plank – on which twentieth century physics rests: a smooth spacetime. But if spacetime were not so, as for example in Quantum Gravity approaches, that is spacetime were non-commutative, then both the anomalies can be explained, as we see below.

2 Superluminal Neutrinos

The OPERA (Oscillation Project with Emulsion Tracking Apparatus) experiment, 1400 meters underground in the Gran Sasso National Laboratory in Italy has detected neutrinos travelling faster than the speed of light, which has been a well acknowledged speed barrier in physics. This limit is 299792,458 meters per second, whereas the experiment has detected a speed of 299,798,454 meters per second. In this experiment neutrinos from the CERN Laboratory 730 kilometers away in Geneva were observed. They arrived 60 nano seconds faster than expected, that is faster than the time allowed by the speed of light. The experiment has been measured to 6σ level of confidence, which makes it a certainty [1]. However it is such an astounding discovery that the OPERA scientists would like further confirmation from other parts of the world. In the meantime they performed the experiment all over again, but this time using smaller bunches of neutrinos, to eliminate certain possible errors. The result was the same. In 2007 the MINOS experiment near Chicago did find hints of this superluminal effect [2]. Nevertheless

scientists wait with bated breath to confirm this earth shattering discovery. The best direct test of Einstein's energy mass formula so far has been made by combining accurate measurements of atomic mass differences Δm and of the γ -ray wavelengths to determine the energy, the nuclear binding energy for isotopes of silicon and sulphur [3]. They found that the energy mass formula can be separately confirmed in two tests yielding a combined result of $1 - \Delta mc^2/E = (-1.4 \pm 1.4) \times 10^{-7}$, indicating that it holds to a level of at least 0.00004%.

It must be reported that the author had predicted such deviations from Einstein's Theory of Relativity, starting from 2000. This work replaces the usual Einstein energy momentum formula with the modified expression (the so called Snyder-Sidharth Hamiltonian),

$$E^2 = p^2 c^2 + m^2 c^4 + \frac{\alpha c^2}{\hbar^2} l^2 p^4 \quad (1)$$

where l is a minimum length like the Planck length and α is positive for fermions or spin half particles like neutrinos [4, 5, 6, 7, 8]. The above formula is based on considerations of a non differentiable spacetime at ultra high energies. In this case, the usual commutative relations of Quantum theory are replaced, as shown a long time ago by Snyder (Cf.refs.[5, 7]) by

$$[x, y] = O(l^2), [x, p_x] = i\hbar[1 + l^2] \quad (2)$$

Equation (1) shows that the energy at very high energies for fermions is greater than that given by the relativity theory so that effectively the speed of the particle is slightly greater than that of light. For example, if in the usual formula, we replace c by $c + c'$, then, comparing with the above we would get:

$$c' = \alpha l^2 \cdot \frac{c^2}{\hbar^2} p^4 [4m^2 c^3 + 2p^2 c]^{-1}$$

The difference is slight, but as can be seen is maximum for the lightest fermions, viz., neutrinos. The above formula simplifies to

$$c' = \frac{\alpha}{6} c,$$

taking the neutrino mass to be $10^{-3} eV$ and l to be the neutrino Compton wavelength. A value, $\alpha \sim 1.2 \times 10^{-4}$ reproduces the Gran Sasso result.

There are other interesting ramifications of this relation, for example the mass of a particle and its antiparticle may differ slightly, and so on (Cf. also [9]).

3 The Higgs Boson

It is well known that in the Standard Model, Peter Higgs and a few others invoked the idea of the Higgs Boson via a BCS mechanism to explain why and how elementary particles acquire a mass. However, as pointed out in the introduction, from the mid sixties the search for the Higgs has as of date failed to locate the particle. We would now like to point out that based on the above fuzzy or non commutative spacetime we get a formulation that mimics the Higgs mechanism to generate mass, without actually requiring a new particle.

It is well known that Hermann Weyl's original phase transformation proposal was generalized, so that the global or constant phase of λ was considered to be a function of the coordinates [7, 10, 11, 12].

As is well known this leads to a covariant gauge derivative. For example, the transformation arising from $(x^\mu) \rightarrow (x^\mu + dx^\mu)$,

$$\psi \rightarrow \psi e^{-i\lambda} \quad (3)$$

leads to the familiar electromagnetic potential gauge,

$$A_\mu \rightarrow A_\mu - \partial_\mu \lambda \quad (4)$$

The above transformation, ofcourse, is a symmetry transformation. In the transition from (3) to (4), we expand the exponential, retaining terms only to the first order in coordinate differentials.

Let us now consider the gauge field in some detail. As is known this could be obtained as a generalization of the above phase function λ to include fields with internal degrees of freedom. For example λ could be replaced by A_μ given by [10]

$$A_\mu = \sum_i A_\mu^i(x) L_i, \quad (5)$$

The gauge field itself would be obtained by using Stoke's Theorem and (5). This is a very well known procedure: considering a circuit, which for simplicity we can take to be a parallelogram of side dx and dy in two dimensions, we can easily deduce the equation for the field, viz.,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - iq[A_\mu, A_\nu], \quad (6)$$

q being the gauge field coupling constant.

In (6), the second term on the right side is typical of a non Abelian gauge

field. In the case of the U(1) electromagnetic field, this latter term vanishes. Further as is well known, in a typical Lagrangian like

$$L = i\bar{\psi}\gamma^\mu D_\mu\psi - \frac{1}{4}F^{\mu\nu}F_{\mu\nu} - m\bar{\psi}\psi \quad (7)$$

D denoting the Gauge covariant derivative, there is no mass term for the field Bosons. Such a mass term in (7) must have the form $m^2 A^\mu A_\mu$ which unfortunately is not Gauge invariant.

This was the shortcoming of the original Yang-Mills Gauge Theory: The Gauge Bosons would be massless and hence the need for a symmetry breaking, mass generating mechanism.

The well known remedy for the above situation has been to consider, in analogy with superconductivity theory, an extra phase of a self coherent system (Cf.ref.[10] for a simple and elegant treatment and also refs. [11] and [13]). Thus instead of the gauge field A_μ , we consider a new phase adjusted gauge field after the symmetry is broken

$$W_\mu = A_\mu - \frac{1}{q}\partial_\mu\phi \quad (8)$$

The field W_μ now generates the mass in a self consistent manner via a Higgs mechanism. Infact the kinetic energy term

$$\frac{1}{2}|D_\mu\phi|^2, \quad (9)$$

where D_μ in (9) denotes the Gauge , now becomes

$$|D_\mu\phi_0|^2 = q^2|W_\mu|^2|\phi_0|^2, \quad (10)$$

Equation (10) gives the mass in terms of the ground state ϕ_0 .

The whole point is as follows: The symmetry breaking of the gauge field manifests itself only at short length scales signifying the fact that the field is mediated by particles with large mass. Further the internal symmetry space of the gauge field is broken by an external constraint: the wave function has an intrinsic relative phase factor which is a different function of spacetime coordinates compared to the phase change necessitated by the minimum coupling requirement for a free particle with the gauge potential. This cannot be achieved for an ordinary point like particle, but a new type of a physical

system, like the self coherent system of superconductivity theory now interacts with the gauge field. The second or extra term in (8) is effectively an external field, though (10) manifests itself only in a relatively small spatial interval. The ϕ of the Higgs field in (8), in analogy with the phase function of Cooper pairs of superconductivity theory comes with a Landau-Ginzburg potential $V(\phi)$.

Let us now consider in the gauge field transformation, an additional phase term, $f(x)$, this being a scalar. In the usual theory such a term can always be gauged away in the U(1) electromagnetic group. However we now consider the new situation of a noncommutative geometry discussed earlier viz.,

$$[dx^\mu, dx^\nu] = \Theta^{\mu\nu} \beta, \beta \sim 0(l^2) \quad (11)$$

where l denotes a minimum spacetime cut off. Equation (11) is infact Lorentz covariant. Then the f phase factor gives a contribution to the second order in coordinate differentials,

$$\begin{aligned} & \frac{1}{2} [\partial_\mu B_\nu - \partial_\nu B_\mu] [dx^\mu, dx^\nu] \\ & + \frac{1}{2} [\partial_\mu B_\nu + \partial_\nu B_\mu] [dx^\mu dx^\nu + dx^\nu dx^\mu] \end{aligned} \quad (12)$$

where $B_\mu \equiv \partial_\mu f$.

As can be seen from (12) and (11), the new contribution is in the term which contains the commutator of the coordinate differentials, and not in the symmetric second term. Effectively, remembering that B_μ arises from the scalar phase factor, and not from the non-Abelian gauge field, in equation (6) A_μ is replaced by

$$A_\mu \rightarrow A_\mu + B_\mu = A_\mu + \partial_\mu f \quad (13)$$

Comparing (13) with (8) we can immediately see that the effect of noncommutativity is precisely that of providing a new symmetry breaking term to the gauge field, instead of the ϕ term, (Cf.refs. [14, 15]) a term not belonging to the gauge field itself.

On the other hand if we neglect in (11) terms $\sim l^2$, then there is no extra contribution coming from (12) or (13), so that we are in the usual non-Abelian gauge field theory, requiring a broken symmetry to obtain an equation like (13).

4 Comments

We note that some of the objections to the superluminal neutrino experiment have included: Energy loss due to Cerenkov radiation and consequently a slowing down of the neutrinos. Or an error in the GPRS determination of emission and arrival times. This will be addressed with fibre optic network instead of GPRS. On the other hand extra dimensions of String theory have been invoked to explain the superluminal feature.

As for the Higgs Bosons, effects like techni-colour have been invoked as a mass generating mechanism.

Appendix

To see the above modified SS Hamiltonian in greater detail, we note that, as shown by the author in early 2000 [16] given a minimum length l , the energy momentum relation gets modified. The usual Quantum Mechanical commutation relations get modified as shown in Section 2 (Cf.(2)) and now become

$$[x, p] = \hbar' = \hbar \left[1 + \left(\frac{l}{\hbar} \right)^2 p^2 \right] \text{ etc} \quad (14)$$

(Cf. also ref.[17]). (14) shows that effectively \hbar is replaced by \hbar' . So, in units, $\hbar = 1 = c$,

$$E = [m^2 + p^2(1 + l^2 p^2)^{-2}]^{\frac{1}{2}}$$

or, the energy-momentum relation leading to the Klein-Gordon Hamiltonian is given by,

$$E^2 = m^2 + p^2 - 2l^2 p^4, \quad (15)$$

neglecting higher order terms. This is the so called Snyder-Sidharth Hamiltonian for Bosons [18]. (It may be mentioned that some other authors have since ad hoc taken a third power of p , and so on [19]. However we should remember that these were mostly phenomenological approaches.)

For Fermions the analysis can be more detailed, in terms of Wilson lattices [20]. The free Hamiltonian now describes a collection of harmonic fermionic oscillators in momentum space. Assuming periodic boundary conditions in

all three directions of a cube of dimension L^3 , the allowed momentum components are

$$\mathbf{q} \equiv \left\{ q_k = \frac{2\pi}{L} v_k; k = 1, 2, 3 \right\}, \quad 0 \leq v_k \leq L - 1 \quad (16)$$

(16) finally leads to

$$E_{\mathbf{q}} = \pm \left(m^2 + \sum_{k=1}^3 a^{-2} \sin^2 q_k \right)^{1/2} \quad (17)$$

where $a = l$ is the length of the lattice, this being the desired result leading to

$$E^2 = p^2 c^2 + m^2 c^4 + \alpha l^2 p^4 \quad (18)$$

(18) shows that α is positive, that is for Fermions the Snyder-Sidharth Hamiltonian is given by (18) as noted in Section 2.

We point out that using the modified dispersion relation (18), for a massless particle, $m = 0$, and identifying the extra term $l^2 p^4$ as being due to a mass δm , we can easily deduce that, restoring proper units,

$$\frac{c^2}{\hbar^2} l^2 p^4 = \Delta E^2 = \delta m^2 c^4,$$

Whence,

$$\delta m = \frac{\hbar}{cl} \quad \text{or} \quad l = \frac{\hbar}{c\delta m}$$

This shows that l is the Compton wavelength for this mass δm or alternatively if l is the Compton wavelength, then we deduce the mass, now generated from the extra effect. This is another demonstration of mass generation from $O(l^2)$ effects as seen in Section 3, without requiring a Higgs mechanism. If, for example, l were the Planck length, then δm would be the Planck mass (and vice versa).

References

- [1] Adam, T. et al., *arXiv 1109.4897*.
- [2] Adamson, P., et al (2007). *Phys.Rev. D* 76, 2007, 072005.

- [3] Rainville, S. et al. (2005). *Nature* Vol.438, No.22, 29 December 2005, pg.1096.
- [4] Sidharth, B.G. (2001). *Chaos, Solitons & Fractals* 12, 2001, pp.1449-1457.
- [5] Sidharth, B.G. (2001). *Chaotic Universe: From the Planck to the Hubble Scale* (Nova Science, New York), p.143.
- [6] Sidharth, B.G. (2008). *Foundation of Physics* 38 (1), 2008, pp89ff.
- [7] Sidharth, B.G. (2005). *The Universe of Fluctuations* (Springer, Netherlands).
- [8] Sidharth, B.G. (2008). *Thermodynamic Universe* (World Scientific, Singapore) and other references therein.
- [9] Sidharth, B.G. (2011). *New Advances in Physics* 5 (2), 2011, p.106.
- [10] Moriyasu, K. (1983). *An Elementary Primer for Gauge Theory* (World Scientific, Singapore).
- [11] Jacob, M. (1974). *Physics Reports, Reprint Volume* (North-Holland, Amsterdam).
- [12] Greiner, W. and Reinhardt, I. (1995). *Gauge Theory of Weak Interactions* (Springer-Verlag, Berlin).
- [13] Taylor, J.C. (1978). *Gauge Theories of Weak Interactions* (Cambridge University Press, Cambridge).
- [14] Sidharth, B.G. (2004). *Proceedings of the Fifth International Symposium on "Frontiers of Fundamental Physics"* (Universities Press, Hyderabad).
- [15] Sidharth, B.G. (2006). *Int.J.SCI.* January 2006, pp.42ff.
- [16] Sidharth, B.G. (2000). *Chaos, Solitons and Fractals* 11, 2000.
- [17] Sidharth, B.G. (2005). *Chaos, Solitons and Fractals* 25, pp.965–968.
- [18] Glinka, L. (2008). *arXiv:hep-th 0812.0551v1*.
- [19] Myers, R.C. and Pospelov, M. (2003). *Phys.Rev.Lett.* 90:211601.

- [20] Montvay, I. and Munster, G. (1994). *Quantum Fields on a Lattice* (Cambridge University Press) pp.174ff.