

# Kinetic Wave Turbulence

Gregory L. Eyink<sup>1,2</sup> and Yi-Kang Shi<sup>1</sup>

(1) *Department of Applied Mathematics & Statistics and (2) Department of Physics & Astronomy,  
The Johns Hopkins University, Baltimore, MD, USA*

## Abstract

We consider a general model of Hamiltonian wave systems with triple resonances, in the standard kinetic limit of a continuum of weakly interacting dispersive waves with random phases. In this asymptotic limit we show that the correct dynamical equation for multimode amplitude distributions is *not* the well-known equation of Peierls, Brout & Prigogine, and Zaslavskii & Sagdeev but is instead a reduced equation with only a subset of the terms in that equation. The equations that we derive are the direct analogue of the Boltzmann hierarchy obtained from the BBGKY hierarchy in the low-density limit for gases. We show that the asymptotic multimode equations possess factorized solutions for factorized initial data, which correspond to preservation in time of the property of “random phases & amplitudes”. The factors satisfy the equations for the 1-mode probability density functions (PDF’s) previously derived by Jakobsen & Newell and Choi et al. Analogous to the Klimontovich density in the kinetic theory of gases, we introduce the concepts of the “empirical spectrum” and the “empirical 1-mode PDF”. We show that the factorization of the hierarchy equations implies that these quantities are self-averaging: they satisfy the wave-kinetic closure equations of the spectrum and 1-mode PDF for almost any selection of phases and amplitudes from the initial ensemble. We show that both of these closure equations satisfy an  $H$ -theorem for an entropy defined by Boltzmann’s prescription  $S = k_B \log W$ . We also characterize the general solutions of our multimode distribution equations, for initial conditions with random phases but with no statistical assumptions on the amplitudes. Analogous to a result of Spohn for the Boltzmann hierarchy, these are “super-statistical solutions” that correspond to ensembles of solutions of the wave-kinetic closure equations with random initial conditions or random forces. On the basis of our results, we discuss possible kinetic explanations of intermittency and non-Gaussian statistics in wave turbulence. In particular, we advance the explanation of a “super-turbulence” produced by stochastic or turbulent solutions of the wave kinetic equations themselves.

*Keywords:* Keywords here.

## 1. Introduction and Outline of Main Results

The theory of wave turbulence has traditionally focused on the wavenumber spectrum, which is expected to satisfy the wave kinetic equation in the limit of a continuum of weakly interacting, phase-incoherent waves [1]. However, recent very interesting works of Jakobsen & Newell (2004) [2] and Choi et al. (2005) [3, 4] have studied higher-order fluctuations of wave amplitudes by deriving equations for probability density functions (PDF’s) in this same kinetic limit. These works generalized the results of a pioneering study of Zaslavskii & Sagdeev (1967) [5], who obtained in some special 3-wave systems a set of evolution equations for multimode PDF’s of wave amplitudes and phases. These are analogues of the equations found even earlier by Peierls [6] and Brout & Prigogine [7] for phonons (sound waves) in anharmonic crystals. The recent works cast new light on some outstanding problems of wave turbulence. On the one hand, the papers [2, 3, 4] have made an important contribution to the foundations of wave kinetic theory, by showing that the “random phase” and “random phases & amplitudes” properties of initial wave fields are preserved in time by their multimode equations. Although there has been some recent progress [8], we are still far from a rigorous mathematical derivation either of the wave kinetic equation or of any of its predictions. Another important result of [2, 3, 4] is a closed equation for the 1-mode PDF’s of the independent wave amplitudes. These equations can have as solutions 1-mode PDFs that are far from Gaussian, especially when the equations are supplemented with boundary conditions and additional terms to represent strongly nonlinear process, such as wave-breaking, that lie outside the validity of weakly-interacting wave kinetics [3, 9]. These results show promise to explain observations

of intermittency and anomalous scaling in some recent experiments [10, 9] and simulations [11] of gravity-capillary wave turbulence, especially for wavenumber regimes where such intermittency was not previously expected [12]

Unfortunately, the Peierls-Brout-Prigogine-Zaslavskii-Sagdeev (PBPZS) equations for multimode PDF's are not correct for wave kinetics. As we shall show in this paper, these equations have no asymptotic validity in the standard limit of a continuum of weakly interacting, incoherent waves. The works of [2, 3, 4] made errors in estimating the sizes of relevant terms, which we shall discuss in detail. Using the same methods as theirs but with a correct estimation of the terms, we shall obtain a novel set of multimode equations distinct from the PBPZS equations. On the other hand, we shall show using these new multimode equations that the 1-mode PDF equations obtained in [2, 3, 4] *are* correct, under the assumption that initial wave modes have statistically-independent random amplitudes as well as phases. More generally, we shall classify all possible realizable solutions of the new multimode equations, for initial conditions with independent (and uniform) random phases but with no assumption on the statistics of initial wave amplitudes. We find that *the most general solutions correspond to “super-ensembles” of solutions of the wave kinetic equations with random initial conditions for the spectrum or with random forcings.* This represents another possible mechanism for intermittency and non-Gaussian distributions by a “super-turbulence” of chaotic or stochastic solutions of the kinetic equations. As we discuss in detail, existing results on linear stability of Kolmogorov cascade solutions of the wave kinetic equations do not rule out transition to such “super-turbulence.” In fact, this possibility was anticipated in a review of the stability theory by Zakharov et al. (1992), ([1], section 4.2.2), who referred to this possibility as “secondary turbulence.” The results of the present work show that this is the *only* possibility for explaining intermittency and anomalous scaling of wave turbulence strictly within the wave kinetics framework.

The failure of the PBPZS equations will be shown below by specific, detailed calculations, but it can be understood on the basis of simple, general considerations. It will be useful to present such arguments here as a preliminary to the concrete calculations. This will also permit us to give an overview of the important new concepts, tools and results of our paper. We must begin with a brief summary of the main results of [2, 4] for a general Hamiltonian dynamics with 3-wave interactions:

$$\dot{a}_{\mathbf{k}}^{\sigma} = \epsilon \sum_{\sigma_1, \sigma_2} \sum_{\mathbf{k}_1, \mathbf{k}_2} L_{\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2}^{\sigma, \sigma_1, \sigma_2} a_{\mathbf{k}_1}^{\sigma_1} a_{\mathbf{k}_2}^{\sigma_2} e^{i(\sigma_1 \omega(\mathbf{k}_1) + \sigma_2 \omega(\mathbf{k}_2) - \sigma \omega(\mathbf{k}))t} \delta_{\sigma_1 \mathbf{k}_1 + \sigma_2 \mathbf{k}_2, \sigma \mathbf{k}}$$

where the wave triplet nonlinear interaction  $L_{\mathbf{k}, \mathbf{k}_2, \mathbf{k}_3}^{\sigma_1, \sigma_2, \sigma_3} = 3i\sigma_1 H_{\mathbf{k}, \mathbf{k}_2, \mathbf{k}_3}^{-\sigma_1, \sigma_2, \sigma_3}$  in terms of the coefficient of the cubic term in the Hamiltonian and where the summations range over wavenumbers  $\mathbf{k} \in \frac{2\pi}{L} \mathbb{Z}^d$  and a degeneracy index  $\sigma = \pm 1$ , with  $L$  the side-length of a periodic box containing the wave system. (For a systematic discussion of the notations used in this paper, see section 1.1 below.) The analysis of [2, 4] is based on action-angle variables  $(J_{\mathbf{k}}, \varphi_{\mathbf{k}})$  for the linear wave dynamics, defined by  $a_{\mathbf{k}}^{\sigma} = \sqrt{J_{\mathbf{k}}} e^{i\sigma \varphi_{\mathbf{k}}}$ . Note that the standard spectral density is related to the action variables as

$$n(\mathbf{k}) = \lim_{L \rightarrow \infty} \left( \frac{L}{2\pi} \right)^d \langle J_{\mathbf{k}} \rangle$$

in the infinite-volume limit, with total wave action per volume given by  $N = \int d^d k n(\mathbf{k})$ . The authors of [2, 4] exploit a generating functional for action and angle variables, defined by

$$\mathcal{Z}_L(\lambda, \mu) = \left\langle \exp \left( \sum_{\mathbf{k}} i\lambda_{\mathbf{k}} J_{\mathbf{k}} + i\mu_{\mathbf{k}} \varphi_{\mathbf{k}} \right) \right\rangle. \quad (1)$$

In the large-box limit ( $L \rightarrow \infty$ ) followed by the weak nonlinearity limit ( $\epsilon \rightarrow 0$ ), for an initial distribution with independent, uniform phases, this generating function is claimed to satisfy the PBPZS equation in the form

$$\frac{d\mathcal{Z}}{d\tau} = -6\pi i \delta_{\mu, 0} \sum_{\underline{\mathbf{k}}, \underline{\sigma}} |H_{\underline{\mathbf{k}}}^{\underline{\sigma}}|^2 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}) \delta(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}})) (\underline{\sigma} \cdot \lambda_{\underline{\mathbf{k}}}) \partial_{\lambda_{\mathbf{k}_1}} \partial_{\lambda_{\mathbf{k}_2}} \partial_{\lambda_{\mathbf{k}_3}} (\underline{\sigma} \cdot \lambda_{\underline{\mathbf{k}}}) \mathcal{Z}$$

with  $\underline{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$  and  $\underline{\mathbf{k}} = (\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$ , where  $\tau = \epsilon^2 t$  is the nonlinear time. From this equation, [2, 4] derived results on the higher-order fluctuations and statistical distributions of the wave mode amplitudes.

To see that the results, as stated above, cannot be correct, consider the limiting behavior of the generating functional (1) when the field  $a_{\mathbf{k}}$  is assumed to be “RPA”, i.e. to have amplitudes and phases for distinct Fourier modes given

by mutually independent random variables. In that case, the generating functional must factorize for all independent variables, in particular for the amplitudes, as:

$$\mathcal{Z}_L(\lambda, \mu = 0) = \left\langle \exp \left( \sum_{\mathbf{k}} i\lambda_{\mathbf{k}} J_{\mathbf{k}} \right) \right\rangle = \prod_{\mathbf{k}} \left\langle \exp(i\lambda_{\mathbf{k}} J_{\mathbf{k}}) \right\rangle, \quad (2)$$

where each factor is a 1-mode generating function  $\mathcal{Z}^{(1)}(\lambda_{\mathbf{k}}; \mathbf{k}) = \left\langle \exp(i\lambda_{\mathbf{k}} J_{\mathbf{k}}) \right\rangle$ . However, the definition of the spectrum implies that the rescaled variable

$$\tilde{J}_{\mathbf{k}} := (L/2\pi)^d J_{\mathbf{k}}$$

must be  $O(1)$  in the limit as  $L \rightarrow \infty$ . For the 1-mode generating function this gives, employing the cumulant expansion, the result:

$$\mathcal{Z}^{(1)}(\lambda_{\mathbf{k}}; \mathbf{k}) = \left\langle \exp \left( i\lambda_{\mathbf{k}} \left( \frac{2\pi}{L} \right)^d \tilde{J}_{\mathbf{k}} \right) \right\rangle = \exp \left( \sum_{p=1}^{\infty} \frac{i^p}{p!} \left( \frac{2\pi}{L} \right)^{pd} \lambda_{\mathbf{k}}^p \langle \tilde{J}_{\mathbf{k}}^p \rangle^c \right),$$

where  $\langle \tilde{J}_{\mathbf{k}}^p \rangle^c$  is the  $p$ th-order cumulant. Inserting back into the product (2), one obtains

$$\mathcal{Z}_L(\lambda, \mu = 0) = \exp \left( \sum_{p=1}^{\infty} \frac{i^p}{p!} \left( \frac{2\pi}{L} \right)^{pd} \sum_{\mathbf{k}} \lambda_{\mathbf{k}}^p \langle \tilde{J}_{\mathbf{k}}^p \rangle^c \right).$$

Now assume that  $\lambda_{\mathbf{k}} = \lambda(\mathbf{k})$  for some smooth function  $\lambda(\mathbf{k})$ . Since for  $L \rightarrow \infty$

$$\left( \frac{2\pi}{L} \right)^{pd} \sum_{\mathbf{k}} \lambda^p(\mathbf{k}) \langle \tilde{J}_{\mathbf{k}}^p \rangle^c \sim \left( \frac{2\pi}{L} \right)^{(p-1)d} \int d^d k \lambda^p(\mathbf{k}) \langle \tilde{J}_{\mathbf{k}}^p \rangle^c,$$

the contributions of  $p$ th-order cumulants are  $O(L^{-d(p-1)})$ . Only the  $p = 1$  contribution  $n(\mathbf{k}) = \langle \tilde{J}_{\mathbf{k}} \rangle$  survives for very large  $L$

$$\mathcal{Z}_L(\lambda, \mu = 0) = \exp \left( i \int d^d k \lambda(\mathbf{k}) n(\mathbf{k}) + O(L^{-d}) \right) \xrightarrow{L \rightarrow \infty} \exp \left( i \int d^d k \lambda(\mathbf{k}) n(\mathbf{k}) \right).$$

Thus, the generating functional  $\mathcal{Z}_L(\lambda, \mu)$  of an RPA field is completely determined by its spectrum  $n(\mathbf{k})$  in the limit as  $L \rightarrow \infty$  and it then contains no information about higher-order fluctuations. This result, which we have derived here assuming existence of all higher-order cumulants, can be proved under much weaker assumptions (see section 2).

There is a simple interpretation of the above limit in terms of the *empirical spectrum*, defined as

$$\widehat{n}_L(\mathbf{k}) = \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \frac{2\pi}{L} \mathbb{Z}^d} \tilde{J}_{\mathbf{k}_1} \delta^d(\mathbf{k} - \mathbf{k}_1).$$

For an RPA field, this quantity is a sum of a large number of independent variables. It is exactly analogous to the ‘‘Klimontovich density’’ or empirical 1-particle density in the kinetic theory of gases [13]. Unlike the usual spectrum  $n(\mathbf{k})$ , the empirical spectrum is a random variable that incorporates the information about amplitude fluctuations. The amplitude generating function  $\mathcal{Z}_L(\lambda) = \mathcal{Z}_L(\lambda, \mu = 0)$  is just the characteristic functional of the empirical spectrum:

$$\mathcal{Z}_L(\lambda) = \left\langle \exp \left( i \int d^d k \lambda(\mathbf{k}) \widehat{n}_L(\mathbf{k}) \right) \right\rangle. \quad (3)$$

The previous limiting result for  $\mathcal{Z}_L(\lambda)$  is mathematically equivalent to the statement that the empirical spectrum has a *deterministic* limit for  $L \rightarrow \infty$  which is the usual spectrum:

$$\lim_{L \rightarrow \infty} \widehat{n}_L(\mathbf{k}) = n(\mathbf{k}).$$

This limit is a probabilistic law of large numbers for the empirical spectrum. There is a precisely analogous law of large numbers for the empirical 1-particle distribution in the low-density limit for the kinetic theory of gases, as first shown in 1975 by O. Lanford [14, 15].

The above results should hold not only for RPA fields at initial times, but also for the evolved fields in the kinetic regime of wave turbulence, if the RPA property is propagated in time as expected. We shall see that when carried out carefully, with due regard to the scaling of various terms with  $\epsilon$  and  $L$ , the standard analytical methods of wave turbulence yield *not* the PBPZS equation in the limit as first  $L \rightarrow \infty$  then  $\epsilon \rightarrow 0$ , but instead the following equation:

$$\begin{aligned} \dot{\mathcal{Z}}[\lambda, \mu] = & -36i\pi\delta_{\mu,0} \sum_{\sigma=(-1,\sigma_2,\sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}) \delta(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}})) |H_{\underline{\mathbf{k}}}^\sigma|^2 \\ & \times \lambda(\mathbf{k}_1) \left\{ \frac{\delta^2 \mathcal{Z}}{\delta\lambda(\mathbf{k}_2)\delta\lambda(\mathbf{k}_3)} - \sigma_2 \frac{\delta^2 \mathcal{Z}}{\delta\lambda(\mathbf{k}_1)\delta\lambda(\mathbf{k}_3)} - \sigma_3 \frac{\delta^2 \mathcal{Z}}{\delta\lambda(\mathbf{k}_1)\delta\lambda(\mathbf{k}_2)} \right\} \end{aligned}$$

This equation will be justified in detail in this work. Here we note only that it indeed differs from the PBPZS equation. It contains only a subset of the terms in the PBPZS equation, lacking in particular all the terms involving third-order derivatives in  $\lambda$ . The important property of the above corrected equation is that it has solutions of exponential form

$$\mathcal{Z}[\lambda, \mu, \tau] = \exp\left(i \int d^d k \lambda(\mathbf{k}) n(\mathbf{k}, \tau)\right)$$

where  $n(\mathbf{k}, \tau)$  solves the classical wave kinetic equation, *if* the initial conditions  $\mathcal{Z}[\lambda, \mu, \tau = 0]$  are also of this exponential form. As we have discussed above, this exponential form indeed holds if the initial conditions are RPA fields. Note that the PBPZS equation does not have such exponential solutions. The property of preserving exponential solutions implies a law of large-numbers for the empirical spectrum  $\widehat{n}_L(\mathbf{k}, \tau)$  at times  $\tau > 0$  and is equivalent to the ‘‘propagation of chaos’’ by the BBGKY hierarchy in the kinetic limit for gases.

The analogy of kinetic wave turbulence with the kinetic theory of gases is in fact quite close. As we shall show below, the above equation for  $\mathcal{Z}$  is equivalent to a hierarchy of equations for  $M$ -point correlation functions of the empirical spectrum, which is exactly analogous to the ‘‘Boltzmann hierarchy’’ obtained from the BBGKY hierarchy for low-density gases [14, 15]. Just as the Boltzmann hierarchy has factorized solutions for factorized initial conditions, so does the kinetic wave hierarchy for all multi-point spectral correlation functions. The factors in both cases solve the relevant kinetic equation, which satisfies an  $H$ -theorem corresponding to positive entropy production. For both hierarchies the general solutions without assuming factorized initial data are ‘‘super-statistical solutions’’ which represent statistical mixtures of factorized solutions. This was first discussed for the Boltzmann hierarchy by Spohn [16]. For wave turbulence these ‘‘super-statistical solutions’’ of the spectral hierarchy correspond to ensembles of solutions  $n(\mathbf{k}, \tau)$  of the classical wave kinetic equation with random initial conditions  $n_0(\mathbf{k})$ . We shall discuss in this work both the mathematical derivation and the physical relevance of the ‘‘super-statistical solutions’’. These are a possible source of non-Gaussian statistics and intermittency not widely appreciated in the wave turbulence literature.

Previous attempts to calculate intermittency effects within wave kinetic theory have employed the PBPZS equation or its reduced forms for the PDFs of a finite number of modes [3, 4, 17]. If the PBPZS equation is not asymptotically valid in the usual wave kinetic limit (first  $L \rightarrow \infty$ , then  $\epsilon \rightarrow 0$ ), then does this mean that the previously claimed results for the finite-mode PDF’s are also incorrect? The answer is yes, but fortunately only partially. The joint PDF of a set of  $M$  wavenumber modes  $\tilde{J}_{\mathbf{k}_1}, \dots, \tilde{J}_{\mathbf{k}_M}$  is characterized by its Fourier transform, the  $M$ -mode generating function:

$$\mathcal{Z}^{(M)}(\lambda_1, \dots, \lambda_M, \mu = 0; \mathbf{k}_1, \dots, \mathbf{k}_M) = \left\langle \exp\left(\sum_{m=1}^M i\lambda_m \tilde{J}_{\mathbf{k}_m}\right) \right\rangle.$$

We find that these objects do not individually satisfy closed equations but instead satisfy a hierarchy of equations that link  $M$ -mode functions to  $(M + 1)$ - and  $(M + 2)$ -mode functions:

$$\begin{aligned} \dot{\mathcal{Z}}^{(M)} = & -36i\pi \sum_{j=1}^M \sum_{\bar{\sigma}_2, \bar{\sigma}_3} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}_j) \delta(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}}_j)) |H_{\underline{\mathbf{k}}_j}^\sigma|^2 \\ & \left\{ (\lambda_j + \lambda_j^2 \frac{\partial}{\partial \lambda_j}) \frac{\partial^2 \mathcal{Z}^{(M+2)}}{\partial \bar{\lambda}_2 \partial \bar{\lambda}_3} \Big|_{\bar{\lambda}_2 = \bar{\lambda}_3 = 0} - \bar{\sigma}_2 \lambda_j \frac{\partial \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_3 \partial \lambda_j} \Big|_{\bar{\lambda}_3 = 0} - \bar{\sigma}_3 \lambda_j \frac{\partial \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_2 \partial \lambda_j} \Big|_{\bar{\lambda}_2 = 0} \right\}. \end{aligned} \quad (4)$$

with the notations  $\underline{\sigma} = (-1, \bar{\sigma}_2, \bar{\sigma}_3)$  and  $\underline{\mathbf{k}}_j = (\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3)$ . The equations in this hierarchy can also be obtained from the PBPZS equation, by setting all except  $M$  of the  $\lambda$ 's equal to 0 and by retaining only a subset of terms. As in the previous equation for the generating functional, no third-order derivatives in  $\lambda$ 's occur but now there are terms involving  $\lambda_j^2 \partial / \partial \lambda_j$  that were absent in that equation. The above hierarchy for the  $M$ -mode generating functions is equivalent to a hierarchy for the joint PDF's  $\mathcal{P}^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M)$  of  $M$  Fourier modes  $\mathbf{k}_1, \dots, \mathbf{k}_M$ :

$$\dot{\mathcal{P}}^{(M)} + \sum_{j=1}^M \frac{\partial}{\partial s_j} \mathcal{F}_j^{(M)} = 0, \quad (5)$$

where, with the same notations as above,

$$\begin{aligned} \mathcal{F}_j^{(M)} = & -36\pi s_j \sum_{\underline{\sigma} = (-1, \bar{\sigma}_2, \bar{\sigma}_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}_j) \delta(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}}_j)) |H_{\underline{\mathbf{k}}_j}^{\underline{\sigma}}|^2 \\ & \left[ \int d\bar{s}_2 d\bar{s}_3 \frac{\partial \mathcal{P}^{(M+2)}}{\partial s_j}(s_1, \dots, s_M, \bar{s}_2, \bar{s}_3) \bar{s}_2 \bar{s}_3 \right. \\ & \left. + \bar{\sigma}_2 \int d\bar{s}_2 \mathcal{P}^{(M+1)}(s_1, \dots, s_M, \bar{s}_3) \bar{s}_3 + \bar{\sigma}_3 \int d\bar{s}_2 \mathcal{P}^{(M+1)}(s_1, \dots, s_M, \bar{s}_2) \bar{s}_2 \right]. \end{aligned} \quad (6)$$

Previously proposed closed equations for multi-mode equations, e.g. for  $M = 2$  in [17], are not asymptotically valid in the usual kinetic limit for wave turbulence.

On the other hand, the one-mode equation that was derived in [2, 4] for the PDF  $P(s, \tau; \mathbf{k}) = \mathcal{P}^{(1)}(s, \tau; \mathbf{k})$  of a single amplitude  $\tilde{J}_{\mathbf{k}}$ :

$$\frac{\partial}{\partial \tau} P = \frac{\partial}{\partial s} \left[ s \left( \eta_{\mathbf{k}} \frac{\partial P}{\partial s} + \gamma_{\mathbf{k}} P \right) \right] \quad (7)$$

is obtained from the above hierarchy for factorized solutions of the form:

$$\mathcal{P}^{(M)}(s_1, \dots, s_M, \tau; \mathbf{k}_1, \dots, \mathbf{k}_M) = \prod_{m=1}^M P(s_m, \tau; \mathbf{k}_m). \quad (8)$$

The coefficients  $\eta_{\mathbf{k}}$ ,  $\gamma_{\mathbf{k}}$  that appear also in the kinetic equation  $\dot{n}_{\mathbf{k}} = \eta_{\mathbf{k}} - \gamma_{\mathbf{k}} n_{\mathbf{k}}$  are obtained from the condition  $n_{\mathbf{k}} = \int ds s P(s; \mathbf{k})$  and the standard wavenumber integrals over products of  $n$ . The one-mode equation is thus a nonlinear Fokker-Planck equation in the sense of McKean [18]. Factorized solutions of the form (8) are obtained for factorized initial data, corresponding to RPA fields. The hierarchy thus preserves in this sense the statistical independence of amplitudes. Just as for the spectral hierarchy, this factorization property of solutions implies a law of large numbers for *empirical 1-mode PDFs*

$$\widehat{P}_L(s, \Delta, t) = \frac{1}{N_L(\Delta)} \sum_{\mathbf{k} \in \frac{2\pi}{L} \mathbb{Z}^d \cap \Delta} \delta(s - \tilde{J}_{\mathbf{k}})$$

with  $\Delta \subset \mathbb{R}^d$  a subset of continuous wavenumbers and with  $N_L(\Delta)$  the number of elements in  $\frac{2\pi}{L} \mathbb{Z}^d \cap \Delta$ . Also as for the spectral hierarchy, the general solutions of the PDF hierarchy are “super-statistical solutions” that correspond to ensembles of solutions of the 1-mode equation (7) with random initial conditions  $P_0(s; \mathbf{k})$ . As we shall discuss, such ensembles may be physically relevant to explain non-Gaussian statistics in weakly nonlinear wave systems if the solutions of (7) themselves become random, either through fluctuating driving forces or internal chaos/turbulence.

The detailed contents of this work are as follows. The introduction concludes with a brief section 1.1 summarizing our model assumptions and notations. The next section 2 discusses the probabilistic properties of “RP” and “RPA” fields. The principal results of the paper are in section 3, which derives the multimode equations and discusses their basic properties. This is done first for the spectral generating functional and correlation functions (section 3.1) and next for the PDF generating functions and correlation functions (section 3.2). Many technical details are provided in Appendix A and Appendix B. Finally, we discuss in section 4 possible kinetic theory mechanisms of intermittency in wave turbulence. The conclusion section 5 summarizes our results.

### 1.1. Model and Notations

We summarize here briefly the notations employed in our work. We consider a system consisting of a complex wavefield  $u(\mathbf{x}, t)$  in a  $d$ -dimensional periodic cube with side  $L$ . As in [2, 4], we assume that there is a maximum wavenumber  $k_{\max}$ , to avoid ultraviolet divergences. This can be achieved by a lattice regularization with spacing  $a = L/M$ , for some large integer  $M$ , so that  $k_{\max} = \pi/a$ . The location variable  $\mathbf{x}$  then ranges over the physical space

$$\Lambda_L = a\mathbb{Z}_M^d,$$

with the usual notation  $\mathbb{Z}_M$  for the field of integers modulo  $M$ . This space has volume  $V = L^d$ . The dual space of wavenumbers is

$$\Lambda_L^* = \frac{2\pi}{L}\mathbb{Z}_M^d$$

with  $k_{\min} = 2\pi/L$ . The total number of modes is  $N = M^d$ , so that spatial volume  $V = Na^d$ . We use the following index notation

$$u^\sigma(\mathbf{x}) = \begin{cases} u(\mathbf{x}) & \sigma = +1 \\ u^*(\mathbf{x}) & \sigma = -1 \end{cases}$$

for  $u$  and its complex-conjugate  $u^*$ , following [2]. Likewise, we adopt their convention for (discrete) Fourier transform

$$A^\sigma(\mathbf{k}) = \frac{1}{N} \sum_{\mathbf{x} \in \Lambda_L} u^\sigma(\mathbf{x}, t) \exp(-i\sigma\mathbf{k} \cdot \mathbf{x})$$

so that  $A^+(\mathbf{k})$  and  $A^-(\mathbf{k})$  are complex conjugates. Notice that this quantity converges to the continuous Fourier transform  $\frac{1}{L^d} \int_{[0, L]^d} d^d x u^\sigma(\mathbf{x}, t) \exp(-i\sigma\mathbf{k} \cdot \mathbf{x})$  in the limit  $a \rightarrow 0$ . The discrete inverse transform is

$$u^\sigma(\mathbf{x}) = \sum_{\mathbf{k} \in \Lambda_L^*} A^\sigma(\mathbf{k}) \exp(i\sigma\mathbf{k} \cdot \mathbf{x}).$$

The dynamics is assumed to be canonical Hamiltonian with a cubic Hamiltonian density (energy per volume) describing 3-wave interactions:

$$H = H_0 + \delta H = \sum_{\sigma, \mathbf{k}} \omega(\mathbf{k}) A_{\mathbf{k}}^\sigma A_{\mathbf{k}}^{-\sigma} + \epsilon \sum_{\underline{\sigma}, \underline{\mathbf{k}}} H_{\underline{\mathbf{k}}}^{\underline{\sigma}} A_{\mathbf{k}_1}^{\sigma_1} A_{\mathbf{k}_2}^{\sigma_2} A_{\mathbf{k}_3}^{\sigma_3} \delta_{\underline{\sigma}, \underline{\mathbf{k}}, \mathbf{0}}.$$

As in [2], we denote triplets of variables by the notation  $\underline{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ ,  $\underline{\mathbf{k}} = (\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$  and define the dot product  $\underline{\sigma} \cdot \underline{\mathbf{k}} = \sigma_1 \mathbf{k}_1 + \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3$ . The Kronecker delta function in the triplet interaction term enforces the symmetry of the dynamics under the group of (discrete) space translations. The interaction coefficients  $H_{\underline{\mathbf{k}}}^{\underline{\sigma}}$  are furthermore assumed to satisfy

$$H_{\underline{\mathbf{k}}}^{\underline{\sigma}*} = H_{\underline{\mathbf{k}}}^{-\underline{\sigma}} \quad \text{and} \quad H_{\underline{\mathbf{k}}}^{\underline{\sigma}} = H_{\pi(\underline{\mathbf{k}})}^{\pi(\underline{\sigma})} \quad (9)$$

for any permutation  $\pi \in S_3$ . The first condition guarantees the reality of the Hamiltonian and the second can always be assumed without loss of generality. The Hamiltonian equations of motion for the Fourier coefficients are

$$\frac{\partial A_{\mathbf{k}}^\sigma}{\partial t} = i\sigma \frac{\partial H}{\partial A_{\mathbf{k}}^{-\sigma}} = i\sigma \omega(\mathbf{k}) A_{\mathbf{k}}^\sigma + \epsilon \sum_{\sigma_1, \sigma_2} \sum_{\mathbf{k}_1, \mathbf{k}_2} L_{\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2}^{\sigma, \sigma_1, \sigma_2} A_{\mathbf{k}_1}^{\sigma_1} A_{\mathbf{k}_2}^{\sigma_2} \delta_{\sigma_1 \mathbf{k}_1 + \sigma_2 \mathbf{k}_2, \sigma \mathbf{k}}. \quad (10)$$

where  $L_{\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2}^{\sigma, \sigma_1, \sigma_2} = 3i\sigma H_{\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2}^{-\sigma, \sigma_1, \sigma_2}$ . It is useful to introduce action-angle variables for the linear dynamics,  $J_{\mathbf{k}} = |A_{\mathbf{k}}^\sigma|^2$  and  $\varphi_{\mathbf{k}} = \sigma \arg(A_{\mathbf{k}}^\sigma)$ , so that  $A_{\mathbf{k}}^\sigma = \sqrt{J_{\mathbf{k}}} \psi_{\mathbf{k}}^\sigma$ , where  $\psi_{\mathbf{k}} = \exp(i\varphi_{\mathbf{k}})$ . In these variables, the Liouville measure  $\mu$  conserved by the Hamiltonian flow can be written variously as

$$d\mu = \prod_{\mathbf{k}} dq_{\mathbf{k}} dp_{\mathbf{k}} = \prod_{\mathbf{k}} \frac{1}{i} dA_{\mathbf{k}}^+ dA_{\mathbf{k}}^- = \prod_{\mathbf{k}} dJ_{\mathbf{k}} d\varphi_{\mathbf{k}}$$

where the canonical momenta and coordinates are given by real and imaginary parts of  $A_{\mathbf{k}}^\sigma = \frac{1}{\sqrt{2}}(p_{\mathbf{k}} + i\sigma q_{\mathbf{k}})$ .

As usual in wave kinetic theory, we introduce the ‘‘interaction representation’’

$$a_{\mathbf{k}}^\sigma = A_{\mathbf{k}}^\sigma e^{-i\sigma\omega(\mathbf{k})t} \quad (11)$$

which removes the rapid wave oscillations. The Liouville measure is invariant under this transformation because of the rotation-invariance of the Haar measure on  $\psi \in S^1$ . We shall often use the shorthand notation  $a_1 = a_{\mathbf{k}_1}^{\sigma_1}$  together with  $\sum_1 := \sum_{\sigma_1, \mathbf{k}_1}$  and

$$\begin{aligned} \omega_{23}^1 &:= -\sigma_1\omega(\mathbf{k}_1) + \sigma_2\omega(\mathbf{k}_2) + \sigma_3\omega(\mathbf{k}_3), \\ \delta_{23}^1 &:= \delta_{\sigma_1\mathbf{k}_1, \sigma_2\mathbf{k}_2 + \sigma_3\mathbf{k}_3}. \end{aligned}$$

The dynamical equation of motion can then be written succinctly as

$$\dot{a}_1 = \epsilon \sum_{2,3} L_{123} a_2 a_3 e^{i\omega_{23}^1 t} \delta_{23}^1. \quad (12)$$

## 2. Fields with Random Phases and Amplitudes

It is often assumed in derivations of wave kinetic equations that initial fields have Fourier coefficients with random (and statistically independent) phases and amplitudes. Furthermore, these properties are expected to be preserved in time, in some suitable sense, in the wave-kinetic limit. See the discussions in [2] and, particularly, [3, 4]. It is therefore important to review here the definitions and properties of such ‘‘RPA’’ fields<sup>1</sup>. Our discussion shall be largely complementary to [4], emphasizing the probabilistic characteristics of such fields.

Consider then  $N$  complex-valued random variables  $a_{\mathbf{k}}$ ,  $\mathbf{k} \in \Lambda_L^*$  taken to be the Fourier coefficients of a random space field:

$$u(\mathbf{x}) = \sum_{\mathbf{k} \in \Lambda_L^*} a_{\mathbf{k}} \exp(i\mathbf{k} \cdot \mathbf{x}).$$

Here  $a_{\mathbf{k}}$  corresponds to  $a_{\mathbf{k}}^+ = A_{\mathbf{k}}^+$  in the previous section (no distinction need be made between the two at time  $t = 0$ ). It will be very important in what follows to work with normalized variables

$$\tilde{a}_{\mathbf{k}} = \left(\frac{L}{2\pi}\right)^{d/2} a_{\mathbf{k}}$$

which are assumed to remain finite in the large-box limit  $L \rightarrow \infty$ . As we shall remind the reader below, this normalization is required so that the spectrum of the random field is well defined in that limit. It is convenient to write the complex variables in polar coordinates  $a_{\mathbf{k}} = \sqrt{J_{\mathbf{k}}} e^{i\varphi_{\mathbf{k}}}$  (action-angle variables) with the normalized action  $\tilde{J}_{\mathbf{k}} = \left(\frac{L}{2\pi}\right)^d J_{\mathbf{k}}$ . We use the notations  $s_{\mathbf{k}}$  and  $\xi_{\mathbf{k}}$  for possible values of the random variables  $\tilde{J}_{\mathbf{k}} \in \mathbb{R}^+$  and  $\psi_{\mathbf{k}} = e^{i\varphi_{\mathbf{k}}} \in S^1$ . The Liouville measure of the previous section becomes

$$d\mu(s, \xi) = \prod_{\mathbf{k} \in \Lambda_L^*} ds_{\mathbf{k}} \frac{|d\xi_{\mathbf{k}}|}{2\pi} \quad (13)$$

suitably normalized. We define the  $N$ -mode joint probability density function  $\mathcal{P}^{(N)}(s, \xi)$  with respect to Liouville measure, such that the average of the random variable  $f(\tilde{J}, \psi)$  is given by

$$\langle f(\tilde{J}, \psi) \rangle = \int d\mu(s, \xi) \mathcal{P}^{(N)}(s, \xi) f(s, \xi)$$

where the integral is over  $(s, \xi)$  in the product space  $(\mathbb{R}^+)^N \times (S^1)^N$ .

<sup>1</sup>More accurate acronyms would be ‘‘IUP’’ for ‘‘independent uniform phases’’ rather than ‘‘RP’’ and ‘‘IPA’’ for ‘‘independent phases & amplitudes’’ rather than ‘‘RPA’’. However, we shall stick here to the abbreviations already employed in the literature.

Following [2, 4] we define  $u(\mathbf{x})$  to be a *random-phase field* (RP) if  $\psi_{\mathbf{k}} = e^{i\varphi_{\mathbf{k}}}$  for all  $\mathbf{k} \in \Lambda_L^*$  are independent and identically distributed (i.i.d.) random variables, uniformly distributed over the unit circle  $S^1$  in the complex plane. In terms of the joint PDF, this is equivalent to the condition that

$$\mathcal{P}^{(N)}(s, \xi) = \mathcal{P}^{(N)}(s)$$

independent of  $\xi$ . It is easy to see that an RP  $u(\mathbf{x})$  is a homogeneous random field on  $\Lambda_L$ , statistically invariant under space-translations by the finite group  $a\mathbb{Z}_M^d$ . This follows from the rotation invariance of the Haar measures  $|d\xi_{\mathbf{k}}|/2\pi$  on the phase variables and by the representation of the translations  $u(\mathbf{x}) \rightarrow u(\mathbf{x} + \mathbf{r})$  for  $\mathbf{r} \in a\mathbb{Z}_M^d$  as phase-rotations:  $\psi_{\mathbf{k}} \rightarrow \psi_{\mathbf{k}} e^{i\mathbf{k}\cdot\mathbf{r}}$ . In the limit  $L \rightarrow \infty$  the field  $u_L(\mathbf{x})$  defined with appropriately chosen  $\tilde{J}_{\mathbf{k},L}$  will converge to a homogeneous random field  $u(\mathbf{x})$  invariant under translations by  $a\mathbb{Z}^d$ . The standard definition of the spectrum  $n(\mathbf{k}) = \lim_{L \rightarrow \infty} (L/2\pi)^d \langle |a_{\mathbf{k},L}|^2 \rangle$  implies that one must choose

$$\lim_{L \rightarrow \infty} \langle \tilde{J}_{\mathbf{k},L} \rangle = n(\mathbf{k}), \quad (14)$$

for  $\mathbf{k} \in \Lambda^* = [-k_{\max}, +k_{\max}]^d$ , where  $\mathbf{k}_L = \frac{\mathbf{k}L}{2\pi} \pmod{M} \cdot \frac{2\pi}{L} \in \Lambda_L^*$  converges to  $\mathbf{k}$  as  $L = aM \rightarrow \infty$  (for fixed  $a$ ). In this case,  $u_L(\mathbf{x})$  converges in distribution as  $L \rightarrow \infty$  to a homogeneous field  $u(\mathbf{x})$  with spectrum  $n(\mathbf{k})$ .

Again following [3, 4], we define  $u(\mathbf{x})$  to be a *random-phase and amplitude field* (RPA) if  $u(\mathbf{x})$  is RP and if also  $\tilde{J}_{\mathbf{k}}$  are mutually independent random variables for all  $\mathbf{k} \in \Lambda_L^*$ . This is equivalent to the factorization of the  $N$ -mode PDF into a product of 1-mode PDF's:

$$\mathcal{P}^{(N)}(s) = \prod_{\mathbf{k} \in \Lambda_k^*} P(s_{\mathbf{k}}; \mathbf{k}).$$

All homogeneous Gaussian random fields are RPA. Conversely, for any sequence of RPA fields satisfying condition (14) the field  $u_L(\mathbf{x})$  converges in distribution to the homogeneous Gaussian field with mean zero and spectrum  $n(\mathbf{k})$  as  $L \rightarrow \infty$ . See [19] for related rigorous results. Here we note only that

$$u_L(\mathbf{x}) = \left( \frac{2\pi}{L} \right)^{d/2} \sum_{\mathbf{k} \in \Lambda_L^*} \sqrt{\tilde{J}_{\mathbf{k},L}} \exp(i\mathbf{k} \cdot \mathbf{x} + i\varphi_{\mathbf{k}})$$

is a sum of  $N$  independent variables scaled by  $1/\sqrt{N}$ . It is important to emphasize that the Fourier coefficients  $\tilde{a}_{\mathbf{k},L}$  remain far from Gaussian in this limit. In physical space also there are non-vanishing cumulants for large but finite  $L$ . A curious property of RPA fields for finite  $L$  is the slow decay of their cumulants in certain directions. For example, it is easy to calculate the 4th-order cumulant as

$$\begin{aligned} & \langle u(\mathbf{x}_1)u(\mathbf{x}_2)u^*(\mathbf{x}_3)u^*(\mathbf{x}_4) \rangle - [\langle u(\mathbf{x}_1)u^*(\mathbf{x}_3) \rangle \langle u(\mathbf{x}_2)u^*(\mathbf{x}_4) \rangle + (1 \leftrightarrow 2)] \\ &= \left( \frac{2\pi}{L} \right)^{2d} \sum_{\mathbf{k} \in \Lambda_L^*} [\langle \tilde{J}_{\mathbf{k}}^2 \rangle - 2\langle \tilde{J}_{\mathbf{k}} \rangle^2] e^{i\mathbf{k} \cdot (\mathbf{x}_1 + \mathbf{x}_2 - \mathbf{x}_3 - \mathbf{x}_4)} \\ &\sim \left( \frac{2\pi}{L} \right)^d \int d^d k [\langle \tilde{J}_{\mathbf{k}}^2 \rangle - 2\langle \tilde{J}_{\mathbf{k}} \rangle^2] e^{i\mathbf{k} \cdot (\mathbf{x}_1 + \mathbf{x}_2 - \mathbf{x}_3 - \mathbf{x}_4)} \end{aligned}$$

asymptotically as  $L \rightarrow \infty$ . As expected, this goes to zero as  $O(L^{-d})$  for  $L \rightarrow \infty$ , but for finite  $L$  the cumulant is constant on the hyperplane  $\mathbf{x}_1 + \mathbf{x}_2 = \mathbf{x}_3 + \mathbf{x}_4$ , even when the pair of points  $\mathbf{x}_1, \mathbf{x}_3$  are separated very far from the pair  $\mathbf{x}_2, \mathbf{x}_4$ , for example. This is in contrast to some more traditional derivations of wave kinetic equations, which assume rapid spatial decay of higher-order cumulants [20].

A most important result for RPA fields is that the *empirical spectrum*

$$\widehat{n}_L(\mathbf{k}) = \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \Lambda_L^*} \tilde{J}_{\mathbf{k}_1,L} \delta^d(\mathbf{k} - \mathbf{k}_1), \quad \mathbf{k} \in \Lambda^*$$

converges under the condition (14) to the deterministic spectrum  $n(\mathbf{k})$  with probability going to 1 in the limit  $L \rightarrow \infty$  (weak law of large numbers). It is worth sketching the simple proof. Note for any continuous function  $\lambda$  on  $\Lambda^*$  that

$$\int_{\Lambda^*} d^d k \lambda(\mathbf{k}) \widehat{n}_L(\mathbf{k}) = \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \Lambda_L^*} \lambda(\mathbf{k}_1) \tilde{J}_{\mathbf{k}_1,L}.$$



Crucially,

$$\begin{aligned} & \left\langle \left| \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \Lambda_L^*} \lambda(\mathbf{k}_1) \tilde{J}_{\mathbf{k}_1} - \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \Lambda_L^*} \lambda(\mathbf{k}_1) n(\mathbf{k}_1) \right|^2 \right\rangle \\ &= \left( \frac{2\pi}{L} \right)^{2d} \sum_{\mathbf{k}_1 \in \Lambda_L^*} \lambda^2(\mathbf{k}_1) \langle |\tilde{J}_{\mathbf{k}_1} - n(\mathbf{k}_1)|^2 \rangle \\ &\sim \left( \frac{2\pi}{L} \right)^d \int d^d k \lambda^2(\mathbf{k}) \langle |\tilde{J}_{\mathbf{k}} - n(\mathbf{k})|^2 \rangle = O(L^{-d}) \end{aligned}$$

under the modest assumption that  $\int d^d k \langle |\tilde{J}_{\mathbf{k}} - n(\mathbf{k})|^2 \rangle < \infty$ . Since also  $\lim_{L \rightarrow \infty} \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \Lambda_L^*} \lambda(\mathbf{k}_1) n(\mathbf{k}_1) = \int d^d k \lambda(\mathbf{k}) n(\mathbf{k})$ , the  $L^2$ -convergence follows

$$\lim_{L \rightarrow \infty} \left\langle \left| \int d^d k \lambda(\mathbf{k}) \widehat{n}_L(\mathbf{k}) - \int d^d k \lambda(\mathbf{k}) n(\mathbf{k}) \right|^2 \right\rangle = 0$$

and thus  $\int d^d k \lambda(\mathbf{k}) \widehat{n}_L(\mathbf{k})$  converges in probability to  $\int d^d k \lambda(\mathbf{k}) n(\mathbf{k})$  for every bounded, continuous  $\lambda$ .<sup>2</sup> This is sufficient to infer that the amplitude generating function defined in the introduction satisfies<sup>3</sup>

$$\lim_{L \rightarrow \infty} \mathcal{Z}_L(\lambda) = \exp \left( i \int d^d k \lambda(\mathbf{k}) n(\mathbf{k}) \right) \quad (15)$$

with  $n(\mathbf{k})$  the deterministic spectrum. The law of large numbers (LLN) derived above means that for RPA fields the empirical spectrum  $\widehat{n}_L(\mathbf{k})$  coincides with  $n(\mathbf{k})$  at large  $L$  for almost every realization of the random phases and amplitudes, not just after averaging over these variables.

Notice that for the above result one does not actually need the full independence assumption in RPA, but only uncorrelated amplitudes:

$$\langle \tilde{J}_{\mathbf{k}_1} \tilde{J}_{\mathbf{k}_2} \rangle = \langle \tilde{J}_{\mathbf{k}_1} \rangle \langle \tilde{J}_{\mathbf{k}_2} \rangle, \quad \mathbf{k}_1 \neq \mathbf{k}_2.$$

An even weaker and more general condition can be stated in terms of the *M-mode correlation functions* of the empirical spectrum, defined as:

$$\mathcal{N}_L^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M) = \langle \widehat{n}_L(\mathbf{k}_1) \cdots \widehat{n}_L(\mathbf{k}_M) \rangle.$$

Note that (14) implies that  $\lim_{L \rightarrow \infty} \mathcal{N}_L^{(1)}(\mathbf{k}) = n(\mathbf{k})$ . A careful examination of the previous proof shows that, in order to obtain the LLN for the empirical spectrum, it suffices that

$$\lim_{L \rightarrow \infty} [\mathcal{N}_L^{(2)}(\mathbf{k}_1, \mathbf{k}_2) - \mathcal{N}_L^{(1)}(\mathbf{k}_1) \mathcal{N}_L^{(1)}(\mathbf{k}_2)] = 0. \quad (16)$$

<sup>2</sup>Technically, the convergence is in the weak- $\star$  topology for  $\widehat{n}_L(\mathbf{k}), n(\mathbf{k})$  as bounded, positive measures on  $\Lambda^*$ . More physically, the function  $\lambda(\mathbf{k})$  may be taken to be a smooth kernel  $G_\delta(\mathbf{k} - \mathbf{k}_0)$  of width  $\delta$  centered around a particular wavenumber  $\mathbf{k}_0$ . The coarse-grained spectrum  $\widehat{n}_{L,\delta}(\mathbf{k}_0) = \int d^d k G_\delta(\mathbf{k} - \mathbf{k}_0) \widehat{n}_L(\mathbf{k})$  can then be interpreted as the result of measuring the spectrum at  $\mathbf{k}_0$  with a finite resolution  $\delta$  in wavenumber. The weak- $\star$  topology implies the convergence of such coarse-grained spectra in the limit  $L \rightarrow \infty$ , when arbitrarily many wavenumbers lie within distance  $\delta$  of  $\mathbf{k}_0$ .

<sup>3</sup>This is another standard result in probability theory, but, for completeness, we here recall the proof. Let  $X_L = \int_{\Lambda^*} d^d k \lambda(\mathbf{k}) \widehat{n}_L(\mathbf{k})$  and  $x = \int_{\Lambda^*} d^d k \lambda(\mathbf{k}) n(\mathbf{k})$ . For  $\epsilon > 0$ , write

$$\left| \langle e^{iX_L} - e^{ix} \rangle \right| \leq \langle |e^{i(X_L-x)} - 1| : |X_L - x| < \epsilon \rangle + \langle |e^{i(X_L-x)} - 1| : |X_L - x| > \epsilon \rangle$$

For any  $\delta > 0$ , one can choose  $\epsilon$  so that  $|e^{iz} - 1| < \delta$  when  $|z| < \epsilon$ . The first term is thus bounded by  $\delta$ . By Chebyshev inequality the second term is bounded as

$$\langle |e^{i(X_L-x)} - 1| : |X_L - x| > \epsilon \rangle \leq 2P(|X_L - x| > \epsilon) \leq \frac{2}{\epsilon^2} \langle |X_L - x|^2 \rangle \rightarrow 0$$

for fixed  $\epsilon$  as  $L \rightarrow \infty$ . Thus,

$$\limsup_{L \rightarrow \infty} \left| \langle e^{iX_L} - e^{ix} \rangle \right| \leq \delta.$$

Since  $\delta$  is arbitrary, it follows that  $\lim_{L \rightarrow \infty} \langle e^{iX_L} \rangle = e^{ix}$ .

This condition is the analogue of the *Stosszahlansatz* invoked by Boltzmann to derive his kinetic equation. Under this assumption, all of the  $M$ -th order correlations that exist will factorize in the large-box limit:

$$\lim_{L \rightarrow \infty} \mathcal{N}_L^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M) = \prod_{m=1}^M n(\mathbf{k}_m). \quad (17)$$

These observations go back to Lanford in his derivation of the Boltzmann equation [14, 15]. The results that we shall present below suggest that properties (14) and (16) for the wave field at the initial time, together with the RP property, are sufficient for the validity of the wave kinetic equation and a LLN for the empirical spectrum at positive times.

RPA fields whose Fourier amplitudes possess the full independence property satisfy an even stronger law of large numbers for the *empirical 1-mode PDF*, which is defined as

$$\widehat{P}_L(s; \mathbf{k}) = \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}_1 \in \Lambda_L^*} \delta(s - \tilde{J}_{\mathbf{k}_1}) \delta^d(\mathbf{k} - \mathbf{k}_1). \quad (18)$$

Assume that the limiting random variables  $\tilde{J}_{\mathbf{k}} = \lim_{L \rightarrow \infty} \tilde{J}_{\mathbf{k},L}$  of an RPA field exist and have PDF's  $P(s; \mathbf{k})$  which are continuous in  $\mathbf{k}$ . Then an analogue of the previous argument implies that the random functions  $\widehat{P}_L(s; \mathbf{k})$  converge to  $P(s; \mathbf{k})$  with probability approaching 1 as  $L \rightarrow \infty$ . This implies the previous LLN for the spectrum, since  $\widehat{n}_L(\mathbf{k}) = \int_0^\infty ds s \widehat{P}_L(s; \mathbf{k})$  and  $n(\mathbf{k}) = \int_0^\infty ds s P(s; \mathbf{k})$ . Although the “empirical PDF” defined in (18) is mathematically very convenient, it is not a PDF for finite  $L$ . It is therefore more intuitive to use an alternative definition

$$\widehat{P}_L(s; \Delta) = \frac{1}{N_L(\Delta)} \sum_{\mathbf{k} \in \Lambda_L^* \cap \Delta} \delta(s - \tilde{J}_{\mathbf{k}}), \quad (19)$$

for any open set  $\Delta \subset \Lambda^*$  and with  $N_L(\Delta)$  the number of elements in  $\Lambda_L^* \cap \Delta$ . This quantity is nearly the same as  $\frac{1}{|\Delta|} \int_{\Delta} d^d k \widehat{P}_L(s; \mathbf{k})$  for large  $L$  but it has the advantage that it defines a probability measure in  $s$  for each fixed  $\Delta$  and  $L$ . Definition (19) also has a simple intuitive meaning, since it represents the instantaneous distribution of amplitudes of the large number of Fourier modes that reside in the set  $\Delta$  for large box-size  $L$ . Under the same assumptions as above, it follows with probability going to 1 that

$$\lim_{L \rightarrow \infty} \widehat{P}_L(s; \Delta) = \frac{1}{|\Delta|} \int_{\Delta} d^d k P(s; \mathbf{k}) \equiv P(s; \Delta).$$

As before, the essential property that is required for the above results to hold is a factorization property of *multi-mode PDF*'s, defined for  $\mathbf{k}_1, \dots, \mathbf{k}_M \in \Lambda^*$  by

$$\mathcal{P}_L^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M) = \langle \delta(s_1 - \tilde{J}_{\mathbf{k}_{1,L}}) \cdots \delta(s_M - \tilde{J}_{\mathbf{k}_{M,L}}) \rangle.$$

The factorization property of the 2-mode PDF's for all pairs of distinct  $\mathbf{k}_1, \mathbf{k}_2 \in \Lambda^*$

$$\lim_{L \rightarrow \infty} [\mathcal{P}_L^{(2)}(s_1, s_2; \mathbf{k}_1, \mathbf{k}_2) - \mathcal{P}_L^{(1)}(s_1; \mathbf{k}_1) \mathcal{P}_L^{(1)}(s_2; \mathbf{k}_2)] = 0 \quad (20)$$

suffices<sup>4</sup> to derive the LLN for the empirical PDF and also the factorization of the multi-mode PDF's

$$\lim_{L \rightarrow \infty} \mathcal{P}_L^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M) = \prod_{m=1}^M P(s_m; \mathbf{k}_m)$$

<sup>4</sup>Since the proof is so similar to the one given previously, we give just a few details here. Because of our assumptions on  $\tilde{J}_{\mathbf{k}} = \lim_{L \rightarrow \infty} \tilde{J}_{\mathbf{k},L}$  it is easy to check that the average of the empirical PDF converges as  $L \rightarrow \infty$ :

$$\langle \widehat{P}_L(s; \mathbf{k}) \rangle = \left( \frac{2\pi}{L} \right)^d \sum_{\mathbf{k}' \in \Lambda_L^*} \mathcal{P}_L^{(1)}(s; \mathbf{k}') \delta^d(\mathbf{k}' - \mathbf{k}) \rightarrow P(s; \mathbf{k}).$$

Therefore, in order to prove that  $\widehat{P}_L(s; \mathbf{k})$  converges in probability to  $P(s; \mathbf{k})$ , with convergence in the weak- $\star$  topology on regular Borel measures,

for all integers  $M > 2$  and distinct  $\mathbf{k}_1, \dots, \mathbf{k}_M \in \Lambda^*$ . The ‘‘asymptotic independence’’ property (20) is considerably weaker than strict RPA, permitting statistical dependence between Fourier modes at finite  $L$ . We shall discuss natural ‘‘microcanonical measures’’ in section 3.2.2 which satisfy the condition (20) but not the more stringent RPA condition.

In the following section we shall show that all of the above properties of RPA (or generalized RPA) initial conditions are preserved in time by the limiting kinetic hierarchies of wave turbulence.

### 3. Multi-Mode Hierarchy Equations

In this section we present our formal derivations of the multimode equations of wave kinetics and a discussion of their main properties. Our derivations are quite similar to those of Choi et al. [4], but our model is somewhat more general in appearance. The major difference between our analysis and theirs, however, lies in a correct accounting of the size of the various terms in the limits as  $L \rightarrow \infty$  and  $\epsilon \rightarrow 0$ .

We shall focus on the multimode generating function for finite box-size  $L$ , defined as in [4] by

$$\mathcal{Z}_L[\lambda, \mu, T] = \left\langle \exp \left( \sum_{\mathbf{k} \in \Lambda_L^*} \lambda_{\mathbf{k}} J_{\mathbf{k}}(T) \right) \prod_{\mathbf{k} \in \Lambda^*} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}}(T) \right\rangle, \quad (21)$$

where  $\lambda_{\mathbf{k}} \in \mathbb{R}$  and  $\mu_{\mathbf{k}} \in \mathbb{Z}$  for all  $\mathbf{k} \in \Lambda_L^*$ . The time  $T$  is free for the moment but will later be chosen to be a time intermediate between the wave-period and the nonlinear time-scale. This generating function is calculated perturbatively in the nonlinearity parameter  $\epsilon$  at finite  $L$ , by expanding the solution of the dynamical equation

$$\dot{a}_1 = \epsilon \sum_{2,3} L_{123} a_2 a_3 e^{i\omega_{23}t} \delta_{23}^1 \quad (22)$$

into a power series

$$a_1(T) = a_1(0) + \epsilon a_1^{(1)}(T) + \epsilon^2 a_1^{(2)}(T) + \mathcal{O}(\epsilon^3). \quad (23)$$

A straightforward calculation gives

$$a_1^{(0)} = a_1(0), \quad (24)$$

$$a_1^{(1)} = \sum_{2,3} L_{123} a_2^{(0)} a_3^{(0)} \Delta_T(\omega_{23}^1) \delta_{23}^1, \quad (25)$$

$$a_1^{(2)} = \sum_{2345} L_{123} L_{245} a_3^{(0)} a_4^{(0)} a_5^{(0)} E_T(\omega_{345}^1, \omega_{23}^1) \delta_{23}^1 \delta_{45}^2 + (2 \leftrightarrow 3) \quad (26)$$

We employ here the standard definitions [20]:

$$\Delta_T(x) = \int_0^T \exp(ixt) dt, \quad E_T(x, y) = \int_0^T \Delta_t(x - y) \exp(iyt) dt.$$

it is enough to show that

$$\lim_{L \rightarrow \infty} \left\langle \left| \int ds \int_{\Lambda^*} d^d k \varphi(s, \mathbf{k}) \widehat{P}_L(s; \mathbf{k}) - \int ds \int_{\Lambda^*} d^d k \varphi(s, \mathbf{k}) \langle \widehat{P}_L(s; \mathbf{k}) \rangle \right|^2 \right\rangle = 0$$

for any continuous function  $\varphi(s, \mathbf{k})$  on  $\mathbb{R}^+ \times \Lambda^*$  which vanishes as  $s \rightarrow \infty$ . Now a direct calculation of the above average shows that it equals

$$\left( \frac{2\pi}{L} \right)^{2d} \sum_{\mathbf{k}'_1, \mathbf{k}'_2 \in \Lambda_L^*} \int ds_1 \int ds_2 \varphi(s_1, \mathbf{k}'_1) \varphi(s_2, \mathbf{k}'_2) [P_L^{(2)}(s_1, s_2; \mathbf{k}'_1, \mathbf{k}'_2) - P_L^{(1)}(s_1; \mathbf{k}'_1) P_L^{(1)}(s_2; \mathbf{k}'_2)].$$

The bracketed expression for  $\mathbf{k}'_1 = \mathbf{k}'_2$  is found to be

$$P_L^{(2)}(s_1, s_2; \mathbf{k}'_1, \mathbf{k}'_1) - P_L^{(1)}(s_1; \mathbf{k}'_1) P_L^{(1)}(s_2; \mathbf{k}'_1) = \delta(s_1 - s_2) P_L^{(1)}(s_1; \mathbf{k}'_1) - P_L^{(1)}(s_1; \mathbf{k}'_1) P_L^{(1)}(s_2; \mathbf{k}'_1)$$

so that its contribution to the average vanishes as  $\sim \left( \frac{2\pi}{L} \right)^d \int_{\Lambda^*} d^d k \left\langle |\varphi(\tilde{\mathbf{k}}, \mathbf{k}) - \langle \varphi(\tilde{\mathbf{k}}, \mathbf{k}) \rangle|^2 \right\rangle$  for  $L \rightarrow \infty$ . Thus, to prove convergence, it must only be shown that the bracket term for  $\mathbf{k}'_1 \neq \mathbf{k}'_2$  vanishes as  $L \rightarrow \infty$ . This is obviously true for RPA fields, when the bracket term for  $\mathbf{k}'_1 \neq \mathbf{k}'_2$  is zero!

The terms in this perturbative solution of the equations of motion can be represented by a version of the Wyld diagram expansion [21]. In this technique the various contributions are represented by tree diagrams, as illustrated in Figs. 1-3 for the zeroth-, first- and second-order terms. In our conventions, similar to those of [4], a solid line labeled with an integer  $j$  represents a factor  $a_j^{(0)}$ , whereas a dashed line indicates the absence of such a factor. An arrow is added to a solid line to indicate  $\sigma_j = +1$  (“source”) when the arrow is pointed away from  $j$  and  $\sigma_j = -1$  (“sink”) when the arrow is pointed toward  $j$ . The triple vertex labelled 1, 2, 3 represents a factor  $L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{\sigma_1, \sigma_2, \sigma_3} e^{\omega_{23}^1 t} \delta_{23}^1$  with  $\sigma_1 = +1$  when the arrow points out of the vertex and  $\sigma_1 = -1$  when the arrow points into the vertex. The times at each vertex are ordered causally, with the latest times at the “root” of the tree, here labelled by 1. When integrations are performed over all times from 0 to  $T$  consistent with this ordering, then the various contributions to the perturbative solution result.



Figure 1: Zeroth-order terms  $a_1^{(0)+}$  and  $a_1^{(0)-}$ .




Figure 2: First-order terms  $a_1^{(1)+}$  and  $a_1^{(1)-}$ .




Figure 3: Second-order term  $a_1^{(2)+}$ .

The generating function is obtained perturbatively by substituting (23) and expanding to obtain

$$\mathcal{Z}_L[\lambda, \mu, T] = \mathcal{X}_L\{\lambda, \mu, T\} + \mathcal{X}_L^*\{\lambda, -\mu, T\} \quad (27)$$

with

$$\mathcal{X}_L\{\lambda, \mu, T\} = \mathcal{X}_L\{\lambda, \mu, 0\} + \left\langle \prod_{\mathbf{k} \in \Lambda^*} e^{\lambda \mathbf{k} J_{\mathbf{k}}^{(0)}} [\epsilon \mathcal{J}_1 + \epsilon^2 (\mathcal{J}_2 + \mathcal{J}_3 + \mathcal{J}_4 + \mathcal{J}_5)] \right\rangle_J + \mathcal{O}(\epsilon^3), \quad (28)$$

where, as in [4],

$$\mathcal{J}_1 = \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}} \sum_1 \left( \lambda_1 + \frac{\mu_1}{2J_1^{(0)}} \right) a_1^{(1)} a_1^{(0)*} \right\rangle_{\psi}, \quad (29)$$

$$\mathcal{J}_2 = \frac{1}{2} \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}} \sum_1 \left( \lambda_1 + \lambda_1^2 J_1^{(0)} - \frac{\mu_1^2}{4J_1^{(0)}} \right) |a_1^{(1)}|^2 \right\rangle_{\psi}, \quad (30)$$

$$\mathcal{J}_3 = \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}} \sum_1 \left( \lambda_1 + \frac{\mu_1}{2J_1^{(0)}} \right) a_1^{(2)} a_1^{(0)*} \right\rangle_{\psi}, \quad (31)$$

$$\mathcal{J}_4 = \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}} \sum_1 \left( \frac{1}{2} \lambda_1^2 + \frac{\mu_1}{4J_1^{(0)2}} \left( \frac{\mu_1}{2} - 1 \right) + \frac{\lambda_1 \mu_1}{2J_1^{(0)}} \right) (a_1^{(1)} a_1^{(0)*})^2 \right\rangle_{\psi}, \quad (32)$$

$$\begin{aligned} \mathcal{J}_5 = \frac{1}{2} \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}} \sum_{1 \neq 2} \left( \lambda_1 \lambda_2 (a_1^{(1)} a_1^{(0)*} + a_1^{(1)*} a_1^{(0)}) a_2^{(1)} a_2^{(0)*} \right. \right. \\ \left. \left. + \left( \lambda_1 + \frac{\mu_1}{4J_1^{(0)}} \right) \frac{\mu_2}{J_2^{(0)}} (a_2^{(1)} a_2^{(0)*} - a_2^{(1)*} a_2^{(0)}) a_1^{(1)} a_1^{(0)*} \right) \right\rangle_{\psi}. \quad (33) \end{aligned}$$

The various contributions before averaging over phases can be represented by diagrams, combining the tree diagrams for each of the factors by joining the trees with the same “root” indices. Each of the integer labels indicates an index to be summed over independently (except for the constraints imposed by delta-functions at the vertices). We illustrate this representation in Figs.4-5 below for the first two contributions:




Figure 4: First-order term  $\sum_1 \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) a_1^{(1)} a_1^{(0)*}$ .




Figure 5: Second-order term  $\sum_1 \left( \lambda_1 + \lambda_1^2 J_1 - \frac{\mu_1^2}{4J_1} \right) |a_1^{(1)}|^2$ .

The only contributions which survive the average over phases must have all phases summing to zero before averaging. This means that every  $a^{(0)}$  factor must either pair with another factor  $a^{(0)}$  so that their phases sum to zero or belong to a set of  $a^{(0)}$ 's that pair with a  $\psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}}$  factor so that the sum of all their phases is zero. The first we call an “internal coupling”, represented graphically by a solid line connecting the paired indices  $i, j$  which contributes a factor  $\delta_{\sigma_i + \sigma_j, 0} \delta_{\mathbf{k}_i, \mathbf{k}_j}$  after phase averaging. The second we call an “external coupling”, represented by joining all of

the solid lines for indices  $i_1, i_2, \dots, i_p$  to a blob  $\bullet$  labelled  $a$  that represents the phase  $\psi_{\mathbf{k}_a}^{(0)\mu_{\mathbf{k}_a}}$  which contributes a factor  $\delta_{\sigma_{i_1} + \dots + \sigma_{i_p} + \mu_a, 0} \prod_{j=1}^p \delta_{\mathbf{k}_j, \mathbf{k}_a}$  after phase averaging. This graphical representation is essentially the same as that employed in [4]. We illustrate the representation in Figs.6 and 7 below for the terms in  $\mathcal{J}_1$  and  $\mathcal{J}_2$  which turn out to give the leading-order contributions to those quantities as  $L \rightarrow \infty$ .




Figure 6: Contributions to  $\mathcal{J}_1$ .




Figure 7: Leading contribution to  $\mathcal{J}_2$ .

Note that solid lines connected to external blobs have their wavenumbers “pinned” at the wavenumbers of those blobs, so that those wavenumbers are no longer summed over. For simplicity we often omit the labels of the blobs, since those play no important role other than the “pinning” described above.

### 3.1. Spectral Hierarchy

We now consider the first of the possible limits of  $\mathcal{Z}_L[\lambda, \mu]$  as  $L \rightarrow \infty$  involving *all* of the  $N$  modes. This leads to a set of equations for the spectral generating function and the spectral correlation functions. We first sketch the derivation of these equations, with more details in Appendix A, and then analyze their basic properties.

#### 3.1.1. Derivation

The crucial observation which leads to our results differing from [4] is that one must keep  $\tilde{J}_{\mathbf{k}} = O(1)$ , *not*  $J_{\mathbf{k}} = O(1)$ , in order to have a finite spectrum in the limit  $L \rightarrow \infty$ . Thus, we take in the generating function of Choi et al.

$$J_{\mathbf{k}} = \left(\frac{2\pi}{L}\right)^d \tilde{J}_{\mathbf{k}}, \quad \lambda_{\mathbf{k}} = i\lambda(\mathbf{k}),$$

where  $\lambda(\mathbf{k})$  is a smooth test function and, as before,  $\mu_{\mathbf{k}}$  are integers. This leads to

$$\mathcal{Z}_L[\lambda, \mu] = \left\langle \exp \left( i \sum_{\mathbf{k} \in \Lambda_L^*} \left(\frac{2\pi}{L}\right)^d \lambda(\mathbf{k}) \tilde{J}_{\mathbf{k}} \right) \prod_{\mathbf{k} \in \Lambda^*} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \right\rangle, \quad (34)$$

We consider the large- $L$  asymptotics of the various terms in the perturbation expansion of this quantity, employing the following standard substitutions:

$$\left(\frac{2\pi}{L}\right)^d \sum_{\mathbf{k}} \Rightarrow \int d^d k, \quad \left(\frac{L}{2\pi}\right)^d \delta_{\mathbf{k},\mathbf{k}'} \Rightarrow \delta^d(\mathbf{k} - \mathbf{k}'), \quad \left(\frac{L}{2\pi}\right)^d \frac{\partial}{\partial \lambda_{\mathbf{k}}} \Rightarrow \frac{\delta}{\delta \lambda(\mathbf{k})} \quad (35)$$

From now on, we suppress superscript (0) when there is no confusion.

*Calculation of  $\mathcal{J}_1$ :* This quantity is represented by the graphs in Fig.6, or analytically:

$$\begin{aligned} \mathcal{J}_1 &= \sum_{1 \neq 2 \neq 3} L_{123} \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) \sqrt{J_1 J_2 J_3} \delta_{\mu_1,1} \delta_{\mu_2+\sigma_2,0} \delta_{\mu_3+\sigma_3,0} \Delta(\omega_{23}^1) \delta_{23}^1 \\ &\quad + \sum_{1 \neq 2} L_{122} \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) \sqrt{J_1 J_2} \delta_{\mu_1,1} \delta_{\mu_2+2\sigma_2,0} \Delta(\omega_{22}^1) \delta_{22}^1 \end{aligned}$$

Note that  $\sigma_1 = +1$  in coefficient  $L_{123}$ . Taking into account the wavenumber delta functions, there are two summations in the first term and one in the second. However, as noted by Choi et al. (2005), these sums contain only a couple of non-zero terms, and then only for special choices of the  $\mu$ 's. In their terminology, the sums are ‘‘pinned’’ by these choices of  $\mu$ . Making the substitutions  $J_i = (2\pi/L)^{d/2} \tilde{J}_i$  and  $\lambda_1 = i\lambda(\mathbf{k}_1)$  into the above gives the leading contribution

$$\begin{aligned} \mathcal{J}_1 &= \left(\frac{2\pi}{L}\right)^{d/2} \sum_{1 \neq 2 \neq 3} L_{123} \frac{1}{2} \mu_1 \sqrt{\frac{\tilde{J}_2 \tilde{J}_3}{\tilde{J}_1}} \delta_{\mu_1,1} \delta_{\mu_2,-\sigma_2} \delta_{\mu_3,-\sigma_3} \Delta(\omega_{23}^1) \delta_{23}^1 \\ &\quad + \left(\frac{2\pi}{L}\right)^{d/2} \sum_{1 \neq 2} L_{122} \frac{1}{2} \mu_1 \frac{\tilde{J}_2}{\sqrt{\tilde{J}_1}} \delta_{\mu_1,1} \delta_{\mu_2,-2\sigma_2} \Delta(\omega_{22}^1) \delta_{22}^1 \propto L^{-d/2} \end{aligned}$$

As we shall see, this term first-order in  $\epsilon$  gives a subleading correction in the limit  $L$  large but finite, larger than many of the terms that Choi et al. retained in their evolution formula for  $L \rightarrow \infty$ .

*Calculation of  $\mathcal{J}_2$ :* Averaging over phases, the leading terms are contained in the contribution from Fig.7 and a similar contribution with 2 and 3 interchanged. Because there are no external couplings, all  $\mu$ 's must vanish. Because of the internal couplings, there are no sums over the wavenumbers 4,5 which the same as wavenumbers 2,3. The final contribution to the generating function is:

$$\begin{aligned} \langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} \mathcal{J}_2} \rangle_J &= \delta_{\mu,0} \sum_{1,2,3} \langle (\lambda_1 + \lambda_1^2 J_1) J_2 J_3 e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \rangle_J |L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3}|^2 |\Delta(\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1)|^2 \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} \\ &= 9 \delta_{\mu,0} \sum_{1,2,3} \left( \lambda_1 + \lambda_1^2 \frac{\partial}{\partial \lambda_1} \right) \frac{\partial^2 \mathcal{Z}}{\partial \lambda_2 \partial \lambda_3} \times |H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3}|^2 |\Delta(\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1)|^2 \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} \end{aligned} \quad (36)$$

Papers [2, 4] retain *both* of the terms in the first factor in (36), i.e. both  $\lambda_1$  and  $\lambda_1^2 J_1$ . However, with the proper scaling,  $\lambda_1 = O(1)$  while  $\lambda_1^2 J_1 = \left(\frac{2\pi}{L}\right)^d \lambda_1 \tilde{J}_1 = O(L^{-d})$ . Thus, the second term is even smaller than the contribution from  $\mathcal{J}_1$  in the limit as  $L \rightarrow \infty$  and should be neglected. This can be seen also substituting  $J_i = \left(\frac{2\pi}{L}\right)^{d/2} \tilde{J}_i$  and  $\lambda_1 = i\lambda(\mathbf{k}_1)$  and taking the limit  $L \rightarrow \infty$  using (35):

$$\begin{aligned} \langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} \mathcal{J}_2} \rangle_J &\sim -9i \delta_{\mu,0} \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \left( \lambda(\mathbf{k}_1) + \left(\frac{2\pi}{L}\right)^d \lambda(\mathbf{k}_1) \frac{\delta}{\delta \lambda(\mathbf{k}_1)} \right) \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_2) \delta \lambda(\mathbf{k}_3)} \\ &\quad \times |H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3}|^2 |\Delta(\underline{\sigma} \cdot \underline{\omega})|^2 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}). \end{aligned} \quad (37)$$

The only surviving term in the limit as  $L \rightarrow \infty$  is the one proportional to  $\lambda(\mathbf{k}_1)$ . Verifying our initial estimation, the term proportional to  $\lambda(\mathbf{k}_1) \delta / \delta \lambda(\mathbf{k}_1)$  is  $O(L^{-d})$  and vanishes in the limit.

A similar analysis may be carried through for the remaining contributions from  $\mathcal{J}_3, \mathcal{J}_4$  and  $\mathcal{J}_5$ . Detailed discussion of all the terms is given in Appendix A, where it is shown that only  $\mathcal{J}_2, \mathcal{J}_3, \mathcal{J}_5$  give  $O(1)$  contributions in the large-box limit  $L \rightarrow \infty$ :

$$\mathcal{X}(T) - \mathcal{X}(0) \sim -9i \delta_{\mu,0} \epsilon^2 \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}) |\Delta_T(\underline{\sigma} \cdot \underline{\omega}(\underline{\mathbf{k}}))|^2$$

$$\begin{aligned}
& \times \lambda(\mathbf{k}_1) |H_{\mathbf{k}}^\sigma|^2 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_2) \delta \lambda(\mathbf{k}_3)} \\
& + \epsilon^2 \left[ 18i \delta_{\mu,0} \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \sigma_2 \int_{(\mathbb{T}^d)^3} d^d k_1 d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) E_T(0, \underline{\sigma} \cdot \omega(\mathbf{k})) \right. \\
& \quad \times \lambda(\mathbf{k}_1) |H_{\mathbf{k}}^\sigma|^2 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_3)} \\
& \quad - 9 \sum_1 \delta_{\mu_1,1} \delta_{\mu-1,1} \prod_{\mathbf{k} \neq \mathbf{k}_1, -\mathbf{k}_1} \delta_{\mu_{\mathbf{k}},0} \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \sigma_2 \int d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \\
& \quad \left. E_T(\omega(\mathbf{k}_1) + \omega(-\mathbf{k}_1), \underline{\sigma} \cdot \omega(\mathbf{k})) H_{\mathbf{k}}^\sigma H_{-\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \left\langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \sqrt{\frac{\tilde{J}_{-1}}{\tilde{J}_1} \tilde{J}_3} \right\rangle_J + (2 \leftrightarrow 3) \right] \\
& - \frac{9}{2} \epsilon^2 \sum_1 \delta_{\mu_1,1} \delta_{\mu-1,1} \prod_{\mathbf{k} \neq \pm \mathbf{k}_1} \delta_{\mu_{\mathbf{k}},0} \sum_{\underline{\sigma}=(-, \sigma_2, \sigma_3)} \int d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \\
& \quad \left[ \Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k})) \Delta_T(-\underline{\sigma} \cdot \omega(\mathbf{k}')) \right] H_{\mathbf{k}}^\sigma H_{-\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \left\langle \frac{\tilde{J}_2 \tilde{J}_3}{\sqrt{\tilde{J}_1 \tilde{J}_{-1}}} e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \right\rangle_J. \tag{38}
\end{aligned}$$

The first term is that already found for  $\mathcal{J}_2$ , the second bracketed term is from  $\mathcal{J}_3$ , and the final term is from  $\mathcal{J}_5$ . Note that the expressions proportional to  $\delta_{\mu_1,1} \delta_{\mu-1,1}$  were missed in [2, 4] (although they are larger than many extra terms that those authors retained in their final equations which actually vanish as  $L \rightarrow \infty$ !) The terms proportional to  $\delta_{\mu_1,1} \delta_{\mu-1,1}$  indeed do not appear in the final equations, not because of the large- $L$  limit but because they are nonsecular.

We therefore consider the limit of weak nonlinearity. The limit is achieved by choosing the time variable  $T$  in the expansion (23) to lie between the wave period and nonlinear timescale  $O(\epsilon^{-2})$  and by then taking  $\epsilon \rightarrow 0$ . For this purpose we use the following standard asymptotic relations for  $T \rightarrow \infty$  [20]:

$$\begin{aligned}
\Delta_T(x) & \sim \tilde{\Delta}(x) = \pi \delta(x) + iP \left( \frac{1}{x} \right), \quad E_T(x; y) \sim \Delta_T(x) \Delta_T(y) \sim \tilde{\Delta}(x) \tilde{\Delta}(y), \\
|\Delta_T(x)|^2 & \sim 2\pi T \delta(x) + 2P \left( \frac{1}{x} \right) \frac{\partial}{\partial x}, \quad E_T(x; 0) \sim \tilde{\Delta}(x) \left( T - i \frac{\partial}{\partial x} \right), \tag{39}
\end{aligned}$$

The terms multiplied by  $\delta_{\mu,0}$  contain secular contributions proportional to  $T$ , while the terms proportional to  $\delta_{\mu_1,1} \delta_{\mu-1,1}$  are nonsecular. We now use (27) to calculate  $\mathcal{Z}(T) - \mathcal{Z}(0)$ , with the observation that it is changed by our replacement  $\lambda_{\mathbf{k}} = i\lambda(\mathbf{k})$  into

$$\mathcal{Z}[\lambda, \mu, T] = \mathcal{X}[\lambda, \mu, T] + \mathcal{X}^*[-\lambda, -\mu, T]. \tag{40}$$

Finally, replacing  $(\mathcal{Z}(T) - \mathcal{Z}(0))/T$  by  $\dot{\mathcal{Z}}$  and using time variable  $\tau = \epsilon^2 t$ , one obtains

$$\begin{aligned}
\frac{d}{d\tau} \mathcal{Z}[\lambda, \mu, \tau] & = -36i\pi \delta_{\mu,0} \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \delta(\underline{\sigma} \cdot \omega(\mathbf{k})) \lambda(\mathbf{k}_1) |H_{\mathbf{k}}^\sigma|^2 \\
& \quad \left\{ \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_2) \delta \lambda(\mathbf{k}_3)} - \sigma_2 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_3)} - \sigma_3 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_2)} \right\}. \tag{41}
\end{aligned}$$

Equation (41) is the main result of this section.

### 3.1.2. Properties

In this section we shall discuss the most basic properties of equation (41) derived in the previous section. As pointed out in [4], the factor  $\delta_{\mu,0}$  implies that the RP property of the initial conditions is preserved in time. Therefore, without loss of generality, we need only consider the characteristic functional for amplitudes, or  $\mathcal{Z}[\lambda, \tau] \equiv \mathcal{Z}[\lambda, \mu = 0, \tau]$ . Its evolution equation is

$$\dot{\mathcal{Z}}[\lambda, \tau] = -36i\pi \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \delta(\underline{\sigma} \cdot \omega(\mathbf{k})) \lambda(\mathbf{k}_1) |H_{\mathbf{k}}^\sigma|^2$$



$$\times \left\{ \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_2) \delta \lambda(\mathbf{k}_3)} - \sigma_2 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_3)} - \sigma_3 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_2)} \right\} \quad (42)$$

Hereafter we consider only this amplitude characteristic functional. Eq.(42) implies a hierarchy of evolution equations for the  $M$ -mode spectral correlation functions defined in section 2, in the wave kinetic limit:

$$\mathcal{N}^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \tau) = \lim_{\epsilon \rightarrow 0} \lim_{L \rightarrow \infty} \mathcal{N}_{L, \epsilon}^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \epsilon^{-2} \tau).$$

The hierarchy is easiest to derive by using the relation (3) between  $\mathcal{Z}[\lambda, \tau]$  and the empirical spectrum, which implies that

$$\mathcal{N}^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \tau) = (-i)^M \frac{\delta^M \mathcal{Z}[\lambda, \tau]}{\delta \lambda(\mathbf{k}_1) \cdots \delta \lambda(\mathbf{k}_M)} \Big|_{\lambda=0}.$$

By taking  $M$  functional derivatives of (42) and setting  $\lambda \equiv 0$ , one derives for each integer  $M = 1, 2, 3, \dots$  the following equation:

$$\begin{aligned} \dot{\mathcal{N}}^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \tau) &= 36\pi \sum_{j=1}^M \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}_j) \delta(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}}_j)) \lambda(\underline{\mathbf{k}}_j) |H_{\underline{\mathbf{k}}_j}^{\underline{\sigma}}|^2 \\ & \left[ \mathcal{N}^{(M+2)}(\mathbf{k}_1, \dots, \mathbf{k}_{j-1}, \mathbf{k}_{j+1}, \dots, \mathbf{k}_M, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3, \tau) - \sigma_2 \mathcal{N}^{(M+1)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \bar{\mathbf{k}}_3, \tau) - \sigma_3 \mathcal{N}^{(M+1)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \bar{\mathbf{k}}_2, \tau) \right]. \end{aligned} \quad (43)$$

which couples the  $M$ th-order correlation functions to the  $(M+1)$ st. We shall refer to the above collection of equations for all  $M = 1, 2, 3, \dots$  as the *spectral hierarchy* of kinetic wave turbulence. It is exactly analogous to the ‘‘Boltzmann hierarchy’’ derived by Lanford from the BBGKY hierarchy in the low-density limit [14, 15]. If the spectral correlation functions satisfy bounds on their growth for large orders  $M$  that allow them to uniquely characterize the distribution of the empirical spectrum, then the spectral hierarchy (43) is not only a consequence of the equation (42) but is in fact equivalent to that equation.

An extremely important property of the equations (42) or (43) is that they possess certain exact solutions. In particular, if the initial functional  $\mathcal{Z}[\lambda, 0]$  is of exponential form (15), as follows for an initial RP field with uncorrelated amplitudes, then an exact solution of (42) is

$$\mathcal{Z}[\lambda, \tau] = \exp\left(i \int d^d \mathbf{k} \lambda(\mathbf{k}) n(\mathbf{k}, \tau)\right), \quad (44)$$

where  $n(\mathbf{k}, \tau)$  satisfies the standard wave kinetic equation

$$\begin{aligned} \dot{n}(\mathbf{k}, \tau) &= 36\pi \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d k_2 d^d k_3 |H_{\underline{\mathbf{k}}}^{\underline{\sigma}}|^2 \delta(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}})) \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}) \\ & \times \left\{ n(\mathbf{k}_2, \tau) n(\mathbf{k}_3, \tau) - \sigma_1 n(\mathbf{k}, \tau) n(\mathbf{k}_3, \tau) - \sigma_3 n(\mathbf{k}, \tau) n(\mathbf{k}_2, \tau) \right\}. \end{aligned} \quad (45)$$

with initial condition  $n(\mathbf{k}, 0) = n(\mathbf{k})$ . This may be checked by direct substitution of (44) into (42). Equivalently, with factorized  $M$ th-order correlation functions (17) as initial data, there is a solution of the spectral hierarchy equations (43) also of factorized form:

$$\mathcal{N}^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \tau) = \prod_{m=1}^M n(\mathbf{k}_m, \tau). \quad (46)$$

Note that such factorized solutions have only power-law growth for large orders  $M$ , so that they uniquely characterize the exponential characteristic functional (44). If it can be proved that solutions of the dynamical equations (42) or (43) are unique for classes of initial data that include the forms (15) and (17), then the equations we have derived imply that spectral correlation functions initially factorized will remain so for  $\tau > 0$ . In this sense, therefore, the property of uncorrelated wave amplitudes is preserved in time by our equations. This is an exact analogue of the ‘‘propagation of chaos’’ property for the Boltzmann hierarchy, which implies that the *Stosszahlansatz* is propagated in time [14, 15]. The uniqueness of solutions of the linear functional equations (42) and (43) is an interesting mathematical problem which we shall address in future work.

The results above have an important implication. As follows from our discussion in section 2, the conditions (44) or (46) imply a law of large numbers for the empirical spectrum at positive times. That is, with probability going to 1 in the kinetic limit (first  $L \rightarrow \infty$ , then  $\epsilon \rightarrow 0$ ), it follows that

$$\widehat{n}_L(\mathbf{k}, \epsilon^{-2}\tau) \simeq n(\mathbf{k}, \tau), \quad \tau > 0$$

where  $n(\mathbf{k}, \tau)$  is the solution of the wave kinetic equation (45). The interesting implication for laboratory and numerical experiments is that the wave kinetic equations will be valid for *typical* initial amplitudes and phases chosen from an RPA ensemble and not just for the spectrum averaged over the RPA ensemble. That is, the empirical spectrum is “self-averaging.” This is the exact analogue of the law of large numbers derived by Lanford for the empirical 1-particle distribution (Klimontovich density) in the low density limit for gases [14, 15].

In order to emphasize the close formal analogy of wave kinetic theory with the kinetic theory of gases, it is worthwhile to make here a few remarks about the role of entropy in both. It is well-known that the wave kinetic equation (45) satisfies an “H-theorem” for the entropy defined, with Boltzmann’s constant  $k_B$ , by

$$S[n] = k_B \int_{\Lambda^*} d^d k \ln n(\mathbf{k}). \quad (47)$$

That is,  $dS/d\tau \geq 0$  for general solutions of (45) and  $dS/d\tau = 0$  for the thermal equilibrium solutions  $n_{\text{eq}}(\mathbf{k}) = k_B T / \omega(\mathbf{k})$  at absolute temperature  $T$ , which maximize the entropy (47) for fixed energy  $E[n] = \int_{\Lambda^*} d^d k \omega(\mathbf{k}) n(\mathbf{k})$ . See [1], section 2.2.2. This is the exact analogue of the  $H$ -theorem originally derived by Boltzmann for his kinetic equation. As pointed out more recently by Spohn [22], the entropy (47) also follows from Boltzmann’s prescription that  $S = k_B \log W$  [23], where  $W$  is the Liouville measure of the set of microstates  $(\tilde{J}, \psi)$  consistent with the given “macrostate” defined by the prescribed spectrum  $n(\mathbf{k})$ . More precisely, let  $\Delta_i, i = 1, \dots, P$  be a partition of  $\Lambda^*$  and let  $\Gamma_{n, \Delta, \eta, L}$  be the set of microstates  $\{(\tilde{J}_{\mathbf{k}}, \psi_{\mathbf{k}}), \mathbf{k} \in \Lambda_L^*\}$  such that  $|\widehat{n}_L(\Delta_i) - n(\Delta_i)| < \eta$ , for  $i = 1, \dots, P$  where

$$\widehat{n}_L(\Delta_i) = \frac{1}{|\Delta_i|} \int_{\Delta_i} d^d k \widehat{n}_L(\mathbf{k}), \quad n(\Delta_i) = \frac{1}{|\Delta_i|} \int_{\Delta_i} d^d k n(\mathbf{k}).$$

Then<sup>5</sup>

$$\lim_{\eta \rightarrow 0} \lim_{L \rightarrow \infty} \left( \frac{2\pi}{L} \right)^d \ln |\Gamma_{n, \Delta, \eta, L}| = \sum_{i=1}^P |\Delta_i| (\ln n(\Delta_i) + 1).$$

The result for each individual cell is the same as the microcanonical entropy  $s(e)$  of a system of non-interacting harmonic oscillators or alternatively as the entropy of an ideal gas in a periodic box, thinking of  $\tilde{J}_{\mathbf{k}}$  as kinetic energies and  $\varphi_{\mathbf{k}}$  as positions of the particles. Furthermore, defining  $|\Delta| = \max_i |\Delta_i|$ , the partition may be refined by taking  $|\Delta| \rightarrow 0$ , so that

$$\lim_{|\Delta| \rightarrow 0} \lim_{\eta \rightarrow 0} \lim_{L \rightarrow \infty} \left( \frac{2\pi}{L} \right)^d \ln |\Gamma_{n, \Delta, \eta, L}| = \int_{\Lambda^*} d^d k (\ln n(\mathbf{k}) + 1).$$

The result agrees, up to constants, with (47). The above argument introduces the “microcanonical measure” obtained by restricting Liouville measure to the set  $\Gamma_{n, \Delta, \eta, L}$  and normalizing by  $|\Gamma_{n, \Delta, \eta, L}|$  to yield a probability measure. By adapting the arguments of [15] it follows that this “microcanonical measure” is a natural example which satisfies asymptotic factorization (17) but *not* the RPA property.

The previous results may appear somewhat disappointing. Equations for the generating function  $\mathcal{Z}[\lambda, \tau]$  like our (42) have been proposed mainly in the hope of developing theories of intermittency of wave turbulence and of higher-order statistics of the wave-amplitudes. It might be concluded from the discussion above that the correct equation for  $\mathcal{Z}[\lambda, \tau]$ —i.e. our equation (42)—is equivalent to the wave kinetic equation (45) and has no more physical content. However, this is not entirely correct. To clarify this point, we now classify all of the realizable solutions of (42).

<sup>5</sup> Note that  $\Gamma_{n, \Delta, \eta, L}$  is a Cartesian product set of the form  $\otimes_{i=1}^P \{(\tilde{J}_{\mathbf{k}}, \psi_{\mathbf{k}}), \mathbf{k} \in \Lambda_L^* \cap \Delta_i : |\widehat{n}_L(\Delta_i) - n(\Delta_i)| < \eta\}$  and for each cell  $\Delta_i$  of the partition,  $\widehat{n}_L(\Delta_i) \approx \frac{1}{2N_L(\Delta_i)} \sum_{\mathbf{k} \in \Delta_i \cap \Lambda_L^*} (\tilde{p}_{\mathbf{k}}^2 + \tilde{q}_{\mathbf{k}}^2)$ . The stated result then follows using the formula  $\frac{\pi^{D/2}}{(D/2)!} R^D$  for the volume of a ball in dimension  $D = 2N(\Delta_i)$  of radius  $R = (nD)^{1/2}$ , dividing by  $(2\pi)^{D/2}$  [eq.(13)], and applying Stirling’s approximation for the factorial  $(D/2)!$  as  $D \rightarrow \infty$ .

That is, we characterize the solutions of the equation (42) for the most general possible initial conditions that can be physically attained in the kinetic limit, assuming RP but *no* form of RPA. For this purpose, note that the characteristic functional  $\mathcal{Z}_L(\lambda, 0)$  defined in (3) is a positive-definite functional, i.e.

$$\sum_{i,j=1}^n c_i c_j^* \mathcal{Z}_L(\lambda_i - \lambda_j, 0) \geq 0$$

for any set of  $n$  fields  $\lambda_1, \dots, \lambda_n$  and complex numbers  $c_1, \dots, c_n$ . It is also normalized so that  $\mathcal{Z}_L(\lambda = 0, 0) = 1$ . Since these properties are preserved under pointwise limits, the physical initial condition

$$\mathcal{Z}(\lambda, 0) = \lim_{L \rightarrow \infty} \mathcal{Z}_L(\lambda, 0)$$

for our eq.(42) also satisfies them. We must assume that the above limit exists, if eq.(42) is to have any validity at all, and we shall assume furthermore that the limiting  $\mathcal{Z}(\lambda, 0)$  is a continuous functional of the  $\lambda$  fields<sup>6</sup>. By the Bochner-Minlos theorem, it therefore has the form

$$\mathcal{Z}(\lambda, 0) = \int d\rho(n_0) \exp\left(i \int d^d \mathbf{k} \lambda(\mathbf{k}) n_0(\mathbf{k})\right)$$

for some probability measure  $\rho$ . E.g. see [24]. That is, the initial condition  $\mathcal{Z}(\lambda, 0)$  is a statistical superposition of exponential initial conditions of the form (15) that arise from RPA fields. Since equation (42) is linear in  $\mathcal{Z}$ , a solution for such superposed initial data is

$$\mathcal{Z}(\lambda, \tau) = \int d\rho(n_0) \exp\left(i \int d^d \mathbf{k} \lambda(\mathbf{k}) n(\mathbf{k}, \tau)\right)$$

where  $n(\mathbf{k}, \tau)$  solves the kinetic equation with initial condition  $n_0(\mathbf{k})$ . This is the only solution if the uniqueness of solutions holds for (42). The conclusion of this argument is that *the most general, statistically realizable solutions<sup>7</sup> of eq.(42) correspond to ensembles of solutions of the wave kinetic equation with random initial conditions  $n_0(\mathbf{k})$ .*

The above argument is a formal analogue of a rigorous result of Spohn for the Boltzmann hierarchy in the kinetic theory of gases [16]. With appropriate technical assumptions, his argument can be carried over to our spectral hierarchy (43), with the conclusion that its general realizable solutions are statistical superpositions of factorized solutions, that is,

$$\mathcal{N}^{(M)}(\mathbf{k}_1, \dots, \mathbf{k}_M, \tau) = \int d\rho(n_0) \prod_{m=1}^M n(\mathbf{k}_m, \tau),$$

where again  $n(\mathbf{k}, \tau)$  solves the kinetic equation with initial condition  $n_0(\mathbf{k})$ . We shall refer to such solutions of the spectral hierarchy (43) or of the equivalent equation (42) as “super-statistical solutions”, since they correspond to random ensembles of solutions of the spectral closure equation. As we shall discuss at length in section 4, such “super-statistical solutions” offer a possibility to explain intermittency and non-Gaussian statistics previously little discussed in the wave turbulence literature. However, we shall first consider the alternative approach based on closed equations for the PDF’s of the wave amplitudes.

### 3.2. PDF Hierarchy

We now consider a second possible limit involving only a fixed number  $M$  of modes  $a_{\mathbf{k}_m}$ ,  $m = 1, \dots, M$  as the total number  $N \rightarrow \infty$ . As before, one must keep  $\tilde{J}_{\mathbf{k}} = O(1)$  for all modes. We thus define the joint characteristic function:

$$\mathcal{Z}_L^{(M)}(\lambda_1, \dots, \lambda_M, \mu_1, \dots, \mu_M, T; \mathbf{k}_1, \dots, \mathbf{k}_M) = \left\langle \exp\left(i \sum_{m=1}^M \lambda_m \tilde{J}_{\mathbf{k}_m}(T)\right) \prod_{m=1}^M \psi_{\mathbf{k}_m}^{\mu_m}(T)\right\rangle, \quad (48)$$

<sup>6</sup>We shall not attempt to identify here the precise topologies on the spaces of  $\lambda$ ’s and  $n$ ’s that would permit a rigorous formulation and proof of our results.

<sup>7</sup>It should be noted that there may be non-realizable solutions of eq.(42) that do not have this form. In general, statistical moment equations may have “parasitic” solutions that do not correspond to realizable solutions of the underlying statistical problem. For an example of this phenomenon in the Kraichnan passive scalar model, see [25].

This is the generating function (21) of [4] with

$$\lambda_{\mathbf{k}_m} = i \left( \frac{L}{2\pi} \right)^d \lambda_m, \quad J_{\mathbf{k}_m} = \left( \frac{2\pi}{L} \right)^d \tilde{J}_{\mathbf{k}_m}, \quad m = 1, \dots, M \quad (49)$$

and all other  $\lambda_{\mathbf{k}} = 0$ . We use the shorthand  $\mathcal{Z}_L^{(M)}(\lambda, \mu, T)$  when there is no possibility of confusion. As we shall see, the limit  $L \rightarrow \infty$ ,  $\epsilon \rightarrow 0$  of this object leads to a *hierarchy* of equations connecting different values of  $M$ . We sketch the derivation of these equations, with more details in Appendix B, and then analyze their basic properties.

### 3.2.1. Derivation

The method is the same as before. We use the perturbation expansion in  $\epsilon$  giving the formula (27) for the generating functions, with the definitions (28) of  $\mathcal{X}_L(\lambda, \mu)$  and (29)-(33) of the  $\mathcal{J}$ 's. As a consequence of (49) all separate terms in the prefactors of the  $\mathcal{J}$ 's are of the same order:

$$\lambda_{\mathbf{k}_1} + \frac{\mu_{\mathbf{k}_1}}{2J_{\mathbf{k}_1}}, \quad \lambda_{\mathbf{k}_1} + \lambda_{\mathbf{k}_1}^2 J_{\mathbf{k}_1} - \frac{\mu_{\mathbf{k}_1}^2}{4J_{\mathbf{k}_1}} = O(L^d)$$

$$\frac{1}{2} \lambda_{\mathbf{k}_1}^2 + \frac{\mu_{\mathbf{k}_1}}{4J_{\mathbf{k}_1}^2} \left( \frac{\mu_{\mathbf{k}_1}}{2} - 1 \right) + \frac{\lambda_{\mathbf{k}_1} \mu_{\mathbf{k}_1}}{2J_{\mathbf{k}_1}}, \quad \lambda_{\mathbf{k}_1} \lambda_{\mathbf{k}_2}, \quad \left( \lambda_{\mathbf{k}_1} + \frac{\mu_{\mathbf{k}_1}}{4J_{\mathbf{k}_1}} \right) \frac{\mu_{\mathbf{k}_2}}{J_{\mathbf{k}_2}} = O(L^{2d}).$$

Hence, we only need to calculate the leading-order graphical contributions. We already analyzed these in the previous section and this discussion carries over here, except that now some wavenumbers are discrete and take on only  $M$  values (mode 1 for  $\mathcal{J}_1 - \mathcal{J}_4$  and modes 1,2 for  $\mathcal{J}_5$ ) whereas all others are continuous in the infinite-box limit. One must consider carefully whether free wavenumbers in graphical summations are discrete or continuous to see whether their contribution is  $O(M)$  or  $O(L^d)$ . This analysis is carried out in Appendix B. It is found that  $\mathcal{X}_L^{(M)}(\lambda, \mu, T)$  in the large-box limit gets no contributions from  $\mathcal{J}_1, \mathcal{J}_4$  and  $O(1)$  contributions from  $\mathcal{J}_2, \mathcal{J}_3, \mathcal{J}_5$ . The result is

$$\begin{aligned} \mathcal{X}^{(M)}(\lambda, \mu, T) - \mathcal{X}^{(M)}(\lambda, \mu, 0) &\sim -9\epsilon^2 i \delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) \left| \Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}_j)) \right|^2 \\ &\quad \times \left| H_{\mathbf{k}_j}^\sigma \right|^2 \left( \lambda_{\mathbf{k}_j} + \lambda_{\mathbf{k}_j}^2 \frac{\partial}{\partial \lambda_{\mathbf{k}_j}} \right) \frac{\partial^2 \mathcal{Z}^{(M+2)}}{\partial \bar{\lambda}_2 \partial \bar{\lambda}_3} \Big|_{\bar{\lambda}_2 = \bar{\lambda}_3 = 0} \\ &+ \epsilon^2 \left[ 18i \delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \sigma_2 \int d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) E_T(0, \underline{\sigma} \cdot \omega(\mathbf{k}_j)) \right. \\ &\quad \times \left| H_{\mathbf{k}_j}^\sigma \right|^2 \lambda_{\mathbf{k}_j} \frac{\partial^2 \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_3 \partial \lambda_j} \Big|_{\bar{\lambda}_3 = 0} + (2 \leftrightarrow 3) \\ &- 9 \sum_{j=1}^M \delta_{\mu_j,1} \delta_{\mu-j,1} \prod_{m \neq j, -j} \delta_{\mu_m,0} \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \sigma_2 \int d^d \bar{k}_2 \int d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) \\ &\quad \times E_T(\omega(\mathbf{k}_j) + \omega(-\mathbf{k}_j), \underline{\sigma} \cdot \omega(\mathbf{k}_j)) H_{\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{-, \sigma_2, \sigma_3} H_{-\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{+, \sigma_2, \sigma_3} \left\langle e^{i \sum_m \lambda_m \tilde{J}_m} \sqrt{\frac{\tilde{J}_{-1}}{\tilde{J}_1} \tilde{J}_3} \right\rangle_J + (2 \leftrightarrow 3) \Big] \\ &+ 18\epsilon^2 \sum_{j=1}^M \delta_{\mu_j,1} \delta_{\mu-j,1} \prod_{m \neq 1, -1} \delta_{\mu_m,0} \left( \lambda_1 \lambda_{-1} - i \frac{\lambda_1}{\tilde{J}_{-1}} - \frac{1}{4\tilde{J}_1 \tilde{J}_{-1}} \right) \sum_{\underline{\sigma}=(-, \sigma_2, \sigma_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) \\ &\quad \times \Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}_j)) \Delta_T(-\underline{\sigma} \cdot \omega(\mathbf{k}'_j)) H_{\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{-, \sigma_2, \sigma_3} H_{-\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{+, \sigma_2, \sigma_3} \left\langle \sqrt{\tilde{J}_1 \tilde{J}_{-1} \tilde{J}_2 \tilde{J}_3} e^{i \sum_m \lambda_m \tilde{J}_m} \right\rangle. \quad (50) \end{aligned}$$

Here  $\underline{\mathbf{k}}_j = (\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3)$  and  $\underline{\mathbf{k}}'_j = (-\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3)$ . Taking the small- $\epsilon$  limit using the asymptotic formulas (39), one finds the terms proportional to  $\delta_{\mu,0}$  contain secular contributions while the terms proportional to  $\delta_{\mu_j,1} \delta_{\mu-j,1}$  are nonsecular. Using (40) to calculate  $\mathcal{Z}^{(M)}(T) - \mathcal{Z}^{(M)}(0)$ , replacing  $(\mathcal{Z}^{(M)}(T) - \mathcal{Z}^{(M)}(0))/T$  by  $\dot{\mathcal{Z}}^{(M)}$  and using time variable  $\tau = \epsilon^2 t$ ,

one finally obtains

$$\begin{aligned} \frac{d}{d\tau} \mathcal{Z}^{(M)}(\lambda, \mu, \tau) = & -36i\pi\delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma}=(-1, \bar{\sigma}_2, \bar{\sigma}_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) \delta(\underline{\sigma} \cdot \omega(\mathbf{k}_j)) |H_{\mathbf{k}_j}^{\underline{\sigma}}|^2 \\ & \left\{ (\lambda_j + \lambda_j^2 \frac{\partial}{\partial \lambda_j}) \frac{\partial^2 \mathcal{Z}^{(M+2)}}{\partial \bar{\lambda}_2 \partial \bar{\lambda}_3} \Big|_{\bar{\lambda}_2 = \bar{\lambda}_3 = 0} - \bar{\sigma}_2 \lambda_j \frac{\partial \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_3 \partial \lambda_j} \Big|_{\bar{\lambda}_3 = 0} - \bar{\sigma}_3 \lambda_j \frac{\partial \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_2 \partial \lambda_j} \Big|_{\bar{\lambda}_2 = 0} \right\}. \end{aligned} \quad (51)$$

Equation (51) is the main result of this section.

### 3.2.2. Properties

We now consider the important properties of the multi-mode equations (51). As for the spectral characteristic functional in the previous section, the factors  $\delta_{\mu,0}$  imply that the RP property is preserved in time. Therefore, we can consider the generating functions for the amplitudes alone, obtained by setting  $\mu = 0$ . These form a *hierarchy* of equations, for  $M = 1, 2, 3, \dots$

$$\begin{aligned} \frac{d}{d\tau} \mathcal{Z}^{(M)}(\lambda, \tau) = & -36i\pi \sum_{j=1}^M \sum_{\underline{\sigma}=(-1, \bar{\sigma}_2, \bar{\sigma}_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) \delta(\underline{\sigma} \cdot \omega(\mathbf{k}_j)) |H_{\mathbf{k}_j}^{\underline{\sigma}}|^2 \\ & \left\{ (\lambda_j + \lambda_j^2 \frac{\partial}{\partial \lambda_j}) \frac{\partial^2 \mathcal{Z}^{(M+2)}}{\partial \bar{\lambda}_2 \partial \bar{\lambda}_3} \Big|_{\bar{\lambda}_2 = \bar{\lambda}_3 = 0} - \bar{\sigma}_2 \lambda_j \frac{\partial \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_3 \partial \lambda_j} \Big|_{\bar{\lambda}_3 = 0} - \bar{\sigma}_3 \lambda_j \frac{\partial \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_2 \partial \lambda_j} \Big|_{\bar{\lambda}_2 = 0} \right\}. \end{aligned} \quad (52)$$

By straightforward Fourier transformation in the  $\lambda$  variables, one can obtain an equivalent hierarchy of equations for the joint PDF's  $\mathcal{P}^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M)$ . For each  $M = 1, 2, 3, \dots$  these are equations for conservation of probability

$$\dot{\mathcal{P}}^{(M)} + \sum_{m=1}^M \frac{\partial}{\partial s_m} \mathcal{F}_m^{(M)} = 0, \quad (53)$$

with the probability flux

$$\begin{aligned} \mathcal{F}_m^{(M)} = & -36\pi s_m \sum_{\underline{\sigma}=(-1, \bar{\sigma}_2, \bar{\sigma}_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_m) \delta(\underline{\sigma} \cdot \omega(\mathbf{k}_m)) |H_{\mathbf{k}_m}^{\underline{\sigma}}|^2 \\ & \left[ \int d\bar{s}_2 d\bar{s}_3 \frac{\partial \mathcal{P}^{(M+2)}}{\partial s_m}(s_1, \dots, s_M, \bar{s}_2, \bar{s}_3) \bar{s}_2 \bar{s}_3 \right. \\ & \left. + \bar{\sigma}_2 \int d\bar{s}_3 \mathcal{P}^{(M+1)}(s_1, \dots, s_M, \bar{s}_3) \bar{s}_3 + \bar{\sigma}_3 \int d\bar{s}_2 \mathcal{P}^{(M+1)}(s_1, \dots, s_M, \bar{s}_2) \bar{s}_2 \right]. \end{aligned} \quad (54)$$

We obtain no closed equations for  $\mathcal{P}^{(M)}$  with any choice of  $M$  but instead an infinite hierarchy, in which the evolution equation for  $\mathcal{P}^{(M)}$  contains  $\mathcal{P}^{(M+1)}$  and  $\mathcal{P}^{(M+2)}$ , for  $M = 1, 2, 3, \dots$ . Our results are therefore not in agreement with previously proposed closed equations for  $\mathcal{P}^{(2)}$  with  $M = 2$  [17].

We do, however, recover the equations for  $\mathcal{P}^{(1)}$  and  $\mathcal{Z}^{(1)}$  with  $M = 1$  which were previously obtained [2, 3, 4], under appropriate conditions. Namely, assume that the initial data for the hierarchy (52) are factorized

$$\mathcal{Z}^{(M)}(\lambda_1, \dots, \lambda_M, 0; \mathbf{k}_1, \dots, \mathbf{k}_M) = \prod_{m=1}^M Z(\lambda_m, 0; \mathbf{k}_m),$$

as would follow from RPA initial conditions, for example. It is then easy to show by substitution into (52) that there are solutions which remain factorized

$$\mathcal{Z}^{(M)}(\lambda_1, \dots, \lambda_M, \tau; \mathbf{k}_1, \dots, \mathbf{k}_M) = \prod_{m=1}^M Z(\lambda_m, \tau; \mathbf{k}_m), \quad \tau > 0,$$

where the factors  $Z(\lambda, \tau; \mathbf{k})$  satisfy the closed equations

$$\frac{\partial}{\partial \tau} Z(\lambda; \mathbf{k}) = i\eta_{\mathbf{k}}\lambda \left(1 + \lambda \frac{\partial}{\partial \lambda}\right) Z(\lambda; \mathbf{k}) - \gamma_{\mathbf{k}}\lambda \frac{\partial Z}{\partial \lambda}(\lambda; \mathbf{k})$$

with

$$\eta_{\mathbf{k}} = 36\pi \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \delta(\underline{\sigma} \cdot \omega(\mathbf{k})) |H_{\mathbf{k}}^{\underline{\sigma}}|^2 n(\mathbf{k}_2) n(\mathbf{k}_3) \geq 0, \quad (55)$$

$$\gamma_{\mathbf{k}} = 36\pi \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \delta(\underline{\sigma} \cdot \omega(\mathbf{k})) |H_{\mathbf{k}}^{\underline{\sigma}}|^2 [\sigma_3 n(\mathbf{k}_2) + \sigma_2 n(\mathbf{k}_3)]. \quad (56)$$

These results are equivalent to the existence of solutions of the PDF hierarchy (53),(54) that remain factorized

$$\mathcal{P}^{(M)}(s_1, \dots, s_M, \tau; \mathbf{k}_1, \dots, \mathbf{k}_M) = \prod_{m=1}^M P(s_m, \tau; \mathbf{k}_m), \quad \tau > 0$$

where the 1-mode pdfs  $P(s, \tau; \mathbf{k})$  satisfy

$$\frac{\partial}{\partial \tau} P = \frac{\partial}{\partial s} \left[ s \left( \eta_{\mathbf{k}} \frac{\partial P}{\partial s} + \gamma_{\mathbf{k}} P \right) \right]. \quad (57)$$

Preservation of RPA therefore follows if one can prove uniqueness of solutions of the hierarchy (52) or, equivalently, (53),(54). The equations for the factors agree with previous results for the 1-mode equations [2, 3, 4].

The equations (57) are not simple linear Fokker-Planck equations, however, but are instead *nonlinear* Markov evolution equations in the sense of McKean [18]. That is, the solutions must satisfy a set of self-consistency conditions,

$$n(\mathbf{k}, \tau) = \int ds s P(s, \tau; \mathbf{k}) \quad (58)$$

where  $n(\mathbf{k}, \tau)$  is the same spectrum that appears in the formulas for the coefficients (55),(56). These equations are the exact solutions of a model of “self-consistent Langevin equations”, like those for the DIA turbulence closure [26, 27]. Here the model equations take the form of the stochastic differential equations

$$ds_{\mathbf{k}} = (\eta_{\mathbf{k}} - \gamma_{\mathbf{k}} s_{\mathbf{k}}) d\tau + \sqrt{2\eta_{\mathbf{k}} s} dW_{\mathbf{k}}, \quad (59)$$

interpreted in the Ito sense and with self-consistent determination of  $n(\mathbf{k}, \tau)$  via (58). Solutions of this stochastic model can be realized by a Monte Carlo procedure of McKean [18], with (59) generalized to

$$ds_{\mathbf{k}}^{(n)} = (\eta_{\mathbf{k}} - \gamma_{\mathbf{k}} s_{\mathbf{k}}^{(n)}) dt + \sqrt{2\eta_{\mathbf{k}} s^{(n)}} dW_{\mathbf{k}}^{(n)}, \quad \mathbf{k} \in \Lambda_L^*, n = 1, \dots, N,$$

with the spectrum obtained by an  $N$ -sample average

$$n_{\mathbf{k}} = \frac{1}{N} \sum_{n=1}^N s_{\mathbf{k}}^{(n)},$$

and with  $\gamma_{\mathbf{k}}, \eta_{\mathbf{k}}$  given by formulas (55),(56) in which wavenumber integrals are discretized as sums. Taking first  $N \gg 1$  and then  $L \gg a$  yields a solution of the PDF equations (57). This procedure works as well to solve the general hierarchy equations (53),(54), without assuming factorized initial data. For the factorized case, a far simpler procedure is to solve first the wave kinetic equation (45) for  $n(\mathbf{k}, \tau)$  and then, using this as input, to solve the 1-mode equation (57) in order to obtain  $P(s, \tau; \mathbf{k})$  for any wavenumber mode  $\mathbf{k}$  of interest. As initial condition one may take any  $P(s, 0; \mathbf{k})$  which satisfies the consistency condition (58) at  $\tau = 0$ , since this condition is preserved in time by the 1-mode equation.

It is interesting that, independent of the initial condition  $P(s, 0; \mathbf{k})$ , the solution  $P(s, \tau; \mathbf{k})$  relaxes as  $\tau$  increases to a Rayleigh distribution

$$Q(s, \tau; \mathbf{k}) = \frac{1}{n(\mathbf{k}, \tau)} \exp(-s/n(\mathbf{k}, \tau)), \quad (60)$$

which corresponds to a Gaussian distribution of the Fourier coefficient  $\tilde{a}_{\mathbf{k}}(\tau)$ . It is easy to check that, for any solution  $n(\mathbf{k}, \tau)$  of the wave kinetic equation (45),  $Q(s, \tau; \mathbf{k})$  defined above solves the 1-mode PDF equation (57). (Since wavenumber  $\mathbf{k}$  appears only as a parameter in our argument, we suppress its appearance for the rest of this paragraph.) The relaxation of a general solution  $P$  to  $Q$  is indicated by an  $H$ -theorem for the *relative entropy*<sup>8</sup>

$$H(P|Q) = \int ds P(s) \ln \left( \frac{P(s)}{Q(s)} \right) = \int ds P(s) \ln P(s) + \ln n + 1.$$

This is a convex function of  $P$ , non-negative, and vanishing only for  $P = Q$  [28]. Taking the time-derivative using (57), it is straightforward to derive

$$\frac{d}{d\tau} H(P(\tau)|Q(\tau)) = -\eta \int ds \frac{s|\partial_s P(s, \tau)|^2}{P(s, \tau)} + \frac{\eta}{n(\tau)},$$

where

$$\int -s\partial_s P(s, \tau) ds = \int P(s, \tau) ds = 1 \quad (61)$$

was used to cancel terms involving the  $\gamma$  coefficient. But note the self-consistency condition  $n(\tau) = \int s P(s, \tau) ds$  implies

$$\frac{d}{d\tau} H(P(\tau)|Q(\tau)) = -\eta \left( \int ds \frac{s|\partial_s P(s, \tau)|^2}{P(s, \tau)} - \frac{1}{\int s P(s, \tau) ds} \right) \leq 0.$$

The inequality follows from the Cauchy-Schwartz inequality applied to (61)

$$1 = \int \sqrt{sP} \cdot \sqrt{\frac{s}{P}} (-\partial_s P) ds \leq \sqrt{\int s P(s) ds} \cdot \sqrt{\int \frac{s|\partial_s P|^2}{P} ds}.$$

Equality holds and entropy production vanishes if and only if  $\sqrt{sP} = c \sqrt{\frac{s}{P}} (-\partial_s P)$  for some constant  $c$ , or  $P = -c\partial_s P$ . The solution of this latter equation gives  $P = Q$  with  $n = 1/c$ . We thus see that  $P(\tau)$  should relax to  $Q(\tau)$  as  $\tau$  increases. For the purpose of later discussion, we emphasize that this argument assumes that the kinetic theory is valid over the entire range of amplitudes  $s \in (0, \infty)$  and it could otherwise fail.

Let us remark that the relative entropy has a simple probabilistic meaning, similar to that discovered by Boltzmann for his entropy function, which involves the empirical PDF  $\tilde{P}_L(s; \Delta)$  defined in (19). For a single cell  $\Delta$  in wavenumber space  $\Lambda^*$ , let  $\mu_{\Delta, n, \eta}$  be the microcanonical measure on the set of microstates  $\Gamma_{\Delta, n, \eta} = \{(\tilde{J}, \psi) : |\tilde{n}_L(\Delta) - n(\Delta)| < \eta\}$ . Then, with respect to this microcanonical measure, the most probable value of  $\tilde{P}_L(s; \Delta)$  as first  $L \rightarrow 0$ , then  $\eta \rightarrow 0$  is  $Q(s; \Delta) = e^{-s/n(\Delta)}/n(\Delta)$ . However, the probability of observing another PDF  $P$  as a rare fluctuation is

$$\lim_{\eta \rightarrow 0} \lim_{L \rightarrow \infty} \frac{1}{N_L(\Delta)} \ln \mu_{\Delta, n, \eta}(\{\tilde{P}_L \approx P\}) = -H(P|Q). \quad (62)$$

This can be shown heuristically by adapting the original argument of Boltzmann [23]. Let  $P(s)$  be a probability density function over possible values of  $s$ , let  $\Sigma = \{\Sigma_j, j = 1, \dots, R\}$  be a finite partition of the positive reals into intervals with  $s_j$  the midpoint values and define the integers  $N_j = \left\lfloor N \int_{\Sigma_j} P(s) ds \right\rfloor$  where  $\lfloor \cdot \rfloor$  denotes integer part and  $N = N_L(\Delta)$ . (Since  $\Delta$  is fixed in this argument, we hereafter omit explicit reference to that quantity in the remainder of this paragraph.) Note that  $\sum_{j=1}^R N_j = N$  for large enough  $L$ . Now let  $\hat{N}_j$  denote the number of modes  $\mathbf{k} \in \Lambda_L^* \cap \Delta$

<sup>8</sup>A related set of observations were made in [2], section 9.

such that  $\tilde{J}_{\mathbf{k}} \in \Sigma_j$  and define the set  $\Gamma_{P,\Sigma,L} = \{(\tilde{J}, \psi) : \hat{N}_j = N_j, j = 1, \dots, R\}$ . The Liouville measure of this set is  $|\Gamma_{P,\Sigma,L}| = \frac{N!}{N_1! \dots N_R!} |\Sigma_1|^{N_1} \dots |\Sigma_R|^{N_R} \sim \exp(-N \sum_j |\Sigma_j| P_j \ln P_j)$  with  $P_j = N_j/N|\Sigma_j|$  and using Stirling's approximation for the factorials, exactly as in the original argument of Boltzmann. Then

$$\lim_{|\Sigma| \rightarrow 0} \lim_{L \rightarrow \infty} \frac{1}{N} \ln |\Gamma_{P,\Sigma,L}| = - \int ds P(s) \ln P(s).$$

Now consider the set  $\Gamma_{n,\eta,L} = \cup_{\{P: |n - \sum_j |\Sigma_j| s_j P_j| < \eta\}} \Gamma_{P,\Sigma,L}$ . Its Liouville measure  $|\Gamma_{n,\eta,L}|$  is dominated in the limits  $L \rightarrow \infty$ ,  $\eta \rightarrow 0$ ,  $|\Sigma| \rightarrow 0$  by  $|\Gamma_{P,\Sigma,L}|$  for the pdf  $P$  satisfying  $\int ds s P(s) = n$  with the largest entropy  $-\int ds P(s) \ln P(s)$ . As is well-known, this maximum entropy distribution is the Rayleigh pdf  $Q(s) = e^{-s/n}/n$ . Thus,

$$\lim_{|\Sigma| \rightarrow 0} \lim_{\eta \rightarrow 0} \lim_{L \rightarrow \infty} \frac{1}{N} \ln |\Gamma_{n,\eta,L}| = - \int ds Q(s) \ln Q(s) = \ln n + 1.$$

Taking  $\mu_{n,\eta}(\{\widehat{P}_L \approx P\}) = \frac{|\Gamma_{P,\Sigma,L}|}{|\Gamma_{n,\eta,L}|}$ , then  $\lim_{|\Sigma| \rightarrow 0} \lim_{\eta \rightarrow 0} \lim_{L \rightarrow \infty} \frac{1}{N} \ln \frac{|\Gamma_{P,\Sigma,L}|}{|\Gamma_{n,\eta,L}|} = -(\int ds P(s) \ln P(s) + \ln n + 1)$ , as claimed. The result (62) becomes the standard Sanov theorem [28], if the measure  $\mu_{\Delta,n,\eta}$  is replaced by the RPA measure for which each mode  $\tilde{J}_{\mathbf{k}}$ ,  $\mathbf{k} \in \Lambda_L^* \cap \Delta$  has the independent density  $Q(s; \Delta)$ .

The above argument motivates a microcanonical measure  $\mu_{P,\Delta,\Sigma,\eta,L}$  defined for a given  $P(s; \mathbf{k})$  by normalizing Liouville measure on the set

$$\Gamma_{P,\Delta,\Sigma,\eta,L} = \left\{ (\tilde{J}, \psi) : \left| \int_{\Sigma_j} ds \widehat{P}_L(s, \Delta_i) - \int_{\Sigma_j} ds P(s, \Delta_i) \right| < \eta, i = 1, \dots, P, j = 1, \dots, R \right\}$$

for partitions  $\Delta$  of  $\Lambda^*$  and  $\Sigma$  of  $\mathbb{R}^+$ . This measure satisfies the RP property, as it contains no dependence on the phases. One can define for this measure the  $M$ -mode correlation functions of the empirical PDF,  $\widehat{P}_L(\varphi) = \left(\frac{2\pi}{L}\right)^d \sum_{\mathbf{k} \in \Lambda_L^*} \varphi(\tilde{J}_{\mathbf{k}}, \mathbf{k})$ :

$$\mathcal{P}_{P,\Delta,\Sigma,\eta,L}^{(M)}(\varphi_1, \dots, \varphi_M) = \mu_{P,\Delta,\Sigma,\eta,L}(\widehat{P}_L(\varphi_1) \cdots \widehat{P}_L(\varphi_M))$$

for any choice of continuous functions  $\varphi_1, \dots, \varphi_M$  on  $\mathbb{R}^+ \times \Lambda^*$  vanishing at infinity. The arguments of [15] can be adapted to show that

$$\lim_{|\Delta|, |\Sigma| \rightarrow 0} \lim_{\eta \rightarrow 0} \lim_{L \rightarrow \infty} \mathcal{P}_{P,\Delta,\Sigma,\eta,L}^{(M)}(\varphi_1, \dots, \varphi_M) = P(\varphi_1) \cdots P(\varphi_M)$$

where  $P(\varphi) = \int_0^\infty ds \int_{\Lambda^*} d^d k \varphi(s, \mathbf{k}) P(s; \mathbf{k})$ . Thus the factorization property of the  $M$ -mode PDFs holds asymptotically for the microcanonical measure  $\mu_{P,\Delta,\Sigma,\eta,L}$ , although it does not satisfy the strict RPA property. Our derivation of the PDF hierarchy equations (53),(54) assumed existence of the  $M$ -mode PDF's in a somewhat stronger sense in the limit as  $L \rightarrow \infty$  (i.e. for fixed values of  $\mathbf{k}_1, \dots, \mathbf{k}_M$ ), but it is reasonable to expect that they remain valid for such "generalized RPA" initial conditions.

Lastly, we can ask what are the solutions of the PDF hierarchy for general initial data which are RP but which are not factorized even asymptotically. One must assume at least that

$$\lim_{L \rightarrow \infty} \langle \widehat{P}_L(\varphi_1, 0) \cdots \widehat{P}_L(\varphi_M, 0) \rangle = \int_0^\infty ds_1 \cdots \int_0^\infty ds_M \int d^d k_1 \cdots d^d k_M \times \varphi_1(s_1, \mathbf{k}_1) \cdots \varphi_M(s_M, \mathbf{k}_M) \mathcal{P}_0^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M),$$

in order to provide suitable initial data  $\mathcal{P}_0^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M)$  for the hierarchy. However, with appropriate technical assumptions, it follows by the argument of Spohn [16] that

$$\mathcal{P}_0^{(M)}(s_1, \dots, s_M; \mathbf{k}_1, \dots, \mathbf{k}_M) = \int d\rho_0(P) P(s_1; \mathbf{k}_1) \cdots P(s_M; \mathbf{k}_M),$$

where  $\rho_0$  is a probability measure on the PDF's. Since the PDF hierarchy equations (53),(54) are linear in the  $\mathcal{P}^{(M)}$ 's, a solution will be provided by

$$\mathcal{P}^{(M)}(s_1, \dots, s_M, \tau; \mathbf{k}_1, \dots, \mathbf{k}_M) = \int d\rho_0(P) P(s_1, \tau; \mathbf{k}_1) \cdots P(s_M, \tau; \mathbf{k}_M).$$

This will be the only solution if existence and uniqueness of solutions holds for the PDF hierarchy. Thus, the most general realizable solutions of the PDF hierarchy equations are again expected to be "super-statistical solutions" that correspond to ensembles of solutions of the 1-mode PDF equations (57) with random initial conditions.



#### 4. Intermittency in Kinetic Wave Turbulence

One of the important potential applications of multimode equations in wave kinetics is the explanation of observed intermittency and anomalous scaling in wave turbulence [10, 9, 11]. On the face of it, wave kinetic theory appears to have few resources to explain such phenomena and instead has all the attributes of a “mean-field theory” [29]. As we have seen in previous sections, it is a theory which ignores fluctuations and in which important quantities in fact become deterministic. All interactions of the infinite collection of modes is through a mean-field  $n(\mathbf{k}, \tau)$ . Furthermore, scaling exponents in the Kolmogorov solutions of the wave kinetic equations are given by dimensional analysis [12]. These are the hallmarks of a mean-field theory, which generally cannot predict anomalous scaling.

Nevertheless, there are at least two approaches based on wave kinetics which seem to have some promise to explain intermittency and non-Gaussian statistics. One is the idea of a “cascade in amplitude space” that was proposed by Choi et al. [3, 9], based on the 1-mode PDF equation (57). Another is the idea of a “super-turbulence” in wave-kinetics, advanced in the present work. We shall now discuss both of these possibilities.

##### 4.1. Cascade in Amplitude Space?

It is well-known that wave kinetic equations generally cannot be uniformly valid over the whole range of wavenumbers, but must break down in either low or high wavenumbers where nonlinearity become strong [12]. Choi et al. [3] (also [9]) have proposed that there is a similar non-uniformity in amplitude space, with equation (57) for the 1-mode PDF  $P(s, \tau; \mathbf{k})$  restricted in validity to  $s < s_{nl}$ , where  $s_{nl}$  is the amplitude for which nonlinear interactions become strong at wavenumber  $\mathbf{k}$ . [For the remainder of this section we shall assume that the wavenumber  $\mathbf{k}$  is fixed and drop it as an explicit label.] For amplitudes  $s > s_{nl}$  strong-interaction processes such as “wave-breaking”, “cresting”, etc. are assumed to occur which are beyond the description of wave kinetics. The hope is that the effects of such nonlinear processes can be modeled by supplementing the equation (57) with suitable boundary conditions and forcing terms. We find this a very intriguing suggestion but, as we now argue, it seems to us to have as yet no successful formulation, either analytically or physically.

Let us review the specific proposals of [3, 9]. They note that eq.(57) is a probability conservation law

$$\partial_t P + \partial_s F = 0 \quad (63)$$

with probability flux in amplitude space given by

$$F(s) = -s(\gamma P + \eta \partial_s P). \quad (64)$$

The general time-independent solution of (63) with constant flux  $F(s) = F_*$  was observed in [3] to be

$$P(s) = C e^{-s/\nu} - \frac{F_*}{\eta} \text{Ei}(s/\nu) e^{-s/\nu}$$

with  $\nu \equiv \eta/\gamma$  and with  $\text{Ei}(x) = \text{P} \int_{-\infty}^x e^t \frac{dt}{t}$  the standard exponential integral. The first term has zero flux  $F(s) \equiv 0$  while the second term has non-vanishing flux  $F(s) \equiv F_*$ . Since  $\text{Ei}(x) > 0$  for  $x > x_* \doteq 0.3725$ , positivity of  $P(s)$  for  $s \gg \nu$  requires  $F_* \leq 0$ . When  $F_* = 0$  then  $P(s) = Q(s)$ , the Rayleigh distribution with spectral density  $n = \nu$ . Not noted in [3] is the fact that one must more generally make a distinction between  $\nu$  and  $n$  given by the self-consistency condition

$$n = \int ds s P(s),$$

which are only the same when  $P = Q$ . The above integral diverges for  $F_* \neq 0$ , if the upper limit extends to infinity, because  $P(s) \sim \frac{|F_*|}{\gamma s}$  due to the asymptotics  $\text{Ei}(x) \sim \frac{e^x}{x} \left(1 + \frac{1!}{x} + \frac{2!}{x^2} + \dots\right)$  for  $x \gg 1$ . The integral is finite if it is cut off at the upper limit  $s_{nl}$ . This assumes that whatever nonlinear processes such as “wave-breaking” occur at  $s > s_{nl}$  lead to a PDF which decays very rapidly for  $s \gtrsim s_{nl}$ .

The papers [3, 9] propose to model the nonlinear effects at  $s > s_{nl}$  by a negative flux  $F(s_{nl}) = F_* < 0$ , which represents a flow of samples back into the kinetic range  $s < s_{nl}$ , at least in some range of wavenumbers. For example, this might occur at low-wavenumbers as wavecaps and cusps in the nonlinear range break up and feed back into

the weak, incoherent background. However, the equation (63) will then no longer conserve probability but instead satisfies  $\frac{d}{dt} \int_0^{s_{nl}} ds P(s) = |F_*| > 0$ . Paper [3] proposes to add to the equation a constant “drag”  $D$

$$\partial_t P + \partial_s F = -D, \quad s < s_{nl} \quad (65)$$

to represent the dilution in the weight of samples at  $s < s_{nl}$  by the flux  $F_*$  of new samples into the ensemble at  $s = s_{nl}$ . It is easy to see that one must choose  $D = |F_*|/s_{nl}$  to conserve total probability<sup>9</sup>. Paper [3] then claims that the general solution to the modified equation is  $P(s) = C e^{-s/n} + \frac{|F_*|}{n} \text{Ei}(s/n - \log(s/n)) e^{-s/n}$ . The picture proposed in [3, 9] is one of an “inverse probability cascade” with flux  $F_* < 0$  in amplitude space, coexisting with the usual forward energy cascade in wavenumber space. The power-law tails  $P(s) \sim \frac{|F_*|}{\gamma s}$  for  $s \gg \nu$  are a source of non-Gaussianity and intermittency in wave turbulence if  $F_* \neq 0$ . A “critical balance” argument is used in [9] to estimate that  $|F_*| \sim \gamma n/s_{nl}$ .

It is easy to see, however, that the proposed cascade picture cannot be correct for the specific model in [3, 9], on both physical and mathematical grounds. A constant “drag”  $D$  is not localized at small and large  $s$  but has effects felt over the whole range of  $s$ . Thus, one would not expect a constant-flux “inertial range” to exist for such a drag. In fact, for any choice of  $D$ , a particular solution of the inhomogeneous equation (65) is

$$P_{part}(s) = D/\gamma.$$

The general solution  $P(s)$  of (65) is a superposition of  $P_{part}(s)$  with any solution of the homogeneous equation (63):

$$P(s) = \frac{D}{\gamma} + \frac{C_1}{\nu} e^{-s/\nu} + \frac{C_2}{\nu} e^{-s/\nu} \text{Ei}\left(\frac{s}{\nu}\right).$$

This has flux  $F(s) = -Ds - C_2\gamma$ . With the b.c.  $F(s_{nl}) = F_*$  and the unique probability-conserving choice  $D = |F_*|/s_{nl}$  one finds  $C_2 = 0$  and thus

$$P(s) = \frac{|F_*|}{\gamma} \cdot \frac{1}{s_{nl}} + \left(1 - \frac{|F_*|}{\gamma}\right) \frac{e^{-s/\nu}}{\nu(1 - e^{-s_{nl}/\nu})}, \quad (66)$$

using normalization. Note the requirement  $|F_*| < \gamma$  for positivity of this solution, which is then a superposition of a Rayleigh distribution and a uniform distribution. As expected, it has not constant flux but instead  $F(s) = F_* s/s_{nl}$ .

We claim that (66) is the correct solution of the model formulated in [3]. It gives an alternative possibility to explain intermittency and non-Rayleigh distributions, with constant tails  $P(s) \propto s^0$  for  $s \gg \nu$  rather than  $P(s) \propto s^{-1}$ . Using the estimate  $|F_*| \sim \gamma n/s_{nl}$  of [9] one finds that the realizability inequality  $|F_*| < \gamma$  is satisfied whenever  $n \lesssim s_{nl}$ . However, it is unclear to us that the flux b.c. and constant drag  $D$  is a physically correct model of wave-breaking. Constant-flux solutions as originally proposed in [3] could be relevant with a more general model of strong nonlinear effects, in which the constant  $D$  is replaced with a function  $D(s)$  such that  $\int_0^{s_{nl}} ds D(s) = |F_*|$  but for which  $D(s)$  nearly vanishes except at the upper and lower limits. This would provide a “transparency window” that would allow constant probability-flux solutions to exist.

Whatever may be the correct model of the strong nonlinear effects at  $s > s_{nl}$ , an important general point is that any modification of the 1-mode PDF eq.(63) requires that the wave kinetic equation must also be modified. This can be illustrated for the specific model in (65) with  $D = |F_*|/s_{nl}$ . Taking the time-derivative of  $n = \int_0^{s_{nl}} ds s P(s)$  gives

$$\dot{n} = -\gamma n + \eta - \eta s_{nl} P(s_{nl}) + \frac{1}{2} |F_*| s_{nl}, \quad (67)$$

which differs from the standard kinetic equation  $\dot{n} = -\gamma n + \eta$ . The stationary form of the “modified kinetic equation” (67) is easily checked to be valid for our solution (66) by calculating  $n = \int_0^{s_{nl}} ds s P(s)$  to be

$$\begin{aligned} n &= \nu - \nu s_{nl} P(s_{nl}) + \frac{|F_*|}{2\gamma} s_{nl} \\ &\simeq \nu \left(1 - \frac{|F_*|}{\gamma}\right) + \frac{|F_*|}{2\gamma} s_{nl}, \end{aligned} \quad (68)$$

<sup>9</sup>In fact, [3] proposed to take  $D = -\gamma P(s_{nl})/s_{nl}$ , which does not conserve probability and which is not even dimensionally correct.

with the latter approximation valid for  $s_{nl} \gg \nu$ . The correction to the usual value  $n = \nu$  predicted by the kinetic equation need not be small. If we use the estimate  $|F_*| \sim \gamma n / s_{nl}$  from “critical balance” [9] then the second term in (68) is comparable to the first term, or even much larger if  $\nu \ll n$ . If such changes in  $n(\mathbf{k})$  occur for a large set of wavenumbers  $\mathbf{k}$ , then the coefficients  $\gamma$  and  $\eta$  defined through the integrals (55),(56) may also be strongly modified. These remarks make clear the nontriviality of constructing a self-consistent hybrid model of wave kinetics and of the strong nonlinear effects.

#### 4.2. Super-Turbulence of Wave Kinetics?

A second possibility to explain intermittency and anomalous scaling entirely within the framework of wave kinetics is by “super-turbulence”. We shall discuss this mechanism here, mainly at a general, qualitative level.

It is useful to begin with a review of the work of Spohn [16], who already discussed turbulence in gas dynamics as a natural motivation for ensembles of Boltzmann solutions. This is easiest to understand in the hydrodynamic regime where length-scale of variations in local equilibrium parameters (temperature, density, velocity) is much larger than the mean-free-path length. The Boltzmann equation reduces then to a hydrodynamic description and one can appeal to the extensive literature on turbulent solutions of hydrodynamic equations. The specific example discussed in [16] is the Rayleigh-Bernard system, considered as a many-particle system subject to thermal boundary conditions. In such a situation driven by the boundary conditions, Boltzmann’s original  $H$ -theorem is no longer valid and the stationary solution of the Boltzmann equation for the pure conducting state is no longer purely Maxwellian but has a small correction corresponding to thermal non-equilibrium [30]. In the turbulent regime at sufficiently high Reynolds and Rayleigh numbers, the laminar purely conducting state is unstable and turbulent convection develops. The temporal dynamics is chaotic so that long-time-averages, for example, are described naturally by invariant measures corresponding to ensembles of Boltzmann solutions.

Such a statistical description is natural also for turbulent situations without driving by body forces or boundaries and with time-dependent statistics. For example, a single realization of decaying, statistically homogeneous turbulence is very spatially complex and heterogeneous. Averages over large volumes —by space-ergodicity—are again described by ensembles of solutions. Note in this example that Boltzmann’s  $H$ -theorem applies, with a monotonic increase of entropy due to heating of the fluid. There is eventual approach to global thermodynamic equilibrium and a space-independent Maxwellian 1-particle distribution, but the turbulent state with strong spatial variations exists as a long transient for an intermediate range of times. The previous examples are in the hydrodynamic regime, but there should be similar turbulence in the fully kinetic regime. For example, in compressible turbulence in gases at sufficiently high Mach numbers the thickness of shocks should be of the order of the mean-free-path length [31] and, for a quantitative description, the Boltzmann equation should be used rather than hydrodynamic equations.

These ideas can carry over from the Boltzmann equation to the wave kinetic equation. As noted in the introduction, Zakharov et al. ([1], section 4.2.2) had already pointed out that strong instability of the Kolmogorov solutions to the wave kinetic equation could lead to such a “secondary turbulence”. The statistical distribution of wave amplitudes that would be obtained in a long time-series from experiment or simulation would then correspond to a random ensemble of solutions of the 1-mode PDF equation (57):

$$\mathcal{P}^{(1)}(s; \mathbf{k}) = \int d\rho(P) P(s; \mathbf{k}).$$

Even if the individual solutions  $P(s; \mathbf{k})$  were close to Rayleigh distributions, the measurable distributions  $\mathcal{P}^{(1)}(s; \mathbf{k})$  could be arbitrarily far from Rayleigh, depending upon the “super-statistical” measure  $\rho$ . To determine this measure becomes a very difficult problem, as daunting as the corresponding problem for Navier-Stokes turbulence! Here we can only identify some of the sources of randomness, including instability, random forcing, and boundary conditions. We consider these in turn.

Instability of the Kolmogorov solution is a natural origin for “super-turbulence”. As a related example, consider the GOY shell model of turbulence, a dynamical system with a stationary Kolmogorov solution supporting a constant energy flux with dimensional scaling [32]. This solution, however, is subject to a rich array of instabilities [33, 34] and the statistical behavior of the GOY model shows strong intermittency and anomalous scaling. There is a detailed theory of linear stability of the Kolmogorov solutions of wave kinetic theory [35, 36, 37, 38], reviewed in [1], Ch. 4. This theory studies perturbations to a scale-homogeneous Kolmogorov solution which, strictly speaking, requires

an infinitely long inertial interval. The conclusion of this theory is that, for most common cases, the Kolmogorov solution is linearly stable. However, the theory, although sophisticated, is not quite definitive. Quoting from [1]:

“It should be noted that a more consistent formulation of the problem of the stability of the Kolmogorov spectrum should be as follows. First of all, the kinetic equation should be supplemented by terms describing the isotropic pumping and damping regions and a stationary solution of this equation should be found that is close to the Kolmogorov spectrum in an interval  $(k_1, k_2)$ ; outside this interval the solution may strongly differ from the Kolmogorov spectrum. Then the kinetic equation must be linearized in the vicinity of this stationary solution and expanded in angular harmonics. . . . Having examined the behavior of the solutions equation, one should clarify the changes that occur when the ranges of the source and sink in  $k$ -space go to zero or to infinity and examine the behavior of the perturbations established in the interval  $(k_1, k_2)$ . Finally, one should analyze in which situations this behavior is independent of the specific type of the source and sink.

This program for examining the stability of Kolmogorov spectra turns out to be too complex. Currently there exists no strict proof of the fact that in general the kinetic equation with a source and sink has a stationary solution close to the Kolmogorov spectrum in some interval.”

In fact, the situation is even less certain than this statement implies. There are hydrodynamic flows such as plane Couette and pipe Poiseuille which, despite being linearly stable for all Reynolds numbers, become turbulent in laboratory experiments and in numerical simulations at moderate Reynolds number [39]. The usual understanding is that these flows are unstable to finite amplitude perturbations, e.g. [40]. Thus, linear stability does not rule out transition to turbulence. For these reasons, we must regard “super-turbulence” as a viable possibility in wave kinetics.

As emphasized in the previous quote, external forcings are required to maintain a stationary energy cascade state. It is important to consider more deeply the origin and role of such forces. The monograph [1] considers the necessary conditions on the force for the existence of a stationary distribution (section 2.2.3) and how to match the Kolmogorov solutions to the wavenumber regions of the force (section 3.4). This discussion assumes a particular idealized model for the forcing, in which to the collision integral  $I(\mathbf{k}, \tau)$  of the wave kinetic equation there is added an additional term:

$$\dot{n}(\mathbf{k}, \tau) = \Gamma(\mathbf{k}, \tau)n(\mathbf{k}, \tau) + I(\mathbf{k}, \tau).$$

This model describes “pumping” for wavenumbers at which  $\Gamma(\mathbf{k}, \tau) > 0$  and “damping” for  $\Gamma(\mathbf{k}, \tau) < 0$ . Validity of this model requires a suitably weak, slowly-changing force in the equations of motion (10):

$$\frac{d}{dt}A_{\mathbf{k}}^\sigma = \frac{1}{2}\epsilon^2\Gamma(\mathbf{k}, \epsilon^2t)A_{\mathbf{k}}^\sigma + \dots$$

Here  $\dots$  denotes the original terms in (10). The addition of this term to the dynamics can be easily accommodated into our derivations, with a new term appearing in eq.(26)

$$a_{\mathbf{k}}^{(2)}(T) = \frac{1}{2}T \cdot \Gamma_{\mathbf{k}} a_{\mathbf{k}}^{(0)} + \dots,$$

in eq.(31)

$$\mathcal{J}_3 = \frac{1}{2}T\delta_{\mu,0} \sum_1 \Gamma_1 \lambda_1 J_1 + \dots,$$

in eq.(42)

$$\frac{d\mathcal{Z}}{d\tau} = \int d^d k \Gamma(\mathbf{k}, \tau) \lambda(\mathbf{k}) \frac{\delta\mathcal{Z}}{\delta\lambda(\mathbf{k})} + \dots,$$

in eq.(51)

$$\frac{d\mathcal{Z}^{(M)}}{d\tau} = \sum_{m=1}^M \Gamma(\mathbf{k}_m, \tau) \lambda_m \frac{\partial\mathcal{Z}^{(M)}}{\partial\lambda_m} + \dots$$

and in eq.(54)

$$\mathcal{F}_m^{(M)} = \Gamma(\mathbf{k}_m, \tau) s_m + \dots$$

Thus, the standard forcing model from [1] is obtained. All of our previous results, e.g. on preservation of factorized solutions, law of large numbers, etc., still carry over. In general, forced wave turbulence can be described by kinetic equations only for suitable assumptions on the forcing, like those above. In the above derivation, the only randomness was in the phases and amplitudes of the waves. However, the forcing amplitudes  $\Gamma(\mathbf{k}, \tau)$  may be chosen to be a realization of a stationary and time-ergodic random process. In that case, long-time averages in the statistical steady-state can be described by averages over ensembles of solutions of the wave kinetic equations with different realizations of the force. In fact, [41] find that in their experiments the forcing is well modelled by an Ornstein-Uhlenbeck process with large fluctuations in the energy input rate. Even without chaos or turbulence in the wave kinetic equations, this will yield “super-randomness”, since the Kolmogorov solution is neutrally stable to change in the energy flux rate [1].

In addition to pumping by an external force, hydrodynamic turbulence may also be (in fact, more commonly, is) driven by boundary conditions. Such a situation can also occur in wave kinetics, as we have discussed in the previous section 4.1, where strongly nonlinear effects such as wave-breaking can provide both sinks and sources to kinetic wave turbulence. These effects restrict the validity of wave kinetics to a “bounded domain” in wavenumbers and amplitudes and could provide suitable forcings at the boundaries to represent those nonlinear effects. In this setting, Kolmogorov cascade solutions which are stable when considered without restriction, might become unstable and “super-turbulence” develop. The two scenarios in this section and the previous one are thus not completely exclusive.

## 5. Conclusions

The major contributions of this paper are the following:

1. We have derived by formal asymptotics the correct multimode equations for wave kinetics in Hamiltonian systems with 3-wave resonances.
2. We have shown that these equations possess factorized solutions for factorized initial conditions, corresponding to “propagation of chaos” or preservation of “random phases & amplitudes”.
3. We introduced the “empirical spectrum” and “empirical 1-mode PDF” and showed that the above factorization implies a law of large numbers, so that these quantities satisfy the wave-kinetic closure equations for nearly every initial realization of random phases and amplitudes, without necessity of averaging.
4. We have demonstrated the close formal relations of wave kinetic theory with the kinetic theory of gases, especially regarding the role of entropy and the 2nd law of thermodynamics in both theories.
5. We have completely classified all realizable solutions of our multimode equations and have shown that they correspond to “super-statistical solutions”, or ensembles of the wave-kinetic closure equations with either random initial conditions or random forces.
6. We have exploited our results to discuss the possibilities to explain intermittency and non-Gaussian statistics of wave turbulence within the kinetic description, in particular by a “super-turbulence” corresponding to chaotic or stochastic solutions of the wave-kinetic closure equations.

Our work solves the problem posed in [42], section 5.2.5 on the continuum limit of finite-dimensional wave turbulence to show “how the natural closure arises in taking the limit  $L \rightarrow \infty$ .” Although we have focused here on 3-wave resonant models, we expect that analogous results should hold also for other closely related wave systems, e.g. those whose first nontrivial resonances are 4-wave.

It is perhaps important to emphasize what we have *not* accomplished in this paper: we have not provided a rigorous mathematical proof of the wave kinetic theory. At this time, no set of sufficient conditions are known that would imply the validity of wave-kinetics for any general class of systems ([42], section 5.2.6). For a recent rigorous proof of validity of wave-kinetic predictions in a 4-wave model problem at thermal equilibrium, see [8].

Our goal here was the more directly physical one of formulating the correct equations of wave kinetics and understanding their content. This should permit a better evaluation of the theory by simulation and experiment. In particular, our most subversive conclusion is that the “mean-field” scaling of the Kolmogorov-Zakharov solutions may not be the true prediction of kinetic theory for scaling exponents of spectra and higher-order statistics in wave turbulence and that wave kinetics may instead allow for intermittency and anomalous scaling.


## Appendix A. Derivation of the Spectral Hierarchy

As discussed in the text, we have verified the results of [4] for the  $\mathcal{J}$ -terms in eqs.(29)-(36), up to minor corrections. The contributions to the spectral generating functional are obtained by making the substitutions  $J_i = (2\pi/L)^d \tilde{J}_i$  and  $\lambda_1 = i\lambda(\mathbf{k}_1)$  and taking the limit  $L \rightarrow \infty$ . The asymptotics of the various terms can be summarized in the following general rule: each action variable  $J$  changes the order by  $L^{-d}$  and each free sum over wavenumbers by  $L^d$ . We illustrate this rule in our detailed calculations below.


*Calculation of  $\mathcal{J}_1$ :* Substituting the definition (23) of  $a_1^{(1)}$ , one obtains

$$\mathcal{J}_1 = \sum_{1,2,3} \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \sqrt{J_1 J_2 J_3} \langle \psi_1^{-1} \psi_2^{\sigma_2} \psi_3^{\sigma_3} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \rangle_{\psi} \Delta_T (\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1) \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3},$$

which can be represented by the following diagram before averaging over phases:



After phase averaging, there are two contributions represented by the following diagrams:



or explicitly written as

$$\begin{aligned} \mathcal{J}_1 &= \sum'_{1,2,3} \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \sqrt{J_1 J_2 J_3} \Delta_T (\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1) \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} \delta_{\mu_1, 1} \delta_{\mu_2, -\sigma_2} \delta_{\mu_3, -\sigma_3} \prod_{m \neq 1, 2, 3} \delta_{\mu_m, 0} \\ &+ \sum'_{1,2} \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_2}^{+, \sigma_2, \sigma_2} \sqrt{J_1 J_2} \Delta_T (2\sigma_2 \omega_2 - \omega_1) \delta_{\mathbf{k}_1, 2\sigma_2 \mathbf{k}_2} \delta_{\mu_1, 1} \delta_{\mu_2, -2\sigma_2} \prod_{m \neq 1, 2} \delta_{\mu_m, 0}. \end{aligned} \quad (\text{A.1})$$

Note that a prime  $\prime$  on a sum indicates that all wavenumbers must be distinct. Here all of the summations are ‘‘pinned.’’ Now making the substitutions  $J_i = (2\pi/L)^d \tilde{J}_i$  and  $\lambda_1 = i\lambda(\mathbf{k}_1)$  into the above and taking the limit  $L \rightarrow \infty$  gives

$$\begin{aligned} \mathcal{J}_1 &= \frac{1}{2} \left( \frac{2\pi}{L} \right)^{d/2} \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \sum_{\underline{\mathbf{k}}} L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \sqrt{\frac{\tilde{J}_2 \tilde{J}_3}{\tilde{J}_1}} \Delta_T (\underline{\sigma} \cdot \omega(\underline{\mathbf{k}})) \delta_{\underline{\sigma}, \underline{\mathbf{k}}, 0} \delta_{\underline{\mu}, -\underline{\sigma}} \prod_{m \neq 1, 2, 3} \delta_{\mu_m, 0} \\ &+ \frac{1}{2} \left( \frac{2\pi}{L} \right)^{d/2} \sum_{\underline{\sigma} = (-1, 2\sigma_2)} \sum_{\underline{\mathbf{k}'}} L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_2}^{+, \sigma_2, \sigma_2} \frac{\tilde{J}_2}{\sqrt{\tilde{J}_1}} \Delta_T (\underline{\sigma} \cdot \omega(\underline{\mathbf{k}'}) ) \delta_{\underline{\sigma}, \underline{\mathbf{k}'}, 0} \delta_{\underline{\mu}, -\underline{\sigma}} \prod_{m \neq 1, 2} \delta_{\mu_m, 0}, \end{aligned} \quad (\text{A.2})$$


where  $\underline{\mu}' = (\mu_1, \mu_2)$  and  $\underline{\mathbf{k}'} = (\mathbf{k}_1, \mathbf{k}_2)$ . Hence  $\mathcal{J}_1$  is  $O(L^{-d/2})$  in the large box limit. In fact, the result easily follows from the aforementioned asymptotic rules. The amplitude factor  $\sqrt{J_1 J_2 J_3}$  changes the order of  $\mathcal{J}_1$  by  $L^{-3d/2}$ . Since

there is no free sum here, the contribution of the term proportional to  $\lambda_1$  is at most  $O(L^{-3d/2})$ . The term proportional to  $\mu_1$  introduces an additional action variable  $1/J_1$  and thus gives the leading contribution  $O(L^{-d/2})$ .


*Calculation of  $\mathcal{J}_2$ :* Again substituting the definition (25) of  $a_1^{(1)}$ , one obtains

$$\mathcal{J}_2 = \frac{1}{2} \sum_{1,2,3,4,5} \left( \lambda_1 + \lambda_1^2 J_1 - \frac{\mu_1^2}{4J_1} \right) L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} L_{\mathbf{k}_1, \mathbf{k}_4, \mathbf{k}_5}^{-, \sigma_4, \sigma_5} \sqrt{J_2 J_3 J_4 J_5} \left\langle \psi_2^{\sigma_2} \psi_3^{\sigma_3} \psi_4^{\sigma_4} \psi_5^{\sigma_5} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \right\rangle_{\psi} \Delta_T(\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1) \Delta_T(\sigma_4 \omega_4 + \sigma_5 \omega_5 + \omega_1) \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} \delta_{-\mathbf{k}_1, \sigma_4 \mathbf{k}_4 + \sigma_5 \mathbf{k}_5}, \quad (\text{A.3})$$


which can be represented by the following diagram before phase-averaging:




After phase averaging,  $\mathcal{J}_2$  has three types of contributions. The leading contributions are two type I diagrams with no external couplings. We show one such diagram here (the other one is obtained by the exchange  $2 \leftrightarrow 3$ ).



These contain three distinct wavenumbers and two vertices, but the wavenumber delta-functions at the two vertices turn out to give the same constraint. Thus the type I diagrams contain two free wavenumbers corresponding to unconstrained sums. Subleading contributions are given by type II diagrams with two wavenumbers externally coupled. We show one diagram here with the wavenumbers 2 and 4 coupled to distinct external blobs. The three others can be obtained by exchanging  $4 \leftrightarrow 5$  and  $2 \leftrightarrow 3$ .




These subleading contributions contain only one free wavenumber. The two delta functions  $\delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3}$  and  $\delta_{-\mathbf{k}_1, \sigma_4 \mathbf{k}_4 + \sigma_5 \mathbf{k}_5}$  along with the internal connection  $\delta_{\mathbf{k}_3, \mathbf{k}_5} \delta_{\sigma_3, -\sigma_5}$  require  $\sigma_2 \mathbf{k}_2 = -\sigma_4 \mathbf{k}_4$ . Therefore,  $\mathbf{k}_2 = -\mathbf{k}_4$  (since  $\mathbf{k}_1$  and  $\mathbf{k}_4$  must be distinct) and  $\sigma_2 = \sigma_4$  in this diagram. In principle, another type II contribution could exist with two wavenumbers coupled to the same external blob, as follows:



together with diagrams resulting from  $4 \leftrightarrow 5$  and  $2 \leftrightarrow 3$ . However, the condition  $\sigma_2 \mathbf{k}_2 = -\sigma_4 \mathbf{k}_4$  in this case requires that  $\mathbf{k}_2 = \mathbf{k}_4$ ,  $\sigma_2 = -\sigma_4$ , and  $\mu_2 = 0$ , which coincides with the type I diagram.

All other contributions are type III where all wavenumbers are pinned, for example,



Now consider large  $L$ . The amplitude factor  $\sqrt{J_2 J_3 J_4 J_5}$  changes the order by  $L^{-2d}$ . Now consider the contributions proportional to the three terms in the prefactor  $\lambda_1 + \lambda_1^2 J_1 - \frac{\mu_1^2}{4J_1}$ . In the term proportional to  $\lambda_1$  the type I diagrams provide two free sums that increase the order by  $L^{2d}$ , giving an  $O(1)$  contribution. The next order contributions from type II diagrams have only one free wavenumber and thus are at most  $O(L^{-d})$ . The term proportional to  $\lambda_1^2 J_1$  is at most  $O(L^{-d})$  because of the additional action variable  $J_1$ . For the term proportional to  $\frac{\mu_1^2}{4J_1}$ , we note for both type I and type II diagrams  $\mathbf{k}_1$  is not pinned, which requires  $\mu_1$  to be zero. Thus the only nonzero contribution here must come from type III graphs, which is at most  $O(L^{-d})$ . Therefore we can conclude that the leading contributions in  $\mathcal{J}_1$  are  $O(1)$ , or explicitly

$$\mathcal{J}_2 \sim 9\delta_{\mu,0} \sum_{1,2,3} \lambda_1 J_2 J_3 |H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3}|^2 |\Delta_T(\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1)|^2 \quad (\text{A.4})$$

Substituting  $J_i = (2\pi/L)^d \tilde{J}_i$  and  $\lambda_1 = i\lambda(\mathbf{k}_1)$  and taking the limit  $L \rightarrow \infty$  using (35), one finds

$$\begin{aligned} \langle e^{\sum \mathbf{k} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \mathcal{J}_2 \rangle_J &\sim -9i\delta_{\mu,0} \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) |\Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}))|^2 \\ &\quad \times \lambda(\mathbf{k}_1) |H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3}|^2 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_2) \delta \lambda(\mathbf{k}_3)} \end{aligned} \quad (\text{A.5})$$

It is worth mentioning here that the term proportional to  $\lambda_1^2$  kept in [4] gives only an  $O(L^{-d})$  contribution in the large box limit, which can be calculated exactly as

$$9i \left(\frac{2\pi}{L}\right)^d \delta_{\mu,0} \sum_{\underline{\sigma}=(-1, \sigma_2, \sigma_3)} \int d^d k_1 d^d k_2 d^d k_3 \lambda^2(\mathbf{k}_1) \frac{\delta^3 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_2) \delta \lambda(\mathbf{k}_3)} |H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3}|^2 |\Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}))|^2 \delta^d(\underline{\sigma} \cdot \mathbf{k}), \quad (\text{A.6})$$

which is even smaller than the  $O(L^{-d/2})$  correction from  $\mathcal{J}_1$  which we calculated before.


*Calculation of  $\mathcal{J}_3$ :* Substituting the definition (26) of  $a_1^{(2)}$  we have

$$\mathcal{J}_3 = \sum_{1,2,3,4,5} \left( \lambda_1 + \frac{\mu_1}{2J_1} \right) L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} L_{\mathbf{k}_2, \mathbf{k}_4, \mathbf{k}_5}^{\sigma_2, \sigma_4, \sigma_5} \sqrt{J_1 J_3 J_4 J_5} \langle \psi_1^{-1} \psi_3^{\sigma_3} \psi_4^{\sigma_4} \psi_5^{\sigma_5} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \rangle_{\psi} \quad (\text{A.7})$$


$$E_T(\omega_{345}^1, \omega_{23}^1) \delta_{23}^1 \delta_{45}^2 \Big|_{\sigma_1=1} + (2 \leftrightarrow 3). \quad (\text{A.8})$$

By symmetry, we only need to consider the first part, which has the following diagram pre-phase-averaged:






After phase averaging, there are three types of contributions. The type I diagrams are two leading contributions which contain two free wavenumbers. We show only one diagram here, the other obtained by exchanging  $4 \leftrightarrow 5$



The type II diagrams are four subleading contributions which contain one free wavenumber. Again we only show one diagram here, with mode 1 externally coupled:



It turns out that the graphs with 1 externally coupled are the most important type II contributions to  $\mathcal{J}_3$ . The delta functions require  $\mathbf{k}_4 = -\mathbf{k}_1$  and  $\sigma_4 = -1$  in this diagram. Additional type II graphs can be obtained by symmetry. All other contributions are type III where all wavenumbers are pinned.

Now consider large  $L$ . The amplitude factor  $\sqrt{J_1 J_3 J_4 J_5}$  changes the order of  $\mathcal{J}_3$  by  $L^{-2d}$ . In the term proportional to  $\lambda_1$ , the type I diagrams increase the order by  $L^{2d}$  since they contain two free sums, giving an  $O(1)$  contribution to  $\mathcal{J}_1$ . In the term proportional to  $\frac{\mu_1}{2J_1}$ , the type I diagrams give zero contribution because they require  $\mu_1$  to vanish. The leading  $O(1)$  contribution now comes from the type II diagrams with mode 1 externally coupled and  $\mu_1 = 1$ . The type III diagrams contain no free wavenumber and give contributions at most  $O(L^{-d})$ . The leading contribution to  $\mathcal{J}_3$  in the large-box limit is thus:

$$\begin{aligned}
\mathcal{J}_3 \sim & 2 \sum_{\sigma_2, \sigma_3} \prod_{\mathbf{k}} \delta_{\mu_{\mathbf{k}}, 0} \sum_{1,2,3} \lambda_1 L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} L_{\mathbf{k}_2, \mathbf{k}_1, \mathbf{k}_3}^{\sigma_2, -1, -\sigma_3} J_1 J_3 E_T(0, \sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1) \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} \\
& + \sum_{\sigma_2, \sigma_3} \sum_{1,2,3} \delta_{\mu_1, 1} \delta_{\mu_{-1}, 1} \prod_{\mathbf{k} \neq \mathbf{k}_1, -\mathbf{k}_1} \delta_{\mu_{\mathbf{k}}, 0} \frac{\mu_1}{2J_1} L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} L_{\mathbf{k}_2, -\mathbf{k}_1, \mathbf{k}_3}^{\sigma_2, -1, -\sigma_3} \sqrt{J_1 J_{-1} J_3} \\
& \times E_T(\omega_1 + \omega_{-1}, \omega_1 - \sigma_2 \omega_2 - \sigma_3 \omega_3) \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} + (2 \leftrightarrow 3)
\end{aligned} \tag{A.9}$$

Substituting  $J_i = (2\pi/L)^d \tilde{J}_i$  and  $\lambda_1 = i\lambda(\mathbf{k}_1)$  and taking the limit  $L \rightarrow \infty$  using (35), one finds


$$\langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \mathcal{J}_3 \rangle_J \sim 18i \delta_{\mu, 0} \sum_{\sigma = (-1, \sigma_2, \sigma_3)} \sigma_2 \int d^d k_1 \int d^d k_2 \int d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) E_T(\underline{\sigma} \cdot \omega(\mathbf{k}), 0)$$

$$\begin{aligned}
& \lambda(\mathbf{k}_1) |H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3}|^2 \frac{\delta^2 \mathcal{Z}}{\delta \lambda(\mathbf{k}_1) \delta \lambda(\mathbf{k}_3)} \\
& -9 \sum_1 \delta_{\mu_1, 1} \delta_{\mu_{-1}, 1} \prod_{\mathbf{k} \neq \mathbf{k}_1, -\mathbf{k}_1} \delta_{\mu_{\mathbf{k}}, 0} \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \sigma_2 \int d^d k_2 \int d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) E_T(\omega(\mathbf{k}_1) + \omega(-\mathbf{k}_1), \underline{\sigma} \cdot \omega(\mathbf{k})) \\
& H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3} H_{-\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \left\langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \sqrt{\frac{\tilde{J}_{-1}}{\tilde{J}_1} \tilde{J}_3} \right\rangle_J + (2 \leftrightarrow 3). \tag{A.10}
\end{aligned}$$


Calculation of  $\mathcal{J}_4$ : Similarly, substituting (25) for  $a_1^{(1)}$

$$\begin{aligned}
\mathcal{J}_4 = & \sum_{1,2,3,4,5} \left( \frac{1}{2} \lambda_1^2 + \frac{\mu_1}{4J_1^2} (\frac{\mu_1}{2} - 1) + \frac{\lambda_1 \mu_1}{2J_1} \right) L_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} L_{\mathbf{k}_1, \mathbf{k}_4, \mathbf{k}_5}^{+, \sigma_4, \sigma_5} J_1 \sqrt{J_2 J_3 J_4 J_5} \left\langle \psi_1^{-2} \psi_2^{\sigma_2} \psi_3^{\sigma_3} \psi_4^{\sigma_4} \psi_5^{\sigma_5} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \right\rangle_{\psi} \\
& \times \Delta_T(\sigma_2 \omega_2 + \sigma_3 \omega_3 - \omega_1) \Delta_T(\sigma_4 \omega_4 + \sigma_5 \omega_5 - \omega_1) \delta_{\mathbf{k}_1, \sigma_2 \mathbf{k}_2 + \sigma_3 \mathbf{k}_3} \delta_{\mathbf{k}_1, \sigma_4 \mathbf{k}_4 + \sigma_5 \mathbf{k}_5}, \tag{A.11}
\end{aligned}$$

which can be represented by the following diagram before phase-averaging:



After phase averaging, there are two types of contributions. Type I diagrams have only the 1 mode externally coupled, corresponding to the following diagram and its partner with  $2 \leftrightarrow 3$ :



These are formally the leading contributions, with one free wavenumber. However, the wavenumber delta functions require  $\mathbf{k}_1 = \mathbf{0}$  and thus this graph vanishes identically. All other nonzero contributions are type II where there are no free wavenumbers.


Now consider large  $L$ . The amplitude factor  $J_1 \sqrt{J_2 J_3 J_4 J_5}$  in  $\mathcal{J}_4$  changes the order by  $L^{-3d}$ . The largest possible contribution comes from the term proportional to  $\frac{\mu_1}{4J_1^2} (\frac{\mu_1}{2} - 1)$ , which is a large factor of  $O(L^{2d})$ . However, such contributions are still only  $O(L^{-d})$ . Hence  $\mathcal{J}_4$  is negligible in the large box limit.

Calculation of  $\mathcal{J}_5$ : Write  $\mathcal{J}_5 \equiv \sum_{1 \neq 2} \left\{ \lambda_1 \lambda_2 (\mathcal{B}_1 + \mathcal{B}_2) + (\lambda_1 + \frac{\mu_1}{4J_1}) \frac{\mu_2}{J_2} (\mathcal{B}_1 - \mathcal{B}_3) \right\}$ , where (25) gives


$$\begin{aligned}
\mathcal{B}_1 & \equiv \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{(0)\mu_{\mathbf{k}}} a_1^{(1)} a_1^{(0)*} a_2^{(1)} a_2^{(0)*} \right\rangle_{\psi} = \sum_{3,4,5,6} L_{\mathbf{k}_1, \mathbf{k}_3, \mathbf{k}_4}^{+, \sigma_3, \sigma_4} L_{\mathbf{k}_2, \mathbf{k}_5, \mathbf{k}_6}^{+, \sigma_5, \sigma_6} \sqrt{J_1 J_2 J_3 J_4 J_5 J_6} \left\langle \psi_1^{-1} \psi_2^{-1} \psi_3^{\sigma_3} \psi_4^{\sigma_4} \psi_5^{\sigma_5} \psi_6^{\sigma_6} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \right\rangle_{\psi} \\
& \quad \times \Delta(\sigma_3 \omega_3 + \sigma_4 \omega_4 - \omega_1) \Delta(\sigma_5 \omega_5 + \sigma_6 \omega_6 - \omega_2) \delta_{\mathbf{k}_1, \sigma_3 \mathbf{k}_3 + \sigma_4 \mathbf{k}_4} \delta_{\mathbf{k}_2, \sigma_5 \mathbf{k}_5 + \sigma_6 \mathbf{k}_6} \\
\mathcal{B}_2 & \equiv \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} a_1^{(1)*} a_1^{(0)} a_2^{(1)} a_2^{(0)*} \right\rangle_{\psi} = \sum_{3,4,5,6} L_{\mathbf{k}_1, \mathbf{k}_3, \mathbf{k}_4}^{-, \sigma_3, \sigma_4} L_{\mathbf{k}_2, \mathbf{k}_5, \mathbf{k}_6}^{+, \sigma_5, \sigma_6} \sqrt{J_1 J_2 J_3 J_4 J_5 J_6} \left\langle \psi_1^{+1} \psi_2^{-1} \psi_3^{\sigma_3} \psi_4^{\sigma_4} \psi_5^{\sigma_5} \psi_6^{\sigma_6} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \right\rangle_{\psi} \\
& \quad \times \Delta(\sigma_3 \omega_3 + \sigma_4 \omega_4 + \omega_1) \Delta(\sigma_5 \omega_5 + \sigma_6 \omega_6 - \omega_2) \delta_{-\mathbf{k}_1, \sigma_3 \mathbf{k}_3 + \sigma_4 \mathbf{k}_4} \delta_{\mathbf{k}_2, \sigma_5 \mathbf{k}_5 + \sigma_6 \mathbf{k}_6}
\end{aligned}$$

$$\mathcal{B}_3 \equiv \left\langle \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} a_1^{(1)} a_1^{(0)*} a_2^{(1)*} a_2^{(0)} \right\rangle_{\psi} = \sum_{3,4,5,6} L_{\mathbf{k}_1, \mathbf{k}_3, \mathbf{k}_4}^{+, \sigma_3, \sigma_4} L_{\mathbf{k}_2, \mathbf{k}_5, \mathbf{k}_6}^{-, \sigma_5, \sigma_6} \sqrt{J_1 J_2 J_3 J_4 J_5 J_6} \left\langle \psi_1^{-1} \psi_2^{+1} \psi_3^{\sigma_3} \psi_4^{\sigma_4} \psi_5^{\sigma_5} \psi_6^{\sigma_6} \prod_{\mathbf{k}} \psi_{\mathbf{k}}^{\mu_{\mathbf{k}}} \right\rangle_{\psi} \times \Delta(\sigma_3 \omega_3 + \sigma_4 \omega_4 - \omega_1) \Delta(\sigma_5 \omega_5 + \sigma_6 \omega_6 + \omega_2) \delta_{\mathbf{k}_1, \sigma_3 \mathbf{k}_3 + \sigma_4 \mathbf{k}_4} \delta_{-\mathbf{k}_2, \sigma_5 \mathbf{k}_5 + \sigma_6 \mathbf{k}_6}. \quad (\text{A.12})$$


Note under the interchanges  $1 \leftrightarrow 2, 3 \leftrightarrow 5, 4 \leftrightarrow 6$ , that  $\mathcal{B}_2 \leftrightarrow \mathcal{B}_3$ . All three of the  $\mathcal{B}$ 's contain the same amplitude factor  $\sqrt{J_1 J_2 J_3 J_4 J_5 J_6}$  that changes the order by  $L^{-3d}$ . We list their diagrams as follows,



After phase averaging, there are three types of diagrams for the  $\mathcal{B}$ 's. The type I diagrams are leading contributions with no external couplings. Shown here is one type I diagram for  $\mathcal{B}_1$  :



with others obtained by  $3 \leftrightarrow 4$  and  $5 \leftrightarrow 6$ . These contain two free wavenumbers. For  $\mathcal{B}_2, \mathcal{B}_3$  there are additional type I contributions with 1 and 2 coupled, but these do not contribute to  $\mathcal{J}_5$  because of the restriction that  $1 \neq 2$ . The type II diagrams are subleading contributions which contain one free wavenumber. Shown here is one such contribution for  $\mathcal{B}_1$  with both 1 and 2 externally coupled:



The delta functions here require  $\mathbf{k}_2 = -\mathbf{k}_1$ . In the corresponding diagrams for  $\mathcal{B}_2, \mathcal{B}_3$  the delta functions instead require  $\mathbf{k}_1 = \mathbf{k}_2$ , so these do not exist. (Contributions to  $\mathcal{B}_2, \mathcal{B}_3$  with 1 and 2 coupled to the *same* external blob coincide with the type I diagrams that do not contribute to  $\mathcal{J}_5$ .) There are also type II contributions for the  $\mathcal{B}$ 's with only one of 1 or 2 externally coupled. For example, one such contribution is obtained for  $\mathcal{B}_1$  by exchanging  $1 \leftrightarrow 4$  in the above graph and then the delta functions require that  $\mathbf{k}_2 = \sigma_4 \mathbf{k}_4$  so that  $\mathbf{k}_2 = -\mathbf{k}_4, \sigma_4 = -1$ . Similar type II contributions exist for  $\mathcal{B}_2, \mathcal{B}_3$ . Other type II contributions with neither 1 nor 2 externally coupled must have both  $\mu_1 = \mu_2 = 0$  and do not contribute at non-vanishing order to  $\mathcal{J}_5$ . Type III contributions with additional external couplings have no free wavenumbers.

Now consider the size of the various terms for large  $L$ . For  $\sum_{1 \neq 2} \lambda_1 \lambda_2 (\mathcal{B}_1 + \mathcal{B}_2)$ , the leading contribution comes from type I diagrams and is  $O(L^{-d})$ . For  $\sum_{1 \neq 2} \frac{\lambda_1 \mu_2}{J_2} (\mathcal{B}_1 - \mathcal{B}_3)$ , the type I diagrams give zero contribution since  $\mu_2 = 0$

and type II diagrams are at most  $O(L^{-d})$ . For  $\sum_{1 \neq 2} \frac{\mu_1 \mu_2}{4J_1 J_2} \mathcal{B}_1$ , type I diagrams give zero contribution and type II diagrams with both 1 and 2 externally coupled give an  $O(1)$  contribution. The corresponding term  $\sum_{1 \neq 2} \frac{\mu_1 \mu_2}{4J_1 J_2} \mathcal{B}_3 = O(L^{-d})$  from type III diagrams, since, as noted above, there do not exist type II diagrams for  $\mathcal{B}_3$  with 1 and 2 externally coupled to distinct blobs. In large box limit, the term  $\mathcal{J}_5$  is given, to leading order, by

$$\begin{aligned} \mathcal{J}_5 \sim & -\frac{9}{2} \sum_1 \delta_{\mu_1,1} \delta_{\mu_{-1},1} \prod_{\mathbf{k} \neq \pm \mathbf{k}_1} \delta_{\mu_{\mathbf{k}},0} \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d k_2 \int d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k})) \Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}')) \\ & \times H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3} H_{-\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \left\langle \frac{\tilde{J}_2 \tilde{J}_3}{\sqrt{\tilde{J}_1 \tilde{J}_{-1}}} e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \right\rangle_J, \end{aligned} \quad (\text{A.13})$$

where  $\underline{\mathbf{k}} = (\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$  and  $\underline{\mathbf{k}}' = (-\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$ . Now making the substitution  $\lambda_{\mathbf{k}} = i\lambda(\mathbf{k})$  and taking the limit, one obtains,

$$\begin{aligned} \left\langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \mathcal{J}_5 \right\rangle_J \sim & -\frac{9}{2} \sum_1 \delta_{\mu_1,1} \delta_{\mu_{-1},1} \prod_{\mathbf{k} \neq \pm \mathbf{k}_1} \delta_{\mu_{\mathbf{k}},0} \sum_{\underline{\sigma} = (-, \sigma_2, \sigma_3)} \int d^d k_2 d^d k_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}) \Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k})) \Delta_T(-\underline{\sigma} \cdot \omega(\mathbf{k}')) \\ & \times H_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{-, \sigma_2, \sigma_3} H_{-\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3}^{+, \sigma_2, \sigma_3} \left\langle \frac{\tilde{J}_2 \tilde{J}_3}{\sqrt{\tilde{J}_1 \tilde{J}_{-1}}} e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \right\rangle_J. \end{aligned} \quad (\text{A.14})$$

## Appendix B. Derivation of the PDF Hierarchy

We here evaluate the  $\mathcal{J}$ -terms that contribute to  $\mathcal{X}^{(M)}(\lambda, \mu, T)$ . Most of the essential work has already been done in Appendix A. As discussed in the text, the main difference is that mode 1 is now discrete for  $\mathcal{J}_1 - \mathcal{J}_4$  and assumes only  $M$  values, while for  $\mathcal{J}_5$  modes 1,2 are both discrete. All other modes are continuous in the infinite-box limit. One must consider carefully whether free wavenumbers in graphical summations are discrete or continuous to see whether their contribution is  $O(M)$  or  $O(L^d)$ .

$\mathcal{J}_1$ : The graphs contain no free wavenumbers, so that  $\mathcal{J}_1$  is still  $O(L^{-d/2})$ .

$\mathcal{J}_2$ : The leading order  $O(1)$  contribution comes again from the type I diagrams, which contain two free wavenumbers. One of these free wavenumbers is mode 1 which is discrete, but this is compensated by the extra factor of  $L^d$  from the prefactors  $\lambda_{\mathbf{k}_1}$  and  $\lambda_{\mathbf{k}_1}^2 J_{\mathbf{k}_1}$ , which now contribute equally. One thus obtains by substituting  $\lambda_{\mathbf{k}_m} = i(L/2\pi)^d \lambda_m$  and  $J_m = (2\pi/L)^d \tilde{J}_m$  the result

$$\begin{aligned} \left\langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \mathcal{J}_2 \right\rangle_J & \sim 9\delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) |\Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}_j))|^2 |H_{\mathbf{k}_j}^{\underline{\sigma}}|^2 \\ & \quad \times \left\langle (i\lambda_j - \lambda_j^2 \tilde{J}_j) \tilde{J}_2 \tilde{J}_3 e^{\sum_m i\lambda_m \tilde{J}_m} \right\rangle_J \\ & = -9i\delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) |\Delta_T(\underline{\sigma} \cdot \omega(\mathbf{k}_j))|^2 |H_{\mathbf{k}_j}^{\underline{\sigma}}|^2 \\ & \quad \times (\lambda_j + \lambda_j^2 \frac{\partial}{\partial \lambda_j}) \frac{\partial^2 \mathcal{Z}^{(M+2)}}{\partial \bar{\lambda}_2 \partial \bar{\lambda}_3} \Big|_{\bar{\lambda}_2 = \bar{\lambda}_3 = 0}. \end{aligned} \quad (\text{B.1})$$

Here  $\underline{\mathbf{k}}_j = (\mathbf{k}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3)$ .

$\mathcal{J}_3$ : There are two contributions, from the two terms in the prefactor,  $\lambda_{\mathbf{k}_1}$  and  $\mu_{\mathbf{k}_1}/2J_{\mathbf{k}_1}$ . For the  $\lambda_{\mathbf{k}_1}$  term, the leading  $O(1)$  contribution comes from the type I diagrams, just as for  $\mathcal{J}_2$ . One of the two free wavenumbers is discrete mode 1, but this is compensated by the  $O(L^d)$  prefactor  $\lambda_{\mathbf{k}_1}$ . For the  $\mu_{\mathbf{k}_1}/2J_{\mathbf{k}_1}$  term, the leading  $O(1)$  contribution comes from the type II diagrams with modes 1 and  $-1$  externally coupled, so that  $\mu_{\mathbf{k}_1} = \mu_{-\mathbf{k}_1} = 1$ . The type II diagram has one free continuous wavenumber. Thus, its contribution is also  $O(1)$  if the set of  $M$  wavenumbers includes at least one pair of opposite wavenumbers. Assuming this is not so, the final result is:

$$\left\langle e^{\sum_{\mathbf{k}} \lambda_{\mathbf{k}} J_{\mathbf{k}}} \mathcal{J}_3 \right\rangle_J^{(d)} \sim -18\delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma} = (-1, \sigma_2, \sigma_3)} \sigma_2 \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \mathbf{k}_j) E_T(0, \underline{\sigma} \cdot \omega(\mathbf{k}_j)) |H_{\mathbf{k}_j}^{\underline{\sigma}}|^2$$

$$\begin{aligned}
& \times \langle i\lambda_j \tilde{J}_j \tilde{J}_3 e^{\sum_m i\lambda_m \tilde{J}_m} \rangle_J + (2 \leftrightarrow 3). \\
= & 18i\delta_{\mu,0} \sum_{j=1}^M \sum_{\underline{\sigma}=(-1,\sigma_2,\sigma_3)} \sigma_2 \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}_j) E_T(0, \underline{\sigma} \cdot \omega(\underline{\mathbf{k}}_j)) \\
& \times \lambda_j |H_{\underline{\mathbf{k}}_j}^\sigma|^2 \frac{\partial^2 \mathcal{Z}^{(M+1)}}{\partial \bar{\lambda}_3 \partial \lambda_j} \Big|_{\bar{\lambda}_3=0} + (2 \leftrightarrow 3). \tag{B.2}
\end{aligned}$$

However, if the  $M$  wavenumbers include pairs of opposites, then there is an additional contribution

$$\begin{aligned}
\langle e^{\sum_k \lambda_k J_k} \mathcal{J}_3 \rangle_J^{(II)} & \sim -9 \sum_{j=1}^M \delta_{\mu_j,1} \delta_{\mu_{-j},1} \prod_{m \neq j,-j} \delta_{\mu_m,0} \sum_{\underline{\sigma}=(-1,\sigma_2,\sigma_3)} \sigma_2 \int d^d \bar{k}_2 \int d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}_j) \\
& \times E_T(\omega(\underline{\mathbf{k}}_j) + \omega(-\underline{\mathbf{k}}_j), \underline{\sigma} \cdot \omega(\underline{\mathbf{k}}_j)) H_{\underline{\mathbf{k}}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{-,\sigma_2,\sigma_3} H_{-\underline{\mathbf{k}}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{+,\sigma_2,\sigma_3} \left\langle e^{\sum_m i\lambda_m \tilde{J}_m} \sqrt{\frac{\tilde{J}_{-1}}{\tilde{J}_1}} \tilde{J}_3 \right\rangle_J \\
& + (2 \leftrightarrow 3). \tag{B.3}
\end{aligned}$$

$\mathcal{J}_4$ : The type I contribution vanishes and the leading contribution from type II diagrams with no free sums remains  $O(L^{-d})$ .

$\mathcal{J}_5$ : The amplitude factors contribute  $O(L^{-3d})$  while the prefactors are  $O(L^{2d})$ . Thus, to give an overall  $O(1)$  contribution, the  $\mathcal{B}$  factors must contain a free continuous wavenumber. For  $\mathcal{B}_1$  the type I diagrams have two free discrete wavenumbers and type II diagrams with mode 2 only pinned have one free discrete wavenumber, but type II diagrams with both 1 and 2 pinned have one free continuous wavenumber. For  $\mathcal{B}_2, \mathcal{B}_3$ , there are no graphs with modes 1 and 2 distinct, contributing to  $\mathcal{J}_5$ , that have a free continuous wavenumber. Type I diagrams have two free wavenumbers, both discrete if 1 and 2 are uncoupled. Type II diagrams with only one of the modes 1 or 2 externally coupled have one discrete free wavenumber.

Thus, the only  $O(1)$  contribution to  $\mathcal{J}_5$  is from  $\sum_{1 \neq 2} \left\{ \lambda_{\mathbf{k}_1} \lambda_{\mathbf{k}_2} + \frac{\lambda_{\mathbf{k}_1} \mu_{\mathbf{k}_2}}{J_{\mathbf{k}_2}} + \frac{\mu_{\mathbf{k}_1} \mu_{\mathbf{k}_2}}{4J_{\mathbf{k}_1} J_{\mathbf{k}_2}} \right\} \mathcal{B}_1$  and type II diagrams for  $\mathcal{B}_1$  with both 1 and 2 pinned. Corrections are at most  $O(M/L^d)$ . The  $O(1)$  contribution is thus like that obtained before for  $\mathcal{J}_5$ , except that the mode-1 sum is over just  $M$  discrete values and all terms in the prefactor contribute equally. This gives

$$\begin{aligned}
\langle e^{\sum_k \lambda_k J_k} \mathcal{J}_5 \rangle_J & \sim 18 \sum_{j=1}^M \delta_{\mu_j,1} \delta_{\mu_{-j},1} \prod_{m \neq 1,-1} \delta_{\mu_m,0} \left( \lambda_1 \lambda_{-1} - i \frac{\lambda_1}{\tilde{J}_{-1}} - \frac{1}{4\tilde{J}_1 \tilde{J}_{-1}} \right) \sum_{\underline{\sigma}=(-,\sigma_2,\sigma_3)} \int d^d \bar{k}_2 d^d \bar{k}_3 \delta^d(\underline{\sigma} \cdot \underline{\mathbf{k}}_j) \\
& \times \Delta_T(\underline{\sigma} \cdot \omega(\underline{\mathbf{k}}_j)) \Delta_T(-\underline{\sigma} \cdot \omega(\underline{\mathbf{k}}'_j)) H_{\underline{\mathbf{k}}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{-,\sigma_2,\sigma_3} H_{-\underline{\mathbf{k}}_j, \bar{\mathbf{k}}_2, \bar{\mathbf{k}}_3}^{+,\sigma_2,\sigma_3} \left\langle \sqrt{\tilde{J}_1 \tilde{J}_{-1} \tilde{J}_2 \tilde{J}_3} e^{i \sum_m \lambda_m \tilde{J}_m} \right\rangle_J. \tag{B.4}
\end{aligned}$$

## References

- [1] V. E. Zakharov, V. S. L'vov, G. Falkovich, Kolmogorov spectra of turbulence I. Wave turbulence, Springer series in nonlinear dynamics, Springer Berlin, 1992. ISBN 3540545336.
- [2] P. Jakobsen, A. C. Newell, Invariant measures and entropy production in wave turbulence, J. Stat. Mech.-Theory E. 2004 (2004) L10002.
- [3] Y. Choi, Y. V. Lvov, S. Nazarenko, B. Pokorni, Anomalous probability of large amplitudes in wave turbulence, Phys. Lett. A 339 (2005) 361–369.
- [4] Y. Choi, Y. V. Lvov, S. Nazarenko, Joint statistics of amplitudes and phases in wave turbulence, Physica D 201 (2005) 121–149.
- [5] G. M. Zaslavskii, R. Z. Sagdeev, Limits of statistical description of a nonlinear wave field, Sov. Phys. JETP 25 (1967) 718–724.
- [6] R. Peierls, Zur kinetischen Theorie der Wärmeleitung in Kristallen, Ann. Phys. 395 (1929) 1055–1101.
- [7] R. Brout, I. Prigogine, Statistical mechanics of irreversible processes Part VIII: general theory of weakly coupled systems, Physica 22 (1956) 621–636.
- [8] J. Lukkarinen, H. Spohn, Weakly nonlinear Schrödinger equation with random initial data, Invent. Math. 183 (2011) 79–188.
- [9] S. Nazarenko, S. Lukaschuk, S. McLelland, P. Denissenko, Statistics of surface gravity wave turbulence in the space and time domains, J. Fluid Mech. 642 (2010) 395–420.
- [10] E. Falcon, S. Fauve, C. Laroche, Observation of intermittency in wave turbulence, Phys. Rev. Lett. 98 (2007) 154501.
- [11] N. Yokoyama, Statistics of gravity waves obtained by direct numerical simulation, J. Fluid Mech. 501 (2004) 169–178.
- [12] C. Connaughton, S. Nazarenko, A. C. Newell, Dimensional analysis and weak turbulence, Physica D 184 (2003) 86–97.
- [13] Y. L. Klimontovich, The Statistical Theory of Non-Equilibrium Processes in a Plasma, International series of monographs in natural philosophy, Pergamon Press, 1967.

- [14] O. E. Lanford, Time evolution of large classical systems, in: J. Moser (Ed.), *Dynamical Systems, Theory and Applications*, volume 38 of *Lecture Notes in Physics*, Springer Berlin / Heidelberg, 1975, pp. 1–111.
- [15] O. E. Lanford, On the derivation of the Boltzmann equation, *Asterisque* 40 (1976) 117–137.
- [16] H. Spohn, Boltzmann hierarchy and Boltzmann equation, in: C. Cercignani (Ed.), *Kinetic Theories and the Boltzmann Equation*, volume 1048 of *Lecture Notes in Mathematics*, Springer Berlin / Heidelberg, 1984, pp. 207–220.
- [17] Y. Choi, S. G. Jo, H. I. Kim, S. V. Nazarenko, Aspects of two-mode probability density function in weak wave turbulence, *J. Phys. Soc. Jap.* 78 (2009) 084403.
- [18] H. P. McKean, A class of Markov processes associated with nonlinear parabolic equations, *P. Natl. Acad. Sci. USA* 56 (1966) 1907–1911.
- [19] O. Kurbanmuradov, Convergence of numerical models for the Gaussian fields., *Russ. J. Numer. Anal. M.* 10 (1995) 311–323.
- [20] D. J. Benney, A. C. Newell, Random wave closures, *Stud. Appl. Math.* 48 (1969) 29–53.
- [21] V. E. Zakharov, V. S. L'vov, Statistical description of nonlinear wave fields, *Radiophys. Quantum El.* 18 (1975) 1084–1097.
- [22] H. Spohn, The phonon Boltzmann equation, properties and link to weakly anharmonic lattice dynamics, *J. Stat. Phys.* 124 (2006) 1041–1104.
- [23] L. Boltzmann, Weitere Studien über das Wärmegleichgewicht unter Gasmolekülen, *Sitzungsberichte Akad. Wiss.* 66 (1872) 275–370.
- [24] Y. Yamasaki, Measures on infinite dimensional spaces, volume 5 of *Pure Mathematics*, Singapore Philadelphia: World Scientific, 1985. ISBN 9971978520.
- [25] G. L. Eyink, J. Xin, Self-similar decay in the Kraichnan model of a passive scalar, *J. Stat. Phys.* 100 (2000) 679–741.
- [26] R. H. Kraichnan, Convergents to turbulence functions, *J. Fluid Mech.* 41 (1970) 189–217.
- [27] C. E. Leith, Atmospheric predictability and two-dimensional turbulence, *J. Atmos. Sci.* 28 (1971) 145–161.
- [28] T. M. Cover, J. A. Thomas, *Elements of Information Theory*, Wiley Series in Telecommunications and Signal Processing, John Wiley & Sons, 1991. ISBN 0471241954.
- [29] N. Goldenfeld, *Lectures on Phase Transitions and the Renormalization Group*, volume 85 of *Frontiers in Physics*, Addison-Wesley, Advanced Book Program, 1992. ISBN 0201554089.
- [30] R. Esposito, R. Marra, J. L. Lebowitz, Solutions to the Boltzmann equation in the Boussinesq regime, *J. Stat. Phys.* 90 (1998) 1129–1178.
- [31] W. C. Griffith, Shock waves, *J. Fluid Mech.* 106 (1981) 81–101.
- [32] L. Biferale, Shell models of energy cascade in turbulence, *Annu. Rev. Fluid. Mech.* 35 (2003) 441–468.
- [33] L. Biferale, A. Lambert, R. Lima, G. Paladin, Transition to chaos in a shell model of turbulence, *Physica D* 80 (1995) 105 – 119.
- [34] L. Kadanoff, D. Lohse, N. Schörghofer, Scaling and linear response in the GOY turbulence model, *Physica D* 100 (1997) 165–186.
- [35] G. E. Falkovich, A. V. Shafarenko, On the stability of Kolmogorov spectra of a weak turbulence, *Physica D* 27 (1987) 399–411.
- [36] A. M. Balk, V. E. Zakharov, Stability of slightly turbulent Kolmogorov spectra, *Sov. Phys. Dokl.* 33 (1988) 270–272.
- [37] A. M. Balk, V. E. Zakharov, Stability of weak turbulence Kolmogorov spectra, in: *Plasma theory and nonlinear and turbulent processes in physics*, Proc. Int. Workshop, Kiev/USSR 1987, volume 1, World Scientific Publishing Co., Singapore, 1988, pp. 359–376.
- [38] A. M. Balk, V. E. Zakharov, Stability of weak-turbulence Kolmogorov spectra, *Amer. Math. Soc. Transl. Ser. 2* 182 (1998) 31–82.
- [39] P. J. Schmid, D. S. Henningson, *Stability and Transition in Shear Flows*, volume 142 of *Applied Mathematical Sciences*, Springer, 2001. ISBN 0387989854.
- [40] B. Dubrulle, J.-P. Zahn, Nonlinear instability of viscous plane Couette flow Part 1. Analytical approach to a necessary condition, *J. Fluid Mech.* 231 (1991) 561–573.
- [41] E. Falcon, S. Aumaître, C. Falcón, C. Laroche, S. Fauve, Fluctuations of energy flux in wave turbulence, *Phys. Rev. Lett.* 100 (2008) 064503.
- [42] A. C. Newell, B. Rumpf, Wave turbulence, *Annu. Rev. Fluid Mech.* 43 (2011) 59–78.