

# Heat Energy, Entropy and The Second Law: A New Perspective in Thermodynamics

Tang Suye\*

(Dated: October 28, 2021)

We present a new state function to define the energy of thermal motion, it is the last one of the undefined forms of the internal energy, combine with the mechanical free energy and Gibbs free energy, we rewrite the equations of the first law, this implies a new advance: we can describe the state changes of the internal energy directly. In a new theoretical structure, we define the concept of the heat partial pressure, rewrite the definition of the entropy, we discuss in detail on the three sources of irreversibility, the compensation mechanism of the heat conversion, and the internal causal principle of the second law, an interesting conclusion shows that the second law itself has already contained the mechanism of evolution.

## INTRODUCTION

Classical thermodynamics is a phenomenological theory, all we can do is to observe the observable behavior from the outside of a given system, and deduce its internal causal mechanism, this mindset start from the first law, and affect profoundly to the theoretical construct of thermodynamics. Until now, there still are some deep problems that cannot be addressed, it shows us that the traditional way of thinking seems to encounter its theoretical limits.

One of the famous problems is the entropy, in the present thermodynamics, "Now entropy is a very strange concept, and without hoping to achieve a complete description." [1] A confusing topic, what is entropy? has been discussed for countless times, no solutions to the problems. The contradiction is,  $\delta Q/T$  is a description observed from the outside of a given system, however, the heat transfer  $\delta Q$  cannot be used to describe the changes of the system state.

It is not an isolated problem, which involves the whole theoretical structure of thermodynamics, and needs for a profound change in the way of thinking: to change our perspective from the outside of a system to the inside of the system, this change depends on the complete classification of the internal energy, and the rediscovery of the first law. In classical theory, the classification for the internal energy was never made clear, and the heat energy is the last one of the undefined forms of the internal energy.

In this research, we present a new state function, the heat energy, the physical meaning of which denotes the energy of thermal motion, thus, the last one of the undefined forms of the internal energy will be made clear, combine with the mechanical free energy and Gibbs free energy, the internal energy can be classified into the three independent forms, this progress will imply a new theoretical structure.

In a new theoretical structure, we rewrite the equations of the first law, the distribution of the heat energy is defined to be the state of thermodynamic system, and the changes of the distribution of the heat energy can be described by an explicit equation combined with the heat transfer and the heat production, it implies an important advance: the energy transfer and the energy conversion can easily be distinguished by different explicit functions, we can describe the state changes of the internal energy directly.

In a new theoretical structure, we have ascertained the state changes when the flux  $\delta Q/T$  to be transferred into a given system, and redefine the concept of the entropy according to these state changes but not by the process, which is the complete description of the physical contents of the entropy. The new definition of the entropy can easily be proven to be a state function in maths, thus, it confirms that the imaginary cycle of heat engine [2] is not an essential requirement for the definition of the physical concept.

In a new theoretical structure, we have ascertained the three sources of irreversibility, the spontaneous processes are driven by different driving forces, the second law of thermodynamics is derived from a fundamental principle: All of the gradients of the thermodynamic forces spontaneously tend to zero, and in an irreversible process, the dissipation of the gradients of the thermodynamic forces cannot be spontaneously recovered. Irreversibility comes from the dissipation of thermodynamic driving forces, and the inequality  $d_i S \geq 0$  is the direction of irreversibility.

In a new theoretical structure, we discussed the relations between dissipation and evolution, we find that the second law itself has already contained the internal mechanism of evolution, this point of view comes from such a fact, in an arbitrary spontaneous process or non-spontaneous process, we always have the total entropy production  $d_i S \geq 0$ , but the partial differentials of  $d_i S$  can be less than zero, it implies that order and evolution can arise spontaneously, therefore, the second law is no longer thought to be only a dissipative law, it is also a law of evolution.

In a new theoretical structure, we establish a new foundation for discussing the relations between thermodynamics and dynamics, in the previous work, we have not found a similar theoretical structure of the second law that can be applied to dynamics. It seems, the master equations of dynamics only involve the conservation of energy, which correspond only to the first law of thermodynamics, it is destined to be a very difficult work in the previous time in that the theoretical structures of the two are different.

We believe, thermodynamics is a self-consistent theory, in a new theoretical structure, we will witness a profound change that will be brought into thermodynamics, in which, all of the concepts and principles are comprehensible.

"The fact that it is comprehensible is a miracle." [3]

## HEAT ENERGY: A NEW STATE FUNCTION

In physics, thermal motion is a state property, and the energy of thermal motion is thought to be a form of energy, J.C.Maxwell[4], R.Feynman[5] explicitly indicated that is the heat energy, and an earlier idea can also be found from D.Bernoulli book[6]: “what we experience as heat is simply the kinetic energy of the great numbers of the particles motion”, as we know, this great numbers of the particles motion now is called as thermal motion.

Now, we consider how to define the energy of thermal motion in thermodynamics. For a given system, using the symbol  $q$  denotes the energy of thermal motion, named as the heat energy, which is the last one of the undefined forms of the internal energy, this seems a bit strange.

By our experience, the heat energy is a non-conserved quantity, so it has the two sources, the energy transfer and the energy conversion, as we have already known,  $\delta Q$  is the heat transfer, and the another source of the energy of thermal motion, named as the heat production, comes from the heat conversion, such as a mechanocaloric effect, the heat effects of chemical reaction or phase transition.

In the present thermodynamics, the heat production is ill defined, and there has been a very strange point of view: the heat is only a form of energy in transfer, it is a common misconception and has been spreading for a long time.

Using  $d_e q$  denotes heat transfer, where the subscript  $e$  denotes the energy transfer, we have  $d_e q = \delta Q$ , using  $d_i q$  denotes the heat production, we obtain

$$dq = d_e q + d_i q. \quad (1)$$

Eq.(1) express the changes in the heat energy for a given system: the total change of the heat energy is equal to the sum of the heat transfer and the heat production.

Now we consider an example, for an ideal monatomic gas, all of the internal energy are the energy of thermal motion, therefore, by the first law, we have

$$dU = dq = \delta Q - pdV, \quad (2)$$

where  $U$  is the internal energy,  $Q$  is the heat transfer,  $p$  is the gas pressure,  $V$  is the gas volume, and  $-pdV = \delta W$  is the reversible volume work.

For an ideal gas, the pressure  $p$  comes from thermal motion, so  $pdV$  denotes work done by the heat energy  $q$ , it signifies the heat energy of the gas to be converted into work, and then exchanged with its surroundings.

$$d_e q = \delta Q, \quad (3)$$

$$d_i q = -pdV. \quad (4)$$

For  $pV$  work of an ideal gas, the work  $-\delta W = pdV$  is the change in the surroundings, and  $d_i q$  is the change in the system, thus, the work done  $-pdV$  denotes not only the energy exchange but also the energy conversion between the system and its surroundings.

In general, the pressure denotes a force acting per unit area, which involves different sources, including thermal motion and the interactions between the particles. Using  $p$  represents the heat partial pressure which comes only from thermal motion, using  $p'$  represents the partial pressure that being the other generalized force  $Y$  is expressed as a force acting per unit area, the different sources of the total pressure therefore can be distinguished by

$$P = p + p', \quad (5)$$

where  $P$  is the total pressure.

The first law thus becomes

$$\begin{aligned} dU &= \delta Q - pdV + Y dx + \sum_j \mu'_j dN_j \\ &= dq + Y dx + \sum_j \mu'_j dN_j, \end{aligned} \quad (6)$$

where  $Y$  is the generalized force,  $dx$  is the generalized displacement,  $Y dx$  includes  $-p' dV$ ,  $\mu'_j$  is the chemical potential of the type  $-j$  particles, and  $N_j$  is the number of such particles. The internal energy  $U$  is the sum of the three independent forms of the energies.

An interesting difference shows that “the pressure  $p$  has different sign from other generalized force, if we increase the pressure, the volume increases, whereas if we increase the force,  $Y$ , for all other cases, the extensive variable,  $x$ , decreases”[7]. This is an interesting criterion that can be used to distinguish the heat partial pressure  $p$  from other generalized force  $Y$ , by this criterion, the heat partial pressure  $p$  will be different from other generalized force.

We find that  $dq$  can be proven to be an exact differential.

Consider  $U=U(T, V, x, N_j)$ , where  $T, V, x$  and  $N_j$  are the four independent state variables. Then we obtain[7]

$$\begin{aligned} dU &= \left( \frac{\partial U}{\partial T} \right)_{V,x,N_j} dT + \left( \frac{\partial U}{\partial V} \right)_{T,x,N_j} dV \\ &+ \left( \frac{\partial U}{\partial x} \right)_{T,V,N_j} dx + \sum_j \left( \frac{\partial U}{\partial N_j} \right)_{T,V,x,N_{i \neq j}} dN_j. \end{aligned} \quad (7)$$

By definition and compare with Eq.(6),  $q$  is the function of the two independent variables  $T$  and  $V$

$$dq = \left( \frac{\partial q}{\partial T} \right)_V dT + \left( \frac{\partial q}{\partial V} \right)_T dV. \quad (8)$$

Compare Eq.(7) with Eq.(8), we get

$$\begin{aligned} dq &= \left( \frac{\partial q}{\partial T} \right)_V dT + \left( \frac{\partial q}{\partial V} \right)_T dV \\ &= \left( \frac{\partial U}{\partial T} \right)_{V,x,N_j} dT + \left( \frac{\partial U}{\partial V} \right)_{T,x,N_j} dV. \end{aligned} \quad (9)$$

Owing to the fact that  $dU$  is an exact differential, by Eq.(7) and Eq.(9), the two partial derivatives of the internal energy satisfy the relation that

$$\left[ \frac{\partial}{\partial V} \left( \frac{\partial U}{\partial T} \right)_{V,x,N_j} \right]_{T,x,N_j} = \left[ \frac{\partial}{\partial T} \left( \frac{\partial U}{\partial V} \right)_{T,x,N_j} \right]_{V,x,N_j} \quad (10)$$

Such that

$$\left[ \frac{\partial}{\partial V} \left( \frac{\partial q}{\partial T} \right) \right]_V \Big|_T = \left[ \frac{\partial}{\partial T} \left( \frac{\partial q}{\partial V} \right) \right]_T \Big|_V. \quad (11)$$

Thus we have proven that  $dq$  is an exact differential.

Using the function  $A$  denotes the mechanical free energy within a given system that

$$A = Yx. \quad (12)$$

The equation of the first law becomes

$$\begin{aligned} dU &= dq + dA + dG \\ &= dq + d(Yx) + \sum_j d(\mu'_j N_j). \end{aligned} \quad (13)$$

The internal energy  $U$  is equal to the sum of the internal heat energy  $q$ , the internal mechanical free energy  $A$ , and the internal Gibbs free energy  $G$ .

By definitions,  $A$  and  $G$  both are the non-conserved quantities, we have the two equations similar to Eq.(1)

$$dA = d_e A + d_i A, \quad (14)$$

where  $d_e A$  denotes the mechanical free energy flux, and  $d_i A$  denotes the mechanical free energy production.

For Gibbs free energy we have

$$dG = d_e G + d_i G, \quad (15)$$

where  $d_e G$  denotes Gibbs free energy flux,  $d_i G$  denotes Gibbs free energy production, similar to Eq.(14).

Thus the concepts “flux” and “production” can be used as a general method to distinguish the two processes: the energy transfer and the energy conversion.

The equation of the first law can be extended that

$$dU = d_e q + d_e A + d_e G, \quad (16)$$

$$d_i q + d_i A + d_i G = 0. \quad (17)$$

Eq.(17) gives the relation

$$d_i q = -(d_i A + d_i G). \quad (18)$$

Eq.(18) denotes the relation of the energy conversions between the different energy forms, the heat production is equal to the negative free energy productions in that energy is a conserved quantity.

Now, we consider some applications.

The first example we consider an equation combined with heat diffusion and heat conversion, which describes the changes of “the distribution of the heat”[8].

$$\frac{\partial q_\rho}{\partial t} = -\nabla \cdot \mathbf{J}_q + \frac{\partial_i q_\rho}{\partial t}, \quad (19)$$

$$\mathbf{J}_q = -k \nabla T, \quad (20)$$

where  $q_\rho$  is  $q$  per unit volume,  $\mathbf{J}_q$  denotes the heat flux density,  $t$  is time, and  $k$  is the thermal conductivity.

The second example is the heat effects of chemical reaction or phase transition, the change in the heat production

$$d_i q = -d_i G, \quad (21)$$

where  $d_i q \geq 0$  or  $d_i G \leq 0$  can be considered as the criterion for a spontaneous reaction or phase transition.

Eq.(1) and Eq.(6) both are the equations of conservation of energy, the two both cannot describe the state changes of a given system completely. We noticed, the state changes of the heat transfer and  $p\bar{V}$  work done are different.

The driving force of heat into work comes from the heat partial pressure, which will be dissipated along with the progress of the conversion, indicates the dissipation of a conversion potential. For a reversible conversion, we have

$$d_i q = -pdV. \quad (22)$$

$-pdV$  has the two meanings: (a) the volume work in a reversible process, (b) the change of a conversion potential. The conversion of heat into work / free energy needs a compensation, which comes from the dissipation of the conversion potential, this is the internal causal mechanism of the heat conversion, and also the compensation mechanism of the conversion of heat into work / free energy.

$$d_i q + pdV = 0. \quad (23)$$

Eq.(23) describes the complete state changes in the heat conversion, the conversion of heat into work / free energy will consume an equal amount of the conversion potential.

In an irreversible conversion process, we have

$$d_i q + pdV \geq 0. \quad (24)$$

The free expansion of a gas is a simple example.

## ENTROPY: EXACT DIFFERENTIAL

In classical thermodynamics, the concept of the entropy was defined by the aid of Carnot cycle that

$$\oint_{rev} \frac{\delta Q}{T} = 0, \quad (25)$$

where  $rev$  denotes the reversible processes. It given a conditional definition with an additional restriction that

$$dS = \left( \frac{\delta Q}{T} \right)_{rev}. \quad (26)$$

In Eq.(26),  $(\delta Q/T)_{rev}$  is an inexact differential in that it is reversible path dependent. For an irreversible path, the definition must combine with the inequality  $dS \geq \delta Q/T$ .

We now consider a new approach.

Consider the heat transfer  $\delta Q$ , it is the increment of an extensive quantity, and can be transferred only from higher temperature to lower temperature, this fact shows a very important property that the heat transfer  $\delta Q$  relates to the intensity distribution of the temperature  $T$ .

For the same  $\delta Q$ , if its temperature distribution is different, the intensive property of  $\delta Q$  will be also different, thus, we can use the function  $\delta Q/T$  to measure  $\delta Q$  and its intensive property of the temperature distribution, and then  $\delta Q/T$  may be named as the entropy of  $\delta Q$ .

$$d_e S = \frac{\delta Q}{T}. \quad (27)$$

In Eq.(27),  $\delta Q/T$  is only a process variable. For a given system, we need to consider when  $\delta Q$  to be transformed into the internal energy of the system, or the internal energy of the system to be transformed into the heat flux  $\delta Q$ , how do we describe the state changes of the system.

Consider a reversible process, combined with Eq.(6) and Eq.(23), it follows that

$$\frac{\delta Q}{T} \Rightarrow \frac{dU}{T} - \frac{Y dx}{T} - \sum_j \frac{\mu'_j dN_j}{T} + \frac{pdV}{T} \Rightarrow \frac{dq}{T} + \frac{pdV}{T}.$$

When  $\delta Q/T$  to be transported into a given system, the right hand side of the equation is the state changes.

Using the following function denotes the state changes

$$dS = \frac{dq}{T} + \frac{pdV}{T}. \quad (28)$$

$S$  is named as the entropy of the given system.

Eq.(28) is a binomial expression, the integral of  $dq/T$  we named as the calorimetric entropy  $S_q$ ,

$$dS_q = \frac{dq}{T}. \quad (29)$$

The integral of  $pdV/T$  we named as the configurational entropy  $S_v$ .

$$dS_v = \frac{pdV}{T}. \quad (30)$$

By Eq.(8) and Eq.(28),  $S$  is the function of the two independent variables  $T$  and  $V$ . According to the mathematical theory of exact differential, if the two partial derivatives of the function  $S$  satisfy the equation

$$\left[ \frac{\partial}{\partial V} \left( \frac{\partial S}{\partial T} \right) \right]_T = \left[ \frac{\partial}{\partial T} \left( \frac{\partial S}{\partial V} \right) \right]_V. \quad (31)$$

$dS$  will be an exact differential.

We now prove this conclusion.

Using Eq.(8) in Eq.(28), it follows that

$$\begin{aligned} dS &= \frac{dq}{T} + \frac{p}{T} dV \\ &= \frac{1}{T} \left( \frac{\partial q}{\partial T} \right)_V dT + \frac{1}{T} \left( \frac{\partial q}{\partial V} \right)_T dV + \frac{p}{T} dV. \end{aligned} \quad (32)$$

By Eq.(32) we have

$$\begin{aligned} \left[ \frac{\partial}{\partial V} \left( \frac{\partial S}{\partial T} \right) \right]_T &= \left\{ \frac{\partial}{\partial V} \left[ \frac{1}{T} \left( \frac{\partial q}{\partial T} \right)_V \right] \right\}_T \\ &= \frac{1}{T} \left[ \frac{\partial}{\partial V} \left( \frac{\partial q}{\partial T} \right)_V \right]_T, \end{aligned} \quad (33)$$

and

$$\begin{aligned} \left[ \frac{\partial}{\partial T} \left( \frac{\partial S}{\partial V} \right) \right]_V &= \left\{ \frac{\partial}{\partial T} \left[ \frac{1}{T} \left( \frac{\partial q}{\partial V} \right)_T + \left( \frac{p}{T} \right) \right] \right\}_V \\ &= \frac{1}{T} \left[ \frac{\partial}{\partial T} \left( \frac{\partial q}{\partial V} \right)_T \right]_V - \frac{1}{T^2} \left( \frac{\partial q}{\partial V} \right)_T \\ &\quad + \frac{1}{T^2} \left[ T \left( \frac{\partial p}{\partial T} \right)_V - p \right]. \end{aligned} \quad (34)$$

Before we compare Eq.(33) with Eq.(34), we need to prove the following relation first

$$\left( \frac{\partial q}{\partial V} \right)_T = T \left( \frac{\partial p}{\partial T} \right)_V - p. \quad (35)$$

From Eq.(32) we have

$$dS = \frac{1}{T} \left( \frac{\partial q}{\partial T} \right)_V dT + \frac{1}{T} \left[ \left( \frac{\partial q}{\partial V} \right)_T + p \right] dV. \quad (36)$$

It follows that

$$dS = \left( \frac{\partial S}{\partial T} \right)_V dT + \left( \frac{\partial S}{\partial V} \right)_T dV. \quad (37)$$

$dq$  can be expressed as the follows

$$dq = T dS - pdV. \quad (38)$$

The pressure  $p$  and the volume  $V$  both are the state variables, so  $d(pV)$  is an exact differential, we consider

$$pV = TS - q. \quad (39)$$

$$d(pV) = d(TS) - dq = T dS + S dT - dq. \quad (40)$$

Using Eq.(38) in Eq.(40), we get

$$d(pV) = d(TS) - dq = S dT + pdV. \quad (41)$$

Where  $d(pV)=d(pV)(T, V)$ , we have from Eq.(41)

$$\left[ \frac{\partial}{\partial V} \left( \frac{\partial(pV)}{\partial T} \right) \right]_T = \left( \frac{\partial S}{\partial V} \right)_T, \quad (42)$$

and

$$\left[ \frac{\partial}{\partial T} \left( \frac{\partial(pV)}{\partial V} \right) \right]_V = \left( \frac{\partial p}{\partial T} \right)_V. \quad (43)$$

Since  $d(pV)$  is an exact differential, so that

$$\left( \frac{\partial S}{\partial V} \right)_T = \left( \frac{\partial p}{\partial T} \right)_V. \quad (44)$$

Using Eq.(44) in Eq.(37), we get

$$dS = \left( \frac{\partial S}{\partial T} \right)_V dT + \left( \frac{\partial p}{\partial T} \right)_V dV. \quad (45)$$

Compare Eq.(45) with Eq.(36), the two partial derivatives satisfy the equation that

$$\left( \frac{\partial p}{\partial T} \right)_V = \frac{1}{T} \left[ \left( \frac{\partial q}{\partial V} \right)_T + p \right]. \quad (46)$$

It follows that

$$\left(\frac{\partial q}{\partial V}\right)_T = T \left(\frac{\partial p}{\partial T}\right)_V - p. \quad (47)$$

Thus we have proven the relation Eq.(35).

Using Eq.(47) in Eq.(34), we get

$$\left[\frac{\partial}{\partial T} \left(\frac{\partial S}{\partial V}\right)\right]_{T,V} = \frac{1}{T} \left[\frac{\partial}{\partial T} \left(\frac{\partial q}{\partial V}\right)\right]_{T,V}. \quad (48)$$

Compare Eq.(33) and Eq.(48) with Eq.(11), we obtain the following conclusion

$$\left[\frac{\partial}{\partial V} \left(\frac{\partial S}{\partial T}\right)\right]_{T,V} = \left[\frac{\partial}{\partial T} \left(\frac{\partial S}{\partial V}\right)\right]_{T,V}. \quad (49)$$

Eq.(49) indicates that  $dS$  is an exact differential,  $S$  is a state function. The conclusion perfectly obeys the general rules of exact differential, and does not need to depend on an imaginary reversible cycle, therefore, there is no exception to the principles of exact differential.

As the conclusion of Eq.(49), the function changes  $dS$  will be path-independent, the integral of  $dS$  around an arbitrary closed path will be always equal to zero

$$\oint dS \equiv 0. \quad (50)$$

In non-equilibrium thermodynamics, the changes in the entropy may be considered as the sum of the entropy flux  $d_e S$  and the entropy production  $d_i S$  that

$$dS = d_e S + d_i S. \quad (51)$$

In this opinion,  $\delta Q/T$  is really the entropy flux  $d_e S$ . The physical meaning of Clausius inequality is that

$$\oint d_e S = \oint \frac{\delta Q}{T} \leq 0, \quad (52)$$

and the second law inequality is actually that

$$\oint (d_e S + d_i S) \geq \oint d_e S. \quad (53)$$

The imaginary cycle of heat engine is only a special case, the definition  $dS = (\delta Q/T)_{rev}$  cannot confirm the physical connotation of the entropy completely.

We now consider the two examples. The first example is a photon gas. Using  $q = \alpha T^4 V$ ,  $p = \alpha T^4 / 3$  in Eq.(32), where  $\alpha$  is a constant, and then we get that

$$dq = d(\alpha T^4 V) = 4\alpha T^3 V dT + \alpha T^4 dV. \quad (54)$$

$$\begin{aligned} dS &= \frac{1}{T} dq + \frac{p}{T} dV \\ &= \frac{1}{T} 4\alpha T^3 V dT + \frac{1}{T} \alpha T^4 dV + \frac{1}{3T} \alpha T^4 dV \\ &= 4\alpha T^2 V dT + \frac{4}{3} \alpha T^3 dV. \end{aligned} \quad (55)$$

We have from Eq.(55)

$$\left[\frac{\partial}{\partial V} \left(\frac{\partial S}{\partial T}\right)\right]_{T,V} = \left[\frac{\partial}{\partial T} \left(\frac{\partial S}{\partial V}\right)\right]_{T,V} = 4\alpha T^2. \quad (56)$$

Thus the entropy of a photon gas is a state function.

The second example is an ideal gas. Using  $dq = C_v dT$  and  $pV = nRT$  in Eq.(32), where  $C_v$  is the heat capacity at constant volume,  $n$  is the number of moles, and  $R$  is the gas constant. We have

$$\begin{aligned} dS &= \frac{C_v}{T} dT + \frac{p}{T} dV \\ &= \frac{C_v}{T} dT + \frac{nR}{V} dV, \end{aligned} \quad (57)$$

and

$$\left[\frac{\partial}{\partial V} \left(\frac{C_v}{T}\right)\right]_{T,V} = \left[\frac{\partial}{\partial T} \left(\frac{nR}{V}\right)\right]_{T,V} = 0. \quad (58)$$

Thus the entropy of an ideal gas is a state function.

For an ideal gas,  $T = 2q / iNk$ ,  $pV = NkT$ , we get

$$dS = \frac{iNk}{2} \frac{dq}{q} + \frac{Nk}{V} dV. \quad (59)$$

where  $i$  is the number of the degrees of freedom of the gas molecules,  $N$  is the number of the gas molecules, and  $k$  is Boltzmann constant.

For a photon gas, we have the photon gas equation

$$pV = \frac{\zeta(4)}{\zeta(3)} NkT = NR_{\{p\}} T, \quad (60)$$

where  $N$  is the number of the photons,  $\zeta(3)$  and  $\zeta(4)$  both are the Riemann zeta functions,  $R_{\{p\}} = [\zeta(4)/\zeta(3)]k$  is the photon gas constant.

Consider  $q = 3pV$  and  $T = q / 3NR_{\{p\}}$ , combine Eq.(60) with Eq.(55), we get

$$dS = \frac{3NR_{\{p\}}}{q} dq + \frac{NR_{\{p\}}}{V} dV. \quad (61)$$

All of the variables in Eq.(59) and Eq.(61) are extensive variables, where  $iN/q$  refers to the average energy levels per degree of freedom of the gas particles, and  $N/V$  is the density function of the gas particles. In an earlier work[9], we presented a mathematical model to express the average degrees for the distributions of the extensive variables  $q$  and  $N$ , for an ideal gas or a photon gas, the Nature of the Entropy is a binomial distribution function.

In Eq.(59) and Eq.(61),  $dS$  has the same function form, it may be the primary equation of the entropy that

$$dS = \frac{\hat{\alpha}NR_{\{k\}}}{q} dq + \frac{NR_{\{k\}}}{V} dV. \quad (62)$$

where  $\hat{\alpha}$  denotes a factor, for an ideal gas  $\hat{\alpha} = i/2$ , for a photon gas  $\hat{\alpha} = i$ , and  $R_{\{k\}}$  denotes the system constant, for an ideal gas  $R_{\{k\}} = k$ , for a photon gas  $R_{\{k\}} = R_{\{p\}}$ .

According to Eq.(49), Eq.(28) is a perfect definition of the state function in the usual sense, and the fundamental equation of thermodynamics is the equivalent equation of the definition of the entropy, so Eq.(28) has the same mathematical results as that can be derived by the fundamental equation, such that, the new approach does not change the mathematical results of classical thermodynamics.

### THE SECOND LAW: THE TOTAL DIFFERENTIAL

The classical inequality of the second law is derived by the aid of heat engine cycle, it is expressed as the increment in the entropy of an isolated system  $dS \geq 0$ , and in non-equilibrium thermodynamics, the increment in the entropy of an isolated system has been reformulated by the entropy production  $d_i S$  and the inequality  $d_i S \geq 0$ [10].

We now discuss the entropy production and the second law, the two both involve the different sources, and need to be carefully distinguished in a depth analysis.

By Eq.(13), for a given system, the internal energy

$$dU = dq + dA + dG, \quad (63)$$

where  $U$  is the internal energy,  $q$  is the heat energy,  $A$  is the mechanical free energy, and  $G$  is Gibbs free energy.

By Eq.(28),  $dq/T$  is the entropy of  $dq$ , but  $dA/T$  and  $dG/T$  make no sense. The mechanical free energy  $A$  and Gibbs free energy  $G$  can be called by a general designation: the free energy. The difference between the two is that the heat energy  $q$  has some quality losses, these quality losses can be measured by the entropy, the free energy has no quality loss, the entropy of which is equal to zero, only the heat energy makes contribution to the entropy.

That is why the entropy can hardly continue to be discussed in the traditional theoretical structure, it is readily apparent that a perfect definition must depend on such a fact: the heat energy has already been well defined.

Assume a given system in locals equilibrium, and some irreversible processes take place between the two locals A and B, the driving forces of the processes are the gradients of thermodynamic forces between the two locals. Using the subscript  $A$  and  $B$  denote the locals A and B, the subscript  $AB$  denote the interactions between the locals A and B, the changes in the internal energy of the processes can be expressed respectively as follows.

The heat flux driven by the gradient in temperature  $\Delta T$  between the two locals A and B satisfies the relation

$$d_e q = d_e q_B = -d_e q_A. \quad (64)$$

The changes in the mechanical free energy

$$d_i A_{AB} = -(Y_B - Y_A)dx = -\Delta Y dx, \quad (65)$$

where the driving force  $\Delta Y \geq 0$  denotes a spontaneous process, and  $d_i A_{AB} \leq 0$  denotes the free energy decreasing.

The changes in Gibbs free energy

$$d_i G_{AB} = -\sum_j (\mu'_{jB} - \mu'_{jA}) dN_j = -\sum_j \Delta \mu'_j dN_j, \quad (66)$$

where the driving forces  $\Delta \mu'_j \geq 0$  denotes a spontaneous process,  $d_i G_{AB} \leq 0$  denotes the free energy decreasing.

If an independent process takes place only in the locals A or B, using the subscripts  $a$  and  $b$  denote the independent changes of the locals A and B, respectively, we have

$$d_i A_A = -Y_A dx_a, \quad (67)$$

$$d_i A_B = -Y_B dx_b, \quad (68)$$

and

$$d_i G_A = -\sum_j \mu'_{jA} dN_{ja}, \quad (69)$$

$$d_i G_B = -\sum_j \mu'_{jB} dN_{jb}. \quad (70)$$

Combine Eqs.(65)–(70), and sum the changes of the two locals A and B, it follows that

$$\begin{aligned} \sum_k d_i A_k &= d_i A_{AB} + d_i A_A + d_i A_B \\ &= -(Y_B - Y_A)dx - (Y_A - 0)dx_a - (Y_B - 0)dx_b \\ &= -\sum_k \Delta Y dx, \end{aligned} \quad (71)$$

$$\begin{aligned} \sum_k d_i G_k &= d_i G_{AB} + d_i G_A + d_i G_B \\ &= -\sum_j (\mu'_{jB} - \mu'_{jA}) dN_j \\ &\quad -\sum_j (\mu'_{jA} - 0) dN_{ja} - \sum_j (\mu'_{jB} - 0) dN_{jb} \\ &= -\sum_k \sum_j \Delta \mu'_j dN_j, \end{aligned} \quad (72)$$

where, the subscript  $k$  denotes the process  $k$ .

Such that, we have

$$d_i A = \sum_k d_i A_k = -\sum_k \Delta Y dx, \quad (73)$$

$$d_i G = \sum_k d_i G_k = -\sum_k \sum_j \Delta \mu'_j dN_j. \quad (74)$$

By Eq.(18), we have the heat production  $d_i q$  that

$$\begin{aligned} d_i q &= -(d_i A + d_i G) \\ &= \sum_k \Delta Y dx + \sum_k \sum_j \Delta \mu'_j dN_j. \end{aligned} \quad (75)$$

Now we consider the total entropy production, which involves the different sources of irreversibility.

Using Eq.(1) in Eq.(28), we get

$$\begin{aligned} dS &= d_e S + d_i S \\ &= \frac{d_e q}{T} + \frac{d_i q}{T} + \frac{p dV}{T}, \end{aligned} \quad (76)$$

where  $d_e S$  represents the entropy flux, and  $d_i S$  represents the internal entropy production.

The sum of the increments in the entropies of the two locals A and B is equal to

$$dS = d_e S + d_i S = dS_A + dS_B. \quad (77)$$

The increments in the entropy of the two locals that

$$dS_A = \frac{d_e q_A}{T_A} + \frac{d_i q_A}{T_A} + \frac{p_A dV_A}{T_A}. \quad (78)$$

$$dS_B = \frac{d_e q_B}{T_B} + \frac{d_i q_B}{T_B} + \frac{p_B dV_B}{T_B}. \quad (79)$$

Assume the external entropy flux  $d_e S = 0$ , and then, the sum of the entropy productions of the two locals will be equal to that

$$\begin{aligned} d_i S &= dS_A + dS_B \\ &= \left( \frac{d_e q_A}{T_A} + \frac{d_e q_B}{T_B} \right) + \left( \frac{d_i q_A}{T_A} + \frac{d_i q_B}{T_B} \right) \\ &\quad + \left( \frac{p_A dV_A}{T_A} + \frac{p_B dV_B}{T_B} \right). \end{aligned} \quad (80)$$

In Eq.(80), the heat fluxes satisfy the relation

$$d_e q_B = -d_e q_A = d_e q. \quad (81)$$

It follows that

$$\left( \frac{d_e q_A}{T_A} + \frac{d_e q_B}{T_B} \right) = \left( \frac{1}{T_B} - \frac{1}{T_A} \right) d_e q = \Delta \left( \frac{1}{T} \right) d_e q. \quad (82)$$

By Eq.(75), we have

$$\left( \frac{d_i q_A}{T_A} + \frac{d_i q_B}{T_B} \right) = \sum_k \frac{1}{T} \Delta Y dx + \sum_k \sum_j \frac{1}{T} \Delta \mu'_j dN_j, \quad (83)$$

where,  $T$  is the temperature of the heat conversion.

Using  $dV_r$  denotes the volume change of the reversible volume work between the two locals A and B, using  $dV_a$  and  $dV_b$  represent the two independent volume changes caused by the pressures  $p_A$  and  $p_B$ , respectively that

$$dV_A = -dV_r + dV_a, \quad (84)$$

$$dV_B = dV_r + dV_b. \quad (85)$$

Using Eq.(84) and Eq.(85) in Eq.(80), we get that

$$\begin{aligned} \left( \frac{p_A dV_A}{T_A} + \frac{p_B dV_B}{T_B} \right) &= \left( \frac{p_B}{T_B} - \frac{p_A}{T_A} \right) dV_r \\ &\quad + \left( \frac{p_A}{T_A} - 0 \right) dV_a + \left( \frac{p_B}{T_B} - 0 \right) dV_b \\ &= \sum_k \Delta \left( \frac{p}{T} \right) dV. \end{aligned} \quad (86)$$

The subscript  $k$  denotes the process  $k$ .

Using Eqs.(82), (83) and (86) in Eq(80), we obtain

$$\begin{aligned} d_i S &= dS_A + dS_B \\ &= \Delta \left( \frac{1}{T} \right) d_e q + \sum_k \frac{1}{T} \Delta Y dx \\ &\quad + \sum_k \sum_j \frac{1}{T} \Delta \mu'_j dN_j + \sum_k \Delta \left( \frac{p}{T} \right) dV. \end{aligned} \quad (87)$$

The heat flux  $d_e q$  can be considered as a micro-quantity  $dq$ , and the sum “ $k$ ” can be simplified, thus we have

$$\begin{aligned} d_i S &= \Delta \left( \frac{1}{T} \right) dq + \frac{1}{T} \Delta Y dx \\ &\quad + \sum_j \frac{1}{T} \Delta \mu'_j dN_j + \Delta \left( \frac{p}{T} \right) dV. \end{aligned} \quad (88)$$

For a non-equilibrium state, consider the spatial distribution of the gradients, it follows that

$$\begin{aligned} d_i S &= \nabla \left( \frac{1}{T} \right) d\mathbf{q} + \frac{1}{T} \nabla Y dx \\ &\quad + \sum_j \frac{1}{T} \nabla \mu'_j dN_j + \nabla \left( \frac{p}{T} \right) dV. \end{aligned} \quad (89)$$

In Eq.(89),  $\mathbf{q}$  is the heat flux,  $dV = dV_i \mathbf{e}_i + dV_j \mathbf{e}_j + dV_k \mathbf{e}_k$ , where  $\mathbf{e}_i$ ,  $\mathbf{e}_j$  and  $\mathbf{e}_k$  are the direction vectors.

Eq.(88) and Eq.(89) are the equations of the total entropy productions for a given system.

Below we discuss the second law.

We consider the independent processes first, if there is only one gradient of thermodynamic force driving the process, the process is called as an independent process.

(1). The first type of the independent process is the heat transfer, a micro-quantity of the heat energy  $dq$  was transferred between different temperatures, the driving force of the heat transfer is the gradient of the temperature, and the direction of the heat flux is determined by the gradient in the temperature  $\Delta T$ , so, the heat energy can only be transferred from higher temperature  $T_h$  to lower temperature  $T_c$ .

$$dq(T_h) \rightarrow dq(T_c).$$

We have the entropy production

$$d_i S = \left( \frac{1}{T_c} - \frac{1}{T_h} \right) dq = \Delta \left( \frac{1}{T} \right) dq \geq 0. \quad (90)$$

The heat transfer obeys a simple principle: the gradient  $\Delta T$ , or  $\Delta(1/T)$  spontaneously tend to zero, the entropy production describes the dissipation of the the driving force  $\Delta(1/T)$  or  $\Delta T$ . In an independent process, the dissipation of the thermodynamic driving force  $\Delta(1/T)$  or  $\Delta T$  cannot be spontaneously recovered.

(2). The second type is the independent process driven by the heat partial pressure  $p$ . Consider a given system, the heat partial pressure  $p$  always points to the normal direction

of the system's surface, thus the volume change of the system always be positive definite in a spontaneous diffusion process driven by the heat partial pressure, so we have

$$d_i S = \left( \frac{p}{T} - 0 \right) dV = \Delta \left( \frac{p}{T} \right) dV \geq 0. \quad (91)$$

The spontaneous diffusion process driven by the heat partial pressure  $p$  obeys such a principle: the gradient  $\Delta p$ , or  $\Delta(p/T)$  spontaneously tends to zero. In an independent process, the dissipation of the driving force  $\Delta(p/T)$  or  $\Delta p$  cannot be spontaneously recovered.

(3). The third type of the independent process is the free energy transport, in such a process, the free energy partly be converted into the heat energy.

By Eq.(75), we have the heat production  $d_i q$  that

$$d_i q = \Delta Y dx + \sum_j \Delta \mu'_j dN_j. \quad (92)$$

The transport process obeys the law of the conservation of energy, so  $d_i q$  cannot be less than zero, it follows that the transport efficiency of the free energy transport process cannot be greater than 100%, thus we get

$$d_i q = \Delta Y dx + \sum_j \Delta \mu'_j dN_j \geq 0, \quad (93)$$

Since  $d_i q \geq 0$  increases the heat partial pressure  $p$ , so meanwhile, we have  $pdV \geq 0$ , the entropy production

$$d_i S = \frac{1}{T} \Delta Y dx + \sum_j \frac{1}{T} \Delta \mu'_j dN_j + \Delta \left( \frac{p}{T} \right) dV \geq 0. \quad (94)$$

where  $T$  is the temperature of the heat conversion.

Eq.(94) represents the free energy dissipations in the free energy transport processes. The free energy dissipations are derived from such a principle: the gradients  $\Delta Y$  and  $\Delta \mu'_j$  spontaneously tend to zero.

The three independent irreversible processes we have discussed above are the three sources of irreversibility, the driving forces of the processes come from the gradients in the temperature  $\Delta(1/T)$ , in the generalized force  $\Delta Y$ , in the chemical potential of the type  $j$  particles  $\Delta \mu'_j$ , and in  $\Delta(p/T)$ , the dissipations of the gradients correspond to  $d\Delta(1/T) \leq 0$ ,  $d\Delta Y \leq 0$ ,  $d\Delta \mu'_j \leq 0$ , and  $d\Delta(p/T) \leq 0$ , the reverse processes cannot progress spontaneously, in that the directions of the processes are determined only by the driving forces, a positive gradient production can only be driven by another gradient and in a coupling process.

In general, a thermodynamic coupling process involves two or more gradients of thermodynamic forces, involves the two basic types: (1) the heat energy is converted into the free energy with some compensations; (2) the free energy is converted into the heat energy via the path of the volume work. The two coupling processes will be discussed below, respectively, the two both involve the energy conversions.

(1). The first type of the coupling process is the conversion of heat into work, the heat energy is converted into the free energy via the path of the volume work

$$dq \rightarrow \delta W_v(-pdV_r) \rightarrow Y dx + \sum_j \mu'_j dN_j.$$

where  $-pdV_r$  denotes the reversible volume work.

Consider an arbitrary coupling process, we have

$$d_i q + pdV_r = \Delta Y dx + \sum_j \Delta \mu'_j dN_j + pdV_r, \quad (95)$$

where we only consider a reversible process.

The conversion process obeys the law of the conservation of energy, so we have

$$d_i q = \Delta Y dx + \sum_j \Delta \mu'_j dN_j = -pdV_r. \quad (96)$$

Combine Eq.(95) and Eq.(96) with Eq.(88), we get

$$\begin{aligned} d_i S &= \frac{1}{T} \Delta Y dx + \sum_j \frac{1}{T} \Delta \mu'_j dN_j + \Delta \left( \frac{p}{T} \right) dV_r \\ &= \left( \frac{1}{T} d_i q + \frac{p}{T} dV_r \right) = 0. \end{aligned} \quad (97)$$

The driving forces of the process is derived from the gradient  $\Delta(p/T)$ , according to Eq.(29) and Eq.(30), the entropy is the sum of the calorimetric entropy  $S_q$  and the configurational entropy  $S_v$ , combine with Eq.(97) that

$$\begin{aligned} d_i S &= d_i S_q + d_i S_v \\ &= \frac{1}{T} d_i q + \frac{p}{T} dV_r = 0. \end{aligned} \quad (98)$$

Since  $d_i q \leq 0$ , so in Eq.(98), we have

$$d_i S_q = \frac{1}{T} d_i q \leq 0, \quad (99)$$

and

$$d_i S_v = \frac{p}{T} dV_r \geq 0. \quad (100)$$

When the heat energy is converted into the free energy, the original quality loss will be retained in another form

$$d_i S_q = \frac{1}{T} d_i q \rightarrow d_i S_v = \frac{p}{T} dV_r. \quad (101)$$

Eq.(101) establishes a compensation mechanism, when the heat energy is converted into the free energy via the path of the volume work done, the calorimetric entropy  $d_i q/T$  will be transformed into the configurational entropy  $pdV_r/T$ , and never disappear. It shows an interesting fact, that the partial differential of the total entropy production can be less than zero.

(2). The second type of the coupling process is the conversion of the free energy into the heat energy via the path



of the volume work, including the case that some part of the free energy is converted directly into the heat energy

$$Y dx + \sum_j \mu'_j dN_j \rightarrow \delta W_v(-pdV_r) \rightarrow dq.$$

The driving forces of the processes come from the gradients in  $\Delta Y$  and / or  $\Delta \mu'_j$ .

Since some part of the free energy be converted directly into the heat energy, so we have

$$d_i q = \Delta Y dx + \sum_j \Delta \mu'_j dN_j \geq -pdV_r. \quad (102)$$

where  $-pdV_r$  denotes the reversible volume work.

Using Eq.(102) in Eq.(88), it follows that

$$\begin{aligned} d_i S &= \frac{1}{T} \Delta Y dx + \sum_j \frac{1}{T} \Delta \mu'_j dN_j + \Delta \left( \frac{p}{T} \right) dV_r \\ &= \frac{1}{T} d_i q + \frac{p}{T} dV_r \geq 0. \end{aligned} \quad (103)$$

Similar to Eq.(98), we have

$$\begin{aligned} d_i S &= d_i S_q + d_i S_v \\ &= \frac{1}{T} d_i q + \frac{p}{T} dV_r \geq 0. \end{aligned} \quad (104)$$

In Eq.(104), we have

$$d_i S_q = \frac{1}{T} d_i q \geq 0, \quad (105)$$

and

$$d_i S_v = \frac{p}{T} dV_r \leq 0. \quad (106)$$

In such a process, the configurational entropy  $pdV_r/T$  is transformed into the calorimetric entropy  $d_i q/T$  that

$$d_i S_v = \frac{p}{T} dV_r \rightarrow d_i S_q = \frac{1}{T} d_i q. \quad (107)$$

In Eq.(98) and Eq.(104), the total entropy productions involve the two terms

$$d_i S = d_i S_q + d_i S_v \geq 0, \quad (108)$$

where  $d_i S_q$  denotes the calorimetric entropy production, and  $d_i S_v$  denotes the configurational or the density entropy production. In a coupling process, the total entropy production will always be positive definite,  $d_i S \geq 0$ , but the partial differentials of  $d_i S$  can be less than zero.

Under more general situations, irreversible processes usually are the mixed behaviors of the above processes that we discussed respectively. Combine with Eqs.(90), (91), (94), (97), and (103), we obtain

$$\begin{aligned} d_i S &= \Delta \left( \frac{1}{T} \right) dq + \frac{1}{T} \Delta Y dx \\ &+ \sum_j \frac{1}{T} \Delta \mu'_j dN_j + \Delta \left( \frac{p}{T} \right) dV \geq 0. \end{aligned} \quad (109)$$

There are three positive definite sources of the entropy production: (1). Heat energy transfer from higher temperature to lower temperature; (2). The independent process driven by the heat partial pressure; (3). The free energy is converted into the heat energy without compensation.

In an irreversible process, the total entropy production is always positive definite, i.e.,  $d_i S \geq 0$ , but the partial differentials of which can be less than zero, the second law itself has already contained the mechanism of evolution, it implies that order and evolution can arise spontaneously.

In non-equilibrium thermodynamics, the increment in the entropy of an open system

$$dS = d_e S + d_i S, \quad (110)$$

has been demonstrated that the total entropy of an open system can be reduced by the aid of the negative entropy flux  $d_e S$ , it is an external mechanism of evolution. From Eq.(99) and Eq.(106), we have found the internal mechanism of evolution, i.e., in the equation

$$d_i S = d_i S_q + d_i S_v \geq 0, \quad (111)$$

the partial differentials  $d_i S_q$  or  $d_i S_v$  can be less than zero.

As an explanation,  $pdV$  is defined as the change of the conversion potential related to the dissipative compensation, and by Eq.(62),  $p/T$  involves a density function or a configurational function.

For a non-equilibrium state, consider the spatial distribution of the gradients, it follows that

$$\begin{aligned} d_i S &= \nabla \left( \frac{1}{T} \right) dq + \frac{1}{T} \nabla Y dx \\ &+ \sum_j \frac{1}{T} \nabla \mu'_j dN_j + \nabla \left( \frac{p}{T} \right) dV \geq 0. \end{aligned} \quad (112)$$

Eq.(109) and Eq.(112) are the total differential expressions of the second law of thermodynamics.

As a conclusion, the second law of thermodynamics indicates such a fact: For all of thermodynamic processes, the total entropy production of the processes  $d_i S \geq 0$ , and as a reason, the second law comes from such a fundamental principle: All of the gradients of the thermodynamic forces spontaneously tend to zero, in an irreversible process, the dissipations of the gradients of the thermodynamic forces cannot be spontaneously recovered.

## DISCUSSION: A NEW FOUNDATION

Now, we are witnessing a profound change that has been brought into thermodynamics, it is a deep change in the way of thinking, we expect to push the restrictions of the traditional way of thinking, and establish a new theoretical structure. Start from the rediscovery of the first law, we need to re-understand thermodynamics, and answer those deep problems that can hardly continue to be discussed in the traditional theoretical structure.

In a new theoretical structure, we have ascertained the concept of the heat energy  $q$ , the last one of the undefined forms of the internal energy, combine with the mechanical free energy and Gibbs free energy, we obtain

$$dU = dq + d(Yx) + \sum_j d(\mu'_j N_j). \quad (113)$$

The first law has been extended into

$$dU = d_e q + d_e A + d_e G, \quad (114)$$

$$d_i q + d_i A + d_i G = 0. \quad (115)$$

The energy transfer and the energy conversion can be easily distinguished by the different explicit equations.

From Eq.(28), we obtain a binomial expression, it is an explicit definition of the entropy

$$dS = \frac{dq}{T} + \frac{p}{T} dV. \quad (116)$$

For an ideal gas or a photon gas, a primary equation of the entropy could be given by

$$dS = \frac{\hat{\alpha} N R_{\{k\}}}{q} dq + \frac{N R_{\{k\}}}{V} dV. \quad (117)$$

It is a binomial distribution function, and the entropy is proportional to the average degrees of the binomial distribution. Perhaps, it represents the real nature of the entropy.

In Eq.(112), we confirm that the driving forces of the spontaneous processes come from the different gradients of the thermodynamic forces that

$$\nabla \left( \frac{1}{T} \right), \quad \nabla Y, \quad \nabla \mu'_j, \quad \nabla \left( \frac{p}{T} \right).$$

The second law of thermodynamics is derived from a fundamental principle: the gradients of the thermodynamic forces spontaneously tend to zero, and in an irreversible process, the dissipations of the gradients of the thermodynamic forces cannot be spontaneously recovered, and in a coupling process, the dissipation of one gradient can drive the production of another gradient, this fact implies, the second law contains the mechanism of evolution.

Thermodynamics involves the two main lines, one is the law of conservation of energy, the another one involves evolution and dissipation, which involves the sum of all microscopic processes, it will be a self-consistent theory, and should be mutually consistent with dynamics.

For an ideal gas or a photon gas, the natures of the temperature  $T$  and the pressure  $p$  can be described by

$$T \rightarrow \frac{1}{R_{\{k\}}} \cdot \frac{q}{\hat{\alpha} N}, \quad p \rightarrow \frac{q}{\hat{\alpha} V}, \quad \left( q = \sum_{j=1}^N \epsilon_j \right)$$

where  $\epsilon_j$  is the kinetic energy of the particle  $j$ .

Temperature  $T$  is the function of  $q/N$ , and the heat partial pressure  $p$  is the function of  $q/V$ .

For Eq.(117), an interesting inference can be made

$$\frac{\hat{\alpha} N R_{\{k\}}}{q} dq \rightarrow R_{\{k\}} \sum_{j=1}^N \sum_{s=1}^s \frac{\hat{\alpha}}{i} d \ln \epsilon_{j\{s\}},$$

where  $s$  denotes the degrees of freedom of the particles.

The master equations of dynamics are established on a fundamental principle: the law of conservation of energy, and the first law of thermodynamics is the total of these master equations, the present dynamics involves only one main line, the law of conservation of energy, so it makes no sense to discuss irreversibility in the present dynamics, it is similar to discuss the same problems only according to the first law of thermodynamics. In the present dynamics, there has no similar theoretical structure of the second law.

This theme depend on such a foundation: the physical property of the entropy has been made clear, we can verify whether the similar theoretical structure of the second law of thermodynamics that can be found in real dynamic processes, before that, there are no solutions to the problems. Now, we are approaching to this foundation.

In a new theoretical structure, we might be able to find a new principle to re-understand being and becoming, and meanwhile, we tend to regard the second law as a law of evolution, and the total entropy production  $d_i S \geq 0$  as the spending of the evolution processes [9, 11, 12].

Such an attempt may be the beginning of a new study, perhaps, the primary structure of theoretical physics might be distinguished into the two main lines according to the different laws [9, 12]: one being the first law physics, the other one being the second law physics [9]. We can do, or cannot do, that is the problem.

---

\* Email: tangsuye@hotmail.com

- [1] I.Prigogine What is entropy? Naturwissenschaften, Vol.76,1. (1989)
- [2] R.Clausius Ann. Phys. Chem. **125** 353 (1865)
- [3] A.Einstein Physics and Reality (1936)
- [4] J.C.Maxwell Theory of Heat. (London: Longmans Green, 1872)
- [5] R.Feynman Lectures on Physics, Vol. 1. (Addison-Welsey, 1963)
- [6] D.Bernoulli Hydrodynamica. (1738)
- [7] L.E.Reichl A Modern Course in Statistical Physics, 2nd ed. (New York: John Wiley & Sons Inc, 1997)
- [8] J.Fourier The Analytical Theory of Heat. (London: Cambridge University Press, 1878)
- [9] Tang Suye Tackling a Century Mystery: Entropy, 2nd ed. (Hefei: University of Science & Technology of China, 2008)
- [10] I.Prigogine Science. **201** 4358 (1978)
- [11] I.Prigogine Structure, Dissipation and Life. Theoretical Physics and Biology, Versailles, 1967, (Amsterdam: North-Holland Publ. Company 1969)
- [12] I.Prigogine From Being to Becoming: Time and Complexity in the Physical Sciences.(New York: W.H.Freeman, 1980)