

E.M. Ovsiyuk, N.V. Gutsko, V.M. Red'kov*

Transitivity in polarization optics and the diagonalization of quadratic forms

Mozyr State Pedagogical University named after I.P. Shamyakin, Belarus

B.I. Stepanov Institute of Physics, NAS of Belarus

November 15, 2018

Abstract

Any one measurement with polarized light makes it possible to fix the Mueller matrices of the Lorentz type with up to four arbitrary numeric parameters $(x, u; z, w)$. These parameters are subject to the quadratic condition. It is demonstrated that the quadratic form can be diagonalized; in the case of partially polarized light four diagonal coefficients turn out to be non-zero and positive; in the case of completely polarized light two diagonal coefficients equal to zero.

In [1, 2] it was shown that any one measurement with polarized light makes it possible to fix the Mueller matrices of the Lorentz type with up to four arbitrary numeric parameters $x, u; z, w$. These parameters are subject to the quadratic condition

$$\begin{aligned} & x^2(A^2 - \mathbf{B}^2) + 2xyA(B^2 - \mathbf{A}^2) + y^2[(\mathbf{A}^2 + \mathbf{B}^2 - B^2)\mathbf{A}^2 - A^2B^2] = \\ & = z^2(B^2 - \mathbf{A}^2) + 2zwB(A^2 - \mathbf{B}^2) + w^2[(\mathbf{A}^2 + \mathbf{B}^2 - A^2)\mathbf{B}^2 - A^2B^2] + 1. \end{aligned} \quad (1)$$

The quantities $A, \mathbf{A}, B, \mathbf{B}$ are defined by the initial and final Stokes 4-vectors

$$A = S + S', \quad B = S - S', \quad \mathbf{A} = \mathbf{S} + \mathbf{S}', \quad \mathbf{B} = \mathbf{S} - \mathbf{S}'$$

*e.ovsiyuk@mail.ru; gutsko-nv@yandex.ru; redkov@dragon.bas-net.by

$$\begin{aligned}
S > 0, \quad S' > 0, \quad S^2 - \mathbf{S}^2 = S'^2 - \mathbf{S}'^2 = \text{inv} = +V^2 \geq 0 \\
A^2 = S_0^2 + S_0'^2 + 2S_0S_0' = s^2 + s'^2 + 2ss', \\
B^2 = S_0^2 + S_0'^2 - 2S_0S_0' = s^2 + s'^2 - 2ss', \\
\mathbf{A}^2 = s^2 + s'^2 - 2V^2 + 2\sqrt{s^2 - V^2}\sqrt{s'^2 - V^2} \cos \phi \\
\mathbf{B}^2 = s^2 + s'^2 - 2V^2 - 2\sqrt{s^2 - V^2}\sqrt{s'^2 - V^2} \cos \phi
\end{aligned} \tag{2}$$

numeric parameters $x, u; z, w$ determine the parameters of possible Mueller matrices of the Lorentzian type according to

$$\begin{aligned}
n_0 = Ax - \mathbf{A}^2y, \quad \mathbf{n} = z\mathbf{A} - w\mathbf{A}\mathbf{B} + y\mathbf{A} \times \mathbf{B}, \\
m_0 = -Bz + \mathbf{B}^2w, \quad \mathbf{m} = x\mathbf{B} - y\mathbf{B}\mathbf{A} + w\mathbf{A} \times \mathbf{B}.
\end{aligned} \tag{3}$$

Note the different dimensions of the coordinates x, y, z, w :

$$[x] = [S]^{-1}, \quad [y] = [S]^{-2}, \quad [z] = [S]^{-1}, \quad [w] = [S]^{-2}.$$

All 6 terms in (1) have such a structure such the dimensions of parameters x, z and y, w are mutually compensated with corresponding dimensions $[S]$ and $[S]^2$ variables formed from the initial and final Stokes 4-vectors, and therefore all the quantities in equation (1) are dimensionless.

The task of the present paper is to explore the geometry of the surface (1) in 4-dimensional space (x, y, z, w) . In equation (1) we can see two quadratic forms: in variables (x, y) and (z, W) , so one can transform them to a diagonal form:

$$\begin{vmatrix} x \\ y \end{vmatrix} = \begin{vmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{vmatrix} \begin{vmatrix} X \\ Y \end{vmatrix}, \quad \begin{vmatrix} z \\ w \end{vmatrix} = \begin{vmatrix} \cos \alpha' & \sin \alpha' \\ -\sin \alpha' & \cos \alpha' \end{vmatrix} \begin{vmatrix} Z \\ W \end{vmatrix}. \tag{4}$$

Substituting (4) into (1) and imposing requirements for the diagonalization of two quadratic forms, we arrive at the following relations

$$\text{tg } 2\alpha = \frac{2a}{c-b}, \quad \text{tg } 2\alpha' = \frac{2a'}{c'-b'}. \tag{5}$$

$$\begin{aligned}
& X^2(b \cos^2 \alpha - a \sin 2\alpha + c \sin^2 \alpha) + Y^2(b \sin^2 \alpha + a \sin 2\alpha + c \cos^2 \alpha) = \\
& = Z^2(b' \cos^2 \alpha' - a' \sin 2\alpha' + c' \sin^2 \alpha') + W^2(b' \sin^2 \alpha' + a' \sin 2\alpha' + c' \cos^2 \alpha') + 1,
\end{aligned} \tag{6}$$

where the notation is used

$$(A^2 - \mathbf{B}^2) = b, \quad A(B^2 - \mathbf{A}^2) = a, \quad (\mathbf{A}^2 + \mathbf{B}^2 - B^2)\mathbf{A}^2 - A^2B^2 = c,$$

$$(B^2 - \mathbf{A}^2) = b', \quad B(A^2 - \mathbf{B}^2) = a', \quad (\mathbf{B}^2 + \mathbf{A}^2 - A^2)\mathbf{B}^2 - A^2B^2 = c'. \quad (7)$$

Further, taking into account the elementary relations

$$\begin{aligned} \cos 2\alpha &= \pm \frac{1}{\sqrt{1 + \operatorname{tg}^2 2\alpha}} = \pm \frac{(c-b)}{\sqrt{(c-b)^2 + 4a^2}}, \\ \sin 2\alpha &= \pm \frac{2a}{\sqrt{(c-b)^2 + 4a^2}}, \\ \cos^2 \alpha &= \frac{\sqrt{(c-b)^2 + 4a^2} \pm (c-b)}{2\sqrt{(c-b)^2 + 4a^2}}, \\ \sin^2 \alpha &= \frac{\sqrt{(c-b)^2 + 4a^2} \mp (c-b)}{2\sqrt{(c-b)^2 + 4a^2}} \end{aligned}$$

we arrive at (let $\delta, \delta' = \pm 1$)

$$\begin{aligned} &X^2 \left(\frac{b+c}{2} - \delta \frac{\sqrt{(b+c)^2 - 4(bc-a^2)}}{2} \right) + \\ &+ Y^2 \left(\frac{b+c}{2} + \delta \frac{\sqrt{(b+c)^2 - 4(bc-a^2)}}{2} \right) - \\ &- Z^2 \left(\frac{b'+c'}{2} - \delta' \frac{\sqrt{(b'+c')^2 - 4(b'c'-a'^2)}}{2} \right) - \\ &- W^2 \left(\frac{b'+c'}{2} + \delta' \frac{\sqrt{(b'+c')^2 - 4(b'c'-a'^2)}}{2} \right) = 1. \quad (8) \end{aligned}$$

First, we assume isotropic Stokes vectors: $V^2 = 0$. This will simplify the analysis, since

$$\begin{aligned} A^2 &= s^2 + s'^2 + 2ss', & B^2 &= s^2 + s'^2 - 2ss', \\ \mathbf{A}^2 &= s^2 + s'^2 + 2ss' \cos \phi, & \mathbf{B}^2 &= s^2 + s'^2 - 2ss' \cos \phi. \end{aligned}$$

We calculate

$$\begin{aligned} b &= 2ss'(1 + \cos \phi), & c &= (s + s')^2 2ss'(1 + \cos \phi). \\ b + c &= 2ss'(1 + \cos \phi) + (s + s')^2 2ss'(1 + \cos \phi) \geq 0. \\ a^2 &= (s^2 + s'^2 + 2ss')(2ss')^2 (1 + \cos \phi)^2, & bc - a^2 &= 0. \quad (9a) \end{aligned}$$

Therefore, if $\delta = +1$, in (8) the coefficient at X^2 is zero and the coefficient at Y^2 is positive ($b + c > 0$). When $\delta = +1$, the coefficient at Y^2 is zero, and the coefficient at X^2 is positive ($b + c > 0$).

In this case (for definiteness, let $\delta = +1$)

$$\begin{aligned}\cos^2 \alpha &= \frac{c}{c+b} = \frac{(s+s')^2}{1+(s+s')^2}, \\ \sin^2 \alpha &= \frac{b}{c+b} = \frac{1}{1+(s+s')^2}.\end{aligned}\quad (9b)$$

Analyzing the coefficients of the Z^2, W^2 , we get

$$\begin{aligned}b' &= -2ss'(1 + \cos \phi) < 0, & c' &= -(s - s')^2 2ss'(1 + \cos \phi) < 0, \\ b' + c' &= -2ss'(1 + \cos \phi) - (s - s')^2 2ss'(1 + \cos \phi) \leq 0, \\ a'^2 &= (s - s')^2 (2ss')^2 (1 + \cos \phi)^2, & b'c' - a'^2 &= 0.\end{aligned}\quad (10a)$$

Thus, in (8) the coefficient at Z^2 is zero and the coefficient at W^2 is negative $b' + c' \leq 0$ (we assume $\delta' = +1$). In this case

$$\begin{aligned}\cos^2 \alpha' &= \frac{c'}{c'+b'} = \frac{(s-s')^2}{1+(s-s')^2}, \\ \sin^2 \alpha' &= \frac{b'}{c'+b'} = \frac{1}{1+(s-s')^2}.\end{aligned}\quad (10b)$$

Therefore, the set of Mueller matrices, connecting two states completely polarized light, are described by the relation

$$0 X^2 + (b+c)Y^2 - 0 Z^2 - (b'+c')W^2 = +1. \quad (11a)$$

The whole set of Mueller matrices, defined by the pair of isotropic Stokes vectors can be specified by following three independent variables:

$$\begin{aligned}X, Z &\text{ are arbitrary,} & Y &= \frac{\cos \Gamma}{\sqrt{+(b+c)}}, \\ W &= \frac{\sin \Gamma}{\sqrt{-(b'+c')}} , & \Gamma &\in [0, 2\pi].\end{aligned}\quad (11b)$$

Let us consider another simple case. Let an initial beam is the natural light, then

$$S_0 = s, \quad \mathbf{S} = 0, \quad V^2 = s^2, \quad s' > s,$$

$$\begin{aligned}
A^2 &= s^2 + s'^2 + 2ss', & B^2 &= s^2 + s'^2 - 2ss', \\
\mathbf{A}^2 &= s'^2 - s^2, & \mathbf{B}^2 &= s'^2 - s^2.
\end{aligned} \tag{12a}$$

We calculate

$$\begin{aligned}
b &= 2s^2 + 2ss' = 2s(s' + s) > 0, & c &= 2s(s' + s)(s' - s)^2 > 0, \\
a^2 &= 4s^2(s' + s)^2(s' - s)^2, & bc - a^2 &= 0.
\end{aligned} \tag{12b}$$

The coefficient at X^2 is zero, and that at Y^2 equals $(b + c)$:

$$b + c = +2s(s' + s)[1 + (s' - s)^2], \tag{12c}$$

in this case

$$\begin{aligned}
\cos^2 \alpha &= \frac{c}{c + b} = \frac{(s' - s)^2}{1 + (s' - s)^2}, \\
\sin^2 \alpha &= \frac{b}{c + b} = \frac{1}{1 + (s' - s)^2}.
\end{aligned} \tag{12d}$$

Similarly, examine the coefficients at Z^2 and W^2 :

$$\begin{aligned}
b' &= -2s(s' - s) < 0, & c' &= -2s(s + s')(s'^2 - s^2) < 0, \\
a'^2 &= 4s^2(s' - s)^2(s' + s)^2, & b'c' - a'^2 &= 0.
\end{aligned} \tag{13a}$$

The coefficient at Z^2 is equal to zero, and the coefficient at W^2 equals $(b' + c')$:

$$b' + c' = -2s(s' - s)[1 + (s' + s)^2], \tag{13b}$$

in this case

$$\begin{aligned}
\cos^2 \alpha' &= \frac{c'}{c' + b'} = \frac{(s' + s)^2}{1 + (s' + s)^2}, \\
\sin^2 \alpha' &= \frac{b'}{c' + b'} = \frac{1}{1 + (s' + s)^2}.
\end{aligned} \tag{13c}$$

The basic quadratic equation takes the form

$$0 X^2 + (b + c)Y^2 - 0 Z^2 - (b' + c')W^2 = +1, \tag{14a}$$

$$\begin{aligned}
+(b + c) &= 2s(s' + s)[1 + (s' - s)^2], \\
-(b' + c') &= 2s(s' - s)[1 + (s' + s)^2].
\end{aligned} \tag{14b}$$

Let us consider another simple case, when an initial beam is partially polarized, and the final beam is the natural light:

$$s' = V, \quad s > s', \quad \mathbf{A}^2 = s^2 - s'^2, \quad \mathbf{B}^2 = s^2 - s'^2,$$

$$\begin{aligned}
A^2 &= s^2 + s'^2 + 2ss', & B^2 &= s^2 + s'^2 - 2ss'. \\
b &= +2s'(s + s') > 0, & b' &= 2s'^2 - 2ss' = -2s'(s - s') < 0, \\
c &= 2s'(s - s')(s^2 - s'^2) > 0, & c' &= -2s'(s + s')(s^2 - s'^2) < 0, \\
a^2 &= 4s'^2(s + s')^2(s' - s)^2, & a'^2 &= 4s'^2(s - s')^2(s' + s)^2. \quad (15a)
\end{aligned}$$

We see that the equalities $bc - a^2 = 0$, $b'c' - a'^2 = 0$. The basic quadratic equation takes the form

$$0 X^2 + (b + c)Y^2 - 0 Z^2 - (b' + c')W^2 = +1. \quad (15b)$$

Let us consider another simple case: a set of Mueller matrices that do not change of light intensity $s' = s$; in this case

$$\begin{aligned}
A^2 &= 4s^2, & B^2 &= 0, \\
\mathbf{A}^2 &= 2(s^2 - V^2)(1 + \cos \phi), \\
\mathbf{B}^2 &= 2(s^2 - V^2)(1 - \cos \phi). \quad (16a)
\end{aligned}$$

The basic quadratic equation takes the form

$$\begin{aligned}
&x^2(A^2 - \mathbf{B}^2) - 2xyAA^2 + y^2(\mathbf{A}^2 + \mathbf{B}^2)\mathbf{A}^2 + \\
&\quad + 2(s^2 - V^2)(1 + \cos \phi) Z^2 + \\
&\quad + 8V^2(s^2 - V^2)(1 - \cos \phi) W^2 = +1. \quad (16b)
\end{aligned}$$

We need only to diagonalize the first quadratic form. After calculation we get

$$\begin{aligned}
b &= 2 \left[V^2(1 - \cos \phi) + s^2(1 + \cos \phi) \right] > 0, \\
c &= 8(s^2 - V^2)^2(1 + \cos \phi) > 0, \\
a^2 &= 16s^2(s^2 - V^2)^2(1 + \cos \phi)^2, \\
bc - a^2 &= 16 V^2 \sin^2 \phi (s^2 - V^2)^2 > 0. \quad (16c)
\end{aligned}$$

It follows that the first quadratic form is also reduced to diagonal form with positive coefficients.

A situation where the initial and final light beams are partially polarized, is the most difficult. The coefficients of a diagonalized quadratic form can be calculated explicitly and they are positive.

Let us detail this subject as well. First, We find

$$b = 2V^2 + 2ss' \left(1 + \sqrt{1 - V^2/s^2} \sqrt{1 - V^2/s'^2} \cos \phi \right) > 2V^2 \geq 0,$$

$$c = [(s + s')^2 - 4V^2] \times \\ \times \left[(s - s')^2 - 2V^2 + 2ss' \left(1 + \sqrt{1 - V^2/s^2} \sqrt{1 - V^2/s'^2} \cos \phi \right) \right] - \\ - (s + s')^2 (s - s')^2 .$$

In turn, we obtain

$$b + c = \left(2ss' + 2 \cos \phi \sqrt{s^2 - V^2} \sqrt{s'^2 - V^2} \right) \left(1 + [(s + s')^2 - 4V^2] \right) + \\ + 2V^2 - 2V^2 [(s + s')^2 - 4V^2] - 4V^2 (s - s')^2 . \quad (17a)$$

At $\phi = \pi$ we have the minimal values for $(b + c)$:

$$(b + c)_{min} = \left(2ss' - 2\sqrt{s^2 - V^2} \sqrt{s'^2 - V^2} \right) \left(1 + [(s + s')^2 - 4V^2] \right) + \\ + 2V^2 \left(1 - [(s + s')^2 - 4V^2] \right) - 4V^2 (s - s')^2 ; \quad (17b)$$

Numerical calculation shows that the quantity $(b + c)_{min}$ is positive (see Fig.1)




Fig. 1: $(b + c)_{min}(V = 1)$

Additionally let us calculate

$$a^2 = (s^2 + s'^2 + 2ss') (2V^2 - 2ss' - 2\sqrt{s^2 - V^2} \sqrt{s'^2 - V^2} \cos \phi)^2 . \quad (18a)$$

and

$$bc - a^2 = -16 V^2 \sin^2 \phi (s^2 V^2 + s'^2 V^2 - s^2 s'^2 - V^4) = \\ = 16 V^2 \sin^2 \phi (s^2 - V^2)(s'^2 - V^2) > 0 \quad (18b)$$

Turning to coefficients at X^2, Y^2 in (8)

$$\begin{aligned}
& X^2 \left(\frac{b+c}{2} - \delta \frac{\sqrt{(b+c)^2 - 4(bc-a^2)}}{2} \right) + \\
& + Y^2 \left(\frac{b+c}{2} + \delta \frac{\sqrt{(b+c)^2 - 4(bc-a^2)}}{2} \right) - \dots = 1 \quad (18c)
\end{aligned}$$

we conclude that they both are positive

Now we analyze coefficient at Z^2, W^2 . After simple calculation we get

$$\begin{aligned}
b' &= -2ss' + 2V^2 - 2 \cos \phi \sqrt{s^2 - V^2} \sqrt{s'^2 - V^2} = \\
c' &= (s^2 + s'^2 - 4V^2 - 2ss') (s^2 + s'^2 - 2V^2 - 2 \cos \phi \sqrt{s^2 - V^2} \sqrt{s'^2 - V^2}) - \\
& \quad - (s^2 + s'^2 + 2ss') (s^2 + s'^2 - 2ss') . \\
b' + c' &= - (2ss' + 2 \cos \phi \sqrt{s^2 - V^2} \sqrt{s'^2 - V^2}) (1 + [(s - s')^2 - 4V^2]) + \\
& \quad + 2V^2 - 2V^2 [(s - s')^2 - 4V^2] - 4V^2 (s + s')^2 . \quad (19a)
\end{aligned}$$

Numerical calculation shows that the quantity $(b' + c')_{max}$ is negative (see Fig.2)




Fig. 2: $(b' + c')_{max}(V = 1)$

Additionally we find

$$\begin{aligned}
a'^2 &= (s^2 + s'^2 - 2ss') (2V^2 + 2ss' + 2 \cos \phi \sqrt{s^2 - V^2} \sqrt{s'^2 - V^2})^2 . \\
b'c' - a'^2 &= -16 V^2 \sin^2 \phi (s^2 V^2 + s'^2 V^2 - s^2 s'^2 - V^4) =
\end{aligned}$$

$$= 16 V^2 \sin^2 \phi (s^2 - V^2)(s'^2 - V^2) > 0 \quad (19b)$$

Therefore, coefficients at $Z^2 W^2$ in (8) are positive

$$\begin{aligned} & \dots - Z^2 \left(\frac{b' + c'}{2} - \delta' \frac{\sqrt{(b' + c')^2 - 4(b'c' - a'^2)}}{2} \right) - \\ & - W^2 \left(\frac{b' + c'}{2} + \delta' \frac{\sqrt{(b' + c')^2 - 4(b'c' - a'^2)}}{2} \right) = 1. \end{aligned} \quad (19c)$$

Conclusion: Any one measurement with polarized light makes it possible to fix the Mueller matrices of the Lorentz type with up to four arbitrary numeric parameters $(x, u); (z, w)$. These parameters are subject to the quadratic condition. It is demonstrated that both quadratic forms, in variables (x, y) and (z, w) , can be diagonalized; in the case of partially polarized light four diagonal coefficients turn out to be positive; in the case of completely polarized light two diagonal coefficient equal to zero.

REFERENCES

1. V.M. Red'kov, E.M. Ovsuiyk. Transitivity in the theory of the Lorentz group and the Stokes - Mueller formalism in optics. // Proceedings of 15th International School & Conference "Foundation & Advances in Nonlinear Science. Eds.: Kuvshinov V.I., Krylov G.G., September 19-22, 2010, Minsk. P. 1-27. - <http://arxiv.org/abs/1005.4212>. - [math-ph; physics.optics]; 22 pages.
2. V.M. Red'kov, E.M. Ovsuiyk. On determining Mueller matrices of an optical element by results of polarization measurements // Proceedings of of III-th International Conference "Optics of inhomogeneous structures-2011", 16-17 February 2011, Mogilev. Eds. B.A. Karpenko and others P. 32-35.