# The DJ method for exact solutions of Laplace equation

M. Yaseen, M. Samraiz, S. Naheed

Department of Mathematics, University of Sargodha, Sargodha, Pakistan.

#### Abstract

In this paper, the iterative method developed by Daftardar-Gejji and Jafari (DJ method) is employed for analytic treatment of Laplace equation with Dirichlet and Neumann boundary conditions. The method is demonstrated by several physical models of Laplace equation. The obtained results show that the present approach is highly accurate and requires reduced amount of calculations compared with the existing iterative methods.

### 1 Introduction

The Daftardar-Jafari method (DJM) developed in 2006 has been extensively used by many researchers for the treatment of linear and nonlinear ordinary and partial differential equations of integer and fractional order [1–7]. An excellent modification of DJM can be found in [8] for solving different models of linear and nonlinear Klein-Gordon equations. The method converges to the exact solution if it exists through successive approximations. For concrete problems, a few number of approximations can be used for numerical purposes with high degree of accuracy. The DJM does not require any restrictive assumptions for nonlinear terms as required by some existing techniques. The aim of this work is to effectively employ DJM to obtain exact solutions for different models of Laplace equation. While the variational iteration method [9–11] requires the determination of Lagrange multiplier in its computational algorithm, DJM is independent of any such requirements. Moreover, unlike the Adomian decomposition method [12–20], where the calculation of the tedious Adomian polynomials is needed to deal with nonlienar terms, DJM handles linear and nonlinear terms in a simple and straightforward manner without any additional requirements.

As mentioned before, we aim to obtain exact solutions of four models of Laplace equation, two with Dirichlet boundary conditions and two with Neumann boundary conditions. We also aim to establish that DJM is powerful, promising and efficient in handling nonlinear engineering problems. In what follows, we give a brief review of DJM.

## 2 The DJ Method

Consider the following general functional equation

(2.1) 
$$u(\overline{x}) = f(\overline{x}) + N(u(\overline{x}))$$

where N is a nonlinear operator from a Banach space  $B \to B$  and f is a known function.  $\overline{x} = (x_1, x_2, \dots, x_n)$ . We are looking for a solution u of (2.1)having the series form

(2.2) 
$$u(\overline{x}) = \sum_{n=0}^{\infty} u_i(\overline{x})$$

The nonlinear operator N can be decomposed as

(2.3) 
$$N(\sum_{i=0}^{\infty} u_i) = N(u_0) + \sum_{i=1}^{\infty} \{N(\sum_{j=0}^{i})u_j - N(\sum_{j=0}^{i-1} u_j)\}.$$

From equations (2.2) and (2.3), equation (2.1) is equivalent to

(2.4) 
$$\sum_{i=0}^{\infty} u_i = f + N(u_0) + \sum_{i=1}^{\infty} \{ N(\sum_{j=0}^i) u_j - N(\sum_{j=0}^{i-1} u_j) \}$$

We define the recurrence relation

(2.5) 
$$\begin{cases} u_0 = f \\ u_1 = N(u_0) \\ u_{m+1} = N(u_0 + \dots + u_m) - N(u_0 + \dots + u_{m-1}) \end{cases}$$

Then

(2.6) 
$$(u_1 + \dots + u_m) = N(u_0 + \dots + u_m)$$

and

(2.7) 
$$\sum_{i=0}^{\infty} u_i = f + N(\sum_{i=0}^{\infty} u_i)$$

The k-term approximate solution of (2.1) is given by  $u = u_0 + u_1 + \cdots + u_{k-1}$ In what follows, we apply DJM to four physical models to demonstrate the strength of the method and to establish exact solutions of these models.

Example 1. Consider the second order laplace equation,

(2.8) 
$$u_{xx} - u_{yy} = 0, \ 0 < x, y < \pi$$

with boundary conditions

(2.9) 
$$u(x,0) = \sinh(x), \ u(x,\pi) = -\sinh(x) u(0,y) = 0, \ u(\pi,y) = \sinh(\pi)\cos(y)$$

The equation (2.8) is equivalent to the following integral equation

$$u = \sinh(x) + yg(x) - \int_{0}^{y} \int_{0}^{y} (u_{xx}(x, y)) dy dy$$

where  $g(x) = u_y(x, 0)$ . Set  $u_0 = \sinh(x) + yg(x)$  and  $N(u) = -\int_0^y \int_0^y (u_{xx}(x, y)) dy dy$ .

Following the algorithm (2.5), the successive approximations are

$$u_{1} = N(u_{0}) = -\frac{1}{2!}y^{2}\sinh(x) - \frac{1}{3!}y^{3}g''(x)$$
  

$$u_{2} = N(u_{0} + u_{1}) - N(u_{0}) = \frac{1}{4!}y^{4}\sinh(x) + \frac{1}{5!}y^{5}g^{iv}(x)$$
  

$$u_{3} = N(u_{0} + u_{1} + u_{2}) - N(u_{0} + u_{1}) = -\frac{1}{6!}y^{6}\sinh(x) - \frac{1}{7!}y^{7}g^{vi}(x)$$
  
:

Combining the above results obtained for the components yields

$$u(x,y) = \sinh(x) + yg(x) - \frac{1}{2!}y^2 \sinh(x) - \frac{1}{3!}y^3 g''(x) + \frac{1}{4!}y^4 \sinh(x) + \frac{1}{5!}y^5 g^{iv}(x) + \cdots$$

Using the inhomogeneous boundary condition,  $u(\pi, y) = \sinh(\pi) \cos(y)$  and the Taylor expansion of  $\cos(y)$ , we obtain

$$\sinh(\pi) + yg(\pi) - \frac{1}{2}y^2 \sinh(\pi) - \frac{1}{6}y^3 g''(\pi) + \frac{1}{24}y^4 \sinh(\pi) + \dots = \sinh(\pi)(1 - \frac{y^2}{2!} + \frac{y^4}{4!} - \dots)$$

Equating the coefficients of like terms on both sides gives

$$g(\pi) = g''(\pi) = g^{iv}(\pi) = \dots = 0$$

This means that g(x) = 0. Consequently, the solution in series form is given by

$$u(x,y) = \sinh(x)(1 - \frac{y^2}{2!} + \frac{y^4}{4!} - \cdots)$$

and in closed form

$$u(x, y) = \sinh(x)\cos(y)$$

which is also the exact solution.

**Example 2.** Consider the second order laplace equation,

$$(2.10) u_{xx} - u_{yy} = 0, \ 0 < x, \ y < \pi$$

with boundary conditions

(2.11) 
$$u(x,0) = 0, \ u(x,\pi) = 0 u(0,y) = \sin(y), u(\pi,y) = \cosh(\pi)\sin(y)$$

The equation (2.10) is equivalent to the following integral equation

$$u(x,y) = yg(x) - \int_{0}^{y} \int_{0}^{y} (u_{xx}(x,y)) dy dy$$

where  $g(x) = u_y(x, 0)$ . Set  $u_0 = yg(x)$  and  $N(u) = -\int_0^y \int_0^y (u_{xx}(x, y)) dy dy$ . Following the algorithm (2.5), the successive approximations are

$$u_{1} = N(u_{0}) = -\frac{1}{3!}y^{3}g''(x)$$
  

$$u_{2} = N(u_{0} + u_{1}) - N(u_{0}) = \frac{1}{5!}y^{5}g^{iv}(x)$$
  

$$u_{3} = N(u_{0} + u_{1} + u_{2}) - N(u_{0} + u_{1}) = -\frac{1}{7!}y^{7}g^{vi}(x)$$
  
:

Combining the above results, we obtain

$$u(x,y) = yg(x) - \frac{1}{3!}y^3g''(x) + \frac{1}{5!}y^5g^{iv}(x) - \frac{1}{7!}y^7g^{vi}(x) \cdots$$

Using the inhomogeneous boundary condition,  $u(\pi, y) = \cosh(\pi) \sin(y)$  and the Taylor expansion of  $\sin(y)$ , we obtain

$$yg(\pi) - \frac{1}{3!}y^3g''(\pi) + \frac{1}{5!}y^5g^{iv}(\pi) - \frac{1}{7!}y^7g^{vi}(\pi) + \dots = \cosh(\pi)(y - \frac{y^3}{3!} + \frac{y^5}{5!} - \dots).$$

Equating the coefficients of like terms on both sides gives

$$g(\pi) = g''(\pi) = g^{iv}(\pi) = \dots = \cosh(\pi)$$

This means that

$$g(x) = \cosh(x)$$

Consequently, the solution in series form is given by

$$u(x,y) = \cosh(x)(y - \frac{y^3}{3!} + \frac{y^5}{5!} - \cdots)$$

and in closed form

$$u(x,y) = \cosh(x)\sin(y)$$

which is also the exact solution.

**Example 3.** Consider the second order laplace equation,

$$(2.12) u_{xx} - u_{yy} = 0, \ 0 < x, y < \pi$$

with boundary conditions

(2.13) 
$$\begin{aligned} u_y(x,0) &= 0, \ u_y(x,\pi) = 2\cos(2x)\sinh(2\pi) \\ u_x(0,y) &= 0, \ u_x(\pi,y) = 0 \end{aligned}$$

The equation (2.12) is equivalent to the following integral equation

$$u = g(x) - \int_{0}^{y} \int_{0}^{y} (u_{xx}(x,y)) dy dy$$

where g(x) = u(x, 0). Set  $u_0 = g(x)$  and  $N(u) = -\int_0^y \int_0^y (u_{xx}(x, y)) dy dy$ .

Following the algorithm (2.5), the successive approximations are

$$\begin{split} & u_1 = N(u_0) = -\frac{1}{2!} y^2 g''(x) \\ & u_2 = N(u_0 + u_1) - N(u_0) = \frac{1}{4!} y^4 g^{iv}(x) \\ & u_3 = N(u_0 + u_1 + u_2) - N(u_0 + u_1) = -\frac{1}{6!} y^6 g^{vi}(x) \\ & \vdots \end{split}$$

Combining the above results, we obtain

(2.14) 
$$u(x,y) = g(x) - \frac{1}{2!}y^2g''(x) + \frac{1}{4!}y^4g^{iv}(x) - \frac{1}{6!}y^6g^{vi}(x) + \cdots$$

Using the inhomogeneous boundary condition,  $u_y(x,\pi) = 2\cos(2x)\sinh(2\pi)$ and the Taylor expansion of  $\sinh(2\pi)$ , we obtain

$$-\pi g''(x) + \frac{1}{3!}\pi^3 g^{iv}(x) - \frac{1}{5!}\pi^5 g^{vi}(x) + \dots = 2\cos(2x)\left(2\pi + \frac{8\pi^3}{3!} + \frac{32\pi^5}{5!} + \dots\right)$$

Equating the coefficients of like terms on both sides gives

$$g''(x) = -4\cos(2x)$$

which gives

$$g(x) = \cos(2x) + C_0$$

Consequently, the solution in series form is given by

$$u(x,y) = \cos(2x)\left(1 + \frac{4y^2}{2!} + \frac{16y^4}{4!} + \cdots\right) + C_0$$

and in closed form

$$u(x,y) = \cos(2x)\cosh(2y) + C_0$$

which is also the exact solution.

**Example 4.** Consider the second order laplace equation,

$$(2.15) u_{xx} - u_{yy} = 0, \ 0 < x, y < \pi$$

with boundary conditions

(2.16) 
$$\begin{aligned} u_y(x,0) &= \cos(x), \ u_y(x,\pi) &= \cosh(\pi)\cos(x) \\ u_x(0,y) &= 0, \ u_x(\pi,y) &= 0 \end{aligned}$$

The equation (2.15) is equivalent to the following integral equation

$$u = g(x) + y\cos(x) - \int_{0}^{y} \int_{0}^{y} (u_{xx}(x,y))dydy$$

where g(x) = u(x, 0). Set  $u_0 = g(x) + y \cos(x)$  and  $N(u) = -\int_0^y \int_0^y (u_{xx}(x, y)) dy dy$ . Following the algorithm (2.5), the successive approximations are

$$u_{1} = N(u_{0}) = -\frac{1}{2!}y^{2}g''(x) + \frac{1}{3!}y^{3}\cos(x)$$
  

$$u_{2} = N(u_{0} + u_{1}) - N(u_{0}) = \frac{1}{4!}y^{4}g^{iv}(x) + \frac{1}{5!}y^{5}\cos(x)$$
  

$$u_{3} = N(u_{0} + u_{1} + u_{2}) - N(u_{0} + u_{1}) = -\frac{1}{6!}y^{6}g^{vi}(x) + \frac{1}{7!}y^{7}\cos(x)$$
  

$$\vdots$$

Combining the above results, we obtain

$$(2.17) \ u(x,y) = \cos(x)\sinh(y) + g(x) - \frac{1}{2!}y^2g''(x) + \frac{1}{4!}y^4g^{iv}(x) - \frac{1}{6!}y^6g^{vi}(x)$$

Using the inhomogeneous boundary condition,  $u_y(x,\pi)=\cosh(\pi)\cos(x)$  , we obtain

$$\cos(x)\cosh(\pi) - \pi g''(x) + \frac{1}{3!!}\pi^3 g^{iv}(x) - \frac{1}{6!}\pi^5 g^{vi}(x) + \dots = \cosh(\pi)\cos(x)$$

Equating the coefficients of like terms on both sides gives

$$g''(x) = g^{iv}(x) = g^{vi}(x) = \dots = 0$$

which gives

$$g(x) = C_0$$

Consequently, the solution in series form is given by

$$u(x,y) = \cos(x)(y + \frac{y^3}{3!} + \frac{y^5}{5!} + \dots) + C_0$$

and in closed form

$$u(x,y) = \cos(x)\sinh(y) + C_0$$

which is also the exact solution.

### References

- Varsha Daftardar-Gejji, Hossein Jafri, An iterative method for solving nonlinear functional equations, J. Math. Appl. **316** (2006) 753.
- [2] Sachin Bhalekar, Varsha Daftardar-Gejji, New iterative method: Application to partial differntial equations, Appl. Math. Comput. 203 (2008) 778.
- [3] V. Daftardar-Gejji and S. Bhalekar, Solving fractional boundary value problems with Dirichlet boundary conditions using a new iterative method, Computers & Mathematics with Applications, 59 (2010) 1801.
- [4] S. Bhalekar and V. Daftardar-Gejji, New iterative method: application to partial differential equations, Applied Mathematics and Computation, 203 (2008) 778.
- [5] V. Daftardar-Gejji and S. Bhalekar, An Iterative method for solving fractional differential equations, Proceedings in Applied Mathematics and Mechanics, 7 (2008) 20500172050018,.
- [6] S. Bhalekar and V. Daftardar-Gejji, Solving evolution equations using a new iterative method, Numerical Methods for Partial Differential Equations, 26 (2010) 906.
- [7] H. Jafari, S. Seifi, A. Alipoor, and M. Zabihi, An Iterative Method for solving linear and nonlinear fractional diffusion-wave equation, International e-Journal of Numerical Analysis and Related Topics, 3 (2009) 20.
- [8] M. Yaseen, M. Samraiz, A modified new iterative method for solving linear and nonlinear Klein-Gordon Equations, Appl. Math. Sci. 6 (2012) 2979.
- [9] A.-M. Wazwaz, The variational iteration method for exact solutions of Laplace equations, Phy. Letters. A. 363 (2007) 260.
- [10] J. H. He, Variational iteration method-a kind of nonlinear analytical technique: some examples, Int. J. Nonlin. Mech. 34 (1999) 699.
- [11] A.-M. Wazwaz, The variational iteration method for rational solutions for KdV, K(2, 2), Burgers, and cubic Boussinesq equations, J. Comput. Appl. Math. 207 (2007) 18.
- [12] G. Adomian, Solving Frontier Problems of Physics: The Decomposition Method, Kluwer, 1994.
- [13] D. Kaya, An implementation of the ADM for generalized one dimensional Klein-Gordon equation, Appl. Math. Comput. 166 (2005) 426.
- [14] K.C. Basak, P.C. Ray, R.K. Bera, Solution of nonlinear Klein-Gordon equation with a quadratic nonlinear term by Adomian decomposition method, Comm. Nonlin. Sci.Num. Simul. 14 (2009) 718.

- [15] A. wazwaz, Partial Differential Equations: Methods and Applications, Netherland Balkema, 2002.
- [16] A. wazwaz, Partial Differential Equations and Solitary Wave Theory, Springer, 2009.
- [17] M. Hussain, M. Khan, A variational iteraive method for solving linear and nonlinear Klein-Gordon Equations, Appl. Math. Sci. 4 (2010) 1931.
- [18] E. Yusufoğla, The variational iteration method for studying the Klein-Gordon equation, Appl. Math. lett. 21 (2008) 669.
- [19] M.A.Abdou, A.A. Soliman, Variational iteraion method for solving Burger's and coupled Burger's equations, J. Comp. Appl. Math. 181 (2005) 245.
- [20] Z. Odibat, S. Momani, A reliable treatment of homotopy pertubation method for Klein-Gordon equations, Phys. Lett. A. 365 (2007) 351.