# Reduction operators and exact solutions of variable coefficient nonlinear wave equations with power nonlinearities

Ding-jiang Huang  $^{\dagger,\ddagger,\S}$ ,Qin-min Yang  $^{\dagger}$  and Shui-geng Zhou  $^{\ddagger,\S}$ 

#### Abstract

Reduction operators, i.e. the operators of nonclassical (or conditional) symmetry of a class of variable coefficient nonlinear wave equations with power nonlinearities is investigated within the framework of singular reduction operator. A classification of regular reduction operators is performed with respect to generalized extended equivalence groups. Exact solutions of some nonlinear wave model which are invariant under certain reduction operators are also constructed.

**Keywords:** symmetry analysis, reduction operators, equivalence group, nonlinear wave equation, exact solutions

Mathematics Subject Classifications (2000): 35L10, 35A22, 35A30

## 1 Introduction

In this paper, we study reduction operators, i.e., the operators of nonclassical (or conditional) symmetry associated with a class of variable coefficient nonlinear wave equations with power nonlinearities of the form

$$f(x)u_{tt} = (q(x)u^{n}u_{x})_{x} + h(x)u^{m}, (1)$$

where f = f(x), g = g(x) and h = h(x) are three arbitrary functions,  $fg \neq 0$ , n and m are arbitrary constants, t is the time coordinate and x is the one-space coordinate. The linear case is excluded from consideration because it was well-investigated. We also assume the variable wave speed coefficient  $u^n$  to be nonlinear, i.e.  $n \neq 0$ . The case n = 0 is quite singular and will be investigated separately.

Many specific nonlinear wave models describing a wide variety of phenomena in Mechanics and Engineering such as the flow of one-dimensional gas, shallow water waves theory, longitudinal wave propagation on a moving threadline, dynamics of a finite nonlinear string, elastic-plastic materials and electromagnetic transmission line and so on, can be reduced to equation (1) (see [1] p.50-52 and [2]). Since 1970s, Lie symmetries and invariant solutions of various kinds of quasi-linear wave equations in two independent variables that intersect class (1) have been investigated [3–20] because of the importance of the wave equation for various applications. Recently, we have present a complete Lie symmetry and conservation law classification of class (1) [21,22]. Classical Lie symmetry reduction and invariant solutions of some variable coefficients wave models which are singled out from the classification results are also investigated [21].

In general, using classical Lie symmetries reduction of partial differential equations can provide part of exact solutions of these equations [23, 24]. In order to find more other types exact

<sup>&</sup>lt;sup>†</sup> Department of Mathematics, East China University of Science and Technology, Shanghai, 200237, China

 $e\text{-}mail:\ djhuang.math@gmail.com,\ djhuang@fudan.edu.cn$ 

<sup>&</sup>lt;sup>‡</sup> School of Computer Science, Fudan University, Shanghai, 200433, China

<sup>§</sup> Shanghai Key Lab of Intelligent Information Processing, Fudan University, Shanghai, 200433, China

solutions, one should generalize Lie's original reduction. The first approach to such generalization was present by Bluman and Cole in 1969 [25] (see also [3]) in which they introduced a wider class of infinitesimal generators than Lie symmetries. Later such infinitesimal generators were named nonclassical symmetries [26] or conditional symmetries [27–31], and were also extended by many authors to some concepts such as weak symmetry [32] or differential constraints, etc., [33–37]. Recently, Popovych et.al named it as 'reduction operators' and present a novel framework, namely singular reduction operators or singular reduction modules [38,39], for finding an optimal way of obtaining the determining equation of conditional symmetries. As application, they have investigated the properties of singular reduction operators for a number of (1+1)-dimensional evolution equations and a specific wave equations [39–46] by using this new framework. However, for more general nonlinear wave equation (1), there exist no general results. In this paper, we employ Popovych's singular reduction operators theory to investigate the properties of nonclassical symmetries of class (1). We propose a complete classification of regular reduction operators for (1) with respect to generalized extended equivalence groups and construct some non-Lie exact solutions for the nonlinear wave model which are invariant under certain reduction operators. Below, following [39] we use the shorter and more natural term 'reduction operators' instead of 'operators of conditional symmetry' or 'operators of nonclassical symmetry'.

The rest of paper is organized as follows. In Section 2, singular reduction operators, and in particular regular reduction operators classification for the class under consideration are investigated. Section 3 contains nonclassical symmetry reduction of some nonlinear wave models. New non-Lie exact solutions of the models are constructed by means the reduction. Conclusions and discussion are given in section 4.

## 2 Nonclassical symmetries

Nonclassical symmetries of class (1) is performed in the framework of the singular reduction operator [39]. All necessary objects (singular and regular reduction operator, etc.) can be found there [39]. Before we proceed the investigation, we can first simplify the class (1). Using the transformation

$$\tilde{t} = t, \quad \tilde{x} = \int \frac{dx}{g(x)}, \quad \tilde{u} = u$$
 (2)

from theorem 1 in [21], we can reduce equation (1) to  $\tilde{f}(\tilde{x})\tilde{u}_{\tilde{t}\tilde{t}}=(\tilde{u}^n\tilde{u}_{\tilde{x}})_{\tilde{x}}+\tilde{h}(\tilde{x})\tilde{u}^m$ , where  $\tilde{f}(\tilde{x})=g(x)f(x), \tilde{g}(\tilde{x})=1$  and  $\tilde{h}(\tilde{x})=g(x)h(x)$ . Thus, without loss of generality we can restrict ourselves to investigation of the equation

$$f(x)u_{tt} = (u^n u_x)_x + h(x)u^m. (3)$$

For convenient, we can further rewrite it as the form

$$L[u] := f(x)u_{tt} - (u^n u_x)_x - h(x)u^m = 0.$$
(4)

All results on symmetries and solutions of class (3) or (4) can be extended to class (1) with transformations (2).

According to the algorithm in [39], we seek a reduction operator of class (4) in the form

$$Q = \tau(t, x, u)\partial_t + \xi(t, x, u)\partial_x + \eta(t, x, u)\partial_u, \quad (\tau, \xi) \neq (0, 0), \tag{5}$$

which is a first-order differential operator on the space  $\mathbb{R}^2 \times \mathbb{R}^1$  with coordinates t, x, and u, where the coefficients  $\tau$  and  $\xi$  do not simultaneously vanish. This operator allows one to construct an

ansatz reducing the original equation (4) to an ordinary differential equation. The conditional invariance criterion [29–31] for equation (4) to be invariant with respect to the operator (5) read as

$$\operatorname{pr}^{(2)} Q(L[u]) \Big|_{\mathcal{L} \cap \mathcal{Q}_{(2)}} = 0, \tag{6}$$

where  $\operatorname{pr}^{(2)}Q$  is the usual second order prolongation [23,24] of the operator (5),  $\mathcal{L}$  is the manifold in the second-order jet space  $J^{(2)}$  determined by the wave equation L[u] = 0, and  $\mathcal{Q}_{(2)} \subset J^{(2)}$  is the the first prolongation of the invariant surface condition

$$Q[u] := \tau u_t + \xi u_x - \eta = 0. \tag{7}$$

The system  $Q_{(2)}$  consists of (7) and the equations obtained by t- and x-differentiation of (7). Below, according to the singular reduction operator theory [39], we first partition the set of reduction operators of class (4) into two subsets, i.e., the singular reduction operator and the regular one. Then we utilize the two kinds of operators to derive determining equations (overdetermined system of nonlinear PDEs with respect to the coefficients of the reduction operator (5)) from the conditional invariance criterion (6) separately. Solving the two systems we can obtain the final reduction operators. In particular, we will present a exhausted classification of the regular operators of class (4) by solving the corresponding determining equations. In general, every Lie symmetry operator is also a reduction operator. Therefore, in this paper we will concentrate on the regular reduction operators which is inequivalent to Lie symmetry operators, called nontrivial one.

#### 2.1 Singular reduction operators

Using the procedure given by Popovych et al. in [39], we can obtain the following assertion.

**Proposition 1.** A vector field  $Q = \tau(t, x, u)\partial_t + \xi(t, x, u)\partial_x + \eta(t, x, u)\partial_u$  is singular for the differential function  $L = f(x)u_{tt} - (u^n u_x)_x - h(x)u^m$  if and only if  $\xi^2 f(x) = \tau^2 u^n$ .

*Proof.* Suppose that  $\tau \neq 0$ . According to the characteristic equation  $\tau u_t + \xi u_x - \eta = 0$ , we can get

$$u_{t} = \frac{\eta}{\tau} - \frac{\xi}{\tau} u_{x},$$

$$u_{tt} = (\frac{\eta}{\tau})_{t} - (\frac{\xi}{\tau})_{t} u_{x} + [(\frac{\eta}{\tau})_{u} - (\frac{\xi}{\tau})_{u} u_{x}](\frac{\eta}{\tau} - \frac{\xi}{\tau} u_{x}) - (\frac{\xi}{\tau})[(\frac{\eta}{\tau})_{x} + (\frac{\eta}{\tau})_{u} u_{x} - (\frac{\xi}{\tau})_{x} u_{x}]$$

$$- (\frac{\xi}{\tau})_{u} u_{x}^{2} - (\frac{\xi}{\tau}) u_{xx}].$$

Substituting the formulas of  $u_{tt}$  from above formulaes into L, we obtain a differential function

$$\tilde{L} = [f(x)(\frac{\xi}{\tau})^2 - u^n]u_{xx} + f(x) \left\{ (\frac{\eta}{\tau})_t - (\frac{\xi}{\tau})_t u_x + [(\frac{\eta}{\tau})_u - (\frac{\xi}{\tau})_u u_x](\frac{\eta}{\tau} - \frac{\xi}{\tau}u_x) - (\frac{\xi}{\tau})[(\frac{\eta}{\tau})_x + (\frac{\eta}{\tau})_u u_x - (\frac{\xi}{\tau})_x u_x - (\frac{\xi}{\tau})_u u_x^2] \right\} - nu^{n-1}u_x^2 - hu^m u_x.$$

According to the definition 4 of singular vector field in [39], we have ord  $\tilde{L} < 2$  if and only if  $f(x)(\frac{\xi}{\tau})^2 - u^n = 0$ .

Therefore, for any f, h, n and m with  $fu^n > 0$  the differential function  $L = f(x)u_{tt} - (u^n u_x)_x - h(x)u^m$  possesses exactly two set of singular vector fields in the reduced form, namely,  $S = \{\partial_t + \sqrt{u^n/f}\partial_x + \hat{\eta}\partial_u\}$  and  $S^* = \{\partial_t - \sqrt{u^n/f}\partial_x + \hat{\eta}\partial_u\}$ , where  $\hat{\eta} = \frac{\eta}{\tau}$ . Any singular vector field of L is equivalent to one of the above fields. The singular sets are mapped to each other by alternating the sign of x and hence one of them can be excluded from the consideration.

**Proposition 2.** For any variable coefficient nonlinear wave equations in the form (4) the differential function  $L = f(x)u_{tt} - (u^n u_x)_x - h(x)u^m$  possesses exactly one set of singular vector fields in the reduced form, namely,  $S = \{\partial_t + \sqrt{u^n/f}\partial_x + \hat{\eta}\partial_u\}$ .

Thus taking into accountant the conditional invariance criterion for an equation from class (4) and the operator  $\partial_t + \sqrt{u^n/f}\partial_x + \eta\partial_u$ , we can get

**Theorem 1.** Every singular reduction operator of an equation from class (4) is equivalent to an operator of the form

$$Q = \partial_t + \sqrt{u^n/f(x)}\partial_x + \eta(t, x, u)\partial_u,$$

where the real-valued function  $\eta(t,x,u)$  satisfies the determining equations

$$(-1/2nhu^{m-1} - 2f\eta_{tu} - 2f\eta\eta_{uu})\sqrt{u^n/f} + (3/4f_x^2/f - 1/2f_{xx})(u^n/f)^{3/2} + (-1/4n^2\eta^2u^{n-2} - n\eta\eta_uu^{n-1} + 1/2n\eta^2u^{n-2})(u^n/f)^{-1/2} - n\eta_xu^{n-1} - 1/2n\eta u^{n-1}f_x/f - 2\eta_{xu}u^n = 0,$$

$$-\eta_{xx}u^n - n\eta\eta_xu^{n-1}(u^n/f)^{-1/2} + h\eta_uu^m + 2f\eta\eta_{tu} + f\eta^2\eta_{uu} - mh\eta u^{m-1} + f\eta_{tt} + (hf_x/f - h_x)u^m\sqrt{u^n/f} = 0.$$
(8)

#### 2.2 Regular reduction operators

The above investigation of singular reduction operators of nonlinear wave equation of the form (4) shows that for these equations the regular case of the natural partition of the corresponding sets of reduction operators is singled out by the conditions  $\xi \neq \pm \sqrt{u^n/f}\tau$ . After factorization with respect to the equivalence relation of vector fields, we obtain the defining conditions of regular subset of reduction operator:  $\tau = 1, \xi \neq \pm \sqrt{u^n/f}$ . Hence we have

**Proposition 3.** For any variable coefficient nonlinear wave equations in the form (4) the differential function  $L = f(x)u_{tt} - (u^n u_x)_x - h(x)u^m$  possesses exactly one set of regular vector fields in the reduced form, namely,  $S = \{\partial_t + \hat{\xi}\partial_x + \hat{\eta}\partial_u\}$  with  $\hat{\xi} \neq \pm \sqrt{u^n/f}$ .

Consider the conditional invariance criterion for an equation from class (4) and the operator  $\partial_t + \xi(t, x, u)\partial_x + \eta(t, x, u)\partial_u$  with  $\xi(t, x, u) \neq \pm \sqrt{u^n/f}$ , we can get the following determining equations for the coefficients  $\xi$  and  $\eta$ :

$$\xi_{u} = 0, \quad 2f\xi_{t} - n\eta u^{n-1} + (2\xi_{x} + \xi f_{x}/f)u^{n} = 0, 
(2n\xi_{x} - n\eta_{u} + n\xi f_{x}/f)u^{n-1} + (n\eta - n^{2}\eta)u^{n-2} - \eta_{uu}u^{n} + f\xi^{2}\eta_{uu} = 0, 
2f\xi_{t}\xi_{x} - 2f\xi_{t}\eta_{u} - 2n\eta_{x}u^{n-1} - f\xi_{tt} - 2f\xi\eta\eta_{uu} - 2f\xi\eta_{tu} + (\xi_{xx} - 2\eta_{xu})u^{n} = 0, 
(\xi hf_{x}/f - \xi h_{x} + h\eta_{u})u^{m} + f\eta^{2}\eta_{uu} + 2f\eta\eta_{tu} - 2f\xi_{t}\eta_{x} - \eta_{xx}u^{n} + f\eta_{tt} - mh\eta u^{m-1} = 0.$$
(9)

From the first two equations of system (9), we have

$$\xi = \xi(t, x), \quad \eta = \frac{1}{n} 2f \xi_t u^{1-n} + \frac{1}{n} (2\xi_x + \xi \frac{f_x}{f})u.$$

Substituting these expression into the last three equations of system (9), we have the following assertion.

**Theorem 2.** Every regular reduction operator of an equation from class (4) is equivalent to an operator of the form

$$Q = \partial_t + \xi(t, x)\partial_x + \eta(t, x, u)\partial_u \quad \text{with} \quad \eta(t, x, u) = \frac{1}{n}2f\xi_t u^{1-n} + \frac{1}{n}(2\xi_x + \xi \frac{f_x}{f})u, \tag{10}$$

where the real-valued function  $\xi(t,x)$  satisfies the overdetermined system of partial differential equations

$$2(1-n)f\xi_{t} = 0, \quad 2(1-n)f^{2}\xi^{2}\xi_{t} = 0, \quad 8(1-n)f^{3}\xi\xi_{t}^{2} = 0,$$

$$4(1-n)[(f\xi_{t})^{2} - f^{2}\xi\xi_{t}(2\xi_{x} + \xi\frac{f_{x}}{f}) + f^{2}\xi\xi_{tt}] = 0,$$

$$\xi_{xx} - 2(1 + \frac{1}{n})(2\xi_{x} + \xi\frac{f_{x}}{f})_{x} = 0,$$

$$2f\xi_{t}\xi_{x} - \frac{2}{n}f\xi_{t}(2\xi_{x} + \xi\frac{f_{x}}{f}) - 4(f\xi_{t})_{x} - f\xi_{tt} - \frac{2}{n}f\xi(2\xi_{x} + \xi\frac{f_{x}}{f})_{t} - \frac{4}{n}(1-n)(f\xi_{t})_{x} = 0,$$

$$(\xi h\frac{f_{x}}{f} - \xi h_{x} + \frac{1}{n}h(1-m)(2\xi_{x} + \xi\frac{f_{x}}{f}))u^{m} + \frac{2}{n}(1-n-m)fh\xi_{t}u^{m-n}$$

$$-\frac{8}{n^{2}}(1-n)f^{4}(\xi_{t})^{3}u^{1-3n} + \frac{8}{n^{2}}(1-n)[f^{2}\xi_{t}(f\xi_{t})_{t} - f^{3}\xi_{t}^{2}(2\xi_{x} + \xi\frac{f_{x}}{f})]u^{1-2n}$$

$$+[\frac{2}{n}f(f\xi_{t})_{tt} - \frac{4}{n}f\xi_{t}(f\xi_{t})_{x} + \frac{4}{n^{2}}f^{2}\xi_{t}(2\xi_{x} + \xi\frac{f_{x}}{f})_{t} - \frac{2}{n^{2}}(1-n)f^{2}\xi_{t}(2\xi_{x} + \xi\frac{f_{x}}{f})^{2}$$

$$+\frac{4}{n^{2}}(1-n)f(f\xi_{t})_{t}(2\xi_{x} + \xi\frac{f_{x}}{f})]u^{1-n} + [\frac{2}{n^{2}}f(2\xi_{x} + \xi\frac{f_{x}}{f})(2\xi_{x} + \xi\frac{f_{x}}{f})_{t}$$

$$+\frac{1}{n}f(2\xi_{x} + \xi\frac{f_{x}}{f})_{tt} - \frac{2}{n}f\xi_{t}(2\xi_{x} + \xi\frac{f_{x}}{f})_{x} - \frac{2}{n}(f\xi_{t})_{xx}]u - \frac{1}{n}(2\xi_{x} + \xi\frac{f_{x}}{f})_{xx}u^{n+1} = 0.$$

Solving the above system with respect to the coefficient functions  $\xi$ , f and h under the equivalence group  $G_1^{\sim}$  of the class (4) which consists of the transformations (see theorem 3 and 4 in [21] for more details): for  $n \neq -1$ 

$$\tilde{t} = \epsilon_1 t + \epsilon_2, \quad \tilde{x} = \frac{\epsilon_3 x + \epsilon_4}{\epsilon_5 x + \epsilon_6} =: X(x), \quad \tilde{u} = \epsilon_7 X_x^{\frac{1}{2n+2}} u,$$

$$\tilde{f} = \epsilon_1^2 \epsilon_7^n X_x^{-\frac{3n+4}{2n+2}} f, \quad \tilde{h} = \epsilon_7^{-m+n+1} X_x^{-\frac{m+3n+3}{2n+2}} h, \quad \tilde{n} = n, \quad \tilde{m} = m,$$

where  $\epsilon_i$   $(j=1,\ldots,7)$  are arbitrary constants,  $\epsilon_1\epsilon_7\neq 0,\epsilon_3\epsilon_6-\epsilon_4\epsilon_5=\pm 1$  and for n=-1

$$\begin{split} \tilde{t} &= \epsilon_1 t + \epsilon_2, \quad \tilde{x} = \epsilon_3 x + \epsilon_4, \quad \tilde{u} = \epsilon_5 e^{\epsilon_6 x} u, \\ \tilde{f} &= \epsilon_1^2 \epsilon_3^{-2} \epsilon_5^{-1} e^{-\epsilon_6 x} f, \quad \tilde{h} = \epsilon_3^{-2} \epsilon_5^{-m} e^{-m\epsilon_6 x} h, \quad \tilde{n} = n, \quad \tilde{m} = m, \end{split}$$

where  $\epsilon_j$  (j = 1, ..., 6) are arbitrary constants,  $\epsilon_1 \epsilon_3 \epsilon_5 \neq 0$ ; we can get a classification of regular reduction operator for the class (4). It is easy to know that some of the regular reduction operator are equivalent to Lie symmetry operators, while some of are nontrivial. Below, we give a detailed investigations for these cases.

In fact, the first three equations of system (11) implies there are two cases should be considered:  $n \neq 1$  or not. (It should be noted that  $\xi = 0$  should be exclude from the consideration because it leads to  $\eta = 0$ ).

Case 1:  $n \neq 1$ . In this case, we have  $\xi_t = 0$ . Thus system (11) can be reduced to

$$(3n+4)\xi_{xx} + 2(n+1)(\xi\frac{f_x}{f})_x = 0,$$
  

$$(\xi h\frac{f_x}{f} - \xi h_x + \frac{1}{n}(1-m)h(2\xi_x + \xi\frac{f_x}{f}))u^m - \frac{1}{n}(2\xi_x + \xi\frac{f_x}{f})_{xx}u^{n+1} = 0.$$
(12)

Thus, there are two cases should be considered:  $m \neq n+1$  or not.

Case 1.1: For  $m \neq n+1$ , from the second equation of (12) we obtain

$$\xi h \frac{f_x}{f} - \xi h_x + \frac{1}{n} (1 - m) h(2\xi_x + \xi \frac{f_x}{f}) = 0, \quad \left(2\xi_x + \xi \frac{f_x}{f}\right)_{xx} = 0.$$
 (13)

The first equation of (12) suggests that  $(3n+4)\xi_x + 2(n+1)\xi\frac{f_x}{f}$  is independent of the variable x, so there exists a constant r such that  $(3n+4)\xi_x + 2(n+1)\xi\frac{f_x}{f} = nr$ . The second equation of

(13) suggests that there exist two constants a and b such that  $2\xi_x + \xi \frac{f_x}{f} = nax + nb$ . By solving the last two equations we obtain

$$\xi_x = 2(n+1)(ax+b) - r, \quad \xi \frac{f_x}{f} = 2r - (3n+4)(ax+b),$$

which together with the first equation of (13) imply

$$\xi = a(n+1)x^{2} + [2b(n+1) - r]x + s,$$

$$f(x) = \exp\left(\int \frac{2r - (3n+4)(ax+b)}{a(n+1)x^{2} + [2b(n+1) - r]x + s} dx\right),$$

$$h(x) = \exp\left(\int \frac{2r - (m+3n+3)(ax+b)}{a(n+1)x^{2} + [2b(n+1) - r]x + s} dx\right),$$
(14)

where a, b, r, s are arbitrary constants. Thus, the corresponding regular reduction operator has the form

$$Q = \partial_t + [a(n+1)x^2 + (2b(n+1) - r)x + s]\partial_x + (ax+b)u\partial_u,$$

which is equivalent to Lie symmetry operator.

Case 1.2: m = n + 1. In this case, system (12) can be rewritten as

$$(3n+4)\xi_{xx} + 2(n+1)(\xi\frac{f_x}{f})_x = 0,$$
  

$$\xi h_x + 2h\xi_x + \frac{1}{n}(2\xi_x + \xi\frac{f_x}{f})_{xx} = 0.$$
(15)

Integrating these two equations with respect to functions f(x) and g(x), we can obtain

$$f(x) = |\xi|^{-\frac{3n+4}{2n+2}} \exp\left(r \int \frac{1}{\xi} dx\right), \quad h(x) = \frac{\xi_x^2 - 2\xi \xi_{xx} - p}{4(n+1)\xi^2}.$$

where p, r are arbitrary constants,  $\xi$  is an arbitrary smooth function and  $n \neq -1$ . In addition,  $\eta = \frac{1}{n}(2\xi_x + \xi \frac{f_x}{f})u = (\frac{r}{n} + \frac{\xi_x}{2n+2})u$ . Thus, we have a nontrivial regular reduction operator

$$Q = \partial_t + \xi(x)\partial_x + \left[\left(\frac{r}{n} + \frac{\xi_x}{2n+2}\right)u\right]\partial_u, \quad n \neq -1.$$
(16)

It should be noted that for n = -1 the reduction operator is also equivalent to Lie symmetry operator.

Case 2: n = 1. In this case, we have  $\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u$ . Thus, system (11) can be reduced to

$$(7\xi_{x} + 4\xi \frac{f_{x}}{f})_{x} = 0, \quad 2\left[\xi_{x} + 2(\xi + 1)\frac{f_{x}}{f}\right]\xi_{t} + 4(\xi + 1)\xi_{tx} + \xi_{tt} = 0,$$

$$\left[2f(2\xi_{x} + \xi \frac{f_{x}}{f})(2\xi_{x} + \xi \frac{f_{x}}{f})_{t} + f(2\xi_{x} + \xi \frac{f_{x}}{f})_{tt} - 2f\xi_{t}(2\xi_{x} + \xi \frac{f_{x}}{f})_{x} - 2(f\xi_{t})_{xx}\right]u$$

$$-(2\xi_{x} + \xi \frac{f_{x}}{f})_{xx}u^{2} + 2f^{2}(2\xi_{t}\xi_{tx} + \xi_{ttt}) + \left[\xi h \frac{f_{x}}{f} - \xi h_{x} + (1 - m)h(2\xi_{x} + \xi \frac{f_{x}}{f})\right]u^{m}$$

$$-2mhf\xi_{t}u^{m-1} = 0.$$

$$(17)$$

After some brief analysis, we find that there are five cases should be considered.

Case 2.1: m = 0. In this case, the third equation of (17) implies

$$2h\xi \frac{f_x}{f} - h_x \xi + 2h\xi_x + 4f^2 \xi_t \xi_{tx} + 2f^2 \xi_{ttt} = 0,$$

$$2f(2\xi_x + \xi \frac{f_x}{f})(2\xi_x + \xi \frac{f_x}{f})_t + f(2\xi_x + \xi \frac{f_x}{f})_{tt} - 2f\xi_t (2\xi_x + \xi \frac{f_x}{f})_x - 2(f\xi_t)_{xx} = 0,$$

$$(2\xi_x + \xi \frac{f_x}{f})_{xx} = 0.$$
(18)

From the last equation of system (18) we can know that there exist two functions a(t) and b(t) such that  $2\xi_x + \xi \frac{f_x}{f} = a(t)x + b(t)$ . On the other hand, the first equation of (17) implies there exists a function c(t) such that  $7\xi_x + 4\xi \frac{f_x}{f} = c(t)$ . Solving the last two equations gives

$$\xi_x = 4a(t)x + 4b(t) - c(t), \quad \xi \frac{f_x}{f} = -7a(t)x - 7b(t) + 2c(t),$$

from which we can get

$$\xi = 2a(t)x^{2} + 4b(t)x - c(t)x + d(t),$$

$$f(x) = \exp\left(\int \frac{-7a(t)x - 7b(t) + 2c(t)}{2a(t)x^{2} + 4b(t)x - c(t)x + d(t)} dx\right).$$
(19)

where d(t) is an arbitrary functions. Since  $\frac{f_x}{f}$  is independent of t, we see that

$$\[\frac{-7a(t)x - 7b(t) + 2c(t)}{2a(t)x^2 + 4b(t)x - c(t)x + d(t)}\]_t = 0,$$

which leads to

$$\begin{cases}
14[a(t)b'(t) - a'(t)b(t)] + 3[a'(t)c(t) - a(t)c'(t)] = 0, \\
[b(t)c'(t) - b'(t)c(t)] + 7[a(t)d'(t) - a'(t)d(t)] = 0, \\
2[c'(t)d(t) - c(t)d'(t)] + 7[b(t)d'(t) - b'(t)d(t)] = 0.
\end{cases}$$
(20)

Now, we multiply both sides of the second equation of (17) by  $\xi$  and substitute (19) into it, then simplify the equation and compare the coefficient of  $x^i (i = 0, 1, ..., 5)$  to obtain

$$\begin{cases} a'(t) = 0, \\ a^{2}[-4b'(t) + c'(t)] = 0, \\ a[8c(t)c'(t) + 20ad'(t) + c''(t) - 4b''(t) - 32c(t)b'(t) + 112b(t)b'(t) - 28b(t)c'(t)] = 0, \\ -4c''(t)b(t) + 32ac(t)d'(t) - 80ab'(t) + 20ac'(t) - 4b''(t)c(t) - 120ab(t)d'(t) \\ +c''(t)c(t) + 2d''(t)a + 2c^{2}(t)c'(t) - 12b(t)c(t)c'(t) + 16b(t)^{2}c'(t) - 8c^{2}(t)b'(t) \\ +48b(t)c(t)b'(t) + 16b''(t)b(t) - 64b^{2}(t)b'(t) - 16ad(t)b'(t) + 4ad(t)c'(t) = 0, \\ [-20ad'(t) + 48b(t)b'(t) + 2c(t)c'(t) - 12b(t)c'(t) - 8c(t)b'(t) + 4b''(t) - c''(t)]d(t) \\ -80b^{2}(t)d'(t) + 4d''(t)b(t) - d''(t)c(t) - 4c(t)c'(t) + 44b(t)c(t)d'(t) - 28ad'(t) \\ -48b(t)b'(t) - 6c^{2}(t)d'(t) + 12b(t)c'(t) + 16c(t)b'(t) = 0, \\ [16b'(t) - 4c'(t)]d^{2}(t) + [-20b(t)d'(t) - 4c'(t) + 6c(t)d'(t) + d''(t) + 16b'(t)]d(t) \\ +8c(t)d'(t) - 28b(t)d'(t) = 0. \end{cases}$$

Note that  $\xi$  is assumed not to be identical with zero, after some simple but lengthy computations, we find that systems (20) and (21) can be reduced to:

$$a'(t) = 0, \quad b'(t) = 0, \quad c'(t) = 0, \quad d'(t) = 0$$
 (22)

or

$$a = 0$$
,  $c(t) = 4b(t)$ ,  $d(t) = qb(t)$ ,  $qb''(t) + 4qb(t)b'(t) + 4b'(t) = 0$  (23)

or

$$a = 0$$
,  $2c(t) = 7b(t)$ ,  $b''(t) = -3b(t)b'(t)$ ,  $b(t)d'(t) + 2b'(t)(d(t) + 1) + d''(t) = 0$  (24)

or

$$a = 0, \quad c(t) = 3b(t), \quad d = qb(t), \quad b''(t) + 2b(t)b'(t) = 0,$$
 (25)

where q is an arbitrary constant.

Case 2.1a: If system (22) is satisfied, then  $\xi_t = 0$ , the second equation of (18) is an identity. The expression (19) can be rewritten as

$$\xi = 2ax^{2} + 4bx - cx + d,$$

$$f(x) = \exp\left(\int \frac{-7ax - 7b + 2c}{2ax^{2} + 4bx - cx + d} dx\right),$$
(26)

where a, b, c and d are arbitrary constants. The first equation of (18) is reduced to

$$\frac{h_x}{h} = \frac{2\left(\xi \frac{f_x}{f} + \xi_x\right)}{\xi}.$$

Substitute the expression of  $\xi$  and f(x) into it and integrate both sides to obtain

$$h(x) = \exp\left(\int \frac{-6ax - 6b + 2c}{2ax^2 + 4bx - cx + d} dx\right).$$

In addition,  $\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u = (ax + b)u$ . Therefore, we have

$$\xi = 2ax^{2} + 4bx - cx + d,$$

$$\eta = (ax + b)u,$$

$$f(x) = \exp\left(\int \frac{-7ax - 7b + 2c}{2ax^{2} + 4bx - cx + d} dx\right),$$

$$h(x) = \exp\left(\int \frac{-6ax - 6b + 2c}{2ax^{2} + 4bx - cx + d} dx\right).$$
(27)

where a,b,c,d are arbitrary constants. Thus, the corresponding regular reduction operator has the form

$$Q = \partial_t + (2ax^2 + 4bx - cx + d)\partial_x + (ax + b)u\partial_u,$$

which is equivalent to Lie symmetry operator.

Case 2.1b: If system (23) is satisfied, then the expression (19) can be rewritten as

$$\xi = qb(t), \quad f(x) = \exp\left(\frac{x}{q}\right).$$
 (28)

Hence,  $\xi_x = 0$ , k = b(t). Substituting these formulaes into the second equation of (18) we obtain

$$qb''(t) + 2qb(t)b'(t) - 2b'(t) = 0.$$

Combine it with the fourth equation of (23) to get b'(t) = 0. Hence a(t), b(t), c(t), d(t) satisfy system (22), and the solution is included in the case 2.1a.

Case 2.1c: If system (24) is satisfied, then the expression (19) can be rewritten as

$$\xi = \frac{1}{2}b(t)x + d(t), \quad f(x) = 1 \mod G_1^{\sim}.$$
 (29)

Substitute it into the second equation of (18) to obtain 2b(t)b'(t) + b''(t) = 0. Combine it with the third equation of (24) to get b'(t) = 0. Substitute it into the fourth equation of (24)

to obtain bd'(t) + d''(t) = 0, which implies  $d(t) = \gamma_1 e^{-bt} + \gamma_0$ , where  $\gamma_1$  and  $\gamma_0$  are arbitrary constants. Therefore  $\xi = \frac{b}{2}x + \gamma_1 e^{-bt} + \gamma_0$ . Substitute it into the first equation of (18) to obtain

$$2\gamma_1(h_x + 2b^3)e^{-bt} + bxh_x + 2\gamma_0h_x - 2bh = 0.$$

Since  $h, \gamma_1, \gamma_0$  are independent of t, the preceding equation suggests that

$$2\gamma_1(h_x + 2b^3) = 0$$
,  $bxh_x + 2\gamma_0h_x - 2bh = 0$ ,

which leads to b = 0. Therefore, we have

$$\xi = d_1 t + d_0, \quad \eta = 2d_0, \quad f = 1, \quad h = h_0 \mod G^{\sim},$$

where  $d_1, d_0, h_0$  are constants. Thus, the corresponding regular reduction operator has the form

$$Q = \partial_t + (d_1t + d_0)\partial_x + 2d_1u\partial_u,$$

which is equivalent to Lie symmetry operator.

Case 2.1d: If system (25) is satisfied, then the expression (19) can be rewritten as

$$\xi = b(t)(x+q), \quad f(x) = \frac{1}{x+q} \mod G_1^{\sim}.$$
 (30)

Substitute it into the second equation of (18) to obtain 2b(t)b'(t) + b''(t) = 0, which is equivalent to the fourth equation of (25), and which leads to  $2b'^2(t) + b'''(t) = -2b(t)b''(t)$ . Substitute (30) into the first equation of (18) to obtain

$$h_x = 2f^2[2b'^2(t) + b'''(t)]/b(t) = 2f^2[-2b(t)b''(t)]/b(t) = -4f^2b''(t).$$

Since h and f are independent of t, there is a constant r such that b''(t) = r. It follows that there exist constants s and w such that  $b(t) = rt^2/2 + st + w$ . Substitute it into the fourth equation of (25) to obtain

$$r^{2}t^{3} + 3rst^{2} + 2(wr + s^{2})t + 2ws + r = 0.$$

Then r=0 and s=0. Hence  $b(t)=w,\,\xi_t=0$ , the solution is included in the case 2.1a.

Case 2.2: When m=1, the third equation of (17) implies

$$\xi h \frac{f_x}{f} - \xi h_x + f(2\xi_x + \xi \frac{f_x}{f})_{tt} + 2f(2\xi_x + \xi \frac{f_x}{f})(2\xi_x + \xi \frac{f_x}{f})_t 
-2f\xi_t(2\xi_x + \xi \frac{f_x}{f})_x - 2(f\xi_t)_{xx} = 0, 
2f\xi_t\xi_{tx} + f\xi_{ttt} - h\xi_t = 0, 
(2\xi_x + \xi \frac{f_x}{f})_{xx} = 0.$$
(31)

Similar to the case of m = 0, from the third equation of (31) and the first two equations of (17) we get the expression of  $\xi$  and f(x) as stated in (19), where a(t), b(t), c(t), d(t) satisfy the condition (22) or (23) or (24) or (25).

Case 2.2a: If system (22) is satisfied, then  $\xi_t = 0$ , the second equation of (31) is an identity. The expression (19) can be rewritten as (26). The first equation of (31) is reduced to  $h_x/h = f_x/f$ , which leads to  $h(x) = \epsilon f(x)$  ( $\epsilon = \pm 1$ ) mod  $G_1^{\sim}$ . In addition,  $\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u = (ax + b)u$ . Thus, we have

$$\xi = 2ax^{2} + 4bx - cx + d, 
\eta = (ax + b)u, 
f(x) = \exp\left(\int \frac{-7ax - 7b + 2c}{2ax^{2} + 4bx - cx + d} dx\right), 
h(x) = \epsilon f(x).$$
(32)

where a, b, c, d are arbitrary constants and  $\epsilon = \pm 1$ . Thus, the corresponding regular reduction operator has the form

$$Q = \partial_t + (2ax^2 + 4bx - cx + d)\partial_x + (ax + b)u\partial_u,$$

which is equivalent to Lie symmetry operator.

Case 2.2b: If system (23) is satisfied, then the expression (19) can be rewritten as (28). Hence,  $\xi_x = 0$ , k = b(t). If b'(t) = 0, then a(t), b(t), c(t), d(t) satisfy system (22), and the solution is included in the case 2.2a. We suppose that  $b'(t) \neq 0$ . From the second equation of (31) we see that  $h = f\xi_{ttt}/\xi_t$ . Substitute it into the first equation of (31) to get  $fk_{tt} + 2fkk_t - 2f_{xx}\xi_t = 0$ . Further it can be reduced to qb''(t) + 2qb(t)b'(t) - 2b'(t) = 0. Combine it with the fourth equation of (23) to get b(t) = -3/q which is contradict to the hypothesis  $b'(t) \neq 0$ .

Case 2.2c: If system (24) is satisfied, then the expression (19) can be rewritten as (29). If b'(t) = d'(t) = 0, then a(t), b(t), c(t), d(t) satisfy both systems (22) and (24), and the solution is included in the case 2.2a. We suppose that  $b'^2(t) + d'^2(t) \neq 0$ . Substitute (29) into the first equation of (31) to obtain

$$h_x = \frac{2[b''(t) + 2b(t)b''(t)]}{b(t)x + 2d(t)}$$
(33)

Substitute (29) into the second equation of (31) to obtain

$$h(x) = b'(t) + \frac{b'''(t)x + 2d'''(t)}{b'(t)x + 2d'(t)}$$
(34)

Then

$$h_x = \frac{2[b'''(t)d'(t) - b'(t)d'''(t)]}{[b'(t)x + 2d'(t)]^2}.$$

Substituting it into (33) yields

$$\frac{b'''(t)d'(t) - b'(t)d'''(t)}{\left[b'(t)x + 2d'(t)\right]^2} = \frac{b''(t) + 2b(t)b'(t)}{b(t)x + 2d(t)}.$$

Compare the coefficient of  $x^2$  to obtain  $b'^2(t)[b''(t) + 2b(t)b'(t)] = 0$ . Substitute the third equation of (24) into it to obtain  $b(t)b'^3(t) = 0$ , hence b'(t) = 0. Thus the fourth equation of (24) can be reduced to bd'(t) + d''(t) = 0. Solving this linear ordinary differential equation gives  $d(t) = \gamma_1 e^{-bt} + \gamma_0$ , where  $\gamma_1$  and  $\gamma_0$  are two arbitrary constants. Therefore the expressions (29) and (34) can be rewritten as

$$\xi = \frac{1}{2}bx + \gamma_1 e^{-bt} + \gamma_0, \quad f(x) = 1, \quad h(x) = b^2 \mod G_1^{\sim}.$$

System (31) is verified to be true. In addition,  $\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u = bu - 2\gamma_1 be^{-bt}$ . Therefore, we have

$$\xi = \frac{1}{2}bx + \gamma_1 e^{-bt} + \gamma_0, \quad \eta = bu - 2\gamma_1 b e^{-bt},$$
  

$$f(x) = 1, \quad h(x) = b^2,$$
(35)

where  $b, \gamma_1, \gamma_0$  are arbitrary constants. Thus, we have a nontrivial regular reduction operator

$$Q = \partial_t + (\frac{1}{2}bx + \gamma_1 e^{-bt} + \gamma_0)\partial_x + (bu - 2\gamma_1 b e^{-bt})\partial_u.$$
(36)

Case 2.2d: If system (25) is satisfied, then the expression (19) can be rewritten as (30). Substitute it into the first equation of (31) to obtain

$$b(t)(x+q)[h + (x+q)h_x] = [b''(t) + 2b(t)b'(t)].$$

Substitute the fourth equation of (25) into it to get  $b(t)(x+q)[h+(x+q)h_x]=0$ . It follows that h(x)=r/(x+q), where r is a nonzero constant. Substitute it and (30) into the second equation of (31) to obtain  $2b'^2(t)+b'''(t)-rb'(t)=0$ . From the fourth equation of (25), we find  $b'''(t)=4b^2(t)b'(t)-2b'^2(t)$ . Substitute it into the preceding equation to get  $b'(t)[4b^2(t)-r]=0$ , which leads to b'(t)=0. Then a(t),b(t),c(t),d(t) satisfy system (22), and the solution is included in the case 2.2a.

Case 2.3: When m=2, system (17) implies

$$(7\xi_{x} + 4\xi \frac{f_{x}}{f})_{x} = 0,$$

$$2(\xi_{x} + 2\xi \frac{f_{x}}{f})\xi_{t} + 4\xi_{t} \frac{f_{x}}{f} + 4\xi\xi_{tx} + 4\xi_{tx} + \xi_{tt} = 0,$$

$$2\xi_{t}\xi_{tx} + \xi_{ttt} = 0,$$

$$2f(2\xi_{x} + \xi \frac{f_{x}}{f})(2\xi_{x} + \xi \frac{f_{x}}{f})_{t} + f(2\xi_{x} + \xi \frac{f_{x}}{f})_{tt} - 4hf\xi_{t} - 2f\xi_{t}(2\xi_{x} + \xi \frac{f_{x}}{f})_{x} - 2(f\xi_{t})_{xx} = 0,$$

$$\xi h_{x} + 2h\xi_{x} + (2\xi_{x} + \xi \frac{f_{x}}{f})_{xx} = 0.$$
(37)

From the first and the last equation of system (37), we can get

$$f(x) = |\xi|^{-7/4} \exp(\alpha(t) \int \frac{\mathrm{d}x}{\xi}), \quad h(x) = \frac{\xi_x^2 - 2\xi \xi_{xx} + q}{8\xi^2},$$

where  $\alpha(t)$  is an arbitrary function, q is a constant. Substituting these expressions into the rest equations of system (37), we can see that  $\xi(t,x)$  and  $\alpha(t)$  satisfy the overdetermined system of partial differential equations

$$2\xi_{t}\xi_{tx} + \xi_{ttt} = 0,$$

$$\xi_{tt} - 3\xi_{tx} - 5\xi_{t}\xi_{x} + 4\xi\xi_{tx} + 4\alpha\xi_{t} + 4\alpha_{t} = 0,$$

$$2\xi^{2}(\frac{1}{4}\xi_{x} + \alpha)_{tt} + 2\xi^{2}[(\frac{1}{4}\xi_{x} + \alpha)^{2}]_{t} - \xi^{2}\xi_{t}\xi_{xx} - \xi_{t}(\xi_{x}^{2} - 2\xi\xi_{xx} + q)$$

$$-4[(\alpha_{t} - \frac{3}{4}\xi_{tx})(\alpha - \frac{7}{4}\xi_{x})\xi - \frac{3}{4}\xi^{2}\xi_{txx}] = 0.$$
(38)

In addition, we have

$$\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u = 2\xi_t |\xi|^{-7/4} \exp(\alpha(t) \int \frac{\mathrm{d}x}{\xi}) + \left[\frac{1}{4}\xi_x + \alpha(t)\right]u.$$

Thus, we have a nontrivial regular reduction operator

$$Q = \partial_t + \xi(t, x)\partial_x + \{2\xi_t |\xi|^{-7/4} \exp(\alpha(t) \int \frac{\mathrm{d}x}{\xi}) + \left[\frac{1}{4}\xi_x + \alpha(t)\right]u\}\partial_u, \tag{39}$$

where  $\xi(t, x)$  and  $\alpha(t)$  satisfy the overdetermined system of partial differential equations (38). In particular, if  $\xi_t = 0$ , from system (37) we can obtain

$$\xi = \xi(x), \quad \eta = \frac{1}{4}(\xi_x + a)u, f(x) = |\xi|^{-7/4} \exp(\frac{a}{4} \int \frac{\mathrm{d}x}{\xi}), \quad h(x) = \frac{\xi_x^2 - 2\xi\xi_{xx} + q}{8\xi^2}.$$
 (40)

where a, q are arbitrary constants. Thus, we have a nontrivial regular reduction operator

$$Q = \partial_t + \xi(x)\partial_x + (\frac{1}{4}\xi_x + a)u\partial_u, \tag{41}$$

which is equivalent to operator (16) with n = 1. Therefore, this special case can be included in case 1.2 and we can impose an additional constraint  $\xi_t \neq 0$  on the regular reduction operator (39).

Case 2.4: When m=3, the third equation of (17) implies

$$2\xi_{t}\xi_{tx} + \xi_{ttt} = 0,$$

$$2f(2\xi_{x} + \xi\frac{f_{x}}{f})(2\xi_{x} + \xi\frac{f_{x}}{f})_{t} + f(2\xi_{x} + \xi\frac{f_{x}}{f})_{tt} - 2f\xi_{t}(2\xi_{x} + \xi\frac{f_{x}}{f})_{x} - 2(f\xi_{t})_{xx} = 0,$$

$$6hf\xi_{t} + (2\xi_{x} + \xi\frac{f_{x}}{f})_{xx} = 0,$$

$$\xi h\frac{f_{x}}{f} - \xi h_{x} - 2h(2\xi_{x} + \xi\frac{f_{x}}{f}) = 0,$$

$$(42)$$

the fourth equation of which can be rewritten as

$$\frac{\xi_x}{\xi} = -\frac{1}{4} \left( \frac{f_x}{f} + \frac{h_x}{h} \right).$$

Since f and h are independent of t, integrate both sides of the preceding equation to obtain  $\xi = r(t)|fh|^{-1/4}$ , where r(t) is a function of t. Substituting it into the first equation of (42) yields the fact that  $r'''(t) = q(x)r'^2(t)$ , where  $q(x) = -2(|fh|^{-1/4})_x$ . It follows that r'(t) = 0 or q'(x) = 0,  $r'''(t) = qr'^2(t)$ .

Case 2.4a: If r'(t) = 0, then  $\xi = r|fh|^{-1/4}$  (r = const),  $\xi_t = 0$ ,  $g_t = 0$ , the second equation of (42) is an identity, so is the second equation of (17). The third equation of (42) reduces to  $\left(2\xi_x + \xi \frac{f_x}{f}\right)_{xx} = 0$ . Combine it with the first equation of (17), and use a progress similar to the case m = 0 (i.e. 2.1a), we get the expression of  $\xi$  and f(x) as stated in (26), where a, b, c, d are constants. From  $\xi = r|fh|^{-1/4}$ , we see that  $h(x) = \pm \frac{1}{f} \left(\frac{r}{\xi}\right)^4$ , where  $r\xi > 0$ . In addition,  $\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u = (ax + b)u$ . Thus, we have

$$\xi = 2ax^{2} + 4bx - cx + d,$$

$$\eta = (ax + b)u,$$

$$f(x) = \exp\left(\int \frac{-7ax - 7b + 2c}{2ax^{2} + 4bx - cx + d} dx\right),$$

$$h(x) = \pm \frac{1}{f} \left(\frac{r}{\xi}\right)^{4}.$$
(43)

where a,b,c,d,r are arbitrary constants. Thus, the corresponding regular reduction operator has the form

$$Q = \partial_t + (2ax^2 + 4bx - cx + d)\partial_x + (ax + b)u\partial_u,$$

which is equivalent to Lie symmetry operator.

Case 2.4b: If q'(x) = 0,  $r'''(t) = qr'^2(t)$ , then  $(|fh|^{-1/4})_x = -\frac{1}{2}q$ . Integration both sides of it gives  $|fh|^{-1/4} = -\frac{1}{2}qx + s$ , where s is a constant. Therefore  $\xi = r(t)|fh|^{-1/4} = r(t)(-\frac{1}{2}qx + s)$ . From the first equation of (17) we see that  $(2\xi_x + \xi \frac{f_x}{f})_x = \frac{1}{4}\xi_{xx} + \frac{1}{4}(7\xi_x + 4\xi \frac{f_x}{f})_x = \frac{1}{4}\xi_{xx}$ . Substituting the last two expressions into the third equation of (42), yields r'(t) = 0 or  $\xi = 0$ , which are the cases have already been discussed.

Case 2.5: When  $m \neq 0, 1, 2, 3$ , the third equation of (17) implies

$$\xi h \frac{f_x}{f} - \xi h_x + (1 - m) h (2\xi_x + \xi \frac{f_x}{f}) = 0,$$
  

$$\xi_t = 0,$$
  

$$(2\xi_x + \xi \frac{f_x}{f})_{xx} = 0.$$
(44)

Notice that the second equation of (44) indicates that  $\xi$  is independent of t, therefore the second equation of (17) is satisfied automatically. Similar to the case of m = 0, from the third equation

of (44) and the first equation of (17) we get the expression of  $\xi$  and f(x) as stated in (26), where a, b, c, d are constants. Substituting the expression of  $\xi$  and f into the first equation of (44) we obtain

$$h(x) = \exp\Big(\int \frac{-7ax - 7b + 2c + (1-m)(ax+b)}{2ax^2 + 4bx - cx + d} dx\Big).$$

In addition,  $\eta = 2f\xi_t + (2\xi_x + \xi \frac{f_x}{f})u = (ax + b)u$ . Therefore, we have

$$\xi = 2ax^{2} + 4bx - cx + d,$$

$$\eta = (ax + b)u,$$

$$f(x) = \exp\left(\int \frac{-7ax - 7b + 2c}{2ax^{2} + 4bx - cx + d} dx\right),$$

$$h(x) = \exp\left(\int \frac{-7ax - 7b + 2c + (1 - m)(ax + b)}{2ax^{2} + 4bx - cx + d} dx\right).$$
(45)

where a, b, c, d are arbitrary constants. Thus, the corresponding regular reduction operator has the form

$$Q = \partial_t + (2ax^2 + 4bx - cx + d)\partial_x + (ax + b)u\partial_u,$$

which is equivalent to Lie symmetry operator.

From the above discussion, we can arrival at the following two theorems.

**Theorem 3.** A complete list of  $G_1^{\sim}$ -inequivalent equations (3) having nontrivial regular reduction operator is exhausted by ones given in table 1.

 $\begin{array}{|c|c|c|c|c|c|}\hline N & n & m & f(x) & h(x) & \text{Regular reduction operator }Q\\ \hline 1 & \neq -1 & n+1 & |\xi|^{-\frac{3n+4}{2n+2}}\exp\left(r\int\frac{1}{\xi}\,\mathrm{d}x\right) & \frac{\xi_x^2-2\xi\xi_{xx}-p}{4(n+1)\xi^2} & \partial_t+\xi(x)\partial_x+\left(\frac{r}{n}+\frac{\xi_x}{2n+2}\right)u\partial_u\\ 2 & 1 & 1 & 1 & b^2 & \partial_t+\left(\frac{1}{2}bx+\gamma_1e^{-bt}+\gamma_0\right)\partial_x+\left(bu-2\gamma_1be^{-bt}\right)\partial_u\\ 3 & 1 & 2 & |\xi|^{-\frac{7}{4}}\exp(\alpha(t)\int\frac{\mathrm{d}x}{\xi}) & \frac{\xi_x^2-2\xi\xi_{xx}+q}{8\xi^2} & \partial_t+\xi(t,x)\partial_x+\left\{2\xi_t|\xi|^{-\frac{7}{4}}\exp(\alpha(t)\int\frac{\mathrm{d}x}{\xi})\\ & +\left[\frac{1}{4}\xi_x+\alpha(t)\right]u\right\}\partial_u \\ \end{array}$ 

Table 1. Results of regular reduction operator classification of class (3)

Here  $r, p, b, \gamma_1, \gamma_0$  are arbitrary constants,  $\xi(x)$  in case 2.1 is an arbitrary functions of the variables  $x, \xi(t, x)$  and  $\alpha(t)$  in case 2.3 satisfy the overdetermined system of partial differential equations (38) and  $\xi_t \neq 0$ .

**Theorem 4.** Any reduction operator of an equations from class (3) having the form (10) with  $\xi_t = 0$ ,  $\xi_{xxx} = 0$  is equivalent to a Lie symmetry operator of this equation.

#### 3 Exact solutions

In this section, we construct nonclassical reduction and exact solutions for the classification models in table 1 by using the corresponding regular reduction operator. Lie reduction and exact solutions of equation from class (3) have been investigated in reference [21]. We choose case 1 in table 1 as an example to implement the reduction, the other cases can be considered in a similar way.

For the first case in table 1, the corresponding equation is

$$\left[\left|\xi\right|^{-\frac{3n+4}{2n+2}}\exp\left(r\int\frac{1}{\xi}\,\mathrm{d}x\right)\right]u_{tt} - (u^n u_x)_x - \frac{\xi_x^2 - 2\xi\xi_{xx} - p}{4(n+1)\xi^2}u^{n+1} = 0,\tag{46}$$

which admit the regular reduction operator

$$Q = \partial_t + \xi(x)\partial_x + (\frac{r}{n} + \frac{\xi_x}{2n+2})u\partial_u.$$

An ansätze constructed by this operator has the form

$$u(t,x) = \varphi(\omega) |\xi|^{\frac{1}{2n+2}} \exp\left(\frac{r}{n} \int \frac{1}{\xi} dx\right), \text{ where } \omega = t - \int \frac{1}{\xi} dx.$$

Substituting this ansätze into equation (46) leads to the reduced ODE

$$[(4r^{2} - p)n^{2} + 4(2n + 1)r^{2}]\varphi^{n+1}(\omega) + 4n(n + 1)[n\varphi''(\omega) - 2(n + 1)r\varphi'(\omega)]\varphi^{n}(\omega) + 4n^{3}(n + 1)\varphi'^{2}(\omega)\varphi^{n-1}(\omega) - 4n^{2}(n + 1)\varphi''(\omega) = 0.$$

$$(47)$$

Because there are higher nonlinear terms, we were not able to completely solve the above equation. Thus, we try to solve this equation under different additional constraints imposed on p and r.

We first rewrite equation (47) as

$$4n^{2}(n+1)[\varphi'(\omega)\varphi^{n}(\omega)]' - 4n^{2}(n+1)\varphi''(\omega) - 8n(n+1)^{2}r\varphi'(\omega)\varphi^{n}(\omega) + [4(n+1)^{2}r^{2} - pn^{2}]\varphi^{n+1}(\omega) = 0.$$
(48)

If we take  $p = 4(1 + \frac{1}{n})^2 r^2$ , then the general solution of (48) can be written in the implicit form

$$\int \frac{n(\varphi^n - 1)}{2r\varphi^{n+1} + c_1} d\varphi = \omega + c_2. \tag{49}$$

Up to similarity of solutions of equation (3), the constant  $c_2$  is inessential and can be set to equal zero by a translation of  $\omega$ , which is always induced by a translation of t.

If we further set n=1, the general solution (49) can be rewritten in the implicit form

$$\omega - \frac{\ln(2r\varphi^2 - c_1)}{4r} - \frac{\sqrt{2}\operatorname{arctanh}(\frac{\sqrt{2}r\varphi}{\sqrt{c_1r}})}{2\sqrt{c_1r}} + c_2 = 0.$$
(50)

Thus we obtain the following solution

$$u(t,x) = \varphi(\omega) |\xi|^{\frac{1}{4}} \exp\left(r \int \frac{1}{\xi} dx\right), \qquad \omega = t - \int \frac{1}{\xi} dx$$

for the equation

$$\left[|\xi|^{-\frac{7}{4}} \exp\left(r \int \frac{1}{\xi} \, \mathrm{d}x\right)\right] u_{tt} - (uu_x)_x - \frac{\xi_x^2 - 2\xi \xi_{xx} - 8r^2}{8\xi^2} u^2 = 0,$$

where  $\varphi$  satisfy the equation (50),  $\xi$  is an arbitrary function and r is a non-zero constant. If we further set r = 0, the general solution (49) can be rewritten in the implicit form

$$\frac{1}{n+1}\varphi^{n+1}(\omega) - \varphi(\omega) = c_1\omega + c_2. \tag{51}$$

Thus we obtain the following solution

$$u(t,x) = \varphi(\omega) |\xi|^{\frac{1}{2n+2}}, \qquad \omega = t - \int \frac{1}{\xi} dx$$

for the equation

$$|\xi|^{-\frac{3n+4}{2n+2}}u_{tt} - (u^n u_x)_x - \frac{\xi_x^2 - 2\xi \xi_{xx}}{4(n+1)\xi^2}u^{n+1} = 0,$$

where  $\varphi$  satisfy the equation (51),  $\xi$  is an arbitrary function. In particular, for n=1 from equation (51) we have

$$\varphi(\omega) = 1 \pm \sqrt{1 + 2(c_1\omega + c_2)}.$$

Thus we obtain an explicit solution

$$u(t,x) = [1 \pm \sqrt{1 + 2(c_1\omega + c_2)}]|\xi|^{\frac{1}{4}}, \qquad \omega = t - \int \frac{1}{\xi} dx$$

for the equation

$$|\xi|^{-\frac{7}{4}}u_{tt} - (uu_x)_x - \frac{\xi_x^2 - 2\xi\xi_{xx}}{8\xi^2}u^2 = 0.$$

If we take different functions for  $\xi$ , then we can obtain a series of solutions for the corresponding equations. In order to avoid tediousness, we do not make a further discussion here.

### 4 Conclusion and Discussion

In this paper we have given a detailed investigation of the reduction operators of the variable coefficient nonlinear wave equations (1) (equivalently to (3)) by using the singular reduction operator theory. A classification of regular reduction operators is performed with respect to generalized extended equivalence groups. The main results on classification for the equation (3) are collected in table 1 where we list three inequivalent cases with the corresponding regular reduction operators. Nonclassical symmetry reduction of a class nonlinear wave model (46) which are singled out the classification models are also performed. This enabled to obtain some non-Lie exact solutions which are invariant under certain conditional symmetries for the corresponding model.

The present paper is a preliminary nonclassical symmetry analysis of the class of hyperbolic type nonlinear partial differential equations (1). Therefore, further investigations of different properties such as nonclassical potential symmetries, and nonclassical potential exact solutions as well as physical application of this class of equations would be extremely interesting. These results will be reported in subsequent publications.

#### Acknowledgements

This work was partially supported by the National Key Basic Research Project of China under Grant No.2010CB126600, the China Postdoctoral Science special Foundation under Grant No.201104247.

### References

- [1] W.F. Ames, Nonlinear partial differential equations in engineering Academic, New York, 1972, p.50-52.
- [2] W.F. Ames, E. Adams and R.J. Lohner, Group properties of  $u_{tt} = [f(u)u_x]_x$ , Int. J. Non-Linear Mech. 16 (1981) 439–447.
- [3] G.W. Bluman, S. Kumei, Symmetries and Differential Equations, Springer, Berlin, 1989.

- [4] N.H. Ibragimov (Editor), Lie group analysis of differential equations symmetries, exact solutions and conservation laws, V.1, CRC Press, Boca Raton, FL, 1994.
- [5] D.J. Arrigo, Group properties of  $u_{xx} u_y^m u_{yy} = f(u)$ , Int. J. Non-Linear Mech., 26 (1991) 619–629.
- [6] G.W. Bluman and A.F. Cheviakov, Nonlocally related systems, linearization and nonlocal symmetries for the nonlinear wave equation, J. Math. Anal. Appl., 333 (2007) 93-111.
- [7] G.W. Bluman, Temuerchaolu and R. Sahadevan, Local and nonlocal symmetries for nonlinear telegraph equation, J. Math. Phys., 46 (2005) 023505.
- [8] S.C. Chikwendu, Non-linear wave propagation solutions by Fourier transform perturbation, Int. J. Non-Linear Mech., 16 (1981) 117–128.
- [9] A. Donato, Similarity analysis and nonlinear wave propagation, Int. J. Non-Linear Mech. 22 (1987) 307–314.
- [10] M.L. Gandarias, M. Torrisi and A. Valenti, Symmetry classification and optimal systems of a non-linear wave equation, Int. J. Non-Linear Mech., 39 (2004) 389–398.
- [11] D.J. Huang and N.M. Ivanova, Group analysis and exact solutions of a class of variable coefficient nonlinear telegraph equations, J. Math. Phys, 48 (2007) 073507. (23 pages)
- [12] D.J. Huang, J.Q. Mei, H.Q. Zhang, Group classification and exact solutions of a class of variable coefficient nonlinear wave equations, Chin. Phys. Lett, 26 (2009) 050202.
- [13] D.J. Huang, S.G. Zhou, Group properties of generalized quasi-linear wave equations, J. Math. Anal. Appl., 366 (2010) 460-472.
- [14] D.J. Huang, S.G. Zhou, Group-theoretical analysis of variable coefficient nonlinear telegraph equations, Acta Appl. Math., 117 (2012) 135-183. (arXiv:[math-ph]1101.4755)
- [15] N.H. Ibragimov, M. Torrisi and A. Valenti, Preliminary group classification of equations  $v_{tt} = f(x, v_x)v_{xx} + g(x, v_x)$ , J. Math. Phys., 32 (1991) 2988–2995.
- [16] A. Oron and P. Rosenau, Some symmetries of the nonlinear heat and wave equations, Phys. Lett. A, 118 (1986) 172–176.
- [17] E. Pucci, Group analysis of the equation  $u_{tt} + \lambda u_{xx} = g(u, u_x)$ , Riv. Mat. Univ. Parma, 12 N 4 (1987) 71–87.
- [18] E. Pucci and M.C. Salvatori, Group properties of a class of semilinear hyperbolic equations, Int. J. Non-Linear Mech., 21 (1986) 147–155.
- [19] M. Torrisi and A. Valenti, Group properties and invariant solutions for infinitesimal transformations of a nonlinear wave equation, Int. J. Non-Linear Mech., 20 (1985) 135–144.
- [20] O.F. Vasilenko and I.A. Yehorchenko, Group classification of multidimensional nonlinear wave equations, Proceedings of Institute of Mathematics of NAS of Ukraine, 36 (2001) 63–66.
- [21] D.J. Huang, Q.M Yang, S.G. Zhou, Lie symmetry classification and equivalence transformation of variable coefficient nonlinear wave equations with power nonlinearities, Chinese Journal of Contemporary Mathematics, 33(3) (2012) 205–214.
- [22] D.J. Huang, Q.M Yang, S.G. Zhou, Conservation law classification of variable coefficient nonlinear wave equation with power Nonlinearity, Chin. Phys. B, 20 (2011) 070202.
- [23] P.J. Olver, Application of Lie Groups to Differential Equations, Springer, New York, 1986.
- [24] L.V. Ovsiannikov, Group analysis of differential equations, Academic Press, New York, 1982.
- [25] G.W. Bluman, J.D. Cole, The general similarity solution of the heat equation, J. Math. Mech., 18 (1969) 1025-1042.
- [26] D. Levi and P. Winternitz, Non-classical symmetry reduction: example of the Boussinesq equation J. Phys. A: Math. Gen. 22 (1989), 2915-2924.
- [27] Fushchych W.I., Shtelen W.M. and Serov N.I., Symmetry Analysis and Exact Solutions of Nonlinear Equations of Mathematical Physics, Dordrecht: Kluwer, (English transl.) 1993.

- [28] Fushchych W. I., Shtelen W.M., Serov M.I. and Popovych R.O., Q-conditional symmetry of the linear heat equation, *Proc. Acad. Sci. Ukraine*, 1992, no. 12, 28C33.
- [29] R.Z. Zhdanov, I.M. Tsyfra, R.O. Popovych, A precise definition of reduction of partial differential equations, J. Math. Anal. Appl., 238 (1999) 101-123. (arXiv:math-ph/0207023)
- [30] W.I. Fushchych and R.Z. Zhdanov, Conditional symmetry and reduction of partial differential equations, Ukr. Math. J., 44 (1992) 970C982.
- [31] M. Kunzinger and R.O. Popovych, Is a nonclassical symmetry a symmetry, Proceedings of 4th Workshop Group Analysis of Differential Equations and Integrability, 2009. (math-ph/0903.0821)
- [32] P.J. Olver and P. Rosenau, Group-invariant solutions of differential equations, SIAM J. Appl. Math. 47 (1987)263-278.
- [33] P.J. Olver, Direct reduction and differential constraints, Proc. R. Soc. Lond. A, 444(1994)509-523.
- [34] P.J. Olver and P. Rosenau, The construction of special solutions to partial differential equations, Phys. Lett. A 114(1986)107-112.
- [35] E. Pucci and G. Saccomandi, On the weak symmetry groups of partial differential equations, J. Math. Anal. Appl. 163(1992)588-598.
- [36] A.F. Sidorov, V.P. Shapeev and N.N. Yanenko, The method of differential constraints and its applications in gas dynamics, Nauka Sibirsk. Otdel., Novosibirsk, 1984. (in Russian).
- [37] N.N. Yanenko, Compatibility theory and methods of integrating systems of nonlinear partial differential equations, in Proc. Fourth All-Union Mathematics Congress, Nauka, Leningrad, 1964, pp. 247-259.
- [38] V.M. Boyko, M. Kunzinger and R.O. Popovych, Singular reduction modules of differential equations, arXiv:1201.3223, 30 pp.
- [39] M. Kunzinger and R.O. Popovych, Singular reduction operators in two dimensions, J. Phys. A, 41 (2008) 505201. (24 pp., [arXiv:0808.3577]).
- [40] O.A. Pocheketa and R.O. Popovych, Reduction operators and exact solutions of generalized Burgers equations, Phys. Lett. A 376 (2012)2847-2850. (arXiv:1112.6394)
- [41] O.A. Pocheketa and R.O. Popovych, Reduction operators of Burgers equation, J. Math. Anal. Appl. 398 (2013)270-277. (arXiv:1208.0232)
- [42] V.M. Boyko and R.O. Popovych, Reduction operators of the linear rod equation, Proceedings of the Sixth International Workshop Group Analysis of Differential Equations and Integrable Systems" (Protaras, Cyprus, June 17-21, 2012), University of Cyprus, Nicosia, 2013, 17-29.
- [43] R.O. Popovych, Reduction operators of linear second-order parabolic equations, J. Phys. A, 41 (2008) 185202. (31 pp., [arXiv:0712.2764])
- [44] R.O. Popovych, O.O Vaneeva and N.M. Ivanova, Potential nonclassical symmetries and solutions of fast diffusion equation, Phys. Lett. A, 362 (2007) 166-173. (arXiv:math-ph/0506067)
- [45] O.O. Vaneeva, R.O. Popovych and C. Sophocleous, Reduction operators of variable coefficient semilinear diffusion equations with a power source, Proceedings of 4th Workshop "Group Analysis of Differential Equations and Integrable Systems", (26-30 October 2008, Protaras, Cyprus), 2009, 191-209. (arXiv:0904.3424)
- [46] O.O. Vaneeva, R.O. Popovych and C. Sophocleous, Reduction operators of variable coefficient semilinear diffusion equations with an exponential source, Proceedings of 5th Workshop "Group Analysis of Differential Equations and Integrable Systems", (June 6-10, 2010, Protaras, Cyprus), 2011, 207-219. (arXiv:1010.2046)