

Multiple-electron losses of highly charged ions colliding with neutral atoms

M S Litsarev¹, V P Shevelko²

¹Department of Physics and Astronomy, Uppsala University, Box 516, 75120 Uppsala, Sweden

²P.N. Lebedev Physical Institute, Leninskii prospect 53, 119991 Moscow, Russia

E-mail: ¹mikhail.litsarev@physics.uu.se, ²shev@sci.lebedev.ru

Abstract. We present calculations of the total and m -fold electron-loss cross sections using the DEPOSIT code for highly charged U^{q+} ions ($q = 10, 31, 33$) colliding with Ne and Ar targets at projectile energies $E = 1.4$ and 3.5 MeV/u. Typical examples of the deposited energy $T(b)$ and m -fold ionization probabilities $P_m(b)$ used for the cross-section calculations as a function of the impact parameter b are given. Calculated m -fold electron-loss cross sections are in a good agreement with available experimental data. Although the projectile charge is rather high, a contribution of multiple-electron loss cross sections to the total electron-loss cross sections is high: about 65 % for the cases mentioned.

PACS numbers: 34.50.Fa 34.50.Bw 29.27.Eg

Submitted to: *Phys. Scr.*

1. Introduction

Multi-electron losses (EL) of projectile positive and negative ions colliding with neutral atoms and molecules play a key role in many problems of plasma physics and physics of accelerators, especially at low and intermediate collision energies. A contribution of multi-electron processes to the total electron-loss cross sections can be substantial and reach up to more than 50 % even for highly charged projectile ions. The total and m -fold EL cross sections can be calculated in the classical approximation using the Classical Trajectory Monte Carlo approach [1, 2, 3] or the energy-deposition model [4, 5, 6, 7] in the impact parameter representation providing an overall agreement with experimental data within a factor of two.

In the present work the DEPOSIT code described in Ref. [8] is used to calculate the total and m -fold EL cross sections for U^{q+} ions, ($q = 10, 31, 33$) colliding with Ne and Ar targets at energies $E = 1.4$ and 3.5 MeV/u. The results are in a good agreement with available experimental data [9, 10].

2. Theoretical approach

The DEPOSIT code is based on the semi-classical energy deposition model introduced by N. Bohr [11] and developed further in Ref. [12] for ionization of *atoms* by ions. Here, this model is used for ionization of *ions* by neutral atoms at energies about a few MeV/u, where a contribution of multiple-electron loss processes to the total loss cross-section is very large

In the energy deposition model, the problem of determination the m -fold ionization probabilities $P_m(b)$ is reduced to calculation of a 3D integral over the ion-coordinate space using projectile electron density $\rho_\gamma(r)$ and the energy gain $\Delta E_\gamma(p)$ obtained in the field $U(R)$ of the target atom

$$T(b) = \sum_{\gamma} \int \rho_{\gamma}(r) \Delta E_{\gamma}(p) d^3 \mathbf{r}. \quad (1)$$

Here, index γ labels nl -shell of the ion, where n is the principal quantum number and l is the orbital quantum number. The summation is over all occupied shells of the projectile ion. The energy gain ΔE_γ is determined depending on the relation between the ion velocity v and orbital velocity $u_\gamma = \sqrt{2I_\gamma}$, I_γ is the binding energy of the projectile γ -shell.

Electron density $\rho_\gamma(r)$ in the present work is calculated using nodeless Slater wave functions, and atomic field at a distance R from its nucleus is given by sum of three Yukawa potentials with five fitting parameters defined from the Dirack-Hartree-Fock-Slater calculations [13].

The total EL cross-section is obtained directly from the equation

$$\sigma_{tot}(v) = \sum_{m=1}^N \sigma_m = \pi b_{\max}^2, \quad (2)$$

where parameter b_{\max} is (numerically) found from the equation

$$T(b_{\max}) = I_1, \quad (3)$$

using the first ionization potential I_1 of the projectile ion and expression (1) for the energy $T(b)$.

The m -fold cross-section is calculated in the following way

$$\sigma_m(v) = 2\pi \int_0^\infty P_m(b) b db, \quad (4)$$

where the m -fold ionization probabilities $P_m(b)$ are taken from the statistical Russek-Meli model [14]. In this model the $P_m(b)$ probabilities are presented analytically and can be found by using of the deposited energy $T(b)$ and the ionization potentials I_k of the projectile.

3. Results and discussions

The calculated deposited energies $T(b)$ and ionization probabilities $P_m(b)$ are presented for U^{31+} ions colliding with Ar target at the energy $E = 3.5$ MeV/u in Fig. 1 and Fig. 2 correspondingly. The calculated m -fold EL cross sections for uranium ions U^{q+} ($q = 10, 31, 33$) colliding with Ne and Ar atoms are given in Fig. 3 and Fig. 4 in comparison with experimental data.

Fig. 1 shows typical example of the $T(b)$ behaviour. The deposited energy for U^{31+} ion is transferred mainly to the $4d^{10}4f^{14}$ shell for all valid values of the impact parameter b . The value $b_{\max} = 0.76 a_0$ is obtained from the equation (3) for the first ionization potential $I_1 = 40.06$ a.u. of U^{31+} ion. For this system the inner-shell electrons does not contribute much to the ionization process.

Ionization probabilities calculated for $U^{31+} + Ar$ system colliding at the energy $E = 3.5$ MeV/u are presented in Fig 2. The Figure shows that for a given collision energy the probability of the 12-fold and more ionization processes is too small. This result is in correspondence with the experimental data [10] available only for number of ejected electrons $m \leq 11$.

In the Fig 3 and Fig 4 the calculated m -fold EL cross sections are shown in comparison with experiment. As one can see, the difference between the two data sets (experimental and theoretical) is small. The figures show that multi-electron losses contribute to the total EL loss significantly.

Finally, we present calculated total EL cross sections for all the considered systems. For $U^{31+} + Ar$ at $E = 3.5$ MeV/u $\sigma_{tot} = 5.18 \times 10^{-17} \text{ cm}^2$ (experimental value $\sigma_{tot}^{exp} = 3.48 \times 10^{-17} \text{ cm}^2$), for $U^{33+} + Ar$ at $E = 3.5$ MeV/u $\sigma_{tot} = 4.27 \times 10^{-17} \text{ cm}^2$ ($\sigma_{tot}^{exp} = 2.63 \times 10^{-17} \text{ cm}^2$), and for $U^{10+} + Ne$ at $E = 1.4$ MeV/u $\sigma_{tot} = 2.27 \times 10^{-16} \text{ cm}^2$ ($\sigma_{tot}^{exp} = 2.09 \times 10^{-16} \text{ cm}^2$). The corresponding experimental data are taken from the Refs. [9, 10], and are in a reasonable agreement with results of our calculations.

4. Conclusions

As seen from the experimental data for m -fold EL cross sections, a contribution of multi-electron losses ($m > 1$) to the total EL cross sections is very large and in the cases considered here reaches about 65 %. Calculations performed by the DEPOSIT code well reproduce this result. It means that multi-electron loss processes should be taken into account in investigations of kinetic of highly charged ions, first of all, in determination of the ion-charge fractions, mean charges of ions passing through the solid, gaseous or plasma targets, in estimations of the vacuum conditions in accelerator machines and many others.

Acknowledgments

The computations were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at Uppsala Multidisciplinary Center for Advanced Computational Science (UPPMAX) under the Project *s00311-8* and on resources provided by SNIC at the National Supercomputer Center (NSC) under the Project *matter2*.

This work was partly supported by the grant #11-02-00526a of the Russian Foundation for Basic Research.

References

- [1] Olson R E *et al* 2004 *J. Phys. B: At. Mol. Opt. Phys.* **37** 4539.
- [2] Olson R E 1989 *Phys. Rev. A* **39** 5572.
- [3] Olson R E 1996 *Atomic, Molecular, and Optical Physics Handbook* (College Park) chapter 56.
- [4] Shevelko V P *et al* 2008 *J. Phys. B: At. Mol. Opt. Phys.* **41** 115204.
- [5] Song M-Y *et al* 2009 *Nucl. Instrum. Methods B* **267** 2369.
- [6] Shevelko V P *et al* 2009 *J. Phys. B: At. Mol. Opt. Phys.* **42** 065202.
- [7] Shevelko V P *et al* 2010 *J. Phys. B: At. Mol. Opt. Phys.* **43** 215202.
- [8] Litsarev M S 2012 (submitted) *Comp. Phys. Com.*, *arXiv:1205.5390*.
- [9] DuBois R D *et al* 2004 *Phys. Rev. A* **70** 032712.
- [10] Perumal A N *et al* 2005 *Nucl. Instrum. Methods B* **227** 251.
- [11] Bohr N 1915 *Phil. Mag.* **30** 581.
- [12] Cocke C L 1979 *Phys. Rev. A* **20** 749.
- [13] Salvat F *et al* 1987 *Phys. Rev. A* **36** 467.
- [14] Russek A and Meli J 1970 *Physica A* **46** 222.
- [15] Rashid K *et al* 1988 *At. Data Nucl. Data Tables* **40** 365.




Figure 1. Calculated energy $T(b)$ deposited to uranium ions by Ar atoms as a function of the impact parameter b , the DEPOSIT code. A contribution of the deposited energy to different subshells of uranium ions are shown. The horizontal line corresponds to the first ionization potential of U^{31+} : $I_1 = 40$ a.u. (1090 eV). The notation for the Slater shells, for example, $2(sp)^8$ means the electron configuration of the $2s^22p^6$ shell.




Figure 2. Calculated probabilities $P_m(b)$ for the m -fold ionization of U^{31+} ions colliding with Ar atoms at energy $E = 1.4$ MeV/u as a function of impact parameter b . In every b -point the condition $\sum_m P_m(b) = 1$ is satisfied.




Figure 3. Multiple-electron loss in $U^{q+} + Ar$ collision at energy $E = 1.4$ MeV/u. The m -fold electron-loss cross sections as a function of ejected electrons m . Experiment – Ref. [10], theory – the DEPOSIT code. The binding energies for U^{q+} ions are taken from Ref. [15].




Figure 4. The m -fold electron-loss cross sections of U^{10+} ions colliding with Ne atoms at the collision energy $E = 1.4$ MeV/u. Experiment – Ref. [9], theory – the DEPOSIT code. The binding energies for U^{10+} ions are taken from Ref. [15].