Alfvénic tornadoes

P. K. Shukla

- ² International Centre for Advanced Studies in Physical Sciences & Institute
- $_{\scriptscriptstyle 3}\,$ for Theoretical Physics, Ruhr University Bochum, D-44780 Bochum,

Germany

- Department of Mechanical and Aerospace Engineering & Center for Energy
- Research, University of California San Diego, La Jolla, CA 92093, USA

P. K. Shukla, International Centre for Advanced Studies in Physical Sciences & Institute for Theoretical Physics, Ruhr University Bochum, 44780 Bochum, Germany (profshukla@yahoo.de)

SHUKLA: FULL-SCALE SIMULATION STUDY

It is shown that a three-dimensional (3D) modified-kinetic Alfvén waves 7 (m-KAWs) can propagate in the form of Alfvénic tornadoes characterized 8 by plasma density whirls or magnetic flux ropes carrying orbital angular mo-9 mentum (OAM). By using the two fluid model, together with Ampère's law, 10 we derive the wave equation for a 3D m-KAWs in a magnetoplasma with $m_e/m_i \ll$ 11 $\beta \ll 1$, where m_e (m_i) is the electron (ion) mass, $\beta = 4\pi n_0 k_B (T_e + T_i)/B_0^2$, 12 n_0 the unperturbed plasma number density, k_B the Boltzmann constant, $T_e(T_e)$ 13 the electron (ion) temperature, and B_0 the strength of the ambient magnetic 14 field. The 3D m-KAW equation admits solutions in the form of a Laguerre-15 Gauss (LG) Alfvénic vortex beam or Alfvénic tornadoes with plasma den-16

¹⁷ sity whirls that support the dynamics of Alfvén magnetic flux ropes.

1. Introduction

The discovery of the Alfvén wave Alfven [1942] had significantly changed the landscape 18 of collective phenomena in magnetized plasmas that frequently occur in interstellar media 19 Shukla and Dawson [1984] and in large laboratory plasma devices [Lennen et al., 1999; 20 Kletzing et al., 2010; Gekelman et al., 2011]. In the Alfvén wave, the restoring force 21 comes from the pressure of the magnetic fields, and the ion mass provides the inertia. 22 The dispersion relation for the low-frequency (in comparison with the ion gyrofrequency) 23 Alfvén wave is deduced from the magnetohydrodynamic (MHD) equations [Alfven, 1942; 24 Cramer, 2001, which do not account for the wave dispersion effects. The dispersion 25 [Stefant, 1970; Brodin and Stenflo, 1990; Morales et al., 1994; Morales and Maggs, 26 1997; Shukla and Stenflo, 1999; Dastgeer and Shukla, 2009; Damino et al., 2009] to 27 the Alfvén wave comes from the finite frequency (ω/ω_{ci}) and the magnetic field-aligned 28 electron inertial force in cold magnetoplasmas, as well as from the finite ion gyroradius 29 effect and the gradient of the electron pressure perturbation in a warm magnetoplasma, 30 Due to non-ideal effects Brodin and Stenflo [1990]; Morales et al. [1994]; Morales and 31 Maggs [1997]; Shukla and Stenflo [1999]; Damino et al. [2009], Alfvén waves thus 32 couple to fast and slow magnetoacoustic waves, the inertial and kinetic Alfvén waves 33 Stefant [1970]; Morales et al. [1994]; Morales and Maggs [1997]; Dastgeer and Shukla 34 [2009]; Damino *et al.* [2009] in a uniform magnetoplasma. The magnetic field-aligned 35 phase speed (ω/k_z) of the inertial (kinetic) Alfvén wave is much larger (smaller) than the 36 electron thermal speed. Accordingly, the inertial (kinetic) Alfvén wave arises in a plasma 37 with $\beta \ll m_e/m_i \ (m_e/m_i \ll \beta \ll 1)$. The dispersive shear Alfvén wave plays a significant 38

X - 4

role in the auroral Chaston *et al.* [2008a, b], magnetospheric Sundkvist *et al.* [2005],39 solar Tripathi and Gekeklman [2010]; Salem et al. [2012], astrophysical and laboratory [Gekelman et al., 2011, 2012] magnetoplasmas with regard to acceleration and heating of 41 the plasma particles Shukla et al. [1998], reconnection of magnetic field lines, wave-wave 42 and wave-particle interactions [Hasegawa and Chen, 1976; Carter et al., 2006; Shukla 43 et al., 2007], and the formation of nonlinear Alfvénic structures [Sundkvist et al., 2005; 44 Stasiewicz et al., 2003; Shukla et al., 2012]. Specifically, it is likely that the dispersive 45 Alfvén waves power the auroral activities Chaston *et al.* [2008a, b] and the solar coronal 46 heating [Tomczyk et al., 2007; Jess et al., 2009; McIntosh et al., 2011], as well as may be 47 responsible for the formation of Alfvénic tornadoes Widemeyer-Böhm et al. [2012] which 48 can transport magneto-convective energy from one region to another in solar plasmas. 49

In this Brief Report, it is shown that a 3D m-KAW can propagate as a twisted vortex beam in a plasma with $m_e/m_i \ll \beta \ll 1$. The present work thus complements a recent investigation Shukla [2012] that focused on examining a 3D twisted inertial Alfvén wave. Twisted m-KAWs can be related with magnetic flux ropes or Alfvénic tornadoes of different scalesizes in observational data from laboratory experiments Gekelman *et al.* [2012], and also from the solar plasmas Widemeyer-Böhm *et al.* [2012]; Salem *et al.* [2012].

2. Mathematical description

We consider a magnetized electron-ion plasma in the presence of low-frequency (in comparison with the electron gyrofrequency $\omega_{ce} = eB_0/m_ec$, where e is the magnitude of the electron charge, B_0 the strength of the external magnetic field $\hat{\mathbf{z}}B_0$, m_e the electron mass, and c the speed of light in vacuum, $\hat{\mathbf{z}}$ the unit vector along the z-axis in a Cartesian coor-

DRAFT

June 21, 2021, 10:07am

dinate system) m-KAWs with the electric and magnetic fields $\mathbf{E} = -\nabla \phi - (1/c) \hat{\mathbf{z}} \partial A_z / \partial t$ and $\mathbf{B}_{\perp} = \nabla A_z \times \hat{\mathbf{z}}$, respectively, where ϕ and A_z are the scalar and magnetic field-aligned vector potentials, respectively. In the m-KAW fields, the electron density perturbation is obtained from

$$\frac{\partial n_{e1}}{\partial t} + \frac{c}{4\pi e} \frac{\partial \nabla_{\perp}^2 A_z}{\partial z} = 0, \tag{1}$$

⁶⁴ where we have used the electron fluid velocity

$$\mathbf{u}_e \approx \frac{c}{B_0} \hat{\mathbf{z}} \times \nabla \phi - \frac{ck_B T_e}{B_0 n_0} \hat{\mathbf{z}} \times \nabla n_{e1} + \hat{\mathbf{z}} \frac{c}{4\pi e n_0} \nabla_{\perp}^2 A_z,$$
(2)

which uses Ampère's law that relates A_z and the magnetic field-aligned electron fluid velocity u_{ez} . Here $n_{e1} \ll n_0$ is a small electron density perturbation in the equilibrium density n_0 .

Since the parallel phase speed of the m-KAWs is much smaller than the electron thermal speed $V_{Te} = (k_B T_e/m_e)^{1/2}$, one can neglect the electron inertia and obtain from the magnetic field aligned electron momentum equation

$$0 = e\frac{\partial\phi}{\partial z} + \frac{e}{c}\frac{\partial A_z}{\partial t} - \frac{k_B T_e}{n_0}\frac{\partial n_{e1}}{\partial z},\tag{3}$$

which dictates that the parallel electric force $-n_0 eE_z$ and $\partial p_1/\partial z$ are in balance, where the magnetic field-aligned electric field $E_z = -\partial \phi - c^{-1} \partial A_z/\partial t$ and $p_1 = k_B T_e n_{e1}$ is the electron pressure perturbation.

⁷⁴ We can now eliminate A_z from Eqs. (1) by using (3), obtaining

$$\left(\frac{\partial^2}{\partial t^2} - c^2 \lambda_{De}^2 \frac{\partial^2 \nabla_{\perp}^2}{\partial z^2}\right) n_{e1} - \frac{c^2}{4\pi e} \frac{\partial^2 \nabla_{\perp}^2 \phi}{\partial z^2} = 0, \tag{4}$$

where $\lambda_{De} = (k_B T_e / 4\pi n_0 e^2)^{1/2}$ is the electron Debye radius.

The perpendicular (to $\hat{\mathbf{z}}$) component of the ion fluid velocity $\mathbf{u}_{i\perp}$ is determined from

$$\left(\frac{\partial^2}{\partial t^2} + \omega_{ci}^2\right)\mathbf{u}_{i\perp} = \frac{c\omega_{ci}^2}{B_0}\hat{\mathbf{z}} \times \nabla\phi - \frac{c\omega_{ci}}{B_0}\frac{\partial\nabla_{\perp}\phi}{\partial t},\tag{5}$$

⁷⁷ which is obtained by manipulating the perpendicular component of the ion momentum ⁷⁸ equation, where $\omega_{ci} = eB_0/m_ic$ is the ion gyrofrequency and m_i the ion mass. We have ⁷⁹ assumed that the m-KAW phase speed is much larger than the ion thermal speed, and ⁸⁰ therefore neglected in Eq. (5) the contribution of the ion pressure gradient. The ions ⁸¹ are assumed to be confined in a two-dimensional (x - y) plane perpendicular to \hat{z} . The ⁸² magnetic field-aligned ion dynamics is unimportant for the m-KAW s, since the phase ⁸³ speed of the latter is much larger than the ion-sound speed.

From the linearized ion continuity equation and Eq. (5), we readily obtain

$$\left(\frac{\partial^2}{\partial t^2} + \omega_{ci}^2\right) n_{i1} - \frac{n_0 c \omega_{ci}}{B_0} \nabla_\perp^2 \phi = 0, \tag{6}$$

where $n_{i1} \ll n_0$ is a small ion number density perturbation.

Invoking the quasi-neutrality condition $n_{e1} = n_{i1} = n_1$, we eliminate n_{i1} from Eq. (6) by using Eq. (5), obtaining the wave equation for a 3D m-KAW after elementary calculation

$$\left[\frac{\partial^2}{\partial t^2} - C_A^2 \rho_s^2 \frac{\partial^2 \nabla_\perp^2}{\partial z^2} - \frac{C_A^2}{\omega_{ci}^2} \left(\frac{\partial^2}{\partial t^2} + \omega_{ci}^2\right) \frac{\partial^2}{\partial z^2}\right] n_1 = 0, \tag{7}$$

June 21, 2021, 10:07am D R A F T

X - 7

where $C_A = B_0/\sqrt{4\pi n_0 m_i}$ is the Alfvén speed, and $\rho_s = C_s/\omega_{ci}$ is the sound gyroradius, with $C_s = (k_B T_e/m_i)^{1/2}$ being the ion-sound speed. The effect of the ion temperature can be incorporated by re-defining $\rho_s = (C_s/\omega_{ci})(1 + 3T_i/T_e)^{1/2}$.

⁹¹ Within the framework of a plane-wave approximation, assuming that n_1 is proportional ⁹² to $\exp(-i\omega t + i\mathbf{k}\cdot\mathbf{r})$, where ω and $\mathbf{k}(=\mathbf{k}_{\perp} + \hat{\mathbf{z}}k_z)$ are the angular frequency and the wave ⁹³ vector, respectively, we Fourier analyze (8) to obtain the frequency spectra Shukla and ⁹⁴ Stenflo [1999] of the modified (by the ω/ω_{ci} effect) KAWs

$$\omega^2 = \frac{k_z^2 V_A^2 (1 + k_\perp^2 \rho_s^2)}{1 + k_z^2 \lambda_i^2},\tag{8}$$

where \mathbf{k}_{\perp} and k_z are the components of \mathbf{k} across and along $\hat{\mathbf{z}}$, and $\lambda_i = c/\omega_{pi}$ the ion inertial scale length, with $\omega_{pi} = (4\pi n_0 e^2/m_i)^{1/2}$ being the ion plasma frequency. The $k_z \lambda_i$ -term in Eq. (8) comes from the ω/ω_{ci} effect.

Let us now study the property of a twisted m-KAW. We seek a solution of Eq. (7) in the form

$$n_1 = N(r) \exp(ik_z z - i\omega_k t), \tag{9}$$

where N(r) is a slowly varying function of z. Here $r = (x^2 + y^2)^{1/2}$ and k is the propagation wave number along the axial (the z- axis) direction. By using Eq. (10) we can write Eq. (7) in a paraxial approximation (viz. $|\partial^2 N/\partial z^2| \ll k_z^2 N$) as

$$\left(2i\frac{\partial}{\partial Z} + \nabla_{\perp}^2\right)N = 0,\tag{10}$$

where $\omega_k = k_z V_A / (1 + k_z^2 \lambda_i^2)^{1/2}$ is the angular frequency of the magnetic field-aligned KAW. Here $Z = (1 + k_z^2 \lambda_i^2) k_z z$ and ∇_{\perp} is in unit of ρ_s . Furthermore, we have denoted the operator $\nabla_{\perp}^2 N = (1/r)(\partial/\partial r)(r\partial N/\partial r) + (1/r^2)\partial^2 N/\partial\theta^2$, and introduced the cylindrical coordinates with $\mathbf{r} = (r, \theta, z)$.

The solution of Eq. (10) can be written as a superposition of Laguerre-Gaussian (LG) modes [Allen *et al.*, 1992, 2003; Mendonça *et al.*, 2009], each of them representing a state of orbital angular momentum, characterized by the quantum number l, such that

$$N = \sum_{pl} N_{pl} F_{pl}(r, Z) \exp(il\theta), \qquad (11)$$

where the mode structure function is $F_{pl}(r, z) = H_{pl}X^{|l|}L_p^{|l|}(X)\exp(-X/2)$, with $X = R^2/W^2(Z)$, $r = r/\rho_s$, and $W = 2\pi(1 + 4\pi^2\lambda_i^2/L_z^2)z/L_z$, $L_z = 2\pi/k_z$, and $W(Z)/L_z$ representing the normalized width of a twisted Alfvénic vortex beam. The normalization factor H_{pl} and the associated Laguerre polynomial $L_p^{|l|}(X)$ are, respectively,

$$H_{pl} = \frac{1}{2\sqrt{\pi}} \left[\frac{(l+p)!}{p!} \right]^{1/2},$$
(12)

114 and

X - 8

$$L_{p}^{|l|}(X) = \frac{\exp(X)}{Xp!} \frac{d^{p}}{dX^{p}} \left[X^{l+p} \exp(-X) \right],$$
(13)

where p and l are the radial and angular mode numbers of the DSAW orbital angular momentum state. In a special case with l = 0 and p = 0, we have a Gaussian beam.

The LG solutions, given by Eq. (11), describe the structure of a twisted 3D Alfvénic 117 vortex beam carrying OAM. In a twisted m-KAW vortex beam, the wavefront rotates 118 around the beam's propagation direction (viz. the z-axis) in a spiral that looks like fusilli 119 pasta (or a bit like a DNA double helix), creating a vortex and leading to the m-KAW 120 vortex (KAWV) beam with zero intensity at its center. A twisted m-KAWV beam (or 121 an Alfvén tornado) can be created with the help of two oppositely propagating 3D m-122 KAWs that are colliding in a magnetoplasma. Twisting of 3D m-KAWs occurs because 123 different sections of the wavefront bounce off different steps, introducing a delay between 124 the reflection of neighboring sections and, therefore, causing the wavefront to be twisted 125 due to an entanglement of the wavefronts. Thus, due to angular symmetry, Noether 126 theorem guards OAM conservation for a m-KAWV beam that is accompanies the parallel 127 electric field, sheared magnetic field, and finite density perturbations. 128

3. Summary and discussions

In summary, we have shown that 3D m-KAWs in a uniform magnetoplasma can prop-129 agate in the form of Alfvénic tornadoes which have axial and radial extents of the order 130 of the ion inertial length c/ω_{pi} and the ion sound radius ρ_s , respectively. The Alfvénic 131 swirls can be identified as observational signatures of rapidly rotating magnetic field ropes 132 [Tripathi and Gekeklman, 2010; Gekelman et al., 2012] (or magnetic field tornadoes), 133 which can provide an alternative mechanism for channelling electromagnetic energy and 134 heat fluxes from the surface of the Sun into the corona Widemever-Böhm et al. [2012] 135 through Alfvénic tornadoes/whirls. Furthermore, the present investigation of twisted m-136

¹³⁷ KAW vortex beams can also be useful for diagnostic purposes, since the m-KAW beam ¹³⁸ frequency can be a fraction of the ion gyrofrequency.

Acknowledgments. This research was partially supported by the Deutsche Forschungsgemeinschaft (DFG), Bonn, through the project SH21/3-2 of the Research Unit 1048.

References

X - 10

- ¹⁴² Alfvén, H. (1942), Existence of electromagnetic hydromagnetic waves, Nature (London)
 ¹⁴³ 150, 405.
- Shukla P. K. and J. M. Dawson (1984), Stimulated scattering of hydromagnetic waves in
 the interstellar medium, Astrophys. J. Lett. 276, L49-L51.
- Leneman, D., W. Gekelman, and J. Maggs (1999), Laboratory observations of shear Alfvén
 waves launched from a small source, Phys. Rev. Lett. 82, 2673-2676.
- Kletzing, C. A., D. J. Thuecks, F. Skiff, S. R. Bounds, and S. Vienna (2010), Measurements
 of inertial limit Alfvén wave dispersion for finite perpendicular wave number, Phys. Rev.
- Lett. 104, 095001.
- ¹⁵¹ Gekelman, W., S. Vincena, B. Van Compernolle, G. J. Morales, J. E. Maggs, P. Pribyl,
 ¹⁵² and T. A. Carter (2011), The many faces of shear Alfvén waves, Phys. Plasmas 18,
- 153 055501.
- ¹⁵⁴ Cramer, N. F. (2001), The Physics of Alfvén Wave, Wiley, Berlin.
- Stefant, R. J. (1970), Alfvén wave damping from finite gyroradius coupling to the ion
 acoustic mode, Phys. Fluids 13, 440-450.

- G. Brodin and L. Stenflo (1990), Coupling coefficients for ion-cyclotron Alfvén waves,
 Contrib. Plasma Phys. 30, 413-419.
- ¹⁵⁹ Morales, G. J., R. S. Loritsch, and J. E. Maggs (1994), Structure of Alfv—'en waves at ¹⁶⁰ the skin-depth scale, Phys. Plasmas *1*, 3765-3774.
- ¹⁶¹ Morales G. J. and J. E. Maggs (1997), Structure of kinetic Alfvén waves with small
 ¹⁶² transverse scale length, Phys. Plasmas 4, 4118-4125.
- ¹⁶³ Shukla P. K. and L. Stenflo (1999), in Nonlinear MHD Waves and Turbulence, Eds. T.
 ¹⁶⁴ Passot and P. L. Sulem, Springer, Berlin), pp. 1-30.
- ¹⁶⁵ Dastgeer D. and P. K. Shukla (2009), 3D simulations of fluctuation spectra of the Hall-
- ¹⁶⁶ MHD plasma, Phys. Rev. Lett. *102*, 045004.
- Damiano, P. A., A. N. Wright, and J. F. McKenzie (2009), Properties of Hall magnetohy drodynamic waves modified by electron inertia and finite Larmor radius effects, Phys.
 Plasmas 16, 062901.
- ¹⁷⁰ Chaston, C. C., C. Salem, J. W. Bonnell, C. W. Carlson, R. E. Ergun, R. J. Stangeway,
 ¹⁷¹ and J. R. McFadden (2008), The turbulent Alfvénic aurora, Phys. Rev. Lett. 100,
 ¹⁷² 175003.
- Koepke, M. E., S. M. Finnegan, S. Vincena, D. J. Knudsen, and Chaston, C. C. (2008),
 Integrated campaign to study the stationary inertial Alfvén wave in the laboratory and
 space regimes, Plasma Phys. Controll. Fusion 50, 07004.
- ¹⁷⁶ Sundkvist, D., V. Krasnoselskikh, P. K. Shukla, A. Vivads, M. Andre, S. Buchert, and H.
- Reme (2005), In situ multi-satellite detection of coherent vortices as a manifestation of
 Alfvénic turbulence, Nature (London) 436, 825-828.

- ¹⁷⁹ Tripathi S. K. P. and W. Gekelman (2010), Laboratory simulation of arched magnetic ¹⁸⁰ flux rope eruption in the solar atmosphere, Phys. Rev. Lett. *105*, 075005.
- ¹⁸¹ Salem, C. S., G. G. Howes, D. Sundkvist, C. C. Chaston, C. H. K. Chen, and F. S. Mozer
- (2012), Identification of kinetic Alfvén wave turbulence in the solar wind, Astrophys. J.
 Lett. 745, L9.
- Gekelman, W., E. Lawrence, and B. Van Compernolle (2012), Three-dimensional recon nection involving magnetic flux ropes, Astrophys J. 753, 131.
- Shukla, P. K., R. Bingham, J. F. McKenzie, and I. Axford (1998), Solar coronal heating
 by high-frequency dispersive Alfvén waves, Solar Phys. 186, 61-66.
- A. Hasegawa and L. Chen (1976), Parametric decay of "Kinetic Alfvén Wave" and its
 applications, Phys. Rev. Lett. 36, 1362-1365.
- ¹⁹⁰ Carter, T. A., B. Brugman, P. Pribyl, and W. L. Lybarger (2006), Laboratory observation
 ¹⁹¹ of a nonlinear interaction between shear Alfvén waves, Phys. Rev. Lett. *96*, 155001.
- ¹⁹² Shukla, P. K., B. Eliasson, L. Stenflo, and R. Bingham (2007), in Recent Research Devel-
- opments in Plasma Physics, Ed. J. Weiland, Transworld Research Network, Trivendrum,
 Kerala, India, pp. 51-74.
- Stasiewicz, K., P. K. Shukla, G. Gustafsson, S. Buchert, B. Lavraud, B. Thide, and Z.
 KLo (2003), Slow magnetosonic solitons detection by the Cluster spacecraft, Phys. Rev.
- Shukla, P. K., B. Eliasson, and L. Stenflo (2012), Alfvénic solitary and shock waves in
 plasmas, in multi-scale Dynamical Processes in Space and Astrophysical Plasmas, Eds.
- M. P. Leubner and Z. Vörös, Astrophys. and Space Sci. Proceedings (Springer, Berlin),

DRAFT

Lett. 90, 085002.

197

33, 129-141.

- ²⁰² Tomczyk, S., S. W. McIntosh, S. L. Keil, P. G. Judge, T. Schad, D. H. Seeley, and J.
- Edmondson (2007), Alfvén waves in the solar corona, Science 317, 1192-1196.
- ²⁰⁴ Jess, D. B., M. Mathioudakis, R. Erdélyi, P. J. Crockett, F. P. Keenan, and D. J. Christian
- ²⁰⁵ (2009), Alfvén waves in the lower solar atmosphere, Science *323*, 1582-1585.
- ²⁰⁶ McIntosh, S. W., B. De Pontieu, M. Carlsson, V. Hansteen, P. Boerner, and M. Goosens
- (2011), Alfvénic waves with sufficient energy to power the quit solar corona and fast
 solar wind, Nature (London) 475, 477-480.
- ²⁰⁹ Widemeyer-Böhm, S., E. Scullion, O. Steiner, L. R. van der Voort, J. de la Cruz Rodriguez,
- V. Fedun, R. Erdélyi (2012), Magnetic tornadoes as energy channels into the solar
 corona, Nature (London) 486, 505-508.
- Shukla, P. K. (2012), Twisted shear Alfvén waves with orbital angular momentum, Phys.
 Lett. A 376, in press, http://dx.doi.org/101016/j.physleta.2012.08.025.
- Allen, L., M. W. Beijersbergen, R. J. C. Spereeuw, and J. P. Woerdman (1992), Orbital
 angular momentum and the transformation of Laguerre-Gaussian laser modes, Phys.
 Rev. A 45, 8185-8189.
- Allen, L., S. M. Barnett, and M. J. Padgett (2003), Orbital Angular Momentum, Institute
 of Physics, Bristol.
- ²¹⁹ Mendonça, J. T., B. Thidé, and H. Then (2009), Stimulated Raman and Brillouin
- backscattering of collimated beams carrying orbital angular momentum, Phys. Rev.
 Lett. 102, 1850050.