

Inverse boundary value problem for the dynamical heterogeneous Maxwell system

Mourad Bellassoued*, Michel Cristofol† and Eric Soccorsi‡

Abstract

We consider the inverse problem of determining the isotropic inhomogeneous electromagnetic coefficients of the non-stationary Maxwell equations in a bounded domain of \mathbb{R}^3 , from a finite number of boundary measurements. Our main result is a Hölder stability estimate for the inverse problem, where the measurements are exerted only in some boundary components. For it, we prove a global Carleman estimate for the heterogeneous Maxwell's system with boundary conditions.

Key words: Inverse problems, Maxwell system, Carleman estimates.

1 Introduction

In this paper, we discuss the uniqueness and stability in determining the isotropic electromagnetic coefficients of the dynamical Maxwell equations, by boundary measurement of their solution. More precisely, given a continuous medium with dielectric permittivity λ^{-1} and magnetic permeability μ^{-1} , occupying an open, bounded and simply connected domain $\Omega \subset \mathbb{R}^3$ with C^∞ boundary $\Gamma = \partial\Omega$, and $T > 0$, we consider the following problem for the linear system of Maxwell's equations

$$\begin{aligned} D' - \operatorname{curl}(\mu B) &= 0, & \text{in } Q := \Omega \times (-T, T), \\ B' + \operatorname{curl}(\lambda D) &= 0, & \text{in } Q, \\ \operatorname{div} D = \operatorname{div} B &= 0, & \text{in } Q, \\ D \times \nu = 0, \quad B \cdot \nu &= 0, & \text{on } \Sigma := \Gamma \times (-T, T), \end{aligned} \tag{1.1}$$

where the prime stands for the time derivative. Here the electric induction D and the magnetic field B are three-dimensional vector-valued functions of the time t and the space variable $x = (x_1, x_2, x_3)$, and $\nu = \nu(x)$ denotes the unit outward normal vector to Γ . Moreover we attach the following initial condition to (1.1):

$$B(x, 0) = B_0(x), \quad D(x, 0) = D_0(x), \quad x \in \Omega. \tag{1.2}$$

Assume that μ and λ are scalar functions in $C^2(\overline{\Omega})$ obeying

$$\mu(x) \geq \mu_0, \quad \lambda(x) \geq \lambda_0, \quad x \in \overline{\Omega}, \tag{1.3}$$

*University of Carthage, Faculty of Sciences of Bizerte, Department of Mathematics, 7021 Jarzouna Bizerte, Tunisia: mourad.bellassoued@fsb.rnu.tn

†Aix-Marseille Université, CNRS, LATP, UMR 7353, 13453 Marseille, France: cristo@cmi.univ-mrs.fr

‡Aix Marseille Université, CNRS, CPT, UMR 7332, 13288 Marseille, France: soccorsi@cpt.univ-mrs.fr

for some $\lambda_0 > 0$ and $\mu_0 > 0$. Next, in view of deriving existence and uniqueness results for (1.1), introduce the following functional space

$$H(\text{curl}, \Omega) := \{u \in L^2(\Omega)^3, \text{curl } u \in L^2(\Omega)^3\},$$

and denote by γ_τ the unique linear continuous application from $H(\text{curl}, \Omega)$ into $H^{-1/2}(\Gamma)^3$, satisfying $\gamma_\tau u = u \wedge \nu$ when $u \in C_0^\infty(\overline{\Omega})^3$ (see [16][Chap. IX A, Theorem 2]). Then, putting

$$H_0(\text{curl}, \Omega) := \{u \in H(\text{curl}, \Omega), \gamma_\tau = 0\},$$

we see that the operator iA , where

$$A\Phi := \begin{pmatrix} 0 & \text{curl}(\mu \cdot) \\ -\text{curl}(\lambda \cdot) & 0 \end{pmatrix}, \quad \Phi = (D, B) \in \text{Dom}(A) := H_0(\text{curl}; \Omega) \times H(\text{curl}; \Omega),$$

is selfadjoint in $\mathcal{H} := L^2(\Omega)^3 \times L^2(\Omega)^3$, endowed with the scalar product

$$\langle \Phi, \tilde{\Phi} \rangle_{\mathcal{H}} := \langle \lambda D, \tilde{D} \rangle_{L^2(\Omega)^3} + \langle \mu B, \tilde{B} \rangle_{L^2(\Omega)^3}, \quad \Phi = (D, B) \in \mathcal{H}, \quad \tilde{\Phi} = (\tilde{D}, \tilde{B}) \in \mathcal{H}.$$

Further, in light of the last line of (1.1), set

$$H(\text{div } 0, \Omega) := \{u \in L^2(\Omega)^3, \text{div } u = 0\} \text{ and } H_0(\text{div } 0, \Omega) := \{u \in H(\text{div } 0, \Omega), \gamma_n u = 0\},$$

where γ_n is the unique linear continuous mapping from $H(\text{div}, \Omega) := \{u \in L^2(\Omega)^3, \text{div } u \in L^2(\Omega)\}$ onto $H^{-1/2}(\Gamma)$, such that $\gamma_n u = u \cdot \nu$ when $u \in C_0^\infty(\overline{\Omega})$ (see [16][Chap. IX A, Theorem 1]). Since $\mathcal{H}_0 := H(\text{div } 0; \Omega) \times H_0(\text{div } 0; \Omega)$ is a closed subspace of \mathcal{H} and that $\mathcal{H}_0^\perp \subset \ker A$, the restriction

$$A_0\Phi = A_{\mathcal{H}_0}\Phi := A\Phi, \quad \Phi \in \text{Dom}(A_0) = \text{Dom}(A) \cap \mathcal{H}_0 := \mathcal{V},$$

is, by Stone's Theorem [17][Chap. XVII A, §4, Theorem 3], the infinitesimal generator of a unitary group of class \mathcal{C}^0 in \mathcal{H}_0 . Thus, by rewriting (1.1)-(1.2) into the equivalent form

$$\begin{cases} \Phi' = A_0\Phi \\ \Phi(0) = \Phi_0, \end{cases} \quad \text{with } \Phi = (D, B)^T \text{ and } \Phi_0 = (D_0, B_0)^T,$$

we get that:

Lemma 1.1 *Given $(D_0, B_0) \in \mathcal{V}$ there exists a unique strong solution (D, B) to (1.1) starting from (D_0, B_0) within the following class*

$$(D, B) \in \mathcal{C}^0(\mathbb{R}; \mathcal{V}) \cap \mathcal{C}^1(\mathbb{R}; \mathcal{H}). \quad (1.4)$$

Moreover it holds true from [16][Chap. IX A, Remark 1] that

$$\mathcal{V} = H_{\tau,0}(\text{curl}, \text{div } 0; \Omega) \times H_{n,0}(\text{curl}, \text{div } 0; \Omega),$$

where

$$H_{*,0}(\text{curl}, \text{div } 0; \Omega) = \{u \in H^1(\Omega)^3, \text{div } u = 0 \text{ and } \gamma_* u = 0\}, \quad * = \tau, n.$$

For further reference we notice from Lemma 1.1 that the solution (D, B) to (1.1)-(1.2) actually satisfies:

$$(D, B) \in \cap_{p=0}^m \mathcal{C}^p([-T, T]; \text{Dom}(A_0^{m-p})) \text{ provided } (D_0, B_0) \in \text{Dom}(A_0^m) \text{ and } \lambda, \mu \in \mathcal{C}^m(\overline{\Omega}), m \geq 1. \quad (1.5)$$

The main purpose of this paper is to study the inverse problem of determining the dielectric permittivity $\lambda^{-1} = \lambda^{-1}(x)$ and the magnetic permeability $\mu^{-1} = \mu^{-1}(x)$ from a finite number of observations on the boundary Γ of the solution (B, D) to (1.1) which corresponds to a realistic physical approach. This is an important problem not only in electromagnetics (see [41]) but also in the identification of cracks/flaws in conductors (see [19]) or the localization of lightning discharges (see [38]). On the other hand, we obtain a reconstruction result which involves only a finite number of measurements which is not the case in most of the existing results.

1.1 Inverse problem

For suitable $B_0^k, D_0^k, k = 1, 2$, we aim to determine $\lambda(x), \mu(x), x \in \Omega$, from the observation of

$$B_\tau^k(x, t), \quad D_\nu^k(x, t), \quad (x, t) \in \Sigma, \quad k = 1, 2,$$

where $B_\tau = B - (B \cdot \nu)\nu$ (resp. $D_\nu = (D \cdot \nu)\nu$) denotes the tangential (resp. normal) component of B (resp. D).

Notice that only a finite number of measurements are needed in the formulation of this inverse problem. For an overview of inverse problems for the Maxwell system, see the monograph [39] by Romanov and Kabanikhin. For actual examples of inverse problems for the dynamical Maxwell system involving infinitely many boundary observations (this is the case when the identification of the electromagnetic coefficients is made from the Dirichlet-to-Neumann map), we refer to Beleshev and Isakov [3], Caro [12], Caro, Ola and Salo [13], Kurylev, Lassas and Somersalo [31], Ola, Paivarinta and Somersalo [35] and Salo, Kenig and Uhlmann [40]. It turns out that a small number of uniqueness and stability results for the inverse problem of determining the electromagnetic parameters of the Maxwell system with a finite number of measurements are available, such as [33, 34]. In both cases, their proof is based on the methodology of [10] or [23], which is by means of a Carleman estimate.

For the formulation with a finite number of observations, Bukhgeim and Klivanov [10] proposed a remarkable method based on a Carleman estimate and established the uniqueness for similar inverse problems for scalar partial differential equations. See also Bellassoued [1], [2], Bellassoued and Yamamoto [4], [5], A. Benabdallah, M. Cristofol, P. Gaitan and M. Yamamoto [7], Bukhgeim [8], Bukhgeim, Cheng, Isakov and Yamamoto [9], Cristofol and Roques [14], Cristofol and Soccorsi [15], Imanuvilov and Yamamoto [22]-[23], Isakov [24], Khaïdarov [26], Klivanov [27], [28], Klivanov and Timonov [29], Klivanov and Yamamoto [30], Li and Yamamoto [33]-[34], Yamamoto [43].

A Carleman estimate is an inequality for a solution to a partial differential equation with weighted L^2 -norm and is a strong tool also for proving the uniqueness in the Cauchy problem or the unique continuation for a partial differential equation with non-analytic coefficients. Moreover Carleman estimates have been applied essentially for estimating the energy (e.g., Kazemi and Klivanov [25]).

As a pioneering work concerning a Carleman estimate, we refer to Carleman's paper [11] which proved what is later called a Carleman estimate and applied it for proving the uniqueness in the Cauchy problem for a two-dimensional elliptic equation. Since [11], the theory of Carleman estimates has been studied extensively. We refer to a general theory by Hörmander [20] in the case where the symbol of a partial differential equation is isotropic and functions under consideration have compact supports (that

is, they and their derivatives of suitable orders vanish on the boundary of a domain). Later Carleman estimates for functions with compact supports have been obtained for partial differential operators with anisotropic symbols by Isakov [24]. Carleman estimates for functions without compact supports, see Imanuvilov [21], Tataru [42]. As for a direct derivation of pointwise Carleman estimates for hyperbolic equations which are applicable to functions without compact supports, see Klivanov and Timonov [29], Lavrent'ev, Romanov and Shishat'skiĭ [32].

The Carleman estimate for the non-stationary Maxwell's system was obtained for functions with compact supports, by Eller, Isakov, Nakamura and Tataru [18]. Li and Yamamoto [33]-[34], prove a Carleman estimate for two-dimensional Maxwell's equations in isomagnetic anisotropic media for functions with compact supports. Lemmas 2.1 and 2.2 are our Carleman estimate for the Maxwell's system whose solutions have not necessarily compact supports.

By the methodology by [10] or [23] with such Carleman estimates, several uniqueness and stability results are available for the inverse problem for the Maxwell's system (1.1). That is, in [33]-[34] Li and Yamamoto established the uniqueness in determining three coefficients, using finite number of measurements.

Li and Yamamoto [34], consider nonstationary Maxwell's equations in an anisotropic medium in the (x_1, x_2, x_3) -space, where equations of the divergences of electric and magnetic flux densities are also unknown. Then they discuss an inverse problem of determining the x_3 -independent components of the electric current density from observations on the plane $x_3 = 0$ over a time interval and prove conditional stability in the inverse problem provided the permittivity and the permeability are independent of x_3 .

In [37], S.Nicaise and C.Pignotti, consider the Heterogenous Maxwell's system defined in an open bounded domain. Under checkable conditions on the coefficients of the principal part they proved a Carleman type estimates where some weighted H^1 -norm of solution is dominated by the L^2 norm of the boundary traces $\partial_\nu U$ and U_t , modulo an interior lower-order term. Once homogeneous boundary conditions are imposed the lower-order term can be absorbed by the standard unique continuation theorem. Unfortunately, to our knowledge, these results may not be applied directly to the linearized inverse problem associated to the original problem.

Our argument is based on a new Carleman estimate. In comparison with [33] and [37], our Carleman estimate is advantageous in the following two points:

- We show a Carleman estimate which holds over the whole domain Q . We need not assume that the functions under consideration have compact supports and so ours is different from the Carleman estimates presented in [33], and we can establish a Hölder estimate.
- We do not need a priori any unique continuation property and compactness/uniqueness argument to absorb the lower-order interior term. In our approach, we establish a Carleman estimates for H^1 -solutions of the hyperbolic equation with variable coefficients. This is essential to the proof of our main result, because here our problem is involved with a source term and we cannot use the standard compactness/ uniqueness argument as in [37].

1.2 Notations and statement of the main result

In this subsection we introduce some notations used throughout this text and state the main result of this article. Pick $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$, set $c(x) = \mu(x)\lambda(x)$ for $x \in \overline{\Omega}$, $c_0 = \mu_0\lambda_0$ where μ_0 and λ_0 are the same as

in (1.3), and assume that the following condition

$$\frac{3}{2} |\nabla \log c(x)| |x - x_0| \leq 1 - \frac{\rho}{c_0}, \quad x \in \overline{\Omega}, \quad (1.6)$$

holds true for some $\rho \in (0, c_0)$. This purely technical condition was imposed by the method we use to solve the inverse problem under study, which is by means of the Carleman estimate stated in Lemma 2.2 for any weight function ψ_0 satisfying the two Assumptions (A1) and (A2). More precisely, in the particular case where

$$\psi_0(x) := |x - x_0|^2, \quad x \in \Omega, \quad (1.7)$$

then (1.6) arises from the classical pseudo-convexity condition expressed by (2.5). The somehow non-natural condition (1.6) is thus closely related to the peculiar expression (1.7) in the sense that another choice of ψ_0 fulfilling (A1) and (A2) may eventually lead to a completely different condition on $c(x)$.

Next, for $M_0 > 0$ and two given functions $\mu^\sharp, \lambda^\sharp \in C^2(\omega)$, where $\omega = \Omega \cap \mathcal{O}$ for some neighbourhood \mathcal{O} of Γ in \mathbb{R}^3 , we define the admissible set of unknown coefficients μ and λ as

$$\Lambda_\omega(M_0) = \left\{ (\mu, \lambda) \text{ obeying (1.3) and (1.6)}; \|(\mu, \lambda)\|_{C^2(\overline{\Omega})} \leq M_0 \text{ and } (\mu, \lambda) = (\mu^\sharp, \lambda^\sharp) \text{ in } \omega \right\}. \quad (1.8)$$

Further, the identification of (λ, μ) imposing, as will appear in the sequel, that (B, D) be observed twice, we consider two sets of initial data (D_0^k, B_0^k) , $k = 1, 2$,

$$D_0^k(x) = \left(d_1^k(x), d_2^k(x), d_3^k(x) \right)^\top, \quad B_0^k(x) = \left(b_1^k(x), b_2^k(x), b_3^k(x) \right)^\top, \quad (1.9)$$

and define the 12×6 matrix

$$\mathcal{K}(x) = \begin{pmatrix} e_1 \times B_0^1 & e_2 \times B_0^1 & e_3 \times B_0^1 & 0 & 0 & 0 \\ 0 & 0 & 0 & e_1 \times D_0^1 & e_2 \times D_0^1 & e_3 \times D_0^1 \\ e_1 \times B_0^2 & e_2 \times B_0^2 & e_3 \times B_0^2 & 0 & 0 & 0 \\ 0 & 0 & 0 & e_1 \times D_0^2 & e_2 \times D_0^2 & e_3 \times D_0^2 \end{pmatrix}, \quad x \in \Omega. \quad (1.10)$$

We then write $(B_i^k(x, t), D_i^k(x, t))$ the solution to (1.1) with initial data (B_0^k, D_0^k) , $k = 1, 2$, where (μ_i, λ_i) , $i = 1, 2$, is substituted for (μ, λ) .

Finally, noting $\mathcal{H}(\Sigma) = H^3(-T, T; L^2(\Gamma)) \cap H^2(-T, T; H^1(\Gamma))$ the Hilbert space equipped with the norm

$$\|u\|_{\mathcal{H}(\Sigma)}^2 = \|u\|_{H^3(-T, T; L^2(\Gamma))}^2 + \|u\|_{H^2(-T, T; H^1(\Gamma))}^2, \quad u \in \mathcal{H}(\Sigma),$$

we now may state the main result of this paper as follows :

Theorem 1 *Let $T > c_0^{-1/2} \max_{x \in \overline{\Omega}} |x - x_0|$ and pick $(B_0^k, D_0^k) \in (H^2(\Omega)^3 \times H^2(\Omega)^3) \cap \mathcal{V}$, $k = 1, 2$, in such a way that there exists a 6×6 minor $m(x)$ of the matrix $\mathcal{K}(x)$ defined in (1.10), obeying:*

$$m(x) \neq 0, \quad x \in \overline{\Omega \setminus \omega}. \quad (1.11)$$

Further, choose $(\mu_i, \lambda_i) \in \Lambda_\omega(M_0)$, $i = 1, 2$, so that

$$\left\| \left(B_i^k, D_i^k \right) \right\|_{C^3(-T, T; W^{2, \infty}(\Omega))} \leq M, \quad k = 1, 2, \quad (1.12)$$

for some $M > 0$. Then there are two constants $C > 0$ and $\kappa \in (0, 1)$, depending on Ω, ω, T, M and M_0 , such that we have:

$$\|\mu_1 - \mu_2\|_{H^2(\Omega)} + \|\lambda_1 - \lambda_2\|_{H^2(\Omega)} \leq C \left(\sum_{k=1}^2 \left(\left\| (B_1^k - B_2^k)_\tau \right\|_{\mathcal{H}(\Sigma)} + \left\| (D_1^k - D_2^k)_\nu \right\|_{\mathcal{H}(\Sigma)} \right) \right)^\kappa.$$

Notice that the condition (1.11), which is independent of the choice of the unknown coefficients μ and λ , actually relates on the initial functions in (1.2). Moreover this condition is stable with respect to perturbations in \mathcal{C}^2 . Namely, if (B_0^k, D_0^k) obeys (1.11) then this is the case for $(\tilde{B}_0^k, \tilde{D}_0^k)$ as well, provided $\max_{k=1,2} \left\| (B_0^k, D_0^k) - (\tilde{B}_0^k, \tilde{D}_0^k) \right\|_{\mathcal{C}(\Omega)}$ is sufficiently small. Furthermore there are actual choices

of (B_0^k, D_0^k) , $k = 1, 2$, satisfying (1.11). This can be seen by taking

$$B_0^1(x) = e_1, \quad D_0^1(x) = e_3, \quad B_0^2(x) = e_2, \quad D_0^2(x) = e_2, \quad x \in \overline{\Omega} \setminus \omega$$

and selecting the 6×6 minor formed by rows 2, 3, 4, 9, 10 and 12.

Theorem 1 asserts Hölder stability in determining the principal part within the class defined by (1.8), under the assumption (1.12). Notice from (1.5) that such a condition is automatically fulfilled for $\lambda, \mu \in \mathcal{C}^7(\overline{\Omega})$ by choosing the initial data (B_0^k, D_0^k) , $k = 1, 2$, in $\text{Dom}(A_0^T)$ (which is a dense in \mathcal{H}).

The proof of Theorem 1 is based on a Carleman estimate stated in Lemma 2.2 under the conditions (1.3) and (1.6). Notice that (1.6), which is essential to our argument, is much stronger than the usual uniform ellipticity condition.

The remainder of the paper is organized as follows: a Carleman estimate for the Maxwell system (1.1) is established in Section 2, while Section 3 contains the proof of Theorem 1.

2 Carleman estimate for Maxwell's system

As already mentioned, this section is devoted to the derivation of a global Carleman estimate for the Maxwell system (1.1).

2.1 The settings

Let us consider the following second order hyperbolic operator

$$Pu = \partial_t^2 u(x, t) - \text{div}(c(x)\nabla u) + \mathcal{R}_1(x, t; \partial)u, \quad x \in \Omega, \quad t \in \mathbb{R}, \quad (2.1)$$

where \mathcal{R}_1 is a first order partial operator with $L^\infty(\Omega \times \mathbb{R})$ coefficients, and $c \in \mathcal{C}^2(\overline{\Omega})$ obeys

$$c(x) \geq c_0, \quad x \in \overline{\Omega}, \quad (2.2)$$

for some positive constant c_0 . Putting

$$a(x, \xi) = c(x) |\xi|^2, \quad x \in \Omega, \quad \xi \in \mathbb{R}^3, \quad (2.3)$$

and recalling the definition of the Poisson bracket of two given symbols p and q ,

$$\{p, q\}(x, \xi) = \frac{\partial p}{\partial \xi} \cdot \frac{\partial q}{\partial x} - \frac{\partial p}{\partial x} \cdot \frac{\partial q}{\partial \xi} = \sum_{i=1}^n \left(\frac{\partial p}{\partial \xi_i} \frac{\partial q}{\partial x_i} - \frac{\partial p}{\partial x_i} \frac{\partial q}{\partial \xi_i} \right),$$

we introduce two assumptions.

Assumption (A1). There exists $\psi_0 \in \mathcal{C}^2(\overline{\Omega}; \mathbb{R}_+^*)$ satisfying

$$\{a, \{a, \psi_0\}\}(x, \xi) > 0, \quad x \in \overline{\Omega}, \quad \xi \in \mathbb{R}^3 \setminus \{0\}, \quad (2.4)$$

where a is given by (2.3).

Since $\overline{\Omega}$ is compact and $a(x, \xi)$ is a homogenous function with respect to ξ , it is clear that (2.4) yields the existence of some constant $\varrho > 0$ such that we have:

$$\frac{1}{4} \{a, \{a, \psi_0\}\}(x, \xi) \geq 2\varrho c(x) |\xi|^2, \quad x \in \overline{\Omega}, \quad \xi \in \mathbb{R}^3 \setminus \{0\}. \quad (2.5)$$

Assumption (A2). The function $\psi_0(x)$ has no critical points on $\overline{\Omega}$:

$$\min_{x \in \overline{\Omega}} |\nabla \psi_0(x)|^2 > 0.$$

Further, ϱ being the same as in (2.5), fix $\delta > 0$ and $\beta \in (0, \varrho)$, in such a way that, upon eventually enlarging T , we have:

$$\beta T^2 > \max_{x \in \overline{\Omega}} \psi_0(x) + \delta. \quad (2.6)$$

Hence, picking $\beta_0 > 0$ and setting

$$\psi(x, t) = \psi_0(x) - \beta t^2 + \beta_0, \quad x \in \overline{\Omega}, \quad t \in [-T, T], \quad (2.7)$$

so that

$$\min_{x \in \overline{\Omega}} \psi(x, 0) \geq \beta_0,$$

we check out from (2.6) that

$$\psi(x, \pm T) \leq \beta_0 - \delta, \quad x \in \overline{\Omega}. \quad (2.8)$$

Notice from (2.7) and (2.8) that

$$\max_{x \in \overline{\Omega}} \psi(x, t) \leq \beta_0 - \frac{\delta}{2}, \quad |t| \in (T - 2\epsilon, T], \quad (2.9)$$

for some constant $\epsilon \in (0, T/2)$.

In view of (2.7) we may now recall the following global Carleman estimate for second order scalar hyperbolic equations, with weight function $\varphi : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$\varphi(x, t) = e^{\gamma \psi(x, t)}, \quad x \in \overline{\Omega}, \quad t \in [-T, T], \quad (2.10)$$

for some fixed $\gamma > 0$.

Theorem 2 Assume (A1)-(A2). Then there exist two constants $C_0 > 0$ and $s_0 > 0$ such that for every $s \geq s_0$ the following Carleman estimate

$$\begin{aligned} & C_0 \int_Q e^{2s\varphi} s \left(|\nabla v|^2 + |\partial_t v|^2 + s^2 |v|^2 \right) dx dt \\ & \leq \int_Q e^{2s\varphi} |Pv(x, t)|^2 dx dt + \int_\Sigma s e^{2s\varphi} \left(\left(|\nabla v|^2 + |\partial_t v|^2 \right) + s^2 |v|^2 \right) d\sigma dt, \end{aligned} \quad (2.11)$$

holds true whenever $v \in H^1(Q)$ verifies $\partial_t^j v(\pm T, \cdot) = 0$ for $j = 0, 1$, and the right hand side of (2.11) is finite. Here P is defined by (2.1)-(2.2) and $d\sigma$ denotes the volume form of Γ .

For the proof see Bellassoued and Yamamoto [6], where this result is obtained from a direct computation based on integration by parts.

It is worth mentioning that there are actual examples of functions ψ_0 fulfilling (A1)-(A2), provided the conductivity function c defined in (2.1) verifies (1.6) for some $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$ and $\varrho \in (0, c_0)$, where c_0 is the constant defined in (2.2). Indeed, by putting $\psi_0(x) = |x - x_0|^2$ and recalling (2.3), we get through an elementary computation that

$$\frac{1}{4} \{a, \{a, \psi_0\}\}(x, \xi) = 2c^2(x) \left(1 - \frac{\nabla c \cdot (x - x_0)}{2c} \right) |\xi|^2 + 2c(\nabla c \cdot \xi)(\xi \cdot (x - x_0)),$$

so (1.6) immediately yields

$$\frac{1}{4} \{a, \{a, \psi_0\}\}(x, \xi) \geq 2\varrho c |\xi|^2.$$

This entails (A1) by (2.2). Moreover (A2) is evidently true as well since $\nabla \psi_0(x) \neq 0$ for every $x \in \overline{\Omega}$.

2.2 Decoupling of the system of equations

Consider now the following Maxwell system

$$\begin{aligned} \mathbf{U}' - \operatorname{curl}(\mu_1 \mathbf{V}) &= \mathbf{f}, & \text{in } Q, \\ \mathbf{V}' + \operatorname{curl}(\lambda_1 \mathbf{U}) &= \mathbf{g}, & \text{in } Q, \\ \operatorname{div} \mathbf{U} = \operatorname{div} \mathbf{V} &= 0, & \text{in } Q, \\ \mathbf{U} \times \nu = 0, \quad \mathbf{V} \cdot \nu &= 0, & \text{on } \Sigma, \end{aligned} \tag{2.12}$$

where the source terms $\mathbf{f}, \mathbf{g} \in H^1(Q; \mathbb{R}^3)$ satisfy the boundary condition

$$\mathbf{f}(x, t) = \mathbf{g}(x, t) = 0, \quad (x, t) \in \omega \times (-T, T). \tag{2.13}$$

For further reference we recall from (2.9) that

$$\begin{aligned} \max_{x \in \overline{\Omega}} \varphi(x, t) &\leq d_0 := e^{\gamma(\beta_0 - \delta/2)}, \quad |t| \in [T - 2\epsilon, T), \\ \min_{x \in \overline{\Omega}} \varphi(x, 0) &\geq d_1 := e^{\gamma\beta_0}, \end{aligned} \tag{2.14}$$

and then state the main result of §2.2:

Lemma 2.1 *Assume (A1)-(A2) and let $\mathbf{h} = (\mathbf{f}, \mathbf{g}) \in H^1(Q; \mathbb{R}^3)^2$ obey (2.13). Then we may find two constants $C_1 > 0$ and $s_1 > 0$, for which the Carleman estimate*

$$\begin{aligned} C_1 \int_Q e^{2s\varphi} s \left(|\nabla_{x,t} \mathbf{W}|^2 + s^2 |\mathbf{W}|^2 \right) dx dt &\leq \int_Q e^{2s\varphi} \left(|\nabla_{x,t} \mathbf{h}|^2 + |\mathbf{h}|^2 \right) dx dt + s^3 e^{2d_0 s} \|\mathbf{W}\|_{H^1(Q)}^2 \\ &\quad + \int_{\Sigma} s e^{2s\varphi} \left(|\nabla \mathbf{W}|^2 + |\mathbf{W}'|^2 + s^2 |\mathbf{W}|^2 \right) d\sigma dt, \end{aligned} \tag{2.15}$$

is true for any $\mathbf{W} = (\mathbf{U}, \mathbf{V})$ solution to the Maxwell system (2.12), whenever $s \geq s_1$.

Proof . The first step of the proof involves bringing (2.12) into two independent systems of decoupled equations. Namely, by differentiating the first line in (2.12) with respect to t , and then substituting $\mathbf{g} - \text{curl}(\lambda_1 U)$ for V' in the obtained equality, we obtain that

$$U'' + \text{curl}(\mu_1 \text{curl}(\lambda_1 U)) = \mathbf{f}' + \text{curl}(\mu_1 \mathbf{g}), \text{ in } Q.$$

This entails $U'' + \text{curl}(\mu_1 \lambda_1 \text{curl} U) + \text{curl}(\mu_1 \nabla \lambda_1 \times U) = \mathbf{f}' + \text{curl}(\mu_1 \mathbf{g})$, and hence

$$U'' + \mu_1 \lambda_1 \text{curl}(\text{curl} U) + \nabla(\mu_1 \lambda_1) \times \text{curl} U + \text{curl}(\mu_1 \nabla \lambda_1 \times U) = \mathbf{f}' + \text{curl}(\mu_1 \mathbf{g}), \text{ in } Q.$$

From this, the well-known identity $\text{curl} \text{curl} U = -\vec{\Delta} U + \nabla \text{div} U$ and the third line of (2.12) then follows that

$$U'' - \mu_1 \lambda_1 \vec{\Delta}(U) + \mathcal{R}_1 U = \mathbf{f}' + \text{curl}(\mu_1 \mathbf{g}), \text{ in } Q, \quad (2.16)$$

where $\mathcal{R}_1 = \mathcal{R}_1(x, \partial)$ is some first order operator with bounded coefficients in Ω .

Arguing in a similar way, we find that

$$V'' - \mu_1 \lambda_1 \vec{\Delta}(V) + \mathcal{S}_1 V = \mathbf{g}' - \text{curl}(\lambda_1 \mathbf{f}), \text{ in } Q, \quad (2.17)$$

for another first order operator $\mathcal{S}_1 = \mathcal{S}_1(x, \partial)$ with bounded coefficients in Ω .

Therefore, putting (2.12) and (2.16)-(2.17) together, we end up getting that any solution $W = (U, V)$ to the Maxwell system (2.12) verifies

$$\begin{cases} U'' - \mu_1 \lambda_1 \vec{\Delta}(U) + \mathcal{R}_1 U = G_1, & \text{in } Q \\ U \times \nu = 0, \quad \text{curl}(\lambda_1 U) \cdot \nu = 0, & \text{on } \Sigma, \end{cases} \text{ and } \begin{cases} V'' - \mu_1 \lambda_1 \vec{\Delta}(V) + \mathcal{S}_1 V = G_2, & \text{in } Q \\ V \cdot \nu = 0, \quad \text{curl}(\mu_1 V) \times \nu = 0, & \text{on } \Sigma, \end{cases} \quad (2.18)$$

where $G_1 = \mathbf{f}' + \text{curl}(\mu_1 \mathbf{g})$ and $G_2 = \mathbf{g}' - \text{curl}(\lambda_1 \mathbf{f})$.

Further, consider a cut-off function $\eta \in C^\infty(\mathbb{R}; [0, 1])$ fulfilling

$$\eta(t) = \begin{cases} 1 & \text{if } |t| < T - 2\epsilon, \\ 0 & \text{if } |t| \geq T - \epsilon, \end{cases} \quad (2.19)$$

where ϵ is the same as in (2.9), and set

$$U_\# = \eta U, \quad V_\# = \eta V, \quad K_1 = \eta G_1 + 2\eta' U' + \eta'' U, \quad K_2 = \eta G_2 + 2\eta' V' + \eta'' V,$$

in such a way that we have

$$\begin{cases} U_\#'' - \mu_1 \lambda_1 \vec{\Delta}(U_\#) + \mathcal{R}_1 U_\# = K_1, & \text{in } Q \\ U_\# \times \nu = 0, \quad \text{curl}(\lambda_1 U_\#) \cdot \nu = 0, & \text{on } \Sigma \end{cases} \text{ and } \begin{cases} V_\#'' - \mu_1 \lambda_1 \vec{\Delta}(V_\#) + \mathcal{S}_1 V_\# = K_2, & \text{in } Q \\ V_\# \cdot \nu = 0, \quad \text{curl}(\mu_1 V_\#) \times \nu = 0, & \text{on } \Sigma, \end{cases}$$

directly from (2.18). Then, each of the two above systems being a principally scalar hyperbolic system, it follows from the two identities

$$U_\#(\cdot, \pm T) = U_\#',(\cdot, \pm T) = 0, \quad V_\#(\cdot, \pm T) = V_\#',(\cdot, \pm T) = 0, \quad (2.20)$$

and Theorem 2, that $W_\# = (U_\#, V_\#)$ obeys the Carleman estimate

$$\begin{aligned} & C_0 \int_Q e^{2s\varphi} s \left(|\nabla_{x,t} W_\#|^2 + s^2 |W_\#|^2 \right) dx dt \\ & \leq \sum_{j=1,2} \int_Q e^{2s\varphi} |K_j(x, t)|^2 dx dt + \int_\Sigma s e^{2s\varphi} \left(|\nabla W_\#|^2 + |W_\#'|^2 + s^2 |W_\#|^2 \right) d\sigma dt, \end{aligned} \quad (2.21)$$

for all $s \geq s_0$. Here we have used Theorem 2 for the diagonal system $U_{\#}'' - \mu_1 \lambda_1 \vec{\Delta}(U_{\#})$ and that we can absorb the non-decoupled first term $\mathcal{R}_1 U_{\#}$.

Moreover, as η' and η'' both vanish in $(-T + 2\epsilon, T - 2\epsilon)$ by (2.14), there is a constant $C > 0$ such that

$$\sum_{j=1,2} \int_Q e^{2s\varphi} |K_j(x,t)|^2 dxdt \leq C \int_Q e^{2s\varphi} (|\mathbf{h}|^2 + |\nabla_{x,t}\mathbf{h}|^2) dxdt + e^{2d_0s} \|W\|_{H^1(Q)}^2,$$

according to (2.20)-(2.21) and since

$$\int_Q e^{2s\varphi} s (|\nabla_{x,t}W|^2 + s^2 |W|^2) dxdt \leq C \int_Q e^{2s\varphi} s (|\nabla_{x,t}W_{\#}|^2 + s^2 |W_{\#}|^2) dxdt + s^3 e^{2d_0s} \|W\|_{H^1(Q)}^2$$

we obtain the result. \square

2.3 Reduction of the boundary terms

The method used to derive a global Carleman estimate for the solution to (1.1) is to replace the local boundary problem (1.1) in $\Omega \times (-T, T)$ by an equivalent one stated on the half space $\mathbb{R}_+^3 \times (-T, T)$. This is possible since the boundary Γ can be represented as the zero level set of some C^∞ function in \mathbb{R}^3 . Namely, Γ being a C^∞ surface, there exist $\theta \in C^\infty(\mathbb{R}^3)$ and some neighbourhood \mathcal{V} of Γ in \mathbb{R}^3 such that $\Gamma = \{x \in \mathcal{V}, \theta(x) = 0\}$. We choose \mathcal{V} so small that $\mathcal{V} \subset \mathcal{O}$, where \mathcal{O} is defined in §1.2, write $y = (y_1, y_2, y_3) = (y', y_3)$ the system of normal geodesic coordinates where $y' = (y_1, y_2)$ are orthogonal coordinates in Γ and $y_3 = \theta(x)$ is the normal coordinate, and call $x = \Phi(y)$, where $\Phi'(y) > 0$ for all $y \in \hat{\mathcal{V}} := \Phi^{-1}(\mathcal{V})$, the corresponding coordinates mapping. As

$$\hat{\Gamma} := \Phi^{-1}(\Gamma) = \{y \in \hat{\mathcal{V}}; y_3 = 0\} \subset \mathbb{R}^2,$$

we may assume that $\hat{\mathcal{V}} = \Phi^{-1}(\mathcal{V})$ is a cylinder of the form $\hat{\Gamma} \times (-r, r)$ with $r > 0$.

Further, the Euclidean metric in \mathbb{R}^3 inducing the Riemannian metric with diagonal tensor g ,

$$g(y) = {}^t\Phi'(y)\Phi'(y) = \text{Diag}(g_1, g_2, g_3), \quad y \in \hat{\mathcal{V}},$$

we use the notations of [36] and note $\left[\frac{1}{\sqrt{g_1}}\frac{\partial}{\partial y_1}, \frac{1}{\sqrt{g_2}}\frac{\partial}{\partial y_2}, \frac{1}{\sqrt{g_3}}\frac{\partial}{\partial y_3}\right]$ the orthonormal basis associated by g to the differential basis of vector fields $\left[\frac{\partial}{\partial y_1}, \frac{\partial}{\partial y_2}, \frac{\partial}{\partial y_3}\right]$. For any vector field $X(x)$ expressed with respect to the Euclidian basis $\left[\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2}, \frac{\partial}{\partial x_3}\right]$ as $X(x) = \sum_{i=1}^3 \alpha^i(x) \frac{\partial}{\partial x_i}$, we have an alternative representation $\hat{X}(y)$ with respect to the new basis vectors $\left[\frac{1}{\sqrt{g_1}}\frac{\partial}{\partial y_1}, \frac{1}{\sqrt{g_2}}\frac{\partial}{\partial y_2}, \frac{1}{\sqrt{g_3}}\frac{\partial}{\partial y_3}\right]$, given by

$$\hat{X}(y) = \sum_{i=1}^3 \hat{\alpha}^i(y) \frac{1}{\sqrt{g_i}} \frac{\partial}{\partial y_i}, \quad \hat{\alpha}(y) = {}^t\Psi(y)\alpha(\Phi(y)), \quad \Psi(y) = \Phi'(y)g^{-1/2}(y), \quad y \in \hat{\mathcal{V}}. \quad (2.22)$$

The divergence (resp. curl) operator of any vector field $\hat{X}(y) = \sum_{i=1}^3 \hat{\alpha}^i(y) \frac{1}{\sqrt{g_i}} \frac{\partial}{\partial y_i}$ with respect to the local coordinates (y_1, y_2, y_3) is denoted by $\text{div}_g \hat{X}$ (resp. $\text{curl}_g \hat{X}$), and can be brought into the form

$$\text{div}_g \hat{X} = \frac{1}{\sqrt{\det g}} \sum_{j=1}^3 \frac{1}{\sqrt{g_j}} \frac{\partial}{\partial y_j} \left(\sqrt{\det g} \hat{\alpha}^j(y) \right) = \sum_{j=1}^3 \frac{1}{\sqrt{g_j}} \frac{\partial \hat{\alpha}^j}{\partial y_j} + A(y) \cdot \hat{X}(y), \quad (2.23)$$

where $A(y)$ is some three dimensional vector (resp.

$$\operatorname{curl}_g \widehat{X} = \frac{1}{2} \sum_{i,j=1}^3 \left(\frac{1}{\sqrt{g_j}} \frac{\partial \widehat{\alpha}^i}{\partial y_j} - \frac{1}{\sqrt{g_i}} \frac{\partial \widehat{\alpha}^j}{\partial y_i} \right) \frac{1}{\sqrt{g_j}} \frac{\partial}{\partial y_j} \times \frac{1}{\sqrt{g_i}} \frac{\partial}{\partial y_i} + M(y) \widehat{X}(y), \quad (2.24)$$

where $M(y)$ is some matrix function), whereas the outward normal vector field to $\widehat{\Gamma}$ at $y \in \widehat{\Gamma}$ is given by

$$\widehat{\nu}(y) = -\frac{1}{\sqrt{g_3}} \frac{\partial}{\partial y_3}. \quad (2.25)$$

In light of (2.22)-(2.25), we find out by performing the change of variable $x = \Phi(y)$ in (2.12), the space variable x being restricted to be in $\Omega \cap \mathcal{V} \subset \omega \cap \mathcal{V}$, that

$$\begin{aligned} \widehat{U}' - \operatorname{curl}_g(\widehat{\mu}_1 \widehat{V}) &= 0, & \text{in } \widehat{\mathcal{V}} \times (-T, T), \\ \widehat{V}' + \operatorname{curl}_g(\widehat{\lambda}_1 \widehat{U}) &= 0, & \text{in } \widehat{\mathcal{V}} \times (-T, T), \\ \operatorname{div}_g \widehat{U} = \operatorname{div}_g \widehat{V} &= 0, & \text{in } \widehat{\mathcal{V}} \times (-T, T), \\ \widehat{U} \times \widehat{\nu} = 0, \quad \widehat{V} \cdot \widehat{\nu} &= 0, & \text{on } \widehat{\Gamma} \times (-T, T), \end{aligned} \quad (2.26)$$

where we have set

$$\widehat{\mu}_1(y) = {}^t \Psi(y) \mu_1(\Phi(y)) \Psi(y), \quad \widehat{\lambda}_1(y) = {}^t \Psi(y) \lambda_1(\Phi(y)) \Psi(y), \quad y \in \widehat{\mathcal{V}} \subset \Phi^{-1}(\omega).$$

Here we used the identity $\mathbf{h}(\Phi(y), t) = (\mathbf{f}(\Phi(y), t), \mathbf{g}(\Phi(y), t)) = 0$ for $t \in (-T, T)$ and $y \in \widehat{\mathcal{V}}$, arising from (2.13). Further, noting $\widehat{U} = (\widehat{u}_1, \widehat{u}_2, \widehat{u}_3)$ and $\widehat{V} = (\widehat{v}_1, \widehat{v}_2, \widehat{v}_3)$, the last equation in (2.26) reads $\widehat{u}_1 = \widehat{u}_2 = \widehat{v}_3 = 0$ on $\widehat{\Gamma}$, so we find that

$$\widehat{U}_{\widehat{\tau}} = 0, \quad \widehat{V}_{\widehat{\nu}} = 0, \quad \text{on } \widehat{\Gamma}, \quad (2.27)$$

where $\widehat{U}_{\widehat{\tau}}$ (resp. $\widehat{V}_{\widehat{\nu}}$) denotes the tangential (resp. normal) component of \widehat{U} (resp. \widehat{V}). From this, (2.23) and the third line in (2.26) then follows that

$$\frac{1}{\sqrt{g_3}} \frac{\partial \widehat{u}_3}{\partial y_3} = - \left(\frac{1}{\sqrt{g_1}} \frac{\partial \widehat{u}_1}{\partial y_1} + \frac{1}{\sqrt{g_2}} \frac{\partial \widehat{u}_2}{\partial y_2} \right) - A(y) \cdot \widehat{U} = -A(y) \cdot \widehat{U},$$

whence

$$\nabla \widehat{u}_3 = \frac{1}{\sqrt{g_1}} \frac{\partial \widehat{u}_3}{\partial y_1} \widehat{e}_1 + \frac{1}{\sqrt{g_2}} \frac{\partial \widehat{u}_3}{\partial y_2} \widehat{e}_2 - (A(y) \cdot \widehat{U}) \widehat{e}_3, \quad \text{on } \widehat{\Gamma}. \quad (2.28)$$

Further, as $\nabla \widehat{u}_1 = \frac{1}{\sqrt{g_3}} \frac{\partial \widehat{u}_1}{\partial y_3} \widehat{e}_3$ and $\nabla \widehat{u}_2 = \frac{1}{\sqrt{g_3}} \frac{\partial \widehat{u}_2}{\partial y_3} \widehat{e}_3$ on $\widehat{\Gamma}$, according to (2.27), (2.28) then yields

$$|\nabla \widehat{U}|^2 = \sum_{j=1}^3 |\nabla \widehat{u}_j|^2 \leq C \left(\left| \frac{\partial \widehat{u}_1}{\partial y_3} \right|^2 + \left| \frac{\partial \widehat{u}_2}{\partial y_3} \right|^2 + \left| \frac{\partial \widehat{u}_3}{\partial y_1} \right|^2 + \left| \frac{\partial \widehat{u}_3}{\partial y_2} \right|^2 + |\widehat{U}_{\widehat{\nu}}|^2 \right), \quad \text{on } \widehat{\Gamma}, \quad (2.29)$$

where, for the sake of notational simplicity, we shall use the generic constant $C > 0$ in the remaining of §2.3. On the other hand, since

$$\operatorname{curl}_g \widehat{U} \times \widehat{\nu} = \left(\frac{1}{\sqrt{g_3}} \frac{\partial \widehat{u}_1}{\partial y_3} - \frac{1}{\sqrt{g_1}} \frac{\partial \widehat{u}_3}{\partial y_1} \right) \frac{1}{\sqrt{g_1 g_3}} \frac{\partial}{\partial y_1} - \left(\frac{1}{\sqrt{g_2}} \frac{\partial \widehat{u}_3}{\partial y_2} - \frac{1}{\sqrt{g_3}} \frac{\partial \widehat{u}_2}{\partial y_3} \right) \frac{1}{\sqrt{g_2 g_3}} \frac{\partial}{\partial y_2} + M(y) \widehat{U}_{\widehat{\nu}}, \quad (2.30)$$

by (2.24)-(2.25), we have

$$\left| \frac{\partial \hat{u}_1}{\partial y_3} \right|^2 + \left| \frac{\partial \hat{u}_2}{\partial y_3} \right|^2 \leq C \left(\left| \operatorname{curl}_g \hat{U} \right|^2 + \left| \hat{U}_{\hat{\nu}} \right|^2 + \left| \frac{\partial \hat{u}_3}{\partial y_1} \right|^2 + \left| \frac{\partial \hat{u}_3}{\partial y_2} \right|^2 \right), \text{ on } \hat{\Gamma}.$$

In view of (2.29), this entails

$$\begin{aligned} \left| \nabla \hat{U} \right|^2 &\leq C \left(\left| \operatorname{curl}_g \hat{U} \right|^2 + \left| \frac{\partial \hat{u}_3}{\partial y_1} \right|^2 + \left| \frac{\partial \hat{u}_3}{\partial y_2} \right|^2 + \left| \hat{U}_{\hat{\nu}} \right|^2 \right) \\ &\leq C \left(\left| \operatorname{curl}_g \hat{U} \right|^2 + \left| \nabla_{\hat{\tau}} \hat{U}_{\hat{\nu}} \right|^2 + \left| \hat{U}_{\hat{\nu}} \right|^2 \right), \text{ on } \hat{\Gamma}. \end{aligned}$$

As a consequence we have

$$\left| \nabla \hat{U} \right|^2 \leq C \left(\left| \operatorname{curl}_g(\hat{\lambda}_1 \hat{U}) \right|^2 + \left| \nabla_{\hat{\tau}} \hat{U}_{\hat{\nu}} \right|^2 + \left| \hat{U}_{\hat{\nu}} \right|^2 \right),$$

whence

$$\left| \nabla \hat{U} \right|^2 \leq C \left(\left| \hat{V}' \right|^2 + \left| \nabla_{\hat{\tau}} \hat{U}_{\hat{\nu}} \right|^2 + \left| \hat{U}_{\hat{\nu}} \right|^2 \right), \text{ on } \hat{\Gamma}, \quad (2.31)$$

by the second line of (2.26).

Similarly, as $\operatorname{div}_g \hat{V} = 0$ in \hat{V} from the third line of (2.26), we get from (2.23) that

$$\frac{1}{\sqrt{g_3}} \frac{\partial \hat{v}_3}{\partial y_3} = - \left(\frac{1}{\sqrt{g_1}} \frac{\partial \hat{v}_1}{\partial y_1} + \frac{1}{\sqrt{g_2}} \frac{\partial \hat{v}_2}{\partial y_2} \right) - A(y) \cdot \hat{V}, \text{ on } \hat{\Gamma}.$$

This, combined with (2.27), yields

$$\left| \nabla \hat{v}_3 \right|^2 \leq C \left(\left| \frac{\partial \hat{v}_1}{\partial y_1} \right|^2 + \left| \frac{\partial \hat{v}_2}{\partial y_2} \right|^2 + \left| \hat{V}_{\hat{\tau}} \right|^2 \right),$$

and consequently

$$\left| \nabla \hat{V} \right|^2 = \sum_{j=1}^3 \left| \nabla \hat{v}_j \right|^2 \leq C \left(\left| \nabla_{\hat{\tau}} \hat{V}_{\hat{\tau}} \right|^2 + \left| \hat{V}_{\hat{\tau}} \right|^2 + \left| \frac{\partial \hat{v}_1}{\partial y_3} \right|^2 + \left| \frac{\partial \hat{v}_2}{\partial y_3} \right|^2 \right), \text{ on } \hat{\Gamma}. \quad (2.32)$$

Furthermore, in light of (2.27) and (2.30) where \hat{V} (resp. \hat{v}_j , $j = 1, 2, 3$) is substituted for \hat{U} (resp. \hat{u}_j , $j = 1, 2, 3$), we see that $\left| \frac{\partial \hat{v}_1}{\partial y_3} \right|^2 + \left| \frac{\partial \hat{v}_2}{\partial y_3} \right|^2$ is upper bounded, up to some multiplicative constant, by $\left| \operatorname{curl}_g \hat{V} \right|^2 + \left| \hat{V}_{\hat{\tau}} \right|^2$, and hence by $\left| \operatorname{curl}_g(\hat{\mu}_1 \hat{V}) \right|^2 + \left| \hat{V}_{\hat{\tau}} \right|^2$, on $\hat{\Gamma}$. From this and the first line of (2.26) then follows that

$$\left| \frac{\partial \hat{v}_1}{\partial y_3} \right|^2 + \left| \frac{\partial \hat{v}_2}{\partial y_3} \right|^2 \leq C \left(\left| \hat{U}' \right|^2 + \left| \hat{V}_{\hat{\tau}} \right|^2 \right), \text{ on } \hat{\Gamma},$$

so, we end up getting with the aid of (2.32):

$$\left| \nabla \hat{V} \right|^2 \leq C \left(\left| \hat{U}' \right|^2 + \left| \nabla_{\hat{\tau}} \hat{V}_{\hat{\tau}} \right|^2 + \left| \hat{V}_{\hat{\tau}} \right|^2 \right) \text{ on } \hat{\Gamma}. \quad (2.33)$$

Finally, putting (2.15), (2.31) and (2.33) together, we may state the main result of §2.3:

Lemma 2.2 *Assume (A1)-(A2) and put $\mathbf{h} = (\mathbf{f}, \mathbf{g})$. Then there are two constants $C_2 > 0$ and $s_2 > 0$ such that the following Carleman estimate*

$$\begin{aligned} & C_2 s \int_Q e^{2s\varphi} \left(s^2 |\mathbf{W}|^2 + |\nabla_{x,t} \mathbf{W}|^2 \right) dx dt \\ & \leq \int_Q e^{2s\varphi} \left(|\mathbf{h}|^2 + |\nabla_{x,t} \mathbf{h}|^2 \right) dx dt + \mathcal{B}_{s,\varphi}(\mathbf{W}) + s^3 e^{2d_0 s} \|\mathbf{W}\|_{H^1(Q)}^2, \end{aligned}$$

where

$$\mathcal{B}_{s,\varphi}(\mathbf{W}) = \int_{\Sigma} s e^{2s\varphi} \left(|\nabla_{\tau} V_{\tau}|^2 + |\nabla_{\tau} U_{\nu}|^2 + |U'_{\nu}|^2 + |V'_{\tau}|^2 + s^2 (|U_{\nu}|^2 + |V_{\tau}|^2) \right) d\sigma dt, \quad (2.34)$$

holds true for every solution $\mathbf{W} = (U, V)$ to (2.12), provided $s \geq s_2$.

3 Inverse problem

This section contains the proof of Theorem 1, which is divided into five steps. Firstly, the unknown parameters λ and μ are brought to the source term of the linearized system associated to (1.1), governing the variation induced on the solution to (1.1) by perturbing the permittivity by λ and the permeability by μ . The second step follows the idea of Bukhgeim and Klivanov presented in [10], which is to differentiate the linearized system with respect to t in order to move the unknown coefficients in the initial condition. The next step is to bound the energy of this system at time $t = 0$ with the aid of the Carleman inequality of Theorem 2. The fourth step involves relating λ and μ to the above mentioned estimate through the Carleman inequality for stationary (div, curl)-systems, stated in Lemma 3.1. This is rather technical and lengthy so we proceed in a succession of the two Lemmas 3.3 and 3.4. The last step, detailed in §3.3, is to derive the desired result from the estimates established in Lemmas 3.3-3.4.

In the remaining of this text, x_0 is a fixed point in $\mathbb{R}^3 \setminus \overline{\Omega}$, we choose as in (1.7) $\psi_0(x) := |x - x_0|^2$ for every $x \in \overline{\Omega}$, and $\varphi_0(x) := \varphi(x, 0)$, where φ denotes the function defined by (2.7) and (2.10). Moreover, for the sake of notational simplicity, we shall use the generic constant $C > 0$ in the various estimates of §3.2-3.3.

3.1 Linearized inverse problem

Given $(\mu_i, \lambda_i) \in \Lambda_{\omega}(M_0)$, $i = 1, 2$, and $(B_0^k, D_0^k) \in H^2(\Omega)^3 \times H^2(\Omega)^3$, $k = 1, 2$, we consider the solution (B_i^k, D_i^k) to the system (1.1) where (λ_i, μ_i) is substituted for (λ, μ) , with initial condition (1.2) where $(B_0, D_0) = (B_0^k, D_0^k)$. Hence, putting

$$\mu = \mu_1 - \mu_2, \quad \lambda = \lambda_1 - \lambda_2,$$

and setting

$$\mathbf{f}_k = \text{curl}(\mu B_2^k), \quad \mathbf{g}_k = -\text{curl}(\lambda D_2^k), \quad (3.1)$$

we find by a straightforward computation that $U_k = D_1^k - D_2^k$ and $V_k = B_1^k - B_2^k$ satisfy the system

$$\begin{aligned} U'_k - \text{curl}(\mu_1 V_k) &= \mathbf{f}_k, & \text{in } Q, \\ V'_k + \text{curl}(\lambda_1 U_k) &= \mathbf{g}_k, & \text{in } Q, \\ \text{div } U_k &= \text{div } V_k = 0, & \text{in } Q, \\ U_k \times \nu &= 0, \quad V_k \cdot \nu = 0, & \text{on } \Sigma, \end{aligned} \quad (3.2)$$

with the initial data

$$U_k(x, 0) = 0, \quad V_k(x, 0) = 0. \quad (3.3)$$

Further, by using the following notations

$$X_{k,j}(x, t) = \partial_t^j X_k(x, t) \text{ for } X = U, V, \mathbf{f}, \mathbf{g} \text{ and } j \in \mathbb{N}^*, \quad (3.4)$$

it turns out by differentiating (3.2) j -times with respect to t that

$$\begin{aligned} U'_{k,j} - \operatorname{curl}(\mu_1 V_{k,j}) &= \mathbf{f}_{k,j}, & \text{in } Q, \\ V'_{k,j} + \operatorname{curl}(\lambda_1 U_{k,j}) &= \mathbf{g}_{k,j}, & \text{in } Q, \\ \operatorname{div} U_{k,j} = \operatorname{div} V_{k,j} &= 0, & \text{in } Q, \\ U_{k,j} \times \nu = 0, \quad V_{k,j} \cdot \nu &= 0, & \text{on } \Sigma, \end{aligned} \quad (3.5)$$

and, due to (3.1)-(3.2), that $U_{k,1}$ and $V_{k,1}$ satisfy the initial condition:

$$U_{k,1}(x, 0) = \operatorname{curl}(\mu B_0^k), \quad V_{k,1}(x, 0) = -\operatorname{curl}(\lambda D_0^k). \quad (3.6)$$

As will appear in §3.2-3.3, the main benefit of dealing with (3.1)-(3.6) in the analysis of the inverse problem of determining λ and μ , is the presence of these two unknown coefficients in the initial condition (3.6).

3.2 Preliminary estimates

Let j and k be in $\{1, 2\}$. As $W_{k,j} = (U_{k,j}, V_{k,j})$ is solution to (3.5), we notice from Lemma 2.2 that

$$\begin{aligned} & C_2 s \int_Q e^{2s\varphi} \left(s^2 |W_{k,j}|^2 + |\nabla_{x,t} W_{k,j}|^2 \right) dx dt \\ & \leq \int_Q e^{2s\varphi} \left(|\mathbf{h}_{k,j}|^2 + |\nabla_{x,t} \mathbf{h}_{k,j}|^2 \right) dx dt + \mathcal{B}_{s,\varphi}(W_{k,j}) + s^3 e^{2d_0 s} \|W_{k,j}\|_{H^1(Q)}^2 := \mathfrak{I}_{k,j}(s), \end{aligned} \quad (3.7)$$

for every $s \geq s_2$, where $\mathbf{h}_{k,j} = (\mathbf{f}_{k,j}, \mathbf{g}_{k,j})$ and $\mathcal{B}_{s,\varphi}$ is given by (2.34).

Moreover by the assumption (1.11) we can derive that

$$\sum_{k=1}^2 |B_0^k(x)|^2 \geq c_*, \quad \text{and} \quad \sum_{k=1}^2 |D_0^k(x)|^2 \geq c_*, \quad x \in \overline{\Omega \setminus \omega} \quad (3.8)$$

for some positive constant c_* . Indeed, if $B_0^1(x) \neq 0$ or $B_0^2(x) \neq 0$ for all $x \in \overline{\Omega \setminus \omega}$ then $\sum_{k=1}^2 |B_0^k(x)|^2 > 0$ in the compact set $\overline{\Omega \setminus \omega}$. Now if there exist $x_1 \in \overline{\Omega \setminus \omega}$ such that $B_0^1(x_1) = 0$ then by (1.11) we have $B_0^2(x) \neq 0$ for any $x \in \overline{\Omega \setminus \omega}$.

Further we recall from [36] the following Carleman estimate for stationary (div, curl)-systems:

Lemma 3.1 *There exist two positive constants s_3 and C_3 depending only on ψ_0 and Ω , such that we have*

$$C_3 s \int_{\Omega} e^{2s\varphi_0} |\mathbf{u}|^2 dx \leq \int_{\Omega} e^{2s\varphi_0} \left(|\operatorname{curl} \mathbf{u}|^2 + |\operatorname{div} \mathbf{u}|^2 \right) dx, \quad (3.9)$$

for every $s \geq s_3$ and $\mathbf{u} \in H_0^1(\Omega)$.

Prior to the proof of Lemma 3.3, we establish the following technical result, needed in the derivation of (3.10)-(3.11).

Lemma 3.2 *There exists a constant $s_* > 0$ depending only on T such that we have*

$$\int_{\Omega} |z(x, 0)|^2 dx \leq 2 \left(s \int_Q |z(x, t)|^2 dx dt + s^{-1} \int_Q |z'(x, t)|^2 dx dt \right),$$

for all $s \geq s_*$ and $z \in H^1(-T, T; L^2(\Omega))$.

Proof . Let $\eta \in C^\infty([-T, T]; [0, 1])$ fulfills (2.19) for some fixed $\epsilon \in (0, T/2)$. Then, the following identity

$$\begin{aligned} \int_{\Omega} |z(x, 0)|^2 dx &= \int_{-T}^0 \frac{d}{dt} \left(\int_{\Omega} \eta^2(t) |z(x, t)|^2 dx \right) dt \\ &= 2\Re \left(\int_{-T}^0 \int_{\Omega} \eta^2(t) \overline{z(x, t)} z'(x, t) dx dt \right) + 2 \int_{-T}^0 \int_{\Omega} \eta(t) \eta'(t) |z(x, t)|^2 dx dt, \end{aligned}$$

holds true for every $z \in H^1(-T, T; L^2(\Omega))$. Applying Young's inequality, this entails

$$\int_{\Omega} |z(x, 0)|^2 dx \leq (s + 2\|\eta'\|_\infty) \int_Q |z(x, t)|^2 dx dt + s^{-1} \int_Q |z'(x, t)|^2 dx dt,$$

for each $s > 0$, so the result follows by taking $s_* = 2\|\eta'\|_\infty$. \square

Having said that, we are now in position to prove the:

Lemma 3.3 *There exist two constants $C_4 > 0$ and $s_4 > 0$ such that the following estimates*

$$s^2 \left(C_4 s \int_{\Omega} e^{2s\varphi_0} (|\mu|^2 + |\lambda|^2) dx - \int_{\Omega} e^{2s\varphi_0} (|\nabla\mu|^2 + |\nabla\lambda|^2) dx \right) \leq \sum_{k=1}^2 \mathfrak{Z}_{k,1}(s), \quad (3.10)$$

and

$$C_4 s \int_{\Omega} e^{2s\varphi_0} (|\nabla\mu|^2 + |\nabla\lambda|^2) dx - \sum_{|\alpha|=2} \int_{\Omega} e^{2s\varphi_0} (|\partial^\alpha\mu|^2 + |\partial^\alpha\lambda|^2) dx \leq \sum_{j,k=1}^2 \mathfrak{Z}_{k,j}(s), \quad (3.11)$$

hold true for $k = 1, 2$, and $s \geq s_4$.

Proof . By applying Lemma 3.2 for $z = e^{s\varphi} U_{k,1}$, we get that

$$C s^2 \int_{\Omega} e^{2s\varphi_0} |U_{k,1}(x, 0)|^2 dx \leq s \int_Q e^{2s\varphi} \left(s^2 |U_{k,1}(x, t)|^2 + |U_{k,2}(x, t)|^2 \right) dx dt,$$

provided s is large enough. In light of (3.6)-(3.7), this entails

$$C s^2 \int_{\Omega} e^{2s\varphi_0} \left| \operatorname{curl} (\mu B_0^k) \right|^2 dx \leq \mathfrak{Z}_{k,1}(s). \quad (3.12)$$

Further, taking into account that $\mu B_0^k \in H_0^1(\Omega)^3$ since μ vanishes in ω and $\operatorname{div} B_0^k = 0$, we have

$$C s^3 \int_{\Omega} e^{2s\varphi_0} \left| \mu B_0^k \right|^2 dx - s^2 \int_{\Omega} e^{2s\varphi_0} |\nabla\mu|^2 dx \leq s^2 \int_{\Omega} e^{2s\varphi_0} \left| \operatorname{curl} (\mu B_0^k) \right|^2 dx,$$

by (3.9), whence

$$Cs^3 \int_{\Omega} e^{2s\varphi_0} \left| \mu B_0^k \right|^2 dx - s^2 \int_{\Omega} e^{2s\varphi_0} |\nabla \mu|^2 dx \leq \mathfrak{Z}_{k,1}(s), \quad (3.13)$$

from (3.12). Similarly, by arguing as above with $z = e^{s\varphi} V_{k,1}$ instead of $e^{s\varphi} U_{k,1}$, we find some constant $C > 0$ for which $Cs^3 \int_{\Omega} e^{2s\varphi_0} \left| \lambda D_0^k \right|^2 dx - s^2 \int_{\Omega} e^{2s\varphi_0} |\nabla \lambda|^2 dx$ can be made smaller than the right hand side of (3.13) by taking s sufficiently large. This, (3.8) and (3.13) entails (3.10).

We turn now to showing (3.11). To do that we apply Lemma 3.2 with $z = e^{s\varphi} \partial_i U_{k,1}$, $i = 1, 2, 3$, getting

$$C \int_{\Omega} e^{2s\varphi_0} |\partial_i U_{k,1}(x, 0)|^2 dx \leq s \int_Q e^{2s\varphi} \left(|\nabla U_{k,1}(x, t)|^2 + s^{-2} |\nabla U_{k,2}(x, t)|^2 \right) dx dt,$$

for s large enough. This yields

$$C \int_{\Omega} e^{2s\varphi_0} \left| \partial_i \operatorname{curl} (\mu B_0^k) \right|^2 dx \leq \sum_{j=1}^2 \mathfrak{Z}_{k,j}(s), \quad (3.14)$$

by (3.6)-(3.7). Further, bearing in mind that $\operatorname{div} B_0^k = 0$ and using that $(\partial_i \mu) B_0^k \in H_0^1(\Omega)^3$, we obtain

$$Cs \int_{\Omega} e^{2s\varphi_0} \left| (\partial_i \mu) B_0^k \right|^2 dx - \sum_{|\alpha|=2} \int_{\Omega} e^{2s\varphi_0} |\partial^\alpha \mu|^2 dx \leq \int_{\Omega} e^{2s\varphi_0} \left| \operatorname{curl} ((\partial_i \mu) B_0^k) \right|^2 dx, \quad (3.15)$$

by (3.9). Moreover, as $\operatorname{curl} ((\partial_i \mu) B_0^k) = \partial_i \operatorname{curl} (\mu B_0^k) - \mu \operatorname{curl} (\partial_i B_0^k) - \nabla \mu \times \partial_i B_0^k$ and $\mu \in H_0^1(\Omega)$, we have by applying the Poincaré inequality

$$C \int_{\Omega} e^{2s\varphi_0} \left| \operatorname{curl} ((\partial_i \mu) B_0^k) \right|^2 dx \leq \int_{\Omega} e^{2s\varphi_0} \left| \partial_i \operatorname{curl} (\mu B_0^k) \right|^2 dx + \int_{\Omega} e^{2s\varphi_0} |\nabla \mu|^2 dx, \quad (3.16)$$

hence

$$Cs \int_{\Omega} e^{2s\varphi_0} \left| (\partial_i \mu) B_0^k \right|^2 dx - \sum_{1 \leq |\alpha| \leq 2} \int_{\Omega} e^{2s\varphi_0} |\partial^\alpha \mu|^2 dx \leq \int_{\Omega} e^{2s\varphi_0} \left| \partial_i \operatorname{curl} (\mu B_0^k) \right|^2 dx, \quad (3.17)$$

by substituting the right hand side of (3.16) for $\int_{\Omega} e^{2s\varphi_0} \left| \operatorname{curl} ((\partial_i \mu) B_0^k) \right|^2 dx$ in (3.15). Putting (3.14) and (3.17) together, and summing up the obtained inequality over $i = 1, 2, 3$, we end up getting that

$$Cs \int_{\Omega} e^{2s\varphi_0} |\nabla \mu|^2 dx - \sum_{1 \leq |\alpha| \leq 2} \int_{\Omega} e^{2s\varphi_0} |\partial^\alpha \mu|^2 dx \leq \sum_{j,k=1}^2 \mathfrak{Z}_{k,j}(s). \quad (3.18)$$

Here we used again (3.8). Finally, by arguing as before with $z = e^{s\varphi} \partial_i V_{k,1}$, $i = 1, 2, 3$, instead of $\partial_i U_{k,1}$, we get that (3.18) remains true with μ replaced by λ . This completes the proof of (3.11). \square

Finally, we establish the:

Lemma 3.4 *There are two constants $C_5 > 0$ and $s_5 > 0$ such that we have*

$$C_5 \sum_{|\alpha|=2} \int_{\Omega} e^{2s\varphi_0} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx - \int_{\Omega} e^{2s\varphi_0} \left(|\nabla \mu|^2 + |\nabla \lambda|^2 \right) dx \leq \sum_{j,k=1}^2 \mathfrak{Z}_{k,j}(s),$$

for all $s \geq s_5$.

Proof . In light of the two following basic identities $U_{k,1}(x, 0) = \mu \operatorname{curl} B_0^k + \nabla \mu \times B_0^k$ and $V_{k,1}(x, 0) = -\mu \operatorname{curl} D_0^k - \nabla \lambda \times D_0^k$, $k = 1, 2$, arising from (3.6), we have

$$\mathcal{K}(x) \begin{pmatrix} \nabla \mu \\ \nabla \lambda \end{pmatrix} = \mathcal{A}(x) \begin{pmatrix} \mu \\ \lambda \end{pmatrix} - \begin{pmatrix} U_{1,1}(x, 0) \\ -V_{1,1}(x, 0) \\ U_{2,1}(x, 0) \\ -V_{2,1}(x, 0) \end{pmatrix}, \text{ with } \mathcal{A}(x) = \begin{pmatrix} \operatorname{curl} B_0^1 & 0 \\ 0 & \operatorname{curl} D_0^1 \\ \operatorname{curl} B_0^2 & 0 \\ 0 & \operatorname{curl} D_0^2 \end{pmatrix},$$

hence

$$\mathcal{K}(x) \begin{pmatrix} \nabla \partial_i \mu \\ \nabla \partial_i \lambda \end{pmatrix} = \partial_i \mathcal{A} \begin{pmatrix} \mu \\ \lambda \end{pmatrix} + \mathcal{A} \begin{pmatrix} \partial_i \mu \\ \partial_i \lambda \end{pmatrix} - \partial_i \mathcal{K} \begin{pmatrix} \nabla \mu \\ \nabla \lambda \end{pmatrix} - \begin{pmatrix} \partial_i U_{1,1}(x, 0) \\ -\partial_i V_{1,1}(x, 0) \\ \partial_i U_{2,1}(x, 0) \\ -\partial_i V_{2,1}(x, 0) \end{pmatrix},$$

for every $i = 1, 2, 3$. From this and (1.11) then follows that

$$\sum_{|\alpha|=2} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) \leq C \left(\sum_{k=1}^2 \left(|\nabla U_{k,1}(x, 0)|^2 + |\nabla V_{k,1}(x, 0)|^2 \right) + \sum_{|\alpha| \leq 1} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) \right). \quad (3.19)$$

Further, by multiplying (3.19) by $e^{s\varphi_0}$, integrating over Ω , and upper bounding $\int_\Omega e^{2s\varphi_0} |\nabla W_{k,1}(x, 0)|^2 dx$, with the aid of Lemma 3.2, we find out that

$$\begin{aligned} & C \sum_{|\alpha|=2} \int_\Omega e^{2s\varphi_0} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx - \int_\Omega e^{2s\varphi_0} \left(|\nabla \mu|^2 + |\nabla \lambda|^2 \right) dx \\ & \leq s \sum_{k=1,2} \left(\int_Q e^{2s\varphi} |\nabla W_{k,1}(x, t)|^2 dx dt + s^{-2} \int_Q e^{2s\varphi} |\nabla W_{k,2}(x, t)|^2 dx dt \right). \end{aligned} \quad (3.20)$$

Here we took advantage of the fact that both μ and λ belong to $H_0^1(\Omega)$ in order to get rid of the integral $\int_\Omega e^{2s\varphi_0} \left(|\mu|^2 + |\lambda|^2 \right) dx$ by applying the Poincaré inequality. Evidently the result now follows from (3.20) and Lemma 2.2. \square

3.3 Completion of the proof of the main result

In light of (3.11) and Lemma 3.4 we may find $C > 0$ such that

$$C \sum_{|\alpha| \leq 2} \int_\Omega e^{2s\varphi_0} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx \leq \sum_{j,k=1}^2 \mathfrak{Z}_{k,j}(s), \quad (3.21)$$

upon taking s sufficiently large. Moreover, due to (3.1), we have

$$\begin{aligned} & \int_Q e^{2s\varphi} \left(|\mathbf{h}_{k,j}|^2 + |\nabla \mathbf{h}_{k,j}|^2 \right) dx dt \\ & \leq \left\| (B_2^k, D_2^k) \right\|_{C^3(-T, T; W^{2, \infty}(\Omega))}^2 \left(\sum_{|\alpha| \leq 2} \int_Q e^{2s\varphi} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx dt \right), \end{aligned}$$

for every $(j, k) \in \{1, 2\}^2$, from where we get

$$\begin{aligned} & \sum_{|\alpha| \leq 2} \left(C \int_{\Omega} e^{2s\varphi_0} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx - \int_Q e^{2s\varphi} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx dt \right) \\ & \leq \sum_{j,k=1}^2 \left(\mathcal{B}_{s,\varphi}(W_{k,j}) + s^3 e^{2d_0 s} \|W_{k,j}\|_{H^1(Q)}^2 \right), \end{aligned} \quad (3.22)$$

by combining (3.7) and (3.21). Further, by recalling (2.7) and (2.10) we see for each $\alpha \in \{1, 2, 3\}^2$ with $|\alpha| = 2$, that

$$\int_Q e^{2s\varphi} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx dt = \int_{\Omega} e^{2s\varphi_0} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) \mathcal{G}_s(x) dx, \quad (3.23)$$

where

$$\mathcal{G}_s(x) := \int_{-T}^T e^{-2s(\varphi_0(x) - \varphi(x,t))} dt \leq \int_{-T}^T e^{-2s(1-\sigma(t))} dt := \mathbf{g}(s), \quad \sigma(t) := e^{-\gamma\beta t^2}. \quad (3.24)$$

As $\lim_{s \rightarrow +\infty} \mathbf{g}(s) = 0$ by Lebesgue's Theorem, we thus obtain from (3.22)-(3.24) that

$$C \sum_{|\alpha| \leq 2} \int_{\Omega} e^{2s\varphi_0} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx \leq \sum_{j,k=1}^2 \left(\mathcal{B}_{s,\varphi}(W_{k,j}) + s^3 e^{2d_0 s} \|W_{k,j}\|_{H^1(Q)}^2 \right), \quad (3.25)$$

for s sufficiently large.

Furthermore, bearing in mind that $x_0 \in \mathbb{R}^3 \setminus \overline{\Omega}$, we notice from (2.7) and (2.10) that

$$\varphi_0(x) \geq \min_{x \in \overline{\Omega}} e^{\gamma(|x-x_0|^2 + \beta_0)} \geq d_1 > d_0, \quad x \in \overline{\Omega}, \quad (3.26)$$

where d_0 is defined in (2.14). From this and (3.25) then follows that

$$C \sum_{|\alpha| \leq 2} \int_{\Omega} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx \leq e^{Cs} \sum_{j,k=1}^2 \mathcal{B}(W_{k,j}) + s^3 e^{-2(d_1-d_0)s} M, \quad (3.27)$$

where M is the same as in (1.12), and

$$\mathcal{B}(W) = \int_{\Sigma} \left(|\nabla_{\tau} V_{\tau}|^2 + |\nabla_{\tau} U_{\nu}|^2 + |U'_{\nu}|^2 + |V'_{\tau}|^2 + (|U_{\nu}|^2 + |V_{\tau}|^2) \right) d\sigma dt, \quad W = (U, V). \quad (3.28)$$

In view of (3.26), (3.27)-(3.28) then yields

$$\sum_{|\alpha| \leq 2} \int_{\Omega} \left(|\partial^\alpha \mu|^2 + |\partial^\alpha \lambda|^2 \right) dx \leq C \left(\sum_{j,k=1}^2 \mathcal{B}(W_{k,j}) \right)^{\kappa},$$

for some $\kappa \in (0, 1)$, proving Theorem 1.

Acknowledgements

Part of this work was done while the first author was visiting the Université d'Aix-Marseille. He gratefully acknowledges the hospitality of the Université d'Aix-Marseille.

References

- [1] M. Bellassoued: *Global logarithmic stability in inverse hyperbolic problem by arbitrary boundary observation*, Inverse Problems 20 (2004), 1033-1052.
- [2] M. Bellassoued: *Uniqueness and stability in determining the speed of propagation of second-order hyperbolic equation with variable coefficients*, Applicable Analysis 83 (2004), 983-1014.
- [3] M.I. Belishev and V.M. Isakov: *On the Uniqueness of the Recovery of Parameters of the Maxwell System from Dynamical Boundary Data*, Journal of Mathematical Sciences Volume 122, Number 5, 3459-3469, (2004).
- [4] M. Bellassoued and M.Yamamoto: *Logarithmic stability in determination of a coefficient in an acoustic equation by arbitrary boundary observation*, J. Math. Pures App. 85(2006), 193-224.
- [5] M. Bellassoued and M.Yamamoto: *Determination of a coefficient in the wave equation with a single measurment*, Applicable Analysis 87 (2008), 901-920.
- [6] M. Bellassoued and M.Yamamoto: *Carleman estimates with second large parameter for second order hyperbolic operators in a Riemannian manifolds*, Applicable Analysis (2011),.
- [7] A. Benabdallah, M. Cristofol, P. Gaitan and M. Yamamoto: *Inverse problem for a parabolic system with two components by measurements of one component*. Appl. Ana. 88 (2009), 683 – 709.
- [8] A.L. Bukhgeim: *Introduction to the Theory of Inverse Problems*, VSP, Utrecht (2000).
- [9] A.L. Bukhgeim, J. Cheng, V. Isakov and M. Yamamoto: *Uniqueness in determining damping coefficients in hyperbolic equations*, S.Saitoh et al. (eds), Analytic Extension Formulas and their Applications, 27-46 (2001).
- [10] A.L. Bukhgeim and M.V. Klibanov: *Global uniqueness of class of multidimensional inverse problems*, Soviet Math. Dokl. 24 (1981), 244-247.
- [11] T. Carleman: *Sur un problème d'unicité pour les systèmes d'équations aux dérivées partielles à deux variables indépendents*, Ark. Mat. Astr. Fys. 2B (1939), 1-9.
- [12] P. Caro : *Stable determination of the electromagnetic coefficients by boundary measurements*. Inverse Problems, 26 (2010), 105014, 25 pp.
- [13] P. Caro, P. Ola and M. Salo: *Inverse boundary value problem for Maxwell equations with local data*. Comm. PDE 34 (2009), no. 11, p. 1425-1464.
- [14] M. Cristofol and L. Roques :*Biological invasions: deriving the regions at risk from partial measurements*. Mathematical Biosciences (2008), 215(2), 158-166.
- [15] M. Cristofol, and E. Soccorsi: *Stability estimate in an inverse problem for non-autonomous magnetic Schrödinger equations*. Applicable Analysis (2011), 90(10), 1499-1520.
- [16] R. Dautray and J.-L. Lions:*Mathematical analysis and Numerical Methods for Science and Technology*, Vol. 3, Spectral theory and applications, Springer Verlag, Berlin Heidelberg, 1992.

- [17] R. Dautray and J.-L. Lions: *Mathematical analysis and Numerical Methods for Science and Technology*, Vol. 5, Evolution problems I, Springer Verlag, Berlin Heidelberg, 1992.
- [18] M. Eller, V. Isakov, G. Nakamura and D. Tataru: *Uniqueness and stability in the Cauchy problem for Maxwell and elasticity systems*, in "Nonlinear Partial Differential Equations and their Applications", Collège de France Seminar, Vol. 14, North-Holland, Amsterdam (2002), 329-349.
- [19] S. He and V. G. Romanov: *Some explicit formulas for crack identification in conductors using boundary measurements of dc fields*, J. Appl. Phys. 85 vol. 9 (1999),
- [20] L. Hörmander: *Linear Partial Differential Operators*, Springer-Verlag, Berlin (1963).
- [21] O. Imanuvilov: *On Carleman estimates for hyperbolic equations*, Asymptotic Analysis 32 (2002), 185-220.
- [22] O. Imanuvilov and M. Yamamoto: *Global Lipschitz stability in an inverse hyperbolic problem by interior observations*, Inverse Problems 17 (2001), 717-728.
- [23] O. Imanuvilov and M. Yamamoto: *Determination of a coefficient in an acoustic equation with single measurement*, Inverse Problems 19 (2003), 157-171.
- [24] V. Isakov: *Inverse Problems for Partial Differential Equations*, Springer-Verlag, Berlin (1998, 2005).
- [25] M.A. Kazemi and M.V. Klibanov: *Stability estimates for ill-posed Cauchy problems involving hyperbolic equations and inequality*, Applicable Analysis 50 (1993), 93-102.
- [26] A. Khaïdarov: *On stability estimates in multidimensional inverse problems for differential equation*, Soviet Math. Dokl. 38 (1989), 614-617.
- [27] M.V. Klibanov: *Inverse problems in the "large" and Carleman bounds*, Differential Equations 20 (1984), 755-760.
- [28] M.V. Klibanov: *Inverse problems and Carleman estimates*, Inverse Problems 8 (1992), 575-596.
- [29] M.V. Klibanov and A. Timonov: *Carleman Estimates for Coefficient Inverse Problems and Numerical Applications*, VSP, Utrecht (2004).
- [30] M.V. Klibanov and M. Yamamoto: *Lipschitz stability of an inverse problem for an acoustic equation*, Applicable Analysis 85 (2006), 515-538.
- [31] Y. Kurylev, M. Lassas and E. Somersalo: *Maxwell's equations with a polarization independent wave velocity: Direct and inverse problems*, J. Math. Pures Appl. , 86 (2006), 237-270.
- [32] M.M. Lavrent'ev, V.G. Romanov and S.P. Shishat'skiĭ: *Ill-posed Problems of Mathematics Physics and Analysis*, American Math. Soc., Providence (1986).
- [33] S. Li and M. Yamamoto: *Carleman estimate for Maxwell's Equations in anisotropic media and the observability inequality*, Journal of Physics: Conference Series 12 (2005) 110-115.
- [34] S. Li and M. Yamamoto: *An inverse source problem for Maxwell's equations in anisotropic media*, Applicable Analysis, Volume 84, Issue 10, (2005).

- [35] P. Ola, L. Paivarinta and E. Somersalo: *An inverse boundary value problem in electrodynamics*, Duke Math. J., 70 (1993), 617-653.
- [36] E. Matthias and M. Yamamoto: *A Carleman inequality for the stationary anisotropic Maxwell system*, Journal de Mathématiques Pures et Appliqués, Volume 86, Issue 6, (2006), 449-462.
- [37] S. Nicaise and C. Pignotti: *Internal and Boundary Observability Estimates for the Heterogeneous Maxwell's System*, Applied Mathematics & Optimization Volume 54, Number 1, 47-70, (2006).
- [38] M. Popov, S. He and R. Thottappillil: *Reconstruction of lightning currents and return stroke model parameters using remote electromagnetic fields*, J. Geophys. Research 105, Iss D19, 24469-24481, (2000).
- [39] V.G. Romanov, S.I. Kabanikhin: *Inverse problems for Maxwell's equations* Walter de Gruyter (1994).
- [40] M. Salo, C. Kenig and G. Uhlmann: *Inverse problems for the anisotropic Maxwell equations*, Duke Math. J. 157 (2011), no. 2, 369-419.
- [41] S. N. Starostenko: *Determining permittivity and permeability from the reflection coefficient of optically thin samples*, Jour. Comm. Tech. & Elec. vol. 46, no2 (2001), 221-228.
- [42] D. Tataru: *Carleman estimates and unique continuation for solutions to boundary value problems*, J.Math.Pures App. 75 (1996), 367-408.
- [43] M. Yamamoto: *Carleman estimates for parabolic equations and applications*, Inverse Problems 25 (2009) 123013 (75pp).