

XXZ-TYPE BETHE ANSATZ EQUATIONS AND QUASI-POLYNOMIALS

JIAN RONG LI* AND VITALY TARASOV*

**Department of Mathematics, Lanzhou University, Lanzhou 730000, P. R. China*

**Department of Mathematical Sciences, Indiana University–Purdue University Indianapolis
402 North Blackford St, Indianapolis, IN 46202-3216, USA*

**St. Petersburg Branch of Steklov Mathematical Institute
Fontanka 27, St. Petersburg, 191023, Russia*

ABSTRACT. We study solutions of the Bethe ansatz equation for the XXZ-type integrable model associated with the Lie algebra \mathfrak{sl}_N . We give a correspondence between solutions of the Bethe ansatz equations and collections of quasi-polynomials. This extends the results of E. Mukhin and A. Varchenko for the XXX-type model and the trigonometric Gaudin model.

1. INTRODUCTION

In this paper we study solutions of the Bethe ansatz equation for the XXZ-type integrable models associated with the Lie algebra \mathfrak{sl}_N , see (2.1). These equations arise in the Bethe ansatz method of computing eigenvalues and eigenvectors of commuting Hamiltonians of integrable models. The method gives the eigenvalues and eigenvectors by evaluating explicit rational functions on solutions of the Bethe ansatz equations, see for instance [KBI] and references therein.

Solutions of the Bethe ansatz equations for the Gaudin model, both rational and trigonometric, as well for the XXX-type model have been studied by E. Mukhin and A. Varchenko in [MV1]–[MV3]. They established a correspondence between solutions of the Bethe ansatz equations and spaces of polynomials, quasi-polynomials or quasi-exponentials with certain properties. In this paper we extend the results of [MV2], [MV3] to the case of the XXZ-type Bethe ansatz equations.

Our method to construct a collection of quasi-polynomials corresponding to a solution of the Bethe ansatz equations is not completely analogous to that of [MV2], [MV3]. As a result, we managed to weaken technical assumptions on solutions of the Bethe ansatz equations compared with those imposed in [MV3]. Another advantage is that our method works without restrictions for the root of unity case.

The plan of the paper is as follows. In Section 2, we describe the XXZ-type Bethe ansatz equations and refine the problem. In particular, we define regular and admissible solutions of the Bethe ansatz equations. In Section 3, we introduce regular collections of quasi-polynomials and show that each such a collection gives a regular solution of the Bethe ansatz equations. We also formulate there the main result of the paper, Theorem 3.4, which

*E-mail: lijrl@lzu.edu.cn, lijrl07@gmail.com

*E-mail: vt@math.iupui.edu, vt@pdmi.ras.ru

*Supported in part by NSF grant DMS-0901616

says that every admissible regular solution of the Bethe ansatz equations comes from a regular collection of quasi-polynomials. We prove Theorem 3.4 in Section 4. In Section 5, for a collection of quasi-polynomials \mathcal{U} we consider the monic difference operator $D_{\mathcal{U}}$ whose kernel is generated by \mathcal{U} . We show that the operator $D_{\mathcal{U}}$ has rational coefficients if and only if there is a regular collection of quasi-polynomials $\tilde{\mathcal{U}}$ such that $D_{\tilde{\mathcal{U}}} = D_{\mathcal{U}}$. Also, for each solution \mathbf{t} of the Bethe ansatz equations we define a difference operator $D^{\mathbf{t}}$ and show that the collection of quasi-polynomials \mathcal{U} associated with \mathbf{t} gives a basis of the kernel of $D^{\mathbf{t}}$, that is, $D^{\mathbf{t}} = D_{\mathcal{U}}$. The Appendix contains necessary technical identities.

The authors thank E. Mukhin for helpful discussions.

2. BETHE ANSATZ EQUATIONS AND QUASI-POLYNOMIALS

2.1. Bethe ansatz equations. Throughout the paper we fix a complex number q such that $q \neq 0, \pm 1$. More precisely, we fix the value of $\log q$ and for any complex number α set $q^\alpha = \exp(\alpha \log q)$.

Fix an integer $N \geq 2$. Given a collection $\mathbf{l} = (l_1, \dots, l_{N-1})$ of nonnegative integers, consider the space $\mathbb{C}^{\mathbf{l}} = \mathbb{C}^{l_1} \oplus \dots \oplus \mathbb{C}^{l_{N-1}}$. We label coordinates on this space by two indices, the superscript enumerating the summands and the subscript enumerating coordinates within a summand. That is, we write

$$\mathbf{t} = (t_1^{(1)}, \dots, t_{l_1}^{(1)}, \dots, t_1^{(N-1)}, \dots, t_{l_{N-1}}^{(N-1)}) \in \mathbb{C}^{\mathbf{l}}.$$

The product of the symmetric groups $S_{\mathbf{l}} = S_{l_1} \times \dots \times S_{l_{N-1}}$ acts on $\mathbb{C}^{\mathbf{l}}$ by permuting coordinates with the same superscript. The $S_{\mathbf{l}}$ -orbit of a single point will be called an *elementary $S_{\mathbf{l}}$ -orbit*.

Let $T_1(x), \dots, T_{N-1}(x)$ be monic polynomials in one variable and $\lambda_1, \dots, \lambda_N$ be complex numbers. Consider a system of equations on the variables $t_b^{(a)}$, $a = 1, \dots, N-1$, $b = 1, \dots, l_a$,

$$(2.1) \quad T_i(t_j^{(i)} q^2) \prod_{r=1}^{l_{i-1}} (t_j^{(i)} q - t_r^{(i-1)} q^{-1}) \prod_{s=1}^{l_{i+1}} (t_j^{(i)} - t_s^{(i+1)}) \prod_{\substack{k=1 \\ k \neq j}}^{l_i} (t_j^{(i)} q^{-1} - t_k^{(i)} q) = \\ = q^{2(\lambda_{i+1} - \lambda_i + l_i) - l_{i-1} - l_{i+1}} T_i(t_j^{(i)}) \prod_{r=1}^{l_{i-1}} (t_j^{(i)} - t_r^{(i-1)}) \prod_{s=1}^{l_{i+1}} (t_j^{(i)} q^{-1} - t_s^{(i+1)} q) \prod_{\substack{k=1 \\ k \neq j}}^{l_i} (t_j^{(i)} q - t_k^{(i)} q^{-1}),$$

$i = 1, \dots, N-1$, $j = 1, \dots, l_i$. Here and later we assume that $l_0 = l_N = 0$, unless otherwise stated. Equations (2.1) are called the XXZ-type Bethe ansatz equations associated with the data $\mathbf{l} = (l_1, \dots, l_{N-1})$, $\mathbf{T} = (T_1, \dots, T_{N-1})$, $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$.

The group $S_{\mathbf{l}}$ acts on solutions of equations (2.1) — for every solution \mathbf{t} of (2.1), all points of the $S_{\mathbf{l}}$ -orbit of \mathbf{t} are solutions of (2.1) as well.

Equations (2.1) arise in the Bethe ansatz method for the quantum integrable model defined on the irreducible finite-dimensional representation of the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_N)$. The polynomials $T_1(x), T_2(xq^{-2}), \dots, T_{N-1}(xq^{-2(N-1)})$ are the Drinfeld polynomials of the

representation and the parameters $q^{2\lambda_1}, \dots, q^{2\lambda_N}$ describe the algebra of commuting Hamiltonians of the model. Notice that in the literature equations (2.1) are usually written in the form

$$(2.2) \quad \frac{T_i(t_j^{(i)} q^2)}{T_i(t_j^{(i)})} \prod_{r=1}^{l_{i-1}} \frac{t_j^{(i)} q - t_r^{(i-1)} q^{-1}}{t_j^{(i)} - t_r^{(i-1)}} \prod_{s=1}^{l_{i+1}} \frac{t_j^{(i)} - t_s^{(i+1)}}{t_j^{(i)} q^{-1} - t_s^{(i+1)} q} \prod_{\substack{k=1 \\ k \neq j}}^{l_i} \frac{t_j^{(i)} q^{-1} - t_k^{(i)} q}{t_j^{(i)} q - t_k^{(i)} q^{-1}} = \\ = q^{2(\lambda_{i+1} - \lambda_i + l_i) - l_{i-1} - l_{i+1}},$$

$$i = 1, \dots, N-1, \quad j = 1, \dots, l_i.$$

2.2. Regular solutions of Bethe ansatz equations. For $i = 1, \dots, N-1$, define the polynomials

$$(2.3) \quad p_i(x; \mathbf{t}) = \prod_{j=1}^{l_i} (x - t_j^{(i)})$$

and

$$(2.4) \quad P_i(x; \mathbf{t}) = q^{2\lambda_{i+1}} p_i(xq^2; \mathbf{t}) p_{i-1}(x; \mathbf{t}) p_{i+1}(xq^{-2}; \mathbf{t}) T_i(x) + \\ + q^{2\lambda_i} p_i(xq^{-2}; \mathbf{t}) p_{i-1}(xq^2; \mathbf{t}) p_{i+1}(x; \mathbf{t}) T_i(xq^2).$$

Then equations (2.1) take the form

$$(2.5) \quad P_i(t_j^{(i)}; \mathbf{t}) = 0, \quad i = 1, \dots, N-1, \quad j = 1, \dots, l_i.$$

A solution \mathbf{t} of equations (2.1) is called *regular* if for any $i = 1, \dots, N-1$, and any subset $I \subset \{1, \dots, l_i\}$, the following condition holds: suppose that all the coordinates $t_j^{(i)}$, $j \in I$ are equal with the common value $t_I^{(i)}$; then

$$(2.6) \quad \frac{d^k}{dx^k} P_i(x; \mathbf{t})|_{x=t_I^{(i)}} = 0, \quad k = 0, \dots, |I| - 1,$$

where and $|I|$ is the cardinality of I . Clearly, if for each $i = 1, \dots, N-1$, all the coordinates $t_j^{(i)}$, $j = 1, \dots, l_i$, are distinct, the solution \mathbf{t} is regular.

If \mathbf{t} is a regular solution of equations (2.1), then all points of the S_I -orbit of \mathbf{t} are regular solutions of equations (2.1) as well.

The next proposition is valid by definition of a regular solution of equations (2.1).

Proposition 2.1. *A point \mathbf{t} is a regular solution of equations (2.1) if and only if for any $i = 1, \dots, N-1$, the polynomial $p_i(x; \mathbf{t})$ divides the polynomial $P_i(x; \mathbf{t})$.*

3. QUASI-POLYNOMIALS

3.1. Quasi-polynomials. A quasi-polynomial $f(x)$ of type α is an expression of the form $f(x) = x^\alpha p(x, \log x)$, where $\alpha \in \mathbb{C}$ and $p(x, s)$ is a polynomial. Call the quasi-polynomial $f(x)$ *log-free* if $p(x, s)$ does not depend on s . Say that a polynomial $r(x)$ divides $f(x)$ if $r(x)$ divides $p(x, s)$. The product of quasi-polynomials of types α and β is supposed to be of type $\alpha + \beta$. Clearly, the product of two nonzero quasi-polynomials is not zero.

We will use only algebraic properties of quasi-polynomials. The key relation is: if $f(x) = x^\alpha p(x, \log x)$, then

$$f(xq^\beta) = q^{\alpha\beta} x^\alpha p(xq^\beta, \log x + \beta \log q).$$

Given functions $g_1(x), \dots, g_k(x)$, their discrete Wronskian $W_k[g_1, \dots, g_k]$ is defined by the rule

$$(3.1) \quad W_k[g_1, \dots, g_k](x) = \det (g_i(xq^{-2(j-1)}))_{i,j=1}^k.$$

Let $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$ be a collection of complex numbers. By definition, a *collection of quasi-polynomials of type $\boldsymbol{\lambda}$* is a sequence of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ such that for every $i = 1, \dots, N$, the quasi-polynomial u_i has type λ_i , and $W_N[u_1, \dots, u_N] \neq 0$. We call the collection \mathcal{U} *semiregular* if $W_N[u_1, \dots, u_N]$ is log-free.

For a semiregular collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ of type $\boldsymbol{\lambda}$, a sequence of monic polynomials $\mathcal{T} = (T_1, \dots, T_N)$ is called a *preframe* of \mathcal{U} if for any $k = 1, \dots, N-1$ and any subset $\{i_1, \dots, i_k\} \subset \{1, \dots, N\}$, the product $\prod_{i=1}^k \prod_{j=0}^{k-i} T_{N-i+1}(xq^{-2j})$ divides the quasi-polynomial $W_k[u_{i_1}, \dots, u_{i_k}](x)$ of type $\lambda_{i_1} + \dots + \lambda_{i_k}$, and

$$(3.2) \quad W_N[u_1, \dots, u_N](x) = \text{const } x^{\lambda_1 + \dots + \lambda_N} \prod_{i=1}^N \prod_{j=0}^{N-i} T_{N-i+1}(xq^{-2j}).$$

Denote

$$(3.3) \quad Q_k^{\mathcal{T}}(x) = \prod_{i=1}^k \prod_{j=0}^{k-i} T_{N-i+1}(xq^{-2j}), \quad k = 1, \dots, N.$$

Say that a preframe \mathcal{T} is stronger than a preframe $\tilde{\mathcal{T}} = (\tilde{T}_1, \dots, \tilde{T}_N)$ if $Q_k^{\tilde{\mathcal{T}}}(x)$ divides $Q_k^{\mathcal{T}}(x)$ for any $k = 1, \dots, N-1$.

Until the end of this section we fix $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$ and assume that every collection of quasi-polynomials is of type $\boldsymbol{\lambda}$.

Lemma 3.1. *Let $\mathcal{U} = (u_1, \dots, u_N)$ be a semiregular collection of quasi-polynomials. There is a preframe $\mathcal{T} = (T_1, \dots, T_N)$ of \mathcal{U} such that for every $k = 1, \dots, N$, the polynomial $Q_k^{\mathcal{T}}(x)$, see (3.3), is the greatest common divisor of the quasi-polynomials $W_k[u_{i_1}, \dots, u_{i_k}](x)$, where $\{i_1, \dots, i_k\}$ runs over all k -element subsets of $\{1, \dots, N\}$ and $W_k[u_{i_1}, \dots, u_{i_k}](x)$ has type $\lambda_{i_1} + \dots + \lambda_{i_k}$.*

Proof. The proof is similar to that of Lemma 4.9 in [MV2]. □

The preframe \mathcal{T} defined in Lemma 3.1 is clearly the strongest preframe of \mathcal{U} . It is called the *frame* of \mathcal{U} .

For a semiregular collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ and a preframe $\mathcal{T} = (T_1, \dots, T_N)$ of \mathcal{U} , define quasi-polynomials y_0, \dots, y_{N-1} by the rule $u_N = y_{N-1} Q_1^{\mathcal{T}}$,

$$(3.4) \quad W_{N-i}[u_{i+1}, \dots, u_N](x) = y_i(x) Q_{N-i}^{\mathcal{T}}(x), \quad i = 0, \dots, N-2,$$

where $Q_1^{\mathcal{T}}, \dots, Q_N^{\mathcal{T}}$ are given by (3.3). Notice that $y_0(x)$ is proportional to $x^{\lambda_1 + \dots + \lambda_N}$. Call the semiregular collection \mathcal{U} *regular* if the quasi-polynomials y_1, \dots, y_{N-1} are log-free.

Assume that the collection \mathcal{U} is regular. For $i = 1, \dots, N-1$, let l_i be the degree of the polynomial $x^{-\lambda_{i+1}-\dots-\lambda_N} y_i(x)$, and $t_1^{(i)}, \dots, t_{l_i}^{(i)}$ be its roots. That is,

$$(3.5) \quad y_i(x) = c_i x^{\lambda_{i+1}+\dots+\lambda_N} \prod_{j=1}^{l_i} (x - t_j^{(i)}), \quad i = 1, \dots, N-1,$$

for some nonzero complex numbers c_1, \dots, c_{N-1} . Set $\mathbf{l} = \mathbf{l}_{\mathcal{U}, \mathcal{T}} = (l_1, \dots, l_{N-1})$, and denote by $X_{\mathcal{U}, \mathcal{T}}$ the $S_{\mathbf{l}}$ -orbit of the point $(t_j^{(i)})_{i=1, \dots, N-1, j=1, \dots, l_i}$ in $\mathbb{C}^{\mathbf{l}}$.

Theorem 3.2. *Let \mathcal{U} be a regular collection of quasi-polynomials of type $\boldsymbol{\lambda}$, $\mathcal{T} = (T_1, \dots, T_N)$ be a preframe of \mathcal{U} , and $\mathbf{l} = \mathbf{l}_{\mathcal{U}, \mathcal{T}} = (l_1, \dots, l_{N-1})$. Then $X_{\mathcal{U}, \mathcal{T}}$ is an elementary $S_{\mathbf{l}}$ -orbit of regular solutions of equations (2.1) associated with the data $l_1, \dots, l_{N-1}, T_1, \dots, T_{N-1}, \lambda_1, \dots, \lambda_N$.*

Proof. Define the quasi-polynomials $\tilde{y}_1, \dots, \tilde{y}_{N-1}$ by the rule

$$(3.6) \quad W_{N-i}[u_i, u_{i+2}, \dots, u_N](x) = \tilde{y}_i(x) Q_{N-i}^{\mathcal{T}}(x).$$

By Lemmas A.1, A.2, and formula (3.3), we have

$$(3.7) \quad W_2[\tilde{y}_i, y_i](x) = y_{i-1}(x) y_{i+1}(xq^{-2}) T_i(x),$$

that is,

$$\tilde{y}_i(x) y_i(xq^{-2}) - \tilde{y}_i(xq^{-2}) y_i(x) = y_{i-1}(x) y_{i+1}(xq^{-2}) T_i(x).$$

Therefore

$$(3.8) \quad \begin{aligned} y_i(x) (\tilde{y}_i(xq^2) y_i(xq^{-2}) - \tilde{y}_i(xq^{-2}) y_i(xq^2)) &= \\ &= y_i(xq^2) y_{i-1}(x) y_{i+1}(xq^{-2}) T_i(x) + y_i(xq^{-2}) y_{i-1}(xq^2) y_{i+1}(x) T_i(xq^2). \end{aligned}$$

If the point $\mathbf{t} = (t_j^{(i)})$ is defined by (3.4), (3.5), the polynomials $p_i(x; \mathbf{t})$, $P_i(x; \mathbf{t})$ are given by (2.3), (2.4), and the polynomial $\tilde{p}_i(x)$ equals $x^{-\lambda_i - \lambda_{i+2} - \dots - \lambda_N} \tilde{y}(x)$, then relation (3.8) is equivalent to

$$P_i(x; \mathbf{t}) = p_i(x) (q^{2(\lambda_i - \lambda_{i+1})} \tilde{p}_i(xq^2) p_i(xq^{-2}) - q^{2(\lambda_{i+1} - \lambda_i)} \tilde{p}_i(xq^{-2}) p_i(xq^2)).$$

By Proposition 2.1, this proves Theorem 3.2. \square

Remark. By Lemma A.1, a collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ has a preframe $\mathcal{T} = (T_1, \dots, T_N)$ if and only if $\tilde{\mathcal{U}} = (u_1/T_N, \dots, u_N/T_N)$ is a collection of quasi-polynomials and has a preframe $\tilde{\mathcal{T}} = (T_1, \dots, T_{N-1}, 1)$. Clearly, $X_{\mathcal{U}, \mathcal{T}} = X_{\tilde{\mathcal{U}}, \tilde{\mathcal{T}}}$, and for both \mathcal{U} and $\tilde{\mathcal{U}}$ equations (2.1) are the same since they involve only the polynomials T_1, \dots, T_{N-1} . Thus discussing relations between collections of quasi-polynomials and solutions of Bethe ansatz equations (2.1) we can restrict ourselves without loss of generality to preframes of the form $(T_1, \dots, T_{N-1}, 1)$.

Say that a point $\mathbf{t} = (t_j^{(i)})$ is *generic* with respect to the polynomials $\mathbf{T} = (T_1, \dots, T_{N-1})$ if $t_j^{(i)} \neq t_k^{(i+1)}$ for all $i = 1, \dots, N-2$, $j = 1, \dots, l_i$, $k = 1, \dots, l_{i+1}$, and $T_i(t_j^{(i)}) \neq 0$ for all $i = 1, \dots, N-1$, $j = 1, \dots, l_i$. Clearly, all points of the $S_{\mathbf{l}}$ -orbit of a generic point \mathbf{t} are generic.

Lemma 3.3. *Let \mathcal{U} be a regular collection of quasi-polynomials of type $\boldsymbol{\lambda}$, $\mathcal{T} = (T_1, \dots, T_N)$ be a preframe of \mathcal{U} , and $\mathbf{l} = \mathbf{l}_{\mathcal{U}, \mathcal{T}} = (l_1, \dots, l_{N-1})$. Assume that $X_{\mathcal{U}, \mathcal{T}}$ is the $S_{\mathbf{l}}$ -orbit of a generic point with respect to the polynomials T_1, \dots, T_{N-1} . Then \mathcal{T} is the frame of \mathcal{U} .*

Proof. Let $p_i(x) = x^{-\lambda_{i+1} - \dots - \lambda_N} y_i(x)$. Suppose $\tilde{\mathcal{T}} = (\tilde{T}_1, \dots, \tilde{T}_N)$ is a preframe of \mathcal{U} strictly stronger than \mathcal{T} . Let i be the largest number such that $\tilde{T}_i \neq T_i$. Notice that $i > 1$. There is a number a such that $\tilde{T}_i(a) = 0$ and $T_i(a) \neq 0$. This implies that $p_{i-1}(a) = 0$ and $p_{i-2}(a)T_{i-1}(a) = 0$, see (3.3), (3.4). However, if $X_{\mathcal{U}, \mathcal{T}}$ is the $S_{\mathbf{l}}$ -orbit of a generic point, then for every $j = 1, \dots, N-1$, the polynomials $p_{j-1}T_j$ and p_j are coprime. The claim follows. \square

3.2. Main result. A point $\mathbf{t} = (t_j^{(i)})_{i=1, \dots, N-1, j=1, \dots, l_i}$ is called *admissible* if

$$(3.9) \quad t_j^{(i)} \neq t_k^{(i)} q^2, \quad i = 1, \dots, N-1, \quad j, k = 1, \dots, l_i.$$

In particular, $t_j^{(i)} \neq 0$ for all $i = 1, \dots, N-1$, $j = 1, \dots, l_i$. Clearly, all points of the $S_{\mathbf{l}}$ -orbit of an admissible point \mathbf{t} are admissible.

Fix collections of complex numbers $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$, nonnegative integers $\mathbf{l} = (l_1, \dots, l_{N-1})$ and monic polynomials $\mathbf{T} = (T_1, \dots, T_{N-1})$. The following theorem is the main result of the paper.

Theorem 3.4. *Let \mathbf{t} be an admissible regular solution of equations (2.1) associated with the data $\mathbf{l}, \mathbf{T}, \boldsymbol{\lambda}$. Then there is a regular collection of quasi-polynomials \mathcal{U} of type $\boldsymbol{\lambda}$ such that $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ is a preframe of \mathcal{U} and the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$.*

Theorem 3.4 will be proved in Section 4. The proof is going in three steps. First we prove the theorem for $N = 2$. The obtained statement is employed then at the second step to construct the required collection of quasi-polynomials \mathcal{U} for general N . The final step is to show that $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ is a preframe of the constructed collection \mathcal{U} .

Corollary 3.5. *Let \mathbf{t} be a generic admissible regular solution of equations (2.1) associated with the data $\mathbf{l}, \mathbf{T}, \boldsymbol{\lambda}$. Then there is a regular collection of quasi-polynomials \mathcal{U} of type $\boldsymbol{\lambda}$ such that $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ is the frame of \mathcal{U} and the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$.*

Proof. The statement follows from Theorem 3.4 and Lemma 3.3. \square

4. PROOF OF THEOREM 3.4

4.1. Proof of Theorem 3.4 for $N = 2$. Let $y(x) = x^\alpha p(x)$ be a log-free quasi-polynomial of type α . We call $y(x)$ admissible if the polynomials $p(x)$ and $p(xq^{-2})$ are coprime. In particular, this implies that $p(0) \neq 0$. Thus for an admissible quasi-polynomial $y(x)$, the number α and the polynomial $p(x)$ are determined uniquely.

Let $y(x) = x^\alpha p(x)$ be an admissible quasi-polynomial of type α . Since $p(x)$ and $p(xq^{-2})$ are coprime, it is known that there are unique polynomials $r(x)$ and $s(x)$ of degree at most $\deg p$ such that $r(x)p(x) + s(x)p(xq^{-2}) = 1$. Define the quasi-polynomials $\mathcal{A}[y]$ and $\mathcal{B}[y]$ of type $-\alpha$ by the rule

$$(4.1) \quad \mathcal{A}[y](x) = x^{-\alpha} r(x), \quad \mathcal{B}[y](x) = x^{-\alpha} s(x),$$

so that

$$(4.2) \quad y(x) \mathcal{A}[y](x) + y(xq^{-2}) \mathcal{B}[y](x) = 1.$$

For a polynomial $P(s)$ and a number c , let $\mathcal{I}[P, c]$ be the unique polynomial such that

$$\mathcal{I}[P, c](s) - c \mathcal{I}[P, c](s - 2 \log q) = P(s),$$

$\deg \mathcal{I}[P, c] = \deg P$ if $c \neq 1$, and $\deg \mathcal{I}[P, 1] = 1 + \deg P$, $\mathcal{I}[P, 1](0) = 0$. For example,

$$\mathcal{I}[1, 1](s) = \frac{s}{2 \log q}, \quad \mathcal{I}[1, c](s) = \frac{1}{1 - c}, \quad c \neq 1.$$

For a quasi-polynomial $f(x) = x^\alpha \sum_i x^i P_i(\log x)$ of type α , define the quasi-polynomial $\mathcal{J}[f]$ of type α by the rule $\mathcal{J}[f](x) = x^\alpha \sum_i x^i \mathcal{I}[P_i, q^{-2\alpha}](\log x)$, so that

$$\mathcal{J}[f](x) - \mathcal{J}[f](xq^{-2}) = f(x).$$

For example,

$$\mathcal{J}[1] = \frac{\log x}{\log q^2}, \quad \mathcal{J}[x^\alpha] = \frac{x^\alpha}{1 - q^{-2\alpha}}, \quad q^{-2\alpha} \neq 1.$$

For a quasi-polynomial $f(x) = x^\alpha p(x, \log x)$ of type α and a log-free quasi-polynomial $g(x) = x^\beta r(x)$ of type β , define the quasi-polynomial $\langle f(x)/g(x) \rangle_+ = x^{\alpha-\beta} h(x, \log x)$ of type $\alpha - \beta$ by requiring that $h(x, s)$ is the polynomial part of the ratio $p(x, s)/r(x)$, that is, $\deg_x(p(x, s) - r(x)h(x, s)) < \deg r(x)$. If $f(x) = g(x)[f(x)/g(x)]_+$, we say that $g(x)$ divides $f(x)$.

For an admissible quasi-polynomial $y(x) = x^\alpha p(x)$ of type α and a quasi-polynomial $V(x)$ of type β , define the quasi-polynomial $\mathcal{F}[y, V]$ of type $\beta - \alpha$ as follows. Let $a = \mathcal{A}[y]$ and $b = \mathcal{B}[y]$, see (4.1). Consider the quasi-polynomial

$$v(x) = \left\langle \frac{a(x)V(x) + b(xq^{-2})V(xq^{-2})}{y(xq^{-2})} \right\rangle_+$$

of type $\beta - 2\alpha$. Set $\mathcal{J}[y, V] = \mathcal{J}[v]$ and

$$(4.3) \quad \mathcal{F}[y, V](x) = V(x) \mathcal{B}[y](x) + y(x) \mathcal{J}[y, V](x).$$

Proposition 4.1. *Let $y(x)$ be an admissible quasi-polynomial of type α , $V(x)$ be a quasi-polynomial of type β . Let $Y = \mathcal{F}[y, V]$. Assume that $y(x)$ divides the quasi-polynomial $y(xq^2)V(x) + y(xq^{-2})V(xq^2)$ of type $\alpha + \beta$. Then $W_2[Y, y] = V$.*

Proof. Let $a = \mathcal{A}[y]$, $b = \mathcal{B}[y]$, $f = \mathcal{J}[y, V]$. By (4.2),

$$(4.4) \quad a(x)y(x) + b(x)y(xq^{-2}) = 1$$

and

$$\begin{aligned} a(xq^2)b(x)(y(xq^2)V(x) + y(xq^{-2})V(xq^2)) &= \\ &= a(xq^2)V(xq^2) + b(x)V(x) - (a(x)a(xq^2)V(xq^2) + b(x)b(xq^2)V(x))y(x), \end{aligned}$$

so $y(x)$ divides the quasi-polynomial $a(xq^2)V(xq^2) + b(x)V(x)$ of type $\beta - \alpha$. Hence

$$y(xq^{-2})(f(x) - f(xq^{-2})) = a(x)V(x) + b(xq^{-2})V(xq^{-2}),$$

and $W_2[Y, y] = V$ by an easy simplification using (4.4). \square

Proof of Theorem 3.4 for $N = 2$. Let $\mathbf{t} = (t_j^{(i)})$ be an admissible regular solution of equations (2.1) and $p_1(x; \mathbf{t})$ be the polynomial given by (2.3). Define an admissible quasi-polynomial $y(x) = x^{\lambda_2} p_1(x; \mathbf{t})$ of type $\alpha = \lambda_2$ and a quasi-polynomial $V(x) = x^{\lambda_1 + \lambda_2} T_1(x)$ of type $\beta = \lambda_1 + \lambda_2$.

Let $u_2 = y$ and $u_1 = \mathcal{F}[y, V]$. Proposition 2.1 shows that $y(x)$ divides the quasi-polynomial $y(xq^2)V(x) + y(xq^{-2})V(xq^2)$ of type $\alpha + \beta$. Hence by Proposition 4.1,

$$W_2[u_1, u_2] = x^{\lambda_1 + \lambda_2} T_1(x),$$

that is, $\mathcal{U} = (u_1, u_2)$ is the required collection of quasi-polynomials of type $\boldsymbol{\lambda} = (\lambda_1, \lambda_2)$. \square

4.2. Construction of a collection of quasi-polynomials. Let $\mathbf{t} = (t_j^{(i)})$ be an admissible regular solution of equations (2.1) associated with the data $\mathbf{l}, \mathbf{T}, \boldsymbol{\lambda}$. Let

$$(4.5) \quad p_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)}), \quad i = 1, \dots, N-1,$$

cf. (2.3), and $p_0(x) = p_N(x) = 1$. Then for every $i = 0, \dots, N$,

$$y_i(x) = x^{\lambda_{i+1} + \dots + \lambda_N} p_i(x)$$

is an admissible quasi-polynomial of type $\lambda_{i+1} + \dots + \lambda_N$. Set $a_i = \mathcal{A}[y_i]$ and $b_i = \mathcal{B}[y_i]$, see (4.1), so that

$$(4.6) \quad a_i(x)y_i(x) + b_i(x)y_i(xq^{-2}) = 1.$$

The next lemma is equivalent to Proposition 2.1.

Lemma 4.2. *For any $i = 1, \dots, N-1$, the quasi-polynomial $y_i(x)$ divides the quasi-polynomial*

$$(4.7) \quad A_i(x) = y_i(xq^2)y_{i-1}(x)y_{i+1}(xq^{-2})T_i(x) + y_i(xq^{-2})y_{i-1}(xq^2)y_{i+1}(x)T_i(xq^2)$$

of type $\lambda_i + 2\lambda_{i+1} + 3(\lambda_{i+2} + \dots + \lambda_N)$.

Set $Q_1(x) = 1$ and

$$(4.8) \quad Q_k(x) = \prod_{i=2}^k \prod_{j=0}^{k-i} T_{N-i+1}(xq^{-2j}), \quad k = 2, \dots, N,$$

cf. (3.3). We will construct a collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ of type $\boldsymbol{\lambda}$ such that $u_N = y_{N-1}$ and

$$(4.9) \quad W_{N-i}[u_{i+1}, \dots, u_N](x) = y_i(x)Q_{N-i}(x), \quad i = 0, \dots, N-2,$$

cf. (3.4). We employ the recursive procedure described below. Given the quasi-polynomials u_{i+1}, \dots, u_N , we obtain the quasi-polynomial u_i by formula (4.16) and verify relation (4.9) in Proposition 4.5.

For the first step of the process, set $u_{N-1} = \mathcal{F}[y_{N-1}, y_{N-2} T_{N-1}]$. Then Lemma 4.2 for $i = N - 1$ and Proposition 4.1 yield

$$(4.10) \quad W_2[u_{N-1}, u_N] = W_2[u_{N-1}, y_{N-1}] = y_{N-2} T_{N-1} = y_{N-2} Q_2,$$

which is relation (4.9) for $i = N - 2$.

Lemma 4.3. *The quasi-polynomial $y_{N-2}(x)$ divides the quasi-polynomial*

$$B(x) = y_{N-2}(xq^2) y_{N-3}(x) u_{N-1}(xq^{-2}) T_{N-2}(x) + y_{N-2}(xq^{-2}) y_{N-3}(xq^2) u_{N-1}(x) T_{N-2}(xq^2)$$

of type $\lambda_{N-2} + 3\lambda_{N-1} + 2\lambda_N$.

Proof. By Lemma 4.2, $y_{N-2}(x)$ divides the quasi-polynomial $A_{N-2}(x)$, see (4.7), of type $\lambda_{N-2} + 2\lambda_{N-1} + 3\lambda_N$. Relations (4.6) and (4.10) yield

$$\begin{aligned} B(x) &= A_{N-2}(x) (a_{N-1}(x) u_{N-1}(x) + b_{N-1}(x) u_{N-1}(xq^{-2})) + \\ &\quad + y_{N-2}(x) T_{N-1}(x) (b_{N-1}(x) - a_{N-1}(x)), \end{aligned}$$

which proves Lemma 4.3. □

Assume that the quasi-polynomials u_{i+1}, \dots, u_N are constructed already, and the following properties hold.

A. For any $j = i, \dots, N$,

$$(4.11) \quad W_{N-j}[u_{j+1}, \dots, u_N](x) = y_j(x) Q_{N-j}(x).$$

B. For any $j = i + 1, \dots, N$, there is a quasi-polynomial $w_{i+1,j}(x)$ of type $\lambda_{i+1} + \dots + \lambda_N - \lambda_j$ such that

$$(4.12) \quad W_{N-i-1}[u_{i+1}, \dots, u_{j-1}, u_{j+1}, \dots, u_N](x) = w_{i+1,j}(x) Q_{N-i-1}(x).$$

In particular, $w_{i+1,i+1}(x) = y_{i+1}(x)$, see (4.11) for $j = i + 1$.

C. For any $j = i + 1, \dots, N$, the quasi-polynomial $y_i(x)$ divides the quasi-polynomial

$$B_{ij}(x) = y_i(xq^2) y_{i-1}(x) w_{i+1,j}(xq^{-2}) T_i(x) + y_i(xq^{-2}) y_{i-1}(xq^2) w_{i+1,j}(x) T_i(xq^2)$$

of type $\lambda_i - \lambda_j + 3(\lambda_{i+1} + \dots + \lambda_N)$. In particular, $B_{i+1,i+1}(x) = A_{i+1}(x)$, see (4.7).

For $i = N - 2$, property A coincide with formula (4.10), property B is straightforward: $w_{N-1,N-1} = u_N$, $w_{N-1,N} = u_{N-1}$, and property C follows from Lemma 4.2 for $i = N - 2$, and Lemma 4.3.

Define the quasi-polynomials $w_{ij}(x)$, $j = i, \dots, N$, by the rule

$$(4.13) \quad w_{ij} = \mathcal{F}[y_i, y_{i-1} \check{w}_{i+1,j} T_i],$$

see (4.3), where $\check{w}_{i+1,j}(x) = \check{w}_{i+1,j}(xq^{-2})$. Property C and Proposition 4.1 yield

$$(4.14) \quad W_2[w_{ij}, y_i](x) = y_{i-1}(x) w_{i+1,j}(xq^{-2}) T_i(x).$$

Lemma 4.4. *The quasi-polynomial $y_i(x)$ divides the quasi-polynomial $\sum_{j=i+1}^N (-1)^j w_{ij}(x) u_j(x)$ of type $\lambda_i + \dots + \lambda_N$.*

Proof. Let $b_i = \mathcal{B}[y_i]$, see (4.1). By (4.13) and (4.3),

$$\begin{aligned} \sum_{j=i+1}^N (-1)^j w_{ij}(x) u_j(x) &= y_i(x) \sum_{j=i+1}^N (-1)^j \mathcal{J}[y_i, y_{i-1} \check{w}_{i+1,j} T_i](x) + \\ &+ b_i(x) \sum_{j=i+1}^N (-1)^j y_{i-1}(x) T_i(x) w_{i+1,j}(xq^{-2}) u_j(x). \end{aligned}$$

Then formula (4.12), Lemma A.3 and formula (4.11) for $j = i$ give

$$(4.15) \quad \sum_{j=i+1}^N (-1)^{j-i-1} w_{i+1,j}(xq^{-2}) u_j(x) = \frac{W_{N-i}[u_{i+1}, \dots, u_N](x)}{Q_{N-i-1}(xq^{-2})} = y_i(x) \prod_{k=i+1}^{N-1} T_k(x),$$

which proves the lemma. □

Set $c_{ij}(x) = w_{ij}(x)/y_i(x)$, $j = i, \dots, N$, and

$$(4.16) \quad u_i(x) = \sum_{j=i+1}^N (-1)^{j-i-1} c_{ij}(x) u_j(x).$$

By Lemma 4.4, $u_i(x)$ is a quasi-polynomial of type λ_i .

Proposition 4.5. $W_{N-i+1}[u_i, \dots, u_N](x) = y_{i-1}(x) Q_{N-i+1}(x)$.

Proof. By (4.14) and (4.12),

$$\begin{aligned} c_{ij}(x) - c_{ij}(xq^{-2}) &= \frac{y_{i-1}(x) w_{i+1,j}(xq^{-2}) T_i(x)}{y_i(x) y_i(xq^{-2})} \\ &= \frac{y_{i-1}(x) T_i(x) W_{N-i-1}[u_{i+1}, \dots, u_{j-1}, u_{j+1}, \dots, u_N](xq^{-2})}{y_i(x) y_i(xq^{-2}) Q_{N-i-1}(xq^{-2})}. \end{aligned}$$

So Lemma A.3 yields

$$\sum_{j=i+1}^N (-1)^j (c_{ij}(x) - c_{ij}(xq^{-2})) u_j(xq^{-2l}) = 0, \quad l = 1, \dots, N - i - 1,$$

and

$$\begin{aligned} \sum_{j=i+1}^N (-1)^{N-j} (c_{ij}(x) - c_{ij}(xq^{-2})) u_j(xq^{2(i-N)}) &= \\ &= \frac{y_{i-1}(x) T_i(x) W_{N-i}[u_{i+1}, \dots, u_N](xq^{-2})}{y_i(x) y_i(xq^{-2}) Q_{N-i-1}(xq^{-2})} = \frac{y_{i-1}(x) Q_{N-i+1}(x)}{W_{N-i}[u_{i+1}, \dots, u_N](x)}, \end{aligned}$$

where the last equality also uses formula (4.11) for $j = i$ and formula (4.8). These relations together with (4.16) give

$$(4.17) \quad u_i(xq^{-2l}) = \sum_{j=i+1}^N (-1)^{j-i-1} c_{ij}(x) u_j(xq^{-2l}), \quad l = 1, \dots, N-i-1,$$

and

$$(4.18) \quad u_i(xq^{2(i-N)}) = \frac{(-1)^{N-i} y_{i-1}(x) Q_{N-i+1}(x)}{W_{N-i}[u_{i+1}, \dots, u_N](x)} + \sum_{j=i+1}^N (-1)^{j-i-1} c_{ij}(x) u_j(xq^{2(i-N)}).$$

Using equalities (4.17), (4.18) in the definition of $W_{N-i+1}[u_i, \dots, u_N]$, see (3.1) completes the proof of the proposition. \square

Proposition 4.6. *For any $j = i, \dots, N$,*

$$W_{N-i}[u_i, \dots, u_{j-1}, u_{j+1}, \dots, u_N](x) = w_{ij}(x) Q_{N-i}(x),$$

and the quasi-polynomial $y_{i-1}(x)$ divides the quasi-polynomial

$$B_{i-1,j}(x) = y_{i-1}(xq^2) y_{i-2}(x) w_{ij}(xq^{-2}) T_{i-1}(x) + y_{i-1}(xq^{-2}) y_{i-2}(xq^2) w_{ij}(x) T_{i-1}(xq^2)$$

of type $\lambda_{i-1} - \lambda_j + 3(\lambda_i + \dots + \lambda_N)$.

Proof. Using (4.17) in the definition of $W_{N-i}[u_i, \dots, u_{j-1}, u_{j+1}, \dots, u_N](x)$, we get

$$W_{N-i}[u_i, \dots, u_{j-1}, u_{j+1}, \dots, u_N](x) = c_{ij}(x) W_{N-i}[u_{i+1}, \dots, u_N](x) = w_{ij}(x) Q_{N-i}(x).$$

By Lemma 4.2, $y_{i-1}(x)$ divides the quasi-polynomial $A_{i-1}(x)$, see (4.7), of type $\lambda_{i-1} + 2\lambda_i + 3(\lambda_{i+1} + \dots + \lambda_N)$. Relations (4.6) and (4.14) yield

$$\begin{aligned} B_{i-1,j}(x) &= A_{i-1}(x) (a_i(x) w_{ij}(x) + b_i(x) w_{ij}(xq^{-2})) + \\ &\quad + y_{i-1}(x) T_i(x) w_{i+1,j}(xq^{-2}) (b_i(x) - a_i(x)), \end{aligned}$$

which proves the second part of the proposition. \square

Propositions 4.5 and 4.6 shows that properties A–C with i replaced by $i - 1$ are valid. So we can construct recursively all quasi-polynomials u_1, \dots, u_N satisfying relations (4.9).

4.3. Proof of Theorem 3.4. In this section we will show that the sequence of monic polynomials $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ is a preframe of the collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ constructed in Section 4.2.

It is shown in Section 4.2 that the quasi-polynomials u_1, \dots, u_N satisfy relations (4.9), see Proposition 4.5. Moreover, for any $1 \leq i \leq j \leq N$, there is a quasi-polynomial $w_{ij}(x)$ of type $\lambda_i + \dots + \lambda_N - \lambda_j$ such that

$$(4.19) \quad W_{N-i}[u_i, \dots, u_{j-1}, u_{j+1}, \dots, u_N](x) = w_{ij}(x) Q_{N-i}(x),$$

see Proposition 4.6, and

$$(4.20) \quad W_2[w_{ij}, y_i](x) = y_{i-1}(x) w_{i+1,j}(xq^{-2}) T_i(x),$$

cf. (4.14).

Lemma 4.7. *For any $1 \leq i \leq j < k \leq N$, there is a quasi-polynomial $\tilde{w}_{i;j,k}(x)$ of type $\lambda_i + \dots + \lambda_N - \lambda_j - \lambda_k$ such that*

$$W_2[w_{ij}, w_{ik}](x) = y_{i-1}(x) \tilde{w}_{i;j,k}(x) T_i(x).$$

Proof. For $j = i$, we have $w_{ii}(x) = y_i(x)$ and $\tilde{w}_{i;i,k}(x) = -w_{i+1,k}(xq^{-2})$ by (4.20), so the statement holds. For $j > i$, by Lemma A.3 and formula (4.6) we have

$$\begin{aligned} W_2[w_{ij}, w_{ik}](x) &= (a_i(x) w_{ik}(x) + b_i(x) w_{ik}(xq^{-2})) W_2[w_{ij}, y_i](x) - \\ &\quad - (a_i(x) w_{ij}(x) + b_i(x) w_{ij}(xq^{-2})) W_2[w_{ik}, y_i](x), \end{aligned}$$

which proves Lemma 4.7. \square

Proposition 4.8. *For any $1 \leq i \leq j_1 < \dots < j_k \leq N$, there is a quasi-polynomial $\tilde{w}_{i;j_1, \dots, j_k}(x)$ of type $\lambda_i + \dots + \lambda_N - \lambda_{j_1} - \dots - \lambda_{j_k}$ such that*

$$W_k[w_{i,j_1}, \dots, w_{i,j_k}](x) = \tilde{w}_{i;j_1, \dots, j_k}(x) \prod_{l=0}^{k-2} \left(y_{i-1}(xq^{-2l}) \prod_{m=i}^{i+l} T_m(xq^{-2l}) \right).$$

Proof. We prove the statement by induction with respect to k . The case $k = 2$ is the base of induction, see Lemma 4.7.

Set $f_{ij} = W[y_i, w_{ij}]$. Assume that $j_1 = i$. By Lemma A.2 and formula (4.20),

$$\begin{aligned} W_k[y_i, w_{i,j_2}, \dots, w_{i,j_k}](x) &= \frac{W_{k-1}[f_{i,j_2}, \dots, f_{i,j_k}](x)}{y_i(xq^{-2}) \dots y_i(xq^{-2k})} = \\ &= (-1)^{k-1} W_{k-1}[w_{i+1,j_2}, \dots, w_{i+1,j_k}](xq^{-2}) \prod_{l=0}^{k-2} \frac{y_{i-1}(xq^{-2l}) T_i(xq^{-2l})}{y_i(xq^{-2(l+1)})} \\ &= (-1)^{k-1} \tilde{w}_{i+1;j_2, \dots, j_k}(x) \prod_{l=0}^{k-2} \left(y_{i-1}(xq^{-2l}) \prod_{m=i}^{i+l} T_m(xq^{-2l}) \right), \end{aligned}$$

where for the last equality we use the induction assumption. For $j_1 > i$, by Lemma A.3 and formula (4.6) we have

$$\begin{aligned} & W_k[w_{i,j_1}, \dots, w_{i,j_k}](x) = \\ &= \sum_{l=1}^k (-1)^{l-1} (a_i(x) w_{i,j_l}(x) + b_i(x) w_{i,j_l}(xq^{-2})) W_k[y_i, w_{i,j_1}, \dots, w_{i,j_{l-1}}, w_{i,j_{l+1}}, \dots, w_{i,j_k}](x), \end{aligned}$$

which proves the proposition. \square

To complete the proof of Theorem 3.4 we should show that for any $k = 1, \dots, N-1$, and any k -element subset $\{i_1, \dots, i_k\} \subset \{1, \dots, N\}$, the polynomial $Q_k(x)$ divides the quasi-polynomial $W_k[u_{i_1}, \dots, u_{i_k}](x)$ of type $\lambda_{i_1} + \dots + \lambda_{i_k}$.

Let $\{j_1, \dots, j_{N-k}\}$ be the complement of $\{i_1, \dots, i_k\}$ in $\{1, \dots, N\}$. Then by formula (4.19), Lemmas A.1, A.4, and formula (4.9),

$$\begin{aligned} W_k[u_{i_1}, \dots, u_{i_k}](x) &= \text{const} \cdot x^{(k+1-N)(\lambda_1 + \dots + \lambda_N)} \times \\ &\quad \times W_{N-k}[w_{1,j_1}, \dots, w_{1,j_{N-k}}](xq^{2(N-k-1)}) Q_{N-1}(x) \prod_{l=1}^{N-k-1} \frac{Q_{N-1}(xq^{2l})}{Q_N(xq^{2l})}. \end{aligned}$$

Then by Proposition 4.8,

$$W_k[u_{i_1}, \dots, u_{i_k}](x) = \text{const} \cdot \tilde{w}_{1;j_1, \dots, j_{N-k}}(xq^{2(N-k-1)}) Q_k(x),$$

where the quasi-polynomial $\tilde{w}_{1;j_1, \dots, j_{N-k}}$ has type $\lambda_{i_1} + \dots + \lambda_{i_k}$. Theorem 3.4 is proved.

5. DIFFERENCE OPERATORS

5.1. Difference operator of a collection of quasi-polynomials. Recall that $q \neq 0, \pm 1$. Denote by τ the multiplicative shift operator that acts on functions of x by the rule

$$(\tau f)(x) = f(xq^{-2}).$$

A function $f(x)$ is called a quasi-constant if $\tau f = f$.

An operator $D = a_0(x) + a_1(x)\tau + \dots + a_N(x)\tau^N$, where a_0, \dots, a_N are functions and a_N is not identically zero, is called a *difference operator of order N* . The functions a_0, \dots, a_N are the *coefficients* of D . If $a_N = 1$, the operator D is called *monic*.

Recall that for any collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ by definition in Section 3.1, $W_N[u_1, \dots, u_N] \neq 0$.

Lemma 5.1. *For any collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ there exists a unique monic difference operator $D_{\mathcal{U}}$ such that $D_{\mathcal{U}} u_i = 0$ for all $i = 1, \dots, N$.*

Proof. Define the operator $D_{\mathcal{U}}$ by the rule

$$(5.1) \quad D_{\mathcal{U}} f = \frac{W_{N+1}[u_1, \dots, u_N, f]}{W_N[u_1, \dots, u_N]}.$$

Then clearly $D_{\mathcal{U}} u_i = 0$ for all $i = 1, \dots, N$. On the other hand, write $D_{\mathcal{U}} = a_0 + \dots + a_{N-1} \tau^{N-1} + \tau^N$. Then equalities $D_{\mathcal{U}} u_i = 0$, $i = 1, \dots, N$, amount to a system of linear equations on a_0, \dots, a_{N-1} :

$$(5.2) \quad a_0 u_i + a_1 \tau u_i + \dots + a_{N-1} \tau^{N-1} u_i = -\tau^N u_i, \quad i = 1, \dots, N.$$

Since $W_N[u_1, \dots, u_N] \neq 0$, the matrix $(\tau^{j-1} u_i)_{i,j=1}^N$ is invertible and solution of system (5.2) has is unique. \square

The operator $D_{\mathcal{U}}$ is called the *fundamental difference operator* of the collection \mathcal{U} . Notice that by (5.1),

$$(5.3) \quad a_0(x) = (-1)^N \frac{W_N[u_1, \dots, u_N](xq^{-2})}{W_N[u_1, \dots, u_N](x)}.$$

Lemma 5.2. *Let $\mathcal{U} = (u_1, \dots, u_N)$ be a collection of quasi-polynomials. Any solution f of the difference equation $D_{\mathcal{U}} f = 0$ is a linear combination of u_1, \dots, u_N with quasi-constant coefficients.*

Proof. Let

$$(5.4) \quad c_i(x) = -\frac{W_N[u_1, \dots, u_{i-1}, f, u_{i+1}, \dots, u_N](x)}{W_N[u_1, \dots, u_N](x)}, \quad i = 1, \dots, N.$$

By Lemma A.4, $f = c_1 u_1 + \dots + c_N u_N$. If $c_i \neq 0$, consider a collection $\tilde{\mathcal{U}}_i = (u_1, \dots, u_{i-1}, f, u_{i+1}, \dots, u_N)$. Lemma 5.1 implies that $D_{\tilde{\mathcal{U}}_i} = D_{\mathcal{U}}$, and by formula (5.3),

$$\frac{W_N[u_1, \dots, u_{i-1}, f, u_{i+1}, \dots, u_N](xq^{-2})}{W_N[u_1, \dots, u_{i-1}, f, u_{i+1}, \dots, u_N](x)} = \frac{W_N[u_1, \dots, u_N](xq^{-2})}{W_N[u_1, \dots, u_N](x)}.$$

Thus $c_i(xq^{-2}) = c_i(x)$ for all $i = 1, \dots, N$. \square

Lemma 5.3. *For a collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$, set $v_N = u_N$ and*

$$(5.5) \quad v_i = \frac{W_{N-i+1}[u_i, \dots, u_N]}{W_{N-i}[u_{i+1}, \dots, u_N]}, \quad i = 1, \dots, N-1.$$

Then

$$(5.6) \quad D_{\mathcal{U}} = \left(\tau - \frac{\tau v_1}{v_1} \right) \dots \left(\tau - \frac{\tau v_N}{v_N} \right).$$

Proof. By applying repeatedly Lemma A.2, we get

$$\left(\tau - \frac{\tau v_i}{v_i} \right) \dots \left(\tau - \frac{\tau v_N}{v_N} \right) f = \frac{W_{N-i+2}[u_i, \dots, u_N, f]}{W_{N-i+1}[u_i, \dots, u_N]}, \quad i = 1, \dots, N.$$

Comparing this formula with formula (5.1) completes the proof. \square

Corollary 5.4. *Let \mathcal{U} be a regular collection of quasi-polynomials. Then the operator $D_{\mathcal{U}}$ has rational coefficients.*

Proof. For a regular collection of quasi-polynomials \mathcal{U} , the expressions $\tau v_i/v_i$ in formula (5.6) are rational functions, which proves the claim. \square

Proposition 5.5. *Let $\mathcal{U} = (u_1, \dots, u_N)$ be a collection of quasi-polynomials such that the operator $D_{\mathcal{U}}$ has rational coefficients. Then the collection \mathcal{U} is semiregular, that is, the quasi-polynomial $W_N[u_1, \dots, u_N]$ is log-free.*

We will prove Proposition 5.5, as well as Propositions 5.6–5.9 below, in Section 5.3.

Proposition 5.6. *Let \mathcal{U} be a collection of quasi-polynomials such that the operator $D_{\mathcal{U}}$ has rational coefficients. Then there is a regular collection of quasi-polynomials $\tilde{\mathcal{U}}$ such that $D_{\tilde{\mathcal{U}}} = D_{\mathcal{U}}$.*

Say that $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$ is *dominance-free* if $q^{2(\lambda_i - \lambda_j)} \neq q^{2s}$ for all $1 \leq i < j \leq N$ and $s \in \mathbb{Z}_{\geq 1}$.

Proposition 5.7. *Let \mathcal{U} be a collection of quasi-polynomials of type $\boldsymbol{\lambda}$ such that the operator $D_{\mathcal{U}}$ has rational coefficients. Assume that $\boldsymbol{\lambda}$ is dominance-free, or q is a root of unity. Then there is a regular collection of quasi-polynomials $\tilde{\mathcal{U}}$ of type $\boldsymbol{\lambda}$ such that $D_{\tilde{\mathcal{U}}} = D_{\mathcal{U}}$.*

Example. Let $\mathcal{U} = (u_1, u_2)$ be the collection of quasi-polynomials of type $\boldsymbol{\lambda} = (1, 0)$:

$$u_1(x) = x, \quad u_2(x) = \frac{1}{1 - q^{-2}} + \frac{q^2 x \log x}{2 \log q}.$$

Then

$$W_2[u_1, u_2](x) = x - x^2, \quad D_{\mathcal{U}} = \tau^2 - \left(q^{-2} + \frac{1 - xq^{-2}}{1 - x} \right) \tau + q^{-2} \frac{1 - xq^{-2}}{1 - x},$$

and $\tilde{\mathcal{U}} = (u_2, u_1)$ is a regular collection of quasi-polynomials such that $D_{\tilde{\mathcal{U}}} = D_{\mathcal{U}}$. However if q is not a root of unity, $D_{\mathcal{U}}$ is not the fundamental difference operator of any regular collection of quasi-polynomials of type $\boldsymbol{\lambda} = (1, 0)$. If $q^{2\ell} = 1$ for a positive integer ℓ , set $u'_2(x) = x^\ell u_2(x)$. Then $\tilde{\mathcal{U}}' = (u'_2, u_1)$ is a regular collection of quasi-polynomials of type $\boldsymbol{\lambda} = (1, 0)$ such that $D_{\tilde{\mathcal{U}}'} = D_{\mathcal{U}}$.

Say that $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$ is *generic* if $q^{2(\lambda_i - \lambda_j)} \neq q^{2s}$ for all $1 \leq i < j \leq N$ and $s \in \mathbb{Z}$.

Proposition 5.8. *Let $\mathcal{U} = (u_1, \dots, u_N)$ be a collection of quasi-polynomials of type $\boldsymbol{\lambda}$ such that the operator $D_{\mathcal{U}}$ has rational coefficients. Assume that $\boldsymbol{\lambda}$ is generic. Then the quasi-polynomials u_1, \dots, u_N are log-free. In particular, the collection \mathcal{U} is regular.*

Proposition 5.9. *Let $\mathcal{U} = (u_1, \dots, u_N)$ and $\tilde{\mathcal{U}} = (\tilde{u}_1, \dots, \tilde{u}_N)$ be collections of quasi-polynomials of type $\boldsymbol{\lambda}$ such that $D_{\mathcal{U}} = D_{\tilde{\mathcal{U}}}$ and the operator $D_{\mathcal{U}}$ has rational coefficients. Assume that $\boldsymbol{\lambda}$ is generic. Then there are quasi-constants c_1, \dots, c_N such that $\tilde{u}_i = c_i u_i$, $i = 1, \dots, N$. If q is not a root of unity, then the quasi-constants c_1, \dots, c_N are constants.*

5.2. Difference operator of a solution of Bethe ansatz equations. Fix collections of complex numbers $\boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_N)$, nonnegative integers $\mathbf{l} = (l_1, \dots, l_{N-1})$ and monic polynomials $\mathbf{T} = (T_1, \dots, T_{N-1})$.

Given $\mathbf{t} \in \mathbb{C}^{\mathbf{l}}$, define the polynomials $p_1(x), \dots, p_{N-1}(x)$ by the rule

$$(5.7) \quad p_i(x) = \prod_{j=1}^{l_i} (x - t_j^{(i)}), \quad i = 1, \dots, N-1,$$

cf. (4.5), and set $p_0(x) = p_N(x) = 1$. Let

$$(5.8) \quad R_i(x) = \frac{p_{i-1}(x)}{p_i(x)} \prod_{j=i}^{N-1} T_j(xq^{2(i-j)}), \quad i = 1, \dots, N,$$

Define the *fundamental difference operator* $D^{\mathbf{t}}$ of the point \mathbf{t} by the rule

$$(5.9) \quad D^{\mathbf{t}} = \left(\tau - q^{-2\lambda_1} \frac{R_1(xq^{-2})}{R_1(x)} \right) \cdots \left(\tau - q^{-2\lambda_N} \frac{R_N(xq^{-2})}{R_N(x)} \right).$$

Theorem 5.10. *Let $\mathbf{t} \in \mathbb{C}^{\mathbf{l}}$ be a solution of equations (2.1) associated with the data $\mathbf{l}, \mathbf{T}, \boldsymbol{\lambda}$. Let \mathcal{U} be a collection of quasi-polynomials of type $\boldsymbol{\lambda}$ such that $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ is a preframe of \mathcal{U} and the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$. Then $D^{\mathbf{t}} = D_{\mathcal{U}}$.*

Proof. Let $\mathcal{U} = (u_1, \dots, u_N)$. Since the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$, formulae (3.3), (3.4) imply

$$x^{\lambda_i} R_i(x) = \frac{W_{N-i+1}[u_i, \dots, u_N](x)}{W_{N-i}[u_{i+1}, \dots, u_N](x)},$$

Thus $D^{\mathbf{t}} = D_{\mathcal{U}}$, see formulae (5.5), (5.6), (5.9). \square

Recall that a point $\mathbf{t} \in \mathbb{C}^{\mathbf{l}}$ is admissible if it satisfies conditions (3.9).

Corollary 5.11. *Let $\mathbf{t} \in \mathbb{C}^{\mathbf{l}}$ be an admissible solution of equations (2.1) associated with the data $\mathbf{l}, \mathbf{T}, \boldsymbol{\lambda}$. Then there exists a collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ of type $\boldsymbol{\lambda}$ such that $D^{\mathbf{t}} = D_{\mathcal{U}}$ and*

$$(5.10) \quad W_N[u_1, \dots, u_N](x) = x^{\lambda_1 + \dots + \lambda_N} \prod_{i=1}^N \prod_{j=0}^{N-i} T_{N-i+1}(xq^{-2j}).$$

Proof. By Theorem 3.4, the collection of quasi-polynomials $\mathcal{U} = (u_1, \dots, u_N)$ constructed in Section 4.2 has a preframe $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ and the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$. Hence, formula (5.10) holds, cf. (3.2), and $D^{\mathbf{t}} = D_{\mathcal{U}}$ by Theorem 5.10. \square

Theorem 5.12. *Let $\mathbf{t} \in \mathbb{C}^{\mathbf{l}}$ be an admissible solution of equations (2.1) associated with the data $\mathbf{l}, \mathbf{T}, \boldsymbol{\lambda}$. Let \mathcal{U} be a collection of quasi-polynomials of type $\boldsymbol{\lambda}$ such that $D^{\mathbf{t}} = D_{\mathcal{U}}$. Assume that q is not a root of unity and $\boldsymbol{\lambda}$ is generic. Then $\mathcal{T} = (T_1, \dots, T_{N-1}, 1)$ is a preframe of \mathcal{U} and the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$.*

Proof. Let $\mathcal{U} = (u_1, \dots, u_N)$, and $\tilde{\mathcal{U}} = (\tilde{u}_1, \dots, \tilde{u}_N)$ be the collection of quasi-polynomials of type $\boldsymbol{\lambda}$ constructed in the proof of Theorem 3.4, see Section 4.2. By Theorem 5.10 $D_{\tilde{\mathcal{U}}} = D^{\mathbf{t}} = D_{\mathcal{U}}$. Then by Lemma 5.9, for any $i = 1, \dots, N$, the quasi-polynomials u_i and \tilde{u}_i are proportional, which proves the theorem. \square

Theorem 5.12 shows that for q being not a root of unity, generic $\boldsymbol{\lambda}$, and an admissible solution \mathbf{t} of the Bethe ansatz equations (2.1) solving the difference equation $D^{\mathbf{t}}f = 0$ allows one to produce a collection of quasi-polynomials \mathcal{U} of type $\boldsymbol{\lambda}$ such that the $S_{\mathbf{l}}$ -orbit of \mathbf{t} equals $X_{\mathcal{U}, \mathcal{T}}$. This is an alternative way to the construction of such a collection of quasi-polynomials given in Section 4.2. Moreover, Theorem 5.12 and Lemma 5.9 imply that such a collection of quasi-polynomials is unique up to rescaling of individual quasi-polynomials.

5.3. Proofs of Propositions 5.5 – 5.9. Recall that $q \neq 0, \pm 1$. Most of the technicalities in this section are related to the fact that q can be a root of unity.

Lemma 5.13. *Let $c(x)$ be a quasi-constant of the form $c(x) = x^\alpha r(x, \log x)$, where $\alpha \in \mathbb{C}$ and $r(x, y)$ is a rational function in x and a polynomial in y . Then $r(x, y)$ does not depend on y .*

Proof. Let $r(x, y) = r_0(x) + r_1(x)y + \dots + r_k(x)y^k$ and $k \geq 1$. The equality $c(x) = c(xq^{-2})$ imply that

$$(5.11) \quad r_k(x) = q^{-2\alpha} r_k(xq^{-2}), \quad r_{k-1}(x) - q^{-2\alpha} r_{k-1}(xq^{-2}) = -2k r_k(x) \log q.$$

Denote by d_i the order of $r_i(x)$ at $x = 0$, that is, $\lim_{x \rightarrow 0} (x^{-d_i} r_i(x)) \neq 0, \infty$. Then the first equality in (5.11) gives $q^{2(\alpha+d_k)} = 1$, which makes impossible matching the orders of the left and right sides in the second equality. \square

For a quasi-polynomial $f(x) = x^\alpha (P_0(x) + P_1(x) \log x + \dots + P_k(x) (\log x)^k)$, where P_0, \dots, P_k are polynomials, $P_k \neq 0$, denote by $\langle f \rangle(x) = x^\alpha P_k(x)$ the top part of $f(x)$.

Lemma 5.14. *Let D be a difference operator with rational coefficients. If a quasi-polynomial $f(x)$ satisfies the equation $Df = 0$, then $D\langle f \rangle = 0$ too.*

Proof. Since $\tau(x^\alpha x^i (\log x)^j) = q^{-2(\alpha+i)} x^\alpha x^i r(\log x)$ for some monic polynomial $r(s)$ of degree j , the claim follows. \square

Proof of Proposition 5.5. Let $c(x) = W_N[u_1, \dots, u_N](x) / \langle W_N[u_1, \dots, u_N] \rangle(x)$. By Lemma 5.14, $c(x)$ is a quasi-constant, and $c(x)$ satisfies the assumption of Lemma 5.13. Since the quasi-polynomial $\langle W_N[u_1, \dots, u_N] \rangle$ is log-free, Proposition 5.5 follows. \square

Proof of Propositions 5.6, 5.7. We will prove the Propositions by induction on the number of quasi-polynomials in the collection $\mathcal{U} = (u_1, \dots, u_N)$.

If the quasi-polynomial u_N is not log-free, let $f = \langle u_N \rangle$. By Lemma 5.14, $D_{\mathcal{U}} f = 0$. If $W_N[u_1, \dots, u_{N-1}, f] \neq 0$, set $\widehat{\mathcal{U}} = (u_1, \dots, u_{N-1}, f)$. Then the collection $\widehat{\mathcal{U}}$ has type $\boldsymbol{\lambda}$, $D_{\widehat{\mathcal{U}}} = D_{\mathcal{U}}$, and the quasi-polynomial f is log-free.

If $W_N[u_1, \dots, u_{N-1}, f] = 0$, then $W_N[u_1, \dots, u_{i-1}, f, u_{i+1}, \dots, u_N] \neq 0$ for some $i = 1, \dots, N-1$, see Lemma 5.4. Then for the collection $\widehat{\mathcal{U}} = (u_1, \dots, u_{i-1}, u_N, u_{i+1}, \dots, u_{N-1}, f)$, $D_{\widehat{\mathcal{U}}} = D_{\mathcal{U}}$, and the quasi-polynomial f is log-free.

The quasi-constants $c_i(x)$ given by (5.4) have the form $c_i(x) = x^{\lambda_N - \lambda_i} r_i(x)$ for some rational functions $r_i(x)$. Let d_i be the order of $r_i(x)$ at $x = 0$. Since $c_i(x)$ is a quasi-constant, $q^{2(\lambda_N - \lambda_i + d_i)} = 1$. For a dominance-free $\boldsymbol{\lambda}$, we have $d_i \leq 0$, and we set $\ell = 0$. If q is a root of unity, we find an integer $\ell \geq d_i$ such that $q^{2\ell} = 1$. Then $x^{\lambda_i - \lambda_N + \ell - d_i}$ is a quasi-constant, and $\hat{u}_i(x) = x^{\lambda_i - \lambda_N} x^{\ell - d_i} u_N(x)$ is a quasi-polynomial of type λ_i . Thus $\widehat{\mathcal{U}}' = (u_1, \dots, u_{i-1}, \hat{u}_i, u_{i+1}, \dots, u_{N-1}, f)$ is a collection of quasi-polynomials of type $\boldsymbol{\lambda}$, $D_{\widehat{\mathcal{U}}'} = D_{\mathcal{U}}$, and the quasi-polynomial f is log-free.

Assume now that the quasi-polynomial u_N is log-free, so $r(x) = u_N(xq^{-2})/u_N(x)$ is a rational function. Let the functions v_1, \dots, v_N be given by (5.5). Set

$$D = \left(\tau - \frac{\tau v_1}{v_1} \right) \dots \left(\tau - \frac{\tau v_{N-1}}{v_{N-1}} \right) = \tau^{N-1} + b_{N-2} \tau^{N-2} + \dots + b_0.$$

By Lemma 5.3, the operator D_U factors,

$$D \cdot (\tau - r(x)) = D_U = \tau^N + a_{N-1} \tau^{N-1} + \dots + a_0,$$

so the coefficients b_0, \dots, b_{N-2} are determined by the equations

$$a_i(x) = b_{i-1}(x) - b_i(x) r(xq^{-2i}), \quad i = 0, \dots, N-1,$$

where $b_{-1}(x) = 0$, and $b_{N-1}(x) = 1$. Therefore, $b_0(x), \dots, b_{N-1}(x)$ are rational functions.

Let $u'_i = W_2[u_i, u_N]$, and $U' = (u'_1, \dots, u'_{N-1})$. By Lemmas 5.3 and 5.1, the operator $D_{U'}$ acts as follows

$$D_{U'} f(x) = \frac{1}{u_N(xq^{2(1-N)})} D(f(x)u_N(x)),$$

and hence, has rational coefficients. By the induction assumption, there is a regular collection of quasi-polynomials $U'' = (u''_1, \dots, u''_{N-1})$ such that $D_{U''} = D_{U'}$. Set

$$(5.12) \quad c_{ij}(x) = -\frac{W_{N-1}[u'_1, \dots, u'_{j-1}, u''_j, u'_{j+1}, \dots, u'_{N-1}](x)}{W_{N-1}[u'_1, \dots, u'_{N-1}](x)}, \quad i, j = 1, \dots, N-1.$$

Similarly to the proofs of Proposition 5.5 and Lemma 5.2, one can show that

$$(5.13) \quad u''_i(x) = c_{i,1}(x) u'_1(x) + \dots + c_{i,N-1}(x) u'_{N-1}(x), \quad i = 1, \dots, N-1,$$

the Wronskians in (5.12) are log-free quasi-polynomials, and $c_{ij}(x)$ are quasi-constants of the form $c_{ij}(x) = x^{\alpha_{ij}} r_{ij}(x)$ for some rational functions $r_{ij}(x)$. Let d_{ij} be the order of $r_{ij}(x)$ at $x = 0$. Since $c_{ij}(x)$ is a quasi-constant, $q^{2(\alpha_{ij} + d_{ij})} = 1$, and the function $x^{-d_{ij}} r_{ij}(x)$ is also a quasi-constant. Denote by $P(x)$ the least common denominator of the functions $x^{-d_{ij}} r_{ij}(x)$, $i, j = 1, \dots, N-1$. The polynomial $P(x)$ is a quasi-constant as well.

Define the collection of quasi-polynomials $\tilde{U} = (\tilde{u}_1, \dots, \tilde{u}_N)$ by the rule: $\tilde{u}_N = u_N$ and

$$\tilde{u}_i(x) = P(x)(c_{i,1}(x) u_1(x) + \dots + c_{i,N-1}(x) u_{N-1}(x)), \quad i = 1, \dots, N-1.$$

Then $W_2[\tilde{u}_i, \tilde{u}_N](x) = P(x) u''_i(x)$, $i = 1, \dots, N-1$, see (5.13). By Lemmas A.1, A.2,

$$W_{N-i+1}[\tilde{u}_i, \dots, \tilde{u}_N](x) \prod_{j=1}^{N-i-1} \tilde{u}_N(xq^{-2j}) = W_{N-i}[u''_i, \dots, u''_{N-1}](x) \prod_{k=0}^{N-i} P(xq^{-2k}),$$

$i = 1, \dots, N-2$, which implies that the quasi-polynomials $W_{N-i+1}[\tilde{u}_i, \dots, \tilde{u}_N]$ are log-free, and \tilde{U} is a regular collection of quasi-polynomials. Since $D_{\tilde{U}} = D_U$ by Lemma 5.1, Proposition 5.6 is proved. \square

Proof of Proposition 5.9. By Proposition 5.5, the quasi-polynomial $W_N[u_1, \dots, u_N]$ is log-free. Since $D_U \tilde{u}_i = 0$, the quasi-polynomials $W_N[u_1, \dots, u_{j-1}, \tilde{u}_i, u_{j+1}, \dots, u_N]$ are log-free as well. Set

$$c_{ij}(x) = -\frac{W_N[u_1, \dots, u_{j-1}, \tilde{u}_i, u_{j+1}, \dots, u_N](x)}{W_N[u_1, \dots, u_N](x)}, \quad i, j = 1, \dots, N.$$

Similarly to the proof of Lemma 5.2, $c_{ij}(x)$ are quasi-constants of the form $c_{ij}(x) = x^{\lambda_i - \lambda_j} r_{ij}(x)$ for some rational functions $r_{ij}(x)$. Since λ is generic, that is, $q^{2(\lambda_i - \lambda_j)} \neq q^{2s}$ for $i \neq j$ and any $s \in \mathbb{Z}$, we get that $c_{ij} = 0$ for $i \neq j$, and $\tilde{u}_i(x) = r_{ii}(x) u_i(x)$, $i = 1, \dots, N$, where $r_{ii}(x)$ are rational functions and quasi-constants.

If q is not a root of unity, then the only rational functions that are quasi-constants are constant functions. \square

Proof of Proposition 5.8. Let $\tilde{u}_i = \langle u_i \rangle$, $i = 1, \dots, N$. Then $\tilde{\mathcal{U}} = (\tilde{u}_1, \dots, \tilde{u}_N)$ is a collection of quasi-polynomials of type λ . By Lemmas 5.14 and 5.1, $D_{\tilde{\mathcal{U}}} = D_{\mathcal{U}}$. By the proof of Proposition 5.9 there are rational functions $r_{ii}(x)$ such that $\tilde{u}_i(x) = r_{ii}(x) u_i(x)$, $i = 1, \dots, N$. Since the quasi-polynomials $\tilde{u}_1, \dots, \tilde{u}_N$ are log-free, the quasi-polynomials u_1, \dots, u_N are log-free as well. \square

APPENDIX A. THE WRONSKIAN IDENTITIES

Lemma A.1. *Given functions $f_1(x), \dots, f_k(x)$ and $g(x)$, we have*

$$W_k[gf_1, \dots, gf_k](x) = W_k[f_1, \dots, f_k](x) \prod_{i=0}^{k-1} g(xq^{-2i}).$$

The proof is straightforward.

Lemma A.2. *Given functions $f_1(x), \dots, f_j(x)$, $g_1(x), \dots, g_k(x)$, let*

$$h_i(x) = W_{j+1}[g_i, f_1, \dots, f_j](x), \quad i = 1, \dots, k.$$

Then

$$W_k[h_1, \dots, h_k](x) = W_{j+k}[g_1, \dots, g_k, f_1, \dots, f_j](x) \prod_{l=1}^{k-1} W_j[f_1, \dots, f_j](xq^{-2l}).$$

The proof is similar to that of Lemma 9.4 in [MV2].

Lemma A.3. *Given functions $f_1(x), \dots, f_k(x)$, we have*

$$\sum_{i=1}^k (-1)^i W_{k-1}[f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_k](x) f_i(xq^{-2l}) = 0, \quad l = 0, \dots, k-2,$$

$$\sum_{i=1}^k (-1)^{k-i} W_{k-1}[f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_k](x) f_i(xq^{2-2k}) = W_k[f_1, \dots, f_k](x).$$

Proof. Consider a $k \times k$ matrix M with entries $M_{ij} = f_i(xq^{-2(j-1)})$ for $j < k$ and $M_{il} = f_i(xq^{-2l})$. If $l = 0, \dots, k-2$, two rows of M are the same, hence $\det M = 0$. If $l = k-1$, then $\det M = W_k[f_1, \dots, f_k]$. Expanding the determinant in the last row yields the claim. \square

Lemma A.4. *Given functions $f_1(x), \dots, f_k(x)$, let*

$$g_i(x) = W_{k-1}[f_1, \dots, f_{i-1}, f_{i+1}, \dots, f_k](x), \quad i = 1, \dots, k.$$

Then

$$W_j[g_j, \dots, g_1](x) = W_{k-j}[f_{j+1}, \dots, f_k](xq^{2-2j}) \prod_{l=0}^{k-2} W_s[f_1, \dots, f_s](xq^{-2l}).$$

The proof is similar to that of Lemma 9.5 in [MV2].

REFERENCES

- [KBI] V. E. Korepin, N. M. Bogoliubov and A. G. Izergin, *Quantum inverse scattering method and correlation functions*, Cambridge University Press, 1993.
- [KS] P. P. Kulish and E. K. Sklyanin, *Quantum spectral transform method. Recent developments* Lect. Notes in Phys. **151** (1982) 61–119.
- [MV1] E. Mukhin, A. Varchenko, *Critical points of master functions and flag varieties*, Commun. Contemp. Math. **6** (2004), no. 1, 111–163.
- [MV2] E. Mukhin, A. Varchenko, *Solutions to the XXX type Bethe ansatz equations and flag varieties*, Cent. Eur. J. Math. **1** (2003), no. 2, 238–271.
- [MV3] E. Mukhin, A. Varchenko, *Quasi-polynomials and the Bethe ansatz*, Geom. Topol. Monogr. **13** (2008), 385–420.