

Spherical null geodesics of rotating Kerr black holes

Shahar Hod

The Ruppin Academic Center, Emeq Hefer 40250, Israel

and

The Hadassah Institute, Jerusalem 91010, Israel

(Dated: November 27, 2024)

Abstract

The non-equatorial spherical null geodesics of rotating Kerr black holes are studied *analytically*. Unlike the extensively studied equatorial circular orbits whose radii are known analytically, no closed-form formula exists in the literature for the radii of generic (non-equatorial) spherical geodesics. We provide here an approximate formula for the radii $r_{\text{ph}}(a/M; \cos i)$ of these spherical null geodesics, where a/M is the dimensionless angular-momentum of the black hole and $\cos i$ is an effective inclination angle (with respect to the black-hole equatorial plane) of the orbit. It is well-known that the equatorial circular geodesics of the Kerr spacetime (the prograde and the retrograde orbits with $\cos i = \pm 1$) are characterized by a *monotonic* dependence of their radii $r_{\text{ph}}(a/M; \cos i = \pm 1)$ on the dimensionless spin-parameter a/M of the black hole. We use here our novel analytical formula to reveal that this well-known property of the equatorial circular geodesics is actually *not* a generic property of the Kerr spacetime. In particular, we find that counter-rotating spherical null orbits in the range $(3\sqrt{3} - \sqrt{59})/4 \lesssim \cos i < 0$ are characterized by a *non-monotonic* dependence of $r_{\text{ph}}(a/M; \cos i = \text{const})$ on the dimensionless rotation-parameter a/M of the black hole. Furthermore, it is shown that spherical photon orbits of rapidly-rotating black holes are characterized by a critical inclination angle, $\cos i = \sqrt{4/7}$, above which the coordinate radii of the orbits approach the black-hole radius in the extremal limit. We prove that this critical inclination angle signals a transition in the physical properties of the spherical null geodesics: in particular, it separates orbits which are characterized by finite proper distances to the black-hole horizon from orbits which are characterized by infinite proper distances to the horizon.

I. INTRODUCTION

The characteristic geodesics of the Kerr black-hole spacetime have been extensively studied since the pioneering work of Carter [1], see also [2–7] and references therein. Of particular importance are spherical null orbits — orbits with constant coordinate radii on which massless particles can orbit the black hole. The spherical null geodesic (“photonsphere”) of a black hole provides valuable information on the structure and geometry of the black-hole spacetime.

The spherical null orbits are especially interesting from both an astrophysical and theoretical points of view. For example, the optical appearance to external observers of a star undergoing gravitational collapse is related to the physical properties of the photonsphere [6, 8, 9]. This surface also determines how the night sky would appear to an observer near a black hole or a very compact star [10]. In addition, spherical null orbits are closely related to the characteristic scattering resonances of black holes in the eikonal limit (the geometric-optics approximation) [6, 11–16]. According to the geometric-optics approximation, these characteristic black-hole quasinormal resonances correspond to massless particles trapped at the unstable null orbit and slowly leaking out [6, 11–16].

Furthermore, it was recently proved that for hairy black-hole configurations, the black-hole photonsphere provides a generic lower bound on the effective length of the hair [17, 18]. In addition, it was recently proved [19] that circular null geodesics provide the fastest way to circle a black hole as measured by asymptotic observers.

It is worth emphasizing that in most cases of physical interest [20, 21] the physical properties of the spherical null geodesics (and in particular, the radii and the energy-to-angular-momentum ratio) must be computed *numerically*. The only known exceptions are the prograde and retrograde circular orbits in the equatorial plane of the rotating Kerr black hole, in which case closed analytical formulae for the physical properties of the orbits have been provided in [2], see Eq. (21) below. The parameters of the null polar orbit are also known in a closed form [5], see Eq. (22) below.

To the best of our knowledge, no closed-form formula exists in the literature for the radii of generic (*non-equatorial*) spherical null geodesics of rotating Kerr black holes. One of the main goals of the present paper is to derive effective *analytical* formulae which describe the physical properties of such non-equatorial spherical geodesics in the Kerr spacetime.

It is well-known that the equatorial (prograde and retrograde) circular orbits of the Kerr spacetime are characterized by a *monotonic* dependence of their radii $r_{\text{ph}}(a)$ on the spin-parameter a of the black hole [2]. In the present study we shall use our new analytical formula [see Eq. (24) below] to show that this well-known property of the equatorial circular geodesics is actually *not* a generic property of the Kerr geometry. In particular, we shall show that there is a significant fraction of counter-rotating spherical orbits for which the function $r_{\text{ph}}(a)$ exhibits a *non-monotonic* behavior.

Recently, Yang et. al. [16] have studied the null spherical orbits of the Kerr spacetime. One of the most remarkable conclusions of [16] is that near-extremal Kerr black holes are characterized by a significant fraction of spherical null geodesics whose radii approach the black-hole radius in the near-extremal limit. In particular, it was observed numerically in [16] that such near-horizon orbits exist for near-extremal black holes in the finite interval

$$\sin \theta_c \equiv 0.731 \lesssim \sin \theta \leq 1 , \quad (1)$$

where θ is the inclination angle of the orbit with respect to the polar axis. Below we shall provide a fully *analytical* explanation for this phenomena. Furthermore, we shall obtain an analytical expression for the exact value of the critical polar angle $\sin \theta_c$ above which the near-horizon spherical null geodesics appear.

The rest of the paper is devoted to the investigation of the physical properties of non-equatorial spherical null geodesics in the rotating Kerr spacetime. In Sec. II we describe the dynamical equations which determine the null geodesics of the Kerr spacetime. In Sec. III we obtain the characteristic equation which determines the radii of the spherical null geodesics. In Sec. IV we solve the characteristic equation using a series expansion of $r_{\text{ph}}(a/M; \cos i)$ in powers of the dimensionless rotation-parameter a/M of the black hole. We then use this analytic approach to reveal the regime in which the function $r_{\text{ph}}(a/M)$ exhibits a non-monotonic behavior. In Sec. V we analyze the spherical null geodesics of rapidly-rotating black holes and discuss the near-horizon orbits of these near-extremal black holes. We conclude in Sec. VI with a summary of the main results.

II. DESCRIPTION OF THE SYSTEM

We shall analyze the spherical null geodesics which characterize the Kerr black-hole spacetime. In Boyer-Lindquist coordinates the metric is given by (we use gravitational units in which $G = c = 1$) [3, 22]

$$ds^2 = -\left(1 - \frac{2Mr}{\rho^2}\right)dt^2 - \frac{4Mar \sin^2 \theta}{\rho^2} dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{2Ma^2 r \sin^2 \theta}{\rho^2}\right) \sin^2 \theta d\phi^2, \quad (2)$$

where M and a are the mass and angular momentum per unit mass of the black hole, respectively. Here $\Delta \equiv r^2 - 2Mr + a^2$ and $\rho \equiv r^2 + a^2 \cos^2 \theta$. The black-hole (event and inner) horizons are located at the zeroes of Δ :

$$r_{\pm} = M \pm (M^2 - a^2)^{1/2}. \quad (3)$$

Null geodesics in the rotating Kerr spacetime are characterized by three constants of the motion [1]. In terms of the covariant Boyer-Lindquist components of the 4-momentum, these conserved quantities are [2]:

$$E \equiv -p_t = \text{total energy}, \quad (4)$$

$$L_z \equiv p_\phi = \text{component of angular momentum parallel to the symmetry axis}, \quad (5)$$

and

$$Q \equiv p_\theta^2 + \cos^2 \theta [-a^2 p_t^2 + p_\phi^2 / \sin^2 \theta]. \quad (6)$$

The geodesics in the black-hole spacetime are governed by the following set of equations [2]

$$\rho \frac{dr}{d\lambda} = \pm \sqrt{V_r}, \quad (7)$$

$$\rho \frac{d\theta}{d\lambda} = \pm \sqrt{V_\theta}, \quad (8)$$

$$\rho \frac{d\phi}{d\lambda} = (L_z / \sin^2 \theta - aE) + aT / \Delta, \quad (9)$$

$$\rho \frac{dt}{d\lambda} = a(L_z - aE \sin^2 \theta) + (r^2 + a^2)T / \Delta, \quad (10)$$

where λ is an affine parameter along the null geodesics. Here [2]

$$T \equiv E(r^2 + a^2) - L_z a, \quad (11)$$

$$V_r \equiv T^2 - \Delta[(L_z - aE)^2 + Q] , \quad (12)$$

$$V_\theta \equiv Q - \cos^2 \theta [-a^2 E^2 + L_z^2 / \sin^2 \theta] . \quad (13)$$

The effective potentials V_r and V_θ determine the orbital motions in the r and θ directions, respectively.

Equatorial orbits are characterized by $Q = 0$ [2]. It is convenient to quantify the deviation of a generic (non-equatorial) orbit from the equatorial plane of the black hole using an effective inclination angle i which is defined by [23–26]

$$\cos i \equiv \frac{L_z}{L} , \quad (14)$$

where

$$L \equiv \sqrt{L_z^2 + Q} . \quad (15)$$

Note that L and i are constants of the motion. For spherical black-hole spacetimes (with $a = 0$), L is the total angular momentum of the particle [2]. The extensively studied simple equatorial geodesics are characterized by $\cos^2 i = 1$, where $\cos i = +1/-1$ correspond to prograde/retrograde orbits, respectively. Polar orbits are characterized by $\cos i = 0$.

III. SPHERICAL NULL ORBITS IN THE BLACK-HOLE SPACETIME

Spherical geodesics in the black-hole spacetime are characterized by the two conditions [2, 3]

$$V_r = 0 \quad \text{and} \quad V_r' = 0 . \quad (16)$$

Substituting (12) into (16), one finds the ratio

$$\frac{L^2}{E^2} = \frac{3r^4 + a^2 r^2}{r^2 - a^2 \sin^2 i} \quad (17)$$

and the characteristic equation

$$\begin{aligned} r^5 - 3Mr^4 + 2a^2 r^3 \sin^2 i - 2Ma^2 r^2 + a^4 r \sin^2 i + Ma^4 \sin^2 i \\ + 2Mar \cos i \sqrt{3r^4 + (1 - 3 \sin^2 i)a^2 r^2 - a^4 \sin^2 i} = 0 , \end{aligned} \quad (18)$$

which determines the radii $r_{\text{ph}}(a/M; \cos i)$ of the null spherical geodesics. It is convenient to define the dimensionless radial coordinate \bar{r} and the dimensionless spin-parameter \bar{a} of the black hole:

$$\bar{r} \equiv r/M \quad \text{and} \quad \bar{a} \equiv a/M , \quad (19)$$

in terms of which the characteristic equation (18) becomes

$$\begin{aligned} \bar{r}^5 - 3\bar{r}^4 + 2\bar{a}^2\bar{r}^3 \sin^2 i - 2\bar{a}^2\bar{r}^2 + \bar{a}^4\bar{r} \sin^2 i + \bar{a}^4 \sin^2 i \\ + 2\bar{a}\bar{r} \cos i \sqrt{3\bar{r}^4 + (1 - 3\sin^2 i)\bar{a}^2\bar{r}^2 - \bar{a}^4 \sin^2 i} = 0. \end{aligned} \quad (20)$$

The exact (analytical) solution of the characteristic equation (20) is only known for the following two limiting simple cases:

- The *equatorial* null geodesics. As mentioned above, these orbits are characterized by $\cos i = \pm 1$, where the upper sign corresponds to the prograde circular geodesic while the lower sign corresponds to the retrograde circular geodesic. In this simple case one finds [2]

$$\bar{r}_{\text{ph}}^{\pm}(\bar{a}) \equiv \bar{r}_{\text{ph}}(\bar{a}; \cos i = \pm 1) = 2\left\{1 + \cos\left[\frac{2}{3} \cos^{-1}(\mp \bar{a})\right]\right\}. \quad (21)$$

- The *polar* null geodesic. As mentioned above, this orbit is characterized by $\cos i = 0$. In this simple case one finds [5]

$$\bar{r}_{\text{ph}}^0(\bar{a}) \equiv \bar{r}_{\text{ph}}(\bar{a}; \cos i = 0) = 1 + 2\sqrt{1 - \frac{1}{3}\bar{a}^2} \cos\left\{\frac{1}{3} \cos^{-1}\left[\frac{1 - \bar{a}^2}{(1 - \frac{1}{3}\bar{a}^2)^{\frac{3}{2}}}\right]\right\}. \quad (22)$$

To the best of our knowledge, no closed-form formula exists in the literature for the radii of generic ($\cos i \neq 0, \pm 1$) spherical null geodesics of the rotating Kerr spacetime. Below we shall derive approximate analytical formulae for the radii of these important orbits.

IV. EXPANSION IN POWERS OF a/M

We shall first consider slowly-rotating black holes. Substituting the power series

$$\bar{r}_{\text{ph}}(\bar{a}; \cos i) = \sum_{k=0}^{\infty} c_k(\cos i) \cdot \bar{a}^k \quad (23)$$

into the characteristic equation (20), one finds the analytical formula

$$\begin{aligned} \bar{r}_{\text{ph}}(\bar{a}; \cos i) = 3 - \frac{2}{\sqrt{3}} \cos i \cdot \bar{a} + \frac{1}{9}(-4 + 2 \cos^2 i) \cdot \bar{a}^2 \\ + \frac{5\sqrt{3}}{81}(\cos^3 i - 2 \cos i) \cdot \bar{a}^3 + O(\bar{a}^4) \end{aligned} \quad (24)$$

for the radii of the null spherical geodesics.

In order to test the accuracy of the approximate formula (24), we present in Table I the dimensionless ratio $r_{\text{ph}}^{\text{approx}}/r_{\text{ph}}^{\text{exact}}$ between the approximate radii as obtained from the *analytical* formula (24) and the exact radii which are obtained by *numerically* solving the characteristic equation (20) [27]. Although the expansion of $\bar{r}_{\text{ph}}(\bar{a}; \cos i)$ in powers of \bar{a} is formally valid for slowly-rotating black holes, one learns from Table I that the agreement between the numerical data and the analytical formula (24) is actually very good in almost the entire range of the dimensionless spin-parameter \bar{a} . As is evident from Table I, the analytical approximation (24) fails only for co-rotating orbits of rapidly-rotating black holes with $\bar{a} \gtrsim 0.9$. Below we shall obtain an alternative (and very accurate) analytical formula for the radii of spherical null geodesics of these near-extremal black holes.

a/M	$\cos i = -1$	$\cos i = -\frac{1}{2}$	$\cos i = 0$	$\cos i = +\frac{1}{2}$	$\cos i = +1$
0	1	1	1	1	1
0.2	1.0000287	1.0000364	1.0000448	1.0000509	1.0000444
0.4	1.0003858	1.0005469	1.0007650	1.0010268	1.0009461
0.6	1.0016708	1.0025501	1.0043633	1.0070245	1.0069312
0.8	1.0045868	1.0076324	1.0168252	1.0349897	1.0376514
0.9	1.0068806	1.0120146	1.0312513	1.0807893	1.0930592
1	1.0098488	1.0181206	1.0585458	1.2674558	1.5161605

TABLE I: Spherical null geodesics of rotating Kerr black holes. We display the dimensionless ratio $r_{\text{ph}}^{\text{approx}}/r_{\text{ph}}^{\text{exact}}$ between the approximate radii as obtained from the *analytical* formula (24) and the exact radii which are obtained by *numerically* solving the characteristic equation (20) [27]. We display results for orbits with $\cos i = -1$ (the retrograde equatorial orbit), $\cos i = -\frac{1}{2}$, $\cos i = 0$ (the polar orbit), $\cos i = +\frac{1}{2}$, and $\cos i = +1$ (the prograde equatorial orbit), and for various values of the dimensionless spin-parameter $\bar{a} \equiv a/M$ of the black hole. The agreement between the numerical data and the analytical formula (24) is very good in almost the entire range of the dimensionless rotation-parameter \bar{a} . There is, however, a large discrepancy between the numerical results and the analytical formula (24) for co-rotating orbits of rapidly-rotating black holes with $\bar{a} \gtrsim 0.9$.

It is well-known that the equatorial circular geodesics of the Kerr spacetime are characterized by a *monotonic* dependence of $\bar{r}_{\text{ph}}(\bar{a}; \cos i = \pm 1)$ on the dimensionless spin-parameter

\bar{a} of the black hole. Specifically, the equatorial prograde/retrograde null circular orbit (with $\cos i = +1/-1$) is a monotonic decreasing/increasing function of the dimensionless spin parameter \bar{a} , see Eq. (21) and also Figure 1 of [2]. We shall now use our novel formula (24) to reveal that this is actually *not* a generic property of the Kerr spacetime.

Inspection of Eq. (24) reveals that, for co-rotating orbits with a given positive value of the conserved inclination angle ($\cos i > 0$) the expansion coefficients are all negative (except from $c_0 = 3$). This implies that the radii $\bar{r}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ of such co-rotating orbits are all monotonic decreasing functions of the dimensionless spin-parameter \bar{a} of the black hole [28]. However, one finds from a second-order expansion of $\bar{r}_{\text{ph}}(\bar{a})$ [see Eq. (24)] that *counter*-rotating orbits in the range $(3\sqrt{3}-\sqrt{59})/4 \lesssim \cos i < 0$ are characterized by a *non*-monotonic dependence of $r_{\text{ph}}(\bar{a}; \cos i = \text{const})$ on the dimensionless spin-parameter of the black hole. In particular, we find that the function $\bar{r}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ has a global *maximum* at

$$\bar{a}_{\text{max}}(\cos i) \simeq \frac{3\sqrt{3} \cos i}{-4 + 2 \cos^2 i} \quad (25)$$

with

$$\bar{r}_{\text{max}}(\cos i) \simeq 3 + \frac{3 \cos^2 i}{4 - 2 \cos^2 i} . \quad (26)$$

Equation (26) is very accurate even for rapidly-rotating black holes. For example, for $\cos i = -0.55$ one finds from (26) $\bar{r}_{\text{max}} \simeq 3.27$. This value agrees to better than 1% with the numerically computed value of ~ 3.3 [29].

V. RAPIDLY-ROTATING (NEAR-EXTREMAL) BLACK HOLES

We have seen that the series expansion (24) provides a good approximation for the radii of the spherical null geodesics in almost the entire range of the dimensionless spin-parameter \bar{a} of the black hole (see Table I). However, this simple formula fails for co-rotating orbits of rapidly-rotating black holes with $\bar{a} \gtrsim 0.9$. We shall now derive an alternative analytical expression for the radii of the spherical null geodesics of these near-extremal black holes. In particular, we shall show that in the near-extremal limit the function $\bar{r}_{\text{ph}}(\bar{a}; \cos i)$ undergoes a “phase transition” at the critical inclination angle $\cos i = \sqrt{4/7}$.

For rapidly-rotating (near-extremal) black holes we define

$$\bar{r}_{\pm} \equiv 1 \pm \epsilon \quad \text{and} \quad \bar{r}_{\text{ph}} \equiv 1 + \delta_{\text{ph}} , \quad (27)$$

where $\epsilon, \delta_{\text{ph}} \ll 1$. From (27) one finds

$$\bar{a} = 1 - \epsilon^2/2 + O(\epsilon^4) . \quad (28)$$

Substituting (27) and (28) into the characteristic equation (20), one obtains

$$\begin{aligned} & 4 \cos^2 i (3 \cos^2 i - 4) (7 \cos^2 i - 4) \delta^2 + (61 \cos^6 i - 220 \cos^4 i + 176 \cos^2 i - 64) \delta^3 \\ & + 16 \cos^4 i (4 - 3 \cos^2 i) \epsilon^2 + O(\delta^4, \epsilon^4) = 0 . \end{aligned} \quad (29)$$

The qualitative behavior of $\delta_{\text{ph}}(\epsilon; \cos i)$ in the near-extremal $\epsilon \rightarrow 0$ limit depends on whether the coefficient of the $O(\delta^2)$ term in Eq. (29) is positive, negative, or zero. Note that this coefficient vanishes at

$$\xi \equiv \cos i - \sqrt{4/7} = 0 . \quad (30)$$

The solution of the characteristic equation (29) is given by [30]

$$\delta_{\text{ph}}(\epsilon; \cos i) = \begin{cases} \sqrt{\frac{4 \cos^2 i}{7 \cos^2 i - 4}} \epsilon + O(\epsilon^2/\xi^2) & \text{for } \cos i - \sqrt{4/7} \gg \epsilon^{2/3} ; \\ 2^{-1/3} \epsilon^{2/3} + O(\epsilon^{4/3}) & \text{for } -\epsilon^{2/3} \ll \cos i - \sqrt{4/7} \ll \epsilon^{2/3} ; \\ \frac{7^{3/2}}{8} (\sqrt{4/7} - \cos i) + O(\epsilon^2/\xi^2) & \text{for } \epsilon^{2/3} \ll \sqrt{4/7} - \cos i \ll 1 . \end{cases} \quad (31)$$

From (31) one learns that the solution $\delta_{\text{ph}}(\epsilon; \cos i)$ of the characteristic equation (29) exhibits a ‘‘phase transition’’ [from a $\delta_{\text{ph}}(\epsilon \rightarrow 0) \rightarrow 0$ behavior to a finite $\delta_{\text{ph}}(\epsilon \rightarrow 0)$ behavior], which occurs in the extremal $\epsilon \rightarrow 0$ limit at the critical inclination angle $\cos i = \sqrt{4/7}$.

We shall now prove that the critical inclination angle, $\cos i = \sqrt{4/7}$, signals a transition from *finite* to *infinite* proper distances of the null orbits from the black-hole horizon. The proper radial distance between the black-hole horizon (at $\bar{r}_+ = 1 + \epsilon$) and the intersection point of a null spherical orbit (of radius $\bar{r}_{\text{ph}} = 1 + \delta_{\text{ph}}$) with the equatorial plane of the black hole is given by [31]

$$\Delta \bar{\ell} = \sqrt{\delta^2 - \epsilon^2} + \ln(\delta + \sqrt{\delta^2 - \epsilon^2}) - \ln \epsilon . \quad (32)$$

Substituting into (32) the value of $\delta_{\text{ph}}(\epsilon; \cos i)$ from Eq. (31), one obtains

$$\Delta \bar{\ell}(\epsilon \rightarrow 0) = \begin{cases} \ln \left(\frac{2 \cos i + \sqrt{4 - 3 \cos^2 i}}{\sqrt{7 \cos^2 i - 4}} \right) + O(\epsilon) & \text{for } \cos i - \sqrt{4/7} \gg \epsilon^{2/3} ; \\ -\frac{1}{3} \ln \epsilon + O(1) & \text{for } -\epsilon^{2/3} \ll \cos i - \sqrt{4/7} \ll \epsilon^{2/3} ; \\ -\ln \epsilon + O(\ln \xi) & \text{for } \epsilon^{2/3} \ll \sqrt{4/7} - \cos i \ll 1 , \end{cases} \quad (33)$$

in the extremal $\epsilon \rightarrow 0$ limit. We therefore find that the critical inclination angle, which occurs at $\cos i = \sqrt{4/7}$ in the extremal limit, separates null orbits which have finite proper distances to the black-hole horizon from null orbits which have infinite proper distances to the horizon [32].

Each spherical null orbit is bounded in some strip $[\theta^-, \theta^+]$ of the polar angle θ , where $\theta^- = \pi - \theta^+$ [16]. The two polar turning-points are determined from the requirement $V_\theta(\theta^\pm) = 0$, see Eq. (13). We shall now evaluate the polar boundaries $\{\theta^-, \theta^+\}$ of the near-horizon orbits discussed above (the orbits whose coordinate radii approach the black-hole horizon in the extremal limit and which have finite proper distances to the horizon in this limit). Taking cognizance of Eqs. (17) and (31), one obtains the dimensionless ratio

$$\frac{L}{ME} = \frac{2}{\cos i} \left[1 + \sqrt{\frac{7 \cos^2 i - 4}{4 \cos^2 i}} \epsilon + O(\epsilon^2) \right] \quad (34)$$

for the null spherical orbits in the regime $\cos i - \sqrt{4/7} \gg \epsilon^{2/3}$. Substituting (34) into (13) with $V_\theta = 0$, one finds that the polar turning-points (which characterize the maximal polar-deviation of a near-horizon orbit from the equatorial plane of the black hole) are given by [33]

$$\cos^2 \theta^\pm = \frac{\cos^2 i - 4 + \sqrt{-15 \cos^4 i + 8 \cos^2 i + 16}}{2 \cos^2 i} . \quad (35)$$

For the equatorial circular orbit with $\cos i = 1$ one finds $\cos^2 \theta^\pm = 0$ ($\theta^\pm = \frac{\pi}{2}$) as expected. For the critical orbit with $\cos i = \sqrt{4/7}$ one finds from (35)

$$\sin \theta^\pm = \sqrt{4 - 2\sqrt{3}} . \quad (36)$$

This *analytically* derived value is in excellent agreement with the *numerical* value of ~ 0.731 computed in [16].

VI. SUMMARY

We have studied *analytically* the non-equatorial spherical null geodesics of rotating Kerr black holes. The main results presented in this paper are:

- (1) We have obtained the analytical expression (24) for the radii $\bar{r}_{\text{ph}}(\bar{a}; \cos i)$ of generic (non-equatorial) spherical null orbits in the Kerr spacetime. This analytical formula is formally valid for slowly-rotating black holes. However, we have demonstrated explicitly (see Table I) that the analytical formula (24) actually agrees very well with the exact (numerically

computed) values of the orbital radii in almost the entire range of the dimensionless spin-parameter \bar{a} of the black hole.

(2) The extensively studied equatorial circular geodesics of the Kerr black hole are known [2] to be characterized by a monotonic dependence of their radii $\bar{r}_{\text{ph}}(\bar{a}; \cos i = \pm 1)$ on the rotation-parameter \bar{a} of the black hole. Our analysis revealed that this is actually *not* a generic property of the Kerr spacetime. In particular, it was shown that counter-rotating orbits in the range $(3\sqrt{3} - \sqrt{59})/4 \lesssim \cos i < 0$ are characterized by a *non-monotonic* dependence of $\bar{r}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ on the dimensionless spin-parameter \bar{a} of the black hole, see Eqs. (25) and (26). Likewise, counter-rotating orbits in the range $3\sqrt{3} - \sqrt{31} \lesssim \cos i < 0$ are characterized by a *non-monotonic* behavior of the dimensionless angular-momentum-to-energy ratio $\mathcal{L}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ which is a constant of the motion.

(3) We have obtained the analytical expression (31) for the radii $\bar{r}_{\text{ph}}(\bar{a}; \cos i)$ of the spherical null geodesics of rapidly-rotating (near-extremal) Kerr black holes. This analytical formula reveals that the spherical photon orbits of rapidly-rotating black holes are characterized by a critical inclination angle, $\cos i = \sqrt{4/7}$, above which the coordinate radii of the orbits approach the black-hole radius in the extremal limit. We have shown that this critical inclination angle is the boundary between spherical null orbits which have finite proper distances to the black-hole horizon and spherical null orbits which have infinite proper distances to the horizon. Our analytical expression (36) for the critical (maximal) polar angle of such near-horizon orbits, $\sin \theta_c = \sqrt{4 - 2\sqrt{3}}$, is remarkably close to the numerically computed [16] value $\sin \theta_c = 0.731$.

ACKNOWLEDGMENTS

This research is supported by the Carmel Science Foundation. I thank Yael Oren, Arbel M. Ongo and Ayelet B. Lata for helpful discussions.

[1] B. Carter, Phys. Rev. **174**, 1559 (1968).

[2] J. M. Bardeen, W. H. Press and S. A. Teukolsky, Astrophys. J. **178**, 347 (1972).

[3] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, (Oxford University Press, New York, 1983).

- [4] S. L. Shapiro and S. A. Teukolsky, *Black holes, white dwarfs, and neutron stars: The physics of compact objects* (Wiley, New York, 1983).
- [5] E. Teo, *Gen. Relativ. and Gravit.* **35**, 1909 (2003).
- [6] V. Cardoso, A. S. Miranda, E. Berti, H. Witek and V. T. Zanchin, *Phys. Rev. D* **79**, 064016 (2009).
- [7] C. M. Will, arXiv:1208.3931 .
- [8] M. A. Podurets, *Astr. Zh.* **41**, 1090 (1964) [English translation in *Sovet Astr.-AJ* **8**, 868 (1965)].
- [9] W. L. Ames and K. S. Thorne, *Astrophys. J.* **151**, 659 (1968).
- [10] R. J. Nemiroff, *Am. J. Phys.* **61**, 619 (1993).
- [11] B. Mashhoon, *Phys. Rev. D* **31**, 290 (1985).
- [12] C. J. Goebel, *Astrophys. J.* **172**, L95 (1972).
- [13] S. Hod, *Phys. Rev. D* **80**, 064004 (2009) [arXiv:0909.0314]; S. Hod, *Phys. Rev. D* **78**, 084035 (2008) [arXiv:0811.3806]; S. Hod, *Phys. Rev. D* **75**, 064013 (2007) [arXiv:gr-qc/0611004]; S. Hod, *Class. Quant. Grav.* **24**, 4235 (2007) [arXiv:0705.2306].
- [14] S. R. Dolan, *Phys. Rev. D* **82**, 104003 (2010).
- [15] Y. Deánini, A. Folacci, and B. Raffaelli, *Phys. Rev. D* **81**, 104039 (2010); *Phys. Rev. D* **84**, 084035 (2011).
- [16] H. Yang, D. A. Nichols, F. Zhang, A. Zimmerman, Z. Zhang and Y. Chen, arXiv:1207.4253.
- [17] S. Hod, *Phys. Rev. D* **84**, 124030 (2011).
- [18] D. Núñez, H. Quevedo, and D. Sudarsky, *Phys. Rev. Lett.* **76**, 571 (1996).
- [19] S. Hod, *Phys. Rev. D* **84**, 104024 (2011).
- [20] It is well-known that realistic black holes generally rotate about their axis. Thus, an astrophysically realistic model of particle-dynamics in black-hole spacetimes must involve a non-spherical Kerr geometry with angular momentum [21].
- [21] S. Hod, *Phys. Rev. Lett.* **84**, 10 (2000) [arXiv:gr-qc/9907096].
- [22] R. P. Kerr, *Phys. Rev. Lett.* **11**, 237 (1963).
- [23] F. D. Ryan, *Phys. Rev. D* **52**, 3159 (1995); F. D. Ryan, *Phys. Rev. D* **53**, 3064 (1996).
- [24] S. A. Hughes, *Phys. Rev. D* **64**, 064004 (2001).
- [25] K. Glampedakis, S. A. Hughes, and D. Kennefick, *Phys. Rev. D* **66**, 064005 (2002).
- [26] R. Grossman, J. Levin, and G. Perez-Giz, *Phys. Rev. D* **85**, 023012 (2012).

- [27] For the special black-hole orbits with $\cos i = 0, \pm 1$ we can use the exact radii which are given by Eqs. (21)-(22).
- [28] Using the expression (22) for the radius of the spherical polar orbit, one can easily verify that $\bar{r}_{\text{ph}}(\bar{a}; \cos i = 0)$ is also a monotonic decreasing function of the spin-parameter \bar{a} .
- [29] Substituting Eq. (24) into Eq. (17), one obtains the dimensionless angular-momentum-to-energy ratio $\mathcal{L}_{\text{ph}} \equiv \frac{L}{ME}(\bar{a}; \cos i) = 3\sqrt{3} - 2 \cos i \cdot \bar{a} + \frac{\cos^2 i - 4}{6\sqrt{3}} \cdot \bar{a}^2 + \frac{2 \cos^3 i - 6 \cos i}{27} \cdot \bar{a}^3 + O(\bar{a}^4)$, which is a *constant* of the motion. Inspection of this relation reveals that, for co-rotating orbits with a given positive value of the conserved inclination angle ($\cos i > 0$) the expansion coefficients are all negative (except from $c_0 = 3\sqrt{3}$). This implies that the dimensionless angular-momentum-to-energy ratio $\mathcal{L}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ of such co-rotating orbits is a monotonic decreasing functions of the spin-parameter \bar{a} . However, one finds from a second-order expansion of $\mathcal{L}_{\text{ph}}(\bar{a})$ that *counter*-rotating orbits in the range $3\sqrt{3} - \sqrt{31} \lesssim \cos i < 0$ are characterized by a *non*-monotonic dependence of $\mathcal{L}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ on the dimensionless spin-parameter of the black hole. In particular, we find that the function $\mathcal{L}_{\text{ph}}(\bar{a}; \cos i = \text{const})$ has a global *maximum* at $\bar{a}_{\text{max}}(\cos i) \simeq \frac{6\sqrt{3}\cos i}{-4+\cos^2 i}$ with $\mathcal{L}_{\text{max}}(\cos i) \simeq 3\sqrt{3} + \frac{6\sqrt{3}\cos^2 i}{4-\cos^2 i}$.
- [30] In obtaining the solution (31) for $\delta_{\text{ph}}(\epsilon; \cos i)$ we have used the observation that, depending of the relative magnitudes of ϵ and ξ , one of the three terms in the characteristic equation (29) can be neglected as compared to the other two terms. It should be emphasized that one of the two remaining (non-negligible) terms must be positive while the other non-negligible term must be negative. Specifically, there are three distinct regimes: (1) For $\xi \gg \epsilon^{2/3}$ the $O(\delta^3)$ term is negligible as compared to the other two terms in Eq. (29). (2) For $-\epsilon^{2/3} \ll \xi \ll \epsilon^{2/3}$ the $O(\xi\delta^2)$ term is negligible as compared to the other two terms in Eq. (29). (3) For $\xi \ll -\epsilon^{2/3}$ the $O(\epsilon^2)$ term is negligible as compared to the other two terms in Eq. (29).
- [31] T. Jacobson, Class. Quant. Grav. **28**, 187001 (2011).
- [32] Note that one finds $\Delta\bar{\ell} = \frac{1}{2} \ln 3$ for the prograde equatorial orbit with $\cos i = 1$, in accord with the result of [2].
- [33] It is worth emphasizing again that the effective inclination angle i is a *constant* of the motion, while the polar angle θ varies along the trajectory within the interval $[\theta^-, \theta^+]$, where the two boundaries are determined by the relation (35).