Comment on "Trouble with the Lorentz Law of Force: Incompatibility with Special Relativity and Momentum Conservation"

F. De Zela

Departamento de Ciencias, Sección Física, Pontificia Universidad Católica del Perú, Ap. 1761, Lima, Peru.

arXiv:1210.7344v2 [physics.gen-ph] 5 Nov 2012

Mansuripur claims that the Lorentz law of force must be abandoned because it violates relativity [1]. To show this, he considers a point charge q and a point magnetic dipole $m_0 \hat{\mathbf{x}}'$, both being at rest in a reference system x'y'z' that moves with constant velocity $\mathbf{V} = V \hat{\mathbf{z}}$ relative to a second system xyz. The magnetization of the point dipole is taken as $\mathbf{M}'(\mathbf{r}', t') =$ $m_0 \widehat{\mathbf{x}}' \delta(x') \delta(y') \delta(z'-d)$ in the x' y' z'-system. By Lorentz transforming \mathbf{M}' one obtains \mathbf{M} and \mathbf{P} as the components of the second-rank tensor describing the point dipole in the xyz-system. Using the Lorentz force expression $\mathbf{F}(\mathbf{r},t) = \rho_{bound} \mathbf{E} + \mathbf{J}_{bound} \times \mathbf{B}$, it can be shown that the net force $\int \mathbf{F}(\mathbf{r},t) d^3 r$ on the point dipole is zero, while the net torque $\mathbf{T} = \int \mathbf{r} \times \mathbf{F}(\mathbf{r}, t) d^3 r = (V q m_0 / 4 \pi d^2) \hat{\mathbf{x}}.$ Mansuripur argues that the appearance of a nonzero torque in the xyz-system – in the absence of a corresponding torque in the x'y'z'-system – is sufficient proof of the inadequacy of the Lorentz law. We believe that Mansuripur's conclusion is premature, for the following reasons.

First of all, the mere appearance of a torque in only one of two inertial systems does not violate relativity. For example, Jackson [2] has recently discussed the case of a point charge q moving in the central field of a second charge Q that remains fixed in an inertial system, the x'y'z'-system, say. As seen from the x'y'z'-system, charge q experiences no torque and moves along the straight line that joins it with Q. However, when seen from the xyz-system – which uniformly moves with respect to the x'y'z'-system – a torque appears, causing a continuous change in the angular momentum of q. Jackson shows that there is no paradox here. Everything is in perfect accordance with relativity. It may exist a torque causing a change of angular momentum in one inertial frame, while there is no angular momentum and no torque in the other frame. Thus, the appearance of a nonzero torque in only one of two inertial systems is not by itself a contradiction. Coming back to Mansuripur's case, if we had obtained that the net force is zero in one system and nonzero in the other, then we would have been faced with a paradox. But the paradox would have not consisted in having two different values for the force. The paradox would have arisen from the observable consequences of this fact. Indeed, in such a case the particle's motion would have appeared as accelerated motion in one inertial frame and as uniform motion in the other. And this would be in conflict with relativity. Can we similarly reason with the torque? The answer is not. Indeed, as we

have a vanishing net force, the torque is an internal one with no effect on the body's center of mass motion. The torque's sole effect must be a rotation of the body around some axis passing through it. However, if the body is a point, such a rotation becomes meaningless. As already said, the fact that a torque appears in the xyz-system and not in the x'y'z'-system is not by itself a paradox [2]. Moreover, in Mansuripur's case such a fact does not even have any observable consequences, as in Jackson's case. Thus, it does not imply a contradiction, like motion that is accelerated in one inertial frame and uniform in the other. Here we are faced with a torque having no observable effects that might violate relativity. Instead, such a result rather hints at some possible inconsistencies that may arise when using models of "point" magnetic dipoles and the like. We analyze this issue in more detail in what follows.

To begin with, quantities like **P** and **M** serve to describe electromagnetic properties of bulk material. A magnetic dipole represents the lowest order multipole contribution from a divergenceless current distribution J within a sample [3]: $\mathbf{M} \propto \int \mathbf{r} \times \mathbf{J}(\mathbf{r}) d^3 r$. In classical electrodynamics, *point* dipoles are no more than convenient idealizations. Dipoles do not appear as fundamental quantities, like charges and fields; but as derived quantities, owing their existence to charge and current distributions. We may certainly consider very small pieces of material whose electromagnetic properties are described by smooth functions of space and time, e.g., **P** and **M**. But there is no sensible physical picture associated with a true *pointlike* object carrying *internal* currents. Idealized entities correspond to real entities whose finite extent we may neglect after having fixed the accuracy of our calculations. We may then employ mathematical tools like Dirac's δ to describe our idealization of the true physical objects. As well known, Dirac's δ is a distribution, not a function. As such, it may be used to describe idealized entities having no extent. We should bear in mind that with the help of Dirac's delta we may derive properties that belong to idealized entities, but not necessarily to real ones [4]. If we give ourselves carte blanche in the usage of Dirac's delta, we risk loosing the self-consistency of our physical model, thereby obtaining contradictory results. As an illustration of this, let us consider instead of the Lorentz force law, the simpler expression for the electric force-density: $\rho \mathbf{E}$. Such an expression follows from the very definition of \mathbf{E} as force per unit charge, and of ρ as charge per unit volume. We have thus an

expression which is valid by definition. Now, following a similar reasoning and employing the same techniques as in [1], we can get contradictory results. Let us see how. Consider the point-dipole's polarization

$$\mathbf{P}(\mathbf{r},t) = \gamma V \varepsilon_0 m_0 \widehat{\mathbf{y}} \delta(x) \delta(y) \delta\left[\gamma(z-Vt) - d\right].$$

It relates to the charge density through

$$\rho_{bound} = -\nabla \cdot \mathbf{P} = \gamma V \varepsilon_0 m_0 \delta(x) \delta'(y) \delta\left[\gamma(z - Vt) - d\right].$$

Hence, the total charge of the dipole is

$$\int \gamma V \varepsilon_0 m_0 \delta(x) \delta'(y) \delta\left[\gamma(z - Vt) - d\right] d^3 r =$$
$$= \int V \epsilon_0 m_0 \delta'(y) dy = -d(V \epsilon_0 m_0)/dy = 0,$$

as it should. Let us assume for a moment that we are using Dirac's deltas for describing not the idealized but the real entities. We would be then describing a neutral and *pointlike* particle. Now, structureless, i.e., truly pointlike particles carrying no electric charge do not couple to the electromagnetic field. If we apply an electric field \mathbf{E} to such a neutral *pointlike* particle, then the field should exert no force upon it. However, if we calculate this force for the Lorentz transformed Coulomb field

$$\mathbf{E} = \left(\frac{\gamma q}{4\pi\varepsilon_0}\right) \frac{x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + (z - Vt)\hat{\mathbf{z}}}{\left(x^2 + y^2 + (z - Vt)^2\right)^{3/2}}$$

we obtain $\int (-\nabla \cdot \mathbf{P}) \mathbf{E}(\mathbf{r}, t) d^3 r = (\gamma V m_0 q / 4\pi d^3) \hat{\mathbf{y}}.$ The same contradiction arises by considering a situation similar to that in [1] but starting with a dipole $\mathbf{P}' = p_0 \widehat{\mathbf{y}}' \delta(x') \delta(y') \delta(z'-d)$ in place of \mathbf{M}' . Such a contradiction does not invalidate the expression for the force-density, $\rho \mathbf{E}$, which follows – as already said – from the very definitions of **E** and ρ . What the contradiction signals is that we cannot use \mathbf{P} in connection with a true point-particle. Endowing a true point-particle with polarization and/or magnetization properties makes no sense in the framework of classical electrodynamics. We can certainly deal with a very small piece of matter that is polarized and/or magnetized, neglect its extension and describe it as a point. Dirac's delta can then be used as an appropriate tool. But it is the right tool only as long as we deal with the idealized model. Not every result that can be obtained with the help of such a tool necessarily applies for the real entities. If we do not distinguish the idealized model from its real counterpart, we can arrive at contradictory results, as illustrated by the above cases. As shown below, other rebuttals of Mansuripur's "paradox" [5–8] illustrate this point as well.

Like in Jackson's example [2], there are other apparent paradoxes that on closer look rather confirm the validity of classical electromagnetism, whenever one deals with cases lying within the scope of this theory. For example, if we properly handle a small current loop as

the source of the magnetic moment $m_0 \hat{\mathbf{x}}'$, everything This has been shown by Saldanha [7], who esfits. sentially reproduces a "paradox" discussed in Griffiths's well-known electrodynamics textbook [9], and by Griffiths and Hnizdo [6] in their rebuttal to [1]. However, as soon as we go from the very small to the *pointlike*. the shortcomings of classical electromagnetism begin to appear. To see this, let us consider in more detail the solution of Mansuripur's "paradox" proposed by Griffiths and Hnizdo [6]. These authors consider two different models for Mansuripur's magnetic dipole. One model is a "Gilbert dipole" (two magnetic monopoles) and the other is an "Ampere dipole" (an electric current loop). Now, the first model lies beyond classical electromagnetism, as it makes use of magnetic monopoles and the corresponding modification of the Maxwell and the Lorentz equations. If we want to prove that a given example does not lead to a true paradox within some theoretical framework, we cannot go beyond that framework. A paradox tries to show the inconsistency of some theoretical construction. By "solving" it with tools taken from outside this theoretical construction, we are doing nothing but confirming the shortcomings of that framework, which is precisely what the paradox intended to do. We should thus consider only the "Ampere dipole". As shown in [6], by considering from the perspectives of two inertial systems a current loop and a charge in relative rest, no paradox arises. The appearance of a torque and the corresponding changing angular momentum in one system and not in the other is in total agreement with relativity. The concept of "hidden momentum" explains all the observed phenomena [6, 10–12]. Though this concept is somewhat controversial, there are cases like those addressed by Griffiths and Hnizdo [6], in which hidden momentum is an incontrovertible relativistic effect that must be taken into account. In all these cases one deals with finite samples of matter. Inconsistencies appear when one goes to the pointlike limit. In the case of an "Ampere dipole", for instance, the hidden momentum is given by $\mathbf{p}_h = \mathbf{m} \times \mathbf{E}/c^2$ in the units and notation of [6]. Considering a case analogous to Mansuripur's, \mathbf{p}_h points perpendicularly to the relative velocity and hence remains unchanged when going from the x'y'z'-system to the xyz-system. The associated hidden angular momentum $\mathbf{L}_h = \mathbf{r} \times \mathbf{p}_h$ has, in turn, a rate of change given by [6] $d\mathbf{L}_h/dt = \mathbf{v} \times (\mathbf{m} \times \mathbf{E})/c^2$, as seen from the *xyz*-system. This is just the value obtained for the torque exerted on the current loop. So, everything appears to be consistent in this case. Inconsistencies show up when we deal with pointlike objects, as Mansuripur did. Grifiths and Hnizdo address this case as well, but in a way that lacks the self-consistency of their previous treatment. Indeed, when dealing with Mansuripur's case, they keep using the expression $\mathbf{p}_h = \mathbf{m} \times \mathbf{E}/c^2$ for the hidden momentum. This is a highly questionable procedure. Such an expression for \mathbf{p}_h is obtained under a series of assumptions which do not hold for the case at hand. Let us briefly review how the hidden momentum appears [11– 13]. One considers the energy-momentum tensor $T^{\mu\nu}$ of a system of particles (a "body"). It satisfies the equation $\partial T^{\mu\nu}/\partial x^{\nu} = f^{\mu}$, with $f^{\mu} = (\mathbf{f} \cdot \mathbf{v}/c, \mathbf{f})$ the fourvector representing the force-density that acts within the body. The $\mu = 0$ component of the above equation reads $\partial u/\partial t + \nabla \cdot (c^2 \mathbf{g}) = \mathbf{f} \cdot \mathbf{v}$, with $u = T^{00}$ being the energy density and $g_i = T^{0i}/c$ the three components of the momentum density **g**. Using these results one can prove that there is a "hidden momentum" $\mathbf{p}_h := \int \mathbf{g} d^3 r$, which shows up even when the body's elements move in a stationary way $(\partial u/\partial t = 0)$ and the body's center of mass remains at rest. In the stationary case, $\mathbf{p}_h = -c^{-2} \int \mathbf{r} (\mathbf{f} \cdot \mathbf{v}) d^3 r$. When the force density is given by $\mathbf{f} = -\rho \nabla \phi$, one obtains [13] $\mathbf{p}_h = -c^{-2} \int \phi \rho \mathbf{v} d^3 r$. And for the special case of a uniform, *external* electric field **E** one derives [13] (using $\phi = -\mathbf{E} \cdot \mathbf{r}$) the alternative expression $\mathbf{p}_h = \mathbf{m} \times \mathbf{E}/c^2$, with $\mathbf{m} = (1/2) \int \mathbf{r} \times \mathbf{j} d^3 r$ denoting the magnetic moment associated to the current density $\mathbf{j} = \rho \mathbf{v}$.

Although the above results can be applied to a small current loop, it is highly questionable that we may keep applying them when dealing with a pointlike object. For, first, the force density f^{μ} referred to above is the total one acting on a given element of the body [13]. That is, it includes external forces as well as the forces originating from the rest of the body. We can neglect internal contributions only under special assumptions. Generally, any such assumptions become invalid when we let the body shrink to sufficiently small dimensions. Classical electromagnetism predicts that below some distance internal forces turn out to be larger than any external forces. But in addition to this objection, there is a second one. We have seen that $\mathbf{p}_h = \mathbf{m} \times \mathbf{E}/c^2$ applies for an external force-density given by $\mathbf{f} = -\rho \nabla \phi$. In Mansuripur's case, however, \mathbf{f} is given (in the notation of [6]) by $\mathbf{f} = -qm_0 v d \left(4\pi\epsilon_0 c^2 R^3\right)^{-1} \delta(x) \delta'(y) \delta\left[z - Vt - d/\gamma\right] \hat{\mathbf{z}},$ with $R = (x^2 + y^2 + (z - Vt)^2)^{1/2}$. Using this **f** in $\mathbf{p}_h = -c^{-2} \int \mathbf{r} (\mathbf{f} \cdot \mathbf{v}) d^3 r$ we do not obtain the desired results. Thus, as stressed by Boyer [14], hidden momentum can lend itself to "explanations of dubious validity or outright error which avoid needed discussions of energy and momentum flow". Such discussions can be carried out when dealing with models that fall in line with the tenets of classical electromagnetism. This is not to say that classical electromagnetism is a complete, selfconsistent theory of electromagnetic phenomena. Even situations that are much simpler than the ones we have discussed so far, do require that we resort to quantum mechanics. Consider for example a point charge q sitting at rest at the origin O of the xyz-system. In accordance with Maxwell's equations, q is the source of a purely electrostatic field, the Coulomb field. Consider now the same situation as seen from the perspective of the x'y'z'-system, whose origin coincides with O while its

axes rotate with respect to the xyz-system. Although the exact space-time transformation linking the two reference frames (in contrast to the *coordinate systems*, t'x'y'z' and txyz) is unknown, we can certainly take for granted that classical electromagnetism predicts that in the rotating frame a magnetic field will be observed, together with an electric field. In other words, in the x'y'z'-system q would be the source of an electric and a magnetic field. Now, given a true *point* charge sitting at rest, how could we tell whether it is rotating or not? We cannot have a consistent model of elementary charges at rest, unless we provide all of them with the property of being sources of both electric and magnetic fields. This is precisely what quantum mechanics does by endowing the electron – and other elementary particles – with a spin. The quantum mechanical electron is thus a source of both electric and magnetic fields. At the same time, even being at rest the pointlike electron can be set in "rotation" by an external magnetic field. But now "rotation" acquires a physical meaning: the observable spin dynamics.

In conclusion, Mansuripur's claim that the Lorentz law should be abandoned seems to be unsubstantiated. The mere existence of a torque that appears in one inertial frame and not in the other does not violate relativity, as it has been illustrated by Jackson's example [2]. Mansuripur's results would rather illustrate the limits of classical models, when it comes to describe some electromagnetic features.

- [1] M. Mansuripur, Phys. Rev. Lett. 108, 193901 (2012).
- [2] J. D. Jackson, Am. J. Phys. 72, 1484 (2004).
- [3] J. D. Jackson, *Classical Electrodynamics* (Wiley, NY, 1999), 3rd ed.
- [4] In physics, as in mathematics, a limiting procedure can bring us out from a given framework. E.g., any finite sum of the form $r_n = 4 - 4/3 + 4/5 - \ldots (-1)^n 4/(2n+1)$ is a rational number $(r_n \in \mathbb{Q})$. However, $\lim_{n\to\infty} r_n = \pi \notin \mathbb{Q}$. Similarly, Dirac's delta, being not a regular function but a distribution, is a limiting case of regular Gaussian functions. Whether such qualitative changes are relevant or not for the corresponding physical models, is something that must be decided in each particular case.
- [5] D. T. A. Vanzella, arXiv:1205.1502
- [6] D. J. Griffiths and V. Hnizdo, arXiv:1205.4646
- [7] P. L. Saldanha, arXiv:1205.6858
- [8] M. Brachet and E. Tirapegui, arXiv:1207.4613
- [9] D. J. Griffiths, Introduction to Electrodynamics (Prentice-Hall, Upper Saddle River, NJ, 1999), 3rd ed.
- [10] W. Shockley and R. P. James, Phys. Rev. Lett. 18, 876 (1967).
- [11] W. H. Furry, Am. J. Phys. 37, 621 (1969).
- [12] S. Coleman and J. H. Van Vleck, Phys. Rev. 171, 1370 (1968).
- [13] V. Hnizdo, Am. J. Phys. 65, 516 (1997).
- [14] T. H. Boyer, Am. J. Phys. 76, 190 (2008).