

SPECTRUM OF THE BURNSIDE TAMBARA FUNCTOR ON A CYCLIC p -GROUP

HIROYUKI NAKAOKA

ABSTRACT. For a finite group G , a Tambara functor on G is regarded as a G -bivariant analog of a commutative ring. In this analogy, previously we have defined an ideal of a Tambara functor. In this article, we will demonstrate a calculation of the prime spectrum of the Burnside Tambara functor, when G is a cyclic p -group for a prime integer p .

1. INTRODUCTION AND PRELIMINARIES

A *Tambara functor* on a finite group G is firstly defined by Tambara [11] in the name ‘TNR-functor’, to treat the multiplicative transfers of Green functors. The terminology *Tambara functor* firstly appeared in Brun’s paper, when he used it to describe the structure of Witt-Burnside rings [2]. It consists of a triplet $T = (T^*, T_+, T_\bullet)$, where the *additive part* (T^*, T_+) forms a Mackey functor, whereas the *multiplicative part* (T^*, T_\bullet) forms a semi-Mackey functor. This is just like a commutative ring consists of an additive abelian group structure and a multiplicative commutative semi-group structure. In fact a Tambara functor is nothing other than a commutative ring when G is trivial. In this sense, this notion is regarded as a Mackey-functor-theoretic analog (or, ‘ G -bivariant analog’) of a commutative ring [14].

In this analogy, some algebraic notions in commutative ring theory find their analogs in Tambara functor theory. We have ideals [5], fractions [4], and polynomials [6] of Tambara functors. These are mutually related, as they should be, and moreover in connection with the celebrated Dress construction [10], [7]. The most typical Tambara functor is the *Burnside Tambara functor* Ω_G (Example 1.4), which plays a role just like \mathbb{Z} in the ordinary commutative ring theory.

Especially, in the G -bivariant analog of the ideal theory, the prime spectrum $\text{Spec } T$ has been defined for any Tambara functor T . In our previous short note [8], we have demonstrated a calculation of the prime spectrum of Ω_G , when G is a group of prime order p . In this article, extending the method in [8], we calculate $\text{Spec } \Omega_G$ when G is a *cyclic p -group* for a prime integer p . To determine the prime ideals of Ω_G , the key observation is that the set of subgroups of G is totally ordered

$$e = H_0 < H_1 < \cdots < H_r = G,$$

The author wishes to thank Professor Fumihito Oda for his suggestions and useful comments. The author also wishes to thank Professor Akihiko Hida for his question and comments. This work is supported by JSPS Grant-in-Aid for Young Scientists (B) 22740005.

and thus Ω_G can be regarded as a sequence of commutative rings equipped with adjacent structure morphisms

$$R_0 \begin{array}{c} \xrightarrow{\text{ind}} \\ \xleftarrow{\text{res}} \\ \xrightarrow{\text{jnd}} \end{array} R_1 \begin{array}{c} \xrightarrow{\text{ind}} \\ \xleftarrow{\text{res}} \\ \xrightarrow{\text{jnd}} \end{array} R_2 \begin{array}{c} \xrightarrow{\text{ind}} \\ \xleftarrow{\text{res}} \\ \xrightarrow{\text{jnd}} \end{array} \cdots \begin{array}{c} \xrightarrow{\text{ind}} \\ \xleftarrow{\text{res}} \\ \xrightarrow{\text{jnd}} \end{array} R_k \begin{array}{c} \xrightarrow{\text{ind}} \\ \xleftarrow{\text{res}} \\ \xrightarrow{\text{jnd}} \end{array} \cdots \begin{array}{c} \xrightarrow{\text{ind}} \\ \xleftarrow{\text{res}} \\ \xrightarrow{\text{jnd}} \end{array} R_r,$$

where $R_k = \Omega_G(G/H_k)$. With this identification, an ideal of Ω_G can be regarded as a sequence $[I_0, \dots, I_r]$ of ideals $I_k \subseteq R_k$. A sequence $[I_0, \dots, I_r]$ forms an ideal of Ω_G if and only if the condition

$$\boxed{\mathcal{I}(k)} \quad \text{ind}_{k-1}^k(I_{k-1}) \subseteq I_k, \quad \text{res}_{k-1}^k(I_k) \subseteq I_{k-1}, \quad \text{jnd}_{k-1}^k(I_{k-1}) \subseteq I_k.$$

is satisfied for each $1 \leq k \leq r$ (Corollary 3.3). The restriction of an ideal $\mathcal{I} = [I_0, \dots, I_k]$ of Ω_{H_k} onto H_i is given by $\mathcal{I}|_{H_i} = [I_0, \dots, I_i]$ for any $0 \leq i \leq k$. On the contrary, we can consider an extension of \mathcal{I} onto H_{k+1} , namely, an ideal \mathcal{I}' of $\Omega_{H_{k+1}}$ satisfying $\mathcal{I}'|_{H_k} = \mathcal{I}$. In particular, the largest and the smallest among such \mathcal{I}' are explicitly given by $\mathcal{L}\mathcal{I}$ and $\mathcal{S}\mathcal{I}$ in Definition 3.5. This allows us an inductive construction of ideals in Ω_G .

Whether an ideal $\mathcal{I} = [I_0, \dots, I_r]$ is prime or not can be also checked inductively on k . In fact, it is prime if and only if the condition

$$\boxed{\mathcal{P}(k)} \quad \text{For any } 0 \leq i \leq k, a \in (\text{res}_{k-1}^k)^{-1}(I_{k-1}) \text{ and } b \in (\text{res}_{i-1}^i)^{-1}(I_{i-1}) \setminus I_i,$$

$$a \cdot \text{jnd}_i^k(b) \in I_k \implies a \in I_k$$

holds.

is satisfied for each $0 \leq k \leq r$ (Proposition 4.4). Consequently, any restriction $\mathcal{I}|_{H_i}$ of a prime ideal $\mathcal{I} = [I_0, \dots, I_k] \subseteq \Omega_{H_k}$ is again prime. Especially I_0 should be a prime ideal in $R_0 \cong \mathbb{Z}$, and thus equal to 0 , (p) , or (q) for some prime integer $q \neq p$. By determining prime ideals over $0, (p), (q)$, namely, prime ideals $\mathcal{I} = [I_0, \dots, I_r]$ satisfying $I_0 = 0, (p), (q)$ respectively, we obtain the following result. In particular, the dimension of Ω_G is calculated as $\dim \Omega_G = r + 1$ (Corollary 6.13).

Theorem 1.1. *Let G be a cyclic p -group of p -rank r . The prime ideals of $\Omega_G \in \text{Ob}(\text{Tam}(G))$ are as follows.*

(i) *Over $(q) \subseteq R_0$, there are $\ell + 1$ prime ideals*

$$\mathcal{S}^r(q) \subsetneq \mathcal{L}\mathcal{S}^{r-1}(q) \subsetneq \cdots \subsetneq \mathcal{L}^r(q).$$

(ii) *Over $(p) \subseteq R_0$, there is only one prime ideal $\mathcal{L}^r(p)$.*

(iii) *Over $0 \subseteq R_0$, there are $\ell + 1$ prime ideals*

$$0 \subsetneq \mathcal{L}(0) \subsetneq \cdots \subsetneq \mathcal{L}^r(0).$$

Thus we have

$$\begin{aligned} \text{Spec } \Omega_G &= \{\mathcal{L}^r(p)\} \cup \{\mathcal{L}^i(0) \mid 0 \leq i \leq r\} \\ &\cup \{\mathcal{L}^i\mathcal{S}^{r-i}(q) \mid 0 \leq i \leq r, q \text{ is a prime different from } p\}. \end{aligned}$$

Throughout this article, the unit of a finite group G will be denoted by e . Abbreviately we denote the trivial subgroup of G by e , instead of $\{e\}$. $H \leq G$ means H is a subgroup of G . ${}_G\text{set}$ denotes the category of finite G -sets and G -equivariant maps. If $H \leq G$ and $g \in G$, then gH denotes the conjugate ${}^gH = gHg^{-1}$. Similarly, $H^g = g^{-1}Hg$. A monoid is always assumed to be unitary and commutative.

Similarly a ring is assumed to be commutative, with an additive unit 0 and a multiplicative unit 1. We denote the category of monoids by Mon , the category of rings by $Ring$. A monoid homomorphism preserves units, and a ring homomorphism preserves 0 and 1. For any category \mathcal{C} and any pair of objects X and Y in \mathcal{C} , the set of morphisms from X to Y in \mathcal{C} is denoted by $\mathcal{C}(X, Y)$.

We briefly recall the definition of a Tambara functor.

Definition 1.2. ([11]) A *Tambara functor* T on G is a triplet $T = (T^*, T_+, T_\bullet)$ of two covariant functors

$$T_+ : {}_G set \rightarrow Set, \quad T_\bullet : {}_G set \rightarrow Set$$

and one contravariant functor

$$T^* : {}_G set \rightarrow Set$$

which satisfies the following. Here Set is the category of sets.

- (1) $T^\alpha = (T^*, T_+)$ is a Mackey functor on G .
- (2) $T^\mu = (T^*, T_\bullet)$ is a semi-Mackey functor on G .
Since T^α, T^μ are semi-Mackey functors, we have $T^*(X) = T_+(X) = T_\bullet(X)$ for each $X \in \text{Ob}({}_G set)$. We denote this by $T(X)$.
- (3) (Distributive law) If we are given an exponential diagram

$$\begin{array}{ccccc} X & \xleftarrow{p} & A & \xleftarrow{\lambda} & Z \\ f \downarrow & & exp & & \downarrow \rho \\ Y & \xleftarrow{q} & & & B \end{array}$$

in ${}_G set$, then

$$\begin{array}{ccccc} T(X) & \xleftarrow{T_+(p)} & T(A) & \xrightarrow{T^*(\lambda)} & T(Z) \\ T_\bullet(f) \downarrow & & \circ & & \downarrow T_\bullet(\rho) \\ T(Y) & \xleftarrow{T_+(q)} & & & T(B) \end{array}$$

is commutative. For the definition and basic properties of exponential diagrams, see [11].

If $T = (T^*, T_+, T_\bullet)$ is a Tambara functor, then $T(X)$ becomes a ring for each $X \in \text{Ob}({}_G set)$ ([11]). Since T^α is a Mackey functor, by definition T is ‘additive’, in the sense that for any $X_1, X_2 \in \text{Ob}({}_G set)$, the inclusions $\iota_1 : X_1 \hookrightarrow X_1 \amalg X_2$ and $\iota_2 : X_2 \hookrightarrow X_1 \amalg X_2$ induce a natural isomorphism of rings $(T^*(\iota_1), T^*(\iota_2)) : T(X_1 \amalg X_2) \xrightarrow{\cong} T(X_1) \times T(X_2)$. For each $f \in {}_G set(X, Y)$,

- $T^*(f) : T(Y) \rightarrow T(X)$ is a ring homomorphism, called the *restriction* along f ,
- $T_+(f) : T(X) \rightarrow T(Y)$ is an additive homomorphism, called the *additive transfer* along f ,
- $T_\bullet(f) : T(X) \rightarrow T(Y)$ is a multiplicative homomorphism, called the *multiplicative transfer* along f .

$T^*(f), T_+(f), T_\bullet(f)$ are often abbreviated to f^*, f_+, f_\bullet .

Remark 1.3. If f is the natural projection $\text{pr}_K^H: G/K \rightarrow G/H$ for some $K \leq H \leq G$, then f^*, f_+, f_\bullet is written as

$$\begin{aligned} \text{res}_K^H &= (\text{pr}_K^H)^* , \\ \text{ind}_K^H &= (\text{pr}_K^H)_+ , \\ \text{jnd}_K^H &= (\text{pr}_K^H)_\bullet . \end{aligned}$$

For a conjugate map $c_g: G/H^g \rightarrow G/H$, we define $c_{g,H}: T(G/H) \rightarrow T(G/H^g)$ (or simply $c_g: T(G/H) \rightarrow T(G/H^g)$) by

$$c_{g,H} = T^*(c_g).$$

If g belongs to the normalizer $N_G(H)$ of H in G , then this gives an automorphism $c_{g,H}: T(G/H) \rightarrow T(G/H)$. With this $N_G(H)$ -action, every $T(G/H)$ becomes an $N_G(H)/H$ -ring.

Since any G -map is a union of compositions of natural projections and conjugate maps, the structure morphisms of a Tambara functor are completely determined by $\text{res}_K^H, \text{ind}_K^H, \text{jnd}_K^H, c_{g,H}$ for $K \leq H \leq G, g \in G$, by virtue of the additivity.

Example 1.4. If we define Ω_G by

$$\Omega_G(X) = K_0(\text{Gset}/X)$$

for each $X \in \text{Ob}(\text{Gset})$, where the right hand side is the Grothendieck ring of the category of finite G -sets over X , then Ω_G becomes a Tambara functor on G [11]. This is called the *Burnside Tambara functor*. For each $f \in \text{Gset}(X, Y)$,

$$f_\bullet: \Omega_G(X) \rightarrow \Omega_G(Y)$$

is the one determined by

$$f_\bullet(A \xrightarrow{p} X) = (\Pi_f(A) \xrightarrow{\varpi} Y) \quad (\forall (A \xrightarrow{p} X) \in \text{Ob}(\text{Gset}/X)),$$

where $\Pi_f(A)$ and ϖ is

$$\Pi_f(A) = \left\{ (y, \sigma) \left| \begin{array}{l} y \in Y, \\ \sigma: f^{-1}(y) \rightarrow A \text{ is a map of sets,} \\ p \circ \sigma = \text{id}_{f^{-1}(y)} \end{array} \right. \right\},$$

$$\varpi(y, \sigma) = y.$$

G acts on $\Pi_f(A)$ by $g \cdot (y, \sigma) = (gy, {}^g\sigma)$, where ${}^g\sigma$ is the map defined by

$${}^g\sigma(x) = g\sigma(g^{-1}x) \quad (\forall x \in f^{-1}(gy)).$$

$f_+: \Omega_G(X) \rightarrow \Omega_G(Y)$ is an additive homomorphism satisfying

$$f_+(A \xrightarrow{p} X) = (A \xrightarrow{f \circ p} Y) \quad (\forall (A \xrightarrow{p} X) \in \text{Ob}(\text{Gset}/X)).$$

f^* is defined by using a fiber product ([11]). Namely, the Mackey-functor structure on the additive part of Ω is the usual one as in [1].

Remark 1.5. For any $K \leq H \leq G$, we have a natural isomorphism (cf. [1]) $\Omega_G(G/K) \cong \Omega_H(H/K)$, and we will identify them through this isomorphism.

We often abbreviate Ω_G to Ω , if the base group is obvious from the context.

2. BURNSIDE TAMBARA FUNCTOR ON A CYCLIC p -GROUP

Throughout this article, we fix a prime number p . Let G be a cyclic p -group of p -rank $r \geq 0$, and let $H_k \leq G$ be its subgroup of order p^k for each $0 \leq k \leq r$. In the following argument, without loss of generality we may assume

$$\begin{aligned} G &= \mathbb{Z}/p^r\mathbb{Z}, & H_k &= p^{r-k}\mathbb{Z}/p^r\mathbb{Z}, \\ e &= H_0 < H_1 \cdots < H_k < \cdots < H_r = G. \end{aligned}$$

Then the G -set G/H_k is canonically isomorphic to $\mathbb{Z}/p^{r-k}\mathbb{Z}$, and the natural projection

$$G/H_k \xrightarrow{\text{pr}_k^\ell} G/H_\ell \quad (k \leq \ell)$$

is identified with a map given by

$$\mathbb{Z}/p^{r-k}\mathbb{Z} \rightarrow \mathbb{Z}/p^{r-\ell}\mathbb{Z}; \quad a \bmod p^{r-k}\mathbb{Z} \mapsto a \bmod p^{r-\ell}\mathbb{Z}$$

for any $a \in \mathbb{Z}$.

Since G is commutative, each $\Omega(G/H_k)$ has a trivial G -action, and admits a natural \mathbb{Z} -basis

$$\{(G/H_i \xrightarrow{\text{pr}_i^k} G/H_k) \mid 0 \leq i \leq k\}.$$

Thus if we denote $(G/H_i \xrightarrow{\text{pr}_i^k} G/H_k)$ by $X_{k,i}$, then it is a free \mathbb{Z} -module

$$\Omega(G/H_k) = \bigoplus_{0 \leq i \leq k} \mathbb{Z}X_{k,i}$$

with a trivial G -action. Therefore, if we put $R_k = \bigoplus_{0 \leq i \leq k} \mathbb{Z}X_{k,i}$, then the Tambara

functor Ω_G is regarded just as a sequence of commutative rings R_k and structure morphisms

$$R_0 \begin{array}{c} \xrightarrow{\text{ind}_0^1} \\ \xleftarrow{\text{res}_0^1} \\ \xrightarrow{\text{jnd}_0^1} \end{array} R_1 \begin{array}{c} \xrightarrow{\text{ind}_1^2} \\ \xleftarrow{\text{res}_1^2} \\ \xrightarrow{\text{jnd}_1^2} \end{array} R_2 \begin{array}{c} \xrightarrow{\text{ind}_2^3} \\ \xleftarrow{\text{res}_2^3} \\ \xrightarrow{\text{jnd}_2^3} \end{array} \cdots \begin{array}{c} \xrightarrow{\text{ind}_{k-1}^k} \\ \xleftarrow{\text{res}_{k-1}^k} \\ \xrightarrow{\text{jnd}_{k-1}^k} \end{array} R_k \begin{array}{c} \xrightarrow{\text{ind}_k^{k+1}} \\ \xleftarrow{\text{res}_k^{k+1}} \\ \xrightarrow{\text{jnd}_k^{k+1}} \end{array} \cdots \begin{array}{c} \xrightarrow{\text{ind}_{r-1}^r} \\ \xleftarrow{\text{res}_{r-1}^r} \\ \xrightarrow{\text{jnd}_{r-1}^r} \end{array} R_r$$

satisfying conditions in Definition 1.2. Here, $\text{ind}_{k-1}^k, \text{res}_{k-1}^k, \text{jnd}_{k-1}^k$ are the abbreviations of $\text{ind}_{H_{k-1}}^{H_k}, \text{res}_{H_{k-1}}^{H_k}, \text{jnd}_{H_{k-1}}^{H_k}$. We use similar abbreviations in the rest. Remark that any structure morphism of Ω_G can be realized as a composition of these morphisms.

Remark 2.1. By virtue of Remark 1.5, the first k -terms

$$(2.1) \quad R_0 \begin{array}{c} \xrightarrow{\text{ind}_0^1} \\ \xleftarrow{\text{res}_0^1} \\ \xrightarrow{\text{jnd}_0^1} \end{array} R_1 \begin{array}{c} \xrightarrow{\text{ind}_1^2} \\ \xleftarrow{\text{res}_1^2} \\ \xrightarrow{\text{jnd}_1^2} \end{array} R_2 \begin{array}{c} \xrightarrow{\text{ind}_2^3} \\ \xleftarrow{\text{res}_2^3} \\ \xrightarrow{\text{jnd}_2^3} \end{array} \cdots \begin{array}{c} \xrightarrow{\text{ind}_{k-1}^k} \\ \xleftarrow{\text{res}_{k-1}^k} \\ \xrightarrow{\text{jnd}_{k-1}^k} \end{array} R_k$$

can be regarded as a sequence representing the Tambara functor Ω_{H_k} . We always work under this identification in this paper. With this identification, forgetting the entire group G , we can regard the Burnside Tambara functor Ω_{H_k} on a cyclic

p -group H_k of p -rank k , simply as a length k sequence of rings (2.1) obtained inductively by adding $R_k, \text{ind}_{k-1}^k, \text{res}_{k-1}^k, \text{jnd}_{k-1}^k$ to the length $k-1$ sequence

$$R_0 \begin{array}{c} \xrightarrow{\text{ind}_0^1} \\ \xleftarrow{\text{res}_0^1} \\ \xrightarrow{\text{jnd}_0^1} \end{array} R_1 \begin{array}{c} \xrightarrow{\text{ind}_1^2} \\ \xleftarrow{\text{res}_1^2} \\ \xrightarrow{\text{jnd}_1^2} \end{array} \cdots \begin{array}{c} \xrightarrow{\text{ind}_{k-2}^{k-1}} \\ \xleftarrow{\text{res}_{k-2}^{k-1}} \\ \xrightarrow{\text{jnd}_{k-2}^{k-1}} \end{array} R_{k-1}$$

corresponding to $\Omega_{H_{k-1}}$. This observation enables us an inductive construction of ideals of in Ω .

We go on to describe R_k and the structure morphisms. As above, we have $R_k = \bigoplus_{0 \leq i \leq k} \mathbb{Z}X_{k,i}$ as a module.

Remark 2.2. Additive transfer $\text{ind}_k^\ell: R_k \rightarrow R_\ell$ is given by

$$\text{ind}_k^\ell(X_{k,i}) = X_{\ell,i} \quad (0 \leq i \leq k)$$

for each $k \leq \ell$.

Proof. This is obvious. □

If we take a fiber product of $\text{pr}_i^k: G/H_i \rightarrow G/H_k$ and $\text{pr}_j^k: G/H_j \rightarrow G/H_k$

$$\begin{array}{ccc} G/H_i \times_{G/H_k} G/H_j & \twoheadrightarrow & G/H_j \\ \downarrow & \square & \downarrow \text{pr}_j^k \\ G/H_i & \xrightarrow{\text{pr}_i^k} & G/H_k \end{array}$$

where $0 \leq i, j \leq k$, then by the Mackey decomposition formula, we have

$$(2.2) \quad G/H_i \times_{G/H_k} G/H_j = \begin{cases} \coprod_{p^{k-j}} G/H_i & (i \leq j) \\ \coprod_{p^{k-i}} G/H_j & (j \leq i). \end{cases}$$

In any case, the cardinality of this G -set is

$$|G/H_i \times_{G/H_k} G/H_j| = p^{r+k-(i+j)}.$$

Equation (2.2) is also written as

$$G/H_i \times_{G/H_k} G/H_j = \coprod_{p^{k-\max(i,j)}} G/H_{\min(i,j)}.$$

As a corollary, we obtain the following.

Corollary 2.3. *For any $0 \leq k \leq r$, the following holds.*

(1) *For any $i, j \leq k$,*

$$X_{k,i} \cdot X_{k,j} = \begin{cases} p^{k-j} X_{k,i} & (i \leq j) \\ p^{k-i} X_{k,j} & (j \leq i) \end{cases}$$

in R_k . In fact, as a residue ring of the polynomial ring $\mathbb{Z}[X_{k,i} \mid 0 \leq i \leq k]$ over indeterminates $X_{k,i}$ ($0 \leq i \leq k$), R_k is written as

$$\mathbb{Z}[X_{k,i} \mid 0 \leq i \leq k] / (\{X_{k,i}X_{k,j} - p^{k-\max(i,j)}X_{k,\min(i,j)} \mid 0 \leq i, j \leq k\}).$$

In particular, $X_{k,k} = 1$ is the unit of R_k .

(2) For any $\ell \geq k$, $\text{res}_k^\ell: R_\ell \rightarrow R_k$ is given by

$$\text{res}_k^\ell(X_{\ell,i}) = \begin{cases} p^{\ell-k} X_{k,i} & (i \leq k) \\ p^{\ell-i} X_{k,k} = p^{\ell-i} & (i \geq k). \end{cases}$$

As for multiplicative transfers, we need some calculation. For the detail, see the Appendix.

Corollary 2.4. For any $\ell \geq k$, $\text{jnd}_k^\ell: R_k \rightarrow R_\ell$ is given by

$$\begin{aligned} \text{jnd}_k^\ell\left(\sum_{0 \leq i \leq k} m_i X_{k,i}\right) &= m_k X_{\ell,\ell} + \sum_{k \leq i < \ell} \frac{(m_k)^{p^{\ell-i}} - (m_k)^{p^{\ell-i-1}}}{p^{\ell-i}} X_{\ell,i} \\ &+ \sum_{0 \leq i < k} \frac{(\sum_{s=i}^k m_s p^{k-s})^{p^{\ell-k}} - (\sum_{s=i+1}^k m_s p^{k-s})^{p^{\ell-k}}}{p^{\ell-i}} X_{\ell,i} \end{aligned}$$

for any $m_0, \dots, m_k \in \mathbb{Z}$.

3. IDEALS OF Ω AS A SEQUENCE

An ideal of a Tambara functor is defined in [5] as follows.

Definition 3.1. Let T be a Tambara functor. An *ideal* \mathcal{I} of T is a family of ideals $\{\mathcal{I}(X) \subseteq T(X)\}_{X \in \text{Ob}(G\text{set})}$ satisfying

- (i) $f^*(\mathcal{I}(Y)) \subseteq \mathcal{I}(X)$,
- (ii) $f_+(\mathcal{I}(X)) \subseteq \mathcal{I}(Y)$,
- (iii) $f_\bullet(\mathcal{I}(X)) \subseteq f_\bullet(0) + \mathcal{I}(Y)$

for any $f \in G\text{set}(X, Y)$.

These conditions also imply

$$\mathcal{I}(X_1 \amalg X_2) \cong \mathcal{I}(X_1) \times \mathcal{I}(X_2)$$

for any $X_1, X_2 \in \text{Ob}(G\text{set})$. From this, an ideal of T is determined by a family

$$\{I_H = \mathcal{I}(G/H)\}_{H \leq G}$$

of ideals $I_H \subseteq T(G/H)$, indexed by the set of subgroups of G , as follows.

Proposition 3.2. Let T be a Tambara functor on G . To give an ideal \mathcal{I} of T is equivalent to give a family $\{I_H\}_{H \leq G}$ of ideals $I_H \subseteq T(G/H)$ satisfying the conditions

- (i) $\text{res}_K^H(I_H) \subseteq I_K$
- (ii) $\text{ind}_K^H(I_K) \subseteq I_H$
- (iii) $\text{jnd}_K^H(I_K) \subseteq I_H$
- (iv) $c_{g,H}(I_H) \subseteq I_{H^g}$

for any $K \leq H \leq G$ and $g \in G$.

Proof. This is straightforward (cf. Corollary 2.2 in [8]). □

In our particular case, an ideal of Ω_G is written as follows.

Corollary 3.3. Let G be a cyclic p -group of p -rank r . An ideal \mathcal{I} of Ω_G is given by a sequence

$$\mathcal{I} = [I_0, \dots, I_r]$$

of ideals $I_k \subseteq R_k$, satisfying the following condition $\mathcal{I}(k)$ for each $1 \leq k \leq r$.

$$\boxed{\mathcal{I}(k)} \quad \text{ind}_{k-1}^k(I_{k-1}) \subseteq I_k, \quad \text{res}_{k-1}^k(I_k) \subseteq I_{k-1}, \quad \text{jnd}_{k-1}^k(I_{k-1}) \subseteq I_k.$$

Remark that for ideals $\mathcal{I} = [I_0, \dots, I_r]$ and $\mathcal{J} = [J_0, \dots, J_r]$, we have $\mathcal{I} \subseteq \mathcal{J}$ if and only if $I_k \subseteq J_k$ holds for any $0 \leq k \leq r$.

In the following, we will simply say “[I_0, \dots, I_k] is an ideal of Ω_{H_k} ” to mean that [I_0, \dots, I_k] satisfies $\mathcal{I}(i)$ for any $1 \leq i \leq k$. This makes sense by virtue of Remark 2.1. The restriction of an ideal $\mathcal{I} = [I_0, \dots, I_k]$ of Ω_{H_k} onto H_i is given by [I_0, \dots, I_i], for each $0 \leq i \leq k$. We denote this by $\mathcal{I}|_{H_i}$. Obviously, restriction preserves inclusions of ideals.

On the contrary, Corollary 3.3 also enables us to extend an ideal [I_0, \dots, I_{k-1}] of $\Omega_{H_{k-1}}$ to an ideal [I_0, \dots, I_k] of Ω_{H_k} , by adding an ideal $I_k \subseteq R_k$ satisfying $\mathcal{I}(k)$. Among the possible extensions of [I_0, \dots, I_{k-1}], the largest and the smallest are given as follows. If an ideal $\mathcal{I}' \subseteq \Omega_{H_k}$ satisfies $\mathcal{I}'|_{H_i} = \mathcal{I}$ for a given ideal $\mathcal{I} \subseteq \Omega_{H_i}$, then we say that \mathcal{I}' is an ideal over \mathcal{I} .

Proposition 3.4. *For a fixed $1 \leq k \leq r$, suppose we are given an ideal $\mathcal{I} = [I_0, \dots, I_{k-1}]$ of $\Omega_{H_{k-1}}$.*

(1) *If we define $L_k(I_{k-1}) = L(I_{k-1}) \subseteq R_k$ by*

$$L(I_{k-1}) = (\text{res}_{k-1}^k)^{-1}(I_{k-1}),$$

then $[I_0, \dots, I_{k-1}, L(I_{k-1})]$ is the largest ideal of Ω_{H_k} over $[I_0, \dots, I_{k-1}]$.

(2) *If we define $S_k(I_{k-1}) = S(I_{k-1}) \subseteq R_k$ to be the ideal of R_k generated by*

$$\{\text{ind}_{k-1}^k(\alpha) \mid \alpha \in I_{k-1}\} \quad \text{and} \quad \{\text{jnd}_{k-1}^k(\alpha) \mid \alpha \in I_{k-1}\},$$

shortly,

$$S(I_{k-1}) = (\text{ind}_{k-1}^k(I_{k-1})) + (\text{jnd}_{k-1}^k(I_{k-1})),$$

then $[I_0, \dots, I_{k-1}, S(I_{k-1})]$ is the smallest ideal of Ω_{H_k} over $[I_0, \dots, I_{k-1}]$.

Definition 3.5. For an ideal $\mathcal{I} = [I_0, \dots, I_{k-1}]$ of $\Omega_{H_{k-1}}$, define ideals $\mathcal{L}_k(\mathcal{I}) = \mathcal{L}\mathcal{I}$ and $\mathcal{S}_k(\mathcal{I}) = \mathcal{S}\mathcal{I}$ of Ω_{H_k} by

$$\begin{aligned} \mathcal{L}\mathcal{I} &= [I_0, \dots, I_{k-1}, L(I_{k-1})], \\ \mathcal{S}\mathcal{I} &= [I_0, \dots, I_{k-1}, S(I_{k-1})], \end{aligned}$$

where $L(I_{k-1})$ and $S(I_{k-1})$ are those in Proposition 3.4. Additionally, we denote the n -times iterations of $L, S, \mathcal{L}, \mathcal{S}$ by L^n, S^n and $\mathcal{L}^n, \mathcal{S}^n$. For example, an ideal $\mathcal{I} = [I_0, \dots, I_k] \subseteq \Omega_{H_k}$ yields $\mathcal{L}^n \mathcal{I} = [I_0, \dots, I_k, L(I_k), \dots, L^n(I_k)] \subseteq \Omega_{H_{k+n}}$.

Proof of Proposition 3.4. (1) To show $\mathcal{L}\mathcal{I}$ is an ideal of Ω_{H_k} , it suffices to confirm $\mathcal{I}(k)$ is satisfied. By definition, $\text{res}_{k-1}^k(L(I_{k-1})) \subseteq I_{k-1}$ is obvious. In addition, by the existence of a pullback diagram

$$\begin{array}{ccc} \prod_p G/H_{k-1} & \xrightarrow{\nabla} & G/H_{k-1} \\ \downarrow & \square & \downarrow \text{pr}_{k-1}^k \\ G/H_{k-1} & \xrightarrow{\text{pr}_{k-1}^k} & G/H_k \end{array}$$

where $\nabla: \amalg_p G/H_{k-1} \rightarrow G/H_{k-1}$ is the folding map, we have

$$\begin{aligned}\text{res}_{k-1}^k \circ \text{ind}_{k-1}^k(\alpha) &= p\alpha, \\ \text{res}_{k-1}^k \circ \text{jnd}_{k-1}^k(\alpha) &= \alpha^p\end{aligned}$$

for any $\alpha \in R_k$. Thus if $\alpha \in I_{k-1}$, then

$$\begin{aligned}\text{ind}_{k-1}^k(\alpha) &\in (\text{res}_{k-1}^k)^{-1}(I_{k-1}), \\ \text{jnd}_{k-1}^k(\alpha) &\in (\text{res}_{k-1}^k)^{-1}(I_{k-1})\end{aligned}$$

holds. This shows $\text{ind}_{k-1}^k(I_{k-1}) \subseteq L(I_{k-1})$ and $\text{jnd}_{k-1}^k(I_{k-1}) \subseteq L(I_{k-1})$, and thus $\mathcal{L}\mathcal{S}$ is an ideal of Ω_{H_k} . Moreover, since any ideal $[I_0, \dots, I_{k-1}, J_k]$ of Ω_{H_k} should satisfy $\text{res}_{k-1}^k(J_k) \subseteq I_{k-1}$, obviously $\mathcal{L}\mathcal{S}$ is the largest.

(2) It suffices to confirm $\mathcal{S}\mathcal{S}$ satisfies $\mathcal{I}(k)$. Since res_{k-1}^k is a ring homomorphism, $\text{res}_{k-1}^k(S(I_{k-1})) \subseteq I_{k-1}$ follows from the fact that any $\alpha \in I_{k-1}$ satisfies

$$\begin{aligned}\text{res}_{k-1}^k \circ \text{ind}_{k-1}^k(\alpha) &= p\alpha \in I_{k-1}, \\ \text{res}_{k-1}^k \circ \text{jnd}_{k-1}^k(\alpha) &= \alpha^p \in I_{k-1}.\end{aligned}$$

The other conditions

$$\begin{aligned}\text{ind}_{k-1}^k(I_{k-1}) &\subseteq S(I_{k-1}), \\ \text{jnd}_{k-1}^k(I_{k-1}) &\subseteq S(I_{k-1})\end{aligned}$$

are obviously satisfied by the definition of $S(I_{k-1})$. Thus $\mathcal{S}\mathcal{S}$ is an ideal of Ω_{H_k} . Moreover, since any ideal $[I_0, \dots, I_{k-1}, J_k]$ of Ω_{H_k} should satisfy $\text{ind}_{k-1}^k(I_{k-1}) \subseteq J_k$ and $\text{jnd}_{k-1}^k(I_{k-1}) \subseteq J_k$, obviously $\mathcal{S}\mathcal{S}$ is the smallest. \square

4. INDUCTIVE CRITERION OF PRIMENESS

In [5], a prime ideal of a Tambara functor is defined as follows.

Definition 4.1. Let G be an arbitrary finite group, and let T be a Tambara functor on G . An ideal $\mathcal{S} \subsetneq T$ is prime if and only if the following two conditions become equivalent, for any transitive $X, Y \in \text{Ob}({}_G\text{set})$ and any $a \in T(X)$, $b \in T(Y)$.

(1) For any $C \in \text{Ob}({}_G\text{set})$ and for any pair of diagrams

$$C \xleftarrow{v} D \xrightarrow{w} X, \quad C \xleftarrow{v'} D' \xrightarrow{w'} Y$$

in ${}_G\text{set}$, $\mathcal{S}(C)$ satisfies

$$(v_\bullet w^*(a) - v_\bullet(0)) \cdot (v'_\bullet w'^*(b) - v'_\bullet(0)) \in \mathcal{S}(C).$$

(2) $a \in \mathcal{S}(X)$ or $b \in \mathcal{S}(Y)$.

Remark that (2) always implies (1).

By a straightforward argument, we may assume C, D, D' are transitive, and this condition can be also written as follows.

Proposition 4.2. Let G be an arbitrary finite group, and let $\mathcal{S} = \{I_H\}_{H \leq G}$ be an ideal of T . Then \mathcal{S} is prime if and only if the following condition is satisfied for any $H, H' \leq G$ and any $a \in T(G/H)$, $b \in T(G/H')$.

(*) If $(\text{jnd}_{K^g}^L \circ c_{g,K} \circ \text{res}_K^H(a)) \cdot (\text{jnd}_{K'^{g'}}^L \circ c_{g',K'} \circ \text{res}_{K'}^{H'}(a)) \in I_L$ is satisfied for any $L, K, K' \leq G$ and $g, g' \in G$ satisfying $L \geq K^g, K \leq H, L \geq K'^{g'}, K' \leq H'$, then

$$a \in I_H \quad \text{or} \quad b \in I_{H'}$$

holds.

In our case, this can be reduced to the following condition.

Corollary 4.3. *Let G be a cyclic p -group of p -rank $r \geq 0$. An ideal $\mathcal{I} = [I_0, \dots, I_r]$ of Ω_G is prime if and only if it satisfies the following condition $\mathcal{P}(k, \ell)$ for each $0 \leq \ell \leq k \leq r$.*

$\boxed{\mathcal{P}(k, \ell)}$ For any $a \in R_k$ and $b \in R_\ell$,

$$\begin{aligned} & (\text{jnd}_i^m \circ \text{res}_i^k(a)) \cdot (\text{jnd}_j^m \circ \text{res}_j^\ell(b)) \in I_m \\ & (0 \leq \forall i \leq k, 0 \leq \forall j \leq \ell, m = \max(i, j)) \end{aligned}$$

implies

$$a \in I_k \quad \text{or} \quad b \in I_\ell.$$

Proof. For any $0 \leq k, \ell \leq r$ and any $a \in R_k, b \in R_\ell$, the condition (*) in Proposition 4.2 is equivalent to the following.

(*) If

$$(4.1) \quad (\text{jnd}_i^m \circ c_g \circ \text{res}_i^k(a)) \cdot (\text{jnd}_j^m \circ c_{g'} \circ \text{res}_j^\ell(b)) \in I_m$$

is satisfied for any $g, g' \in G$ and any $m \geq i \leq k, m \geq j \leq \ell$, then

$$a \in I_k \quad \text{or} \quad b \in I_\ell$$

holds.

Remark that we have $c_g = \text{id}, c_{g'} = \text{id}$. Besides, by (iii) of Proposition 3.2, assumption (4.1) is only have to be confirmed for $m = \max(i, j)$. Moreover, by the symmetry in k and ℓ , we may assume $\ell \leq k$. \square

Furthermore, this condition can be checked on each k -th step, as follows.

Proposition 4.4. *Let G be as above. An ideal $\mathcal{I} = [I_0, \dots, I_r]$ of Ω_G is prime if and only if it satisfies the following condition $\mathcal{P}(k)$ for each $0 \leq k \leq r$.*

$\boxed{\mathcal{P}(k)}$ For any $0 \leq i \leq k$ and any $a \in L_k(I_{k-1}), b \in L_i(I_{i-1}) \setminus I_i$,

$$(4.2) \quad a \cdot \text{jnd}_i^k(b) \in I_k \implies a \in I_k$$

holds.

Here, when $k = 0$, we define $L_0(I_{-1})$ to be R_0 . Namely, $\mathcal{P}(0)$ is as follows.

$\boxed{\mathcal{P}(0)}$ For any $a \in R_0$ and $b \in R_0 \setminus I_0$,

$$ab \in I_0 \implies a \in I_0$$

holds. (This is saying $I_0 \subseteq R_0$ is prime in the ordinary ring-theoretic meaning.)

Proof. For each $0 \leq k \leq r$, we define condition $\mathcal{Q}(k)$ as follows.

$\boxed{\mathcal{Q}(k)}$ $\mathcal{P}(k, \ell)$ holds for all $0 \leq \ell \leq k$.

It suffices to show that $\mathcal{Q}(k)$ holds for any $0 \leq k \leq r$ if and only if $\mathcal{P}(k)$ holds for any $0 \leq k \leq r$. This follows from:

Claim 4.5. For any $0 \leq k \leq r$, the following holds.

- (1) $\mathcal{Q}(k)$ implies $\mathcal{P}(k)$.
- (2) If $k \geq 1$, then $\mathcal{Q}(k-1)$ and $\mathcal{P}(k)$ imply $\mathcal{Q}(k)$.
- (3) $\mathcal{Q}(0)$ is equivalent to $\mathcal{P}(0)$.

In fact if this is shown, then by an induction on k , we can easily show that the following are equivalent for each $0 \leq k \leq r$.

- $\mathcal{Q}(k')$ holds for any $0 \leq k' \leq k$.
- $\mathcal{P}(k')$ holds for any $0 \leq k' \leq k$.

This proves Proposition 4.4. Thus it remains to show Claim 4.5.

Proof of Claim 4.5. (3) When $k = 0$, then the condition

$\mathcal{Q}(0) \iff \mathcal{P}(0,0)$ holds. Namely, for any $a, b \in R_0$,

$$ab \in I_0 \implies a \in I_0 \text{ or } b \in I_0$$

holds.

is obviously equivalent to $\mathcal{P}(0)$.

(1) Fix k , suppose we are given $0 \leq \ell \leq k$ and $a \in L_k(I_{k-1}), b \in L_\ell(I_{\ell-1}) \setminus I_\ell$ satisfying

$$(4.3) \quad a \cdot \text{jnd}_\ell^k(b) \in I_k.$$

It suffices to show $a \in I_k$. Since $\mathcal{Q}(k)$ (in particular $\mathcal{P}(k, \ell)$) is assumed, it is enough to confirm that

$$(4.4) \quad (\text{jnd}_i^m \circ \text{res}_i^k(a)) \cdot (\text{jnd}_j^m \circ \text{res}_j^\ell(b)) \in I_m$$

is satisfied for any $0 \leq i \leq k, 0 \leq j \leq \ell$ and $m = \max(i, j)$. However, when $i < k$ or $j < \ell$, (4.4) follows from $\text{res}_i^k(a) \in I_i$ and $\text{res}_j^\ell(b) \in I_j$, since we have $a \in L_k(I_{k-1})$ and $b \in L_\ell(I_{\ell-1})$. In the remaining case of $i = k$ and $j = \ell$, (4.4) is also satisfied by the assumption (4.3).

(2) Fix $k \geq 1$, and assume $\mathcal{Q}(k-1)$. Under this assumption, we show $\mathcal{P}(k)$ implies $\mathcal{P}(k, \ell)$ for any $0 \leq \ell \leq k$. By an induction on ℓ , this is reduced to the following.

Claim 4.6. For any $1 \leq k \leq r$ and $0 \leq \ell \leq k$, we have

$$\mathcal{Q}(k-1), \mathcal{P}(k), \mathcal{P}(k, \ell-1) \implies \mathcal{P}(k, \ell).$$

(Here, for $\ell = 0$, $\mathcal{P}(k, -1)$ is regarded as an empty condition.)

We only have to show this claim in the rest. Suppose $a \in R_k$ and $b \in R_\ell$ satisfy

$$(4.5) \quad (\text{jnd}_i^m \circ \text{res}_i^k(a)) \cdot (\text{jnd}_j^m \circ \text{res}_j^\ell(b)) \in I_m$$

for any $0 \leq i \leq k, 0 \leq j \leq \ell, m = \max(i, j)$. Claim 4.6 will follow immediately, if the following are shown.

- (A) If $a \in R_k \setminus I_k$, then $b \in L_\ell(I_{\ell-1})$. (This is trivial when $\ell = 0$, since we have defined as $L_0(I_{-1}) = R_0$.)
- (B) If $b \in R_\ell \setminus I_{\ell-1}$, then $a \in L_k(I_{k-1})$.

In fact, if (A) and (B) are shown, then the above a and b will satisfy

- (i) $a \in I_k$ or

- (ii) $b \in I_\ell$ or
- (iii) $a \in L_k(I_{k-1})$ and $b \in L_\ell(I_{\ell-1}) \setminus I_\ell$.

In the third case, since $a \cdot \text{jnd}_\ell^k(b) \in I_k$ is satisfied by (4.5) for $i = k$ and $j = \ell$, it follows $a \in I_k$ by $\mathcal{P}(k)$.

Thus it remains to show (A) and (B).

(A) By applying $\mathcal{P}(k, \ell - 1)$ to

$$a \in R_k \quad \text{and} \quad \text{res}_{\ell-1}^\ell(b) \in R_{\ell-1},$$

we obtain $\text{res}_{\ell-1}^\ell(b) \in I_{\ell-1}$, namely $b \in L_\ell(I_{\ell-1})$.

(B) When $\ell < k$, by applying $\mathcal{Q}(k - 1)$, in particular $\mathcal{P}(k - 1, \ell)$ to

$$\text{res}_{k-1}^k(a) \in R_{k-1} \quad \text{and} \quad b \in R_\ell,$$

we obtain $\text{res}_{k-1}^k(a) \in I_{k-1}$, namely $a \in L_k(I_{k-1})$.

When $\ell = k$, by applying $\mathcal{P}(k, k - 1)$ to

$$b \in R_k \quad \text{and} \quad \text{res}_{k-1}^k(a) \in R_{k-1},$$

we obtain $\text{res}_{k-1}^k(a) \in I_{k-1}$, namely $a \in L_k(I_{k-1})$. □

□

By Proposition 4.4, whether an ideal is prime or not can be checked inductively on k using $\mathcal{P}(k)$. This is applied to restrictions and extensions of prime ideals as follows.

Corollary 4.7. *For any $0 \leq i \leq k \leq r$, if an ideal $\mathcal{S} = [I_0, \dots, I_k] \subseteq \Omega_{H_k}$ is prime, then its restriction $\mathcal{S}|_{H_i} = [I_0, \dots, I_i]$ onto H_i is also prime.*

Proof. This immediately follows from Proposition 4.4. □

Corollary 4.8. *For $k \geq 1$, let $\mathcal{S} = [I_0, \dots, I_{k-1}]$ be an ideal of Ω_{H_k} . If \mathcal{S} is prime, then $\mathcal{L}\mathcal{S} \subseteq \Omega_{H_k}$ is also prime.*

Proof. It suffices to show $\mathcal{P}(k)$ is satisfied. However for $\mathcal{L}\mathcal{S} = [I_0, \dots, I_{k-1}, I_k = L(I_{k-1})]$, this condition becomes trivial as follows.

- For any $0 \leq i \leq k$, $a \in L_k(I_{k-1})$ and $b \in L_i(I_{i-1}) \setminus I_i$,

$$(4.6) \quad a \cdot \text{jnd}_i^k(b) \in I_k \implies a \in I_k$$

holds.

Of course this is satisfied, since a belongs to $I_k = L_k(I_{k-1})$ from the first. □

5. IDEALS $J_{\ell,k}(x) \subseteq R_\ell$

In this section, we introduce ideals $J_{\ell,k}(x)$ of R_ℓ , which will perform an essential role in determining the prime ideals of Ω .

First we prepare another \mathbb{Z} -basis for R_ℓ , which is more suitable for calculation.

Definition 5.1. For any $0 \leq i \leq \ell \leq r$, define $F_{\ell,i} \in R_\ell$ by

$$F_{\ell,i} = X_{\ell,i} - p^{\ell-i}.$$

Obviously, each R_ℓ admits a \mathbb{Z} -basis

$$\{1, F_{\ell,0}, F_{\ell,1}, \dots, F_{\ell,\ell-1}\}$$

for $0 \leq \ell \leq r$, and thus any element $\alpha \in R_\ell$ can be written as

$$\alpha = m_\ell + \sum_{i=0}^{\ell-1} m_i F_{\ell,i}$$

for some uniquely determined $m_0, \dots, m_\ell \in \mathbb{Z}$.

This basis behaves well with the multiplication and the structure morphisms as follows.

Proposition 5.2. *The following holds.*

(1) *For any $0 \leq i, j \leq \ell$, we have*

$$X_{\ell,j} \cdot F_{\ell,i} = \begin{cases} p^{\ell-j} F_{\ell,i} - p^{\ell-i} F_{\ell,j} & (i \leq j), \\ 0 & (j \leq i). \end{cases}$$

(2) *For any $0 \leq i, k \leq \ell$, we have*

$$\text{res}_k^\ell(F_{\ell,i}) = \begin{cases} p^{\ell-k} F_{k,i} & (i \leq k), \\ 0 & (k \leq i). \end{cases}$$

In particular, we have

$$\text{res}_{\ell-1}^\ell(\alpha) = m_\ell + \sum_{i=0}^{\ell-2} m_i p F_{\ell-1,i}$$

for any $\alpha = m_\ell + \sum_{i=0}^{\ell-1} m_i F_{\ell,i} \in R_\ell$.

(3) *When $\ell \geq 1$, for any*

$$\alpha = n_{\ell-1} + \sum_{i=0}^{\ell-2} m_i F_{\ell-1,i} \in R_{\ell-1},$$

we have

$$\text{jnd}_{\ell-1}^\ell(\alpha) = (n_{\ell-1})^p + \sum_{i=0}^{\ell-1} u_i F_{\ell,i}$$

for some $u_0, \dots, u_{\ell-1} \in \mathbb{Z}$.

(4) *When $\ell \geq 2$, for any $0 \leq i \leq \ell - 1$, we have*

$$\text{ind}_{\ell-1}^\ell(F_{\ell-1,i}) = F_{\ell,i} - p^{\ell-i-1} F_{\ell,\ell-1}.$$

Moreover, $F_{\ell,\ell-1}$ is calculated as

$$F_{\ell,\ell-1} = \text{jnd}_{\ell-1}^\ell(F_{\ell-1,\ell-2}) + \frac{(-p)^p}{p^2} \text{ind}_{\ell-1}^\ell(F_{\ell-1,\ell-2}).$$

Proof. (1) and (2) follows immediately from Corollary 2.3.

(3) For any $\alpha = n_{\ell-1} + \sum_{i=0}^{\ell-2} m_i F_{\ell-1,i} \in R_{\ell-1}$, if we put $m_{\ell-1} = n_{\ell-1} - \sum_{i=0}^{\ell-2} m_i p^{\ell-i-1}$,

then we have

$$\alpha = m_{\ell-1} + \sum_{i=0}^{\ell-2} m_i X_{\ell-1,i}$$

and thus by Corollary 2.4, we obtain

$$\begin{aligned}
\text{jnd}_{\ell-1}^{\ell}(\alpha) &= m_{\ell-1} + \frac{(m_{\ell-1})^p - m_{\ell-1}}{p} X_{\ell, \ell-1} \\
&+ \sum_{i=0}^{\ell-2} \frac{(\sum_{s=i}^{\ell-1} m_s p^{\ell-1-s})^p - (\sum_{s=i+1}^{\ell-1} m_s p^{\ell-1-s})^p}{p^i} X_{\ell, i} \\
&= m_{\ell-1} + \{(m_{\ell-1})^p - m_{\ell-1}\} + \sum_{i=0}^{\ell-2} \left\{ (\sum_{s=i}^{\ell-1} m_s p^{\ell-1-s})^p - (\sum_{s=i+1}^{\ell-1} m_s p^{\ell-1-s})^p \right\} \\
&+ \frac{(m_{\ell-1})^p - m_{\ell-1}}{p} F_{\ell, \ell-1} + \sum_{i=0}^{\ell-2} \frac{(\sum_{s=i}^{\ell-1} m_s p^{\ell-1-s})^p - (\sum_{s=i+1}^{\ell-1} m_s p^{\ell-1-s})^p}{p^i} F_{\ell, i} \\
&= (n_{\ell-1})^p + \frac{(m_{\ell-1})^p - m_{\ell-1}}{p} F_{\ell, \ell-1} \\
&+ \sum_{i=0}^{\ell-2} \frac{(n_{\ell-1} - \sum_{s=0}^{i-1} m_s p^{\ell-1-s})^p - (n_{\ell-1} - \sum_{s=0}^i m_s p^{\ell-1-s})^p}{p^i} F_{\ell, i}.
\end{aligned}$$

(4) By Remark 2.2, we have

$$\begin{aligned}
&\text{ind}_{\ell-1}^{\ell}(F_{\ell-1, i}) + p^{\ell-i-1} F_{\ell, \ell-1} \\
&= (X_{\ell, i} - p^{\ell-i-1} X_{\ell, \ell-1}) + (p^{\ell-i-1} X_{\ell, \ell-1} - p^{\ell-i}) \\
&= F_{\ell, i}.
\end{aligned}$$

Moreover, by Remark 2.2 and Corollary 2.4, we have

$$\begin{aligned}
&\text{jnd}_{\ell-1}^{\ell}(F_{\ell-1, \ell-2}) + \frac{(-p)^p}{p^2} \text{ind}_{\ell-1}^{\ell}(F_{\ell-1, \ell-2}) \\
&= (-p + \frac{(-p)^p + p}{p} X_{\ell, \ell-1} - \frac{(-p)^p}{p^2} X_{\ell, \ell-2}) + \frac{(-p)^p}{p^2} (X_{\ell, \ell-2} - p X_{\ell, \ell-1}) \\
&= X_{\ell, \ell-1} - p = F_{\ell, \ell-1}.
\end{aligned}$$

□

Lemma 5.3. For $1 \leq \ell \leq r$ and $n \in \mathbb{Z}$, we have the following.

(1) If n is not divisible by p , then we have

$$n = \text{jnd}_{\ell-1}^{\ell}(n) - \frac{n^{p-1} - 1}{p} \text{ind}_{\ell-1}^{\ell}(n).$$

Remark that $\frac{n^{p-1}-1}{p}$ is an integer, by assumption.

(2) If $n = pu$ for some $u \in \mathbb{Z}$, then we have

$$n = \text{ind}_{\ell-1}^{\ell}(u) - u F_{\ell, \ell-1}.$$

Proof. (1) By Remark 2.2 and Corollary 2.4, we have

$$\text{jnd}_{k-1}^k(n) - \frac{n^{p-1} - 1}{p} \text{ind}_{k-1}^k(n) = n + \frac{n^p - n}{p} X_{k, k-1} - \frac{n^{p-1} - 1}{p} n X_{k, k-1} = n.$$

(2) This follows from

$$\begin{aligned}
\text{ind}_{\ell-1}^{\ell}(u) - u F_{\ell, \ell-1} &= u X_{\ell, \ell-1} - u (X_{\ell, \ell-1} - p) \\
&= pu = n.
\end{aligned}$$

□

Definition 5.4. For any $0 \leq \ell \leq r$, $0 \leq k \leq \ell$ and $x \in \mathbb{Z}$, we define an ideal $J_{\ell,k}(x) \subseteq R_\ell$ by

$$J_{\ell,k}(x) = \begin{cases} (x, F_{\ell,k}, F_{\ell,k+1}, \dots, F_{\ell,\ell-1}) & (k \leq \ell - 1), \\ (x) & (k = \ell). \end{cases}$$

$J_{\ell,k}(x)$ can be calculated as follows.

Proposition 5.5. For any $1 \leq \ell \leq r$, $0 \leq k \leq \ell - 1$ and $x \in \mathbb{Z}$, we have

$$\begin{aligned} J_{\ell,k}(x) &= \left\{ x \cdot \left(n_\ell + \sum_{i=0}^{k-1} n_i F_{\ell,i} \right) + \sum_{i=k}^{\ell-1} n_i F_{\ell,i} \mid n_0, \dots, n_\ell \in \mathbb{Z} \right\} \\ &= \left\{ m_\ell + \sum_{i=0}^{\ell-1} m_i F_{\ell,i} \mid \begin{array}{l} m_0, \dots, m_\ell \in \mathbb{Z}, \\ m_\ell \in \mathbb{Z}x, \\ m_i \in \mathbb{Z}x \quad (0 \leq i \leq k-1) \end{array} \right\}. \end{aligned}$$

Proof. Obviously, $J_{\ell,k}(x)$ contains any element of the form

$$(5.1) \quad x \cdot \left(n_\ell + \sum_{i=0}^{k-1} n_i F_{\ell,i} \right) + \sum_{i=k}^{\ell-1} n_i F_{\ell,i}.$$

To show the converse, since any element in $J_{\ell,k}(x)$ can be written as an R_ℓ -coefficient sum of

$$x, F_{\ell,k}, \dots, F_{\ell,\ell-1},$$

it suffices to show that any element

- (i) αx
- (ii) $\alpha F_{\ell,i}$ ($k \leq i \leq \ell - 1$)

can be written in the form of (5.1) for any $\alpha \in R_\ell$.

- (i) For any $\alpha = m_\ell + \sum_{i=0}^{\ell-1} m_i F_{\ell,i}$ ($m_i \in \mathbb{Z}$), we have

$$\alpha x = x \cdot \left(m_\ell + \sum_{i=0}^{k-1} m_i F_{\ell,i} \right) + \sum_{i=k}^{\ell-1} x m_i F_{\ell,i}.$$

- (ii) For any $\alpha = \sum_{j=0}^{\ell} m_j X_{\ell,j}$ ($m_j \in \mathbb{Z}$), we have

$$\begin{aligned} \alpha F_{\ell,i} &= \sum_{j=0}^i m_j X_{\ell,j} F_{\ell,i} + \sum_{j=i+1}^{\ell} m_j X_{\ell,j} F_{\ell,i} \\ &= \sum_{j=i+1}^{\ell} m_j (p^{\ell-j} F_{\ell,i} - p^{\ell-i} F_{\ell,j}) \\ &= \left(\sum_{j=i+1}^{\ell} m_j p^{\ell-j} \right) F_{\ell,i} - \sum_{j=i+1}^{\ell} m_j p^{\ell-i} F_{\ell,j} \end{aligned}$$

for any $k \leq i \leq \ell - 1$ by Proposition 5.2. □

Proposition 5.6. *For any $1 \leq \ell \leq r$, $0 \leq k \leq \ell - 1$ and any $n \in \mathbb{Z}$, we have the following.*

- (1) *If $n = 0$ or $n = q$ for some prime integer q different from p , then we have*
- (i) $L(J_{\ell-1,k}(n)) = J_{\ell,k}(n)$,
 - (ii) $S(J_{\ell-1,k}(n)) = \begin{cases} J_{\ell,k}(n) & (k \leq \ell - 2), \\ J_{\ell,\ell}(n) = (n) & (k = \ell - 1). \end{cases}$

Moreover, when $n = q$ is a prime different from p , then there is no ideal between $J_{\ell,k+1}(q) \subsetneq J_{\ell,k}(q)$.

- (2) *For $k = 0$ and $n = p^{e+1}$ for some $e \in \mathbb{N}_{\geq 0}$, we have*

- (i) $L(J_{\ell-1,0}(p^{e+1})) = J_{\ell,0}(p^{e+1})$,
- (ii) $S(J_{\ell-1,0}(p^{e+1})) = J_{\ell,0}(p^{e+2})$.

Moreover, there is no ideal between $J_{\ell,0}(p^{e+2}) \subsetneq J_{\ell,0}(p^{e+1})$.

Proof. (1) (i) By definition,

$$L(J_{\ell-1,k}(n)) = \{\alpha \in R_\ell \mid \text{res}_{\ell-1}^\ell(\alpha) \in J_{\ell-1,k}(n)\}.$$

For any $\alpha = m_\ell + \sum_{i=0}^{\ell-1} m_i F_{\ell,i}$ ($m_i \in \mathbb{Z}$), since we have

$$\begin{aligned} \text{res}_{\ell-1}^\ell(\alpha) &= m_\ell + \sum_{i=0}^{\ell-1} m_i p F_{\ell-1,i} \\ &= m_\ell + \sum_{i=0}^{k-1} m_i p F_{\ell-1,i} + \sum_{i=k}^{\ell-1} m_i p F_{\ell-1,i}, \end{aligned}$$

this satisfies $\text{res}_{\ell-1}^\ell(\alpha) \in J_{\ell-1,k}(n)$ if and only if

$$\begin{aligned} m_\ell &\in n\mathbb{Z}, \\ m_i p &\in n\mathbb{Z} \quad (0 \leq i \leq k-1), \end{aligned}$$

by Proposition 5.5. Since n is 0 or a prime different from p , this is equivalent to

$$m_\ell \in n\mathbb{Z} \quad \text{and} \quad m_i \in n\mathbb{Z} \quad (0 \leq i \leq k-1),$$

namely, to $\alpha \in J_{\ell,k}(n)$.

(ii) When $k \leq \ell - 2$, it suffices to show

$$J_{\ell,k}(n) \subseteq S(J_{\ell-1,k}(n)).$$

In fact, this implies

$$J_{\ell,k}(n) \subseteq S(J_{\ell-1,k}(n)) \subseteq L(J_{\ell-1,k}(n)) = J_{\ell,k}(n),$$

and thus $S(J_{\ell-1,k}(n)) = L(J_{\ell-1,k}(n)) = J_{\ell,k}(n)$ follows.

Thus it remains to show $J_{\ell,k}(n) = (n, F_{\ell,k}, \dots, F_{\ell,\ell-1}) \subseteq S(J_{\ell-1,k}(n))$. However, this immediately follows from Proposition 5.2 and Lemma 5.3, since we have

$$\begin{aligned} n &= \text{jnd}_{\ell-1}^\ell(n) - \frac{n^{p-1} - 1}{p} \quad (\text{for } n \neq 0), \\ F_{\ell,\ell-1} &= \text{jnd}_{\ell-1}^\ell(F_{\ell-1,\ell-2}) + \frac{(-p)^p}{p^2} \text{ind}_{\ell-1}^\ell(F_{\ell-1,\ell-2}), \\ F_{\ell,i} &= \text{ind}_{\ell-1}^\ell(F_{\ell-1,i}) + p^{\ell-i-1} F_{\ell,\ell-1} \quad (k \leq i \leq \ell-1). \end{aligned}$$

When $k = \ell - 1$, remark that we have $J_{\ell-1, \ell-1}(q) = (q) \subseteq R_{\ell-1}$. If $n = 0$, $(0) \subseteq S(J_{\ell-1, \ell-1}(0))$ is trivial. If $n = q$ is a prime different from p , then by Lemma 5.3 we have

$$q = \text{jnd}_{\ell-1}^{\ell}(q) - \frac{q^{p-1} - 1}{p} \text{ind}_{\ell-1}^{\ell}(q) \in S((q)),$$

which means $(q) \subseteq S(J_{\ell-1, \ell-1}(q))$.

Conversely, if $n = q$ is a prime different from p , then for any $\alpha \in R_{\ell-1}$, we have

$$\begin{aligned} \text{jnd}_{\ell-1}^{\ell}(\alpha q) &= \text{jnd}_{\ell-1}^{\ell}(\alpha) \cdot \text{jnd}_{\ell-1}^{\ell}(q), \\ &= \text{jnd}_{\ell-1}^{\ell}(\alpha) \cdot \left(q + \frac{q^p - q}{p} X_{\ell, \ell-1} \right), \\ &= \text{jnd}_{\ell-1}^{\ell}(\alpha) \cdot \left(1 + \frac{q^{p-1} - 1}{p} X_{\ell, \ell-1} \right) \cdot q \in (q), \\ \text{ind}_{\ell-1}^{\ell}(\alpha q) &= \text{ind}_{\ell-1}^{\ell}(\alpha) \cdot q \in (q), \end{aligned}$$

which imply $S(J_{\ell-1, \ell-1}(q)) \subseteq (q)$. Similarly, $S(J_{\ell-1, \ell-1}(0)) \subseteq (0)$ follows from $\text{ind}_{\ell-1}^{\ell}(0) = \text{jnd}_{\ell-1}^{\ell}(0) = 0$.

Thus it remains to show there is no ideal between $J_{\ell, k+1}(q) \subsetneq J_{\ell, k}(q)$ for a prime $q \neq p$. Suppose there is an ideal

$$J_{\ell, k+1}(q) \subsetneq I \subseteq J_{\ell, k}(q).$$

By Proposition 5.5, I should contain an element

$$\alpha = q\beta + \sum_{i=k}^{\ell-1} n_i F_{\ell, i}$$

in $J_{\ell, k}(q)$, for some $\beta \in R_{\ell}$ and $n_i \in \mathbb{Z}$ ($k \leq i \leq \ell - 1$), which does not belong to $J_{\ell, k+1}(q)$. Then I should contain

$$n_k F_{\ell, k} = \alpha - \left\{ q\beta + \sum_{i=k+1}^{\ell-1} n_i F_{\ell, i} \right\}.$$

Since α does not belong to $J_{\ell, k+1}(q)$, it follows that q does not divide n_k , and thus I contains an element of the form

$$n_k F_{\ell, k} \quad (n_k \in \mathbb{Z}, \text{ not divisible by } q).$$

On the other hand, $q \in J_{\ell, k+1}(q) \subseteq I$ implies

$$q F_{\ell, k} \in I.$$

Since q and n_k are coprime, it follows $F_{\ell, k} \in I$, which means $I = J_{\ell, k}(q)$.

(2) (i) For an element $\alpha = m_{\ell} + \sum_{i=0}^{\ell-1} m_i F_{\ell, i}$, since we have

$$\text{res}_{\ell-1}^{\ell}(\alpha) = m_{\ell} + \sum_{i=0}^{\ell-2} m_i p F_{\ell-1, i}$$

by Proposition 5.2, it satisfies $\alpha \in L(J_{\ell-1, 0}(p^{e+1}))$ if and only if $m_{\ell} \in p^{e+1}\mathbb{Z}$, namely $\alpha \in J_{\ell, 0}(p^{e+1})$.

(ii) By Proposition 5.2 and Lemma 5.3, we have

$$\begin{aligned} F_{\ell, \ell-1} &= \text{jnd}_{\ell-1}^{\ell}(F_{\ell-1, \ell-2}) + \frac{(-p)^p}{p^2} \text{ind}_{\ell-1}^{\ell}(F_{\ell-1, \ell-2}), \\ F_{\ell, i} &= \text{ind}_{\ell-1}^{\ell}(F_{\ell-1, i}) + p^{\ell-i-1}(F_{\ell, \ell-1}) \quad (0 \leq i \leq \ell-1), \\ p^{e+2} &= \text{ind}_{\ell-1}^{\ell}(p^{e+1}) - p^{e+1}F_{\ell, \ell-1}, \end{aligned}$$

which imply $J_{\ell, 0}(p^{e+2}) \subseteq S(L_{\ell-1, 0}(p^{e+1}))$.

To show the converse, by Proposition 5.5, it suffices to show any element

$$\alpha = p^{e+1}n_{\ell-1} + \sum_{i=0}^{\ell-2} m_i F_{\ell-1, i} \quad (n_{\ell-1}, m_i \in \mathbb{Z} \quad (0 \leq i \leq \ell-2))$$

in $J_{\ell-1, 0}(p^{e+1})$ satisfies $\text{ind}_{\ell-1}^{\ell}(\alpha) \in J_{\ell, 0}(p^{e+2})$ and $\text{jnd}_{\ell-1}^{\ell}(\alpha) \in J_{\ell, 0}(p^{e+2})$. However, these follow from

$$\begin{aligned} \text{ind}_{\ell-1}^{\ell}(\alpha) &= p^{e+1}n_{\ell-1}X_{\ell, \ell-1} + \sum_{i=0}^{\ell-2} m_i (F_{\ell, i} - p^{\ell-i-1}F_{\ell, \ell-1}) \\ &= p^{e+2}n_{\ell-1} + \sum_{i=0}^{\ell-2} m_i F_i + (p^{e+1}n_{\ell-1} - \sum_{i=0}^{\ell-2} p^{\ell-i-1}m_i)F_{\ell, \ell-1} \end{aligned}$$

and

$$\text{jnd}_{\ell-1}^{\ell}(\alpha) = (p^{e+1}n_{\ell-1})^p + \sum_{i=0}^{\ell-1} u_i F_{\ell, i}$$

for some $u_0, \dots, u_{\ell-1} \in \mathbb{Z}$, by Proposition 5.2.

It remains to show there is no ideal between $J_{\ell, 0}(p^{e+2}) \subsetneq J_{\ell, 0}(p^{e+1})$. Suppose there is an ideal

$$J_{\ell, 0}(p^{e+2}) \subsetneq I \subseteq J_{\ell, 0}(p^{e+1}).$$

Then by Proposition 5.5, I should contain an element

$$\alpha = p^{e+1}n_{\ell} + \sum_{i=0}^{\ell-1} m_i F_{\ell, i}$$

in $J_{\ell, 0}(p^{e+1})$, for some $n_{\ell}, m_i \in \mathbb{Z}$ ($0 \leq i \leq \ell-1$), which does not belong to $J_{\ell, 0}(p^{e+2})$. Then I should contain

$$p^{e+1}n_{\ell} = \alpha - \sum_{i=0}^{\ell-1} m_i F_{\ell, i}.$$

Since α is not in $J_{\ell, 0}(p^{e+2})$, it follows that p does not divide n_{ℓ} . On the other hand, we have

$$p^{e+2} \in J_{\ell, 0}(p^{e+2}) \subseteq I.$$

These imply $p^{e+1} \in I$, which means $I = J_{\ell, 0}(p^{e+1})$. □

6. STRUCTURE OF $\text{Spec } \Omega$

Let $0 \leq \ell \leq r$ be any integer. For any ideal $\mathcal{I} = [I_0, \dots, I_\ell] \subseteq \Omega_{H_\ell}$, define $F(\mathcal{I})$ to be the ideal $F(\mathcal{I}) = I_0$ of R_0 . Since I_0 becomes prime if \mathcal{I} is prime, this gives a map

$$F: \text{Spec } \Omega_G \rightarrow \text{Spec } R_0.$$

To determine $\text{Spec } \Omega$, it suffices to investigate each fiber of F over the points of $\text{Spec } R_0$. Since $\text{Spec } R_0 = \text{Spec } \mathbb{Z} = \{0\} \cup \{(q) \mid q \text{ is a prime integer}\}$, in the rest we will study the following three cases.

- (i) Prime ideals over (q) , where q is a prime integer different from p .
- (ii) Prime ideals over (0) .
- (iii) Prime ideals over (p) .

6.1. Prime ideals over $q \neq p$. Let q be any prime integer different from p .

Proposition 6.1. *We have the following.*

- (1) For any $k \geq 0$, we have

$$S^k(q) = (q) \subseteq R_k,$$

and thus $S^k(q) = [(q), (q), \dots, (q)]$.

- (2) For any $1 \leq \ell \leq r$ and $0 \leq k \leq \ell$, we have

$$L^{\ell-k} S^k(q) = J_{\ell,k}(q) \subseteq R_\ell.$$

- (3) For any $1 \leq \ell \leq r$ and $0 \leq k \leq \ell - 2$, we have

$$SL^{\ell-k-1} S^k(q) = L^{\ell-k} S^k(q).$$

Proof. This follows from Proposition 5.6, by an induction on k and ℓ . □

Proposition 6.2. *For any $0 \leq k \leq r$,*

$$S^k(q) = [(q), \dots, (q)] \subseteq \Omega_{H_k}$$

is a prime ideal.

Proof. We show this by an induction on k . If $k = 0$, then $(q) \subseteq \Omega_e \cong \mathbb{Z}$ is indeed a prime ideal.

When $k \geq 1$, suppose $S^{k-1}(q) \subseteq \Omega_{H_{k-1}}$ is prime. It suffices to show $S^k(q) = [(q), \dots, (q)]$ satisfies $\mathcal{P}(k)$. Namely, we only have to show the following.

Claim 6.3. For any $0 \leq i \leq k$, any $a \in L_k(q)$ and any $b \in L_i(q) \setminus (q)$,

$$a \cdot \text{jnd}_i^k(b) \in (q) \implies a \in (q)$$

holds.

Proof. Since $L_k(q) = J_{k,k-1}(q)$, a can be written as

$$a = q\beta + mF_{k,k-1} \quad (\beta \in R_k, m \in \mathbb{Z})$$

by Proposition 5.5.

Similarly by Proposition 5.5, $b \in L_i(q) = J_{i,i-1}(q)$ can be written in the form

$$\begin{aligned} b &= q(n_i + \sum_{j=0}^{i-2} n_j F_{i,j}) + n_{i-1} F_{i,i-1} \\ &= q(n_i - \sum_{j=0}^{i-2} n_j p^{i-j}) - pn_{i-1} + \sum_{j=0}^{i-2} n_j q X_{i,j} + n_{i-1} X_{i,i-1} \end{aligned}$$

for some $n_j \in \mathbb{Z}$ ($0 \leq j \leq i$). If we put $n'_i = n_i - \sum_{j=0}^{i-2} n_j p^{i-j}$, then we have

$$\text{jnd}_i^k(b) = (qn'_i - pn_{i-1}) + \sum_{t=0}^{k-1} u_t X_{k,t}$$

for some $u_t \in \mathbb{Z}$ ($0 \leq t < k$), by Corollary 2.4.

Now we have

$$a \cdot \text{jnd}_i^k(b) = q(\beta \cdot \text{jnd}_i^k(b)) + (qn'_i - pn_{i-1})mF_{k,k-1}.$$

This satisfies $a \cdot \text{jnd}_i^k(b) \in (q)$ if and only if

$$pn_{i-1}mF_{k,k-1} \in (q),$$

which is equivalent to $pn_{i-1}m \in q\mathbb{Z}$ by Proposition 5.2. Since b is not in (q) , n_{i-1} is not divisible by q . Thus it follows $m \in q\mathbb{Z}$, which means $a \in (q)$. \square

\square

Proposition 6.4. *In Ω_{H_ℓ} , there are exactly $(\ell + 1)$ ideals over $(q) \subseteq R_0$*

$$(6.1) \quad \mathcal{S}^\ell(q) \subsetneq \mathcal{L}\mathcal{S}^{\ell-1}(q) \subsetneq \cdots \subsetneq \mathcal{L}^{\ell-k}\mathcal{S}^k(q) \subsetneq \cdots \subsetneq \mathcal{L}^\ell(q),$$

all of which are prime.

Proof. By Proposition 6.2 and Corollary 4.8, these are prime.

We show the proposition by an induction on ℓ . Suppose we have done for $\ell - 1$, and take any ideal $\mathcal{I} = [I_0, \dots, I_\ell] \subseteq \Omega_{H_\ell}$ over $(q) \subseteq R_0$. By the assumption of the induction, there exists some $0 \leq k \leq \ell - 1$ satisfying

$$\mathcal{I}|_{H_{\ell-1}} = \mathcal{L}^{\ell-k-1}\mathcal{S}^k(q).$$

If $k \leq \ell - 2$, then I_ℓ should satisfy

$$I_\ell = SL^{\ell-k-1}\mathcal{S}^k(q) = L^{\ell-k}\mathcal{S}^k(q) = J_{\ell,k}(q)$$

by Proposition 6.1, and thus $\mathcal{I} = \mathcal{L}^{\ell-k}\mathcal{S}^k(q)$.

If $k = \ell - 1$, then I_ℓ should satisfy

$$\mathcal{S}^\ell(q) \subseteq I_\ell \subseteq LS^{\ell-1}(q).$$

Since there is no ideal between $\mathcal{S}^\ell(q) = (q)$ and $LS^{\ell-1}(q) = J_{\ell,\ell-1}(q)$ by Proposition 5.6, we have

$$I_\ell = \mathcal{S}^\ell(q) \quad \text{or} \quad LS^{\ell-1}(q),$$

namely

$$\mathcal{I} = \mathcal{S}^\ell(q) \quad \text{or} \quad \mathcal{L}\mathcal{S}^{\ell-1}(q).$$

\square

6.2. Prime ideals over p .

Proposition 6.5. *Consider an ideal in R_ℓ , obtained from $(p) \subseteq R_0$ by an iterated application of L and S*

$$S^{a_1} L^{b_1} S^{a_2} L^{b_2} \dots S^{a_s} L^{b_s} (p) \subseteq R_\ell$$

for $a_i, b_i \in \mathbb{N}_{\geq 0}$ satisfying $\sum_{i=1}^s a_i + \sum_{i=1}^s b_i = \ell$. Then this depends only on $k = \sum_{i=1}^s a_i$ ($0 \leq k \leq \ell$), and is equal to $J_{\ell,0}(p^{k+1})$. Namely we have

$$\begin{aligned} S^{a_1} L^{b_1} S^{a_2} L^{b_2} \dots S^{a_s} L^{b_s} (p) &= S^k L^{\ell-k} (p) \\ &= L^{\ell-k} S^k (p) = J_{\ell,0}(p^{k+1}). \end{aligned}$$

Proof. By an induction on ℓ , this immediately follows from Proposition 5.6. \square

Corollary 6.6. *For any $1 \leq \ell \leq r$, any ideal $\mathcal{I} = [I_0 = (p), I_1, \dots, I_\ell]$ of Ω_{H_ℓ} over (p) satisfies*

- (1) $I_i = S(I_{i-1})$ or $I_i = L(I_{i-1})$
- (2) $I_i = J_{i,0}(p^{k+1})$ for some $0 \leq k \leq i$

for each $1 \leq i \leq \ell$.

Proof. We proceed by an induction on ℓ . Suppose we have shown for $\ell - 1$. Then, any ideal $\mathcal{I} = [(p), I_1, \dots, I_\ell]$ should satisfy

$$I_{\ell-1} = J_{\ell-1,0}(p^{k+1})$$

for some $0 \leq k \leq \ell - 1$. Then by Proposition 5.6, I_ℓ should lie between $S(I_{\ell-1}) = J_{\ell,0}(p^{k+2})$ and $L(I_{\ell-1}) = J_{\ell,0}(p^{k+1})$, while there is no ideal between them. Thus I_ℓ should agree with $S(I_{\ell-1}) = J_{\ell,0}(p^{k+2})$ or $L(I_{\ell-1}) = J_{\ell,0}(p^{k+1})$. \square

Proposition 6.7. *Among all the ideals of Ω_{H_ℓ} over (p) determined in Corollary 6.6,*

$$\mathcal{L}^\ell(p) = [(p), J_{1,0}(p), \dots, J_{\ell,0}(p)]$$

is the only one prime ideal.

Proof. $\mathcal{L}^\ell(p)$ is prime by Corollary 4.8. Remark that any other ideal $\mathcal{I} = [(p), I_1, \dots, I_\ell]$ should satisfy $I_k = S(I_{k-1})$ for some $1 \leq k \leq \ell$. If we take the smallest such k , then it satisfies

$$I_{k-1} = J_{k-1,0}(p) \quad \text{and} \quad I_k = J_{k,0}(p^2).$$

Then $\mathcal{P}(k)$ fails for \mathcal{I} . In fact, for

$$a = b = p \in L(I_{k-1}) = J_{k,0}(p),$$

we have $ab = p^2 \in J_{k,0}(p^2)$, while neither a nor b belong to I_k . \square

6.3. Prime ideals over 0.

Proposition 6.8. *We have the following.*

(1) *For any $k \geq 0$, we have*

$$S^k(0) = (0).$$

(2) *For any $1 \leq \ell \leq r$ and $0 \leq k \leq \ell$, we have*

$$L^{\ell-k}S^k(0) = J_{\ell,k}(0) \subseteq R_\ell.$$

(3) *For any $1 \leq \ell \leq r$ and $0 \leq k \leq \ell - 2$, we have*

$$SL^{\ell-k-1}S^k(0) = L^{\ell-k}S^k(0).$$

Proof. Similarly to Proposition 6.1, this follows from Proposition 5.6 by an induction. \square

Proposition 6.9. *For any $0 \leq k \leq r$,*

$$S^k(0) = [0, \dots, 0] = 0 \subseteq \Omega_{H_k}$$

is prime.

Proof. This is shown in a similar way as in Proposition 6.2. More generally, for any finite group G , it was shown that the zero ideal $0 \subseteq \Omega_G$ is prime (Theorem 4.40 in [5]). \square

Lemma 6.10. *For any $1 \leq \ell \leq r$ and $0 \leq k \leq \ell - 1$, any ideal I satisfying*

$$J_{\ell,k+1}(0) \subseteq I \subseteq J_{\ell,k}(0)$$

is of the form

$$\begin{aligned} I &= J_{\ell,k+1}(0) + (nF_{\ell,k}) \\ &= (nF_{\ell,k}, F_{\ell,k+1}, F_{\ell,k+2}, \dots, F_{\ell,\ell-1}) \end{aligned}$$

for some $n \in \mathbb{Z}$.

Proof. By Proposition 5.5, we have

$$\begin{aligned} J_{\ell,k}(0) &= \left\{ \sum_{i=k}^{\ell-1} n_i F_{\ell,i} \mid n_i \in \mathbb{Z} \ (k \leq i \leq \ell-1) \right\}, \\ J_{\ell,k+1}(0) &= \left\{ \sum_{i=k+1}^{\ell-1} n_i F_{\ell,i} \mid n_i \in \mathbb{Z} \ (k+1 \leq i \leq \ell-1) \right\}. \end{aligned}$$

When $I \neq J_{\ell,k+1}(0)$, if we put

$$n = \min\{n \in \mathbb{N}_{>0} \mid nF_{\ell,k} \in I\},$$

then we can show easily

$$I = J_{\ell,k+1}(0) + (nF_{\ell,k}).$$

\square

Proposition 6.11. *In Ω_{H_ℓ} , there are exactly $\ell + 1$ prime ideals*

$$0 \subsetneq \mathcal{L}(0) \subsetneq \mathcal{L}^2(0) \subsetneq \dots \subsetneq \mathcal{L}^\ell(0)$$

over $(0) \subseteq R_0$.

Here, $\mathcal{L}^k(0)$ is the ideal obtained by a k -times application of \mathcal{L} to $0 \subseteq \Omega_{H_{\ell-k}}$, for each $0 \leq k \leq \ell$.

Proof. By Corollary 4.8 and Proposition 6.9, each $\mathcal{L}^k(0)$ is prime.

We show the proposition by an induction on ℓ . Suppose we have done for $\ell - 1$, and take any prime ideal $\mathcal{S} = [I_0, \dots, I_\ell] \subseteq \Omega_{H_\ell}$ over $0 \subseteq R_0$. Since $\mathcal{S}|_{H_{\ell-1}}$ is prime by Corollary 4.7, we have $\mathcal{S}|_{H_{\ell-1}} = \mathcal{L}^k(0)$ on $H_{\ell-1}$ for some $0 \leq k \leq \ell - 1$, namely

$$[I_0, \dots, I_{\ell-1}] = [0, \dots, 0, L(0), \dots, L^k(0)].$$

If $k \geq 1$, then Proposition 6.8 shows

$$S(I_{\ell-1}) = SL^k(0) = L^{k+1}(0) = L(I_{\ell-1}),$$

and thus I_ℓ should be equal to $L^{k+1}(0)$. This means $\mathcal{S} = \mathcal{L}^{k+1}(0)$.

If $k = 0$, then I_ℓ should satisfy

$$0 \subseteq I_\ell \subseteq L_\ell(0) = (F_{\ell, \ell-1}).$$

By Lemma 6.10, there exists $n \in \mathbb{N}_{\geq 0}$ satisfying $I_\ell = (nF_{\ell, \ell-1})$. Thus it suffices to show that $\mathcal{S} = [0, \dots, 0, (nF_{\ell, \ell-1})] \subseteq \Omega_{H_\ell}$ is not prime unless $n = 0$ or 1 .

Suppose \mathcal{S} is prime for some $n \neq 0$. Then for $a = F_{\ell, \ell-1}$ and $b = n \in R_0 \setminus (0)$, since we have

$$\begin{aligned} a \cdot \text{jnd}_0^\ell(\ell) &= F_{\ell, \ell-1} \cdot \left(n + \sum_{i=0}^{\ell-1} m_i X_{\ell, i} \right) \quad (\text{for some } m_i \in \mathbb{Z}) \\ &= nF_{\ell, \ell-1} \in I_\ell, \end{aligned}$$

we obtain $F_{\ell, \ell-1} \in I_\ell$ by $\mathcal{P}(\ell)$. This implies $I_{\ell-1} = J_{\ell, \ell-1}(0)$ and $n = 1$. \square

6.4. Total picture. Putting Propositions 6.4, 6.7, 6.11 together, we obtain the following.

Theorem 6.12. *Let G be a cyclic p -group of p -rank r . The prime ideals of $\Omega_G \in \text{Ob}(\text{Tam}(G))$ are as follows.*

(i) *Over $(q) \subseteq R_0$, there are $\ell + 1$ prime ideals*

$$\mathcal{S}^r(q) \subsetneq \mathcal{L}\mathcal{S}^{r-1}(q) \subsetneq \dots \subsetneq \mathcal{L}^r(q).$$

(ii) *Over $(p) \subseteq R_0$, there is only one prime ideal $\mathcal{L}^r(p)$.*

(iii) *Over $0 \subseteq R_0$, there are $\ell + 1$ prime ideals*

$$0 \subsetneq \mathcal{L}(0) \subsetneq \dots \subsetneq \mathcal{L}^r(0).$$

Thus we have

$$\begin{aligned} \text{Spec } \Omega_G &= \{\mathcal{L}^r(p)\} \cup \{\mathcal{L}^i(0) \mid 0 \leq i \leq r\} \\ &\quad \cup \{\mathcal{L}^i\mathcal{S}^{r-i}(q) \mid 0 \leq i \leq r, q \text{ is a prime different from } p\}. \end{aligned}$$

For each $0 \leq i \leq r$, there is an inclusion $\mathcal{L}^i(0) \subsetneq \mathcal{L}^i\mathcal{S}^{r-i}(q)$, while we have $\mathcal{L}^j(0) \not\subseteq \mathcal{L}^i\mathcal{S}^{r-i}(q)$ for any $j > i$. Moreover, we have

$$\mathcal{L}^r(0) \subseteq \mathcal{L}^r(p), \quad \mathcal{L}^r(p) \not\subseteq \mathcal{L}^r(q), \quad \mathcal{L}^r(q) \not\subseteq \mathcal{L}^r(p).$$

Corollary 6.13. *The longest sequence of prime ideals of Ω_G is of length $r + 1$, such as*

$$0 \subsetneq \mathcal{S}^r(q) \subsetneq \mathcal{L}\mathcal{S}^{r-1}(q) \subsetneq \dots \subsetneq \mathcal{L}^r(q),$$

and thus the dimension of $\text{Spec } \Omega_G$, which we denote simply by $\dim \Omega_G$, is

$$\dim \Omega_G = r + 1.$$

Let us draw some picture of $\text{Spec } \Omega_G$, by indicating the closures of points. If $r = 1$ and G is a group of prime order p , then the picture will become as follows (§3.7 in [8]).

Example 6.14. Let G be the group of order p . Then we have

$$\begin{aligned} \text{Spec } \Omega &= \{\mathcal{L}(p)\} \cup \{0\} \cup \{\mathcal{L}(0)\} \\ &\cup \{\mathcal{S}(q) \mid q \in \mathbb{Z} \text{ is prime, } q \neq p\} \cup \{\mathcal{L}(q) \mid q \in \mathbb{Z} \text{ is prime, } q \neq p\}. \end{aligned}$$

Inclusions are

$$\begin{aligned} (0) &\subsetneq \mathcal{L}(0) \subsetneq \mathcal{L}(p) \\ \mathcal{S}(q) &\subsetneq \mathcal{L}(q) \quad (q \neq p). \end{aligned}$$

Especially, the dimension is 2. $\mathcal{L}(p)$ and $\mathcal{L}(q)$'s are the closed points, and 0 is the generic point in $\text{Spec } \Omega$. If we represent the points in $\text{Spec } \Omega$ by their closures, $\text{Spec } \Omega$ with fibration F can be depicted as in Figure 1.





FIGURE 1. $\text{Spec } \Omega$ for $G = \mathbb{Z}/p\mathbb{Z}$

A similar description is also possible for an arbitrary r . $\mathcal{L}^r(p)$ and $\mathcal{L}^r(q)$'s are the closed points, and 0 is the generic point in $\text{Spec } \Omega$. The picture becomes as in Figure 2.


 FIGURE 2. $\text{Spec } \Omega$ for $G = \mathbb{Z}/p^r\mathbb{Z}$

 7. APPENDIX: AN INDUCTIVE CALCULATION OF jnd , WHEN G IS ABEL

We demonstrate how to calculate the multiplicative transfers of Ω inductively, when G is an abelian group. This will show Corollary 2.4. A detailed investigation of an exponential map will be found in [12]. A more clear argument using Möbius inversion will be found in [9].

Since $\text{jnd}_H^L: \Omega_L(L/H) \rightarrow \Omega_L(L/L)$ and $\text{jnd}_H^L: \Omega_G(G/H) \rightarrow \Omega_G(G/L)$ are essentially equal for $H \leq L \leq G$, we may assume $L = G$ from the first, and calculate $\text{jnd}_H^G: \Omega_G(G/H) \rightarrow \Omega_G(G/G)$ for any $H \leq G$. Remark that any element in $\Omega_H(H/H)$ is of the form $\sum_{I \leq H} m_I H/I$ for some $m_I \in \mathbb{Z}$ ($I \leq H$), and is identi-

fied with $(\sum_{I \leq H} m_I G/I \xrightarrow{\text{Pr}} G/H) \in \Omega_G(G/H)$, where Pr is the sum of the natural

projections $G/I \xrightarrow{\text{Pr}_I^H} G/H$.

Remark that $\text{jnd}_H^G: \Omega_G(G/H) \rightarrow \Omega_G(G/G)$ is a polynomial map [13] as in the following. (As for polynomial maps, see [3].)

Fact 7.1. There exist polynomials

$$P_K \in \mathbb{Z}[m_I \mid I \leq H] \quad (K \leq G)$$

which satisfy

$$(7.1) \quad \text{jnd}_H^G(\sum_{I \leq H} m_I G/I \xrightarrow{\text{Pr}} G/H) = \sum_{K \leq G} P_K(\{m_I\}_{I \leq H}) G/K$$

for any $\{m_I\}_{I \leq H}$.

In particular if we obtain polynomials P_K which satisfy (7.1) whenever $\{m_I\}_I$ satisfies $m_I \geq 0$ ($\forall I \leq H$), then (7.1) should hold for any $\{m_I\}_{I \leq H}$ with these P_K 's. Now we calculate P_K , for each $K \leq G$. Let $\sum_{I \leq H} m_I H/I \in \Omega_H(H/H)$ be any element satisfying $m_I \geq 0$ ($\forall I \leq H$), and denote the corresponding G -set $\coprod_{I \leq H} (\coprod_{m_I} G/I)$ by

A. By the definition of $\text{jnd}_H^G = \Omega_\bullet(\text{pr}_H^G)$, the G -set $S = \text{jnd}_H^G(A \xrightarrow{\text{Pr}} G/H)$ is given by

$$\begin{aligned} S &= \{(y, s) \mid y \in G/G, s \in \text{Map}((\text{pr}_H^G)^{-1}(y), A), \text{Pr} \circ s = \text{id}_{G/H}\} \\ &= \{s \in \text{Map}(G/H, A) \mid \text{Pr} \circ s = \text{id}_{G/H}\} \\ &= \{s \in \text{Map}(G/H, A) \mid s \text{ is a section of Pr}\} \end{aligned}$$

(remark that G/G consists of only one element). S is equipped with a G -action

$${}^g s(x) = gs(g^{-1}x) \quad (\forall g \in G, \forall s \in S, \forall x \in G/H).$$

Definition 7.2. Let G, H, A, S be as above. For each $K \leq G$, define $c(K) \in \mathbb{Z}$ by

$$c(K) = \#\{s \in S \mid G_s = K\}.$$

Then, since G is abel, the number of G -orbits in S isomorphic to G/K should be equal to $\frac{c(K)}{|G:K|}$, and thus we have

$$S = \sum_{K \leq G} \frac{c(K)}{|G:K|} G/K,$$

namely, $P_K = \frac{c(K)}{|G:K|}$. Thus it remains to calculate $c(K)$.

Since G is abel, we have the following.

Remark 7.3. If we decompose G/K into K -orbits

$$G/H = X_1 \amalg \cdots \amalg X_{r(K)},$$

then $r(K) = |K \backslash G/H| = |G:KH|$, and each X_i is isomorphic to

$$(7.2) \quad KH/H \cong K/(K \cap H).$$

Fix an element $x_i \in X_i$ for each $1 \leq i \leq r(K)$. Then, we have the following.

Proposition 7.4. *For any element $s \in S$, the following are equivalent.*

- (1) $K \leq G_s$.
- (2) For any $1 \leq i \leq r(K)$ and any $k \in K$, we have

$$(7.3) \quad ks(x_i) = s(kx_i).$$

Proof. By definition, $K \leq G_s$ holds if and only if for any $1 \leq i \leq r(K)$, any $x \in X_i$ and any $k \in K$,

$$(7.4) \quad {}^k s(x) = s(x)$$

is satisfied. Since X_i is K -transitive, there is some $k' \in K$ satisfying $x = k'x_i$. Then (7.4) is written as

$$ks(k^{-1}k'x_i) = s(k'x_i).$$

This is easily shown to be equivalent to (2). \square

Corollary 7.5. *Let G, H, A, S be as above. For each $K \leq G$, there is a bijection*

$$\{s \in S \mid G_s \geq K\} \cong \prod_{1 \leq i \leq r(K)} \left(\prod_{I \leq H} \left(\prod_{m_I} \{s_i \in G/I \mid \text{pr}_I^H(s_i) = x_i, K_{s_i} = K_{x_i}\} \right) \right).$$

Proof. By Proposition 7.4, the set of $s \in S$ satisfying $G_s \geq K$ is determined by the values $s(x_i)$ ($1 \leq i \leq r(K)$). In fact, to a set of elements $s_1, \dots, s_{r(K)} \in A$, we may associate a map $s: G/H \rightarrow A$ satisfying $s(x_i) = s_i$, by using (7.3). This map s becomes well-defined if and only if $K_{x_i} \leq K_{s_i}$ is satisfied for any i . Moreover, we see that the following are equivalent.

- (i) s is a section of $\text{Pr}: A \rightarrow G/H$, and satisfies $G_s \geq K$.
- (ii) $\text{Pr}(s_i) = x_i$ ($1 \leq i \leq r(K)$), and $K_{s_i} = K_{x_i}$.

Thus, to give an element in $\{s \in S \mid G_s \geq K\}$ is equivalent to give a set of elements $s_1, \dots, s_{r(K)} \in A$ satisfying (ii). Namely, we have a bijection

$$\begin{aligned} \{s \in S \mid G_s \geq K\} &\cong \prod_{1 \leq i \leq r(K)} \{s_i \in A \mid \text{Pr}(s_i) = x_i, K_{s_i} = K_{x_i}\} \\ &= \prod_{1 \leq i \leq r(K)} \left(\prod_{I \leq H} \left(\prod_{m_I} \{s_i \in G/I \mid \text{pr}_I^H(s_i) = x_i, K_{s_i} = K_{x_i}\} \right) \right). \end{aligned}$$

□

Corollary 7.6. *For any $K \leq G$, we have*

$$\#\{s \in S \mid G_s \geq K\} = \left(\sum_{(K \cap H) \leq I \leq H} m_I |H : I| \right)^{r(K)}.$$

Consequently, we can calculate $c(K)$ inductively by

$$c(K) = \left(\sum_{(K \cap H) \leq I \leq H} m_I |H : I| \right)^{r(K)} - \sum_{K < L \leq G} c(L).$$

Proof. Since G is abelian, we have $K_{x_i} = K \cap H$ by (7.2). Similarly, for each $s_i \in G/I$, we have $K_{s_i} = K \cap I$. Since $I \leq H$, we obtain

$$K_{x_i} = K_{s_i} \Leftrightarrow K \cap H = K \cap I \Leftrightarrow K \cap H \leq I.$$

As a consequence, the bijection in Corollary 7.5 is reduced to

$$\{s \in S \mid G_s \geq K\} \cong \prod_{1 \leq i \leq r(K)} \left(\prod_{(K \cap H) \leq I \leq H} \left(\prod_{m_I} \{s_i \in G/I \mid \text{pr}_I^H(s_i) = x_i\} \right) \right).$$

Since the fiber $(\text{pr}_I^H)^{-1}(x_i)$ has $|H : I|$ elements, it follows

$$\begin{aligned} \#\{s \in S \mid G_s \geq K\} &= \prod_{1 \leq i \leq r(K)} \left(\sum_{(K \cap H) \leq I \leq H} (m_I |H : I|) \right) \\ &= \left(\sum_{(K \cap H) \leq I \leq H} m_I |H : I| \right)^{r(K)}. \end{aligned}$$

□

Example 7.7. When $G = \mathbb{Z}/p^\ell \mathbb{Z}$, let $H_k \leq G$ be the subgroup of order p^k ;

$$e = H_0 < H_1 < \dots < H_\ell = G.$$

For $H = H_k$, we can determine $c(H_j)$ ($0 \leq j \leq \ell$) of $\text{jnd}_H^G \left(\sum_{0 \leq i \leq k} m_i H/H_i \right)$ as follows.

By Corollary 7.6, for each $0 \leq j \leq \ell$ we have

$$c(H_j) = \begin{cases} m_k & j = r, \\ (m_k)^{p^{\ell-j}} - (m_k)^{p^{\ell-j-1}} & k \leq j < r, \\ \left(\sum_{s=j}^k m_s p^{k-s} \right)^{p^{\ell-k}} - \left(\sum_{s=j+1}^k m_s p^{k-s} \right)^{p^{\ell-k}} & 0 \leq j < k, \end{cases}$$

which leads to Corollary 2.4.

REFERENCES

- [1] Bouc, S.: *Green functors and G-sets*, Lecture Notes in Mathematics, 1671, Springer-Verlag, Berlin (1977).
- [2] Brun, M.: *Witt vectors and Tambara functors*. Adv. in Math. **193** (2005) 233–256.
- [3] Dress, A.W.M.: *Operations in representation rings. Representation theory of finite groups and related topics* (Proc. Sympos. Pure Math., Vol. XXI, Univ. Wisconsin, Madison, Wis., 1970), pp. 39–45. Amer. Math. Soc., Providence, R.I., 1971.
- [4] Nakaoka, H.: *On the fractions of semi-Mackey and Tambara functors*. J. of Alg. **352** (2012) 79–103.
- [5] Nakaoka, H.: *Ideals of Tambara functors*. Adv. in Math. **230** (2012) 2295–2331.
- [6] Nakaoka, H.: *Tambaraization of a Mackey functor and its application to the Witt-Burnside construction*. Adv. in Math. **227** (2011) 2107–2143.
- [7] Nakaoka, H.: *A generalization of the Dress construction for a Tambara functor, and its relation to polynomial Tambara functors*. Adv. in Math. **235** (2013) 237–260.
- [8] Nakaoka, H.: *On the calculation of the spectra of Burnside Tambara functors*. Sūrikaiseikikenkyūsho Kōkyūroku No. 1811 (2012), 78–92.
- [9] Oda, F.: *Tensor induction for Burnside rings of the cyclic p-groups*, to appear.
- [10] Oda, F.; Yoshida, T.: *Crossed Burnside rings III: The Dress construction for a Tambara functor*, J. of Alg. **327** (2011) 31–49.
- [11] Tambara, D.: *On multiplicative transfer*. Comm. Algebra **21** (1993), no. 4, 1393–1420.
- [12] Yaman, A.: *On the exponential map of the Burnside ring*. Thesis, Bilkent University.
- [13] Yoshida, T.: *On the unit groups of Burnside rings*. J. Math. Soc. Japan **42** (1990), no. 1, 31–64.
- [14] Yoshida, T.: *Polynomial rings with coefficients in Tambara functors*. (Japanese) Sūrikaiseikikenkyūsho Kōkyūroku No. 1466 (2006), 21–34.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, KAGOSHIMA UNIVERSITY, 1-21-35
KORIMOTO, KAGOSHIMA, 890-0065 JAPAN

E-mail address: nakaoka@sci.kagoshima-u.ac.jp