Mobile Product and Zariski decomposition

MARIO PRINCIPATO

Università degli studi di Roma "Tor Vergata", Via della Ricerca Scientifica 1, 0133 Rome, Italy *email:principa@mat.uniroma2.it* phone-number: +39 06 72594648

October 17, 2018

Abstract

We explain the relationship between $\alpha_1 \cdots \alpha_q$ (standard cohomology product) and $(\alpha_1 \cdots \alpha_q)$ (mobile intersection product) of pseudo-effective classes $\alpha_1, \ldots, \alpha_q$ on a compact Kähler manifold. We also show how to use this relationship for proving some holomorphic Morse inequalities. Then we prove a result concerning the direct image of Lelong numbers under a modification in dimension 3, deriving a continuity property for the Lelong numbers of the wedge of (1, 1)-currents.

Introduction

Since the work of O. Zariski [Zar62] the study of the ring

$$R(X,D) := \bigoplus_{k \ge 0} H^0(X, \mathcal{O}(kD)),$$

where D is an effective divisor on a projective surface X, became very important.

In particular the main result is that any Q-divisor D on a projective surface X can be decomposed into a sum D = P + N where P is a nef Q-divisor, $N = \sum a_j D_j$ is an effective Q-divisor such that $(D_i \cdot D_j)$ is negative definite, and P is orthogonal to N with respect to the intersection form. Then Zariski showed that $H^0(kP) \hookrightarrow H^0(kD)$ is an isomorphism in that case. Using the metrics with minimal singularities on a pseudo-effective line bundle L introduced by J.-P. Demailly, S. Boucksom [Bou04] defines the divisorial Zariski decomposition of a pseudo-effective class $\alpha = (\alpha) + N(\alpha)$ where $N(\alpha)$ is an effective \mathbb{R} -divisor which is "exceptional" in some sense. In this work we are going to study the relationship between the product α^i and (α^i) with respect the non-nef locus of $\alpha E_{nn}(\alpha)$ as defined in ([Bou04]), where (α^i) is the mobile intersection product introduced in ([BDPP12]). Let \mathcal{E} be the cone of pseudo-effective classes, we can prove **Proposition 1** Let (X, ω) be a compact complex Kähler manifold of complex dimension n and let $\alpha_1, \ldots, \alpha_q \in \mathcal{E}$ such that $cod(E_{nn}(\alpha_1) \cup \cdots \cup E_{nn}(\alpha_q)) = m.$

Then if s < m and $j_1 < j_2 < \cdots < j_s \in \{1, \cdots, q\}$ we have

$$\alpha_{j_1} \cdots \alpha_{j_s} = (\alpha_{j_1} \cdots \alpha_{j_s}) \tag{1}$$

In particular when $\alpha = \alpha_1 = \cdots = \alpha_q$ and $cod(E_{nn}(\alpha)) = q$ then

$$\alpha^{j} = \langle \alpha^{j} \rangle = (\alpha^{j}) \quad \forall j = 1, \dots, q-1.$$
⁽²⁾

Furthermore we can study the extremal case of Proposition above, and introducing for $0 \le q \le n$ the cone

$$\mathcal{M}^q := \{ \alpha \in \mathcal{E} : \dim(E_{nn}(\alpha)) \le n - q \},\$$

then we have the following

Proposition 2 $\mathcal{M}^q \subseteq \mathcal{E}$ is a closed convex cone.

Moreover one has

Proposition 3 $E_{nn}(\alpha)$ doesn't have irreducible components of dimension zero.

Thus we can prove the following

Theorem 1 Let X be a compact Kähler manifold and be $\alpha_1, \ldots, \alpha_q \in \mathcal{M}^q$.

Let $\{Y_h\}_{h\in H}$ the family of codimension q components of $\bigcap_{j=1}^{r} E_{nn}(\alpha_j)$. Then we

have the following decomposition:

$$i) \ \alpha_1 \cdots \alpha_q = \langle \alpha_1 \cdots \alpha_q \rangle + \left\{ \sum_{h \in H} \nu(\alpha_1 \cdots \alpha_q, Y_h) [Y_h] \right\}, \ \nu(\alpha_1 \cdots \alpha_q, Y_h) > 0$$

where $\nu(\alpha_1 \cdots \alpha_q, Y_h)$ are the multiplicities of $\alpha_1 \cdots \alpha_q$ (see Definition 1.16), and the set H is at most countable;

ii) if $\alpha_j \in int(\mathcal{M}^q)$ for $1 \leq j \leq q$, then the set H is finite and $\nu(\alpha_1 \cdots \alpha_q, Y_h) = \nu(T_{min,\alpha_1} \wedge \cdots \wedge T_{min,\alpha_q}, Y_h)$ for all $h \in H$, where T_{min,α_j} is a positive current with minimal singularities in α_j .

As an immediate consequence of **Theorem 1** we have

Corollary 1 If $\alpha \in int(\mathcal{M}^q)$ then $\nu(T_{min}^q, Y) = 0$ for every irreducible analytic set of codimension q not completely contained in $E_{nn}(\alpha)$.

We can give a partial converse of this Theorem (1.6 (ii))

Proposition 4 Let (X, ω) be a compact Kähler manifold of complex dimension n, and let $\alpha \in \mathcal{E}$. Then $\alpha \in \mathcal{M}^{q+1}$ if and only if $\alpha^j = (\alpha^j)$ for all $j = 1, \ldots, q$.

In particular α is nef if and only if $\alpha^j = (\alpha^j)$ for all $j = 1, \dots, n-1$.

Then we focused our attention to the Holomorphic Morse Inequalities, a theory initiated by J.-P Demailly in the '80's.

Let's suppose we have an hermitian line bundle L over a compact Kähler manifold and we want to study the asymptotic behavior (for $k \to +\infty$) of the partial alternating sum of $h^q(X, kL)$; the complete sum is simply the Euler characteristic. In general the behavior is controlled by an estimate involving the integral of the top wedge power of the Chern curvature of L, extended over suitable subsets of X. One difficulty in the application of these inequalities is that the curvature integral is in general quite uneasy to compute, it is neither a topological nor an algebraic invariant. However a special case of the Morse inequalities can be reformulated in a more algebraic setting in which only algebraic invariants are involved, see e.g. [Tra95], [Siu93]. Now by using the decomposition

$$\alpha_1 \cdots \alpha_q - \langle \alpha_1 \cdots \alpha_q \rangle = \left\{ \sum_{h=1}^t \nu(\alpha_1 \cdots \alpha_q, Y_h)[Y_h] \right\}$$

we can prove

Theorem 2 Let X be a compact projective manifold of complex dimension n. Let L and F be two line bundles over X with L nef and $F \in \mathcal{M}^s$ with $\dim(E_{nn}(F)) = n - s$. Let $\{Y_t\}_{t \in T}$ be the irreducible components (possibly infinite) of codimension s of $E_{nn}(\alpha)$, and let ν_t and ν'_t be the multiplicities of F^s and the multiplicity of F along Y_t respectively. Then we have the following strong Morse inequalities

$$\sum_{j=0}^{s} (-1)^{s-j} h^{j}(X, k(L-F)) \leq \frac{k^{n}}{n!} (\sum_{j=0}^{s} \binom{n}{j} (-1)^{s-j} L^{n-j}(F^{j}) + \sum_{t=1}^{+\infty} \binom{n}{s} (L + \nu_{t}^{'}\{u\})^{n-s} \nu_{t}[Y_{t}]) + o(k^{n}).$$

And as a consequence we have

Corollary 2 Let X be a compact projective manifold of complex dimension n. Let L and F be two line bundles over X with L nef and $F \in \mathcal{M}^s$ with $\dim(E_{nn}(F)) = n - s$. Then we have the following Morse inequalities

$$\sum_{j=0}^{s} (-1)^{s-j} h^{j}(X, k(L-F)) \leq \frac{k^{n}}{n!} \sum_{j=0}^{s} {n \choose j} (-1)^{s-j} L^{n-j}(F^{j}) + {n \choose s} (L+b\{u\})^{n-s} (F^{s} - \langle F^{s} \rangle) + o(k^{n}),$$

where $b := \max_{t=1,\ldots,N} \nu_t$.

In the last Section we restrict to the dimension 3 case and we prove the following result about the direct image of Lelong numbers under a modification,

Theorem 3 Let X be a complex compact manifold with $\dim_{\mathbb{C}}(X) = 3$. Let $\tilde{\mu} : \tilde{X} \to X$ be a modification of X and Ω is a smooth, positive form on \tilde{X} of bidimension 1. Then $\nu(\tilde{\mu}_*(\tilde{\Omega}), Y) = 0 \ \forall Y$ irreducible curve on X, where ν is the generic Lelong number.

Thus we can prove the following continuity property for Lelong numbers

Proposition 5 Let $\alpha, \beta \in int(\mathcal{M}^2)$. Then

$$\lim_{k \to +\infty} \nu(T_{k,\alpha} \wedge T_{k,\beta}, x) = \nu(T_{\min,\alpha} \wedge T_{\min,\beta}, x) \ \forall \ x \in X,$$

where $\{T_{k,\alpha}\}$ and $\{T_{k,\beta}\}$ are sequences of currents with analytic singularities obtained in [Dem92] which converge weakly to $T_{min,\alpha}$ and $T_{min,\beta}$ respectively.

Acknowledgements

I would like thank my advisor Stefano Trapani for introducing me to the topic of this article, for his great knowledge and inspiration, and for his continuous support.

1 **Decomposition(s) in** $H^{k,k}_{>0}(X)$

We need two basic types of regularizations (inside a fixed cohomology class) for (1, 1)-current, both due to J.-P. Demailly [Dem92]

Theorem 1.1 Let $T = \alpha + i\partial\bar{\partial}\varphi$ be a closed almost positive (1,1)-current on a compact complex manifold X and fix an Hermitian form ω . Suppose that $T \ge \gamma$ for some smooth real (1,1)-form γ on X. Then: i) There exists a sequence of smooth forms θ_k in $\{T\}$ (cohomology class of T) which converges weakly to T and such that $\theta_k \ge \gamma - C\lambda_k \omega$ where C is a constant depending on the curvature of (T_X, ω) only, and λ_k is a decreasing sequence of continuous functions such that $\lambda_k(x) \to \nu(T, x)$ for every $x \in X$. ii) There exists a sequence $T_k = \alpha + i\partial\bar{\partial}\varphi_k$ of closed currents such that:

• φ_k (and thus T_k) is smooth on the complement $X \setminus Z_k$ of an analytic set Z_k such that there is an increasing sequence

$$Z_0 \subset Z_1 \subset \cdots \subset X.$$

- There is a uniform estimate $T_k \ge \gamma \varepsilon_k \omega$ with $\lim \downarrow \varepsilon_k = 0$ as k tends to $+\infty$.
- The sequence (φ_k) is non increasing, and we have lim ↓ φ_k = φ. As a consequence, T_k → T.
- Near Z_k, the potential φ_k has logarithmic poles, namely, for every x₀ ∈ Z_k, there is a neighborhood U of x₀ such that φ_k(z) = λ_k log Σ_l |g_{k,l}|² + O(1) for suitable holomorphic functions (g_{k,l}) on U and λ_k > 0. Moreover, there is a (global) proper modification μ_k: X_k → X of X, obtained as a sequence of blow-ups with smooth centers, such that φ_k ∘ μ_k can be written locally on X_k as

$$\varphi_k \circ \mu_k(w) = \lambda_k \left(\sum n_l |\tilde{g}_l|^2 + f(w) \right)$$

where $(\tilde{g}_l = 0)$ are local generators of suitable (global) divisor D_l on X_k such that $\sum D_l$ has normal crossings, n_l are positive integers and the f's are smooth functions on X_k .

It is important to observe that such μ_k are obtained blowing up along the multiplier ideal sheaves $\mathcal{I}(k\varphi)$ defined as

$$\mathcal{I}(k\varphi)_x := \left\{ f \in \mathcal{O}_{\Omega,x} : \ |f|^2 e^{-2k\varphi} \in \mathcal{L}^1_{loc} \right\}$$

where $\Omega \subseteq X$ is open.

Following [Bou04], when $\alpha \in \mathcal{E}$ one can introduce a measure of nefness of α . Let ψ_1, ψ_2 be almost plurisubharmonic functions on X we say that ψ_1 is less singular than ψ_2 (and write $\psi_1 \leq \psi_2$) if $\psi_2 \leq \psi_1 + C$ for some constant C. When S_1 and S_2 are closed almost positive (1,1)-currents on X, we can compare their singularities by comparing those of their local potentials ψ_1, ψ_2 . For each $\varepsilon > 0$ let $T_{min,\varepsilon}$ be a current with minimal singularities in $\alpha[-\varepsilon\omega]$ which is the set of closed almost positive (1,1)-currents T lying in α with $T \geq -\varepsilon\omega$.

Definition 1.2 The minimal multiplicity at $x \in X$ of the pseudo-effective class α is defined as

$$\nu(\alpha, x) = \sup_{\varepsilon > 0} \nu(T_{\min,\varepsilon}, x).$$

When D is a prime divisor, we define the generic minimal multiplicity of α along D as

$$\nu(\alpha, D) = \inf\{\nu(\alpha, x), x \in D\}.$$

We then have $\nu(\alpha, D) = \sup_{\varepsilon > 0} \nu(T_{\min,\varepsilon}, D)$, and $\nu(\alpha, D) = \nu(\alpha, x)$ for the very general $x \in D$.

One can give the following

Definition 1.3 The non-nef locus of a pseudo-effective class α is defined by

$$E_{nn}(\alpha) := \{ x \in X, \ \nu(\alpha, x) > 0 \}.$$

Remark 1.4 Let α be a pseudo-effective cohomology class and be ε small positive rational number. Let's fix a smooth hermitian form ω on X so that $T_{\min,\varepsilon} + \varepsilon \omega \ge a$ positive current representing $\alpha + \varepsilon \omega$. Then $T_{\min,\varepsilon} + \varepsilon \omega \ge T_{\min,\alpha+\varepsilon \omega}$ where $T_{\min,\alpha+\varepsilon \omega}$ is a positive current with minimal singularities in $\alpha + \varepsilon \omega$. On the other hand $T_{\min,\alpha+\varepsilon \omega} - \varepsilon \omega$ is a current representing α such that $T_{\min,\alpha+\varepsilon \omega} - \varepsilon \omega \ge -\varepsilon \omega$ then $T_{\min,\alpha+\varepsilon \omega} - \varepsilon \omega \ge T_{\min,\varepsilon}$. Hence we infer that $T_{\min,\varepsilon} + \varepsilon \omega$ is a positive current with minimal singularities representing $\alpha + \varepsilon \omega$.

Now let X be a compact Kähler manifold and let $Y \subseteq X$ be an analytic set of dimension p and let $\alpha_1, \ldots, \alpha_p \in H^{1,1}(X, \mathbb{R})$ be pseudo-effective classes. Then one can define a "positive" number of intersection $(\alpha_1 \cdots \alpha_p \cdot Y)_{>0}$ which intuitively should be equal to the usual intersection number $(\beta_1 \cdots \beta_p \cdot Y)$, where β_i is the nef part of α_i in its Zariski decomposition. But in general the Zariski decomposition does not exist, however using the currents with analytic singularities, one can solve the problem. More generally one can consider a closed positive current Θ instead of Y, then one can give the following

Definition 1.5 ([Bou04]) The mobile intersection number $(\alpha_1 \cdots \alpha_p \cdot \Theta)_{>0}$ of α_i 's and Θ is defined as

$$\inf_{\varepsilon>0} \left(\sup \int_{X\setminus F} (T_1 + \varepsilon \omega) \wedge \dots \wedge (T_p + \varepsilon \omega) \wedge \Theta \right),$$

where $T_i \in \alpha_i[-\varepsilon\omega]$ are currents with analytic singularities and F is the union of its unbounded-loci.

Let's observe that the integrals in the definition above are all convergent and the supremum is finite because the integrals can be bounded in terms only of cohomology classes of currents, moreover the definition does not depend on the choice of the Kähler form ω . Finally the supremum increases with ε so that the limit exist and it is equal to the infimum for $\varepsilon > 0$. Then by duality we have

Theorem 1.6 ([BDPP12]) Let X be a compact Kähler manifold. We denote here by $H_{\geq 0}^{k,k}(X)$ the cone of cohomology classes of type (k,k) which have non-negative intersection with all closed semi-positive smooth forms of bidegree (n-k, n-k).

(i)For each k = 1, ..., n, there exists a canonical mobile intersection product

$$\mathcal{E} \times \cdots \times \mathcal{E} \to H^{k,k}_{>0}(X), \ (\alpha_1, \dots, \alpha_k) \to \langle \alpha_1 \cdot \alpha_2 \cdots \alpha_k \rangle$$

(ii) The product is increasing, homogeneous of degree 1 and superadditive in each argument, i.e.

$$\langle \alpha_{1}\cdots(\alpha_{j}^{'}+\alpha_{j}^{''})\cdots\alpha_{k}\rangle\geq \langle \alpha_{1}\cdots\alpha_{j}^{'}\cdots\alpha_{k}\rangle+\langle \alpha_{1}\cdots\alpha_{j}^{''}\cdots\alpha_{k}\rangle.$$

It coincides with the ordinary intersection product when the α_j are nef classes.

Remark 1.7 Let us note that by construction $(\alpha_1 \cdots \alpha_p) \cdot u = (\alpha_1 \cdots \alpha_p \cdot u)_{>0}$ for all closed semi-positive forms u.

Now using the definition of non-pluripolar product $\langle u_1 \cdots u_p \rangle$ ([BEGZ10]) for u_1, \ldots, u_p psh-function on an open set of X (here X is a compact complex manifold), one can define a cohomology class $\langle \alpha_1 \cdots \alpha_p \rangle \in H^{p,p}(X, \mathbb{R})$ as follows

Definition 1.8 Let $\alpha_1, \ldots, \alpha_p \in H^{1,1}(X, \mathbb{R})$ be big cohomology classes and let $T_{min,i} \in \alpha_i$ be a positive current with minimal singularities. Then the cohomology class of the non-pluripolar product $\langle T_{min,1} \wedge \cdots \wedge T_{min,p} \rangle$ is indipendent of the choice of $T_{min,i} \in \alpha_i$ with minimal singularities. It will be denoted by

$$\langle \alpha_1 \cdots \alpha_p \rangle \in H^{p,p}(X,\mathbb{R})$$

and called the **non** – **pluripolar product** of the α_j . If $\alpha_1, \ldots, \alpha_p \in \mathcal{E}$ one sets

 $\langle \alpha_1 \cdots \alpha_p \rangle = \lim_{\varepsilon \to 0} \langle (\alpha_1 + \varepsilon \beta) \cdots (\alpha_p + \varepsilon \beta) \rangle$

where β is an arbitrary Kähler class, using the continuity of the non-pluripolar product.

Now on a compact Kähler manifold there exist two possible products (different a priori) : the mobile intersection product and the positive product of pseudo-effective cohomology classes; actually, as it is remarked in [BEGZ10] with no proof, these products are equal. For the reader convenience we give a proof of the following well-known

Lemma 1.9 Let T be any closed positive (p, p)-current on X compact complex Kähler manifold. Then the Lelong number $\nu(T, x)$ of T can be bounded by a constant depending only on the $\partial \overline{\partial}$ -cohomology class of T.

Proof. Let ω be a Kähler form on X, one has by definition that $\nu(T, x)$ is (up to a constant depending on ω near x) the limit for $r \to 0^+$ of

$$\nu(T,x,r) := \frac{(n-p)!}{(\pi r^2)^{n-p}} \int_{B(x,r)} T \wedge \omega^{n-p},$$

known to be an increasing function of r. Thus if we choose r_0 small enough to ensure that each ball $B(x, r_0)$ is contained in a coordinate chart, we get $\nu(T, x) \leq \nu(T, x, r_0) \leq C \int_X T \wedge \omega^{n-p}$, a quantity depending only on the cohomology class of T.

Proposition 1.10 Let X be a compact Kähler manifold and let $\alpha_1, \ldots, \alpha_p \in H^{1,1}(X, \mathbb{R})$ be pseudo-effective classes. Then

$$(\alpha_1 \cdots \alpha_p) = \langle \alpha_1 \cdots \alpha_p \rangle. \tag{3}$$

Proof. To keep notations simple we assume $\alpha = \alpha_1 = \cdots = \alpha_p$. Let's suppose first that α is big. By remark (1.7) this is equivalent to showing that:

$$(\alpha^p \cdot u)_{>0} = \int_X \langle T^p_{min} \rangle \wedge u$$

for all closed semi-positive smooth forms u of bidegree n - p, where T_{min} is a positive current with minimal singularities in α . Let $T_k \in \alpha[-\varepsilon_k \omega]$ be a sequence of currents with analytic singularities such that $T_k \rightharpoonup T_{min}$. Set $S_k := (1 - \varepsilon_k)T_k + \varepsilon_k S$ where S is a Kähler current with analytic singularities such that $E_+(S) = E_{nK}(\alpha)$, where

$$E_{nK}(\alpha) := \bigcap_{T \ge \varepsilon \omega, T \in \alpha} E_+(T)$$

is the non-Kähler locus of α as in [Bou04] and $E_+(S) = \bigcup_{c>0} \{x \in X : \nu(S, x) \ge c\}$. Then $\langle S_k^p \rangle \rightharpoonup \langle T_{min}^p \rangle$ on X from which we infer

$$\int_{X \setminus F_k} S_k^p \wedge u \le (\alpha^p \cdot u)_{>0}$$

and letting $k \to +\infty$

$$\int_X \langle T^p_{min} \rangle \wedge u \le (\alpha^p \cdot u)_{>0}.$$

But by the proof of theorem (1.6) we can find a sequence of Kähler currents $R_k \in \alpha$ with analytic singularities such that

$$\int_{X \setminus G_k} R_k^p \wedge u \xrightarrow{k \to +\infty} (\alpha^p \cdot u)_{>0}$$

where G_k is the unbounded-locus of R_k . Then we find that

$$\int_{X \backslash G_k} R_k^p \wedge u \leq \int_X \langle T_{min}^p \rangle \wedge u$$

and letting $k \to +\infty$ we find the other inequality.

If $\alpha_1, \ldots, \alpha_p$ are merely pseudo-effective since $(\alpha_1 \cdots \alpha_p)$ and $\langle \alpha_1 \cdots \alpha_p \rangle$ depend continuously on the *p*-tuple $\alpha_1, \ldots, \alpha_p$ of big classes the statement holds.

Now let $\alpha \in H^{1,1}(X, \mathbb{R})$ be a pseudo-effective class, we want to study the relation between α^p (the standard product in cohomology) and $\langle \alpha^p \rangle = (\alpha^p)$ when the codimension of $E_{nn}(\alpha) > p$. In general we have

Proposition 1.11 Let $\alpha_1, \ldots, \alpha_q$ be pseudo-effective classes on compact complex Kähler manifold (X, ω) of complex dimension n, such that $cod(E_{nn}(\alpha_1) \cup \cdots \cup E_{nn}(\alpha_q)) = m$ for all $s_1 < \cdots < s_t \in \{1, \ldots, q\}$. Then

$$\alpha_{s_1} \cdots \alpha_{s_t} = (\alpha_{s_1} \cdots \alpha_{s_t}) \tag{4}$$

for all t < m. In particular when $\alpha = \alpha_1 = \cdots = \alpha_q$ and $cod(E_{nn}(\alpha)) = q$ then

$$\alpha^{j} = \langle \alpha^{j} \rangle = (\alpha^{j}) \quad \forall j = 1, \dots, q-1.$$
(5)

Proof. Let $\{T_{k,s_i}\} \in \alpha_{s_i}$ be a sequence of currents with analytic singularities as in Theorem (1.1) such that $T_{k,s_i} \rightharpoonup T_{min,s_i}$ for $i = 1, \ldots, t$, where $T_{min,s_i} \in \alpha_{s_i}$ is a current with minimal singularities with local potentials φ_{min,s_i} . Let's define for all $i = 1, \ldots, t$

$$S_{k,s_i} := T_{k,s_i} + \varepsilon_{k,s_i} \omega \ge 0$$

and let $\mu_k : X_k \to X$ a common log-resolution for S_{k,s_i} obtained by blowing-up along $V = V(\mathcal{I}(k\varphi_{min,s_1})) \cup \cdots \cup V(\mathcal{I}(k\varphi_{min,s_t}))$, such that for all $i = 1, \ldots, t$

$$\mu_k^* S_{k,s_i} = [E_{k,s_i}] + \beta_{k,s_i}$$

where $\beta_{k,s_i} \geq 0$ and smooth. Then

$$S_{k,s_1} \wedge \dots \wedge S_{k,s_t} - (\mu_k)_* (\beta_{k,s_1} \wedge \dots \wedge \beta_{k,s_t}) = 0 \text{ on } X \setminus V,$$

and by using the support theorem for currents we get

$$S_{k,s_1} \wedge \dots \wedge S_{k,s_t} - (\mu_k)_* (\beta_{k,s_1} \wedge \dots \wedge \beta_{k,s_t}) = \sum_{h=1}^{+\infty} \lambda_{h,k} [A_{h,k}]$$

where $[A_{h,k}]$ are the irreducible components of codimension t in V, but $V \subseteq \bigcup_{i=1}^{t} E_{nn}(\alpha_{s_i})$ thus

$$S_{k,s_1} \wedge \dots \wedge S_{k,s_t} = (\mu_k)_* (\beta_{k,s_1} \wedge \dots \wedge \beta_{k,s_t})$$
on X.

So we can write the last equality as follows

$$T_{k,s_1} \wedge \dots \wedge T_{k,s_t} + O(\varepsilon_k) = (\mu_k)_* (\beta_{k,s_1} \wedge \dots \wedge \beta_{k,s_t}) \tag{6}$$

thanks to uniform control of the mass by cohomology classes. Passing in cohomology in (6) and letting k to $+\infty$, the statement holds.

Now we want to study the extremal case of Proposition (1.11), i.e. one can consider the difference $\alpha_1 \cdots \alpha_q - \langle \alpha_1 \cdots \alpha_q \rangle$. The case q = 1 is given by [Bou04]. We define for every $0 \le q \le n$

$$\mathcal{M}^q := \{ \alpha \in \mathcal{E} : \dim(E_{nn}(\alpha)) \le n - q \},\$$

then we have the following

Proposition 1.12 \mathcal{M}^q is a convex closed cone of \mathcal{E} for all $q = 0, \ldots, n$.

Proof. \mathcal{M}^q is convex as it follows from the convexity of the map $\mathcal{E} \to \mathbb{R}$, $\alpha \to \nu(\alpha, x)$. Now let α be in the closure of \mathcal{M}^q , for $\varepsilon > 0$ small enough $\alpha + \varepsilon \omega$ is in the interior of \mathcal{M}^q . By using Remark (1.4) one has that

$$E_{nn}(\alpha) = \bigcup_{\varepsilon > 0, \varepsilon \in \mathbb{Q}} E_{nn}(\alpha + \varepsilon \omega)$$

thus dim $(E_{nn}(\alpha)) \leq n-q$ then \mathcal{M}^q is closed. Finally we observe that for q = 0, 1 $\mathcal{M}^q = \mathcal{E}$, while for q = n $\mathcal{M}^n = \mathcal{N}$ the cone of nef classes.

As a consequence of the definition of non-pluripolar product one has

Corollary 1.13 If T_1, \ldots, T_q are closed positive (1, 1) – currents such that the union F of their unbounded locus is contained in an analytic set Y of pure dimension n - q. Then

$$\nu(\langle T_1 \wedge \dots \wedge T_q \rangle, Z) = 0 \tag{7}$$

for all irreducible components Z of dimension n - q in Y.

Proof. Let's consider the Siu decomposition of $\langle T_1 \wedge \cdots \wedge T_q \rangle = R + \sum_h \nu(\langle T_1 \wedge \cdots \wedge T_q \rangle, Z_h)[Z_h]$ then $Z_h \subseteq Y$ and we find that $\chi_{X-Y}\langle T_1 \wedge \cdots \wedge T_q \rangle = \chi_{X-Y}R$, then the extension by zero of $\chi_{X-Y}\langle T_1 \wedge \cdots \wedge T_q \rangle$ on Y is $\langle T_1 \wedge \cdots \wedge T_q \rangle$ then $R \ge \langle T_1 \wedge \cdots \wedge T_q \rangle$ and since $\nu(R, Z) = 0$ for every analytic set of dimension n - q we infer the statement. \Box

Proposition 1.14 Let T_j, T'_j be positive closed currents for j = 1, ..., q with $T_j \leq T'_j$, such that $T_1 \wedge \cdots \wedge T_q$ and $T'_1 \wedge \cdots \wedge T'_q$ are defined. Then

$$\nu(T_1 \wedge \dots \wedge T_q, x) \le \nu(T_1^{'} \wedge \dots \wedge T_q^{'}, x)$$
(8)

for all $x \in X$.

In particular if $T_{\min,j}$ and $T'_{\min,j}$ are currents with minimal singularities in α_j then $\nu(T_{\min,1} \wedge \cdots \wedge T_{\min,q}, x)$ does not depend in the chosen currents with minimal singularities.

Proof. We can choose local coordinates $z = (z_1, \dots, z_n)$ centered in x. Let $v_j \leq 0$ and $u_j \leq 0$ be the local potentials for T_j and T'_j respectively, then $v_j + C_j \geq u_j$ for some constants $C_j \geq 0$. Let's define

- $v'_{i} := v_{j} + \lambda \log |z|$
- $u'_{i} := u_{i} + \lambda \log |z|$

for $\lambda > 0$. Thus we find

$$v_j' + C_j \ge u_j'$$

then dividing by $u_{i}^{'}$ and passing to the \limsup we get

$$\limsup_{z \to x} \frac{v'_j(z)}{u'_j(z)} \le 1.$$
(9)

Applying ([Dem 93], Theorem 5.9) we find

$$\nu(\bigwedge_{j=1}^{q} dd^{c}v_{j}^{'}, x) \leq \nu(\bigwedge_{j=1}^{q} dd^{c}u_{j}^{'}, x).$$
(10)

In the calculation of the left-side hand in (10) we get a polynomial $p(\lambda)$ whose constant term a_0 is just q-times the wedge product of $dd^c v_j$, while for the right-side hand we get a polynomial $q(\lambda)$ whose constant term b_0 is just q-times the wedge product of $dd^c u_i$. Then letting $\lambda \to 0$ equation (8) holds. \Box

Remark 1.15 If $\alpha_1, \ldots, \alpha_q \in \mathcal{M}^q$ with then $\beta_i = \alpha_i + \varepsilon \omega$ are in the $int(\mathcal{M}^q)$ for all $1 \leq i \leq q$. If $0 < \varepsilon^{'} < \varepsilon$ then $T_{min,\alpha_j+\varepsilon\omega} \preceq T_{min,\alpha_j+\varepsilon^{'}\omega} + (\varepsilon - \varepsilon^{'})\omega$ then by Proposition (1.14) $\nu(\bigwedge_{j=1}^{q} T_{min,\alpha_j+\varepsilon\omega}, x)$ increases when $\varepsilon \to 0$ for all $x \in X$ and it is bounded in terms of $\alpha_1 \cdots \alpha_q \cdot \{\omega\}$.

Then we can give the following

Definition 1.16 The "multiplicity" of $\alpha_1, \ldots, \alpha_q$ at $x \in X$ is defined as

$$\nu(\alpha_1 \cdots \alpha_q, x) := \sup_{\varepsilon > 0} \nu(\bigwedge_{j=1}^q T_{\min,\alpha_j + \varepsilon\omega}, x)$$
(11)

and we define $\nu(\alpha_1 \cdots \alpha_q, Y) = \inf\{\nu(\alpha_1 \cdots \alpha_q, x), x \in Y\}.$

Proposition 1.17 If $\alpha_1, \ldots, \alpha_q$ are in the interior of \mathcal{M}^q . Then

$$\nu(\alpha_1 \cdots \alpha_q, x) = \nu(T_{\min,\alpha_1} \wedge \cdots \wedge T_{\min,\alpha_q}, x)$$

where T_{\min,α_i} is a positive current with minimal singularities in α_i .

Proof. It is clear that $\nu(\alpha_1 \cdots \alpha_q, x) \leq \nu(T_{\min,\alpha_1} \wedge \cdots \wedge T_{\min,\alpha_q}, x)$. For the other inequality let $S_j \in \alpha_j$ be a Kähler current such that $S_j \geq \omega$ and let $T_{\min,\varepsilon,j}$ be a minimal current in $\alpha_j[-\varepsilon\omega]$. Then by remark (1.4) we have that

$$\nu(\lambda S_j + (1 - \lambda)T_{\min,\varepsilon,j}, x) = \nu(\lambda S_j + (1 - \lambda)(T_{\min,\alpha_j + \varepsilon\omega} - \varepsilon\omega), x)$$

where the current $K_j = \lambda S_j + (1 - \lambda)(T_{\min,\alpha_j + \varepsilon \omega} - \varepsilon \omega) \ge 0$ is a positive current representing α_j , with $\lambda = \frac{\varepsilon}{1+\varepsilon}$. By using Proposition (1.14) we infer that

$$\nu(K_1 \wedge \dots \wedge K_q, x) \ge \nu(T_{\min,\alpha_1} \wedge \dots \wedge T_{\min,\alpha_q}, x).$$

Now by using the uniform bound of mass in terms of cohomology classes and letting $\varepsilon \to 0$ (hence $\lambda \to 0$), we obtain the other inequality.

Before proving **Theorem 1**, we recall that

Lemma 1.18 If $\alpha_1, \ldots, \alpha_q$ are in \mathcal{M}^q then

$$\bigcup_{\varepsilon>0,\varepsilon\in\mathbb{Q}}\bigcap_{j=1}^{q}E_{nK}(\alpha_j+\varepsilon\omega)=\bigcap_{j=1}^{q}\bigcup_{\varepsilon>0,\varepsilon\in\mathbb{Q}}E_{nK}(\alpha_j+\varepsilon\omega)=\bigcap_{j=1}^{q}E_{nn}(\alpha_j).$$
 (12)

Proof. For $\varepsilon > 0$ $\alpha_j + \varepsilon \omega \in \operatorname{int}(\mathcal{M}^q)$ (interior of \mathcal{M}^q) for all $j = 1, \ldots, q$, in particular $\alpha_j + \frac{\varepsilon}{2}\omega \in \operatorname{int}(\mathcal{M}^q)$. Let $\{T_{k,j}\} \in \alpha_j + \frac{\varepsilon}{2}\omega$ be a sequence of currents with analytic singularities as in Theorem (1.1) such that $T_{k,j} \rightharpoonup T_{\min,\alpha_j + \frac{\varepsilon}{2}\omega}$ where $T_{\min,\alpha_j + \frac{\varepsilon}{2}\omega}$ is a positive current with minimal singularities in $\alpha_j + \frac{\varepsilon}{2}\omega$. Now for k >> 0 $T_{k,j} + \frac{\varepsilon}{2}\omega$ are Kähler currents with analytic singularities, representing (as currents) $\alpha_j + \varepsilon \omega$, then we have the following inclusions

$$E_{nn}(\alpha_j + \varepsilon\omega) \subseteq E_{nK}(\alpha_j + \varepsilon\omega) \subseteq E_+(T_{k,j} + \frac{\varepsilon}{2}\omega) \subseteq E_{nn}(\alpha_j + \frac{\varepsilon}{2}\omega)$$
(13)

thus it follows

$$E_{nn}(\alpha_j) = \bigcup_{\varepsilon > 0, \varepsilon \in \mathbb{Q}} E_{nK}(\alpha_j + \varepsilon \omega).$$
(14)

Now thanks to (14) the equality (12) is equivalent to the following

$$\bigcup_{\varepsilon>0,\varepsilon\in\mathbb{Q}}\bigcap_{j=1}^{q}E_{nK}(\alpha_j+\varepsilon\omega)=\bigcap_{j=1}^{q}\bigcup_{\varepsilon>0,\varepsilon\in\mathbb{Q}}E_{nn}(\alpha_j+\varepsilon\omega).$$
(15)

Let's define A and B the left-hand side and the right-hand side of (15). Now if $x \in A \Rightarrow \exists \varepsilon_0 > 0$ such that $\forall j = 1, \ldots, q, x \in \bigcap_{j=1}^q E_{nK}(\alpha_j + \varepsilon_0 \omega)$ but $E_{nK}(\alpha_j + \varepsilon_0 \omega) \subseteq E_{nn}(\alpha_j + \frac{\varepsilon_0}{2}\omega) \; \forall j = 1, \ldots, q$ thanks to (13). Then $x \in \bigcup_{\varepsilon > 0} E_{nn}(\alpha_j + \varepsilon \omega) \; \forall j = 1, \ldots, q$, i.e. $x \in B$.

Viceversa if $x \in B$ then there exist $\varepsilon_1, \ldots, \varepsilon_q > 0$ such that $x \in E_{nn}(\alpha_j + \varepsilon_j \omega)$ for $j = 1, \ldots, q$. Let $\overline{\varepsilon} = \min_j \varepsilon_j$ then again thanks to (13) we have

$$E_{nn}(\alpha_j + \varepsilon_j \omega) \subseteq E_{nn}(\alpha_j + \overline{\varepsilon}\omega) \subseteq E_{nK}(\alpha_j + \overline{\varepsilon}\omega).$$

It follows that $x \in E_{nk}(\alpha_j + \bar{\varepsilon}\omega)$ for all $j = 1, \ldots, q$ then $x \in A$.

Using the previous Lemma we give a proof of the following well-known

Proposition 1.19 The set $E_{nn}(\alpha)$ does not have irreducible components of dimension zero if $\alpha \in \mathcal{E}$.

Proof. Let $x \in E_{nn}(\alpha)$ be an irreducible zero dimensional component. By Lemma (1.18) we have

$$E_{nn}(\alpha) = \bigcup_{\varepsilon > 0, \varepsilon \in \mathbb{Q}} E_{nK}(\alpha + \varepsilon \omega)$$

then x is an isolated point in $E_{nK}(\alpha + \varepsilon \omega)$ for all $\varepsilon > 0$ rational since $E_{nK}(\alpha + \varepsilon \omega)$ are analytic sets. Now let $\{T_k\} \in \alpha$ be a sequence of currents with analytic singularities as in Theorem (1.1) such that $T_k \rightarrow T_{min}$ weakly where T_{min} is a positive current with minimal singularities in α . Then for k >> 0 $T_k + \varepsilon \omega$ are Kähler currents with analytic singularities only in x representing $\alpha + \varepsilon \omega$ (as currents); hence by the gluing property of plurisubharmonic functions we can find new Kähler currents $S_k + \varepsilon \omega$ still representing $\alpha + \varepsilon \omega$ as currents, but with $\nu(S_k + \varepsilon \omega, x) = 0$, a contradiction. \Box

Now we can prove

Theorem 1.20 Let X be a compact Kähler manifold and be $\alpha_1, \ldots, \alpha_q \in \mathcal{M}^q$. Let $\{Y_h\}_{h \in H}$ the family of codimension q components of the set $\bigcap_{j=1}^q E_{nn}(\alpha_j)$. Then we have the following decomposition:

i)
$$\alpha_1 \cdots \alpha_q = \langle \alpha_1 \cdots \alpha_q \rangle + \left\{ \sum_{h \in H} \nu(\alpha_1 \cdots \alpha_q, Y_h) [Y_h] \right\}, with$$

 $\nu(\alpha_1 \cdots \alpha_q, Y_h) > 0 \ \forall h \in H;$

ii) if $\alpha_j \in int(\mathcal{M}^q)$ for $1 \leq j \leq q$, then the set H is finite and $\nu(\alpha_1 \cdots \alpha_q, Y_h) = \nu(T_{min,\alpha_1} \wedge \cdots \wedge T_{min,\alpha_q}, Y_h)$ for all $h \in H$.

Proof. ii) By the proof of Lemma (1.18) for $\varepsilon > 0$ small enough $\alpha_j - \varepsilon \omega \in int(\mathcal{M}^q)$ and let $\{T_{k,j}\} \in \alpha_j - \varepsilon \omega$ be a sequence of currents with analytic singularities as in Theorem (1.1) such that $T_{k,j} \rightharpoonup T_{min,\alpha_j-\varepsilon\omega}$ then

$$\nu(T_{k,j} + \varepsilon\omega, x) = \nu(T_{k,j}, x) \le \nu(T_{\min,\alpha_j - \varepsilon\omega}, x)$$

But $\alpha_j - \varepsilon \omega$ is still in $\operatorname{int}(\mathcal{M}^q)$ thus $E_+(T_{k,j} + \varepsilon \omega)$ is an analytic set of dimension $\leq n-q$ so the set of components of dimension n-q of $E_{nn}(\alpha_j)$ and of $E_{nK}(\alpha_j)$ is finite . Let $T_{min,j}$ be a positive current with minimal singularities in α_j , since $\dim_{\mathcal{H}} \left(\bigcup_i^q E_{nK}(\alpha_j) \right) \leq 2n - 2q$ (dim_{\mathcal{H}} is the Hausdorff dimension), then $T := T_{min,1} \wedge \cdots \wedge T_{min,q}$ is well defined on Xaccording to ([Dem92]) and by Proposition (1.17) $\nu(\alpha_1 \cdots \alpha_q, x) = \nu(T_{min,1} \wedge \cdots \wedge T_{min,q}, x)$ for all $x \in X$. Let's consider the positive current $T - \langle T \rangle$: this difference is identically zero outside $\bigcup_j^q E_{nK}(\alpha_j)$ and by the support theorem of positive currents it follows:

$$T - \langle T \rangle = \sum_{s=1}^{t} \lambda_s [Y_s]$$

where Y_s are the irreducible components of codimension q of $\bigcup_{j}^{q} E_{nK}(\alpha_j)$ and $\lambda_s \geq 0$. Now by Corollary (1.13)

$$T = \langle T \rangle + \sum_{h=1}^{t} \nu(T, Y_h)[Y_h]$$
(16)

is the Siu decomposition of T. Now if Y_{s_0} is contained in $\bigcap_{j=1}^{q} E_{nn}(\alpha_j)$ then by [Dem93]

$$\nu(T_{min,1} \wedge \dots \wedge T_{min,q}, Y_{s_0}) \ge \prod_{j=1}^q \nu(T_{min,j}, Y_{s_0}) > 0.$$

On the other hand let $Y_s \subseteq \bigcup_{j=1}^q E_{nK}(\alpha_j)$ be an irreducible component of codimension q not entirely contained in $\bigcap_{j=1}^q E_{nn}(\alpha_j)$. Then by Lemma (1.18)

there exists j_0 such that for all $\varepsilon > 0$ Y_s is not entirely contained in $\bigcup_{\varepsilon > 0, \varepsilon \in \mathbb{Q}} E_{nK}(\alpha_{j_0} + \varepsilon \omega)$. It is not restrictive to suppose $j_0 = 1$ then the local potential $\varphi_{min,1}$ of $T_{min,\alpha_1+\varepsilon\omega}$ is locally bounded in a neighborhood of a generic point $x \in Y_s$.

Let B = B(x, r) be an open ball of center x and radius r > 0 small enough such that B is contained in a coordinate patch with local coordinates $z = (z_1, \ldots, z_n)$; set $\beta = dd^c |z|^2$ and $S = \bigwedge_{i=2}^q dd^c \varphi_{min,i}$. Let χ be a smooth function with compact support in B such that $0 \le \chi \le 1$. Then

$$\int_{B} \chi dd^{c} \varphi_{min,1} \wedge S \wedge \beta^{n-q} = \int_{B} \varphi_{min,1} \cdot S \wedge dd^{c} \chi \wedge \beta^{n-q}$$

where the equality is obtained integrating by parts and using that S and β are closed.

By using Chern-Levine-Nirenberg inequalities ([Dem93], Proposition 1.3), we get

$$\int_{B} \varphi_{min,1} \cdot S \wedge dd^{c} \chi \wedge \beta^{n-q} \leq \operatorname{Vol}(B) ||S||_{X} C$$

where C is the product of the bound for the coefficients of the smooth form $dd^c \chi \wedge \beta^{n-q}$ and of the bound for $\varphi_{min,1}$, $\operatorname{Vol}(B)$ is the volume of B and $||S||_X$ is the mass of the current S that can be bounded in terms of $\alpha_2 \cdots \alpha_q \cdot \{\omega\}$. Using the fact that $\operatorname{Vol}(B) \sim r^{2n}$ and that

$$\nu(dd^c\varphi_{min,1} \wedge S, x) = \lim_{r \to 0} \frac{1}{2^{2(n-q)}r^{2(n-q)}} \int_{B(x,r)} dd^c\varphi_{min,1} \wedge S \wedge \beta^{n-q}$$

we can infer that $\nu(T, Y_s) = 0$. Then in equation (16) Y_h are actually irreducible components of codimension q of $\bigcap_{j=1}^q E_{nn}(\alpha_j)$. Then passing in cohomology and letting $\varepsilon \to 0$ the statement holds. To prove i) it is sufficient to observe that $\alpha_j + \varepsilon \omega \in \mathring{\mathcal{M}}^q$ for $1 \leq j \leq q$, applying ii) to $\alpha_j + \varepsilon \omega$ take the limit as $\varepsilon \to 0$ and using Lemma (1.18). \Box

Let's observe that if one of α_i 's is big but it is on the boundary of \mathcal{M}^q , it may be not true that the number of irreducible components of codimension q of $E_{nn}(\alpha_i)$ is finite. Since it could happen that $E_{nK}(\alpha_i)$ have irreducible components of codimension q-1, hence an infinite number of components of codimension q of $E_{nn}(\alpha_i)$ could be contained in one of these components. An immediate consequence of Theorem (1.20) is the following

Corollary 1.21 If $\alpha \in int(\mathcal{M}^q)$ then $\nu(T^q_{min}, Y) = 0$ for every irreducible analytic set of codimension q not completely contained in $E_{nn}(\alpha)$.

Another consequence of Theorem (1.20) is a partial converse of Theorem (1.6 (ii))

Proposition 1.22 Let (X, ω) be a compact Kähler manifold of complex dimension n. Then:

$$\alpha \in \mathcal{M}^q$$
 if and only if $\alpha^j = (\alpha^j)$ for all $1 \leq j \leq q-1$.

In particular, α is nef if and only if $\alpha^j = (\alpha^j)$ for all $1 \leq j \leq n-1$.

Proof. If $\alpha \in \mathcal{M}^q$ by Proposition (1.11) $\alpha^j = (\alpha^j)$ for all $1 \leq j < q$. To prove the vice versa we go by induction on q.

For q = 2, the statement follows from [Bou04]. Assume now the statement for a given q and let $\alpha^j = (\alpha^j)$ for $1 \le j \le q+1$, then by induction we know that $\alpha \in \mathcal{M}^{q+1}$, the by Theorem (1.20) we know that $\operatorname{cod}(E_{nn}(\alpha)) \ge q+2$, i.e. $\alpha \in \mathcal{M}^{q+2}$.

The last statement follows from Proposition (1.19).

2 Algebraic Morse inequalities in codimension "s"

Let X be a compact Kähler manifold, E a holomorphic vector bundle of rank r and L a line bundle over X. If L is equipped with a smooth metric of curvature form $\Theta(L)$, one defines the s-index set of L to be the open subset

$$X(s,L) = \begin{cases} x \in X : i\Theta(L)_x \text{ has } s \text{ negative eigenvalues} \\ n-s \text{ positive eigenvalues} \end{cases}$$

for $0 \leq s \leq n = \dim_{\mathbb{C}}(X)$. Hence X admits a partition $X = \Delta \cup \bigcup_{s} X(s, L)$ where $\Delta = \{x \in X : \det(\Theta(L)_x) = 0\}$. One also introduces

$$X(\leq s, L) = \bigcup_{0 \leq j \leq s} X(j, L).$$

It is shown in [Dem85] that the cohomology groups $H^s(X, E \otimes \mathcal{O}(kL))$ satisfy the following asymptotic weak Morse inequalities as $k \to +\infty$

$$h^{s}(X, E \otimes \mathcal{O}(kL)) \leq r \frac{k^{n}}{n!} \int_{X(s,L)} (-1)^{s} \left(\frac{i}{2\pi} \Theta(L)\right)^{n} + o(k^{n}).$$
(17)

A sharper form is given by the strong Morse inequalities

$$\sum_{j=0}^{s} (-1)^{s-j} h^{j}(X, E \otimes \mathcal{O}(kL)) \le r \frac{k^{n}}{n!} \int_{X(\le s,L)} (-1)^{s} \left(\frac{i}{2\pi} \Theta(L)\right)^{n} + o(k^{n}).$$
(18)

One difficulty in the application of these inequalities is that the curvature integral is in general quite uneasy to compute, it is neither a topological nor an algebraic invariant. However a special case of the Morse inequalities can be reformulated in a more algebraic setting in which only algebraic invariants are involved, se e.g. [Tra95], [Siu93].

Here we give an *algebraic* reformulation for the Holomorphic Morse inequalities proved by J.-P. Demailly ([Dem85]) in a more general setting.

Theorem 2.1 ([Dem00]) Let V = L - F be a holomorphic line bundle over a compact Kähler manifold X, where L and F are nef line bundle. Then for every s = 0, ..., n, there is an asymptotic strong Morse inequality

$$\sum_{0 \le j \le s} (-1)^{s-j} h^j(X, kV) \le \frac{k^n}{n!} \sum_{0 \le j \le s} (-1)^{s-j} \binom{n}{j} L^{n-j} F^j + o(k^n).$$

If F is not nef but it is just pseudo-effective with $\operatorname{codim}(E_{nn}(F)) \leq s+1$, then

Theorem 2.2 ([Tra11]) Let L and F be holomorphic line bundle over X a projective compact manifold, with L nef and $F \in \mathcal{E}$ such that $\dim(E_{nn}(F)) \leq n - (s+1)$. Then for $0 \leq m \leq s$ we have the following holomorphic Morse inequalities

$$\sum_{0 \le j \le m} (-1)^{m-j} h^j(X, k(L-F)) \le \frac{k^n}{n!} \sum_{0 \le j \le m} (-1)^{m-j} \binom{n}{j} L^{n-j} F^j + o(k^n)$$
(19)

However, when s = 1 there is a version of the algebraic Morse inequalities which uses the full divisorial Zariski decomposition of F

Theorem 2.3 ([Tra11]) Let L and F be line bundles over X, assume L nef and F pseudo-effective, let $F = (F) + \{N(F)\}$ be the divisorial Zariski decomposition of F with $N(\{F\}) = \sum_{j=1}^{N} \nu(F, D_j)[D_j]$, and let $\{u\}$ be a nef cohomology class in $H^2(X, \mathbb{R})$ such that $c_1(\mathcal{O}_{T_X}(1)) + \pi^*\{u\}$ is a nef cohomology class in $H^2(\mathbb{P}(T_X^*), \mathbb{R})$. Then

$$\limsup_{k \to +\infty} \frac{n!}{k^n} h^0(X, k(L-F)) \ge L^n - nL^{n-1}(F) - n\sum_{j=1}^N (L + \nu(F, D_j)\{u\})^{n-1} \nu(F, D_j) [D_j]$$
(20)

We can also treat the extremal case, i.e. when the codimension of $E_{nn}(F) = s$. We give two formulations.

Theorem 2.4 (First Formulation) Let X be a compact projective manifold of complex dimension n. Let L and F be two line bundles over X with L nef and $F \in \mathcal{M}^s$ with $\dim(E_{nn}(F)) = n - s$. Let $\{Y_t\}_{t \in T}$ be the irreducible components (possibly infinite) of codimension s of $E_{nn}(F)$, and let ν_t and ν'_t be the multiplicities of F^s and the multiplicity of F along Y_t respectively. Then we have the following Morse inequalities

$$\sum_{j=0}^{s} (-1)^{s-j} h^{j}(X, k(L-F)) \leq \frac{k^{n}}{n!} (\sum_{j=0}^{s} \binom{n}{j} (-1)^{s-j} L^{n-j}(F^{j}) + \sum_{t=1}^{+\infty} \binom{n}{s} (L + \nu_{t}^{'}\{u\})^{n-s} \nu_{t}[Y_{t}]) + o(k^{n}).$$
(21)

Proof. We first assume L to be ample and $F \in \operatorname{int}(\mathcal{M}^s)$. Since $F \in \operatorname{int}(\mathcal{M}^s)$, in particular F is big, then if $T_{min} \in c_1(F)$ is a current with minimal singularities then we have $\nu'(c_1(F), x) = \nu'(T_{min}, x)$ for all $x \in X$. We recall that there is a finite number of Y_j 's thanks to Theorem (1.20). Set $0 < \nu'_1 < \cdots < \nu'_h$, where ν'_1 is the minimum of ν'_j positive and so on. Let's choose c_i real positive numbers such that

$$0 < c_0 < \nu_1' < c_1 < \nu_2' < \dots < \nu_{h-1}' < c_{h-1} < \nu_h' < c_h < \nu := \max_{x \in X} (\nu(T_{min}, x)) + 1$$
(22)

Choose a positive closed (1,1)-form ω which is the curvature of a smooth metric on the ample line bundle L. Now by Theorem (1.1) we know that there exist a sequence of closed smooth forms T_k in the cohomology class of $c_1(F)$, a decreasing sequence of positive functions $\lambda_k(x)$, and a decreasing sequence of positive real numbers ε_k , with the following properties

- $T_k \rightharpoonup T_{min}$,
- $\lambda_k(x)$ converges to $\nu(T_{min}, x)$ for all $x \in X$,
- ε_k converges to 0,
- $T_k > -\lambda_k u 2\varepsilon_k \omega$.

Let c be a positive real number, then one can defines $O_{c} = \left\{ x \in V, x \in V \} \right\}$

 $\Omega_{k,c} = \{x \in X : \lambda_k(x) < c\}, v_{k,c} = 2\varepsilon_k + cu \text{ and } w_{k,c} = v_{k,c} + \omega.$ Then on $\Omega_{k,c}$ the forms $T_k + v_{k,c}$ and $w_{k,c}$ are positive. Now $\omega - T_k$ is the curvature of smooth metric on L - F. Let $\alpha_1 \leq \cdots \alpha_n$ be the eigenvalues of $\omega - T_k$ with respect to $w_{k,c}$, so that $\alpha_j \leq 1$ for all j. Let X(p) be the p-index set for L - F and $X(\leq p)$ be the set of points of index at most p. Then on X(p) we have $(-1)^p(\omega - T_k)^n \leq (-1)^p\alpha_1 \cdots \alpha_n w_{k,c}^n$. Now

$$\binom{n}{p}w_{k,c}^{n-p}\wedge(T_k+v_{k,c})^p = \binom{n}{p}w_k^{n-p}\wedge(w_{k,c}-(\omega-T_k))^p = \sigma_p(1-\alpha)w_{k,c}^n,$$

where $\sigma_p(1-\alpha)$ is the *p*-th elementary symmetric function in $1-\alpha_1,\ldots,1-\alpha_n$. However, since $\alpha_j < 1$ for $1 \leq j \leq n$, it follows that on X(p) we have

$$\sigma_p(1-\alpha) \ge (1-\alpha_1)\cdots(1-\alpha_p) \ge (-1)^p \alpha_1\cdots\alpha_p.$$

Furthermore one can easily prove by induction on n ([Dem00]) that

$$\sum_{j=0}^{p} (-1)^{p-j} \sigma_p(1-\alpha) \le \chi_{X(\le p)}(-1)^p \alpha_1 \cdots \alpha_n,$$

where $\chi_{X(\leq p)}$ is the characteristic function of the set $X(\leq p)$. Now we want to estimate $\int_{X(< s)} (-1)^s (\omega - T_k)^n$.

$$\int_{X(\leq s)} (-1)^s (\omega - T_k)^n = \int_{X(\leq s) \cap \Omega_{k,c_0}} (-1)^s (\omega - T_k)^n + \int_{X(\leq s) \cap \Omega_{k,c_0}^c} (-1)^s (\omega - T_k)^n.$$
(23)

Estimate of
$$\int_{X(\leq s)\cap\Omega_{k,c_0}} (-1)^s (\omega - T_k)^n$$

We get

$$\int_{X(\leq s)\cap\Omega_{k,c_0}} (-1)^s (\omega - T_k)^n \leq \int_{X(\leq s)\cap\Omega_{k,c_0}} \sum_{j=0}^s \binom{n}{j} (-1)^{s-j} w_{k,c_0}^{n-j} \wedge (T_k + v_{k,c_0})^j,$$
(24)

Set $S_k := \chi_{\Omega_{k,c_0}} w_{k,c_0}^{n-j} \wedge (T_k + v_{k,c_0})^j$ then

$$S_k \rightharpoonup \chi_{E_{c_0}(T_{min})^c} (\omega + c_0 u)^{n-j} \wedge (T_{min} + c_0 u)^j$$
 (25)

where $E_{c_0}(T_{min})^c$ is the complementary set of $E_{c_0}(T_{min})$. Now for $j = 0, \ldots, s - 1$, since $\operatorname{cod}(E_{c_0}(T_{min})) = s$ and $T_{min}^j = \langle T_{min}^j \rangle$ by Proposition (1.11), then

$$\int_{X} \chi_{E_{c_0}(T_{min})^c} (\omega + c_0 u)^{n-j} \wedge (T_{min} + c_0 u)^j =$$

$$\int_{X} \chi_{E_{c_0}(T_{min})^c} (\omega + c_0 u)^{n-j} \wedge \left(\sum_{h=0}^j {j \choose h} \langle T_{min}^h \rangle \wedge (c_0 u)^{j-h} \right) =$$

$$\int_{X} (\omega + c_0 u)^{n-j} \wedge \left(\sum_{h=0}^j {j \choose h} \langle T_{min}^h \rangle \wedge (c_0 u)^{j-h} \right). \quad (26)$$

For j = s, by Theorem (1.20) one has $T_{min}^s = \langle T_{min}^s \rangle + \sum_{t=1}^N \nu(T_{min}^s, Y_t)[Y_t]$, where Y_t are the irreducible components of codimension s in $E_{nn}(T_{min})$. Then we obtain the same identity for j = s as in (26). So first letting $k \to +\infty$ and then $c_0 \to 0$ and using the continuity of mobile-product the following holds

$$\limsup_{k \to +\infty} \int_{X(\leq s) \cap \Omega_{k,c}} \sum_{j=0}^{s} \binom{n}{j} (-1)^{s-j} w_{k,c}^{n-j} \wedge (T_k + v_{k,c})^j \leq \sum_{j=0}^{s} \binom{n}{j} (-1)^{s-j} L^{n-j} \langle F^j \rangle \quad (27)$$

For the second addendum in (23), we intersect with Ω_{k,c_1} thus

$$\int_{X(\leq s)\cap\Omega_{k,c_0}^c} (-1)^s (\omega - T_k)^n = \int_{X(\leq s)\cap\Omega_{k,c_0}^c\cap\Omega_{k,c_1}} (-1)^s (\omega - T_k)^n + \int_{X(\leq s)\cap\Omega_{k,c_0}^c\cap\Omega_{k,c_1}^c} (-1)^s (\omega - T_k)^n.$$
(28)

Then for the first addendum in (28), letting $k \to +\infty$ and $c_1 \to \nu'_1$, we get

$$\limsup_{k \to +\infty} \int_{X(\leq s) \cap \Omega_{k,c_0}^c \cap \Omega_{k,c_1}} (-1)^s (\omega - T_k)^n \le \sum_{t=1}^{M_1} (L + \nu_t^{'} \{u\})^{n-s} \nu_t [Y_t]$$

where the sum is over all $Y_t \subseteq E_{nn}(F)$ irreducible components of codimension s such that $\nu(T_{min}, Y_t) = \nu'_1$. Now it is clear how to study the second addendum in (28), one can intersect with Ω_{k,c_2} . Then in general we have the following situation:

$$\limsup_{k \to +\infty} \int_{X(\leq s) \cap \left(\bigcap_{i=0}^{j-1} \Omega_{k,c_{i}}^{c}\right) \cap \Omega_{k,c_{j}}} (-1)^{s} (\omega - T_{k})^{n} \leq \sum_{t=M_{j-1}}^{M_{j}} (L + \nu_{t}^{'} \{u\})^{n-s} \nu_{t}[Y_{t}]$$

$$\tag{29}$$

where for all j = 2, ..., h, $Y_t \subseteq E_{nn}(F)$ are the irreducible components of codimension s with $\nu(T_{min}, Y_t) = \nu'_j$ for $M_{j-1} \leq t \leq M_j$. While one has that

$$\int_{X(\leq s)\cap\left(\bigcap_{i=1}^{N}\Omega_{k,c_{i}}^{c}\right)} (-1)^{s} (\omega - T_{k})^{n} \leq \int_{X(\leq s)\cap\Omega_{k,\nu}} (-1)^{s} (\omega - T_{k})^{n} \xrightarrow{k \to +\infty} 0$$

$$(30)$$

for dimensional reasons.

Then putting together the inequalities of (29), we finally have

$$\limsup_{k \to +\infty} \int_{X(\leq s)} (-1)^s (\omega - T_k)^n \leq \sum_{j=0}^s \binom{n}{j} (-1)^{s-j} L^{n-j} \langle F^j \rangle + \sum_{t=1}^N \binom{n}{s} (L + \nu_t^{'} \{u\})^{n-s} \nu_t [Y_t].$$
(31)

And using now the standard holomorphic Morse inequalities the statement holds.

If F is not in the interior of \mathcal{M}^s and L is not ample, we consider $F + \varepsilon A$, $L + \varepsilon A$ and then pass to the limit by using the very definition of multicity (Definition 1.16).

Corollary 2.5 (Second formulation) Let X be a compact projective manifold of complex dimension n. Let L and F be two line bundles over X with L nef and $F \in \mathcal{M}^s$ with $\dim(E_{nn}(F)) = n - s$. Then we have the following Morse inequalities

$$\sum_{j=0}^{s} (-1)^{s-j} h^{j}(X, k(L-F)) \leq \frac{k^{n}}{n!} \sum_{j=0}^{s} {n \choose j} (-1)^{s-j} L^{n-j}(F^{j}) + {n \choose s} (L+b\{u\})^{n-s} (F^{s}-\langle F^{s} \rangle) + o(k^{n}) \quad (32)$$

where $b = \max_{j=1,\dots,N} \nu_{j}^{'}$.

Proof. It is sufficient to notice that $b \ge \nu'_j$ for all $j = 1, \dots, N$ and $F^s - \langle F^s \rangle = \sum_{j=1}^N \nu_j[Y_j]$, then one can applies Theorem (2.4).

Finally, by using remark 1.4 we can infer that Theorem (2.3) is a particular case of Theorem (2.4).

3 Transformation of Lelong numbers by Direct Images

3.1 The push-forward of Lelong numbers by a modification

We want to prove the following:

Proposition 3.1 Let X be a complex compact manifold with $\dim_{\mathbb{C}}(X) = 3$. Let $\tilde{\mu} : \tilde{X} \to X$ be a modification of X and Ω is a smooth, positive form on \tilde{X} of bidimension 1. Then $\nu(\tilde{\mu}_*(\tilde{\Omega}), Y) = 0 \ \forall Y$ irreducible curve on X, where ν is the generic Lelong number.

Since a modification can be decomposed as a finite sequence of blow-ups with smooth centers, we can write $\tilde{\mu}$ as follows:

$$\tilde{\mu}: \tilde{X} := X_s \to X_{s-1} \to \dots \to X_1 \to X_0 = X.$$
(33)

with $\forall i = 1, ..., s$ $Z_i, E_i \subset X_i$ are the smooth centers and the exceptional divisors of the *i*-th blow-up respectively.

Let $y \in Y$ be a generic point, without loose of generality one can suppose that y is the origin for a local chart with coordinates $z = (z_1, z_2, z_3)$. One has the following:

$$\nu(\tilde{\mu_*}(\tilde{\Omega}), 0) = \lim_{r \to 0} \frac{1}{r^2} \int_{B(0,r)} \tilde{\mu_*}(\tilde{\Omega})(z) \wedge \frac{i}{2\pi} \partial \bar{\partial} |z|^2 =$$
$$= \lim_{r \to 0} \frac{1}{r^2} \int_{\tilde{\mu}^{-1}(B(0,r))} \tilde{\Omega} \wedge \tilde{\mu^*}\left(\frac{i}{2\pi} \partial \bar{\partial} |z|^2\right)$$

Let's note that the integrand in the last equality is smooth so one wants to estimate the volume of $\tilde{\mu}^{-1}(B(0,r))$ and to compare with the parameter r^2 . Using the fact that $\tilde{\mu}$ can be decomposed as a finite sequence of blow-ups, $\tilde{\mu}^{-1}(B(0,r))$ can be expressed explicitly by a local expression of the composition of these blow-ups. Thus, one can associate to this local expression a *psh*-function φ such that the volume of $\tilde{\mu}^{-1}(B(0,r))$ can be studied in therms of asymptotic estimates for the volume of sublevel sets $\{\varphi < \log r\}$. Explicitly, let $B(0,r) = \{|z_1|^2 + |z_2|^2 + |z_3|^2 < r\}$ be the local expression on X with coordinates (z_1, z_2, z_3) , so $\tilde{\mu}^{-1}(B(0,r)) = \{|h_1|^2 + |h_2|^2 + |h_3|^2 < r\}$ where $h_i := h_i(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3)$ are holomorphic functions on \tilde{X} and $(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3)$ are local coordinates on \tilde{X} .

Then one defines $\varphi = \frac{1}{2} \log \sum_{i=1}^{3} |h_i|^2$, so that $\tilde{\mu}^{-1}(B(0,r)) = \{\varphi < \log r\}$. Let us give the following

Definition 3.2 ([DK01]) Let $K \subset X$ be a compact set, $U \Subset X$ a relatively compact neighborhood of K and let θ_U be the Lebesgue measure on U associated with some choice of hermitian metric ω on X. Then the **log-canonical threshold** of φ is defined as:

$$c_K(\varphi) := \sup \{ c \ge 0 : r^{-2c} \theta_U(\{\varphi < \log r\}) \text{ is bounded as } r \to 0 \text{ for } U \supset K \}.$$

Now the crucial fact is that each h_i can be expressed as a sum of a holomorphic monomial in the \tilde{z}_i and another holomorphic function, so that looking at $\tilde{\mu}(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3) = (h_1(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3), h_2(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3), h_3(\tilde{z}_1, \tilde{z}_2, \tilde{z}_3))$ as an ideal "almost" monomial, one can calculate $c_K(\varphi)$ using Howald's theorem [How01].

Remark 3.3 One may assume that for all *i* the centers Z_i are connected by considering a longer sequence of blow-ups if necessary.

Remark 3.4 If Y is not entirely contained in Z_0 then the statement is trivial. In fact let's suppose $Y \cap Z_0 = \{a_1, \ldots, a_k\}$ is a finite set of points or possibly the empty set, let $\mu_1 \circ \cdots \circ \mu_s : X_s \to X_{s-1} \to \cdots \to X_1 \to X_0$ be a finite sequence of blow-ups, let $\tilde{\Omega}$ be a smooth positive form of bidimension 1 on X_s , and let E_1 be the exceptional divisor of μ_1 then $\mu_1(X_1 \setminus E_1) = X_0 \setminus \mu_1(E_1) = X_0 \setminus Z_0$ and $\mu_1|_{X_1 \setminus E_1} : X_1 \setminus E_1 \to X_0 \setminus Z_0$ is an isomorphism. Thus $\nu(\tilde{\mu}_*(\tilde{\Omega}), Y) = \nu((\mu_2 \circ \cdots \circ \mu_s)_* \tilde{\Omega}, \mu_1^{-1}(Y))$, and now if $\mu_1^{-1}(Y)$ is not entirely contained in Z_1 , the center of μ_2 , one can omit μ_2 in the calculation of ν and so on. So we may assume $Y = Z_0$.

Let $\tilde{\mu} = \mu_1 \circ \cdots \circ \mu_s$ be a finite sequence of blow-ups with smooth connected centers Z_i for $= 1, \ldots, s - 1$ (Z_0 is given by (3.4)). Let's define $Exc(\mu_1 \circ \cdots \circ \mu_i) := \bigcup_{j=1}^i (\mu_j \circ \cdots \circ \mu_i)^{-1} (Z_{j-1})$ then the centers can just satisfy one of these conditions:

a) Z_i is a point and $Z_i \in Exc(\mu_1 \circ \cdots \circ \mu_i)$,


b) Z_i is a point and $Z_i \notin Exc(\mu_1 \circ \cdots \circ \mu_i)$,

c) Z_i is a curve such that $Z_i \cap Exc(\mu_1 \circ \cdots \circ \mu_i) = \emptyset$,

d) Z_i is a curve such that $Z_i \cap Exc(\mu_1 \circ \cdots \circ \mu_i)$ is a finite set of points,

e) Z_i is a curve such that $Z_i \subset Exc(\mu_1 \circ \cdots \circ \mu_i)$ entirely.

Then one can associate to this sequence a rooted tree \mathcal{T} , where the root is the vertex X_0 and the edges $\mu_i^a, \mu_i^b, \mu_i^c, \mu_i^d, \mu_i^e$ denote the *i*-th blow-up (for $i \ge 2$) where the centers Z_i satisfy respectively \mathbf{a}), \mathbf{b}), \mathbf{c}), \mathbf{d}), \mathbf{e}).



Since we are interested in calculating $\nu(\tilde{\mu}_* \tilde{\Omega}, Z_0)$, where $\tilde{\mu} = \mu_1 \circ \cdots \circ \mu_s$ is a finite sequence of blow-ups with smooth centers, we want to understand which path of the tree \mathcal{T} represents the more general sequence according to the following

Definition 3.5 The sequence of blow-ups $\mu_1 \circ \cdots \circ \mu_s$ is called minimal if there does not exist another sequence $\mu'_1 \circ \cdots \circ \mu'_s$ such that $\nu((\mu_1 \circ \cdots \circ \mu_s)_*(\tilde{\Omega}), Z_0) = \nu((\mu'_1 \circ \cdots \circ \mu'_s)_*(\tilde{\Omega}), Z_0)$ where there exists an *i* for which μ'_i is a local biholomorphism.

Thus we want to understand which path of the tree doesn't have this property because it reflects the general sequence of s blow-ups. The following lemma shows that if Z_i satisfies one of conditions \mathbf{a}), \mathbf{b} , \mathbf{c}), \mathbf{d}) then the corresponding path is not minimal.

Lemma 3.6 Let \mathcal{T} be the rooted tree associated to a finite sequence of blow-ups. The following path

 $\tilde{\mu}^{J_1,J_2,\ldots,J_s} := \mu_1^{J_1} \circ \mu_2^{J_2} \circ \cdots \circ \mu_s^{J_s} \text{ where } J_1,\ldots,J_s \in \{a,b,c,d\}$

is not minimal.

Proof. Second blow-up. Let $\tilde{\mu} = \mu_1^J \circ \mu_2 \circ \mu_3 \circ \cdots \circ \mu_s$ be a sequence of s blow-ups such that $J \in \{a, b, c, d\}$.

<u>Case J=a.</u> Let $p_0 \in Z_0$ such that $\mu_1^{-1}(p_0) \ni Z_1$ then for all $p \neq p_0$ with $p \in Z_0$ there exists r > 0 small enough with $Z_1 \nsubseteq \mu_1^{-1}(B(p,r))$. Then near p

$$\mu_2: \mu_2^{-1}(\mu_1^{-1}(B(p,r))) \xrightarrow{\simeq} \mu_1^{-1}(B(p,r))$$
(34)

is an isomorphism. Now if we consider $\tilde{\mu}' = \mu_1 \circ \mu'_2 \circ \mu_3 \circ \cdots \circ \mu_s$ with μ'_2 any isomorphism, as above we have that $\nu(\tilde{\mu}_*\tilde{\Omega}, Z_0) = \nu(\tilde{\mu}'_*\tilde{\Omega}, Z_0)$.

<u>Case J=b</u>. The argument above is still valid.

<u>Case J=c</u> For all points $p \in Z_0$ one can find an r > 0 sufficiently small such that (6) still holds, then one can repeat the argument of case $J = \mathbf{a}$.

<u>Case J=d</u>. Let $\{a_1, \ldots, a_k\} = Z_1 \cap E_1 = Exc(\mu_1)$ be a finite set of points. For all $p \in Z_0$ such that $p \neq \mu_1(a_i)$ for all $i = 1, \ldots, k$ one can find again an r > 0 small enough such that $a_i \notin \mu_1^{-1}(B(p, r)) \quad \forall i = 1, \ldots, k$. Repeating the argument as before one can find another sequence of s - 1 blow-ups $\tilde{\mu}'$ such that $\nu(\tilde{\mu}_*\tilde{\Omega}, Z_0) = \nu(\tilde{\mu}'_*\tilde{\Omega}, Z_0)$.

Third blow-up. Now let's suppose $\tilde{\mu} = \mu_1^e \mu_2^J \circ \mu_3 \circ \cdots \circ \mu_s$ with $J \in \{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}\}$ and we may assume that $\mu_1 : Z_1 \to Z_0$ is surjective. <u>Case $J = \mathbf{a}, \mathbf{b}$ </u>. Let p be a point of Z_0 such that $(\mu_1 \circ \mu_2)(Z_2) = p$ then for all $p' \neq p$ and $p' \in Z_0$ we can find a small enough r > 0 such that

$$\mu_3: \mu_3^{-1}\left(\mu_2^{-1} \circ \mu_1^{-1}(B(p', r))\right) \xrightarrow{\simeq} \mu_2^{-1} \circ \mu_1^{-1}(B(p', r)).$$
(35)

Then for any isomorphism μ_3 as above one can change the sequence $\tilde{\mu}$ with $\tilde{\mu}' = \mu_1^e \circ \mu_2^J \circ \mu_3' \circ \cdots \circ \mu_s$ and we have $\nu(\tilde{\mu}_*\tilde{\Omega}, Z_0) = \nu(\tilde{\mu}'_*\tilde{\Omega}, Z_0)$. <u>Case $J=\mathbf{c}$ </u>. The argument above is still valid.

<u>Case J=d</u>. Let $\{a_1, \dots, a_k\} = Z_2 \cap Exc(\mu_1^e \circ \mu_2)$ be a finite set of points in X_2^d , then there exist $p_1, \dots, p_k \in Z_0$ such that $(\mu_1^e \circ \mu_2^d)(a_i) = p_i$. Then for all $p' \in Z_0$ with $p' \neq p_i$ $1 \leq i \leq k$ we can find r > 0 sufficiently small so that (35) is again true. Therefore one finds a sequence of s - 1 blow-ups as in the cases above.

Now it's clear how to study the general case.

i-th blow-up. Let $\tilde{\mu} = \mu_1^e \circ \cdots \circ \mu_{i-1}^e \circ \mu_i^J \circ \mu_{i+1} \circ \cdots \circ \mu_s$ be a finite sequence of blow-ups with $J \in \{\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}\}$.

<u>Case J=a</u>. Let Z_i be the center of μ_{i+1} with $Z_i \subset Exc(\mu_1^e \circ \cdots \circ \mu_{i-i}^e \circ \mu_i^a)$. and let $(\mu_1^e \circ \cdots \circ \mu_{i-i}^e \circ \mu_i^a)(Z_i) = p$. Thus for all $p' \neq p$ in Z_0 we can find a small positive r such that the following hold

- 1) $(\mu_1^e \circ \cdots \circ \mu_i^a)^{-1}(B(p', r)) \cap Z_i = \emptyset$
- 2) $\mu_{i+1} : \mu_{i+1}^{-1} \left((\mu_1^e \circ \dots \circ \mu_{i-i}^e \circ \mu_i^a)^{-1} B(p', r) \right) \xrightarrow{\simeq} (\mu_1^e \circ \dots \circ \mu_{i-i}^e \circ \mu_i^a)^{-1} B(p', r).$

So $\tilde{\mu}' = \mu_1^e \circ \cdots \cap \mu_{i-1}^e \circ \mu_i^a \circ \mu_{i+1}' \circ \cdots \cap \mu_s$, where μ_{i+1}' is an isomorphism, satisfies $\nu(\tilde{\mu}_*\tilde{\Omega}, Z_0) = \nu(\tilde{\mu}_*'\tilde{\Omega}, Z_0)$.

<u>Case $J=\mathbf{b}$ </u>. As in the case $J=\mathbf{a}$ 1) and 2) are still valid, but now they are valid for all $p' \in Z_0$.

<u>Case J=c</u>. Again as in the case J=a, b we can find an r > 0 sufficiently small such that for all $p' \in Z_0$ 1) and 2) are still valid and then the statement holds. <u>Case J=d</u>. $Z_i \cap Exc(\mu_1^e \circ \cdots \circ \mu_{i-i}^e \circ \mu_i^a) = \{a_1, \ldots, a_k\}$, where a_i are points of Z_i . Now there exist $p_1, \ldots, p_k \in Z_0$ such that $(\mu_1^e \circ \cdots \circ \mu_{i-i}^e \circ \mu_i^a)(a_i)p_i$. So we can find r > 0 small so that for $p \neq p_i$ $1 \leq i \leq k$ such that

$$a_i \notin (\mu_1^e \circ \cdots \circ \mu_{i-i}^e \circ \mu_i^a)^{-1} B(p,r),$$

so we can repeat the same reasoning as in the previous cases. According to Lemma (3.6) it remains to analyze just the path of the tree \mathcal{T} corresponding to a sequence of blow-ups $\tilde{\mu} = \mu_1^e \circ \cdots \circ \mu_{s-1}^e \circ \mu_s^e$ where for all $i = 1, \ldots, s - 1$ $Z_i \subset Exc(\mu_1^e \circ \cdots \circ \mu_i^e)$ and $\mu_1^e : Z_1 \to Z_0$ is surjective. Now let's distinguish two possible cases:

1)
$$Z_i \subseteq Exc(\mu_1^e \circ \cdots \circ \mu_i^e) := Exc(\mu_1^e \circ \cdots \circ \mu_i^e) \setminus E_i;$$
 2) $Z_i \subseteq E_i$ entirely. (36)

Notations 3.7 We write

$$\tilde{\mu}^e = \tilde{\mu}_1^{e_{i_1}} \circ \tilde{\mu}_2^{e_{i_2}} \cdots \circ \tilde{\mu}_s^{e_{i_s}} \tag{37}$$

with $i_1, \ldots, i_s \in \{1, 2\}$, where 1,2 correspond to the two conditions of (36) for the center of the following blow-up. (For example $\mu_j^{e_1}$ means that the center of the j + 1-th blow-up satisfies condition 1 of (36).)

Let's note that for the first blow-up it is not necessary to indicate which condition of (36) Z_1 satisfies thanks to Lemma 3.6 and to the following remark.

Remark 3.8 If $Z_i \subseteq E_i$ for some $i = 1, \ldots, s - 1$. Since

 $\mu_i|_{E_i}: E_i \to Z_{i-1}$

is a holomorphic fiber bundle isomorphic to the projectivized normal bundle $\mathbb{P}(N_{Z_{i-1}}) \to Z_{i-1}$. We can suppose that $\mu_i|_{Z_i} : Z_i \to Z_{i-1}$ is surjective.

Now let's fix some notations. Since everything is local, without loose of generality, one can suppose that $Z_0 = \{x_0 = y_0 = 0\}$ where (x_0, y_0, z_0) are local coordinates on $X_0 = X$.

Remark 3.9 Let Y be a n-dimensional smooth variety and B a closed smooth subvariety with $\operatorname{codim}_Y(B) = t$ and $\sigma : \tilde{Y} \to Y$ the blow-up of Y with center B. For every point $b_0 \in B$ there exists local coordinates (u_1, \ldots, u_n) on Y centered at b_0 such that $B = \{u_1 = \cdots = u_t = 0\}$ and local coordinates (w_1, \ldots, w_n) on \tilde{Y} such that the map σ is given by:

$$\sigma(w_1, \dots, w_n) = (w_1 w_j, \dots, w_{j-1} w_j; w_j; w_{j+1} w_j, \dots, w_t w_j, w_{t+1}, \dots, w_n)$$
(38)
$$\forall j = 1, \dots, t.$$

Lemma 3.10 If the first condition of (36) is satisfied for some *i* with $1 \le i \le s$ the $\tilde{\mu}$ is not minimal.

Proof. Let's consider the *i*-th blow-up $\mu_i = X_i \to X_{i-1}$ by using (3.9) we see that there exists local coordinates (x_i, y_i, z_i) on X_i such that

$$\mu_i(x_i, y_i, z_i) = \begin{cases} (x_i, y_i x_i, z_i) \\ (x_i y_i, y_i, z_i) \end{cases}$$

where we used the same coordinates on two different charts to keep notations simple. By Lemma (3.6) one has that $Z_i \subseteq Exc(\mu_1^e \circ \cdots \circ \mu_i^e)$. Now let us distinguish the following two cases:

- 1. $Z_i \subseteq E_i$ entirely,
- 2. $Z_i \subseteq \widetilde{Exc}(\mu_1^e \circ \cdots \circ \mu_i^e)$ entirely.

<u>Case 1.</u> Repeating the argument of (3.8) we get that

$$Z_i = \{x_i = 0 = f_i(y_i, z_i)\}$$
 or $Z_i = \{y_i = 0 = g_i(x_i, z_i)\}$

depending on which chart we are considering where f_i, g_i are holomorphic functions. By surjectivity of the map $\mu_i : Z_i \to Z_{i-1}$ we deduce that on the generic point of Z_i

$$\frac{\partial f_i}{\partial y_i} \neq 0 \quad \text{or} \quad \frac{\partial g_i}{\partial x_i} \neq 0$$

then by implicit function theorem there exist holomorphic functions ${\cal F}_i, {\cal G}_i$ such that

$$Z_i = \{x_i = 0, y_i = Fz_i\}$$
 or $Z_i = \{y_i = 0, x_i = G_i(z_i)\}.$

Now considering the following holomorphic changes of coordinates on the two different charts of μ_i

$$\begin{cases} U_i = x_i \\ V_i = y_i - F_i(z_i) \\ W_i = z_i \end{cases} \quad \text{or} \quad \begin{cases} U_i = x_i - G_i(z_i) \\ V_i = y_i \\ W_i = z_i \end{cases}$$
(39)

where we have still used the same coordinates on the two different charts. So the equation of Z_i by (39) becomes $Z_i = \{U_i = V_i = 0\}$. Now we can use (3.9) blowing-up along Z_i and repeating the same operations as above.

<u>Case 2.</u> Let $i \in \{1, \ldots, s-1\}$ be the first index such that $Z_i \nsubseteq E_i$ entirely, i.e. $Z_i \cap E_i$ is a finite set of points. Hence for all $j < i \ Z_j \subseteq E_j$ entirely. Now to keep notations simple we denote by E'_j for j < i the pre-image of $E_j \setminus Z_j$ in X_i by $(\mu_j \circ \cdots \circ \mu_i)$; then $Z_i \subseteq \bigcup_{j=1}^{i-1} E'_j$. Since Z_i is connected then we can suppose

that $Z_i \subseteq E_{j_0}$ entirely for some $1 \le j_0 \le i-1$. The expression in local coordinates of μ_i is given by

$$\mu_i(x_i, y_i, z_i) = \begin{cases} (x_i, y_i x_i, z_i) \\ (x_i y_i, y_i, z_i) \end{cases}$$

where (x_i, y_i, z_i) depends on $(U_{i-1}, V_{i-1}, W_{i-1})$ on X_{i-1} . But now $Z_i \not\subseteq E_i$ entirely means that $(\mu_{j_0+1} \circ \cdots \circ \mu_i)|_{Z_i \setminus \{p_1, \dots, p_k\}}$ is an isomorphism and furthermore $(\mu_{j_0+1} \circ \cdots \circ \mu_i)(Z_i \setminus \{p_1, \dots, p_k\}) \subseteq E_{j_0}$ entirely thus in local coordinates $(x_{j_0}, y_{j_0}, z_{j_0})$ on X_{j_0} we have that

$$\mu_{j_0}(x_{j_0}, y_{j_0}, z_{j_0}) = \begin{cases} (x_{j_0}, y_{j_0} x_{j_0}, z_{j_0}) \\ (x_{j_0} y_{j_0}, y_{j_0}, z_{j_0}) \end{cases}$$

and $(\mu_{j_0+1} \circ \cdots \circ \mu_i)(Z_i)$ is equal to

$$\{x_{j_0} = 0 = f_i(y_{j_0}, z_{j_0})\} \text{ or } \{y_{j_0} = 0 = g_i(x_{j_0}, z_{j_0})\}.$$
(40)

So performing the (i + 1)-blow-up along Z_i on X_i or along $(\mu_{j_0+1} \circ \cdots \circ \mu_i)(Z_i)$ on X_{j_0} it is essentially equivalent. By remark (3.8) it is not restrictive to suppose that

$$\mu_{j_0}|_{(\mu_{j_0+1}\circ\cdots\circ\mu_i)(Z_i)}:(\mu_{j_0+1}\circ\cdots\circ\mu_i)(Z_i)\twoheadrightarrow Z_{j_0-1}$$

and this means that in the local expressions (40) one can suppose that

$$\frac{\partial f_i}{\partial y_{j_0}} \neq 0 \text{ or } \frac{\partial g_i}{\partial x_{j_0}} \neq 0$$

depending which local chart we are considering. So we can define new coordinates on X_{j_0} using (39). Namely:

$$\begin{cases} U_{i} = x_{j_{0}} \\ V_{i} = y_{j_{0}} - F_{i}(z_{j_{0}}) \\ W_{i} = z_{j_{0}} \end{cases} \text{ or } \begin{cases} U_{i} = x_{j_{0}} - G_{i}(z_{j_{0}}) \\ V_{i} = y_{j_{0}} \\ W_{i} = z_{j_{0}} \end{cases}$$
(41)

for which $(\mu_{j_0+1} \circ \cdots \circ \mu_i)(Z_i) = \{U_i = V_i = 0\}$ and we can now perform the (i+1)-blow-up. The relation with the coordinates system $(U_{j_0}, V_{j_0}, W_{j_0})$ on X_{j_0} is straightforward by using:

$$\left\{ \begin{array}{ll} U_{j_0} = x_{j_0} \\ V_{j_0} = y_{j_0} - F_{j_0}(z_{j_0}) \\ W_{j_0} = z_{j_0} \end{array} \right. \quad \text{or} \quad \left\{ \begin{array}{ll} U_{j_0} = x_{j_0} - G_{j_0}(z_{j_0}) \\ V_{j_0} = y_{j_0} \\ W_{j_0} = z_{j_0}. \end{array} \right. \right.$$

Then the expression in local coordinates of the composition $(\mu_{j_0} \circ \cdots \circ \mu_{i+1})$ is the following:

$$(\mu_{j_0} \circ \dots \circ \mu_{i+1}) = \begin{cases} (x_{i+1}, x_{i+1}(x_{i+1}y_{i+1}F_i(z_{i+1})), z_{i+1}) \\ (x_{i+1}y_{i+1}, x_{i+1}y_{i+1}(y_{i+1} + F_i(z_{i+1})), z_{i+1}) \\ ((x_{i+1} + G_i(z_{i+1}))x_{i+1}y_{i+1}, x_{i+1}y_{i+1}, z_{i+1}) \\ ((x_{i+1}y_{i+1} + G_i(z_{i+1}))y_{i+1}, y_{i+1}, z_{i+1}) \end{cases}$$
(42)

Let's observe that in the <u>Case 2</u> we can disreguard the intersection of Z_i with E_i since it has dimension zero and it has no effect on the generic Lelong number.

3.2 Estimate for $c_K(\varphi)$

Let's now compute the log-canonical threshold of the ideals rising from $\tilde{\mu} = \mu_1 \circ \cdots \circ \mu_s$ a composition of s blow-ups. By the previous section without

loss of generality one can suppose that each center Z_i is entirely contained in the exceptional divisor E_i , since we have already notice (<u>Case 2</u>) that the expression of the composition of blow-ups is equivalent to a composition of blow-ups in which each center Z_i is entirely contained in E_i .

Proposition 3.11 Let $\tilde{\mu} = \mu_1 \circ \cdots \circ \mu_s : \tilde{X} \to X$ be a finite sequence of blow-ups with smooth centers Z_i such that $\dim_{\mathbb{C}}(Z_i) = 1$ and Z_i is entirely contained in the exceptional divisor of μ_i and it's surjective $\forall i = 1, \ldots, s$. Then considering the ideal generated by all possible 2^k expressions of $\tilde{\mu}$ in local coordinates, one can reduce them to ideals of the forms

$$\mathcal{I} = (x_s^h y_s^k, z_s)$$
 with $h, k \ge 0$ with $max(h, k) > 0$.

Proof. Finite induction on s.

Case s=1. $\tilde{\mu} = \mu_1 : X_1 \to X_0$, there exist local coordinates (x_0, y_0, z_0) on X_0 such that $Z_0 = \{x_0 = y_0 = 0\}$, by (3.9) the expressions of μ_1 in local coordinates are:

$$\mu_1(x_1, y_1, z_1) = (x_1, x_1y_1, z_1)$$
 and $\mu_1(x_1, y_1, z_1) = (x_1y_1, y_1, z_1)$

so one has that

$$(x_1, x_1y_1, z_1) = (x_1, z_1)$$
 and $(x_1y_1, y_1, z_1) = (y_1, z_1)$.

Case s=2. $\tilde{\mu} = \mu_1 \circ \mu_2 : X_2 \xrightarrow{\mu_2} X_1 \xrightarrow{\mu_1} X_0$. Let $Z_0 = \{x_0 = y_0 = 0\}$ be the center of μ_1 then there exists local coordinates on X_1 such that

$$\mu_1(x_1, y_1, z_1) = (x_1, x_1y_1, z_1)$$
 and $\mu_1(x_1, y_1, z_1) = (x_1y_1, y_1, z_1)$

Let's consider first the expression $\mu_1(x_1, y_1, z_1) = (x_1, x_1y_1, z_1)$; by (39) let $Z_1 = \{x_1 = 0 = f_1(y_1, z_1)\} = \{x_1 = 0, y_1 = F_1(z_1)\}$ be the center of μ_2 . So that $Z_1 = \{U_1 = V_1 = 0\}$. Then there exist local coordinates on X_2 such that:

$$\mu_2(x_2, y_2, z_2) = (x_2, x_2y_2, z_2)$$
 and $\mu_2(x_2, y_2, z_2) = (x_2y_2, y_2, z_2),$

repeating the argument used in the case of $\mu_1(x_1, y_1, z_1) = (x_1y_1, y_1, z_1)$, the following holds:

1. $(\mu_1 \circ \mu_2)(x_2, y_2, z_2) = (x_2, x_2(x_2y_2 + F_1(z_2)), z_2)$ 2. $(\mu_1 \circ \mu_2)(x_2, y_2, z_2) = (x_2y_2, x_2y_2(y_2 + F_1(z_2)), z_2)$ 3. $(\mu_1 \circ \mu_2)(x_2, y_2, z_2) = ((x_2 + G_1(z_2))x_2y_2, x_2y_2, z_2)$ 4. $(\mu_1 \circ \mu_2)(x_2, y_2, z_2) = ((x_2y_2 + G_1(z_2))y_2, y_2, z_2).$

Considering the simmetry of these expressions, it's enough to analyze 1) and 2), thus these local expressions can be reduced to the ideals respectively:

$$(x_2, z_2)$$
 and (x_2y_2, z_2) .

Case s=3. One would have to consider the 2^3 possible expressions in local coordinates but by simmetry it's enough to study four of them, in particular:

$$\begin{split} \mathfrak{a} &= (x_3, x_3((x_3y_3 + F_2(z_3))x_3 + F_1(z_3)), z_3) \\ \mathfrak{b} &= (x_3y_3, x_3y_3((y_3 + F_2(z_3))x_3y_3 + F_1(z_3)), z_3) \\ \mathfrak{c} &= (x_3y_3(x_3 + G_2(z_3)), x_3y_3(x_3 + G_2(z_3))(x_3y_3 + F_1(z_3)), z_3) \\ \mathfrak{d} &= (y_3(y_3x_3 + G_2(z_3)), y_3(y_3x_3 + G_2(z_3))(y_3 + F_1(z_3)), z_3). \end{split}$$

The ideals \mathfrak{a} and \mathfrak{b} can be reduced in the desired form, i.e.:

 $a = (x_3, z_3)$ and $b = (x_3y_3, z_3)$.

While for the last two ideals one obtains:

 $\mathbf{c} = (x_3y_3(x_3 + G_2(z_3)), z_3) = (x_3^2y_3 + x_3y_3G_2(z_3), z_3)$ $\mathbf{d} = (y_3(y_3x_3 + G_2(z_3)), z_3) = (y_3^2x_3 + y_3G_2(z_3), z_3).$

Let's study \mathfrak{c} , in particular notice that $H_3(x_3, y_3, z_3) := x_3 y_3 \tilde{G}_2(z_3) z_3^i$ is an holomorphic function which is an element of (z_3) where $\tilde{G}_2(0) \neq 0$ (here we are using the fact that $\mu_2 : Z_2 \twoheadrightarrow Z_1$ and $G_2(0) = 0$). Thus

$$x_3^2 y_3 = x_3^2 y_3 + x_3 y_3 G_2(z_3) - H_3(x_3, y_3, z_3),$$

so $\mathbf{c} = (x_3^2 y_3, z_3).$

Repeating the same argument for the case of \mathfrak{d} one obtains $\mathfrak{d} = (y_3^2 x_3, z_3)$. Now let's suppose the statement true for all $j = 1, \ldots, s - 1$. Let $\tilde{\mu} = \mu_1 \circ \cdots \circ \mu_s$ be a composition of s blow-ups, then μ_s has two possible expressions in local coordinates (x_s, y_s, z_s) which depend on the coordinates on X_{s-1} , but by induction the local expression of $\mu_1 \circ \cdots \circ \mu_{s-1}$ is of the form $(x_{s-1}^i y_{s-1}^j, z_{s-1})$ so composing with local expressions of μ_s one obtains the statement.

Now let's recall Howald's theorem [How01] for multiplier ideals of monomial ideals.

Definition 3.12 Let $\mathfrak{a} \subset \mathbb{C}[x_1, \ldots, x_n]$ be a monomial ideal. We will regard \mathfrak{a} as a subset of the lattice $L = \mathbb{N}^n$ of monomials. The **Newton Polygon** P of \mathfrak{a} is the convex hull of this subset of L, considered as a subset of $L \otimes \mathbb{R} = \mathbb{R}^n$. It is an unbounded region.

Notations 3.13 We write **1** for the vector (1, 1, ..., 1), which is identified with the monomial $x_1x_2 \cdots x_n$. We use Greek letters $(\lambda \in L)$ for elements of Lor $L \otimes \mathbb{R}$, and exponent notation x^{λ} for the associated monomial. For any subset P of $L \otimes \mathbb{R}$, we define rP "pointwise"

$$rP = \{r\lambda : \lambda \in P\}.$$

We write Int(P) for the topoogical interior of P.

Theorem 3.14 ([How01]) (Howald's theorem.) Let $\mathfrak{a} \subset \mathcal{O}_{\mathcal{A}^n}$ be a monomial ideal. Let P be its Newton polygon. Then $\mathcal{J}(r \cdot \mathfrak{a})$ is a monomial ideal, and contains exactly the following monomials:

$$\mathcal{J}(r \cdot \mathfrak{a}) = \{ x^{\lambda} : \lambda + \mathbf{1} \in Int(P) \cap L \}.$$

where $\mathcal{J}(r \cdot \mathfrak{a})$ is the multiplier ideal associated to r and \mathfrak{a} (Cfr. [Laz04]).

Let \mathfrak{a} be a monomial ideal and let P be its Newton polygon. The log canonical threshold $c(\mathfrak{a})$ of \mathfrak{a} is defined to be

$$c(\mathfrak{a}) = \sup\{r : \mathcal{J}(r \cdot \mathfrak{a}) \neq \mathcal{O}_X\}.$$
(43)

Howald's theorem shows that this must be equal to $\sup\{r : \mathbf{1} \notin rP\}$. Thus the log canonical threshold is the reciprocal of the (unique) number m such that the boundary of P intersects the diagonal in \mathbb{R}^n at the point $m\mathbf{1}$. In other words, in order to calculate the threshold, we need only find where Pintersects the diagonal.

Proposition 3.15 ([DK01]) Let X, Y be complex manifolds of respective dimension n, m, let $\mathcal{I} \subset \mathcal{O}_X$, $\mathcal{J} \subset \mathcal{O}_Y$ be coherent ideals, and let $K \subset X$, $L \subset Y$ be compact set. Put $\mathcal{I} \oplus \mathcal{J} := pr_1^*\mathcal{I} + pr_2^*\mathcal{J} \subset \mathcal{O}_{X \times Y}$. Then

$$c_{K \times L}(\mathcal{I} \oplus \mathcal{J}) = c_K(\mathcal{I}) + c_L(\mathcal{J}).$$

Proposition 3.16 Let $\mathcal{I} = (x^h y^k, z)$ be an ideal as (3.11) then $c(\mathcal{I}) > 1$.

Proof. Using Howald's theorem and (3.15)

$$c_K(\mathcal{I}) = c_K(x^h y^k) + c_K(z) = \frac{h+k}{hk} + 1 > 1.$$
(44)

3.3 Proof of Proposition (3.1)

Now we can give the proof for the main proposition

Proof of (3.1). By (3.11) and (3.15) one can define $\varphi := \log(|x_s^h y_s^k| + |z_s|)$ thus $c_K(\varphi) = c_K((x_s^h y_s^k, z_s)) > 1$. Then by (3.16) and by the definition of Lelong number the following holds:

$$\nu(\tilde{\mu_*}(\tilde{\Omega}), 0) = \lim_{r \to 0} \frac{1}{r^2} \int_{\tilde{\mu}^{-1}(B(0, r))} \tilde{\Omega} \wedge \tilde{\mu^*}\left(\frac{i}{2\pi} \partial \bar{\partial} |z|^2\right) \le \lim_{r \to 0} C_1 r^{2(c_K(\varphi) - 1)} = 0,$$

where C_1 is a positive constant.

3.4 Relation between mobile intersection and positive product in dimension 3

By using Proposition (3.1) we obtain a similar decomposition as in Theorem (1.20), using the definition of mobile product via modifications. In fact we have the following

Proposition 3.17 Let X be a compact Kähler manifold of complex dimension 3 and let $\alpha_1, \alpha_2 \in \mathcal{M}^2$ then

$$\alpha_1 \alpha_2 - \langle \alpha_1 \alpha_2 \rangle = \left\{ \sum_{t=1}^{+\infty} \nu_{\infty,t} [Y_{\infty,t}] \right\}$$

where $\nu_{\infty,t} \ge 0$ and Y_t are the irreducible components of codimension 2 of $\bigcup_{i=1}^{2} E_{nn}(\alpha_i)$.

Proof. For i = 1, 2 Let $T_{k,i} \in \alpha_i$ be a sequence of currents with analytic singularities as in the Theorem (1.1) such that $T_{k,i} \geq -\varepsilon_{k,i}\omega$ and $T_{k,i} \rightharpoonup T_{min,i}$ where $T_{min,i}$ is a positive current with analytic singularities in α_i . Now let $\mu_k : X_k \rightarrow X$ be a common modification for $T_{k,i}$ such that $\mu_k^* T_{k,i} = [E_{k,i}] + \beta_{k,i}$, where $\beta_{k,i}$ are smooth. Now we have that

$$T_{k,1} \wedge T_{k,2} - (\mu_k)_* (\beta_{k,1} \wedge \beta_{k,2}) = 0 \text{ on } X \setminus (\bigcup_{i=1}^2 V(\mathcal{I}(k\varphi_{min,i})))$$

thus by using the support theorem for currents and the inclusion $\cup_{i=1}^2 V(\mathcal{I}(k\varphi_{min,i})) \subseteq \cup_{i=1}^2 E_{nn}(\alpha_i)$ we obtain

$$T_{k,1} \wedge T_{k,2} - (\mu_k)_* (\beta_{k,1} \wedge \beta_{k,2}) = \sum_{t=1}^{+\infty} \nu_{k,t} [Y_{k,t}]$$
(45)

where $Y_{k,t}$ are the irreducible components of $E_{nn}(\alpha_1) \cup E_{nn}(\alpha_2)$ and $\nu_{k,t} \in \mathbb{R}$. Now let $\nu_{k,t}^+$ and $\nu_{k,t}^-$ be the positive and the negative coefficients in the series of (45), thus we have

$$T_{k,1} \wedge T_{k,2} + \sum_{t} (-\nu_{k,t}^{-})[Y_{k,t}] = (\mu_k)_* (\beta_{k,1} \wedge \beta_{k,2}) + \sum_{t} \nu_{k,t}^{+}[Y_{k,t}].$$
(46)

If there exists t_0 such that $\nu_{k,t_0}^- < 0$ i.e. $-\nu_{k,t_0}^- > 0$ then by calculating the generic Lelong number along Y_{k,t_0} , using equality (46), we find

$$\nu(T_{k,1} \wedge T_{k,2}, Y_{k,t_0}) - \nu_{k,t_0}^- = \nu((\mu_k)_*(\beta_{k,1} \wedge \beta_{k,2}), Y_{k,t_0}) + 0$$

and thanks to Proposition (3.1) $\nu((\mu_k)_*(\beta_{k,1} \land \beta_{k,2}), Y_{k,t_0}) = 0$ then we have $\nu(T_{k,1} \land T_{k,2}, Y_{k,t_0}) = \nu_{k,t_0}^- < 0$, a contradiction. Hence we have that $\nu_{k,t} = \nu(T_{k,1} \land T_{k,2}, Y_{k,t})$ for all t. By using Lemma (1.9) the sequence $\{\nu_{k,t}\}$ is bounded, and $T_{k,1} \land T_{k,2}$ are bounded in mass since they are in the same cohomology class for all k, while $\{(\mu_k)_*(\beta_{k,1} \land \beta_{k,2})\}$ are bounded in mass thanks to Theorem (1.6). Hence we can extract a common subsequence such that passing in cohomology we obtain the statement.

And as a consequence we have

Corollary 3.18 Assume dim(X) = 3 and $\alpha, \beta \in \mathring{\mathcal{M}}^2$. Let $T_{\min,\alpha}, T_{\min,\beta}$ be positive closed currents with minimal singularities in α and β respectively. Then for all $x \in X$ one has

$$\lim_{k \to +\infty} \nu(T_{k,\alpha} \wedge T_{k,\beta}, x) = \nu(T_{\min,\alpha} \wedge T_{\min,\beta}, x), \tag{47}$$

where $\{T_{k,\alpha}\}$ and $\{T_{k,\beta}\}$ are sequences of currents as in Theorem (1.1) which weakly converge to $T_{min,\alpha}$ and $T_{min,\beta}$ respectively.

Proof. By using Theorem (1.20) we get the following decomposition, i.e. the Siu decomposition of $T_{min,\alpha} \wedge T_{min,\beta}$:

$$T_{\min,\alpha} \wedge T_{\min,\beta} - \langle T_{\min,\alpha} \wedge T_{\min,\beta} \rangle = \sum_{h=1}^{N} \nu(T_{\min,\alpha} \wedge T_{\min,\beta}, Y_h)[Y_h] \quad (48)$$

On the other hand by Proposition (3.17) we also have the following decomposition

$$T_{k,1} \wedge T_{k,2} - (\mu_k)_* (\beta_{k,1} \wedge \beta_{k,2}) = \sum_{h=1}^N \nu(T_{k,1} \wedge T_{k,2}, Y_h)[Y_h]$$
(49)

where in both decompositions Y_h are the irreducible components of dimension 1 of $E_{nn}(\alpha) \cap E_{nn}(\beta)$.

Then by choosing a common subsequence we obtain by uniform bound of mass and by Proposition (1.14) that

 $\nu(T_{k,1} \wedge T_{k,2}, Y_{h,k}) \rightarrow \nu_{\infty,h} \leq \nu(T_{min,\alpha} \wedge T_{min,\beta}, Y_h)$. Hence we have positive current

$$S = T_{\min,\alpha} \wedge T_{\min,\beta} - \langle T_{\min,\alpha} \wedge T_{\min,\beta} \rangle - \sum_{h=1}^{N} \nu_{\infty,h}[Y_h]$$
(50)

whose cohomology class is the class zero therefore ${\cal S}=0$ then the statement holds.

We can also conclude that

$$(\mu_k)_*(\beta_{k,1} \wedge \beta_{k,2}) \rightharpoonup \langle T_{min,\alpha} \wedge T_{min,\beta} \rangle.$$
(51)

References

- [BDPP12] S. Boucksom, J.P. Demailly, M. Păun, and T. Peternell, The pseudo-effective cone of a compact Kähler manifold and varieties of negative Kodaira dimension., J. Algebraic Geom (2012).
- [BEGZ10] S. Boucksom, P. Eyssidieux, V. Guedj, and A. Zeriahi, Monge-ampere equations in big cohomology classes, Acta Math. 205 (2010), 199–265, http://fr.arxiv.org/abs/0812.3674.
- [Bou04] S. Boucksom, Divisorial Zariski decompositions on compact complex manifolds., Ann. Sci. Ecole Norm. Sup. 37 (2004), no. 1, 45–76.
- [Dem85] J.P. Demailly, Champs magnétiques et inégalités de Morse pour la ā-cohomologie, Ann. Inst. Fourier (Grenoble) (1985), no. 35, 189–229.
- [Dem92] _____, Regularization of closed positive currents and intersection theory., J. Algebraic Geom. 1 (1992), no. 3, 361–409.
- [Dem93] _____, Monge-Ampère operators, Lelong numbers and intersection theory., Complex analysis and geometry (1993), 115–193, Univ. Ser. Math., Plenum, New York.
- [Dem00] _____, Multiplier ideal sheaves and analytic methods in algebraic geometry, Geometry (Trieste, 2000), ICTP Lect. Notes 6 (2000), 1–148.
- [DK01] J.P. Demailly and J. Kollár, Semi-continuity of complex singularity exponents and Kähler Einstein metrics on fano orbifolds., Ann. Ec. Norm. Sup 34 (2001), 525–556.
- [How01] J.A. Howald, Multiplier ideals of monomial ideals, Trans. Amer. Math. Soc. 7 (2001), no. 353, 2665–2671.
- [Laz04] R.K. Lazarsfeld, Positivity in Algebraic Geometry ii: Positivity for Vector Bundles, and Multiplier Ideals, Springer, 2004.
- [Siu93] Y.T. Siu, An effective matsusaka big theorem, Ann. Inst. Fourier (Grenoble) 43 (1993), 1387–1405.
- [Tra95] S. Trapani, Numerical criteria for the positivity of the difference of ample divisors, Math. Z. 219 (1995), no. 3, 387–401.
- [Tra11] _____, Divisorial Zariski decomposition and algebraic Morse inequalities., Proc. Amer. Math. Soc. **139** (2011), 177–183.
- [Zar62] O. Zariski, The theorem of Riemann-Roch for high multiples of an effective divisor on an algebraic surface, Ann. of Math 76 (1962), 560-615.