

EQUIVARIANT ALGEBRAIC kk -THEORY AND ADJOINTNESS THEOREMS

EUGENIA ELLIS

ABSTRACT. We introduce an equivariant algebraic kk -theory for G -algebras and G -graded algebras. We study some adjointness theorems related with crossed product, trivial action, induction and restriction. In particular we obtain an algebraic version of the Green-Julg Theorem which gives us a computational tool.

1. INTRODUCTION

Algebraic kk -theory has been introduced by G. Cortiñas and A. Thom in [2]. This is a bivariant K -theory on the category of ℓ -algebras where ℓ is a commutative ring with unit. For each pair (A, B) of ℓ -algebras a group $\mathrm{kk}(A, B)$ is defined. A category \mathfrak{Kk} is obtained whose objects are ℓ -algebras and where the morphisms from A to B are the elements of the group $\mathrm{kk}(A, B)$. The category \mathfrak{Kk} is triangulated and there is a canonical functor $j : \mathrm{Alg}_\ell \rightarrow \mathfrak{Kk}$ with universal properties. These properties are algebraic homotopy invariance, matrix invariance and excision.

The definition of algebraic kk -theory was inspired by the work of J. Cuntz [4] and N. Higson [7] on the universal properties of Kasparov KK -theory [8]. The KK -theory of separable C^* -algebras is a common generalization both of topological K -homology and topological K -theory as an additive bivariant functor. Let A, B be separable C^* -algebras. Then

$$(1.1) \quad KK_*(\mathbb{C}, B) \simeq K_*^{top}(B) \quad KK^*(A, \mathbb{C}) = K_{hom}^*(A)$$

here $K_*^{top}(B)$ denotes the K -theory of B and $K_{hom}^*(A)$ the topological K -homology of A . J. Cuntz in [3] gave another equivalent definition of the original one given in [8]. This new approach allowed to put bivariant K -theory in algebraic context. Higson in [7] stated the universal property of KK whose algebraic analogue is studied in [2], where also an analogue of (1.1) is proved. On the algebraic side, if A is an ℓ -algebra then

$$\mathrm{kk}(\ell, A) \simeq \mathrm{KH}(A)$$

here KH is Weibel's homotopy K -theory defined in [12]. We can start to build a dictionary between Kasparov's KK -theory and algebraic kk -theory in the following way

The author was partially supported by CSIC-Uruguay; by the MathAmSud network U11MATH-05 partially financed by ANII (Uruguay) and MINCYT (Argentina) and by grants UBACyT 20020100100386, and MTM2007-64704 (FEDER funds).

Kasparov's KK -theory	\leftrightarrow	algebraic kk -theory
bivariant K -theory on separable C^* -algebras $C^*\text{-Alg}$	\leftrightarrow	bivariant K -theory on ℓ -algebras Alg_ℓ
$k : C^*\text{-Alg} \rightarrow KK$	\leftrightarrow	$j : \text{Alg}_\ell \rightarrow \mathfrak{K}\mathfrak{K}$
k is stable with respect to compact operators \mathcal{K}	\leftrightarrow	j is stable with respect to $M_\infty = \bigcup_{n \in \mathbb{N}} M_n$
k is continuous homotopy invariant	\leftrightarrow	j is polynomial homotopy invariant
k is split exact	\leftrightarrow	j is excisive
k is universal for the properties described above	\leftrightarrow	j is universal for the properties described above
$KK_*(\mathbb{C}, A) \simeq K_*^{\text{top}}(A)$	\leftrightarrow	$kk_*(\ell, A) \simeq KH_*(A)$

In this paper we obtain an equivariant version of the dictionary stated above in the following sense

Equivariant Kasparov's KK -theory	\leftrightarrow	Equivariant algebraic kk -theory
bivariant K -theory on separable G - C^* -algebras	\leftrightarrow	bivariant K -theory on G -algebras
$k : G\text{-}C^*\text{-Alg} \rightarrow KK^G$	\leftrightarrow	$j^G : G\text{-Alg} \rightarrow \mathfrak{K}\mathfrak{K}^G$
k is stable with respect to $\mathcal{K}(\ell^2(G \times \mathbb{N}))$	\leftrightarrow	j^G is G -stable
k is continuous homotopy invariant	\leftrightarrow	j^G is polynomial homotopy invariant
k is split exact	\leftrightarrow	j^G is excisive
k is universal for the properties described above	\leftrightarrow	j^G is universal for the properties described above
$KK_*^G(\mathbb{C}, A) \simeq K_*^{\text{top}}(A \rtimes G)$ with G compact	\leftrightarrow	$kk^G(\ell, A) \simeq KH(A \rtimes G)$ with G finite and $1/ G \in \ell$

We also introduce a dual theory $\widehat{\mathfrak{K}\mathfrak{K}}^G$ for G -graded algebras and establish a duality result similar to that proved by Baaj and Skandalis in [1].

In Section 2 we recall some results from [2]. We take special care in the definition of equivariant matrix invariance. We introduce the concept of G -stable functor in Section 3, the rest of the concepts which appears in the equivariant dictionary are straightforward. The definition of G -stability was inspired by the definition of equivariant stability for G - C^* -algebras (see [11]). In sections 2 and 3 G is a group without any other assumption but from Section 4 and for the rest of the paper G is a countable group.

In Section 4 we introduce the appropriate brand of equivariant algebraic kk -theory in each case and establish its universal properties. For a countable group G , we

define an equivariant algebraic kk -theory $\mathfrak{K}\mathfrak{K}^G$ for the category of G -algebras and $\widehat{\mathfrak{K}\mathfrak{K}}^G$ for the category of G -graded algebras.

We study adjointness theorems in equivariant kk -theory. We put in an algebraic context some of the adjointness theorems which appear in Kasparov KK-theory. In Section 5, we define the functors of trivial action and crossed product between $\mathfrak{K}\mathfrak{K}$ and $\mathfrak{K}\mathfrak{K}^G$. The first adjointness theorem is Theorem 5.2.1 which is an algebraic version of the GREEN-JULG THEOREM. This result gives us the first computation related with homotopy K-theory. If G is a finite group, A is a G -algebra, B is an algebra and $\frac{1}{|G|} \in \ell$ then there is an isomorphism

$$\psi_{GJ} : \mathfrak{K}\mathfrak{K}^G(B^\tau, A) \rightarrow \mathfrak{K}\mathfrak{K}(B, A \rtimes G).$$

In particular, if $B = \ell$ then

$$\mathfrak{K}\mathfrak{K}^G(\ell, A) \simeq \mathfrak{K}\mathfrak{H}(A \rtimes G).$$

In Section 6 we consider H a subgroup of G . We define induction and restriction functors between $\mathfrak{K}\mathfrak{K}^G$ and $\mathfrak{K}\mathfrak{K}^H$ and study the adjointness between them. If B is an H -algebra and A is a G -algebra then there is an isomorphism

$$\psi_{IR} : \mathfrak{K}\mathfrak{K}^G(\text{Ind}_H^G B, A) \rightarrow \mathfrak{K}\mathfrak{K}^H(B, \text{Res}_G^H A).$$

This result gives us another computation. Taking H the trivial group and $B = \ell$ we obtain that

$$\mathfrak{K}\mathfrak{K}^G(\ell^{(G)}, A) \simeq \mathfrak{K}\mathfrak{H}(A) \quad \forall A \in G\text{-Alg}.$$

Here $\ell^{(G)} = \bigoplus_{g \in G} \ell$ with the regular action of G . More general, if H is a finite subgroup of G and $1/|H| \in \ell$ we combine ψ_{GJ} and ψ_{IR} and obtain

$$\mathfrak{K}\mathfrak{K}^G(\ell^{(G/H)}, A) \simeq \mathfrak{K}\mathfrak{H}(A \rtimes H) \quad \forall A \in G\text{-Alg}.$$

In Section 7 we obtain an algebraic version of the Baaž-Skandalis duality theorem. We show that the functors

$$\rtimes G : \mathfrak{K}\mathfrak{K}^G \rightarrow \widehat{\mathfrak{K}\mathfrak{K}}^G \quad G \widehat{\rtimes} : \widehat{\mathfrak{K}\mathfrak{K}}^G \rightarrow \mathfrak{K}\mathfrak{K}^G$$

are inverse category equivalences.

Acknowledgements. The results here are part of my PhD Thesis. I am thankful to Willie Cortiñas for his orientation and comments about this paper. I would like also to thank the referee for useful comments. This paper was finished during a visit to the Freie Universität Berlin (FU), financed by Berlin Mathematical School (BMS) and International Mathematical Union (IMU). I am also thankful to Freie Universität for its hospitality and to BMS-IMU for its support.

2. ALGEBRAIC KK-THEORY

In this section we recall some results from [2] and we adapt them to our setting. Let ℓ be a commutative ring with unit. We consider an ℓ -bimodule A such that $x \cdot a = a \cdot x$, with $x \in \ell$ and $a \in A$. An ℓ -bimodule A is an ℓ -algebra if it is an associative and not necessarily unital algebra. Let G be a group. A G -algebra A is an ℓ -algebra with an action of G , i.e. with a group homomorphism $\alpha : G \rightarrow \text{End}_\ell(A)$, where $\text{End}_\ell(A)$ denote the group of ℓ -linear endomorphisms of A . We shall denote by $g(a)$ or $g \cdot a$ the element $\alpha(g)(a)$. An equivariant morphism $f : A \rightarrow B$ between

G -algebras is a G -equivariant ℓ -linear map. A G -graded algebra A is an ℓ -algebra with a family of ℓ -submodules $\{A_s\}_{s \in G}$ such that

$$A = \bigoplus_{s \in G} A_s \quad A_s A_t \subseteq A_{st} \quad s, t \in G.$$

We write $|a| = s$ if $a \in A_s$. An homogeneous morphism $f : A \rightarrow B$ of G -graded algebras is an algebra morphism such that $f(A_s) \subseteq B_s$, for all $s \in G$. We consider the category of G -algebras with equivariant morphisms and the category of G -graded algebras with homogeneous morphisms.

In this section we define an algebraic kk -theory for the categories of G -algebras and G -graded algebras. We write \mathcal{C} to refer to either of these categories.

2.1. Homotopy invariance. Let A be an object of \mathcal{C} . Put $A^{\Delta^1} := A[t] = A \otimes_{\mathbb{Z}} \mathbb{Z}[t]$. Consider the trivial action of G (trivial G -grading) on $\mathbb{Z}[t]$ and the diagonal structure on $A[t]$. Then $A[t]$ is an object of \mathcal{C} . Let us write $c_A : A \rightarrow A[t]$ for the inclusion of A as constant polynomials in $A[t]$ and $\text{ev}_i : A[t] \rightarrow A$ for the evaluation of t at i ($i = 0, 1$). Note that these maps are morphisms in \mathcal{C} and that c_A is a section of ev_i .

Let $f_0, f_1 : A \rightarrow B$ be morphisms in \mathcal{C} . We call f_0 and f_1 *elementarily homotopic*, and write $f_0 \sim_e f_1$, if there exists a morphism $H : A \rightarrow B[t]$ such that $\text{ev}_i H = f_i$, $i = 0, 1$. It is easy to check that elementary homotopy is a reflexive and symmetric relation but in general it is not transitive. Let $f, g : A \rightarrow B$ be morphisms in \mathcal{C} . We call f and g *homotopic*, and write $f \sim g$, if they can be connected by a finite chain of elementary homotopies,

$$f \sim_e h_0 \sim_e \dots \sim_e h_n \sim_e g.$$

We denote the set of homotopy classes by $[A, B]_{\mathcal{C}}$. A morphism $f : A \rightarrow B$ is an *elementary homotopy equivalence* if there exists a morphism $g : B \rightarrow A$ such that $f \circ g \sim_e \text{id}_B$ and $g \circ f \sim_e \text{id}_A$. We say A is *elementarily contractible* if the null morphism and the identity morphism are elementarily homotopic.

The category of ind-objects of \mathcal{C} is the category $\text{ind-}\mathcal{C}$ of directed diagrams in \mathcal{C} . An object in $\text{ind-}\mathcal{C}$ is described by a filtering partially ordered set (I, \leq) and a functor $A : I \rightarrow \mathcal{C}$. The set of homomorphisms in $\text{ind-}\mathcal{C}$ is defined by

$$\text{hom}_{\text{ind-}\mathcal{C}}((A, I), (B, J)) := \lim_{i \in I} \text{colim}_{j \in J} \text{hom}_{\mathcal{C}}(A_i, B_j).$$

Let $A = (A, I)$ and $B = (B, J)$ be objects of $\text{ind-}\mathcal{C}$, we have a map

$$(2.1.1) \quad \text{hom}_{\text{ind-}\mathcal{C}}(A, B) \mapsto [A, B]_{\mathcal{C}} = \lim_{i \in I} \text{colim}_{j \in J} [A_i, B_j]_{\mathcal{C}}.$$

We say two morphisms in $\text{ind-}\mathcal{C}$ are *homotopic* if their images by (2.1.1) are equal. An object of $\text{ind-}\mathcal{C}$ is *contractible* if the null morphism and the identity morphism are homotopic. Sometimes we shall omit the poset in the notation.

Let \mathcal{D} be an arbitrary category. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is *homotopy invariant* if it maps the inclusion $c_A : A \rightarrow A[t]$ to an isomorphism. It is easy to check that $F : \mathcal{C} \rightarrow \mathcal{D}$ is a homotopy invariant functor if and only if $F(f) = F(g)$ when $f \sim g$ (or equivalently when $f \sim_e g$). If \mathcal{C} is the category of G -algebras we say that a homotopy invariant functor is *equivariantly homotopy invariant* and if \mathcal{C} is the category of G -graded algebras we say that a homotopy invariant functor is *graded homotopy invariant*.

2.2. Matrix stability. Consider M_n the algebra of $n \times n$ -matrices with coefficients in \mathbb{Z} with the trivial action (grading) of G and $M_\infty = \cup_{n \geq 0} M_n$. Let A be an object in \mathcal{C} . We define

$$M_n A = M_n \otimes_{\mathbb{Z}} A \quad M_\infty A = M_\infty \otimes_{\mathbb{Z}} A$$

which are objects in \mathcal{C} with the diagonal action (grading). Denote by $\iota_n : \mathbb{Z} \rightarrow M_n$ and $\iota_\infty : \mathbb{Z} \rightarrow M_\infty$ the inclusions at the upper left corner. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is M_n -stable (M_∞ -stable) if $F(\iota_n \otimes \text{id}_A)$ ($F(\iota_\infty \otimes \text{id}_A)$) is an isomorphism for all $A \in \mathcal{C}$.

2.3. Algebra of polynomial functions. Consider the following simplicial ring

$$(2.3.1) \quad \mathbb{Z}^\Delta : [n] \mapsto \mathbb{Z}^{\Delta^n} \quad \mathbb{Z}^{\Delta^n} := \mathbb{Z}[t_0, \dots, t_n] / \langle 1 - \sum_i t_i \rangle$$

$$\begin{aligned} \Theta : [n] \rightarrow [m] &\mapsto \Theta^* : \mathbb{Z}^{\Delta^m} \rightarrow \mathbb{Z}^{\Delta^n} \\ \Theta^*(t_i) &= \begin{cases} 0 & \text{si } \Theta^{-1}(i) = \emptyset \\ \sum_{j \in \Theta^{-1}(i)} t_j & \text{si } \Theta^{-1}(i) \neq \emptyset \end{cases} \end{aligned}$$

Let A be an object in \mathcal{C} . Define

$$A^\Delta : [n] \mapsto A^{\Delta^n} \quad A^{\Delta^n} := A \otimes_{\mathbb{Z}} \mathbb{Z}^{\Delta^n}$$

Note A^Δ is a simplicial \mathcal{C} -object. Let X be a simplicial set. Define

$$A^X := \text{map}_{\mathbb{S}}(X, A^\Delta)$$

If (K, \star) is a pointed simplicial set, put

$$A^{(K, \star)} := \text{map}_{\mathbb{S}_*}((K, \star), A^\Delta) = \ker(\text{map}_{\mathbb{S}}(K, A^\Delta) \rightarrow \text{map}_{\mathbb{S}}(\star, A^\Delta)) = \ker(A^K \rightarrow A)$$

If A is a G -algebra (G -graded algebra) and K a finite simplicial set, by [2, Lemma 3.1.3] we can consider A^K and $A^{(K, \star)}$ as G -algebras (G -graded algebras) taking the diagonal structure with the trivial structure in \mathbb{Z}^K and $\mathbb{Z}^{(K, \star)}$.

We will denote by $\text{sd}^\bullet X$ the following pro-simplicial set

$$\text{sd}^\bullet X : \dots \rightarrow \text{sd}^n X \xrightarrow{h} \text{sd}^{n-1} X \rightarrow \dots \rightarrow \text{sd} X \xrightarrow{h} X$$

where $\text{sd} X$ is the subdivision of X and h is the last vertex map, see [5, III.4]. If A is an object in \mathcal{C} we consider the following object in $\text{ind-}\mathcal{C}$

$$A^{\text{sd}^\bullet X} : A^X \rightarrow A^{\text{sd} X} \rightarrow \dots \rightarrow A^{\text{sd}^{n-1} X} \rightarrow A^{\text{sd}^n X} \rightarrow \dots$$

2.4. Extensions and classifying map. A sequence of morphisms in $\text{ind-}\mathcal{C}$

$$(2.4.1) \quad A \xrightarrow{f} B \xrightarrow{g} C$$

is called an extension if f is a kernel of g and g is a cokernel of f . Let $\mathcal{U}(\mathcal{C})$ be the category of modules with linear and equivariant maps (graded maps), i.e. we forget the multiplication in \mathcal{C} keeping the module structure. Let $F : \mathcal{C} \rightarrow \mathcal{U}(\mathcal{C})$ be the forgetful functor. This functor can be extended to $F : \text{ind-}\mathcal{C} \rightarrow \text{ind-}\mathcal{U}(\mathcal{C})$. We will call an extension (2.4.1) weakly split if $F(g)$ has a section in $\text{ind-}\mathcal{U}(\mathcal{C})$.

Let M be an object in $\mathcal{U}(\mathcal{C})$. Consider in

$$\tilde{T}(M) = \bigoplus_{n \geq 1} M^{\otimes n} \quad M^{\otimes n} = \underbrace{M \otimes \dots \otimes M}_{n\text{-times}}$$

the usual structure of ℓ -algebra. If \mathcal{C} is the category of G -algebras, we consider in $M^{\otimes n}$ the following action,

$$g \cdot (m_1 \otimes m_2 \otimes \dots \otimes m_n) = g \cdot m_1 \otimes g \cdot m_2 \otimes \dots \otimes g \cdot m_n$$

which gives to $\tilde{T}(M)$ a G -algebra structure. If \mathcal{C} is the category of G -graded algebras, take in $M^{\otimes n}$ the following G -graduation

$$|m_1 \otimes m_2 \otimes \dots \otimes m_n| = |m_1| |m_2| \dots |m_n|.$$

By this way $\tilde{T}(M)$ is a G -graded algebra. Both constructions are functorial hence we consider the functor $\tilde{T} : \mathcal{U}(\mathcal{C}) \rightarrow \mathcal{C}$. Put

$$T := \tilde{T} \circ F : \mathcal{C} \rightarrow \mathcal{C}$$

If A is an object in \mathcal{C} there exists an morphism in \mathcal{C}

$$\eta_A : T(A) \rightarrow A \quad \eta_A(a_1 \otimes \dots \otimes a_n) = a_1 \dots a_n$$

and a morphism in $\mathcal{U}(\mathcal{C})$ $\mu_A : A \rightarrow T(A)$ which is the inclusion at the first summand of $T(A)$. Let A, B be objects in \mathcal{C} , it is easy to check that

$$(2.4.2) \quad \text{hom}_{\mathcal{C}}(T(A), B) \simeq \text{hom}_{\mathcal{U}(\mathcal{C})}(F(A), F(B)).$$

Hence if we have a morphism $A \rightarrow B$ in $\mathcal{U}(\mathcal{C})$, we can extend it to a morphism $T(A) \rightarrow B$ in \mathcal{C} . It shows that T is the left adjoint of F . The counit of the adjunction is $\eta_A : T(A) \rightarrow A$ and it is surjective (see [10, IV.3 Thm 1]). We define

$$J(A) := \ker \eta_A.$$

The universal extension of A is

$$J(A) \xrightarrow{\iota_A} T(A) \xrightarrow{\eta_A} A.$$

Let $A \xrightarrow{f} B \xrightarrow{g} C$ be a weakly split extension. Let s be a section of $F(g)$ and define $\hat{\xi} = \eta_B \circ \tilde{T}(s)$, then

$$\eta_C = \eta_C \circ T(g) \circ \tilde{T}(s) = g \circ \eta_B \circ \tilde{T}(s) = g \circ \hat{\xi}.$$

Define $\xi : J(C) \rightarrow A$ as the restriction of $\hat{\xi}$ to $J(C)$. We obtain a commutative diagram of extensions

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ \xi \uparrow & & \hat{\xi} \uparrow & & \uparrow \text{id}_C \\ J(C) & \xrightarrow{\iota_C} & T(C) & \xrightarrow{\eta_C} & C \end{array}$$

This construction of ξ depends on which s is chosen. If $\xi_1, \xi_2 : J(C) \rightarrow A$ are morphisms constructed taking different sections of $F(g)$ we will show ξ_1, ξ_2 are homotopic. Define a linear equivariant (graded) map

$$H : C \rightarrow A[t] \quad H(c) = (1-t)\xi_1(c) + t\xi_2(c).$$

By (2.4.2) it extends to a homomorphism and there exists a morphism $H : T(C) \rightarrow A[t]$ in \mathcal{C} such that

$$\text{ev}_0 \circ H|_{J(C)} = \xi_1 \quad \text{ev}_1 \circ H|_{J(C)} = \xi_2.$$

Then the map ξ is unique up to elementary homotopy. We call ξ the classifying map of the extension $A \xrightarrow{f} B \xrightarrow{g} C$. This construction is functorial, that means if we have a diagram of weakly split extensions,

$$\begin{array}{ccccc} A & \longrightarrow & B & \longrightarrow & C \\ f \downarrow & & \downarrow h & & \downarrow g \\ A' & \longrightarrow & B' & \longrightarrow & C' \end{array}$$

then there is a diagram

$$\begin{array}{ccc} J(C) & \longrightarrow & A \\ J(g) \downarrow & & \downarrow f \\ J(C') & \longrightarrow & A' \end{array}$$

of classifying maps, which is commutative up to elementary homotopy.

Let L be a ring and A an object in \mathcal{C} . Then the extension

$$(2.4.3) \quad J(A) \otimes_{\mathbb{Z}} L \rightarrow T(A) \otimes_{\mathbb{Z}} L \rightarrow A \otimes_{\mathbb{Z}} L$$

is weakly split, and there is a choice for classifying map

$$\phi_{A,L} : J(A \otimes_{\mathbb{Z}} L) \rightarrow J(A) \otimes_{\mathbb{Z}} L$$

of (2.4.3), which is natural in both variables. In particular, if K is a finite pointed simplicial set. There is a homotopy class of maps

$$(2.4.4) \quad \phi_{A,K} : J(A^K) \rightarrow J(A)^K$$

natural with respect to K , which is represented by a classifying map of the following extension

$$J(A)^K \xrightarrow{\iota_A^*} T(A)^K \xrightarrow{\eta_A^*} A^K$$

Let S^1 be the simplicial circle $\Delta^1/\partial\Delta^1$, we define

$$\Omega := \mathbb{Z}^{(S^1, \star)} \quad P := \mathbb{Z}^{(\Delta^1, \star)}$$

The path extension of A is the extension induced by the cofibration $\partial\Delta^1 \subset \Delta^1$ (see [2, Lemma 3.1.2]),

$$(2.4.5) \quad \Omega A \longrightarrow PA \xrightarrow{(ev_0, ev_1)} A \oplus A$$

The extension (2.4.5) is weakly split because we have a linear and equivariant (graded) section of (ev_0, ev_1)

$$(2.4.6) \quad (a, b) \mapsto (1-t)a + tb.$$

The loop extension of A is

$$(2.4.7) \quad \Omega A \rightarrow PA \xrightarrow{ev_1} A$$

Note $a \mapsto at$ is a natural section of $F(ev_1)$. Thus we can pick a natural choice for the classifying map of (2.4.7). We call it

$$\rho_A : J(A) \rightarrow \Omega A$$

Let $\mathcal{P} := \mathbb{Z}^{(sd^\bullet \Delta^1, \star)}$ we have an extension in $\text{ind-}\mathcal{C}$

$$A^{S^1} := A^{(sd^\bullet S^1, \star)} \quad A^{S^1} \rightarrow \mathcal{P}A \xrightarrow{ev_1} A$$

which is naturally weakly split. The classifying map $J(A) \rightarrow A^{\mathcal{S}^1}$ is the following composition

$$(2.4.8) \quad J(A) \xrightarrow{\rho_A} \Omega A \xrightarrow{h} A^{\mathcal{S}^1}$$

where h is induced by the last vertex map. We will sometimes abuse notation and write ρ_A for the map (2.4.8).

Let $f : A \rightarrow B$ be a morphism in \mathcal{C} . The mapping path extension of f is the extension obtained from the loop extension of B by pulling it back to A

$$P_f := PB \times_B A \quad \begin{array}{ccccc} \Omega B & \xrightarrow{\iota} & PB \times_B A & \xrightarrow{\pi_f} & A & E' \\ \downarrow & & \downarrow & & \downarrow f & \\ \Omega B & \xrightarrow{\iota} & PB & \xrightarrow{\text{ev}_1} & B & E \end{array}$$

We call P_f the **path algebra** of f . Note the extension E' is naturally weakly split because $\tilde{s}(a) = (s \circ f(a), a)$ is a natural section of π_f where s is the natural section of E . We define

$$(2.4.9) \quad \mathcal{P}_f := \mathcal{P}B \times_B A \quad B^{\mathcal{S}^1} \rightarrow \mathcal{P}B \times_B A \xrightarrow{\pi_f} A.$$

Note $\rho_f := \rho_B J(f)$ is the classifying map of the extension (2.4.9).

2.5. Excisive homology theories. We consider triangulated categories in terms of a loop functor Ω . A triangulated category $(\mathcal{T}, \Omega, \mathcal{Q})$ is an additive category \mathcal{T} with an equivalence $\Omega : \mathcal{T} \rightarrow \mathcal{T}$ and a class \mathcal{Q} of sequences in \mathcal{T} called **distinguished triangles**

$$(T) \quad \Omega C \rightarrow A \rightarrow B \rightarrow C$$

satisfying some axioms, see [9], [2]. A triangle functor from $(\mathcal{T}_1, \Omega_1, \mathcal{Q}_1)$ to $(\mathcal{T}_2, \Omega_2, \mathcal{Q}_2)$ is a pair consisting of an additive functor $R : \mathcal{T}_1 \rightarrow \mathcal{T}_2$ and a natural transformation $\alpha : \Omega_2 R \rightarrow R \Omega_1$ such that

$$\Omega_2 R(C) \xrightarrow{R(f) \circ \alpha_C} R(A) \xrightarrow{R(g)} R(B) \xrightarrow{R(h)} R(C)$$

is a distinguished triangle in \mathcal{T}_2 for each triangle $\Omega_1 C \xrightarrow{f} A \xrightarrow{g} B \xrightarrow{h} C$.

Let $(\mathcal{T}, \Omega, \mathcal{Q})$ be a triangulated category. An excisive homology theory for \mathcal{C} with values in \mathcal{T} is an \mathcal{E} -excisive homology, as defined in [2, Sec 6.6], when \mathcal{E} is the class of weakly split extensions.

Let P be a property for functors defined on \mathcal{C} to some triangulated category. A functor $u : \mathcal{C} \rightarrow \mathcal{V}$ is a **universal functor** with P if it has the property P and if $F : \mathcal{C} \rightarrow \mathcal{T}$ is another functor with P there exists a unique triangle functor $G : \mathcal{V} \rightarrow \mathcal{T}$ such that the following diagram is commutative

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{u} & \mathcal{V} \\ & \searrow F & \downarrow G \\ & & \mathcal{T} \end{array}$$

2.6. The category $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$. Let A and B be objects in \mathcal{C} . Consider \mathcal{M}_{∞} the ind-ring defined in [2, Sec 4.1] and define inductively $B^{\mathcal{S}^{n+1}} := (B^{\mathcal{S}^n})^{\mathcal{S}^1}$. Define

$$E_n(A, B)_{\mathcal{C}} := [J^n(A), \mathcal{M}_{\infty} B^{\mathcal{S}^n}]_{\mathcal{C}}.$$

Consider the following morphism $\iota_n : E_n(A, B)_{\mathcal{C}} \rightarrow E_{n+1}(A, B)_{\mathcal{C}}$ such that

$$\begin{array}{c} J^n(A) \xrightarrow{f} \mathcal{M}_{\infty} B^{\mathcal{S}^n} \\ \downarrow \\ J^{n+1}(A) \xrightarrow{J(f)} J(\mathcal{M}_{\infty} B^{\mathcal{S}^n}) \xrightarrow{\phi_{\mathcal{M}_{\infty}, B^{\mathcal{S}^n}}} \mathcal{M}_{\infty} J(B^{\mathcal{S}^n}) \xrightarrow{\rho_{B^{\mathcal{S}^n}}} \mathcal{M}_{\infty} B^{\mathcal{S}^{n+1}} \end{array}$$

Define

$$\mathrm{kk}_{\mathcal{C}}(A, B) = \mathrm{colim}_{n \in \mathbb{N}} E_n(A, B)_{\mathcal{C}}.$$

Let A, B and C be objects in \mathcal{C} . There exists an associative product

$$\circ : \mathrm{kk}_{\mathcal{C}}(B, C) \times \mathrm{kk}_{\mathcal{C}}(A, B) \rightarrow \mathrm{kk}_{\mathcal{C}}(A, C)$$

which extends the composition of algebra homomorphisms. If $[\alpha] \in \mathrm{kk}_{\mathcal{C}}(B, C)$ is an element represented by $\alpha : J^n(B) \rightarrow C^{\mathcal{S}^n}$ and $[\beta] \in \mathrm{kk}_{\mathcal{C}}(A, B)$ is an element represented by $\beta : J^m(A) \rightarrow B^{\mathcal{S}^m}$ then $[\alpha] \circ [\beta]$ is an element $\mathrm{kk}_{\mathcal{C}}(A, C)$ represented by

$$(2.6.1) \quad J^{n+m}(A) \xrightarrow{J^n(\beta)} J^n(B^{\mathcal{S}^m}) \rightarrow J^n(B)^{\mathcal{S}^m} \xrightarrow{\alpha^{\mathcal{S}^m}} C^{\mathcal{S}^{n+m}}.$$

This product allows us to define a composition in the category $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ whose objects are the same objects of \mathcal{C} and the morphisms from A to B are the elements of $\mathrm{kk}_{\mathcal{C}}(A, B)$. Denote by

$$j_{\mathcal{C}} : \mathcal{C} \rightarrow \mathfrak{K}\mathfrak{K}_{\mathcal{C}}$$

the functor which at the level of objects is the identity and at level of morphism sends $f : A \rightarrow B$ to $[f] \in \mathrm{kk}_{\mathcal{C}}(A, B)$.

Remark 2.6.2. A morphism $f : C \rightarrow M_{\infty} C$ in \mathcal{C} represents an element $[f]$ in $\mathrm{kk}_{\mathcal{C}}(C, \mathcal{M}_{\infty} C)$. But also represents an element in $\mathrm{kk}_{\mathcal{C}}(C, C)$ because

$$\mathrm{kk}_{\mathcal{C}}(C, C) = \mathrm{colim} E_n(C, C)_{\mathcal{C}} \quad \text{and} \quad E_0(C, C)_{\mathcal{C}} = [C, \mathcal{M}_{\infty} C]_{\mathcal{C}}.$$

Consider the functor $\Omega : \mathfrak{K}\mathfrak{K}_{\mathcal{C}} \rightarrow \mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ which sends an object A of \mathcal{C} to the path object ΩA . Let $[\alpha]$ be an element of $\mathrm{kk}_{\mathcal{C}}(A, B)$ represented by $\alpha : J^n(A) \rightarrow B^{\mathcal{S}^n}$. The class of $\Omega[\alpha]$ is represented by

$$(2.6.3) \quad J^n(A^{\mathcal{S}^1}) \rightarrow J^n(A)^{\mathcal{S}^1} \xrightarrow{\alpha^{\mathcal{S}^1}} B^{\mathcal{S}^{n+1}}$$

Note (2.6.3) represents an element of $\mathrm{kk}_{\mathcal{C}}(A^{\mathcal{S}^1}, B^{\mathcal{S}^1})$ (see Lemma 6.3.8 [2] to check it is well defined). As $\iota : \Omega A \rightarrow A^{\mathcal{S}^1}$ is a $\mathrm{kk}_{\mathcal{C}}$ -equivalence (see corollary of Lemma 6.3.2 [2]), we have the natural isomorphism

$$\mathrm{kk}_{\mathcal{C}}(A^{\mathcal{S}^1}, B^{\mathcal{S}^1}) \xrightarrow{\sim} \mathrm{kk}_{\mathcal{C}}(\Omega A, \Omega B).$$

A diagram

$$\Omega C \rightarrow A \rightarrow B \rightarrow C$$

of morphisms in $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ is called a **distinguished triangle** if it is isomorphic in $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ to the path sequence

$$\Omega B' \xrightarrow{j(\iota)} P_f \xrightarrow{j(\pi_f)} A' \xrightarrow{j(f)} B'$$

associated with a homomorphism $f : A' \rightarrow B'$ in \mathcal{C} . Denote the class of distinguished triangles by \mathcal{Q} . The category $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ is triangulated with respect to the

endofunctor $\Omega : \mathfrak{K}\mathfrak{K}_{\mathcal{C}} \rightarrow \mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ and the class \mathcal{Q} of distinguished triangles, see [2, Theorem 6.5.2].

Let $E : A \xrightarrow{f} B \xrightarrow{g} C$ be a weakly split extension and let $c_E \in \text{kk}_{\mathcal{C}}(J(C), A)$ be the classifying map of E . As the natural map $\rho_A : J(A) \rightarrow \Omega A$ induces a $\text{kk}_{\mathcal{C}}$ -equivalence (see Lemma 6.3.10, [2]) we can consider the following morphisms in $\text{kk}_{\mathcal{C}}(\Omega C, A)$

$$(2.6.4) \quad \partial_E := c_E \circ \rho_C^{-1}.$$

The functor $j_{\mathcal{C}} : \mathcal{C} \rightarrow \mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ with the morphisms $\{\partial_E : E \in \mathcal{E}\}$ is an excisive homology theory, homotopy invariant and M_{∞} -stable.

Theorem 2.6.5. *The functor $j_{\mathcal{C}} : \mathcal{C} \rightarrow \mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ is universal with the properties defined above. In other words, if \mathcal{T} is a triangulated category and $G : \mathcal{C} \rightarrow \mathcal{T}$ together a class of morphisms $\{\bar{\partial}_E : E \in \mathcal{E}\}$ is an excisive, homotopy invariant and M_{∞} -stable functor, then there exists a unique triangle functor $\bar{G} : \mathfrak{K}\mathfrak{K}_{\mathcal{C}} \rightarrow \mathcal{T}$ such that the following diagram commutes*

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{j_{\mathcal{C}}} & \mathfrak{K}\mathfrak{K}_{\mathcal{C}} \\ & \searrow G & \downarrow \bar{G} \\ & & \mathcal{T} \end{array}$$

Proof. By definition $j_{\mathcal{C}}$ is homotopy invariant and M_{∞} -stable. Let us show that $j_{\mathcal{C}} : \mathcal{C} \rightarrow \mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ is an excisive homology theory with the family $\{\partial_E : E \in \mathcal{E}\}$. Let $E : A \xrightarrow{f} B \xrightarrow{g} C$ be a weakly split extension. Take the path sequence associated to g and the following diagram in $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$

$$\begin{array}{ccccccc} \Omega C & \xrightarrow{j(\iota)} & P_g & \xrightarrow{j(\pi_g)} & B & \xrightarrow{j(g)} & C & T \\ \text{id} \uparrow & & \iota_f \uparrow & & \text{id} \uparrow & & \text{id} \uparrow & \\ \Omega C & \xrightarrow{\partial_E} & A & \xrightarrow{j(f)} & B & \xrightarrow{j(g)} & C & T' \end{array}$$

The first square commutes because $\iota_f \circ c_E$ is elementarily homotopic to $\iota \circ \rho_C$. By [2, Lemma 6.3.2], the morphism ι_f is a $\text{kk}_{\mathcal{C}}$ -equivalence. Finally T and T' are isomorphic in $\mathfrak{K}\mathfrak{K}_{\mathcal{C}}$ and T' is a distinguished triangle. For the rest of the proof see [2, Theorem 6.6.2]. \square

Algebraic K -theory is not excisive, nor is it M_{∞} -stable or homotopy invariant. However the homotopy algebraic K -theory, defined in [12] and denoted by KH , have all these properties. Consider the functor $\text{KH} = \text{KH}_0 : \text{Alg}_{\ell} \rightarrow \text{Ab}$. Because it satisfies [2, Theorem 6.6.6] we have a natural map

$$\text{kk}_{\text{Alg}_{\ell}}(\ell, A) \rightarrow \text{hom}_{\text{Ab}}(\text{KH}(\ell), \text{KH}(A))$$

As ℓ is unital, there is a map $\mathbb{Z} \rightarrow \ell$ which induce a map

$$\text{hom}_{\text{Ab}}(\text{KH}(\ell), \text{KH}(A)) \mapsto \text{hom}_{\text{Ab}}(\text{KH}(\mathbb{Z}), \text{KH}(A)) \simeq \text{KH}(A).$$

Composing both maps we obtain a homomorphism $\text{kk}_{\text{Alg}_{\ell}}(\ell, A) \rightarrow \text{KH}(A)$ and the main theorem from [2] prove that it is an isomorphism.

Theorem 2.6.6. *Consider $\mathcal{C} = \text{Alg}_{\ell}$ the category of ℓ -algebras, then*

$$\text{kk}_{\text{Alg}_{\ell}}(\ell, A) \simeq \text{KH}(A).$$

Proof. See [2, Theorem 8.2.1]. \square

3. EQUIVARIANT MATRIX INVARIANCE

In the equivariant setting we replace the property of M_∞ -stability by a stability condition depending on \mathcal{C} . We consider the different cases of \mathcal{C} separately.

3.1. G -equivariant stability. Regard

$$M_G = \{f : G \times G \rightarrow \ell : \text{supp}(f) < \infty\}$$

as the algebra of matrices with coefficients in ℓ indexed by $G \times G$, with translation action of G :

$$g \cdot e_{s,t} = e_{gs,gt}.$$

We are going to identify a G -algebra A with the G -algebra $M_G \otimes A$ with the diagonal action. Note that the map $a \rightarrow a \otimes e_{1_G, 1_G}$ is not equivariant then we can not define G -stability as in Section 2.2. For this reason we define G -stability in terms of G -modules.

A pair (\mathcal{W}, B) is a G -module with basis if \mathcal{W} is a G -module, free as an ℓ -module and B is a basis of \mathcal{W} . A pair (\mathcal{W}', B') is a submodule with basis of (\mathcal{W}, B) if \mathcal{W}' is a submodule of \mathcal{W} and $B' \subset B$. Note that if (\mathcal{W}_1, B_1) and (\mathcal{W}_2, B_2) are G -modules with basis then $(\mathcal{W}_1 \oplus \mathcal{W}_2, B_1 \sqcup B_2)$ is a G -module with basis.

Let (\mathcal{W}, B) be a G -module with basis B . We define

$$\mathcal{L}(\mathcal{W}, B) := \{\psi : B \times B \rightarrow \ell : \{v : \psi(v, w) \neq 0\} \text{ is finite for all } w\}$$

Note that $\mathcal{L}(\mathcal{W}, B)$ and $\text{End}_\ell(\mathcal{W}) = \{f : \mathcal{W} \rightarrow \mathcal{W} : f \text{ is } \ell\text{-linear}\}$ are isomorphic; indeed we have inverse isomorphisms

$$(3.1.1) \quad \text{End}_\ell(\mathcal{W}) \rightarrow \mathcal{L}(\mathcal{W}, B) \quad f \mapsto \psi_f \quad \psi_f(v, w) = p_v(f(w))$$

here $p_v : \mathcal{W} \rightarrow \ell$ is the projection to the submodule of \mathcal{W} generated by v

$$(3.1.2) \quad \mathcal{L}(\mathcal{W}, B) \rightarrow \text{End}_\ell(\mathcal{W}) \quad \psi \mapsto f_\psi \quad f_\psi(w) = \sum_{v \in B} \psi(v, w)v.$$

Define

$$\begin{aligned} \mathcal{C}(\mathcal{W}, B) &:= \{\psi \in \mathcal{L}(\mathcal{W}, B) : \{w : \psi(v, w) \neq 0\} \text{ is finite for all } v\} \\ \mathcal{F}(\mathcal{W}, B) &:= \{\psi \in \mathcal{L}(\mathcal{W}, B) : \{(v, w) : \psi(v, w) \neq 0\} \text{ is finite}\} \\ \text{End}_\ell^F(\mathcal{W}, B) &:= \{f \in \text{End}_\ell(\mathcal{W}) : \psi_f \in \mathcal{F}(\mathcal{W}, B)\} \\ \text{End}_\ell^C(\mathcal{W}, B) &:= \{f \in \text{End}_\ell(\mathcal{W}) : \psi_f \in \mathcal{C}(\mathcal{W}, B)\} \end{aligned}$$

Note that $\mathcal{C}(\mathcal{W}, B)$ is a ring with the matrix product and $\text{End}_\ell^C(\mathcal{W}, B)$ is a ring with the composition. These rings are isomorphic.

Let (\mathcal{W}, B) be a G -module with basis. Consider the representation

$$\rho : G \rightarrow \text{End}_\ell(\mathcal{W}) \quad \rho_g(w) = g \cdot w$$

We say that (\mathcal{W}, B) is a G -module by finite automorphisms if $\rho(G) \subset \text{End}_\ell^F(\mathcal{W}, B)$.

We say that (\mathcal{W}, B) is a G -module by locally finite automorphisms if $\rho(G) \subset \text{End}_\ell^C(\mathcal{W}, B)$.

If (\mathcal{W}, B) is a G -module by locally finite automorphisms, $\text{End}_\ell^C(\mathcal{W}, B)$ and $\text{End}_\ell^F(\mathcal{W}, B)$ are G -algebras with the following action

$$g \cdot f = \rho(g)f(\rho(g))^{-1}$$

Note that $\text{End}_\ell^F(\mathcal{W}, B)$ is an ideal of $\text{End}_\ell^C(\mathcal{W}, B)$.

Example 3.1.3. Let $\mathcal{W} = \ell G$ be the group algebra considered as a G -module via the regular representation with basis $B = \{\delta_g : g \in G\}$,

$$(3.1.4) \quad g \cdot \left(\sum_{h \in G} a_h \delta_h \right) = \sum_{h \in G} a_h \delta_{gh}, \quad a_h \in \ell.$$

Note

$$\rho : G \rightarrow \text{End}_\ell(\ell G) \simeq \mathcal{L}(\ell G, B) \quad g \mapsto M_g = \sum_{t \in G} e_{gt,t}$$

As $M_g \in \mathcal{C}(\ell G, B)$ for all $g \in G$, $(\ell G, B)$ is a G -module by locally finite automorphisms. Moreover we have $(M_g)^{-1} = M_{g^{-1}} = (M_g)^t$ and $M_G = \mathcal{F}(\ell G, B)$.

Let A be a G -algebra. Consider the tensor product $M_G A = M_G \otimes A$ with the diagonal action of G .

Remark 3.1.5. Let \mathcal{W} be a G -module. Let \mathcal{W}^τ be \mathcal{W} considered as a G -module with trivial action. Recall that $\ell G \otimes \mathcal{W} \simeq \ell G \otimes \mathcal{W}^\tau$. Inverse isomorphisms are given by

$$\begin{aligned} T : \ell G \otimes \mathcal{W}^\tau &\rightarrow \ell G \otimes \mathcal{W} & S : \ell G \otimes \mathcal{W} &\rightarrow \ell G \otimes \mathcal{W}^\tau \\ T(\delta_g \otimes h) &= \delta_g \otimes g(h) & S(\delta_g \otimes h) &= \delta_g \otimes g^{-1}(h) \end{aligned}$$

If (\mathcal{W}, B) is a G -module with basis we will write $\text{End}_\ell^C(\mathcal{W})$ and $\text{End}_\ell^F(\mathcal{W})$ omitting the basis when there is no confusion.

Remark 3.1.6. Let (\mathcal{W}, B) be a G -module with basis. Let us check that

$$\text{End}_\ell^F(\ell G \otimes \mathcal{W}) \simeq \text{End}_\ell^F(\ell G) \otimes \text{End}_\ell^F(\mathcal{W}).$$

Define a G -algebra homomorphism $T : \text{End}_\ell^F(\ell G) \otimes \text{End}_\ell^F(\mathcal{W}) \rightarrow \text{End}_\ell^F(\ell G \otimes \mathcal{W})$ by

$$T(e_{g,h} \otimes e_{v,w}) = e_{g,v,h,w} \quad v, w \in B \quad g, h \in G$$

As T is a bijection between the basis, T is an isomorphism.

Let (\mathcal{W}_1, B_1) and (\mathcal{W}_2, B_2) be G -modules by locally finite automorphisms such that $\text{card}(B_i) \leq \text{card}(G) \times \text{card}\mathbb{N}$, $i = 1, 2$. The inclusion $\iota : \mathcal{W}_1 \rightarrow \mathcal{W}_1 \oplus \mathcal{W}_2$ induces a morphism of G -algebras

$$(3.1.7) \quad \tilde{\iota} : \text{End}_\ell^F(\mathcal{W}_1) \rightarrow \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \quad f \mapsto \begin{pmatrix} f & 0 \\ 0 & 0 \end{pmatrix}$$

Let A be a G -algebra and consider

$$\tilde{\iota} \otimes 1 : \text{End}_\ell^F(\mathcal{W}_1) \otimes A \rightarrow \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A.$$

A functor $F : G\text{-Alg} \rightarrow \mathcal{D}$ is G -stable if for (\mathcal{W}_1, B_1) , (\mathcal{W}_2, B_2) and A as above $F(\tilde{\iota} \otimes 1)$ is an isomorphism in \mathcal{D} .

Let us show that if $F : G\text{-Alg} \rightarrow \mathcal{D}$ is a G -stable functor then F is M_∞ -stable. Consider $(\mathcal{W}_1, B_1) = (\ell, \{1\})$ and $(\mathcal{W}_2, B_2) = (\ell^{(\mathbb{N})}, \{e_i : i \in \mathbb{N}\})$ with $\ell^{(\mathbb{N})} = \bigoplus_{i=1}^\infty \ell$, $\{e_i : i \in \mathbb{N}\}$ is the canonical basis and both modules have the trivial action of G . Note

$$\text{End}_\ell^F(\ell) = \text{End}_\ell(\ell) = \ell \quad \text{End}_\ell^F(\ell \oplus \ell^{(\mathbb{N})}) = \text{End}_\ell^F(\ell^{(\mathbb{N})}) = M_\infty$$

and $\tilde{\iota}_\infty : \ell \rightarrow M_\infty$ is the inclusion at the upper left corner. Then $\tilde{\iota} \otimes 1 = \iota : A \rightarrow M_\infty(A)$ and $F(\iota)$ is an isomorphism. Observe that if $G = \{e\}$, F is G -stable if and only if F is M_∞ -stable.

Let A, B be G -algebras and $F : G\text{-Alg} \rightarrow \mathcal{D}$ a functor. A zig-zag between A and B by F is a diagram in $G\text{-Alg}$

$$A \xrightarrow{f_1} C_1 \xleftarrow{g_1} \dots \xrightarrow{f_n} C_n \xleftarrow{g_n} B$$

such that $F(g_i)$, $i = 1, \dots, n$, are isomorphisms in \mathcal{D} .

Example 3.1.8. Let A be a G -algebra and F a G -stable functor. There exists a zig-zag between A and $M_G A$ by F . Consider $\mathcal{W}_1 = (\ell G, B)$ as in the example 3.1.3 and consider $\mathcal{W}_2 = (\ell, \{1\})$ with the trivial action of G . Put $\mathcal{W} = \mathcal{W}_1 \oplus \mathcal{W}_2$ and $C = \text{End}_\ell^F(\mathcal{W})$ with the induced action, then

$$\iota : A = A \otimes \ell = A \otimes \text{End}_\ell^F(\ell) \rightarrow A \otimes C \leftarrow A \otimes M_G : \iota'$$

is a zig-zag between A and $M_G A$ by F .

Proposition 3.1.9. *Suppose G is countable. Let $F : G\text{-Alg} \rightarrow \mathcal{D}$ be an M_∞ -stable functor. Define*

$$\hat{F} : G\text{-Alg} \rightarrow \mathcal{D} \quad A \mapsto F(M_G \otimes A)$$

Then \hat{F} is G -stable.

Proof. Let $(\mathcal{W}_1, B_1), (\mathcal{W}_2, B_2)$ be G -modules by locally finite automorphisms and let A be a G -algebra. Consider

$$\tilde{\iota} \otimes 1 : \text{End}_\ell^F(\mathcal{W}_1) \otimes A \rightarrow \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A.$$

We have to prove that

$$(3.1.10) \quad \hat{F}(\tilde{\iota} \otimes 1) : F(M_G \otimes \text{End}_\ell^F(\mathcal{W}_1) \otimes A) \rightarrow F(M_G \otimes \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A)$$

is an isomorphism. By remarks 3.1.6 and 3.1.5 we know that

$$F(M_G \otimes \text{End}_\ell^F(\mathcal{W}_1) \otimes A) \simeq F(\text{End}_\ell^F(\mathcal{W}_1^\tau) \otimes M_G \otimes A)$$

and

$$F(M_G \otimes \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \otimes A) \simeq F(\text{End}_\ell^F((\mathcal{W}_1 \oplus \mathcal{W}_2)^\tau) \otimes M_G \otimes A).$$

Note that $\text{End}_\ell^F(\mathcal{W}_1^\tau)$ and $\text{End}_\ell^F((\mathcal{W}_1 \oplus \mathcal{W}_2)^\tau)$ are equivariantly isomorphic to M_n or M_∞ . As F is M_∞ -stable, (3.1.10) is an isomorphism. \square

Remark 3.1.11. Suppose G is a finite group of order n such that $1/n \in \ell$. Then the element $\xi = (1/n) \sum_{g \in G} \delta_g$ in ℓG is idempotent. The map $s : \ell \rightarrow \ell G$, $s(1) = \xi$, is a G -equivariant section of the canonical augmentation $\varphi : \ell G \rightarrow \ell$. Thus the sequence of G -modules

$$(3.1.12) \quad 0 \longrightarrow I \longrightarrow \ell G \xrightarrow{\varphi} \ell \longrightarrow 0$$

splits. Hence $\ell G = \ell \xi \oplus I$. Notice that I is a G -module with basis $\{\delta_e - \delta_g : g \neq e\}$. Define

$$\lambda_g = \begin{cases} \xi & g = e \\ \delta_e - \delta_g & g \neq e \end{cases}$$

The set $\Lambda = \{\lambda_g : g \in G\}$ is a basis of ℓG and the relations with the elements of $B = \{\delta_g : g \in G\}$ are the following

$$\begin{aligned} \lambda_e &= \frac{1}{n} \sum_{g \in G} \delta_g & \lambda_h &= \delta_e - \delta_h & h &\neq e \\ \delta_e &= \lambda_e + \frac{1}{n} \sum_{g \neq e} \lambda_g & \delta_h &= \lambda_e - \lambda_h + \frac{1}{n} \sum_{g \neq e} \lambda_g \end{aligned}$$

Consider $\mathcal{W}_1 = \ell = (\ell\xi, \{\xi\})$ and $\mathcal{W}_2 = (I, \{\lambda_g\}_{g \neq e})$, in this case the morphism (3.1.7) is

$$\iota : \ell \rightarrow M_G \simeq \text{End}(\ell G, \Lambda) \quad 1 \mapsto \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{pmatrix}$$

If we write it in the canonical basis B we have

$$\bar{\iota} : \ell \rightarrow M_G = \text{End}(\ell G, B) \quad 1 \mapsto \begin{pmatrix} \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \\ \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{n} & \frac{1}{n} & \dots & \frac{1}{n} \end{pmatrix}$$

If $F : G\text{-Alg} \rightarrow \mathcal{D}$ is a G -stable functor then $F(\bar{\iota})$ is an isomorphism in \mathcal{D} .

3.2. G -graded stability. In this section we consider a dual notion of G -equivariant stability. We want to identify a G -graded algebra A with the G -graded matrix algebra $M_G A$. The definition of G -graded stability is easier than that of G -equivariant stability because the morphism $A \rightarrow M_G A$, $a \mapsto e_{1_G, 1_G} \otimes a$, is homogeneous.

Write $G_{gr}\text{-Alg}$ for the category of G -graded algebras. Let A be a G -graded algebra. Define the following grading in $M_G A$

$$(3.2.1) \quad (M_G A)_g := \langle e_{s,t} \otimes a : g = s|a|t^{-1} \rangle.$$

Depending on the context $M_G A$ will be considered as an algebra, a G -algebra or a G -graded algebra. Let be

$$\iota_A : A \rightarrow M_G A \quad a \mapsto e_{1_G, 1_G} \otimes a,$$

note ι_A is homogeneous. A functor $F : G_{gr}\text{-Alg} \rightarrow \mathcal{D}$ is G_{gr} -stable if $F(\iota_A)$ is an isomorphism in \mathcal{D} for all $A \in G_{gr}\text{-Alg}$.

Proposition 3.2.2. *Suppose G is countable. Let $F : G_{gr}\text{-Alg} \rightarrow \mathcal{D}$ be an M_∞ -stable functor. Define*

$$\hat{F} : G_{gr}\text{-Alg} \rightarrow \mathcal{D} \quad A \mapsto F(M_G A)$$

with $M_G A$ as in (3.2.1). Then \hat{F} is G_{gr} -stable.

Proof. Denote by $M_{|G|}$ to M_G with the trivial grading. Because G is countable we have that $M_\infty M_{|G|}$ and M_∞ are isomorphic in $G_{gr}\text{-Alg}$. We also have the following isomorphisms of G -graded algebras

$$\begin{aligned} \eta : M_G M_G &\rightarrow M_G M_{|G|} & e_{s,g} \otimes e_{t,r} &\mapsto e_{st,gr} \otimes e_{t,r} \\ \mu : M_G M_{|G|} &\rightarrow M_{|G|} M_G & e_{s,g} \otimes e_{t,r} &\mapsto e_{t,r} \otimes e_{s,g} \end{aligned}$$

Consider the following commutative diagram

$$\begin{array}{ccc} & M_G M_G A & \xrightarrow{\eta} & M_G M_{|G|} A \\ \text{id}_{M_G} \otimes \iota_A \nearrow & & & \searrow \mu \\ M_G A & & & M_{|G|} M_G A \\ \iota_\infty \searrow & & & \swarrow \iota_\infty \\ & M_\infty M_G A & \xrightarrow{\simeq} & M_\infty M_{|G|} M_G A \end{array}$$

It follows that $\hat{F}(\iota_A) = F(\text{id}_{M_G} \otimes \iota_A)$ is an isomorphism in \mathcal{D} because F is M_∞ -stable. \square

4. EQUIVARIANT ALGEBRAIC KK-THEORY

From now to the rest of the paper we suppose G is a countable group.

4.1. The category $\mathfrak{K}\mathfrak{K}^G$. In this section we introduce an equivariant algebraic kk-theory for G -algebras. Let A, B be G -algebras, we define

$$\text{kk}^G(A, B) := \text{kk}_{G\text{-Alg}}(M_G \otimes A, M_G \otimes B).$$

Consider the category $\mathfrak{K}\mathfrak{K}^G$ whose objects are the G -algebras and where the morphisms between A and B are the elements of $\text{kk}^G(A, B)$. Let $j^G : G\text{-Alg} \rightarrow \mathfrak{K}\mathfrak{K}^G$ be the functor defined as the identity on objects and which sends each morphism of G -algebras $f : A \rightarrow B$ to its class $[\text{id}_{M_G} \otimes f] \in \text{kk}^G(A, B)$. The composition law in $\mathfrak{K}\mathfrak{K}^G$ is the same that in (2.6.1) taking $M_G \otimes A, M_G \otimes B$ and $M_G \otimes C$ instead of A, B and C . As in Section 2.6 we have an equivalence $\Omega : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}^G$ and distinguished triangles

$$\Omega C \rightarrow A \rightarrow B \rightarrow C$$

in $\mathfrak{K}\mathfrak{K}^G$ which gives to $\mathfrak{K}\mathfrak{K}^G$ a triangulated category structure.

Theorem 4.1.1. *The functor $j^G : G\text{-Alg} \rightarrow \mathfrak{K}\mathfrak{K}^G$ is an excisive, equivariantly homotopy invariant, and G -stable functor. Moreover, it is the universal functor for these properties. In other words, if \mathcal{T} is a triangulated category and $R : G\text{-Alg} \rightarrow \mathcal{T}$ together a class of morphisms $\{\bar{\partial}_E : E \in \mathcal{E}\}$ is an excisive, equivariantly homotopy invariant and G -stable functor, then there exists a unique triangle functor $\bar{R} : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathcal{T}$ such that the following diagram commutes*

$$\begin{array}{ccc} G\text{-Alg} & \xrightarrow{j^G} & \mathfrak{K}\mathfrak{K}^G \\ & \searrow R & \downarrow \bar{R} \\ & & \mathcal{T} \end{array}$$

Proof. Let E be a weakly split extension. Define

$$\partial_E^G \in \text{hom}_{\mathfrak{K}\mathfrak{K}^G}(\Omega C, A) = \text{hom}_{\mathfrak{K}\mathfrak{K}_{G\text{-Alg}}}(\Omega M_G \otimes C, M_G \otimes A)$$

as the morphism $\partial_{E'}$ defined in (2.6.4) associated to the following weakly split extension

$$M_G \otimes A \rightarrow M_G \otimes B \rightarrow M_G \otimes C \quad (E')$$

By theorem 2.6.5 and proposition 3.1.9 the functor $j^G : G\text{-Alg} \rightarrow \mathfrak{K}\mathfrak{K}^G$ with the family $\{\partial_E^G : E \in \mathcal{E}\}$ is an excisive, homotopy invariant and G -stable functor. Let us check it is universal for these properties. Let $X : G\text{-Alg} \rightarrow \mathcal{T}$ be a functor which has the mentioned properties with a family $\{\bar{\partial}_E : E \in \mathcal{E}\}$. By theorem 2.6.5 there exists a unique triangle functor $\bar{X} : \mathfrak{K}\mathfrak{K}_{G\text{-Alg}} \rightarrow \mathcal{T}$ such that the following diagram

commutes

$$(4.1.2) \quad \begin{array}{ccc} G\text{-Alg} & \xrightarrow{j^G} & \mathfrak{K}\mathfrak{K}^G \\ & \searrow^{j_{G\text{-Alg}}} & \nearrow \\ & \mathfrak{K}\mathfrak{K}_{G\text{-Alg}} & \\ & \downarrow \overline{X} & \\ & \mathcal{T} & \end{array}$$

X X'

We will define $X' : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathcal{T}$. We know that $X' = X$ on objects. As X is G -stable the following morphisms are a zig-zag between A and $M_G \otimes A$ by X , (see Example 3.1.8)

$$(4.1.3) \quad A \xrightarrow{\iota_A} M_{G \sqcup \{*\}} \otimes A \xleftarrow{\iota'_A} M_G \otimes A$$

Let $\alpha \in \text{kk}^G(A, B)$ and define

$$X'(\alpha) := X(\iota_B)^{-1} X(\iota'_B) \overline{X}(\alpha) X(\iota'_A)^{-1} X(\iota_A).$$

Note this definition is the unique possibility to make the diagram (4.1.2) commutative. \square

4.2. The category $\hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$. In this section we introduce the equivariant algebraic kk -theory for G -graded algebras. Let A, B be G -graded algebras. We define

$$\hat{\text{kk}}^G(A, B) := \text{kk}_{G_{gr}\text{-Alg}}(M_G A, M_G B).$$

Consider the category $\hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$ whose objects are the G -graded algebras and the morphisms between A and B are the elements of $\hat{\text{kk}}^G(A, B)$. Let $j^{G_{gr}} : G_{gr}\text{-Alg} \rightarrow \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$ be the functor defined as the identity on objects and which sends each morphism of G -graded algebras $f : A \rightarrow B$ to its class $[\text{id}_{M_G} f] \in \hat{\text{kk}}^G(A, B)$. As in Section 4.1 we can consider a triangulated category structure on $\hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$.

Theorem 4.2.1. *The functor $j^{G_{gr}} : G_{gr}\text{-Alg} \rightarrow \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$ is an excisive, graded homotopy invariant, and G -graded stable functor. Moreover, it is the universal functor for these properties. In other words, if \mathcal{T} is a triangulated category and $R : G_{gr}\text{-Alg} \rightarrow \mathcal{T}$ together a class of morphisms $\{\overline{\partial}_E : E \in \mathcal{E}\}$ is an excisive, graded homotopy invariant and G -graded stable functor, then there exists a unique triangle functor $\overline{R} : \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G \rightarrow \mathcal{T}$ such that the following diagram commutes*

$$\begin{array}{ccc} G_{gr}\text{-Alg} & \xrightarrow{j^{G_{gr}}} & \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G \\ & \searrow R & \downarrow \overline{R} \\ & & \mathcal{T} \end{array}$$

Proof. The proof is similar to Theorem 4.1.1. \square

5. ALGEBRAIC GREEN-JULG THEOREM

5.1. Crossed product and trivial action. Let A be an ℓ -algebra. Write A^τ for A with the trivial action of G . This gives us a functor $\tau : \text{Alg} \rightarrow G\text{-Alg}$. It is easy to check that $j^G \circ \tau$ satisfies excision, is homotopy invariant and is M_∞ -stable. Write $\mathfrak{K}\mathfrak{K} = \mathfrak{K}\mathfrak{K}_{\text{Alg}}$. By Theorem 2.6.5 there exists a unique functor $\tau : \mathfrak{K}\mathfrak{K} \rightarrow \mathfrak{K}\mathfrak{K}^G$ such that the following diagram is commutative

$$\begin{array}{ccc} \text{Alg} & \xrightarrow{\tau} & G\text{-Alg} \\ j \downarrow & & \downarrow j^G \\ \mathfrak{K}\mathfrak{K} & \xrightarrow{\tau} & \mathfrak{K}\mathfrak{K}^G \end{array}$$

Let A be a G -algebra. The crossed product algebra $A \rtimes G$ is the ℓ -module $A \otimes \ell G$ with the following multiplication

$$(a \rtimes g)(b \rtimes h) = a(g \cdot b) \rtimes gh \quad a, b \in A, g, h \in G.$$

Proposition 5.1.1. *Let A be a G -algebra and \mathcal{W} a G -module by locally finite automorphisms. The following algebras are naturally isomorphic*

$$(A \rtimes G) \otimes \text{End}_\ell^F(\mathcal{W}) \simeq (A \otimes \text{End}_\ell^F(\mathcal{W})) \rtimes G.$$

Proof. Let $\rho : G \rightarrow (\text{End}_\ell^K(\mathcal{W}))^\times$ be the structure map. Note that the homomorphisms

$$\begin{aligned} \phi : (A \rtimes G) \otimes \text{End}_\ell^F(\mathcal{W}) &\rightarrow (A \otimes \text{End}_\ell^F(\mathcal{W})) \rtimes G & \phi(a \rtimes g \otimes \varphi) &= a \otimes \varphi \rho(g^{-1}) \rtimes g \\ \psi : (A \otimes \text{End}_\ell^F(\mathcal{W})) \rtimes G &\rightarrow (A \rtimes G) \otimes \text{End}_\ell^F(\mathcal{W}) & \psi(a \otimes \varphi \rtimes g) &= a \rtimes g \otimes \varphi \rho(g) \end{aligned}$$

are inverse of each other. \square

Proposition 5.1.2. *There exists a unique functor $\rtimes G : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}$ such that the following diagram is commutative*

$$\begin{array}{ccc} G\text{-Alg} & \xrightarrow{\rtimes G} & \text{Alg} \\ j^G \downarrow & & \downarrow j \\ \mathfrak{K}\mathfrak{K}^G & \xrightarrow{\rtimes G} & \mathfrak{K}\mathfrak{K} \end{array}$$

Proof. We shall show $j(- \rtimes G)$ is excisive, homotopy invariant and G -stable. Because $\rtimes G$ maps split exact sequences to split exact sequences and j is excisive, then $j(- \rtimes G)$ is excisive.

That $j(- \rtimes G)$ is homotopy invariant follows from the fact that

$$A[t] \rtimes G = (A \rtimes G)[t].$$

Let $(\mathcal{W}_1, B_1), (\mathcal{W}_2, B_2)$ be G -modules by locally finite automorphisms and A a G -algebra. Consider the isomorphism ψ defined in Proposition 5.1.1. Note that the following diagram is commutative

$$\begin{array}{ccc} (A \otimes \text{End}_\ell^F(\mathcal{W}_1)) \rtimes G & \xrightarrow{(1 \otimes \bar{i}) \rtimes G} & (A \otimes \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2)) \rtimes G \\ \psi \downarrow & & \downarrow \psi \\ (A \rtimes G) \otimes \text{End}_\ell^F(\mathcal{W}_1) & \xrightarrow{(1 \rtimes G) \otimes \bar{i}} & (A \rtimes G) \otimes \text{End}_\ell^F(\mathcal{W}_1 \oplus \mathcal{W}_2) \end{array}$$

Because j is M_∞ -stable, $j((1 \times G) \otimes \tilde{i})$ is an isomorphism. Hence $j(- \times G)(1 \otimes \tilde{i})$ is an isomorphism by the diagram above. \square

Remark 5.1.3. Let $[\iota_\infty \alpha] \in \text{kk}^G(A, B)$ be an element represented by $\alpha : J^n(M_G A) \rightarrow (M_G B)^{\text{sd}^p \mathcal{S}^n}$ which is a morphism in $[J^n(M_G A), \mathcal{M}_\infty(M_G B)^{\text{sd}^p \mathcal{S}^n}]$. Consider the classifying map

$$J^n(M_G A \rtimes G) \rightarrow J^n(M_G A) \rtimes G$$

The element $[\alpha] \rtimes G$ is represented by the following composition

$$J^n(M_G A \rtimes G) \rightarrow J^n(M_G A) \rtimes G \xrightarrow{\alpha \rtimes G} (M_G B)^{\text{sd}^p \mathcal{S}^n} \rtimes G.$$

Proposition 5.1.4. *The functor $\rtimes G : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}$ is a triangle functor.*

Proof. A distinguished triangle in $\mathfrak{K}\mathfrak{K}^G$ is a diagram isomorphic to

$$\Omega B \xrightarrow{j^G(\iota)} P_f \xrightarrow{j^G(\pi_f)} A \xrightarrow{j^G(f)} B$$

for some morphism of G -algebras $f : A \rightarrow B$. That means, if it is isomorphic to

$$(5.1.5) \quad M_G \Omega B \xrightarrow{j_{G\text{-Alg}}(\iota)} M_G P_f \xrightarrow{j_{G\text{-Alg}}(\pi_f)} M_G A \xrightarrow{j_{G\text{-Alg}}(f)} M_G B$$

in $\mathfrak{K}\mathfrak{K}_{G\text{-Alg}}$. The functor $\rtimes G : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}$ sends the triangle (5.1.5) to

$$(5.1.6) \quad (M_G \Omega B) \rtimes G \rightarrow (M_G P_f) \rtimes G \rightarrow (M_G A) \rtimes G \rightarrow (M_G B) \rtimes G$$

which by Proposition 5.1.1 and M_∞ -stability is isomorphic to

$$(5.1.7) \quad \Omega(B \rtimes G) \rightarrow P_f \rtimes G \rightarrow A \rtimes G \rightarrow B \rtimes G$$

As $P_f \rtimes G \simeq P_{f \rtimes G}$, (5.1.7) is a distinguished triangle in $\mathfrak{K}\mathfrak{K}$. \square

5.2. Green-Julg Theorem for $\mathfrak{K}\mathfrak{K}^G$. In this section we shall see an algebraic version of the GREEN-JULG THEOREM, see [11] and [6] for versions of this result in Kasparov KK-theory and E -theory setting.

Theorem 5.2.1. *Let G be a finite group of n elements and $1/n \in \ell$. The functors $\tau : \mathfrak{K}\mathfrak{K} \rightarrow \mathfrak{K}\mathfrak{K}^G$ and $\rtimes G : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}$ are adjoint functors. Hence*

$$\text{kk}^G(A^\tau, B) \simeq \text{kk}(A, B \rtimes G) \quad A \in \text{Alg} \quad B \in G\text{-Alg}.$$

Proof. By [10, Theorem 2, pag 81], it is enough to prove that there exist natural transformations $\bar{\alpha}_A \in \text{kk}(A, A^\tau \rtimes G)$ and $\bar{\beta}_B \in \text{kk}^G((B \rtimes G)^\tau, B)$ such that the following compositions

$$A^\tau \xrightarrow{\tau(\bar{\alpha}_A)} (A^\tau \rtimes G)^\tau \xrightarrow{\bar{\beta}_{\tau(A)}} A^\tau \quad B \rtimes G \xrightarrow{\bar{\alpha}_{B \rtimes G}} (B \rtimes G)^\tau \rtimes G \xrightarrow{\bar{\beta}_{B \rtimes G}} B \rtimes G$$

are the identities in $\text{kk}^G(A^\tau, A^\tau)$ and $\text{kk}(B \rtimes G, B \rtimes G)$ respectively.

Put $\epsilon = 1/n \sum_{g \in G} g$ in ℓG and define

$$(5.2.2) \quad \alpha_A : A \rightarrow A^\tau \rtimes G = A \otimes \ell G \quad \alpha(a) = a \otimes \epsilon.$$

Note α_A is an algebra morphism since ϵ is idempotent. Consider the element $\bar{\alpha}_A \in \text{kk}(A, A^\tau \rtimes G)$ represented by α_A . Let

$$\beta_B : (B \rtimes G)^\tau \rightarrow M_G B \quad \beta_B(b \rtimes g) = \sum_{s \in G} s(b) e_{s, sg}$$

One can check that β_B is an equivariant algebra morphism. Let $\bar{\beta}_B \in \text{kk}^G((B \rtimes G)^\tau, B)$ be represented by β_B . The composite $\beta_{\tau(A)} \tau(\alpha_A)$ is $\text{id}_{A^\tau} \otimes \bar{\iota}$ where $\bar{\iota}$ is the map defined in 3.1.11. As j^G is G -stable, $j^G(\text{id}_{A^\tau} \otimes \bar{\iota})$ is the identity in $\text{kk}^G(A^\tau, A^\tau)$.

Let ψ be the morphism defined in the proof of the Proposition 5.1.1. By Remark 5.1.3, $\overline{\beta}_B \rtimes G$ is represented by $\psi \circ (\beta_B \rtimes G)$. We want to prove that $\psi \circ (\beta_B \rtimes G) \circ \alpha_{B \rtimes G}$ represents the identity in $\text{kk}(B \rtimes G, B \rtimes G)$. Note that

$$(\psi \circ (\beta_B \rtimes G) \circ \alpha_{B \rtimes G})(b \rtimes g) = \frac{1}{n} \sum_{h, s \in G} (s(b) \rtimes h) e_{s, h^{-1}sg}.$$

Put $t = h^{-1}sg$ and note

$$(5.2.3) \quad \frac{1}{n} \sum_{h, s \in G} (s(b) \rtimes h) e_{s, h^{-1}sg} = \frac{1}{n} \sum_{t, s \in G} (s(b) \rtimes sgt^{-1}) e_{s, t}$$

and

$$s(b) \rtimes sgt^{-1} = (\mathbf{1} \rtimes s)(b \rtimes g)(\mathbf{1} \rtimes t^{-1}) \quad \text{in } \tilde{B} \rtimes G$$

We can write (5.2.3) as $TA_{b \rtimes g}T^{-1}$, where

$$A_{b \rtimes g} = \frac{1}{n} \sum_{t, s \in G} (b \rtimes g) e_{s, t} \quad T = \sum_{t \in G} (\mathbf{1} \rtimes t) e_{t, t}$$

Because $b \rtimes g \mapsto A_{b \rtimes g}$ represents the identity, the same is true for $b \rtimes g \mapsto TA_{b \rtimes g}T^{-1}$, see [2, Proposition 5.1.2]. \square

Example 5.2.4. We give an example to show that the adjointness between of τ and $\rtimes G$ of Theorem 5.2.1 fails to hold at the algebra level. Let $G = \mathbb{Z}_2 = \{1, \sigma\}$, $A = \ell$ and $B = (\ell G)^*$ the dual algebra of ℓG with the regular action. Note $\text{hom}_{G\text{-Alg}}(A^\tau, B)$ has two elements only:

$$\varphi_i : \ell \rightarrow (\ell G)^* \quad \varphi_0(1) = 0 \quad \varphi_1(1) = \chi_1 + \chi_\sigma$$

One the other hand $\text{hom}_{\text{Alg}}(A, B \rtimes G) = \text{hom}_{\text{Alg}}(\ell, (\ell G)^* \rtimes G)$ has at least as many elements as ℓ . For each $\lambda \in \ell$ we can define

$$\varphi_\lambda : \ell \rightarrow (\ell G)^* \rtimes G \quad \varphi_\lambda(1) = \chi_1 \rtimes 1 + \lambda(\chi_1 \rtimes \sigma) \quad \lambda \in \ell$$

Note φ_λ is an algebra morphism because $\chi_1 \rtimes 1 + \lambda(\chi_1 \rtimes \sigma)$ is an idempotent element.

Write $\psi_{G,J}$ for the isomorphism of the Theorem 5.2.1

$$(5.2.5) \quad \psi_{G,J} : \text{kk}^G(B^\tau, A) \rightarrow \text{kk}(B, A \rtimes G) \quad \psi_{G,J} = \alpha^* \circ \rtimes G$$

where α is the morphism defined in (5.2.2).

Corollary 5.2.6. *Let G be a finite group such that $1/|G| \in \ell$. Let A be a G -algebra, then*

$$\text{kk}^G(\ell, A) \simeq \text{kk}(\ell, A \rtimes G) \simeq \text{KH}(A \rtimes G)$$

\square

6. INDUCTION AND RESTRICTION

In this section we study the adjointness property of the functors of induction and restriction between $\mathfrak{K}\mathfrak{R}^H$ and $\mathfrak{K}\mathfrak{R}^G$ where G is a group and H is a subgroup of G .

Let A be a G -algebra and $H \subset G$ a subgroup. If we restrict the action to H we obtain an H -algebra $\text{Res}_G^H(A)$. It is clear this construction defines a functor

$\text{Res}_G^H : G\text{-Alg} \rightarrow H\text{-Alg}$. It is easily seen that we can extend $\text{Res}_G^H : G\text{-Alg} \rightarrow H\text{-Alg}$ to a triangle functor $\text{Res}_G^H : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}^H$ so that the following diagram commutes

$$\begin{array}{ccc} G\text{-Alg} & \xrightarrow{\text{Res}_G^H} & H\text{-Alg} \\ j^G \downarrow & & \downarrow j^H \\ \mathfrak{K}\mathfrak{K}^G & \xrightarrow{\text{Res}_G^H} & \mathfrak{K}\mathfrak{K}^H \end{array}$$

Let $\pi : G \rightarrow G/H$ the projection and A an H -algebra. Consider

$$A^{(G,H)} := \{f : G \rightarrow A : \#\pi(\text{supp}(f)) < \infty\}$$

and define

$$\text{Ind}_H^G(A) = \{f \in A^{(G,H)} : f(s) = h(f(sh)) \quad \forall s \in G, \quad h \in H\}.$$

One checks that $\text{Ind}_H^G(A)$ is a G -algebra with pointwise multiplication and the following action of G

$$(6.1) \quad (g \cdot f)(s) = f(g^{-1}s) \quad f \in \text{Ind}_H^G(A) \quad g, s \in G.$$

Observe this construction is functorial, if $\varphi : A \rightarrow B$ is a morphism of H -algebras then $\text{Ind}_H^G(\varphi)(f) = \varphi \circ f$.

If $g \in G$, write $\chi_g : G \rightarrow \mathbb{Z}$ for the characteristic function. If $a \in A$ and $g \in G$, define

$$(6.2) \quad \xi_H(g, a) = \sum_{h \in H} \chi_{gh} h^{-1}(a) \quad \xi_H(g, a)(s) = \begin{cases} h^{-1}(a) & s = gh \\ 0 & s \notin gH \end{cases}$$

It is easy to check that these elements belong to $\text{Ind}_H^G(A)$ and every element $\phi \in \text{Ind}_H^G(A)$ can be written as a finite sum

$$(6.3) \quad \phi = \sum_{g \in \mathcal{R}} \xi_H(g, \phi(g))$$

where $r : G/H \rightarrow G$ is a pointed section and $\mathcal{R} = r(G/H)$. Note that we have the following relations

$$(6.4) \quad s \cdot \xi_H(g, a) = \xi_H(sg, a)$$

$$(6.5) \quad \xi_H(g, a)\xi_H(\tilde{g}, \tilde{a}) = \begin{cases} \xi_H(\tilde{g}, \tilde{g}^{-1}g(a)\tilde{a}) & \tilde{g}^{-1}g \in H \\ 0 & \tilde{g}^{-1}g \notin H \end{cases}$$

$$(6.6) \quad \xi_H(g, a) = \xi_H(gh, h^{-1} \cdot a) \quad h \in H$$

Proposition 6.7. *Let A be a G -algebra and B be an H -algebra, then*

$$\text{Ind}_H^G(B \otimes \text{Res}_G^H A) \simeq \text{Ind}_H^G(B) \otimes A$$

Proof. The isomorphisms are given by

$$S : \begin{array}{ccc} \text{Ind}_H^G(B) \otimes A & \rightarrow & \text{Ind}_H^G(B \otimes \text{Res}_G^H A) \\ \xi_H(g, b) \otimes a & \mapsto & \xi_H(g, b \otimes g^{-1} \cdot a) \end{array}$$

$$T : \begin{array}{ccc} \text{Ind}_H^G(B \otimes \text{Res}_G^H A) & \rightarrow & \text{Ind}_H^G(B) \otimes A \\ \xi_H(g, b \otimes a) & \mapsto & \xi_H(g, b) \otimes g \cdot a \end{array}$$

□

Corollary 6.8. *Let A be a G -algebra. Then*

$$\mathrm{Ind}_H^G \mathrm{Res}_G^H A \rightarrow \ell^{(G/H)} \otimes A \quad \xi_H(s, b) \mapsto \chi_{sH} \otimes s \cdot b$$

is an isomorphism of G -algebras.

□

Proposition 6.9. *Let $\mathrm{Ind} : H\text{-Alg} \rightarrow G\text{-Alg}$ be the following functor*

$$\mathrm{Ind}(A) = \mathrm{Ind}_H^G(M_H \otimes A).$$

There exists a functor $\mathrm{Ind}_H^G : \mathfrak{K}\mathfrak{K}^H \rightarrow \mathfrak{K}\mathfrak{K}^G$ such that the following diagram is commutative

$$(6.10) \quad \begin{array}{ccc} H\text{-Alg} & \xrightarrow{\mathrm{Ind}} & G\text{-Alg} \\ j^H \downarrow & & \downarrow j^G \\ \mathfrak{K}\mathfrak{K}^H & \xrightarrow{\mathrm{Ind}_H^G} & \mathfrak{K}\mathfrak{K}^G \end{array}$$

Proof. Straightforward. □

Proposition 6.11. *The functor $\mathrm{Ind}_H^G : \mathfrak{K}\mathfrak{K}^H \rightarrow \mathfrak{K}\mathfrak{K}^G$ is a triangle functor.*

Proof. Let $f : A \rightarrow B$ be a morphism of H -algebras. The following is an isomorphism of G -algebras

$$(6.12) \quad \Theta : \mathrm{Ind}_H^G(P_f) \rightarrow P_{\mathrm{Ind}_H^G(f)} \quad \xi_H(g, (tp(t), a)) \mapsto (\xi_H(g, tp(t)), \xi_H(g, a)).$$

The image of the induction functor applied to the path extension of f is

$$(6.13) \quad \mathrm{Ind}_H^G(\Omega B) \xrightarrow{\mathrm{Ind}_H^G \iota} \mathrm{Ind}_H^G(P_f) \xrightarrow{\mathrm{Ind}_H^G \pi_f} \mathrm{Ind}_H^G A \xrightarrow{\mathrm{Ind}_H^G f} \mathrm{Ind}_H^G B$$

By Proposition 6.7 and (6.12), the extension (6.13) is isomorphic to

$$\Omega \mathrm{Ind}_H^G(B) \xrightarrow{\mathrm{Ind}_H^G \iota} P_{\mathrm{Ind}_H^G f} \xrightarrow{\pi_{\mathrm{Ind}_H^G f}} \mathrm{Ind}_H^G A \xrightarrow{\mathrm{Ind}_H^G f} \mathrm{Ind}_H^G B.$$

Then $\mathrm{Ind}_H^G : \mathfrak{K}\mathfrak{K}^H \rightarrow \mathfrak{K}\mathfrak{K}^G$ is a triangle functor. □

Theorem 6.14. *Let G be a group and H a subgroup of G . Then the functors*

$$\mathrm{Ind}_H^G : \mathfrak{K}\mathfrak{K}^H \rightarrow \mathfrak{K}\mathfrak{K}^G \quad \mathrm{Res}_G^H : \mathfrak{K}\mathfrak{K}^G \rightarrow \mathfrak{K}\mathfrak{K}^H$$

are adjoint. Hence

$$\mathrm{kk}^G(\mathrm{Ind}_H^G(B), A) \simeq \mathrm{kk}^H(B, \mathrm{Res}_G^H(A)) \quad \forall B \in H\text{-Alg} \quad A \in G\text{-Alg}$$

Proof. Let $A \in G\text{-Alg}$ and $B \in H\text{-Alg}$. We need natural transformations

$$\alpha_A \in \mathrm{kk}^G(\mathrm{Ind}_H^G \mathrm{Res}_G^H A, A) \quad \beta_B \in \mathrm{kk}^H(B, \mathrm{Res}_G^H \mathrm{Ind}_H^G B)$$

which verify the unit and counit condition, respectively.

Define $\varphi_A : \mathrm{Ind}_H^G(\mathrm{Res}_G^H(A)) \rightarrow M_{G/H} \otimes A$ such that

$$\varphi_A(\xi_H(s, b)) = e_{sH, sH} \otimes s \cdot b.$$

One checks that φ_A is a G -equivariant algebra morphism. Put

$$(6.15) \quad \psi_B : B \rightarrow \mathrm{Res}_G^H \mathrm{Ind}_H^G(B) \quad \psi_B(b) = \xi_H(e, b)$$

It is easy to check that ψ_B is well-defined and is a map of H -algebras. Let $\alpha_A \in \mathrm{kk}^G(\mathrm{Ind}_H^G \mathrm{Res}_G^H A, A)$ the element represented by φ_A and $\beta_B \in \mathrm{kk}^H(B, \mathrm{Res}_G^H \mathrm{Ind}_H^G B)$ the element represented by ψ_B .

The composite $\text{Res}_G^H(\alpha_A) \circ \beta_{\text{Res}_G^H A}$ is represented by $\text{Res}_G^H(\varphi_A) \circ \psi_{\text{Res}_G^H A}$ which is kk^H -equivalent to the identity in the sense of remark 2.6.2. The element $\alpha_{\text{Ind}_H^G B} \circ \text{Ind}_H^G(\beta_B) \in \text{kk}^G(\text{Ind}_H^G B, \text{Ind}_H^G B)$ is represented by

$$\gamma : \text{Ind}_H^G B \rightarrow M_{G/H} \otimes \text{Ind}_H^G B \quad \xi_H(g, b) \mapsto e_{gH, gH} \otimes \xi_H(g, b)$$

The following morphism of H -algebras

$$\theta : C \rightarrow M_{G/H} \otimes C \quad \theta(c) = e_{H, H} \otimes c$$

represents to the identity in the sense of remark 2.6.2. Then $\text{Ind}_H^G(\theta)$ is kk^G -equivalent to the identity. It is easy to check $\text{Ind}_H^G(\theta) = \gamma$ with $C = B$. \square

Write ψ_{IR} for the isomorphism

$$(6.16) \quad \psi_{IR} : \text{kk}^G(\text{Ind}_H^G B, A) \rightarrow \text{kk}^H(B, \text{Res}_G^H A) \quad \psi_{IR} = \psi_B^* \circ \text{Res}_G^H$$

where ψ_B is the morphism defined (6.15).

Corollary 6.17. *Let G be a group, H a finite subgroup of G and A a G -algebra then*

$$\text{kk}^G(\ell^{(G/H)}, A) \simeq \text{kk}(\ell, A \rtimes H) \simeq \text{KH}(A \rtimes H)$$

Proof. The isomorphism is the composition of ψ_{GJ} and ψ_{IR} defined in (5.2.5) and in (6.16). \square

7. BAAJ-SKANDALIS DUALITY

In this section we define crossed product functors between the categories $G\text{-Alg}$ and $G_{gr}\text{-Alg}$. We prove that they extend to equivalences between $\mathfrak{K}\mathfrak{K}^G$ and $\hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$. In this way we obtain an algebraic duality theorem similar to the duality given by Baaj-Skandalis in [1].

Let A be a G -algebra. Then

$$A \rtimes G = \bigoplus_{s \in G} A \rtimes s \quad \text{and} \quad (A \rtimes s)(A \rtimes t) \subset A \rtimes st$$

thus $A \rtimes G$ is a G -graded algebra. If $f : A \rightarrow B$ is a homomorphism of G -algebras then $f \rtimes G : A \rtimes G \rightarrow B \rtimes G$ is a graded homomorphism. Hence we have a functor

$$\rtimes G : G\text{-Alg} \rightarrow G_{gr}\text{-Alg}$$

We can also define a functor

$$G \hat{\rtimes} : G_{gr}\text{-Alg} \rightarrow G\text{-Alg}$$

as follows. Let B be a G -graded algebra. Let $G \hat{\rtimes} B$ be the algebra which as a module is $\ell^{(G)} \otimes B$ and the product is the following

$$(7.1) \quad (\chi_g \rtimes a)(\chi_h \rtimes b) := \chi_g \rtimes a_{g^{-1}h} b.$$

Here b_g is the homogeneous element associated to g in the decomposition

$$b = \sum_{g \in G} b_g.$$

One checks that the product (7.1) is associative and the action of G , $s \cdot \chi_g \rtimes a = \chi_{sg} \rtimes a$, makes it into a G -algebra. The crossed product $G \hat{\rtimes} B \rtimes G$ is a G -graded algebra which contains B as a graded subalgebra by

$$(7.2) \quad b \mapsto \sum_{g \in G} \chi_{1_G} \rtimes b_g \rtimes g.$$

In (3.2.1) we defined $M_G B$ a G -graded matrix algebra associated to B . Through the inclusion $b \mapsto e_{1_G, 1_G} \otimes b$ we can see B as a G -graded subalgebra of it. In Proposition 7.4 below, we prove that $G \hat{\rtimes} B \rtimes G$ is isomorphic to $M_G B$ as a G -graded algebras.

If $f : A \rightarrow B$ is a homogeneous homomorphism we define a G -algebra homomorphism $(G \hat{\rtimes} f)$ in the obvious way. Thus we have a functor

$$(7.3) \quad \hat{\rtimes} G : G_{gr}\text{-Alg} \rightarrow G\text{-Alg}.$$

Proposition 7.4. *Let A be a G -algebra and let B be a G -graded algebra.*

a) *There are natural isomorphisms of G -algebras*

$$G \hat{\rtimes} (A \rtimes G) \simeq M_G \otimes A$$

b) *There are natural isomorphisms of G -graded algebras*

$$(G \hat{\rtimes} B) \rtimes G \simeq M_G \otimes B$$

Proof. a) Define $T : G \hat{\rtimes} (A \rtimes G) \rightarrow M_G \otimes A$ as

$$T(\chi_g \rtimes a \rtimes s) = g \cdot a \otimes e_{g, gs}.$$

It is easy to check that T is an equivariant algebra isomorphism with inverse given by

$$S(a \otimes e_{r, t}) := \chi_r \rtimes r^{-1} \cdot a \rtimes r^{-1} t$$

b) Define $T : (G \hat{\rtimes} B) \rtimes G \rightarrow M_G \otimes B$ as

$$T(\chi_h \rtimes b \rtimes s) = \sum_{r \in G} e_{h, s^{-1} hr} \otimes b_r.$$

It is easy to check that T is a graded algebra isomorphism with inverse given by

$$(7.5) \quad S(e_{r, s} \otimes b_q) = \chi_r \rtimes b_q \rtimes rqs^{-1}$$

□

Theorem 7.6. *The functors $\rtimes G$ and $G \hat{\rtimes}$ extend to inverse equivalences*

$$-\rtimes G : \mathfrak{K}\mathfrak{K}^G \longrightarrow \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G \quad G \hat{\rtimes} - : \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G \longrightarrow \mathfrak{K}\mathfrak{K}^G$$

Hence if A and B are G -algebras and C and D are G -graded algebras then

$$\text{kk}^G(A, B) \simeq \hat{\text{kk}}^G(A \rtimes G, B \rtimes G) \quad \text{kk}^G(C, D) \simeq \text{kk}^G(G \hat{\rtimes} C, G \hat{\rtimes} D)$$

Proof. As $\rtimes G$ maps split sequences to split sequences, $j^G(- \rtimes G)$ is excisive. By Proposition 5.1.2 $j^G(- \rtimes G)$ is G -stable and homotopy invariant, whence it extends to $-\rtimes G : \mathfrak{K}\mathfrak{K}^G \rightarrow \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G$ by universality. Similarly, as $G \hat{\rtimes}$ maps split exact sequences to split exact sequences then $j^G(G \hat{\rtimes} -)$ is excisive. Because $G \hat{\rtimes}$ maps graded homotopies to equivariant homotopies and $j^G(G \hat{\rtimes} -)$ is M_∞ -stable, $j^G(G \hat{\rtimes} -)$ extends to $G \hat{\rtimes} - : \hat{\mathfrak{K}}\hat{\mathfrak{K}}^G \rightarrow \mathfrak{K}\mathfrak{K}^G$ by universality. To finish we must show that the maps

$$\text{kk}^G(A, B) \rightarrow \hat{\text{kk}}^G(A \rtimes G, B \rtimes G) \quad \text{and} \quad \text{kk}^G(C, D) \rightarrow \text{kk}^G(G \hat{\rtimes} C, G \hat{\rtimes} D)$$

are isomorphisms. This is true by Proposition 7.4. □

REFERENCES

- [1] S. Baaĵ and G. Skandalis, C^* -algèbres de Hopf et th orie de Kasparov  quivariante (Hopf C^* -algebras and equivariant Kasparov theory). *K-Theory*, 2 (1989), no. 6, 683-721.
- [2] G. Corti nas and A. Thom. Bivariant algebraic K -theory. *J. Reine und Angew. Math.* 610 (2007) 71-123.
- [3] J. Cuntz. A new look at KK -theory. *K-theory* 1 (1987) no. 1, 31-51.
- [4] J. Cuntz. Bivariant K -theory and the Weyl algebra. *K-theory* 35 (2005), no. 1-2, 93-137.
- [5] P.G. Goerss and J.F. Jardine. Simplicial homotopy theory. *Progress in Mathematics*, 174. Birkh user Verlag, Basel, 1999.
- [6] E. Guentner, N. Higson and J Trout. Equivariant E -theory for C^* -algebras. *Mem. Amer. Math. Soc.* 148 (2000), no. 703.
- [7] N. Higson. A characterization of KK -theory. *Pacific J. Math.* 126 (1987), no. 2, 253-276.
- [8] G.G. Kasparov The operator K -functor and extensions of C^* -algebras. *Izv. Akad. Nauk. SSR Ser. Mat.* 44 (1980), no. 3, 571-636, 719.
- [9] B. Keller Derived categories and their uses. *Handbook of algebra*, Vol. 1, 671701, North-Holland, Amsterdam, 1996
- [10] S. Mac Lane. Categories for the working mathematician. *Second edition. Graduate Texts in Mathematics*, 5. Springer-Verlag, New York, 1998.
- [11] R. Meyer. Categorical aspects of bivariant K -theory. *K-theory and noncommutative geometry*, 1-39, *EMS Ser. Congr. Rep., Eur. Math. Soc., Z rich*, 2008.
- [12] C. Weibel. Homotopy Algebraic K -theory. Algebraic K -theory and algebraic number theory (Honolulu, HI, 1987), 461-488, *Contemp. Math.*, 83, Amer. Math. Soc., Providence, RI, 1989.
E-mail address: eugenia@cmat.edu.uy - eugenia.ellis@fu-berlin.de

CMAT, FACULTAD DE CIENCIAS-UDELAR, IGU  4225, 11400 MONTEVIDEO, URUGUAY
 INSTITUT F R MATHEMATIK, FU, ARNIMALLEE 7, RAUM 212, 14195 BERLIN, GERMANY.