# ON THE COHOMOLOGY OF POLYHEDRAL PRODUCTS WITH SPACE PAIRS $(D^1, S^0)$

### LI CAI

ABSTRACT. In this note we prove that  $H(\mathbb{RZ}_K)$ , namely the integer cohomology ring of *polyhedral products* with space pairs  $(D^1, S^0)$ , can also be described explicitly by a multiplicative Hochster's formula.

The objective of this note is to understand  $H(\mathbb{RZ}_K)$ , in a way parallelly to the work related to the cohomology of *moment-angle complexes*  $\mathcal{Z}_K$  (see e.g. [3], [2], [4]) by Buchstaber, Panov and Baskakov.

After some constructions, an explicit formula for  $H(\mathbb{R}\mathcal{Z}_K)$  is obtained in Theorem 1.6, the whole proof of which is completed later after Proposition 2.1. Some applications are discussed in Remark 1.8.

## 1. On the Additive Structure

In this note let  $\mathbb{Z}$  be the ring of integers and let all coefficients be from  $\mathbb{Z}$ .

For a non-negative integer m, we write [m] for the set  $\{1, \dots, m\}$ . A simplicial complex K on [m] means the vertice set of K is identified with [m], such that

(1) there is an unique emptyset which belongs to every simplex of K;

(2) as a subset of [m],  $\sigma$  is a simplex of K implies that any face (subset) of  $\sigma$  is a simplex. For instance, if K contains [m] as a simplex, then every subset  $\sigma = \{i_0, \ldots, i_p\}$  is a p-simplex of K.

The geometric realization of K will be denoted by |K|. By  $\mathbb{R}\mathcal{Z}_K$  we denote a specific polyhedral product (we follow [3] on this notation) defined as the union

(1) 
$$\mathbb{R}\mathcal{Z}_K = \bigcup_{\sigma \in K} \{ (x_1, \cdots, x_m) \in (D^1)^m | x_i \in S^0, \text{ when } i \notin \sigma \},$$

thus  $\mathbb{R}\mathcal{Z}_K$  can be embedded in the product  $(D^1)^m$  in an obvious way.

**Example 1.1.** It is known that if K is a simplicial (n-1)-sphere (a simplicial complex homeomorphic to the (n-1)-sphere), then  $\mathbb{RZ}_K$  is an n-dimensional manifold (see [3, p. 98]). Let K be the pentagon on [5], namely it is a 1-dimensional simplicial complex with

edges  $\{i, i + 1\}$  for i = 1, 2, 3, 4 and  $\{1, 5\}$ . For the five 1-simplices (resp. five 0-simplices) of K, let each 1-simplex (resp. 0-simplex) correspond to the  $2^3$  squares (resp. the  $2^4$  edges)  $(D^1)^2 \times (S^0)^3$  (resp.  $D^1 \times (S^0)^4$ ) and let  $(S^0)^5$  be the  $2^5$  vertices as in definition (1). Then by counting the Euler number we can identify this  $\mathbb{RZ}_K$  as a surface with genus 5.

1.1. A Cell Decomposition. Now we give  $(D^1)^m$  a CW structure by a simplicial decomposition to each component  $D_i^1$  (i = 1, ..., m), which descends to  $\mathbb{R}\mathcal{Z}_K$  as a subcomplex: let  $\underline{01}_i$  be the 1-simplex and  $0_i$ ,  $1_i$  be two 0-simplices of the unit interval  $[0, 1]_i \cong D_i^1$  respectively, then each oriented cell of  $(D^1)^m$  can be written as

$$e_1 \times \cdots \times e_m, \ e_i = 0_i, \ 1_i \text{ or } \underline{01}_i.$$

With the decomposition above, we find

**Lemma 1.2.** A cell  $e = e_1 \times \cdots \times e_m$  of  $(D^1)^m$  belongs to its subspace  $\mathbb{R}\mathcal{Z}_K$  if and only if

$$\sigma_e := \{i | e_i = \underline{01}_i\}$$

is a simplex of K.

In what follows we will abuse the notation  $e_i$ : it may mean a simplex or the corresponding simplicial chain, and we will declare what it means depending on different situations. For a topological space X, we denote S(X) (resp.  $S^*(X)$ ) to be its singular chains (resp. cochains);  $C_e(X)$  (resp.  $C_e^*(X)$ ) will mean the cellular chains (resp. cochains) when X is a CW complex, and C(X) (resp.  $C^*(X)$ ) will be denoted for the simplicial chains (resp. cochains) when X is a simplicial complex. The notation H(X) (resp.  $H^*(X)$ ) means the singular homology (resp. cohomology) of X.

1.2. A Brief Review. Let  $X = \prod_{i=1}^{m} |K_i|$  be a product space by topologized finite simplicial complexes  $K_i$ , note that now each  $C^*(K_i)$  is finitely generated. Let  $(\bigotimes_{i=1}^{m} C^*(K_i); d)$  be the differential graded abelian group with generators

$$e_1^* \otimes \cdots \otimes e_m^*$$

where  $e_i^*$  is the dual of the simplicial chain generated by  $e_i$  in  $K_i$ , and whose differential d satisfies

(2) 
$$d(e_1^* \otimes \cdots \otimes e_m^*) = \sum_{i=1}^m (-1)^{\sum_{j=1}^{i-1} \deg(e_j^*)} e_1^* \otimes \cdots \otimes e_{i-1}^* \otimes de_i^* \otimes e_{i+1}^* \otimes \cdots \otimes e_m^*$$

on each generator (on the right hand side the differential d is the usual simplicial coboundary operator). For instance, assume that i = 1, 2, 3, from  $d0_i^* = -\underline{01}_i^*$ ,  $d1_i^* = \underline{01}_i^*$  and  $d\underline{01}_i^* = 0$ ,

one has

$$\mathbf{d}(\mathbf{0}_1^* \otimes \underline{\mathbf{01}}_2^* \otimes \mathbf{1}_3^*) = -\underline{\mathbf{01}}_1 \otimes \underline{\mathbf{01}}_2 \otimes \mathbf{1}_3^* - \mathbf{0}_1^* \otimes \underline{\mathbf{01}}_2^* \otimes \underline{\mathbf{01}}_3^*.$$

 $C_e^*(X)$  means the set of cellular cochains generated by dual cells

$$(e_1 \times \cdots \times e_m)^*,$$

on which we define a morphism

(3) 
$$f: C^*(K_1) \otimes \cdots \otimes C^*(K_m) \longrightarrow C^*_e(X)$$
$$e_1^* \otimes \cdots \otimes e_m^* \longrightarrow (e_1 \times \cdots \times e_m)^*,$$

which is a 1-1 correspondence by a comparison of basis, thus we can equip  $C_e^*(X)$  with a differential d such that f is a cochain map, and it will be denoted as  $(C_e^*(X); d)$ .

**Proposition 1.3.** Consider  $X = \prod_{i=1}^{m} |K_i|$  as a CW complex with cellular cochains  $(C_e^*(X); d)$ , and assume that A is a subcomplex of X subject to this cell decomposition, then there is an isomprphism

$$H^*(A) \cong H(C^*_e(A); d)$$

preserving degrees. For instance, we have isomorphisms

$$H(\bigoplus_{\sum_{i=1}^{m} p_i = p} C^{p_1}(K_1) \otimes \cdots \otimes C^{p_m}(K_m); d) \xrightarrow{f} H^p(C_e^*(X); d) \cong H^p(X)$$

where  $p_i$  are non-negative integers.

A standard cellular (co)homology argument (see e.g. [6, Theorem 57.1, p. 339]) will provide a proof without involving cup products. We will give full version of this proposition later (i.e. Proposition 2.1), but here the simplicial cochain algebra of  $C^*(D_i^1)$  with simplicial cup products will be presented, because it motivates the algebraic construction later. Henceforth let us work with the following basis

(4) 
$$\{\mathbf{1}_i = \mathbf{1}_i^* + \mathbf{0}_i^*, t_i = \mathbf{1}_i^*, u_i = \underline{\mathbf{0}}_{i_i}\}$$

of  $C^*(D_i^1)$  rather than the original  $\{0_i^*, 1_i^*, \underline{01}_i^*\}$ . By definition (see e.g. [6, p. 292]), we have the following relations on simplicial cup products:

$$\mathbf{1}_i \cup t_i = t_i \cup \mathbf{1}_i, \quad \mathbf{1}_i \cup u_i = u_i \cup \mathbf{1}_i, \quad t_i \cup t_i = t_i, \quad u_i \cup u_i = 0, \quad t_i \cup u_i = 0, \quad u_i \cup t_i = u_i.$$

1.3. An Algebraic Description. In what follows,  $x_{\sigma}$  will mean the monomial  $x_{i_0} \dots x_{i_p}$ when  $\sigma = \{i_0, \dots, i_p\}$  is a subset of [m]. By  $\mathbb{Z}[u_1, \dots, u_m; t_1, \dots, t_m]$  we denote the differential graded free  $\mathbb{Z}$ -algebra with 2m generators, such that

(5) 
$$\deg u_i = 1, \quad \deg t_i = 0; \quad \mathrm{d}u_i = 0, \quad \mathrm{d}t_i = u_i,$$

where the differential d follows the rule (2) on each monomial. Let R be the quotient algebra of  $\mathbb{Z}[u_1, \ldots, u_m; t_1, \ldots, t_m]$  subject to the following relations

(6) 
$$u_i t_i = u_i, \quad t_i u_i = 0, \quad u_i t_j = t_j u_i, \quad t_i t_i = t_i, \quad u_i u_i = 0, \quad u_i u_j = -u_j u_i, \quad t_i t_j = t_j t_i,$$

for i, j = 1, ..., m such that  $i \neq j$ , in which the Stanley-Reisner ideal  $\mathcal{I}_K$  is defined to be generated by all square-free monomials  $u_{\sigma}$  such that  $\sigma$  is not a simplex of K.

One can check that the relations (6) listed above are compatible with the differential d, for instance,

$$\mathbf{d}(t_i t_i) = u_i t_i + t_i u_t = u_i t_i = \mathbf{d} t_i$$

**Theorem 1.4.** There is a (degree-preserving) ring isomorphism

(7) 
$$H(R/\mathcal{I}_K; \mathbf{d}) \cong H(\mathbb{R}\mathcal{Z}_K).$$

*Proof.* Here we only prove the additive part of this theorem, and the whole proof involving products will be completed later after Proposition 2.1.

We start from a cochain map, which is generated by mapping each square-free monomial to a tensor product of simplicial cochains, namely

$$\theta: R \longrightarrow C^*(D_i^1) \otimes \cdots \otimes C^*(D_m^1)$$
$$x_1 \dots x_m \longrightarrow e_1^* \otimes \cdots \otimes e_m^*,$$

in which (we are using the basis (4))

$$e_i^* = \begin{cases} \mathbf{1}_i, & \text{if } x_i = 1, \\ t_i, & \text{if } x_i = t_i, \\ u_i, & \text{if } x_i = u_i. \end{cases}$$

The additive generators of R are square-free monomials follows from the relations (6), and by definition  $\theta$  induces an isomorphism between abelian groups such that  $d \circ \theta = \theta \circ d$ . Then from the following diagram about cochain maps (in which q is the quotient and i<sup>\*</sup> is induced by the inclusion)

together with Lemma 1.2, we observe that the quotient part in the Stanley-Reisner ideal corresponds to the cells not contained in  $\mathbb{RZ}_K$ , thus by Proposition 1.3, this part of proof is completed.

1.4. A Combinatorial Description (Hochster's Formula). With the notation  $x_{\sigma}$ , we can write each square-free monomial of  $R/\mathcal{I}_K$  in the form

$$u_{\sigma}t_{\tau}, \quad \sigma \cap \tau = \emptyset, \quad \sigma \in K$$

(for instance,  $u_2 t_3 u_4 t_5 t_6 = u_{2,4} t_{3,5,6}$ ).

When  $\omega$  is a subset of [m] (may be empty), we denote  $K_{\omega}$  to be the full subcomplex of K in  $\omega$ , namely it is the subcomplex

$$\{\sigma \cap \omega | \sigma \in K\},\$$

and  $R/\mathcal{I}_K|_{\omega}$  will be denoted as a subgroup of R (as a group) generated by

$$\{u_{\sigma}t_{\tau}|\sigma\in K \ s.t. \ \sigma\sqcup\tau=\omega, \ \sigma\cap\tau=\emptyset\}.$$

We observe that  $R/\mathcal{I}_K|_{\omega}$  is closed under the differential d and  $R/\mathcal{I}_K = \bigoplus_{\omega \subset [m]} R/\mathcal{I}_K|_{\omega}$ , therefore

$$H(R/\mathcal{I}_K; \mathbf{d}) \cong \bigoplus_{\omega \subset [m]} H(R/\mathcal{I}_K|_{\omega}; \mathbf{d}).$$

Each  $\omega \subset [m]$  yields a 1-1 cochain map (on the right hand side we use the usual simplicial coboundary operator)

(9) 
$$\lambda_{\omega} : R/\mathcal{I}_{K}|_{\omega} \xrightarrow{\cong} C^{*}(K_{\omega})$$
$$u_{\sigma}t_{\tau} \longrightarrow \sigma^{*},$$

which induces an additive isomorphism of cohomology

$$H^p(R/\mathcal{I}_K|_{\omega}) \xrightarrow{\lambda_{\omega}} \widetilde{H}^{p-1}(K_{\omega}),$$

because

$$d(t_1 \dots t_m) = \sum_{i=1}^m t_1 \dots t_{i-1} u_i t_{i+1} \dots t_m$$

behaves as the augmentation and  $\lambda_{\emptyset}^*$  maps  $H^0(R)$  isomorphically onto  $\widetilde{H}^{-1}(\emptyset)$ .

**Remark 1.5.** Indeed,  $\lambda$  induces cochain maps since every time we jump over an index in  $\sigma$ , we will get a -1. Here the construction of  $\lambda$  is motivated by Baskakov's idea in [1].

Therefore we can equip  $\bigoplus_{\omega \subset [m]} \widetilde{H}^{p-1}(K_{\omega})$  with a product structure described by simplicial cochains, such that cochain map  $\lambda$  induces an isomorphism between cochain algebras. Together with Theorem 1.4, we have the following theorem, the full proof of which will be completed later.

**Theorem 1.6.** There are isomorphisms

$$\bigoplus_{\omega \subset [m]} \widetilde{H}^{p-1}(K_{\omega}) \xrightarrow{\cong} H^p(R/\mathcal{I}_K) \xrightarrow{\cong} H^p(\mathbb{R}\mathcal{Z}_K)$$

for all positive integers p. Moreover, after endowing  $\bigoplus_{\omega \subset [m]} \widetilde{H}^{p-1}(K_{\omega})$  the product structure induced by  $\lambda$ , they are isomorphisms between  $\mathbb{Z}$ -algebras.

Thus the cohomology ring of  $\mathbb{R}\mathcal{Z}_K$  will be clear once we write down all cochains representing the (reduced) simplicial cohomology for each  $K_{\omega}$ .

**Example 1.7.** Assume that K is the pentagon described in Example 1.1, let us compute the cohomology of  $\mathbb{RZ}_K$ . First we can write down all the full subcomplexes of K with non-trivial cohomology, namely

$$1: H^{-1}(K_{\emptyset})|1;$$

$$\alpha_{1} - \alpha_{5}: \widetilde{H}^{0}(K_{1,3})|u_{1}t_{3}, \widetilde{H}^{0}(K_{1,4})|u_{1}t_{4}, \widetilde{H}^{0}(K_{2,4})|u_{2}t_{4}, \widetilde{H}^{0}(K_{2,5})|u_{2}t_{5}, \widetilde{H}^{0}(K_{3,5})|u_{3}t_{5};$$

$$\beta_{1} - \beta_{5}: \widetilde{H}^{0}(K_{1,3,4})|u_{1}t_{3,4}, \widetilde{H}^{0}(K_{2,4,5})|u_{2}t_{4,5}, \widetilde{H}^{0}(K_{3,5,1})|u_{3}t_{1,5}, \widetilde{H}^{0}(K_{4,1,2})|u_{4}t_{1,2}, \widetilde{H}^{0}(K_{5,2,3})|u_{5}t_{2,3};$$

$$\gamma: \widetilde{H}^{1}(K_{1,2,3,4,5})|u_{1,2}t_{3,4,5},$$

in which we write a generator for each group, via  $\lambda$ . Thus (by Theorem 1.6)  $H(\mathbb{RZ}_K)$  has ten generators in degree 1 (i.e.  $\alpha_i$ ,  $\beta_i$ , i = 1, ..., 5), and one generator in degree 0 (i.e. 1 the unit) and degree 2 (i.e.  $\gamma$ ) respectively. Note that these generators may not be unique: for instance,

 $d(u_1t_{2,3,4,5}) = -u_{1,2}t_{3,4,5} - u_{1,5}t_{2,3,4} + \text{terms equivalent to } 0,$ 

and in the same way  $-u_1t_{3,4}$  is equivalent to  $u_3t_{1,4} + u_4t_{1,3}$ . A little computation shows, up to equivalences in cohomology, all non-trivial cup products are listed as follows (where we omit the notation  $\cup$ )

$$\gamma = \alpha_1 \beta_2 = \alpha_2 \beta_5 = \alpha_3 \beta_3 = \alpha_4 \beta_1 = \alpha_5 \beta_4 = \beta_1 \beta_2 = \beta_2 \beta_3 = \beta_3 \beta_4 = \beta_4 \beta_5 = \beta_5 \beta_1.$$

We can use the construction (8) in the proof of Theorem 1.4 to identify all corresponding cellular cochains.

**Remark 1.8.** From Theorem 1.6, after a comparison with the result in [3], it follows that there is an additive isomprphism between  $H(\mathcal{Z}_K)$  and  $H(\mathbb{R}\mathcal{Z}_K)$  if we forget about degrees. For instance, we can also find arbitrary torsions in  $H(\mathbb{R}\mathcal{Z}_K)$  when K are special simplicial spheres (see [5, Theorem 11.11]). As algebras, the difference between  $H(\mathcal{Z}_K)$  and  $H(\mathbb{R}\mathcal{Z}_K)$ comes from defferent sign rules and square rules.

## 2. On the Cup Product

In this section we will describe cup products in Proposition 1.3, and then finish the proofs of Theorem 1.4 and Theorem 1.6.

Assume that X and Y are two topological spaces, we define a map  $g: S^*(X) \otimes S^*(Y) \to$ Hom $(S(X) \otimes S(Y), \mathbb{Z})$  generated by

(10) 
$$g(c_X^* \otimes c_Y^*)(c_X' \otimes c_Y') = c_X^*(c_X')c_Y^*(c_Y'), \quad c_X^* \in S^*(X), \quad c_Y^* \in S^*(Y),$$

(the evaluation is trivial when degrees do not match) for each singular trangles  $c'_X$  and  $c'_Y$ .

By Eilenberg-Zilber theorem, there is a natural chain equivalence  $\mu$  between singular chain complexes (unique up to chain homotopy, see [8]; the defferential on tensors follows the rule (2))

$$S(X \times Y) \xrightarrow{\mu_{X \times Y}} S(X) \otimes S(Y),$$

and the cup product is defined via the following

$$S^*(X) \otimes S^*(X) \xrightarrow{\mu^*_{X \times X} \circ g} S^*(X \times X) \xrightarrow{d^*} S^*(X)$$

in which  $d^*$  is induced by the diagonal map  $X \to X \times X$  (the *cross product* is defined by  $\mu^*_{X \times X} \circ g$ ).

From now on let  $\text{Top}^m$  be the category with *m*-products of toplogical spaces as objects, whose morphisms are *m*-products of continuous maps, and let  $\mathcal{C}$  be the category of chain complexes equiped with chain maps.

Consider  $X = \prod_{i=1}^{m} |K_i|$  as an object of Top<sup>*m*</sup>: a product by finite topologized simplicial complexes.

Thus in the following diagram, all morphisms can be considered as natural transformations between functors from Top<sup>m</sup> to C, where  $d_i$  is the diagonal chain map from  $S(|K_i|)$  to  $S(|K_i|) \otimes S(|K_i|)$ , and  $\tilde{\mu}$  is defined by using  $\mu$  for m-1 times, factoring out one component each time:

(11)  

$$S(\prod_{i=1}^{m} |K_{i}|) \otimes S(\prod_{i=1}^{m} |K_{i}|) \stackrel{\widetilde{\mu}_{X} \otimes \widetilde{\mu}_{X}}{\longrightarrow} \otimes_{i=1}^{m} S(|K_{i}|) \otimes \otimes_{i=1}^{m} S(|K_{i}|)$$

$$S(\prod_{i=1}^{m} |K_{i}|) \stackrel{\widetilde{\mu}_{X}}{\longrightarrow} S(|K_{1}|) \otimes \cdots \otimes S(|K_{m}|) \stackrel{\otimes_{i=1}^{m} \mu_{|K_{i}| \times |K_{i}|} \circ d_{i}}{\longrightarrow} \bigotimes_{i=1}^{m} S(|K_{i}|) \otimes S(|K_{i}|),$$

in which T is defined as follows. For each tensor by singular triangles  $\otimes_i c_i \otimes c'_i \in \bigotimes_{i=1}^m S(|K_i|) \otimes S(|K_i|)$ ,

$$T(c_1 \otimes c'_1 \otimes \cdots \otimes c_m \otimes c'_m) = (-1)^{\varepsilon(c,c')} c_1 \otimes \cdots \otimes c_m \otimes c'_1 \otimes \cdots \otimes c'_m$$

where the mod 2 invariant  $\varepsilon(c, c')$  is generated by this rule: each operation of changing the order of any adjacent two terms will produce a multiplication of their degrees, thus we obtain  $\varepsilon(c, c')$  by summing up these multiplications for an arbitrary sequence of operations, for instance one choice for  $\varepsilon(c, c')$  can be

$$\varepsilon(c,c') = \sum_{i=1}^{m} \deg c'_i \sum_{j>i} \deg c_j.$$

It is straightforward to check that T is a chain map. From the method of *acyclic model* with the model category as *m*-products of topologized simplices (see [7], the case for 2-products can be found in [9, p. 252]), it follows that there is a natural chain homotopy between

$$\widetilde{\mu}_X \otimes \widetilde{\mu}_X \circ \mu_{X \times X} \circ d$$
 and  $T \circ (\bigotimes_{i=1}^m \mu_{|K_i| \times |K_i|} \circ d_i) \circ \widetilde{\mu}_X.$ 

Therefore after taking Hom to diagram (11), we obtain the following (where  $g_1, g_2$  are defined from (10))

in which  $\widetilde{T}^*$  is well defined since  $\operatorname{Im}(T^* \circ g_1) \subset \operatorname{Im}(g_2)$ , and here every  $C^*(K_i)$  is considered as a subgroup of  $S^*(|K_i|)$   $(i = 1, \ldots, m)$ , in which only simplicial chains are non-trivially evaluated.

8

**Proposition 2.1.** Assume that  $e^* = \bigotimes_{i=1}^m e_i^*$  and  $e^{*'} = \bigotimes_{i=1}^m e_i^{*'}$  are tensors of dual simplices in  $(\bigotimes_{i=1}^m C^*(K_i) \subset) \bigotimes_{i=1}^m S^*(|K_i|)$ , then as elements in  $S^*(\prod_{i=1}^m |K_i|)$  on both sides, we have

(12) 
$$\widetilde{\mu}_X^* \circ g(\bigotimes_{i=1}^m e_i^*) \cup \widetilde{\mu}_X^* \circ g(\bigotimes_{i=1}^m e_i^{*\prime}) = (-1)^{\varepsilon(e^*, e^{*\prime})} \widetilde{\mu}_X^* \circ g(\bigotimes_{i=1}^m e_i^* \cup e_i^{*\prime}).$$

up to a cochain homotopy. Moreover, consider  $X = \prod_{i=1}^{m} |K_i|$  as a CW complex whose cells are m-products of (topologized) simplices with i-th component in  $K_i$ , and suppose that A is a subcomplex of X with  $e^*$  and  $e^{*'}$  in  $C_e^*(A)$ , then as elements in  $S^*(A)$ , (12) holds up to a cochain homotopy.

*Proof.* After a little chasing on the diagram (11), the first statement follows from

$$\begin{split} \widetilde{\mu}_X^* \circ g(\otimes_{i=1}^m e_i^*) \cup \widetilde{\mu}_X^* \circ g(\otimes_{i=1}^m e_i^{*\prime}) &= d^* \circ \mu_{X \times X}^* \circ \widetilde{\mu}_X^* \otimes \widetilde{\mu}_X^* \circ g_1(\otimes_{i=1}^m e_i^* \otimes \otimes_{i=1}^m e_i^{*\prime}) \\ &= \widetilde{\mu}_X^* \circ (\otimes_{i=1}^m \mu_{|K_i| \times |K_i|} \circ d_i)^* \circ T^* \circ g_1(\otimes_{i=1}^m e_i^* \otimes \otimes_{i=1}^m e_i^{*\prime}) \\ &= (-1)^{\varepsilon(e^*, e^{*\prime})} \widetilde{\mu}_X^* \circ (\otimes_{i=1}^m \mu_{|K_i| \times |K_i|} \circ d_i)^* \circ g_2(\otimes_{i=1}^m e_i^* \otimes e_i^{*\prime}) \\ &= (-1)^{\varepsilon(e^*, e^{*\prime})} \widetilde{\mu}_X^* \circ g(\otimes_{i=1}^m e_i^* \cup e_i^{*\prime}), \end{split}$$

in which from the second line to the third we use the (co)chain homotopy and change  $\widetilde{T}^*$  for the sign  $(-1)^{\varepsilon(e^*,e^{*'})}$ . Now we prove the part about the subcomplex A. Denote all cells of A by  $E_A$ , we can consider  $E_A$  as a subcategory of Top<sup>m</sup> with inclusions as morphisms, then there is a chain equivalence (see e.g. [6, Theorem 31.5], together with the fact that each ein  $E_A$  is a neighborhood deformation retract of  $A = \bigcup_{e \in E_A} e$ )

$$S(A) \cong \operatorname{colim}_{e \in E_A} S(\prod_{i=1}^m e_i)$$

where  $e = \prod_{i=1}^{m} e_i$  with each simplex  $e_i$  in  $K_i$ . Next, from the naturality of  $\mu$  we have the diagram

$$S(X) \qquad \xrightarrow{\mu_X} \qquad \otimes_{i=1}^m S(|K_i|)$$

$$\uparrow \qquad \qquad \uparrow$$

 $S(A) \cong \operatorname{colim}_{e \in E_A} S(\prod_{i=1}^m e_i) \xrightarrow{\widetilde{\mu}_X|_A} \operatorname{colim}_{e \in E_A} \otimes_{i=1}^m S(e_i),$ 

to which after taking Hom, it is not difficult to observe

where  $g_A : C_e^*(A) \to \operatorname{Hom}(\operatorname{colim}_{e \in E_A} \otimes_{i=1}^m S(e_i), \mathbb{Z})$  is defined by evaluating each  $\tilde{e}^* \in C_e^*(A)$ non-trivially on its dual  $\tilde{e} \in \operatorname{colim}_{e \in E_A} \otimes_{i=1}^m S(e_i)$  only. Then we can replace everything of diagram (11) by colimits, and repeat the process for X.

It is not difficult to find the sign rule for commutativity defined by  $\varepsilon$  coincides with the one in  $R/\mathcal{I}_K$ , which is defined in (6). Therefore we have completed all proofs.

#### References

- I. Baskakov, Cohomology of K-powers of spaces and the combinatorics of simplicial divisions, Russian Math. Surveys 57 (2002), no. 5, 9899990.
- [2] I. Baskakov, V. Buchstaber and T. Panov, Algebras of cellular cochains, and torus actions, Russian Math. Surveys 59 (2004), no. 3, 5625563.
- [3] V. Buchstaber and T. Panov, Torus actions and their applications in topology and combinatorics, AMS University Lecture Series, volume 24, (2002).
- [4] T. Panov, Cohomology of face rings and torus actions, arXiv:math.AT/0506526.
- [5] Frédéric Bosio and Laurent Meersseman, Real quadrics in C<sup>n</sup>, complex manifolds and convex polytopes, Acta Mathematica 197 (2006), no. 1, 53-127. MR 2285318
- [6] J. Munkres, *Elements of algebraic topology*, the Benjamin/Cummings Publishing Company, California, 1984.
- [7] S. Eilenberg and S. MacLane, Acyclic models, American Journal of Mathematics, vol. 75 (1953), pp. 189-199.
- [8] , On the groups  $H(\prod, n)$ , I, Annals of Mathematics, vol. 88, No. 1, (1953), pp. 55-106.
- [9] E. Spanier, Algebraic topology, Springer-Verlag, New York, 1966.

GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, 744, MOTOOKA, NISHI-KU, FUKUOKA-CITY 819-0395, JAPAN

E-mail address: l-cai@math.kyushu-u.ac.jp