Harmonic bundles and Toda lattices with opposite sign

Takuro Mochizuki

Abstract

We study a certain type of wild harmonic bundles in relation with a Toda equation. We explain how to obtain a classification of the real valued solutions of the Toda equation in terms of their parabolic weights, from the viewpoint of the Kobayashi-Hitchin correspondence. Then, we study the associated integrable variation of twistor structure. In particular, we give a criterion for the existence of an integral structure. It follows from two results. One is the explicit computation of the Stokes factors of a certain meromorphic flat bundle. The other is an explicit description of the associated meromorphic flat bundle. We use the opposite filtration of the limit mixed twistor structure with an induced torus action.

Keywords: harmonic bundle, Toda lattice with opposite sign, Kobayashi-Hitchin correspondence MSC2010: 53C07, 14H60, 34M40, 32G20

1 Introduction

According to M. Guest and C.-S. Lin [13], the following equation is called the Toda lattice with opposite sign:

$$2\overline{\partial}_z \partial_z w_i - e^{2(w_i - w_{i-1})} + e^{2(w_{i+1} - w_i)} = 0 \quad (i = 1, \dots, r)$$
(1)

Here, we use the convention $w_{r+i} = w_i$ ($\forall i$), and w_i are \mathbb{R} -valued functions on \mathbb{C}^* .

S. Cecotti and C. Vafa found that the equation (1) appeared from the tt^* -equation for several significant models in physics [6], [7]. Moreover, they observed that, if n=2, the equation (1) can be reduced to a Painlevé III equation under some natural requirements. Based on the analysis in [24] and [25], they mathematically verified the existence and uniqueness of the solutions adapted to the conditions required by the models. They found that the solutions have abundant information of the models, and called them magical solutions. Among their rich studies, they proposed a problem to classify the solutions of the equation whose associated $\frac{\infty}{2}$ -variations of Hodge structure have \mathbb{Z} -structure, which are expected to be significant in physics.

As far as the author knows, Guest and Lin are the first mathematicians who systematically studied such issues. They classified the \mathbb{R} -valued global solutions of the equation (1), by imposing an additional symmetry, if it is reduced to the equation for two unknown functions. Moreover, collaborating with A. Its, they studied the Stokes structure of the associated $\frac{\infty}{2}$ -VHS, and they determined when it has a \mathbb{Z} -structure.

In this paper, we study the general case. We shall give a classification of the \mathbb{R} -valued solutions in terms of the parabolic weights, and give a purely algebraic criterion for the existence of \mathbb{Z} -structure.

As is well known, at least implicitly, the equation (1) is closely related to the Hitchin equation for Higgs bundles on curves. Indeed, the solutions of (1) are equivalent to Toda-like harmonic bundles. (See §3.2.2.) Hence, it is quite natural to study the problem from the viewpoint of the Kobayashi-Hitchin correspondence and the asymptotic behaviour of harmonic bundles. We prove that the solutions (1) are classified by the parabolic weights. (See Theorem 3.34 and the remark right after it.) We also describe their asymptotic behaviour and additional symmetry.

Remark 1.1 The harmonic bundles considered in this paper are close to cyclotomic harmonic bundles of C. Simpson in [46]. The author found a more related work due to D. Baraglia [2], who clarified the relation of his cyclic harmonic bundles and affine Toda equations without poles on smooth compact Riemann surfaces. (See also [1].) He has already used efficiently the idea to deduce the symmetry of a harmonic metric from the symmetry of the underlying Higgs bundle.

Remark 1.2 There are many researches on the Toda equation. However, at this moment, the author does not know whether we can deduce the classification of \mathbb{R} -valued solutions (1) from the previous results (for example [47], [48] and [50]). Because our method gives only the existence and uniqueness, it would be significant to obtain explicit descriptions of the solutions by another method.

As for \mathbb{Z} -structure, our argument proceeds as follows. First, we closely study the Stokes structure of a certain meromorphic flat bundle (25). In general, it is rather difficult to understand the Stokes structure of a given meromorphic flat bundle. But, in our case, it is not difficult. The Stokes factors are related to the monodromy in a simple way, and we obtain that the non-zero entries of the Stokes factors are described by the coefficients of the characteristic polynomial of the monodromy. In particular, we obtain that, the meromorphic flat bundle has a \mathbb{Z} -structure, if and only if the coefficients of the characteristic polynomial are integers.

Second, to apply the result in the previous paragraph, we need to know the precise form of the meromorphic flat bundle associated to a Toda-like harmonic bundle. One of the important ingredients is the limit mixed twistor structure associated to the harmonic bundle with homogeneity. It is naturally equipped with a torus action, and hence it comes from a mixed Hodge structure. (The mixed Hodge structure seems to be related with that in [18]. See Remark 5.11.) Once it is obtained, we can apply an argument of M. Saito which is familiar in the construction of Frobenius manifold. ([42]. See [10], [38], [39].) Namely, by using the canonical decomposition of the mixed Hodge structure due to P. Deligne, we can find an opposite filtration to the Hodge filtration. Then, we can extend the associated meromorphic flat bundle to tr-TLE-structure in the sense of C. Hertling [17]. Then, it is not difficult to deduce the more precise form of the connection in the case of Toda-like harmonic bundle by using its high symmetry. Getting together the above two results, we obtain the desired criterion.

It is reasonable to divide this paper into two parts, Part I: the classification of \mathbb{R} -valued solutions (§2–3), and Part II: the study on the Stokes structure of the associated $\frac{\infty}{2}$ -VHS (§4–6). They are rather independent, although related, and we use rather different techniques.

In §2, we give a review of a general theory of filtered Higgs bundles and wild harmonic bundles on curves for the convenience of the readers. In §3, we apply it to a class of filtered Higgs bundles and harmonic bundles on $(\mathbb{P}^1, \{0, \infty\})$, and we explain how to deduce a classification of the \mathbb{R} -valued solutions of the Toda lattice with opposite sign by parabolic weights.

In §4, after a review of \mathbb{Z} -structure of a meromorphic flat bundle possibly with irregular singularity of pure slope, we will study the Stokes structure of a certain meromorphic flat bundle (25). In particular, we give a criterion for the existence of a \mathbb{Z} -structure on the meromorphic flat bundle or its pull back by a ramified covering. In §5, we describe a rather general theory of harmonic bundle with homogeneity and the associated integrable variation of twistor structure, which can be regarded as a variant of [17], [31], [41] and [45]. We would like to give details on the role of Euler field in our situation. We explain how a torus action is induced on the limit mixed twistor structure in some situation, and how we obtain a tr-TLE-structure. In §6, we apply the results in §4 and §5 to Toda-like harmonic bundles. In §6.1, we give a complement on the tr-TLE structure associated to a Toda-like harmonic bundle, in particular, around the irregular singular point ∞ . We use it to obtain a precise description of the associated meromorphic flat bundle. Then, we obtain in §6.2 the criterion when they have \mathbb{Z} -structure.

Acknowledgement This study is inspired by the intriguing works [6], [7], [8], [13], [14] and [15]. I am heartily grateful to Martin Guest for his detailed and useful comments on the earlier version of this manuscript. I also thank Hiroshi Iritani who attracted my attention to the real structure of a quantum D-module. I am grateful to Akira Ishii and Yoshifumi Tsuchimoto for their constant encouragements. I thank Claude Sabbah for numerous discussion and his kindness for years. A part of this paper is based on my talk in the conference "Various Aspects on the Painlevé Equations" held in November 2012. It is my great pleasure to express my gratitude to the organizers. This study is supported by Grant-in-Aid for Scientific Research (C) 10315971.

Part I

2 Preliminary

2.1 Filtered Higgs bundle on a curve

2.1.1 Filtered Higgs bundle on a disc with a marked point

We give a review of filtered Higgs bundle on curves. We shall explain it in the case that X is a disc $\{z \in \mathbb{C} \mid |z| < 1\}$ with a point $D = \{0\}$.

Filtered bundle A filtered bundle on (X, D) is a locally free $\mathcal{O}_X(*D)$ -module \mathcal{V} of finite rank with an increasing filtration by locally free \mathcal{O}_X -submodules $\mathcal{P}_a\mathcal{V}\subset\mathcal{V}$ $(a\in\mathbb{R})$ satisfying the following conditions, (i) $\mathcal{P}_a\mathcal{V}_{|X\setminus D}=\mathcal{V}_{|X\setminus D}$, (ii) $\mathcal{P}_a\mathcal{V}=\bigcap_{b>a}\mathcal{P}_b\mathcal{V}$, (iii) $\mathcal{P}_{a-1}\mathcal{V}=\mathcal{P}_a\mathcal{V}\otimes\mathcal{O}(-D)=z\mathcal{P}_a\mathcal{V}$. Let $\mathcal{P}_*\mathcal{V}$ denote the $\mathcal{O}_X(*D)$ -module \mathcal{V} with the filtration $\{\mathcal{P}_a\mathcal{V}\mid a\in\mathbb{R}\}$, and we call it a filtered bundle on (X,D). We say that \mathcal{V} is the underlying $\mathcal{O}_X(*D)$ -module of $\mathcal{P}_*\mathcal{V}$, and that $\mathcal{P}_*\mathcal{V}$ is a filtered bundle over the $\mathcal{O}_X(*D)$ -module \mathcal{V} . The filtration is also denoted by $\mathcal{P}_*\mathcal{V}$.

We set $\mathcal{P}_{< a}\mathcal{V} := \bigcup_{c < a} \mathcal{P}_c \mathcal{V}$ and $\operatorname{Gr}_a^{\mathcal{P}}(\mathcal{V}) := \mathcal{P}_a \mathcal{V}/\mathcal{P}_{< a}\mathcal{V}$. It is naturally regarded as a finite dimensional vector space over \mathbb{C} . We set $\mathcal{P}ar(\mathcal{P}_*\mathcal{V}) := \{a \in \mathbb{R} \mid \operatorname{Gr}_a^{\mathcal{P}}(\mathcal{V}) \neq 0\}$ which is discrete in \mathbb{R} . If $a \in \mathcal{P}ar(\mathcal{P}_*\mathcal{V})$, then $a + \mathbb{Z} \subset \mathcal{P}ar(\mathcal{P}_*\mathcal{V})$.

The sheaf $\mathcal{O}_X(*D)$ is naturally a filtered bundle, i.e., $\mathcal{P}_a\mathcal{O}_X(*D) = \mathcal{O}([a]D)$ for $a \in \mathbb{R}$, where [a] denote the integer such that $a-1 < [a] \le a$. The filtered bundle is denoted just by $\mathcal{O}_X(*D)$.

A morphism of filtered bundles $\psi: \mathcal{P}_*\mathcal{V}_1 \longrightarrow \mathcal{P}_*\mathcal{V}_2$ is defined to be a morphism of $\mathcal{O}_X(*D)$ -modules $\psi: \mathcal{V}_1 \longrightarrow \mathcal{V}_2$ such that $\psi(\mathcal{P}_a\mathcal{V}_1) \subset \mathcal{P}_a\mathcal{V}_2$ for any $a \in \mathbb{R}$.

Let $\mathcal{P}_*\mathcal{V}_i$ (i=1,2) be filtered bundles on (X,D). The direct sum $\mathcal{P}_*\mathcal{V}_1 \oplus \mathcal{P}_*\mathcal{V}_2$ is defined as the $\mathcal{O}_X(*D)$ module $\mathcal{V}_1 \oplus \mathcal{V}_2$ with the filtration $\mathcal{P}_a(\mathcal{V}_1 \oplus \mathcal{V}_2) := \mathcal{P}_a\mathcal{V}_1 \oplus \mathcal{P}_a\mathcal{V}_2$ $(a \in \mathbb{R})$. We naturally have $\operatorname{Gr}_a^{\mathcal{P}}(\mathcal{V}_1 \oplus \mathcal{V}_2) \simeq$ $\operatorname{Gr}_a^{\mathcal{P}}(\mathcal{V}_1) \oplus \operatorname{Gr}_a^{\mathcal{P}}(\mathcal{V}_2)$. The tensor product $\mathcal{P}_*\mathcal{V}_1 \otimes \mathcal{P}_*\mathcal{V}_2$ is defined as the $\mathcal{O}_X(*D)$ -module $\mathcal{V}_1 \otimes \mathcal{V}_2$ with the filtration $\mathcal{P}_a(\mathcal{V}_1 \otimes \mathcal{V}_2)$ $(a \in \mathbb{R})$:

$$\mathcal{P}_a(\mathcal{V}_1 \otimes \mathcal{V}_2) = \sum_{b+c \leq a} \mathcal{P}_b \mathcal{V}_1 \otimes \mathcal{P}_c \mathcal{V}_2 = \sum_{\substack{b+c \leq a \\ -1 < c \leq 0}} \mathcal{P}_b \mathcal{V}_1 \otimes \mathcal{P}_c \mathcal{V}_2 \quad (a \in \mathbb{R})$$

We define $\mathcal{H}om(\mathcal{P}_*\mathcal{V}_1, \mathcal{P}_*\mathcal{V}_2)$ as the $\mathcal{O}_X(*D)$ -module $\mathcal{H}om(\mathcal{V}_1, \mathcal{V}_2)$ with the filtration $\mathcal{P}_a\mathcal{H}om(\mathcal{V}_1, \mathcal{V}_2)$ $(a \in \mathbb{R})$:

$$\mathcal{P}_a\mathcal{H}om(\mathcal{V}_1,\mathcal{V}_2) := \left\{ f \in \mathcal{H}om(\mathcal{V}_1,\mathcal{V}_2) \,\middle|\, f\left(\mathcal{P}_b\mathcal{V}_1\right) \subset \mathcal{P}_{b+a}\mathcal{V}_2 \right\}$$

As a special case, we have the dual $\mathcal{P}_*\mathcal{V}^\vee$ of $\mathcal{P}_*\mathcal{V}$ as $\mathcal{H}om(\mathcal{P}_*\mathcal{V},\mathcal{O}_X(*D))$.

Pull back, push-forward, descent We set $X_1 := \{w \in \mathbb{C} \mid |w| < 1\}$ and $D_1 := \{w = 0\}$, and let $\varphi : X_1 \longrightarrow X$ be given by $\varphi(w) = w^m$. Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle on (X, D). We have an induced filtered bundle $\varphi^*\mathcal{P}_*\mathcal{V}$, called the pull back of $\mathcal{P}_*\mathcal{V}$ by φ . It is the $\mathcal{O}_{X_1}(*D_1)$ -module $\varphi^*\mathcal{V}$ with the filtrations $(\mathcal{P}_a\varphi^*\mathcal{V} \mid a \in \mathbb{R})$:

$$\mathcal{P}_a \varphi^* \mathcal{V} = \sum_{mb+n \le a} \varphi^* \mathcal{P}_b \mathcal{V} \otimes \mathcal{O}(nD_1)$$

Let $\mathcal{P}_*\mathcal{V}_1$ be a filtered bundle on (X_1, D_1) . We have an induced filtered bundle $\varphi_*\mathcal{P}_*\mathcal{V}_1$ on (X, D), called the push-forward of $\mathcal{P}_*\mathcal{V}_1$ by φ . It is the $\mathcal{O}_X(*D)$ -module $\varphi_*\mathcal{V}$ with the filtration $(\mathcal{P}_a(\varphi_*\mathcal{V}) \mid a \in \mathbb{R})$, where $\mathcal{P}_a(\varphi_*\mathcal{V}_1) = \varphi_*\mathcal{P}_{am}\mathcal{V}_1$.

Let $\operatorname{Gal}(\varphi)$ be the Galois group of the ramified covering φ . In this case, we have $\operatorname{Gal}(\varphi) = \{t \in \mathbb{C} \mid t^m = 1\}$ with the natural action ρ on X_1 given by $\rho(t, w) = tw$. If $\mathcal{P}_*\mathcal{V}_1$ is $\operatorname{Gal}(\varphi)$ -equivariant, then $\varphi_*\mathcal{P}_*\mathcal{V}_1$ is equipped with an induced $\operatorname{Gal}(\varphi)$ -action. The invariant part is called the descent of $\mathcal{P}_*\mathcal{V}_1$.

If $\mathcal{P}_*\mathcal{V}$ is a filtered bundle on (X, D), the pull back $\varphi^*\mathcal{P}_*\mathcal{V}$ is naturally $\operatorname{Gal}(\varphi)$ -equivariant, and the descent of $\varphi^*\mathcal{P}_*\mathcal{V}$ is naturally isomorphic to $\mathcal{P}_*\mathcal{V}$.

Subobject Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle on (X, D). A subobject of $\mathcal{P}_*\mathcal{V}$ is an $\mathcal{O}_X(*D)$ -submodule $\mathcal{V}' \subset \mathcal{V}$ with a filtration $\{\mathcal{P}_a\mathcal{V}' \mid a \in \mathbb{R}\}$ such that $\mathcal{P}_a\mathcal{V}' \subset \mathcal{P}_a\mathcal{V}$. It is called strict, if $\mathcal{P}_a\mathcal{V}' = \mathcal{P}_a\mathcal{V} \cap \mathcal{V}'$ for any a. Note that if we are given an $\mathcal{O}_X(*D)$ -submodule $\mathcal{V}' \subset \mathcal{V}$, it has an induced filtration $\{\mathcal{P}_a\mathcal{V}'\}$ given by $\mathcal{P}_a\mathcal{V}' := \mathcal{P}_a\mathcal{V} \cap \mathcal{V}'$, and such $\mathcal{P}_*\mathcal{V}'$ is strict. In practice, we have only to consider such strict subobjects.

Remark 2.1 Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle on (X, D). Let $\mathcal{V} = \bigoplus \mathcal{V}_i$ be a direct sum of $\mathcal{O}_X(*D)$ -modules. Then, we have the induced subobjects $\mathcal{P}_*\mathcal{V}_i$ as above. In general, we may have $\mathcal{P}_*\mathcal{V} \not\simeq \bigoplus \mathcal{P}_*\mathcal{V}_i$. If $\mathcal{P}_*\mathcal{V} \simeq \bigoplus \mathcal{P}_*\mathcal{V}_i$, we say that the filtration $\mathcal{P}_*\mathcal{V}$ is compatible with the decomposition $\mathcal{V} = \bigoplus \mathcal{V}_i$.

Filtered Higgs bundle Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle on (X, D). A Higgs field of a filtered bundle $\mathcal{P}_*\mathcal{V}$ is an \mathcal{O}_X -morphism $\theta: \mathcal{V} \longrightarrow \mathcal{V} \otimes \Omega^1_X$. The pair $(\mathcal{P}_*\mathcal{V}, \theta)$ is called a filtered Higgs bundle on (X, D). We say that (\mathcal{V}, θ) is the underlying meromorphic Higgs bundle of $(\mathcal{P}_*\mathcal{V}, \theta)$, and that $(\mathcal{P}_*\mathcal{V}, \theta)$ is a filtered Higgs bundle over (\mathcal{V}, θ) .

A morphism of filtered Higgs bundles $\psi: (\mathcal{P}_*\mathcal{V}_1, \theta_1) \longrightarrow (\mathcal{P}_*\mathcal{V}_2, \theta_2)$ is defined to be a morphism of filtered bundles $\psi: \mathcal{P}_*\mathcal{V}_1 \longrightarrow \mathcal{P}_*\mathcal{V}_2$ such that $\psi \circ \theta_1 = \theta_2 \circ \psi$. A subobject of $(\mathcal{P}_*\mathcal{V}, \theta)$ is a subobject $\mathcal{P}_*\mathcal{V}' \subset \mathcal{P}_*\mathcal{V}$ such that $\theta(\mathcal{V}') \subset \mathcal{V}' \otimes \Omega_X^1$.

Let $(\mathcal{P}_*\mathcal{V}_i, \theta_i)$ (i = 1, 2) be filtered Higgs bundles on (X, D). We have the naturally induced Higgs fields of $\theta^{(1)}$, $\theta^{(2)}$ and $\theta^{(3)}$ on $\mathcal{P}_*\mathcal{V}_1 \oplus \mathcal{P}_*\mathcal{V}_2$, $\mathcal{P}_*\mathcal{V}_1 \otimes \mathcal{P}_*\mathcal{V}_2$ and $\mathcal{H}om(\mathcal{P}_*\mathcal{V}_1, \mathcal{P}_*\mathcal{V}_2)$, respectively: $\theta^{(1)}(v_1, v_2) = (\theta_1(v_1), \theta_2(v_2)), \theta^{(2)}(v_1 \otimes v_2) = \theta_1(v_1) \otimes v_2 + v_1 \otimes \theta_2(v_2), \theta^{(3)}(f) = \theta_2 \circ f - f \circ \theta_1$. In particular, we have the dual $(\mathcal{P}_*\mathcal{V}, \theta)^{\vee} = (\mathcal{P}_*\mathcal{V}, -\theta^{\vee})$ of a filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}, \theta)$.

Let $\varphi: X_1 \longrightarrow X$ be a morphism given by $\varphi(w) = w^m$. For a filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}, \theta)$ on (X, D), we have its pull back $\varphi^*(\mathcal{P}_*\mathcal{V}, \theta) = (\varphi^*\mathcal{P}_*\mathcal{V}, \varphi^*\theta)$ on (X_1, D_1) . For a filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}_1, \theta_1)$ on (X_1, D_1) , we have its push-forward $\varphi_*(\mathcal{P}_*\mathcal{V}_1, \theta_1) = (\varphi_*\mathcal{P}_*\mathcal{V}_1, \varphi_*\theta_1)$ on (X, D). If $(\mathcal{P}_*\mathcal{V}_1, \theta_1)$ is $Gal(\varphi)$ -equivariant, we have its descent as the $Gal(\varphi)$ -invariant part of $\varphi_*(\mathcal{P}_*\mathcal{V}_1, \theta_1)$.

Good filtered Higgs bundle and regular filtered Higgs bundle Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a filtered Higgs bundle. It is called regular, if $\theta(\mathcal{P}_a\mathcal{V}) \subset \mathcal{P}_a\mathcal{V} \otimes \Omega^1_X(\log D) = \mathcal{P}_{a+1}\mathcal{V} \otimes \Omega^1_X$. In that case, θ is called logarithmic. A filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}, \theta)$ is called unramifiedly good, if there exist a subset $\mathcal{I} \subset z^{-1}\mathbb{C}[z^{-1}]$ and a decomposition of filtered bundles $\mathcal{P}_*\mathcal{V} = \bigoplus_{a \in \mathcal{I}} \mathcal{P}_*\mathcal{V}_a$ on a neighbourhood \mathcal{U} of D, such that the following holds for each \mathfrak{a} :

- $\theta(\mathcal{V}_{\mathfrak{a}}) \subset \mathcal{V}_{\mathfrak{a}}$. The restriction of θ to $\mathcal{V}_{\mathfrak{a}}$ is denoted by $\theta_{\mathfrak{a}}$.
- $\theta_{\mathfrak{a}} d\mathfrak{a}$ id $\mathcal{V}_{\mathfrak{a}}$ is logarithmic with respect to $\mathcal{P}_*\mathcal{V}_{\mathfrak{a}}$.

A filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}, \theta)$ is called good, if there exists a ramified covering $\varphi : (X_1, D_1) \longrightarrow (X, D)$ given by $\varphi(w) = w^m$ such that $\varphi^*(\mathcal{P}_*\mathcal{V}, \theta)$ is unramifiedly good. The following lemma is easy to see.

Lemma 2.2 Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a good filtered Higgs bundle on (X, D). Assume that θ is tame, i.e., for the expression $\theta = f \, dz/z$, the coefficients of the characteristic polynomial of f are holomorphic at 0. Then, $(\mathcal{P}_*\mathcal{V}, \theta)$ is regular.

2.1.2 Filtered Higgs bundles on a curve

Let X be a complex curve with a finite discrete subset $D \subset X$. The notions of filtered bundle and filtered Higgs bundle are naturally generalized in this global situation. We take small neighbourhoods U_P of any $P \in D$. A filtered bundle $\mathcal{P}_*\mathcal{V}$ on (X, D) is a locally free $\mathcal{O}_X(*D)$ -module \mathcal{V} of finite rank with the filtrations $\mathcal{P}_*^{(P)}(\mathcal{V}_{|U_P}) = (\mathcal{P}_a^{(P)}(\mathcal{V}_{|U_P}) \mid a \in \mathbb{R})$ by $\mathcal{O}_{U_P}(*P)$ -modules for each $P \in U_P$, such that $\mathcal{P}_*^{(P)}(\mathcal{V}_{|U_P})$ is a filtered bundle on (U_P, P) . For each $P \in D$, we obtain the set $\mathcal{P}ar(\mathcal{P}_*\mathcal{V}, P) := \mathcal{P}ar(\mathcal{P}_*^{(P)}(\mathcal{V}_{|U_P}))$. For each $P \in D$, we have an \mathcal{O}_X -submodule $\mathcal{P}_a\mathcal{V} \subset \mathcal{V}$ determined by $\mathcal{P}_a\mathcal{V}_{|U_P} = \mathcal{P}_{a_P}^{(P)}(\mathcal{V}_{|U_P})$ for any $P \in D$. Obviously, the tuple $\{\mathcal{P}_a\mathcal{V} \mid \mathbf{a} \in \mathbb{R}^D\}$ determines $\mathcal{P}_*\mathcal{V}$. A filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}, \theta)$ is a filtered bundle $\mathcal{P}_*\mathcal{V}$ with a Higgs field θ of \mathcal{V} .

Direct sum, tensor product and inner homomorphism, subobjects and morphisms are defined as in the case of filtered bundles on the disc. A filtered Higgs bundle $(\mathcal{P}_*\mathcal{V}, \theta)$ is called logarithmic (resp. good) if $(\mathcal{P}_*\mathcal{V}, \theta)_{|U_P|}$ is logarithmic (resp. good) for each $P \in D$. The other operations are also naturally generalized in the global case.

2.1.3 Parabolic degree and stability condition

Let X be a projective curve with a finite subset $D \subset X$. Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle. Recall that the parabolic degree of $\mathcal{P}_*\mathcal{V}$ is given as follows:

$$\operatorname{par-deg}(\mathcal{P}_*\mathcal{V}) := \operatorname{deg}(\mathcal{P}_a\mathcal{V}) - \sum_{P \in D} \sum_{\substack{c \in \mathcal{P}ar(\mathcal{P}_*\mathcal{V}, P) \\ a_P - 1 < c < a_P}} c \operatorname{dim} \operatorname{Gr}_c^{\mathcal{P}^{(P)}}(\mathcal{V}_{|U_P})$$

$$\tag{2}$$

It is independent of the choice of $\mathbf{a} = (a_P \mid P \in D) \in \mathbb{R}^D$ in (2). We define $\mu(\mathcal{P}_*\mathcal{V}) := \operatorname{par-deg}(\mathcal{P}_*\mathcal{V})/\operatorname{rank}\mathcal{V}$. Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a filtered Higgs bundle on (X, D). It is called stable, if $\mu(\mathcal{P}_*\mathcal{V}_1) < \mu(\mathcal{P}_*\mathcal{V})$ holds for any subobject $(\mathcal{P}_*\mathcal{V}_1, \theta_1) \subset (\mathcal{P}_*\mathcal{V}, \theta)$ such that $0 < \operatorname{rank}\mathcal{V}_1 < \operatorname{rank}\mathcal{V}$. It is called semistable, if the condition " $\mu(\mathcal{P}_*\mathcal{V}_1) < \mu(\mathcal{P}_*\mathcal{V})$ " is replaced with " $\mu(\mathcal{P}_*\mathcal{V}_1) \le \mu(\mathcal{P}_*\mathcal{V})$ ". It is called poly-stable, if we have a decomposition $(\mathcal{P}_*\mathcal{V}, \theta) = \bigoplus (\mathcal{P}_*\mathcal{V}_i, \theta_i)$ such that each $(\mathcal{P}_*\mathcal{V}_i, \theta_i)$ is stable with $\mu(\mathcal{P}_*\mathcal{V}_i) = \mu(\mathcal{P}_*\mathcal{V})$. The following standard fact will be used implicitly.

Lemma 2.3 Suppose that $(\mathcal{P}_*\mathcal{V}_i, \theta_i)$ (i = 1, 2) are stable. The space of morphisms $(\mathcal{P}_*\mathcal{V}_1, \theta_1) \longrightarrow (\mathcal{P}_*\mathcal{V}_2, \theta_2)$ is at most one dimensional, and any non-zero morphism is an isomorphism.

2.2 Harmonic bundles on curves

Let Y be a complex curve. Let $(E, \overline{\partial}_E)$ be a holomorphic vector bundle on Y. Let θ be a Higgs field of $(E, \overline{\partial}_E)$. Let h be a hermitian metric of E. We have the Chern connection, whose (1,0)-part is denoted by ∂_E . Let θ^{\dagger} denote the adjoint of θ . The metric h is called harmonic, if the connection $\nabla^1 := \overline{\partial}_E + \theta^{\dagger} + \partial_E + \theta$ is flat. In that case, $(E, \overline{\partial}_E, \theta, h)$ is called a harmonic bundle. Note that $(E, \overline{\partial}_E, \alpha\theta, h)$ is also a harmonic bundle if α is a complex number with $|\alpha| = 1$. If the base space is higher dimensional, such a metric is called pluri-harmonic.

2.2.1 Prolongation of a harmonic bundle on a punctured disc

Let $X:=\{z\in\mathbb{C}\,|\,|z|<1\}$ and $D:=\{0\}$. Let us consider a harmonic bundle $(E,\overline{\partial}_E,\theta,h)$ on $X\setminus D$. We have the expression $\theta=f\,dz/z$ where f is a holomorphic section of $\operatorname{End}(E)$. We have the characteristic polynomial $P=\det(t\operatorname{id}_E-f)=\sum a_j(z)t^j$. The harmonic bundle is called tame if $a_j(z)$ are holomorphic on X, and it is called wild if $a_j(z)$ are meromorphic on X. It is called unramifiedly good wild, if there exist a subset $\mathcal{I}\subset z^{-1}\mathbb{C}[z^{-1}]$ and a decomposition $(E,\theta)=\bigoplus_{\mathfrak{a}\in\mathcal{I}}(E_{\mathfrak{a}},\theta_{\mathfrak{a}})$ such that $\theta_{\mathfrak{a}}-d\mathfrak{a}$ id $_{E_{\mathfrak{a}}}$ are tame for each $\mathfrak{a}\in z^{-1}\mathbb{C}[z^{-1}]$. Note that, if $(E,\overline{\partial}_E,\theta,h)$ is wild, there exists a ramified covering $\varphi:(X_1,D_1)\longrightarrow (X,D)$, such that $\varphi^{-1}(E,\overline{\partial}_E,\theta,h)$ is unramifiedly good wild.

Let $(E, \overline{\partial}_E, \theta, h)$ be a wild harmonic bundle on (X, D). For any $a \in \mathbb{R}$, we have the \mathcal{O}_X -module $\mathcal{P}_a E$ given as follows. Let U be an open subset of X. If $0 \notin U$, let $\mathcal{P}_a E(U)$ denote the space of holomorphic sections of E on U. If $0 \in U$, let $\mathcal{P}_a E(U)$ denote the space of holomorphic sections f of E on $U \setminus D$ such that $|f|_h = O(|z|^{-a-\epsilon})$ for any $\epsilon > 0$. We define $\mathcal{P}E := \bigcup_{a \in \mathbb{R}} \mathcal{P}_a E$, which is an $\mathcal{O}_X(*D)$ -module. The following proposition was proved by Simpson [44] in the tame case, and by the author in the general case [36].

Proposition 2.4 (\mathcal{P}_*E, θ) is a good filtered Higgs bundle on (X, D). If the harmonic bundle is tame, (\mathcal{P}_*E, θ) is regular. If the harmonic bundle is unramifiedly good wild, then (\mathcal{P}_*E, θ) is unramifiedly good.

Let $(\mathcal{P}_*\mathcal{V},\theta)$ be a filtered Higgs bundle on (X,D). Let h be a harmonic metric of $(E,\theta):=(\mathcal{V},\theta)_{|X\setminus D}$. We obtain a filtered bundle \mathcal{P}_*E as above. If the identity $E\simeq\mathcal{V}_{|X\setminus D}$ on $X\setminus D$ is extended to an isomorphism $\mathcal{P}_*E\simeq\mathcal{P}_*\mathcal{V}$ on (X,D), we say that h is adapted to the filtered bundle $\mathcal{P}_*\mathcal{V}$.

2.2.2 Harmonic bundles on projective curves

Let X be a projective curve with a finite subset $D \subset X$. We take small neighbourhoods U_P of any $P \in D$. Let $(E, \overline{\partial}_E, \theta, h)$ be a harmonic bundle on $X \setminus D$. It is called tame (resp. wild, unramifiedly good wild) if $(E, \overline{\partial}_E, \theta, h)_{|U_P \setminus P|}$ is tame (resp. wild, unramifiedly good wild) for any $P \in D$. By applying the procedure in §2.2.1 to each $(E, \overline{\partial}_E, \theta, h)_{|U_P \setminus P|}$, we obtain a good filtered Higgs bundle (\mathcal{P}_*E, θ) on (X, D). **Proposition 2.5** (\mathcal{P}_*E, θ) is polystable and satisfies par-deg $(\mathcal{P}_*E) = 0$.

Proof The tame case was proved by Simpson (Theorem 5 in [44]). The general case can be shown similarly.

Proposition 2.6 Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a filtered Higgs bundle on (X, D). Let h_j (j = 1, 2) be two harmonic metrics of $(E, \theta) := (\mathcal{V}, \theta)_{|X \setminus D}$ adapted to $\mathcal{P}_*\mathcal{V}$. Then, there exists a decomposition $(\mathcal{P}_*\mathcal{V}, \theta) = \bigoplus_{i \in I} (\mathcal{P}_*\mathcal{V}_i, \theta_i)$ such that the following holds:

- We set $(E_i, \theta_i) := (\mathcal{V}_i, \theta_i)_{|X \setminus D}$. Then, the decomposition $(E, \theta) = \bigoplus (E_i, \theta_i)$ is orthogonal with respect to both of h_j (j = 1, 2).
- There exist positive numbers a_i such that $h_{1|E_i} = a_i h_{2|E_i}$.

In particular, if $(\mathcal{P}_*\mathcal{V}, \theta)$ is stable, the adapted harmonic metric is unique up to the multiplication of positive constants.

Proof The argument for the tame case is given in §2.2 of [30], for example. The general case can be shown similarly.

Conversely, the following theorem holds. The tame case is due to Simpson [44]. The wild case is due to Biquard-Boalch [5], although some inessential assumption is imposed on the residues. We may also prove it as in [36] where a similar statement is established for filtered flat bundles by directly using the method in [44]. We are planning to give more details elsewhere, including the higher dimensional case.

Theorem 2.7 Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a μ -stable Higgs bundle on (X, D) with $\mu(\mathcal{P}_*\mathcal{V}) = 0$. Then, there exists a harmonic metric h of $(E, \theta) := (\mathcal{V}, \theta)_{|X \setminus D}$ such that $\mathcal{P}_{\mathbf{a}}E = \mathcal{P}_{\mathbf{a}}\mathcal{V}$ for any $\mathbf{a} \in \mathbb{R}^D$. Such a metric is unique up to the multiplication of a positive constant.

We can apply it if the Higgs bundle is irreducible, for example.

2.2.3 Symmetry

By the uniqueness in Kobayashi-Hitchin correspondence, the symmetry of a filtered Higgs bundle is inherited to a harmonic bundle. For the explanation, we give a typical statement. Let X be a smooth projective curve with a finite subset $D \subset X$. Let f be a holomorphic automorphism of (X, D). Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a stable good filtered Higgs bundle on (X, D), with an isomorphism $\tilde{f}: f^*\mathcal{P}_*\mathcal{V} \simeq \mathcal{P}_*\mathcal{V}$. Suppose $f^*\theta = \alpha\theta$ for a complex number α with $|\alpha| = 1$ under the isomorphism.

Proposition 2.8 There exists a > 0 such that $f^*h = ah$ under \tilde{f} .

Proof Put $(E, \theta) := (\mathcal{V}, \theta)_{|X \setminus D}$. Because h and f^*h are harmonic metrics of $(E, \overline{\partial}_E, \alpha\theta)$ adapted to $\mathcal{P}_*\mathcal{V}$, we obtain $f^*h = ah$ by the uniqueness in Theorem 2.7.

We can obtain the number a by comparing det(h) and $f^* det(h)$.

3 Toda-like harmonic bundles

3.1 Toda-like filtered Higgs bundles on $(\mathbb{P}^1, \{0, \infty\})$

3.1.1 Meromorphic Higgs bundles

Let r and m be positive integers. Let q be a variable. Let $\mathcal{K}(r,m)(q)$ be the matrix whose (i,j)-entries $\mathcal{K}(r,m)_{i,j}(q)$ are 1 if i=j+1, q^m if (i,j)=(1,r), or 0 otherwise. We may regard $\mathcal{K}(r,m)$ as a matrix-valued holomorphic function on $\mathbb{C}_q = \{q \in \mathbb{C}\}$, which is meromorphic at $q = \infty$.

holomorphic function on $\mathbb{C}_q = \{q \in \mathbb{C}\}$, which is meromorphic at $q = \infty$. We set $D := \{0, \infty\} \subset \mathbb{P}^1$. Let \mathcal{V}_r be a free $\mathcal{O}_{\mathbb{P}^1}(*D)$ -module $\bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}(*D)e_i$. Let $\theta_{r,m}$ be the Higgs field determined by $\theta_{r,m}e = e \mathcal{K}(r,m) \, m \, dq/q$, where $e = (e_1, \ldots, e_r)$. **Remark 3.1** Let \mathcal{J} be the r-square matrix whose (i,j)-entries are $\alpha_i q^{\ell_i}$ if $i-j \equiv 1$ modulo r, or 0 otherwise. Let θ be the Higgs field of \mathcal{V}_r determined by $\theta e = e \mathcal{J} dq/q$. If $m = \sum \ell_i > 0$, (\mathcal{V}_r, θ) is transformed to $(\mathcal{V}_r, \theta_{r,m})$ by an appropriate gauge transform and an appropriate coordinate change of \mathbb{P}^1 .

Let $\tau = e^{2\pi\sqrt{-1}/r}$. Let σ be the endomorphism of \mathcal{V}_r determined by $\sigma(e_i) = \tau^i e_i$. We have $\sigma^*\theta := \sigma \circ \theta_{r,m} \circ \sigma^{-1} = \tau \theta_{r,m}$. Hence, σ induces an isomorphism of meromorphic Higgs bundles $\sigma : (\mathcal{V}_r, \theta_{r,m}) \simeq (\mathcal{V}_r, \tau \theta_{r,m})$.

We have a \mathbb{C}^* -action $\mathbb{C}^* \times \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ given by $(t,q) \longmapsto t^r q$. The bundle \mathcal{V}_r is \mathbb{C}^* -equivariant by the isomorphism $\widetilde{t}: t^* \mathcal{V}_r \simeq \mathcal{V}_r$, $\widetilde{t}(t^* e_i) = t^{im} e_i$. We have $t^* \theta = t^m \theta$.

Let ℓ be a factor of m. We set $\mu_{\ell} := \{ \kappa \in \mathbb{C} \mid \kappa^{\ell} = 1 \}$. We have a natural μ_{ℓ} -action on \mathbb{P}^1 given by $(\kappa, q) \longmapsto \kappa q$. Let $\widetilde{\kappa} : \kappa^* \mathcal{V}_r \simeq \mathcal{V}_r$ be the isomorphism given by $\widetilde{\kappa}(\kappa^* e_i) = e_i$. It induces a μ_{ℓ} -action on (\mathcal{V}_r, θ) . The descent is isomorphic to $(\mathcal{V}_r, \theta_{r,m/\ell})$.

Let $\varphi: \mathbb{P}^1_w \longrightarrow \mathbb{P}^1_q$ be given by $\varphi^*(q) = w^r$. For $j = 1, \ldots, r$, we set

$$v_j := \sum_{i=1}^r (\tau^j w^m)^{r-i} \varphi^* e_i.$$
 (3)

We have the following formulas:

$$\varphi^*(\theta_{r,m})v_j = \left(r \, m \, w^m \tau^j\right) v_j \frac{dw}{w} \tag{4}$$

We also have the following:

$$\sigma(v_i) = v_{i-1} \quad (1 \le j \le r - 1), \quad \sigma(v_r) = v_1.$$
 (5)

We have a natural \mathbb{C}^* -action on \mathbb{P}^1_w given by $(t, w) \longmapsto tw$. The morphism φ is \mathbb{C}^* -equivariant, and $\varphi^* \mathcal{V}_r$ is naturally \mathbb{C}^* -equivariant. We have $\widetilde{t}(t^*v_i) = t^{mr}v_i$.

We have a μ_m -action on \mathbb{P}^1_w given by $(\kappa, w) \longmapsto \kappa w$. (The morphism φ is not necessarily μ_m -equivariant.) We have a μ_m -action on $\varphi^* \mathcal{V}_r$ by $\widetilde{\kappa}(\kappa^* e_i) = e_i$. We have $\widetilde{\kappa}(\kappa^* v_j) = v_j$. Indeed, we have $\widetilde{\kappa}(\kappa^* v_j) = \sum_i (\tau^j w^m)^{r-i} \widetilde{\kappa}(\kappa^* e_i) = v_j$.

We have the description of e_k in terms of v_j as follows:

$$e_k = \frac{1}{rw^{m(r-k)}} \sum_{i=1}^r \tau^{ki} v_i$$
 (6)

Indeed, $\sum_{j=1}^{r} \tau^{kj} v_j = \sum_{i=1}^{r} \sum_{j=1}^{r} \tau^{j(k-i)} (w^m)^{(r-i)} e_i = rw^{m(r-k)} e_k$.

3.1.2 Filtrations around 0

Let $U_0 = \mathbb{P}^1 \setminus \{\infty\}$. We shall consider filtered bundles on $(U_0, 0)$ over $\mathcal{V} := \mathcal{V}_{r,m|U_0}$. The induced sections $e_{i|U_0}$ are denoted by e_i for simplicity. If we are given a tuple of real numbers $\mathbf{a} = (a_1, \dots, a_r) \in \mathbb{R}^r$, \mathcal{V} is equipped with the filtration \mathcal{P}^a_* given by $\mathcal{P}^a_a \mathcal{V} = \bigoplus \mathcal{O}([a-a_i]0)e_i$. Here, $[a] := \max\{c \in \mathbb{R} \mid c \in \mathbb{Z}, c \leq a\}$ for any $a \in \mathbb{R}$. The filtration is compatible with the decomposition $\mathcal{V} = \bigoplus \mathcal{O}_{U_0}(*0)e_i$. Conversely, if a filtration \mathcal{P}_* is compatible with the decomposition, it is \mathcal{P}^a_* for some \mathbf{a} .

Lemma 3.2 Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle over \mathcal{V} . The automorphism σ of \mathcal{V} gives an automorphism of $\mathcal{P}_*\mathcal{V}$, i.e., $\sigma(\mathcal{P}_a\mathcal{V}) \subset \mathcal{P}_a\mathcal{V}$, if and only if $\mathcal{P}_*\mathcal{V} \simeq \mathcal{P}_*^a\mathcal{V}$ for some $\mathbf{a} \in \mathbb{R}^r$.

Proof The "if" part is clear. To prove the "only if" part, put $a_i := \min\{a \in \mathbb{R} \mid e_i \in \mathcal{P}_a \mathcal{V}\}$, and $\mathbf{a} := (a_i)$. We clearly have $\mathcal{P}_a^a \mathcal{V} \subset \mathcal{P}_a \mathcal{V}$. A section $f \in \mathcal{P}_a \mathcal{V}$ has the expression $\sum f_i e_i$. Because $\sigma^j(f) = \sum \tau^{ij} f_i e_i \in \mathcal{P}_a \mathcal{V}$, we obtain $f_i e_i \in \mathcal{P}_a \mathcal{V}$. Hence, we obtain $f_i \in \mathcal{O}([a-a_i])$, which means $f \in \mathcal{P}_a^a \mathcal{V}$.

The following lemma is clear by the construction of \mathcal{P}^a_* .

Lemma 3.3 If the conditions in Lemma 3.2 hold, \mathcal{P}_* is \mathbb{C}^* -equivariant.

Note that, in general, the converse of the claim in Lemma 3.3 does not hold. For example, in the case r=m=2 and $0=a_1=a_2+1$, e_1 and $q^{-1}e_2$ are \mathbb{C}^* -invariant sections. By using any one dimensional subspace of $\langle e_1, q^{-1}e_2 \rangle$, we obtain a filtration $\mathcal{P}_*\mathcal{V}_2$ such that it is homogeneous with respect to the \mathbb{C}^* -action, but it is not isomorphic to $\mathcal{P}_*^a\mathcal{V}_2$.

The following lemma is also clear by construction.

Lemma 3.4 If the conditions in Lemma 3.2 hold, \mathcal{P}_* is μ_m -equivariant.

Let $\theta := \theta_{r,m|U_0}$. We consider the condition such that θ is logarithmic with respect to the filtered bundle $\mathcal{P}_*\mathcal{V}$. The following lemma is clear.

Lemma 3.5 Let $\mathcal{P}_*\mathcal{V}$ be a filtered bundle over \mathcal{V} . We set $a_i := \min\{a \mid e_i \in \mathcal{P}_a\mathcal{V}\}$. If θ is logarithmic, then we have $a_i \geq a_{i+1}$ $(i = 1, \ldots, r-1)$ and $a_r \geq a_1 - m$. If $\mathcal{P}_*\mathcal{V} = \mathcal{P}_*^a\mathcal{V}$ for $\mathbf{a} = (a_i)$, the converse is also true.

Lemma 3.6 Assume that $(\mathcal{P}_*\mathcal{V}, \theta)$ is regular. After a canonical change of frame, we may assume that $a_r > a_1 - m$.

Proof Assume that $a_r = a_1 - m$. We have j such that $a_j > a_{j+1} = a_r = a_1 - m$. We set $e'_i := q^{-m}e_{j+i}$ for $i \le r - j$, and $e'_i := e_{i-r+j}$ for i > r - j. Then, we have $\theta e' = e' \mathcal{K}(r, m)$, and we have $a'_r > a'_1 - m$.

The case m = 1 is rather special.

Lemma 3.7 Assume that m = 1, and that $(\mathcal{P}_*\mathcal{V}, \theta)$ is regular. Then, the conditions in Lemma 3.2 hold. In particular, $\mathcal{P}_*\mathcal{V}$ is \mathbb{C}^* -equivariant.

Proof Let us consider the case $a_r > a_1 - 1$. Each e_i gives a section of $\mathcal{P}_{a_i}\mathcal{V}$. The induced elements of $\operatorname{Gr}_{a_i}^{\mathcal{P}}(\mathcal{V})$ are denoted by u_i . For each $a_1 - 1 < c \le a_1$, we set $S(c) := \{i \mid a_i = c\}$. We have only to prove that $\{u_i \mid i \in S(c)\}$ is a base of $\operatorname{Gr}_c^{\mathcal{P}}(\mathcal{V})$.

Assume that u_i $(i \in S(c))$ are linearly dependent in $\operatorname{Gr}_c^{\mathcal{P}}(\mathcal{V})$, and we shall deduce a contradiction. Let $i_0 := \min\{i \in S(c)\}$ and $i_1 := \max\{i \in S(c)\}$. We have $\operatorname{Res}(\theta)u_i = u_{i+1}$ for $i_0 \leq i < i_1$ and $\operatorname{Res}(\theta)u_{i_1} = 0$. By the assumption, we have a non-trivial linear relation $\sum \alpha_j u_j = 0$. We have i_2 such that $\alpha_{i_2} \neq 0$ and $\alpha_j = 0$ for $j > i_2$. We may assume that u_i $(i = i_0, \ldots, i_2 - 1)$ are linearly independent. Then, because $u_{i_2} = \sum_{j=i_0}^{i_2-1} \beta_j u_j$, $\operatorname{Res}(\theta)$ preserves the subspace H generated by u_j $(i_0 \leq j < i_2)$. But, $\operatorname{Res}(\theta)_{|H}$ must have a non-zero eigenvalue. On the contrary, because the eigenvalues of $\mathcal{K}(r,1)_{|q=0}$ are 0, $\operatorname{Res}(\theta)$ is nilpotent on $\operatorname{Gr}_c^{\mathcal{P}}(\mathcal{V})$ for any $c \in \mathbb{R}$. Thus, we arrive at a contradiction. Hence, we obtain that u_j $(j \in S(c))$ are linearly independent. Then, we can prove that they give a base of $\operatorname{Gr}_c^{\mathcal{P}}$, by using $\sum_{a_1-1 < c \leq a_1} \dim \operatorname{Gr}_c = r$.

Let us consider the case $a_r = a_1 - 1$. We use the frame e' as in the proof of Lemma 3.6. Then, by applying the result in the case $a_r > a_1 - 1$, we obtain that $\mathcal{P}_*\mathcal{V}$ is compatible with the decomposition $\mathcal{V} = \bigoplus \mathcal{O}_{U_0}(*0)e_i$, i.e., $\mathcal{P}_*\mathcal{V} \simeq \mathcal{P}_*^a\mathcal{V}$.

Corollary 3.8 Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a filtered Higgs bundle over $(\mathcal{V}_r, \theta_{r,m})_{|U_0}$. The conditions in Lemma 3.2 hold, if and only if $\mathcal{P}_*\mathcal{V}$ is μ_m -equivariant.

Proof The "only if" part is clear. Let us prove the "if" part. We have the descent $(\mathcal{P}_*\mathcal{V}', \theta')$ of $(\mathcal{P}_*\mathcal{V}, \theta)$ with respect to $\varphi: U_0 \longrightarrow U_0$ given by $\varphi(q) = q^m$, which is a filtered Higgs bundle over $(\mathcal{V}_r, \theta_{r,1})_{|U_0}$. Hence, the condition in Lemma 3.2 holds for $(\mathcal{P}_*\mathcal{V}', \theta')$. Because $(\mathcal{P}_*\mathcal{V}, \theta)$ is isomorphic to $\varphi^*(\mathcal{P}_*\mathcal{V}', \theta')$, the condition in Lemma 3.2 holds also for $(\mathcal{P}_*\mathcal{V}, \theta)$.

3.1.3 Filtration around ∞

Let $U_{\infty} := \mathbb{P}^1 \setminus \{0\}$. Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a good filtered Higgs bundle on (U_{∞}, ∞) over $(\mathcal{V}, \theta) := (\mathcal{V}_r, \theta_{r,m})_{|U_{\infty}}$. The morphism $\varphi : \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ induces $\varphi : U_{\infty} \longrightarrow U_{\infty}$. We have the induced filtered Higgs bundle $\varphi^*(\mathcal{P}_*\mathcal{V}, \theta)$. Because $(\mathcal{P}_*\mathcal{V}, \theta)$ is good, the filtration of $\varphi^*\mathcal{V}$ is compatible with the decomposition $\varphi^*\mathcal{V} = \bigoplus_{i=1}^r \mathcal{O}_{U_{\infty}}(*\infty)v_i$, where v_i are as in §3.1.1. Hence, $\mathcal{P}_*\mathcal{V}$ is determined by $b_i := \min\{b \mid v_i \in \mathcal{P}_b\varphi^*\mathcal{V}\}$ $(i = 1, \dots, r)$.

Lemma 3.9 Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a good filtered Higgs bundle on (U_∞, ∞) over (\mathcal{V}, θ) . The following conditions are equivalent:

- The filtered bundle $\mathcal{P}_*\mathcal{V}$ is compatible with the decomposition $\mathcal{V} = \bigoplus \mathcal{O}_{U_\infty}(*\infty)e_i$.
- The automorphism σ of V gives an automorphism of \mathcal{P}_*V .
- The automorphism of σ of $\varphi^* \mathcal{V}$ gives an automorphism of $\varphi^* \mathcal{P}_* \mathcal{V}$.
- b_i are independent of i.

Proof The equivalence of the first and second conditions can be proved by the argument in the proof of Lemma 3.2. The equivalence of the second and third conditions follows from the construction of the pull back and the descent. The third condition imply the fourth condition due to (5). The fourth condition implies the third condition, because $\varphi^*\mathcal{P}_*\mathcal{V}$ is compatible with the decomposition $\mathcal{V} = \bigoplus \mathcal{O}_{U_\infty}(*\infty)v_i$ by the assumption that $(\mathcal{P}_*\mathcal{V},\theta)$ is good.

The following lemma is easy to see.

Lemma 3.10 Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a good filtered Higgs bundle on (U_∞, ∞) over (\mathcal{V}, θ) . If the conditions in Lemma 3.9 hold, the filtration $\mathcal{P}_*\mathcal{V}$ is preserved by the \mathbb{C}^* -action.

Lemma 3.11 Assume that m and r are relatively prime. Let $(\mathcal{P}_*\mathcal{V}, \theta)$ be a good filtered Higgs bundle on (U_{∞}, ∞) over (\mathcal{V}, θ) . Then, the conditions in Lemma 3.9 hold.

Proof Because $Gal(\varphi)$ acts on $\{v_i\}_{i=1,\dots,r}$ transitively, b_i are independent of i.

Lemma 3.12 The conditions in Lemma 3.9 hold, if and only if $\mathcal{P}_*\mathcal{V}$ is μ_m -equivariant

Proof The "if" part is clear. Suppose that $\mathcal{P}_*\mathcal{V}$ is μ_m -equivariant. By Lemma 3.11, the conditions in Lemma 3.9 hold for the descent. Hence, they hold for $\mathcal{P}_*\mathcal{V}$.

3.1.4 Filtered Higgs bundle over $(\mathcal{V}_r, \theta_{r,m})$

Let $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,m})$ be a good filtered Higgs bundle on (\mathbb{P}^1, D) over $(\mathcal{V}_r, \theta_{r,m})$. Note that it is logarithmic at 0, according to Lemma 2.2. We have the following. (See Lemmas 3.2, 3.9, 3.3 and 3.10.)

Proposition 3.13 The automorphism σ of \mathcal{V}_r gives an automorphism of $\mathcal{P}_*\mathcal{V}_r$, if and only if $\mathcal{P}_*\mathcal{V}_r$ is compatible with the decomposition $\mathcal{V}_r = \bigoplus \mathcal{O}_{\mathbb{P}^1}(*D)e_i$. In that case, $\mathcal{P}_*\mathcal{V}_r$ is \mathbb{C}^* -equivariant.

Proposition 3.14 If m = 1, the conditions in Proposition 3.13 always hold. If $m \neq 1$, $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,m})$ satisfies the conditions in Proposition 3.13 if and only if it is μ_m -equivariant.

Proof The first claim follows from Lemmas 3.7 and 3.9. Note that the equivariance with respect to the Galois covering $w \longmapsto w^r$ ensures the 4-th condition in Lemma 3.9. The second claim follows from Corollary 3.8 and Lemma 3.12.

A Toda-like filtered Higgs bundle is a good filtered Higgs bundles $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,m})$ such that $\mathcal{P}_*\mathcal{V}_r$ is σ -invariant. Let $\mathcal{F}(r,m)$ denote the set of such Toda-like filtered Higgs bundles over $(\mathcal{V}_r, \theta_{r,m})$. Let $\varphi_m : \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ be given by $\varphi_m(q) = q^m$. We have the pull back $\varphi_m^* : \mathcal{F}(r,1) \longrightarrow \mathcal{F}(r,m)$. We obtain the following from Proposition 3.14.

Corollary 3.15 The map $\varphi_m^* : \mathcal{F}(r,1) \longrightarrow \mathcal{F}(r,m)$ is a bijection.

If m is relatively prime to r, the Higgs bundle $(\mathcal{V}_r, \theta_{r,m})$ is irreducible, and hence any good filtered Higgs bundle is stable. In general, we have the following.

Proposition 3.16 Any $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,m}) \in \mathcal{F}(r,m)$ is poly-stable. (We shall give the decomposition into the stable components in §3.1.5.)

Proof It is easily reduced to the case par-deg($\mathcal{P}_*\mathcal{V}_r$) = 0. Let $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,m}) \in \mathcal{F}(r,m)$. Because it is μ_m -equivariant, it is isomorphic to the pull back of some object $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,1})$ in $\mathcal{F}(r,1)$. Then, the claim follows from the Kobayashi-Hitchin correspondence. (See Theorem 2.7.)

Let $(\mathcal{P}_*\mathcal{V}_r,\theta) \in \mathcal{F}(r,m)$. The filtration around 0 (resp. ∞) is denoted by $\mathcal{P}_*^{(0)}$ (resp. $\mathcal{P}_*^{(\infty)}$). Let $\varphi: \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ be given by $\varphi(q) = q^r$. Recall that the parabolic structure at ∞ is determined by $b = \min\{c \mid v_i \in \mathcal{P}_c^{(\infty)}\varphi^*\mathcal{V}\}$ ($\forall i$). It is easy to observe that, if the condition par-deg $(\mathcal{P}_*\mathcal{V}) = 0$ is imposed, b is uniquely determined by $a_i := \min\{c \mid e_i \in \mathcal{P}_c^{(0)}\mathcal{V}\}$ ($i = 1, \ldots, r$). Indeed, by (6), we have the following for each k:

$$\min\left\{c \mid e_k \in \mathcal{P}_c^{(\infty)}\mathcal{V}\right\} = \frac{b}{r} - \frac{m(r-k)}{r} \tag{7}$$

The condition par-deg($\mathcal{P}_*\mathcal{V}$) = 0 is equivalent to

$$\sum_{j=1}^{r} a_j + \sum_{k=1}^{r} \left(\frac{b}{r} - \frac{m(r-k)}{r} \right) = \sum_{j=1}^{r} a_j + b - \frac{m(r-1)}{2} = 0.$$

Hence, we have par-deg($\mathcal{P}_*\mathcal{V}$) = 0 if and only if $b = -\sum a_j + m(r-1)/2$. We have

$$b - m(r - k) = -\sum a_j - \frac{m(r+1)}{2} + mk.$$
(8)

Let $\mathcal{F}_0(r,m) \subset \mathcal{F}(r,m)$ be determined by the condition that the parabolic degree is 0. We obtain the following proposition by the above consideration.

Proposition 3.17 $\mathcal{F}_0(r,m)$ naturally bijective to the following:

$$\mathfrak{R}_{r,m} := \{(a_1, \dots, a_r) \in \mathbb{R}^r \mid a_1 \ge a_2 \ge \dots \ge a_r \ge a_1 - m\}$$

In the following, let $\mathcal{P}_*^a \in \mathcal{F}_0(r,m)$ denote the filtration corresponding to $a \in \mathfrak{R}_{r,m}$.

3.1.5 Decomposition into the stable components

Let $\mathbf{a} = (a_i) \in \mathfrak{R}_{r,m}$. We set $b_i := ra_i + mi$. We define b_ℓ for any $\ell \in \mathbb{Z}$ by using the convention $b_{r+\ell} = b_\ell$. We set $r_1 := r/g.c.d.(m,r)$. We define

$$\mathcal{S}(\boldsymbol{a}, m) := \{ 1 \le j \le r \mid b_{i+j} = b_i \ (\forall i), \ j \equiv 0 \mod r_1 \}.$$

Proposition 3.18 $(\mathcal{P}^{\boldsymbol{a}}_*\mathcal{V}_r, \theta_{r,m})$ is stable, if $\mathcal{S}(\boldsymbol{a}, m) = \{r\}$.

Proof According to Proposition 3.16, it is polystable. Hence, we have only to show that any endomorphism of $(\mathcal{P}_*\mathcal{V}_r, \theta_{r,m})$ is a scalar multiplication. Let $\varphi: \mathbb{P}^1_w \longrightarrow \mathbb{P}^1_q$ be given by $\varphi(w) = w^r$. We set $u_i := w^{-mi}\varphi^*e_i$ for $i = 1, \ldots, r$. We define u_i for any $i \in \mathbb{Z}$ by the convention $u_i = u_{r+i}$. We have the parabolic structure \mathcal{P}_* on $\varphi^*\mathcal{V}_r$ obtained as the pull back of $\mathcal{P}^a_*\mathcal{V}_r$, that is $\mathcal{P}_*\varphi^*\mathcal{V}_r = \bigoplus_{i=1}^r \mathcal{O}_{\mathbb{P}^1}\left([c_0 - b_i] \cdot 0 + [c_\infty - d] \cdot \infty\right)u_i$ for any $(c_0, c_\infty) \in \mathbb{R}^2$, where $d \in \mathbb{R}$ is independent of i.

Let g be the endomorphism of $\varphi^* \mathcal{V}_r$ determined by $g(u_i) = u_{i+1}$. We have $\varphi^* \theta_{r,m} = g \cdot (rmw^{m-1}dw)$. Let τ be an n-th primitive root. Let ι_τ denote the map $\mathbb{P}^1_w \longrightarrow \mathbb{P}^1_w$ given by $\iota_\tau(w) = \tau w$. We have $\iota_\tau^* g = \tau^{-m} g$.

Any endomorphism F of $\varphi^*(\mathcal{V}_r, \theta_{r,m})$ is expressed as $F = \sum_{j=0}^{r-1} F_j(w) g^j$. If F preserves the parabolic structure at ∞ , then we obtain $F_j(w) \in \mathbb{C}[w^{-1}]$. If it preserves the parabolic structure at 0, then each $F_j g^j$ also preserves the parabolic structure at 0.

Assume $F_j \neq 0$ for some $1 \leq j \leq r-1$, and we shall derive a contradiction. Let $\ell = \deg_{z^{-1}} F_j \geq 0$. We obtain that $z^{-\ell}g^j$ preserves the parabolic structure at 0. Then, we obtain $b_i \geq b_{i+j} + \ell$ for any i, which implies that $\ell = 0$ and that $b_{i+j} = b_i$ for any i. Note that endomorphisms of $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m})$ bijectively correspond to endomorphisms F of $\varphi^*(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m})$ such that $\iota_*^*F = F$. Hence, we have $mj \equiv 0$ modulo r, which implies $j \equiv 0$

modulo r_1 , i.e., j is an element of $S(\boldsymbol{a}, m)$. It contradicts with $S(\boldsymbol{a}, m) = \{r\}$. Thus, we obtain $F_j = 0$ for any $j \neq 0$.

Let r_0 be the minimum of $\mathcal{S}(\boldsymbol{a}, m)$. If we regard $\mathcal{S}(\boldsymbol{a}, m)$ as a subset of μ_r by $j \longmapsto e^{2\pi\sqrt{-1}j/r}$, then $\mathcal{S}(\boldsymbol{a}, m)$ is a subgroup of μ_r . Hence, r_0 is a divisor of any $i \in \mathcal{S}(\boldsymbol{a}, m)$, in particular, a divisor of r. By construction, we have $r_0 m \equiv 0$ modulo r.

We put $j_0 := r/r_0 \in \mathbb{Z}$ and $m_0 := m/j_0 = mr_0/r \in \mathbb{Z}$. Let τ_0 be a primitive j_0 -th root of 1. For $i = 1, \ldots, r_0$ and $s = 0, \ldots, j_0 - 1$, we set $x_i^{(s)} := \sum_{k=0}^{j_0-1} \tau_0^{sk} q^{-km_0} e_{i+kr_0}$. Let $f \in \text{End}(\mathcal{V}_r)$ be determined by $f e = e \mathcal{K}(r, m)$. We have

$$f(x_i^{(s)}) = x_{i+1}^{(s)} \ (i = 1, \dots, r_0 - 1), \qquad f(x_{r_0}^{(s)}) = q^{m_0} \tau_0^{-s} x_1^{(s)}.$$

For $s=0,\ldots,j_0-1$, we define $\mathcal{V}^{(s)}:=\bigoplus_{j=1}^{r_0}\mathcal{O}_{\mathbb{P}^1_q}(*D)x_j^{(s)}$. We obtain a decomposition $\mathcal{V}_r=\bigoplus_{s=0}^{j_0-1}\mathcal{V}^{(s)}$ preserved by $\theta_{r,m}$. Let $\theta^{(s)}:=\theta_{r,m|\mathcal{V}^{(s)}}$. We have $\theta^{(s)}(x_i^{(s)})=x_{i+1}^{(s)}\,mdq/q$ for $i=1,\ldots,r_0-1$, and $\theta^{(s)}(x_{r_0}^{(s)})=q^{m_0}\tau_0^{-s}x_1^{(s)}\,mdq/q$. It is easy to see that there is no common eigenvalues of $\theta^{(s)}$ and $\theta^{(s')}$ if $s\neq s'$. In particular, any morphism $(\mathcal{V}^{(s)},\theta^{(s)})\longrightarrow (\mathcal{V}^{(s')},\theta^{(s')})$ is 0.

We set $\mathbf{a}_0 = (a_1, \dots, a_{r_0}) \in \mathbb{R}^{r_0}$. Because $a_{1+r_0} = a_1 - m_0$, we have $a_1 \geq \dots \geq a_{r_0} \geq a_1 - m_0$. We have the parabolic structure $\mathcal{P}^{\mathbf{a}_0}_*$ of $\mathcal{V}^{(s)}$ for the frame $\mathbf{x}^{(s)} = (x_1^{(s)}, \dots, x_{r_0}^{(s)})$. By construction, we have the following lemma.

Lemma 3.19 The parabolic structure $\mathcal{P}_*^{\mathbf{a}}\mathcal{V}_r$ is compatible with the decomposition $\mathcal{V}_r = \bigoplus \mathcal{V}^{(s)}$. The induced parabolic structure of $\mathcal{V}^{(s)}$ is the same as $\mathcal{P}_*^{\mathbf{a}_0}\mathcal{V}^{(s)}$.

For each s, we can choose a complex number β_s such that $\psi_s^*(\mathcal{P}_*^{\boldsymbol{a}_0}\mathcal{V}^{(s)},\theta^{(s)}) \simeq (\mathcal{P}_*^{\boldsymbol{a}_0}\mathcal{V}_{r_0},\theta_{r_0,m_0})$, where $\psi_s: \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ is given by $\psi_s(q) = \beta_s q$.

According to Lemma 3.19, we have the decomposition

$$(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m}) = \bigoplus_{s=0}^{j_0-1} (\mathcal{P}_*^{\boldsymbol{a}_0}\mathcal{V}^{(s)}, \theta^{(s)}). \tag{9}$$

By construction, we have $S(a_0, m_0) = \{r_0\}$. By Proposition 3.18 and the above remark, each $(\mathcal{P}^{a_0}\mathcal{V}^{(s)}, \theta^{(s)})$ is stable. We also have the description of the endomorphisms.

Proposition 3.20 Let f be as above, i.e., $f e = e \mathcal{K}(r, m)$. Then, the space of endomorphisms of $(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m})$ is $\left\{ \sum_{s=0}^{j_0-1} \beta_s \left(q^{-m_0} f^{r_0} \right)^s \mid \beta_s \in \mathbb{C} \right\}$.

Proof Let g be as in the proof of Proposition 3.18. We have $g = w^{-m}\varphi^*f$, and $g^{r_0} = \varphi^*(q^{-m_0}f^{r_0})$. Then, the claim is clear from the proof of Proposition 3.18.

3.1.6 Symmetric pairing

Let \mathcal{V}_r^{\vee} denote the dual of \mathcal{V}_r , i.e., $\mathcal{H}om_{\mathcal{O}_{\mathbb{P}^1}(*D)}\big(\mathcal{V}_r,\mathcal{O}_{\mathbb{P}^1}(*D)\big)$. It is naturally equipped with the dual $\theta_{r,m}^{\vee}$ of $\theta_{r,m}$. It is also equipped with the dual frame e_i^{\vee} $(i=1,\ldots,r)$. Let $\Psi:\mathcal{V}_r\simeq\mathcal{V}_r^{\vee}$ be given by $\Psi(e_i)=e_{r+1-i}^{\vee}$. It gives an isomorphism of meromorphic Higgs bundles $\Psi:(\mathcal{V}_r,\theta_{r,m})\simeq(\mathcal{V}_r^{\vee},\theta_{r,m}^{\vee})$. Let $f\in \operatorname{End}(\mathcal{V}_r)$ be determined by $\theta_{r,m}=f(mdq/q)$. Let $\mathbf{a}\in\mathfrak{R}_{r,m}$. Let $\mathcal{H}(\mathcal{P}_*^{\mathbf{a}}\mathcal{V}_r,\theta_{r,m})$ denote the space of morphisms $(\mathcal{P}_*^{\mathbf{a}}\mathcal{V}_r,\theta_{r,m})\longrightarrow((\mathcal{P}_*^{\mathbf{a}}\mathcal{V}_r)^{\vee},\theta_{r,m}^{\vee})$.

Proposition 3.21 We have $\mathcal{H}(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m}) \neq 0$ if and only if there exist $k_0, \ell \in \mathbb{Z}$ with $0 \leq k_0 \leq r-1$ such that the following holds:

$$a_i + a_j = \begin{cases} -\ell & (i+j=r+1-k_0) \\ -\ell - m & (i+j=2r+1-k_0) \end{cases}$$
 (10)

In that case, $\mathcal{H}(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m}) = \left\{ \sum_{s=0}^{j_0-1} \beta_s q^{\ell-sm_0} \Psi \circ f^{k_0+sr_0} \,\middle|\, \beta_s \in \mathbb{C} \right\}.$

Proof We use the symbols \mathcal{V} and θ instead of \mathcal{V}_r and $\theta_{r,m}$, respectively. We argue the "only if" part of the first claim. We put $a_i^{\vee} := -a_{r+1-i}$. They give $\mathbf{a}^{\vee} \in \mathfrak{R}_{r,m}$, and Ψ gives an isomorphism $(\mathcal{P}_*^{\mathbf{a}^{\vee}}\mathcal{V}, \theta) \simeq ((\mathcal{P}_*^{\mathbf{a}}\mathcal{V})^{\vee}, \theta^{\vee})$. Any morphism $F: (\mathcal{V}, \theta) \longrightarrow (\mathcal{V}^{\vee}, \theta^{\vee})$ is described as $F = \sum_{k=0}^{r-1} \alpha_k(q) \Psi \circ f^k$. It is easy to see that F gives a morphism $\mathcal{P}_*^{\mathbf{a}}\mathcal{V} \longrightarrow (\mathcal{P}_*^{\mathbf{a}}\mathcal{V})^{\vee}$ if and only if each $\alpha_k(q)\Psi \circ f^k$ gives a morphism $\mathcal{P}_*^{\mathbf{a}}\mathcal{V} \longrightarrow (\mathcal{P}_*^{\mathbf{a}}\mathcal{V})^{\vee}$.

Suppose that $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta)$ is stable. If $\mathcal{H}(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta) \neq 0$, by Lemma 2.3, there exist k_0 with $0 \leq k_0 \leq r-1$ such that any non-zero element of $\mathcal{H}(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta)$ is described as $\alpha(q)\Psi \circ f^{k_0}$. If $\alpha(q) \neq 0$, then $\alpha(q)\Psi \circ f^{k_0}$ has to be an isomorphism by Lemma 2.3 again. Hence, we obtain that there exists an integer ℓ such that any morphism is described as $\beta q^{\ell}\Psi \circ f^{k_0}$ for some $\beta \in \mathbb{C}$, and moreover $q^{\ell}\Psi \circ f^{k_0}$ is an isomorphism. Then, we obtain (10), and thus the "only if" part in the case that $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta)$ is stable.

Let us consider the case that $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta)$ is not necessarily stable. We have the decomposition (9) into the stable components. Any morphism $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta) \longrightarrow ((\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V})^\vee, \theta^\vee)$ is the direct sum of morphisms $(\mathcal{P}_*^{\boldsymbol{a}_0}\mathcal{V}^{(s)}, \theta^{(s)}) \longrightarrow ((\mathcal{P}_*^{\boldsymbol{a}_0}\mathcal{V}^{(s)})^\vee, (\theta^{(s)})^\vee)$. By using the "only if" part in the stable case, we obtain that there exist integers k_0 and ℓ_0 with $0 \le k_0 \le r_0 - 1$ such that the following holds for $1 \le i, j \le r_0$:

$$a_i + a_j = \begin{cases} -\ell_0 & (i+j = r_0 + 1 - k_0) \\ -\ell_0 - m_0 & (i+j = 2r_0 + 1 - k_0) \end{cases}$$

$$(11)$$

If (11) holds, (10) holds for k_0 and $\ell = \ell_0 + r - m_0$. Thus, the "only if" part is proved.

Conversely, if (10) holds for k_0 and ℓ , we can easily check that $q^{\ell}\Psi \circ f^{k_0}$ gives a morphism $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}, \theta) \longrightarrow ((\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V})^{\vee}, \theta^{\vee})$. Hence, we obtain the "if" part. Note that $q^{\ell}\Psi \circ f^{k_0}$ is an isomorphism. The second claim follows from Proposition 3.20.

The condition in Proposition 3.21 is equivalent to that the following holds for any $s \in \mathbb{Z}$ satisfying $0 \le k_0 + sr_0 \le r - 1$:

$$a_i + a_j = \begin{cases} -\ell + sm_0 & (i + j = r + 1 - (k_0 + sr_0)) \\ -\ell - m + sm_0 & (i + j = 2r + 1 - (k_0 + sr_0)) \end{cases}$$
(12)

It is also equivalent to that the following holds if $i + j - 1 + k_0 \equiv 0$ modulo r_0

$$a_i + a_j = -\ell + m - \frac{m_0}{r_0}(i + j - 1 + k_0)$$
(13)

Corollary 3.22 We have $\mathcal{H}(\mathcal{P}^a_*\mathcal{V}_r,\theta_{r,m}) \neq 0$, if and only if there exists a half-integer ν such that $\prod_{i=1}^r (T - e^{2\pi\sqrt{-1}(a_i+\nu)/m}) \in \mathbb{R}[T]$.

Proof If $\mathcal{H}(\mathcal{P}^{\boldsymbol{a}}_*\mathcal{V}_r,\theta_{r,m})\neq 0$, we obtain $\prod_{i=1}^r(T-e^{2\pi\sqrt{-1}(a_i+\nu)/m})\in\mathbb{R}[T]$ for $\nu=\ell/2$ from (10) some k_0 and ℓ . Conversely, suppose $\prod_{i=1}^r(T-e^{2\pi\sqrt{-1}(a_i+\nu)/m})\in\mathbb{R}[T]$ for a half-integer ν . We put $c_i:=a_i+\nu$. We may assume that $0\geq c_1\geq\cdots c_r>-m$ by a successive use of the gauge transform and the inverse transform in the proof of Lemma 3.6. There exists s such that $0=c_1=\cdots=c_s>c_{s+1}$. We have $c_{s+j}+c_{r+1-j}=-m$ for $j=1,\ldots,r-s$. By Proposition 3.21, we have $\mathcal{H}(\mathcal{P}^{\boldsymbol{a}}_*\mathcal{V}_r,\theta_{r,m})\neq 0$.

For $\mathcal{O}_{\mathbb{P}^1}(*D)$ -modules \mathcal{F}_i (i=1,2), a pairing $\mathcal{F}_1 \times \mathcal{F}_2 \longrightarrow \mathcal{O}_{\mathbb{P}^1}(*D)$ means a bi- $\mathcal{O}_{\mathbb{P}^1}(*D)$ -homomorphism. The natural pairing $\mathcal{V}_r \times \mathcal{V}_r^{\vee} \longrightarrow \mathcal{O}_{\mathbb{P}^1}(*D)$ is denoted by $\langle \cdot, \cdot \rangle$. A pairing $C: \mathcal{V}_r \times \mathcal{V}_r \longrightarrow \mathcal{O}_{\mathbb{P}^1}(*D)$ induces a morphism $\Phi_C: \mathcal{V}_r \longrightarrow \mathcal{V}_r^{\vee}$ by $\langle \Phi_C(u), v \rangle = C(u, v)$ for local sections u, v of \mathcal{V}_r . We have $C(\theta_{r,m}u, v) = C(u, \theta_{r,m}v)$ if and only if $\Phi_C \circ \theta_{r,m} = \theta_{r,m}^{\vee} \circ \Phi_C$.

Let C_k denote the pairing of $\mathcal{V}_r \times \mathcal{V}_r \longrightarrow \mathcal{O}_{\mathbb{P}^1}(*D)$ given by $C_k(u,v) := \langle \Psi \circ f^k(u), v \rangle$. For $0 \leq k \leq r-1$, we have the following:

$$C_k(e_i, e_j) = \begin{cases} 1 & (i+j = r+1-k) \\ q^m & (i+j = 2r+1-k) \\ 0 & \text{otherwise} \end{cases}$$

In particular, C_k are symmetric, i.e., $C_k(u, v) = C_k(v, u)$.

Let $\mathfrak{P}(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m})$ denote the space of pairings $C: \mathcal{V}_r \times \mathcal{V}_r \longrightarrow \mathcal{O}_{\mathbb{P}^1}(*D)$ such that Φ_C gives a morphism $(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m}) \longrightarrow ((\mathcal{P}_*^a \mathcal{V}_r)^{\vee}, \theta_{r,m}^{\vee})$.

Corollary 3.23 We have $\mathfrak{P}(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m}) \neq 0$, if and only if there exist integers k_0 and ℓ with $0 \leq k_0 \leq r-1$ such that (10) holds. In that case, (13) holds if $i+j-1+k_0 \equiv 0$ modulo r_0 , and we have

$$\mathfrak{P}(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m}) = \Big\{ \sum_{s=0}^{j_0-1} \beta_s q^{\ell-sm_0} C_{k_0+sr_0} \, \Big| \, \beta_s \in \mathbb{C} \Big\}.$$

In particular, any $C \in \mathfrak{P}(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m})$ satisfies $t^*C = t^{r\ell - m(r+1-k_0)}C$.

Proof The first two claims follows from Proposition 3.21. The last claim can be checked directly.

Corollary 3.24 Suppose $(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m})$ is stable. We have $\mathfrak{P}(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m}) \neq 0$, if and only if there exist integers k_0 and ℓ with $0 \leq k_0 \leq r-1$ such that (10) holds. In that case, we have $\mathfrak{P}(\mathcal{P}_*^a \mathcal{V}_r, \theta_{r,m}) = \{\beta q^{\ell} C_{k_0} \mid \beta \in \mathbb{C}\}$.

As remarked in Corollary 3.22, the existence of k_0 and ℓ as above can be replaced with the existence of a half-integer ν such that $\prod_{i=1}^r (T - e^{2\pi\sqrt{-1}(a_i + \nu)/m}) \in \mathbb{R}[T]$.

3.2 Toda-like Harmonic bundle

3.2.1 Preliminaries

We set $E_r := \mathcal{V}_{r|\mathbb{P}^1 \setminus D}$, which is equipped with the base $e_{i|\mathbb{P}^1 \setminus D}$ and the Higgs field $\theta_{r,m|\mathbb{P}^1 \setminus D}$. We will omit " $|\mathbb{P}^1 \setminus D$ ". Let h be any harmonic metric of $(E_r, \theta_{r,m})$. Clearly, $(E_r, \theta_{r,m}, h)$ is tame at 0, and wild at ∞ . We have the associated good filtered Higgs bundle denoted by $(\mathcal{P}_*^h E_r, \theta_{r,m})$ and the meromorphic Higgs bundle $(\mathcal{P}^h E_r, \theta_{r,m})$. (We add the superscript h to emphasize the dependence on h.)

3.2.2 Toda-like harmonic metric

In this paper, a harmonic metric h of $(E_r, \theta_{r,m})$ is called Toda-like, if $h(e_i, e_j) = 0$ for $i \neq j$. In that case, $(E_r, \theta_{r,m}, h)$ is called a Toda-like harmonic bundle. Clearly, the condition is equivalent to $\sigma^* h = h$. The following lemma is clear.

Lemma 3.25 If h is a Toda-like harmonic metric of $(E_r, \theta_{r,m})$, then σ gives an automorphism of \mathcal{P}^h_*E .

A Toda-like harmonic metric is uniquely determined by the parabolic structure in the sense of the following lemma.

Lemma 3.26 Let h_i (i = 1, 2) be Toda-like harmonic metrics of $(E_r, \theta_{r,m})$ such that $\mathcal{P}_*^{h_1}E = \mathcal{P}_*^{h_2}E$. Then, $h_1 = a h_2$ for some a > 0.

Proof We have an automorphism g of $(\mathcal{P}_*^{h_1}E_r, \theta_{r,m})$ such that (i) g is self-adjoint with respect to both of h_i (i = 1, 2), (ii) $h_1 = h_2 g$. Because $\sigma^* h_i = h_i$, we have $\sigma \circ g = g \circ \sigma$. We also have $g \circ \theta = \theta \circ g$. Then, it is easy to prove that g = a id $_{E_r}$.

3.2.3 Adaptedness

A harmonic metric h of $(E_r, \theta_{r,m})$ is called adapted to \mathcal{V}_r , if $\mathcal{P}^h E_r = \mathcal{V}_r$. The case m = 1 is special.

Proposition 3.27 Any harmonic metric of $(E_r, \theta_{r,1})$ adapted to \mathcal{V}_r is Toda-like.

Proof Because $(E_r, \theta_{r,1})$ is irreducible, the filtered Higgs bundle $(\mathcal{P}_*^h E_r, \theta_{r,1})$ is stable. Because $\sigma^*\theta = \tau\theta$ with $|\tau| = 1$, σ^*h is also a harmonic metric of $(E_r, \theta_{r,1})$. By Proposition 3.14, we have $\mathcal{P}_*^{\sigma^*h} E_r \simeq \sigma^* \mathcal{P}_*^h E_r \simeq \mathcal{P}_*^h E_r$. Hence, we have $\sigma^*h = ah$ for some a > 0 by the uniqueness in Theorem 2.7. Because we clearly have $\det(\sigma^*h) = \det(h)$, we obtain $\sigma^*h = h$.

We give a complement. We set $\Omega := e_1 \wedge \cdots \wedge e_r$.

Lemma 3.28 A harmonic metric h of $(E_r, \theta_{r,m})$ is adapted to \mathcal{V}_r , if and only if $|e_i|_h = O((|q| + |q|^{-1})^N)$ for some i and some N. Suppose moreover that h is Toda-like. Then, h is adapted to \mathcal{V}_r , if and only if $|\Omega|_h = O((|q| + |q|^{-1})^N)$ for some N.

Proof If h is adapted, any e_i are sections of $\mathcal{P}^h E_r$, and hence we have $|e_i|_h = O((|q| + |q|^{-1})^N)$ for some N. Conversely, assume $|e_1|_h = O((|q| + |q|^{-1})^N)$. For the expression $\theta = f \cdot (m \, dq/q)$, f naturally gives an endomorphism of $\mathcal{P}^h E_r$, according to [5], [36] and [44]. Because $e_j = f^{j-1}(e_1)$ for $j = 1, \ldots, r$, we obtain that e_j are also sections of $\mathcal{P}^h E_r$. Then, we obtain $\mathcal{V}_r \subset \mathcal{P}^h E_r$, which implies $\mathcal{V}_r = \mathcal{P}^h E_r$ because both of them are $\mathcal{O}_{\mathbb{P}^1}(*D)$ -locally free. Because $f(e_i) = e_{i+1}$ $(i \leq r-1)$ and $f(e_r) = q^m e_1$, we may replace the role of e_1 with the other e_i . Hence, we obtain the first claim.

Let us prove the second claim. The "only if" part is clear. Assume that h is Toda-like, and that $|\Omega|_h = O((|q| + |q|^{-1})^N)$ for some N. We have $|\Omega|_h = C \prod_{i=1}^r |e_i|_h$ for some C > 0. By the relation $e_r = f^{r-j}(e_j)$, we have $|e_j|_h \ge (|q| + |q|^{-1})^{-N} |e_r|_h$. Hence, $|e_r|_h^r \le (|q| + |q|^{-1})^M$ for some M. We obtain that h is adapted.

It is easy to see that, if h is adapted, we have $|\Omega|_h = C|q|^a$ for some C > 0 and $a \in \mathbb{R}$. Indeed, Ω is a section of $\mathcal{P}\det(E)$, and we have $\partial_q \partial_{\overline{q}} \log |\Omega|_h = 0$. Because the parabolic degree of $\mathcal{P}_* \det(E)$ is 0, there exists $a \in \mathbb{R}$ such that $\log |\Omega|_h - a \log |q|$ is bounded from above on $\mathbb{P}^1 \setminus D$. Then, we obtain that $\log |\Omega|_h - a \log |q|$ is constant.

Corollary 3.29 Let $(E_r, \theta_{r,m}, h)$ be any Toda-like harmonic bundle. There exists a holomorphic function g on \mathbb{C}^* such that the Toda-like harmonic metric $e^{\operatorname{Re} g}h$ of $(E_r, \theta_{r,m})$ is adapted to \mathcal{V}_r .

Proof We have a holomorphic function g_1 on \mathbb{C}^* such that (i) g_1 is nowhere vanishing on \mathbb{C}^* , (ii) $g_1 \Omega$ is a section of $\det \mathcal{P}^h E$. It means $\det h(\Omega, \Omega)|g_1|^2 = (|q| + |q|^{-1})^N$ for some N. We have an integer ℓ such that $\int_{|q|=1} (dg_1/g_1 + \ell dq/q) = 0$. We can find a holomorphic function g_2 on \mathbb{C}^* such that $e^{g_2} = g_1 q^{\ell}$. We obtain that $e^{2\operatorname{Re}(g_2)} \det h(\Omega, \Omega) = O((|q| + |q|^{-1})^N)$ for some N. Hence, $e^{2\operatorname{Re}(g_2)/r}h$ is adapted to \mathcal{V}_r , according to Lemma 3.28.

3.2.4 Homogeneity and μ_m -equivariance

We have the symmetry of Toda-like harmonic bundles, inherited to the symmetry of the underlying filtered Higgs bundles. By the restriction of the \mathbb{C}^* -action ρ , we obtain the S^1 -action on E_r , i.e., $\rho: S^1 \times \mathbb{C}^* \longrightarrow \mathbb{C}^*$ is given by $(t,q) \longmapsto t^r q$, and $\widetilde{t}: t^*E_r \simeq E_r$ is given by $\widetilde{t}(t^*e_i) = t^{im}e_i$. We have $t^*\theta = t^m\theta$.

Lemma 3.30 Let h be a Toda-like harmonic metric of $(E_r, \theta_{r,m})$ adapted to \mathcal{V}_r . Then, $t^*h = h$ for $t \in S^1$.

Proof According to Lemma 3.25, the conditions in Proposition 3.13 hold. In particular, \mathcal{P}_*E_r is \mathbb{C}^* -equivariant. We clearly have $\det(t^*h) = \det(h)$ for any $t \in S^1$. Because t^*h is also Toda-like, the claim follows from Lemma 3.26.

We have the natural action of μ_m on $(E_r, \theta_{r,m})$ given by $\widetilde{\kappa}(\kappa^* e_i) = e_i$ for $\kappa \in \mu_m$. The following lemma follows from Lemma 3.30, or it can be also proved easily by the argument in the proof of Lemma 3.30.

Lemma 3.31 Any Toda-like harmonic metric of $(E_r, \theta_{r,m})$ is μ_m -equivariant, if it is adapted to \mathcal{V}_r .

3.2.5 Correspondence

Let $a \in \mathbb{R}$. Let $\mathfrak{H}_{r,m}(a)$ be the set of Toda-like harmonic metrics of $(E_r, \theta_{r,m})$ such that $|\Omega|_h = |q|^{-a}$. Let $\mathfrak{R}_{r,m}(a)$ be the set of $a = (a_i) \in \mathbb{R}^r$ such that $a_1 \geq a_2 \geq \cdots \geq a_r \geq a_1 - m$ and $\sum a_i = a$. From $h \in \mathfrak{H}_{r,m}(a)$, we obtain $\Phi_{r,m}(h) = (a_i(h)) \in \mathbb{R}^r$ such that $a_i(h) := \min\{c \mid e_i \in \mathcal{P}_c^h E \text{ at } 0\}$. They satisfy $a_i(h) \geq a_{i+1}(h)$, $a_r(h) \geq a_1(h) - m$ and $\sum a_i(h) = a$. Thus, we obtain a map $\Phi_{r,m} : \mathfrak{H}_{r,m}(a) \longrightarrow \mathfrak{R}_{r,m}(a)$.

Theorem 3.32 $\Phi_{r,m}$ is bijective.

Proof It is injective by Lemma 3.26. Let us consider the case m=1. Let $\mathbf{a} \in \mathfrak{R}_{r,1}(a)$. Let $(\mathcal{P}_*^{\mathbf{a}}\mathcal{V}_r, \theta_{r,1})$ be the associated filtered Higgs bundle. It is always stable. By the Kobayashi-Hitchin correspondence, we have a unique harmonic metric h of $(E_r, \theta_{r,1})$ such that $\mathcal{P}_*^h E_r \simeq \mathcal{P}_*^a \mathcal{V}_r$ and $|\Omega|_h = |q|^{-a}$. It is automatically Toda-like as remarked in Proposition 3.27. It means that $\Phi_{r,1}$ is a bijection.

Let us consider the general case. Let $\mathbf{a} \in \mathfrak{R}_{r,m}(a)$. Let $(\mathcal{P}_*^{\mathbf{a}}\mathcal{V}, \theta_{r,m})$ be the associated filtered Higgs bundle. It is μ_m -equivariant. Any Toda-like harmonic metrics are also μ_m -equivariant as remarked in Lemma 3.31. Then, the bijectivity of $\Phi_{r,m}$ follows from the bijectivity of $\Phi_{r,1}$. Thus, we obtain Theorem 3.32.

Asymptotic behaviour around 0 and ∞ We have the norm estimate for wild harmonic bundles ([5], [31], [36], [44]). In our situation, it is described as follows. Suppose that $h \in \mathfrak{H}_{r,m}(a)$ corresponds to $\mathbf{a} \in \mathfrak{R}_{r,m}(a)$. We obtain the following around ∞ from (7) and (8):

$$\log|e_i|_h + \frac{1}{r}\left(a + \frac{m(r+1)}{2} - mi\right) \times \log|q| = O(1)$$
(14)

Let us describe the asymptotic behaviour of $\log |e_i|_h$ around 0. For $c \in \mathbb{R}$, we set $S(\boldsymbol{a}, c) = \{i \mid a_i = c\}$. First, let us consider the case $a_r > a_1 - m$. For i, we define

$$k_i := |S(\mathbf{a}, a_i)| - 2|\{j \mid a_j = a_i, j \le i\}| + 1.$$
(15)

Then, we have the following around 0:

$$\log|e_i|_h + a_i \log|q| - \frac{k_i}{2} \log(-\log|q|) = O(1)$$
(16)

Let us consider the case $a_r = a_1 - m$. If $a_i \neq a_j$ (j = 1, r), then the behaviour of $\log |e_i|_h$ around 0 is given by (16). We have j_0 such that $a_{j_0} > a_{j_0+1} = a_r$. We have j_1 such that $a_1 = a_{j_1} > a_{j_1+1}$. We put

$$S'(\boldsymbol{a}, a_1) := S(\boldsymbol{a}, a_1 - 1) \cup S(\boldsymbol{a}, a_1) = \{j_0 + 1, j_0 + 2, \dots, r, 1, \dots, j_1\}.$$

Let \leq' be the order given by $j_0+1, j_0+2, \ldots, r, 1, \ldots, j_1$. For $i \in S'(\boldsymbol{a}, a_1)$, we put

$$k'_i := |S'(\boldsymbol{a}, a_1)| - 2|\{j \in S'(\boldsymbol{a}, a_1) | j \le i\}| + 1.$$

Then, for $i \in S'(\boldsymbol{a}, a_1)$, we have the following around 0:

$$\log|e_i|_h + a_i \log|q| - \frac{k_i'}{2} \log(-\log|q|) = O(1)$$
(17)

3.2.6 Additional symmetry

We use the notation in §3.1.5–3.1.6. Let $\mathbf{a} \in \mathfrak{R}_{r,m}(a)$. Suppose $\mathfrak{P}(\mathcal{P}_*^{\mathbf{a}}\mathcal{V}_r, \theta_{r,m}) \neq 0$. Let k_0 and ℓ be as in Proposition 3.21. See also the remark right after the proposition.

Proposition 3.33 Let $h \in \mathfrak{H}_{r,m}(a)$ be the adapted Toda-like harmonic metric of $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m})$. If $i+j+k_0-1 \equiv 0$ modulo r_0 , then we have

$$\log|e_i|_h + \log|e_j|_h = \left(\ell - m + \frac{m_0}{r_0}(i+j+k_0-1)\right)\log|q|. \tag{18}$$

Proof Let h^{\vee} denote the induced Toda-like harmonic metric of $(E_r^{\vee}, \theta_{r,m}^{\vee})$, which is adapted to $(\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r)^{\vee}$. Recall that we have an isomorphism $\Phi := q^{\ell-sm_0}\Psi \circ f^{k_0+sr_0} : (\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r, \theta_{r,m}) \longrightarrow ((\mathcal{P}_*^{\boldsymbol{a}}\mathcal{V}_r)^{\vee}, \theta_{r,m}^{\vee})$. By the uniqueness of the Toda-like harmonic metric, $\Phi : (E_r, h) \longrightarrow (E_r^{\vee}, h^{\vee})$ is isometric up to the multiplication of the positive constant. We have $\det \Phi(e_1 \wedge \cdots \wedge e_r) = \pm q^{b_1} e_1^{\vee} \wedge \cdots \wedge e_r^{\vee}$ for some $b_1 \in \mathbb{Z}$. We have $|e_1^{\vee} \wedge \cdots \wedge e_r^{\vee}|_{h^{\vee}} = |q|^{b_2}$ for some $b_2 \in \mathbb{Z}$. Hence, we obtain that $\det \Phi : \det(E_r, h) \longrightarrow \det(E_r, h)^{\vee}$ is an isometry, and thus Φ is an isometry. It implies

$$h(e_i, e_i) = \begin{cases} h^{\vee}(e_j^{\vee}, e_j^{\vee})|q|^{2(\ell - s m_0)} & (i + j = r + 1 - (k_0 + s r_0)) \\ h^{\vee}(e_j^{\vee}, e_j^{\vee})|q|^{2(\ell + m - s m_0)} & (i + j = 2r + 1 - (k_0 + s r_0)). \end{cases}$$

Thus, we obtain (18).

3.3 Toda lattice with opposite sign

According to [13], the Toda lattice with opposite sign is the following equation for \mathbb{R}^r -valued function $\mathbf{w} = (w_i | i = 1, ..., r)$ on \mathbb{C}^* :

$$2\overline{\partial}_z \partial_z w_i - e^{2(w_i - w_{i-1})} + e^{2(w_{i+1} - w_i)} = 0$$

Here, we use the convention $w_{r+i} := w_i$ for any i. By the coordinate change z = rq, it is transformed to

$$2\overline{\partial}_{q}\partial_{q}w_{i} - r^{2}e^{2(w_{i} - w_{i-1})} + r^{2}e^{2(w_{i+1} - w_{i})} = 0$$
(19)

Recall that the solutions of the equation (19) naturally correspond to the Toda-like harmonic metrics of the Higgs bundle $(E_r, \theta_{r,r})$. Let h be a Toda-like harmonic metric of $(E_r, \theta_{r,r})$. We set $u_i := q^{-(i-1)}e_i$. Note that θ is represented by A dq, where $A_{ij} = r$ if $i - j \equiv 1$ modulo r, and $A_{ij} = 0$ otherwise. Let w_i be the functions determined by $e^{2w_i} = h(u_i, u_i)$. Then, it is direct to check that $(w_i | i = 1, ..., r)$ satisfies the equation (19). Conversely, if we are given a solution of $(w_i | i = 1, ..., r)$ of (19), we define the metric h by $h(u_i, u_i) := e^{2w_i}$ and $h(u_i, u_i) = 0$ $(i \neq j)$. Then, h is a Toda-like harmonic metric of $(E_r, \theta_{r,r})$.

Let $\mathfrak{S}_r(a)$ be the set of solutions $\boldsymbol{w}=(w_i)$ of (19) such that

$$\sum_{i=1}^{r} w_i = \left(-\frac{r(r-1)}{2} - a\right) \log|q|.$$

We obtain the following theorem from Theorem 3.32 and Lemma 3.30.

Theorem 3.34 We have a natural bijection $\mathfrak{S}_r(a)$ and $\mathfrak{R}_{r,r}(a)$. For any $\mathbf{w} \in \mathfrak{S}_r(a)$, the functions w_i depend only on |q|.

Let $\mathbf{w} = (w_i)$ be any solution of the equation (19). Then, $\sum w_i$ is an \mathbb{R} -valued harmonic function on \mathbb{C}^* . Hence, it is described as $A \log |z|^2 + \operatorname{Re} f$ for a real number A and a holomorphic function f. If we put $\widetilde{w}_i := w_i - \operatorname{Re}(f)/r$, then $\widetilde{\mathbf{w}}_i$ is a solution of (19) such that $\sum \widetilde{w}_i = A \log |z|^2$. In this sense, Theorem 3.34 gives a classification of the \mathbb{R} -valued solutions of (19).

Asymptotic behaviour Let $\mathbf{a} \in \mathfrak{R}_{r,r}(a)$. Let $\mathbf{w} \in \mathfrak{S}_r(a)$ and $h \in \mathfrak{H}_{r,r}(a)$ corresponding to \mathbf{a} . Let us describe the asymptotic behaviour of \mathbf{w} . Note $w_i = \log |u_i|_h = \log |e_i|_h - (i-1) \log |q|$. We obtain the following around $q = \infty$ from (14):

$$w_i = \log|q| \times \left(-\frac{a}{r} - \frac{r-1}{2}\right) + O(1)$$

Let k_i be as in (15). Around q=0, if $a_r>a_1-r$, we obtain the following for any $i=1,\ldots,r$:

$$w_i = -(a_i + i - 1)\log|q| + \frac{k_i}{2}\log(-\log|q|) + O(1)$$

If $a_r = a_1 - r$, the numbers k_i should be modified to k'_i as in (17).

Additional symmetry Let $\mathbf{w} \in \mathfrak{S}_r(a)$ corresponding to $\mathbf{a} \in \mathfrak{R}_{r,r}$. Suppose that there exist integers k_0 and ℓ with $0 \le k_0 \le r - 1$ such that the following holds:

$$a_i + a_j = \begin{cases} -\ell & (i+j=r+1-k_0) \\ -\ell - r & (i+j=2r+1-k_0) \end{cases}$$
 (20)

We use the notation in $\S 3.1.5 - 3.1.6$.

Theorem 3.35 $w_i + w_j = (\ell - r + 1 + k_0) \log |q|$ if $i + j + k_0 - 1 \equiv 0$ modulo r_0 .

Proof Let $(E_r, \theta_{r,r}, h)$ be the corresponding Toda-like harmonic bundle. If $i + j + k_0 - 1 \equiv 0$ modulo r_0 , we have $w_i + w_j = \log|e_i|_h - (i-1)\log|q| + \log|e_j|_h - (j-1)\log|q| = (\ell - m + k_0 + 1)\log|q|$.

We state some special cases as corollaries.

Corollary 3.36 Assume that there exist an integer ℓ satisfying the following:

$$a_i + a_j = \begin{cases} -\ell & (\text{if } i + j = 2 + \ell) \\ -\ell - r & (\text{if } i + j = 2 + \ell + r) \end{cases}$$
 (21)

Then, we have $w_i + w_j = 0$ $(i + j \equiv 2 + \ell \text{ modulo } r)$. In particular, if $a_i + a_j = -r + 1$ holds for (i, j) with i + j = r + 1, we have $w_i + w_j = 0$.

Part II

4 Stokes factor and integral structure

4.1 Preliminary

4.1.1 Stokes structure

Let us recall the concept of Stokes structure in the local and unramified case. Let $X := \{z \in \mathbb{C} \mid |z| < 1\}$ and $D := \{0\}$. Let (V, ∇) be a meromorphic flat bundle on (X, D). For simplicity, we assume that it is unramified, i.e., its formal completion $(V, \nabla) \otimes_{\mathcal{O}_X} \mathbb{C}[\![z]\!]$ has a formal decomposition

$$(V, \nabla) \otimes_{\mathcal{O}_X} \mathbb{C}[\![z]\!] = \bigoplus_{\mathfrak{a} \in z^{-1} \mathbb{C}[z^{-1}]} (\widehat{V}_{\mathfrak{a}}, \widehat{\nabla}_{\mathfrak{a}}), \tag{22}$$

where $\widehat{\nabla}_{\mathfrak{a}} - d\mathfrak{a}$ are regular singular. We set $\mathcal{I}(\nabla) := \{ \mathfrak{a} \in z^{-1} \mathbb{C}[z^{-1}] \mid \widehat{V}_{\mathfrak{a}} \neq 0 \}$.

Remark 4.1 In general, according to the Hukuhara-Levelt-Turrittin theorem, we have such a decomposition of $(V, \nabla) \otimes_{\mathcal{O}_X} \mathbb{C}[\![z^{1/e}]\!]$.

In this paper, a sector $S \subset X \setminus D$ means a subset of the form $\{z \in X \setminus D \mid 0 < |z| < R_0, \ \theta_0 \le \arg(z) \le \theta_1\}$. We set $\partial S = \{z \in S \mid \arg(z) = \theta_0 \text{ or } \theta_1\}$. For each $\mathfrak{a}, \mathfrak{b} \in \mathcal{I}(V)$ with $\mathfrak{a} \ne \mathfrak{b}$, put $S(\mathfrak{a}, \mathfrak{b}) := \{z \in S \mid \operatorname{Re}(\mathfrak{a} - \mathfrak{b})(z) = 0\}$. A sector S is called adapted to (V, ∇) , if the following holds for each $\mathfrak{a}, \mathfrak{b} \in \mathcal{I}(V)$ with $\mathfrak{a} \ne \mathfrak{b}$:

• $S(\mathfrak{a},\mathfrak{b})$ is connected or empty, and we have $S(\mathfrak{a},\mathfrak{b})\cap\partial S=\emptyset$.

Such a sector is called strictly adapted to (V, ∇) , if moreover each $S(\mathfrak{a}, \mathfrak{b})$ is non-empty.

According to the classical asymptotic analysis for meromorphic flat bundles on curves, if S is adapted, then there exists a flat decomposition

$$(V, \nabla)_{|S} = \bigoplus_{\mathfrak{a} \in \mathcal{I}(V)} (V_{\mathfrak{a}, S}, \nabla_{\mathfrak{a}, S})$$
(23)

which is asymptotic to (22). If S is strictly adapted, then the decomposition (23) is unique.

A flat frame $\mathbf{y} = (y_i)$ of $(V, \nabla)_{|S}$ is called adapted, if it is compatible with a decomposition (23) in the sense that, for each y_i , there exists $\mathfrak{a} \in \mathcal{I}(V)$ such that $y_i \in V_{\mathfrak{a},S}$. Let $S^{(i)}$ (i = 1,2) be adapted sectors such that $S^{(1)} \cap S^{(2)} \neq \emptyset$. Suppose we are given adapted flat frames $\mathbf{y}^{(i)}$ of $(V, \nabla)_{|S^{(i)}}$. We obtain the matrix A determined by $\mathbf{y}^{(1)} = \mathbf{y}^{(2)}A$. The matrix A is called the Stokes matrix or Stokes factor with respect to $\mathbf{y}^{(i)}$ (i = 1, 2). (If $S^{(1)} \cap S^{(2)}$ is not connected, we consider the matrix for each connected component.)

Remark 4.2 Although we fix a coordinate z, the above notions are independent of z. It is better to replace the index set $z^{-1}\mathbb{C}[z^{-1}]$ with $\mathcal{O}_X(*D)/\mathcal{O}_X$.

Remark 4.3 For a general meromorphic flat bundle on (X, D), we take an appropriate ramified covering $\varphi: (X, D) \longrightarrow (X, D)$ given by $\varphi(z) = z^e$ such that the pull back $\varphi^*(V, \nabla)$ is unramified, for which the Stokes structure is considered as above.

Remark 4.4 Let C be a general complex curve with a discrete subset $Z \subset C$. Let (V, ∇) be a meromorphic flat bundle on (C, Z). We can consider the Stokes structure of (V, ∇) on a neighbourhood of each point of Z.

Lift of a formal frame Let (V, ∇) be an unramified meromorphic flat bundle with the decomposition (23). For each $\mathfrak{a} \in \mathcal{I}(V)$, let $r(\mathfrak{a}) := \operatorname{rank}(\widehat{V}_{\mathfrak{a}})$. We have a frame $\widehat{\boldsymbol{v}}_{\mathfrak{a}}$ of $\widehat{V}_{\mathfrak{a}}$ with the following property:

$$abla \widehat{oldsymbol{v}}_{\mathfrak{a}} = \widehat{oldsymbol{v}}_{\mathfrak{a}} \Big(d \mathfrak{a} I_{\mathfrak{a}} + A_{\mathfrak{a}} rac{dz}{z} \Big)$$

Here, $I_{\mathfrak{a}}$ denote the $r(\mathfrak{a})$ -th identity matrix, and $A_{\mathfrak{a}} \in M_{r(\mathfrak{a})}(\mathbb{C})$. They give a frame $\widehat{\boldsymbol{v}} = (\widehat{\boldsymbol{v}}_{\mathfrak{a}} \mid \mathfrak{a} \in \mathcal{I}(V))$ of $V \otimes \mathbb{C}[\![z]\!]$ for which we have $\nabla \widehat{\boldsymbol{v}} = \widehat{\boldsymbol{v}} \Big(\bigoplus (d\mathfrak{a}I_{\mathfrak{a}} + A_{\mathfrak{a}}dz/z) \Big)$. Recall the following.

Lemma 4.5 Let S be an adapted sector. Take a flat decomposition (23). We have a holomorphic frame $\mathbf{v}_{\mathfrak{a},S}$ of $V_{\mathfrak{a},S}$ such that (i) $\mathbf{v}_{\mathfrak{a},S}$ is asymptotic to $\widehat{\mathbf{v}}_{\mathfrak{a}}$, (ii) $\nabla_{\mathfrak{a},S}\mathbf{v}_{\mathfrak{a},S} = \mathbf{v}_{\mathfrak{a},S}(d\mathfrak{a}I_{\mathfrak{a}} + A_{\mathfrak{a}}dz/z)$. Such $\mathbf{v}_{\mathfrak{a},S}$ is uniquely determined, once we fix the decomposition (23).

In particular, if S is strictly adapted, there uniquely exists a frame v_S of $V_{|S}$ such that (i) v_S is asymptotic to \hat{v} , (ii) $\nabla v_S = v_S \left(\bigoplus (d\mathfrak{a} I_{\mathfrak{a}} + A_{\mathfrak{a}} dz/z) \right)$. If we choose a branch of $\log z$ on S, we obtain an adapted flat frame

$$\widetilde{\boldsymbol{v}}_S := \boldsymbol{v}_S \, \exp\Bigl(\bigoplus \bigl(-\mathfrak{a} \, I - A_{\mathfrak{a}} \log z\bigr)\Bigr).$$
 (24)

Pure slope Let (V, ∇) be an unramified meromorphic flat bundle on (X, D) with the decomposition (23). Let d be a positive integer. In this paper, we say that (V, ∇) has pure slope d, if (i) $\deg_{z^{-1}}(\mathfrak{a}) = d$ for each $\mathfrak{a} \in \mathcal{I}(\nabla)$, (ii) $\deg_{z^{-1}}(\mathfrak{a} - \mathfrak{b}) = d$ for each pair $\mathfrak{a}, \mathfrak{b} \in \mathcal{I}(\nabla)$ with $\mathfrak{a} \neq \mathfrak{b}$. We shall use the following lemma implicitly.

Lemma 4.6 Suppose that (V, ∇) has pure slope d. Let S be a sector adapted to (V, ∇) whose angle is larger than π/d . Then, S is strictly adapted. In particular, there exist strictly adapted sectors S_i (i = 1, ..., N) such that $\bigcup S_i = X \setminus D$.

Let (V, ∇) be a meromorphic flat bundle on (X, D), which is not necessarily unramified. Let μ be a positive rational number. In this paper, we say that (V, ∇) has pure slope μ , if there exists a ramified covering $\varphi: (X, D) \longrightarrow (X, D), \ \varphi(z) = z^e$ such that $\varphi^*(V, \nabla)$ is unramified with pure slope $e\mu \in \mathbb{Z}$.

4.1.2 *R*-structure

Let X and D be as in §4.1.1. Let (V, ∇) be a meromorphic flat bundle on (X, D). Let \mathcal{L} be the local system on $X \setminus D$ associated to (V, ∇) , i.e., it is the local system of flat sections of $(V, \nabla)_{|X\setminus D}$. Let R be a subring of \mathbb{C} . (We will be mainly interested in the cases that R is \mathbb{Z} or \mathbb{Q} .) An R-structure of $(V, \nabla)_{|X\setminus D}$ is an R-local system \mathcal{L}_R with an isomorphism $\mathcal{L}_R \otimes_R \mathbb{C} \simeq \mathcal{L}$. In this paper, an R-local system means a locally constant sheaf of finitely generated projective R-modules. We shall often regard \mathcal{L}_R as a subsheaf of \mathcal{L} . We recall a compatibility condition of R-structure with Stokes structure in the case that (V, ∇) has pure slope.

Definition 4.7 Let (V, ∇) be a meromorphic flat bundle on (X, D) such that it is unramified with pure slope. We say that an R-structure of $(V, \nabla)_{|X \setminus D}$ is compatible with the Stokes structure, if the following holds:

• Let S be a sector strictly adapted to (V, ∇) . Then the decomposition (23) of $(V, \nabla)_{|S}$ is induced by a decomposition of $\mathcal{L}_{R|S}$.

In that case, the R-structure of $(V, \nabla)_{|X \setminus D}$ is called an R-structure of (V, ∇) .

For each $\mathfrak{a} \in \mathcal{I}(\nabla)$, we can take a meromorphic flat bundle $\operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V,\nabla)$ on (X,D) with an isomorphism $\operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V,\nabla) \otimes_{\mathcal{O}_X} \mathbb{C}[\![z]\!] \simeq (\widehat{V}_{\mathfrak{a}},\widehat{\nabla}_{\mathfrak{a}})$, where $(\widehat{V}_{\mathfrak{a}},\widehat{\nabla}_{\mathfrak{a}})$ are as in (22). Recall that an R-structure of (V,∇) induces an R-structure of $\operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V,\nabla)$. Indeed, for any strictly adapted sectors S, (23) gives a unique isomorphism $(V,\nabla)_{|S} \simeq \bigoplus \operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V,\nabla)_{|S}$, which induces an R-structure of $\operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V,\nabla)_{|S}$. It is easy to observe that the R-structures can be glued when the strictly adapted sectors are varied.

Lemma 4.8 Let (V, ∇) be a meromorphic flat bundle on (X, D) which is unramified with pure slope. The following conditions are equivalent.

- (V, ∇) has an R-structure \mathcal{L}_R such that the induced R-structure $\mathcal{L}_{\mathfrak{a},R}$ of $\mathrm{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V, \nabla)$ are local systems of free R-modules.
- There exists a frame \hat{v} of $V \otimes \mathbb{C}[\![z]\!]$ such that, for any strictly adapted sectors S_i (i = 1, 2) with $S_1 \cap S_2 \neq \emptyset$, the entries of the Stokes matrix with respect to \tilde{v}_{S_i} (i = 1, 2) are elements of R. (See (24) for \tilde{v}_{S_i} .)
- We can associate an adapted flat frame $\widetilde{\boldsymbol{v}}_S$ of $(V,\nabla)_{|S}$ to each strictly adapted sector S such that, if $S \cap S' \neq \emptyset$, then the entries of the Stokes matrix with respect to $\widetilde{\boldsymbol{v}}_S$ and $\widetilde{\boldsymbol{v}}_{S'}$ are elements of R.

Proof The second condition implies the third. Suppose the third. By using the adapted flat frames \tilde{v}_S for any strictly adapted sectors S, we can define an R-structure of the local system. It is clearly compatible with the Stokes structure by construction, i.e., the first condition is satisfied.

Let us prove that the first condition implies the second. We take a holomorphic frame $\mathbf{v}_{\mathfrak{a}}$ of $\mathrm{Gr}_{\mathfrak{a}}^{\mathcal{F}}(V, \nabla)$ such that $\mathbf{v}_{\mathfrak{a}} = \mathbf{v}_{\mathfrak{a}} \Big(d\mathfrak{a} \, I_{r(\mathfrak{a})} + A_{\mathfrak{a}} \, dz/z \Big)$, where $A_{\mathfrak{a}}$ is a constant matrix. We may assume that the multi-valued flat frame $\mathbf{v}_{\mathfrak{a}} \exp \left(-\mathfrak{a} I_{r(\mathfrak{a})} - A_{\mathfrak{a}} \log z \right)$ is a frame of $\mathcal{L}_{\mathfrak{a}R}$. We have a natural isomorphism $\mathrm{Gr}^{\mathcal{F}}(V) \otimes \mathbb{C}[\![z]\!] \simeq V \otimes \mathbb{C}[\![z]\!]$. Let $\widehat{\boldsymbol{v}}_{\mathfrak{a}}$ be induced by $\boldsymbol{v}_{\mathfrak{a}}$ and the isomorphism. For any strictly adapted sector S, we take a flat frame $\widetilde{\boldsymbol{v}}_{S}$ from $\widehat{\boldsymbol{v}} = (\widehat{\boldsymbol{v}}_{\mathfrak{a}})$. It naturally gives a frame of $\mathcal{L}_{R|S}$. Then, the entries of the Stokes matrices with respect to the lift of $\widehat{\boldsymbol{v}}$ are elements of R.

Remark 4.9 The conditions in Lemma 4.8 are equivalent to the existence of a covering $X \setminus D = \bigcup_{i=1}^{N} S_i$ by strictly adapted sectors, such that (i) each S_i is equipped with an adapted flat frame \mathbf{v}_{S_i} of $(V, \nabla)_{|S_i}$ \mathbf{v}_{S_i} , (ii) if $S_i \cap S_j \neq \emptyset$, the entries of the Stokes matrix with respect to $\widetilde{\mathbf{v}}_{S_i}$ and $\widetilde{\mathbf{v}}_{S_j}$ are elements of R. It will be explained elsewhere.

Let us consider the case where (V, ∇) is not necessarily unramified but has pure slope. We have an appropriate ramified covering $\varphi: (X', D') \longrightarrow (X, D)$ such that $\varphi^*(V, \nabla)$ is unramified with pure slope.

Definition 4.10 We say that an R-structure of $(V, \nabla)_{|X \setminus D}$ is an R-structure of (V, ∇) , if the induced R-structure of $\varphi^*(V, \nabla)_{|X \setminus D}$ is an R-structure of $\varphi^*(V, \nabla)$.

Suppose we have a decomposition $(V, \nabla) = \bigoplus_{i=1}^{s} (V_i, \nabla_i)$ such that $\mathcal{I}(\nabla_i) \cap \mathcal{I}(\nabla_j) = \emptyset$ if $i \neq j$. Let \mathcal{L}_i denote the local systems corresponding to (V_i, ∇_i) .

Proposition 4.11 If (V, ∇) has an R-structure \mathcal{L}_R , it is compatible with the decomposition $\mathcal{L} = \bigoplus \mathcal{L}_i$. Namely, each \mathcal{L}_i has an R-structure \mathcal{L}_{Ri} compatible with the Stokes structure such that $\mathcal{L}_R = \bigoplus \mathcal{L}_{Ri}$.

Proof Let S be any sector strictly adapted to (V, ∇) . It is also adapted to each (V_i, ∇_i) . The decomposition (23) is compatible with the R-structure of \mathcal{L}_R . Because (23) is a refinement of the decomposition $V = \bigoplus V_i$, we obtain that \mathcal{L}_R is compatible with $V = \bigoplus V_i$.

Let us give a remark on the uniqueness of R-structure in the case that R is a subfield of \mathbb{C} . Let (V, ∇) be a meromorphic flat bundle on (X, D) with pure slope. Let \mathcal{L} be the corresponding local system.

Proposition 4.12 Suppose that (V, ∇) is irreducible, and that it has R-structures \mathcal{L}_{Ri} (i = 1, 2). Then, there exists a complex number α such that $\mathcal{L}_{R1} = \alpha \mathcal{L}_{R2}$ in \mathcal{L} . The number α is well defined in $\mathbb{C}^*/\mathbb{Q}^*$, where k^* denotes $k \setminus \{0\}$ for a field k.

Proof Let $\varphi: (X', D') \longrightarrow (X, D)$ be a ramified covering $\varphi(z) = z^e$ such that $\varphi^*(V, \nabla)$ is unramified. It has the R-structures $\varphi^{-1}\mathcal{L}_{Ri}$, which are compatible with the Stokes structure of $\varphi^*(V, \nabla)$. Let $\operatorname{Hom}^{\operatorname{St}}(\mathcal{L}_{R1}, \mathcal{L}_{R2})$ denote the space of R-homomorphisms $f: \mathcal{L}_{R1} \longrightarrow \mathcal{L}_{R2}$ such that, for any sectors $S \subset X' \setminus D'$ adapted $\varphi^*(V, \nabla)$, the restriction $\varphi^{-1}f_{|S}$ preserves the decompositions of $\varphi^{-1}\mathcal{L}_{Ri|S}$. It is naturally an R-vector space. Similarly, let $\operatorname{Hom}^{\operatorname{St}}(\mathcal{L}, \mathcal{L})$ denote the space of \mathbb{C} -homomorphisms $f: \mathcal{L} \longrightarrow \mathcal{L}$ such that, for any strictly adapted sectors $S \subset X' \setminus D'$, the restriction $\varphi^{-1}f_{|S}$ preserves the decompositions of $\varphi^{-1}\mathcal{L}_{|S}$. By the Riemann-Hilbert correspondence, $\operatorname{Hom}^{\operatorname{St}}(\mathcal{L}, \mathcal{L})$ is naturally isomorphic to the space of flat morphisms $(V, \nabla) \longrightarrow (V, \nabla)$. By the irreducibility of (V, ∇) , we have $\operatorname{Hom}^{\operatorname{St}}(\mathcal{L}, \mathcal{L}) = \mathbb{C}$. The following naturally induced morphism is an isomorphism:

$$\operatorname{Hom}^{\operatorname{St}}(\mathcal{L}_{R1},\mathcal{L}_{R2}) \otimes_R \mathbb{C} \longrightarrow \operatorname{Hom}^{\operatorname{St}}(\mathcal{L},\mathcal{L})$$

We take a non-zero element $f \in \text{Hom}^{\text{St}}(\mathcal{L}_{R1}, \mathcal{L}_{R2})$. Then, there exists a complex number β such that $\beta f = \text{id}_{\mathcal{L}}$. It implies the claim of the proposition.

4.1.3 Appendix: Stokes filtration

We also have a formulation of R-structure in terms of the Stokes filtrations, which works even in the case that (V, ∇) does not have pure slope, although we shall not use it in this paper. For an adapted sector S, which is not necessarily strictly adapted, the decomposition (23) is not uniquely determined. For simplicity, suppose (V, ∇) is unramified. We consider the filtration \mathcal{F}^S given by $\mathcal{F}^S_{\mathfrak{a}}(V_{|S}) = \bigoplus_{\mathfrak{b} \leq_S \mathfrak{a}} V_{\mathfrak{b},S}$. Here, $\mathfrak{a} \leq_S \mathfrak{b}$ is defined to be $-\operatorname{Re}(\mathfrak{a}) \leq -\operatorname{Re}(\mathfrak{b})$ on S. The filtration is given also by a growth order of flat sections of (V, ∇) . In particular, it is independent of the choice of the decomposition (23). It is called the Stokes filtration of (V, ∇) . It induces a filtration of $\mathcal{L}_{|S|}$ denoted by \mathcal{F}^S .

Remark 4.13 Suppose that (V, ∇) is unramified with pure slope. Then, (V, ∇) has an R-structure, if and only if the associated local system has an R-structure \mathcal{L}_R satisfying the following conditions:

- For each adapted sector S, the filtration $\mathcal{F}^S(\mathcal{L}_{|S})$ is induced by a filtration $\mathcal{F}^S(\mathcal{L}_{R|S})$ of $\mathcal{L}_{R|S}$.
- The filtration $\mathcal{F}^S(\mathcal{L}_{R|S})$ has a splitting, i.e., $\mathcal{F}^S_{\mathfrak{a}}(\mathcal{L}_{R|S}) \longrightarrow \mathcal{F}^S_{\mathfrak{a}}(\mathcal{L}_{R|S})/\mathcal{F}^S_{<\mathfrak{a}}(\mathcal{L}_{R|S})$ has a splitting.

Indeed, the "only if" part is clear. We shall explain the "if" part elsewhere. The condition for the Stokes filtration makes sense, even if (V, ∇) does not have pure slope.

4.2 Integral structure of a certain meromorphic flat bundle

4.2.1 Statement

Let $\mathbf{a} = (a_i \mid i = 1, \dots, r) \in \mathbb{C}^r$. We set $P_{\mathbf{a}}(T) := \prod_{i=1}^r (T - e^{2\pi\sqrt{-1}a_i})$. Let diag $[a_1, \dots, a_r]$ denote the diagonal matrix whose (i, i)-entries are a_i . We consider the following meromorphic connection $\nabla_{\mathbf{a}}$ of \mathcal{V}_r :

$$\nabla_{\mathbf{a}}\mathbf{e} = \mathbf{e}\left(\mathcal{K}(r,1) - \operatorname{diag}[a_1, \dots, a_r]\right) \frac{dq}{q}$$
(25)

It is irregular singular at ∞ , and regular singular at 0. Let R be any subring of \mathbb{C} , and let R^* denote the set of the invertible elements in R.

Theorem 4.14 The meromorphic flat bundle $(\mathcal{V}_r, \nabla_a)$ has an R-structure if and only if $P_a(T) \in R[T]$ and $P_a(0) \in R^*$. In that case, the induced R-structure on $\operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(\mathcal{V}_r, \nabla_a)$ are local systems of free R-modules. (See the remark right before Lemma 4.8 for the induced R-structure of $\operatorname{Gr}_{\mathfrak{a}}^{\mathcal{F}}(\mathcal{V}_r, \nabla_a)$.)

Let us prove the "only if" part. Suppose $(\mathcal{V}_r, \nabla_{\mathbf{a}})$ has an R-structure \mathcal{L}_R . Let $R_f := R[f^{-1}]$. We may assume that R is finitely generated over \mathbb{Z} . Take any $f \in R$ such that $\mathcal{L}_R \otimes_R R_f$ is a locally constant sheaf of free R_f -module. We have $P_{\mathbf{a}}(T) \in R_f[T]$, because the polynomial $P_{\mathbf{a}}(T)$ is the characteristic polynomial of the monodromy along the loop around q = 0 in the counter clock-wise direction. Because \mathcal{L}_R is a locally constant sheaf of finitely generated projective R-modules, we can take $f_i \in R$ (i = 1, ..., N) as above such that $\operatorname{Spec}(R) = \bigcup \operatorname{Spec}(R_{f_i})$. Hence, we have $P_{\mathbf{a}}(T) \in R[T]$.

To prove the "if" part, we shall observe that the non-zero entries of the Stokes factors with respect to some natural frames can be described in terms of the coefficients of the polynomial $P_a(T)$ (Proposition 4.21).

Before going to the proof, we shall give some consequences.

Corollary 4.15 Let \mathcal{R} denote the subring of \mathbb{C} generated by $e^{\pm 2\pi\sqrt{-1}a_i}$ $(i=1,\ldots,r)$ over \mathbb{Z} . Then, (\mathcal{V}_r,∇_a) has an \mathcal{R} -structure.

Corollary 4.16 For the expression $P_{\mathbf{a}}(T) = T^r + \sum_{j=0}^{r-1} P_j T^j$, let \mathcal{R} denote the subring of \mathbb{C} generated by P_i $(i=0,\ldots,r-1)$ and P_0^{-1} over \mathbb{Z} , Then, $(\mathcal{V}_r,\nabla_{\mathbf{a}})$ has an \mathcal{R} -structure.

Corollary 4.17 Suppose $\mathbf{a} \in \mathbb{Q}^r$. Then, $(\mathcal{V}_r, \nabla_{\mathbf{a}})$ has a \mathbb{Q} -structure if and only if $P_{\mathbf{a}}(T) \in \mathbb{Z}[T]$. In that case, $(\mathcal{V}_r, \nabla_{\mathbf{a}})$ has a \mathbb{Z} -structure.

Proof If $(\mathcal{V}_r, \nabla_a)$ has a \mathbb{Q} -structure, we have $P_a(T) \in \mathbb{Q}[T]$. Because $e^{2\pi\sqrt{-1}a_i}$ are algebraic integers, we obtain $P_a(T) \in \mathbb{Z}[T]$. The converse is clear from Theorem 4.14.

4.2.2 Preliminary

In the following, $\nabla_{\boldsymbol{a}}$ will be denoted by ∇ . Let $\varphi: \mathbb{P}^1_w \longrightarrow \mathbb{P}^1_q$ be the ramified covering given by $\varphi(w) = w^r$. The pull back $\varphi^*(\mathcal{V}_r, \nabla)$ is denoted by $(\mathcal{V}^\circ, \nabla^\circ)$. We put $\tau := \exp(2\pi \sqrt{-1}/r)$. We set $v_j := \sum_i \tau^{ij} w^{-i+1} \varphi^* e_i$.

Lemma 4.18 We have $\nabla^{\circ} \mathbf{v} = \mathbf{v} \left(\operatorname{diag}[\tau^{-1}, \dots, \tau^{-r}] \, r \, dw + B \, dw/w \right)$ for some constant matrix B. For any i, the (i, i)-entry of B is $b := -\sum a_j - (r-1)/2$.

Proof It can be checked by a direct computation. We give an indication. Let $v'_i := w^{-i+1}e_i$. Let C be a cyclic matrix such that the (i, j)-entry is 1 if $i - j \equiv 1$ modulo r, or 0 otherwise. Then, we have

$$\nabla^{\circ} \boldsymbol{v}' = \boldsymbol{v}' \Big(Crdw - \big(\operatorname{diag}[a_1, \dots, a_r]r + [0, 1, \dots, r-1] \big) dw/w \Big).$$

Let Υ be the $r \times r$ -matrix whose (i, j)-entries are τ^{ij} . We have $\mathbf{v} = \mathbf{v}' \Upsilon$. We have $\Upsilon^{-1} = r^{-1}(\tau^{-ij})$. We set $B := -\Upsilon^{-1}(r \operatorname{diag}[a_1, \dots, a_r] + [0, 1, \dots, r-1])\Upsilon$. Then, we obtain the formula for $\nabla^{\circ} \mathbf{v}$. Moreover, the (i, i)-entries of B are $\sum r^{-1}\tau^{-ij}(-a_jr+1-j)\tau^{ji} = -\sum a_j - (r-1)/2$.

By a standard argument as in [27], at the formal completion $\widehat{\infty}$, we have a formal frame \widehat{p} such that

$$\nabla^{\circ} \widehat{\boldsymbol{p}} = \widehat{\boldsymbol{p}} \left(\operatorname{diag}[\tau^{-1}, \dots, \tau^{-r}] r \, dw + \operatorname{diag}[b, \dots, b] \frac{dw}{w} \right)$$
 (26)

In particular, we have $\mathcal{I}(\nabla^{\circ}) = \{r\tau^{-i}w \mid i=1,\ldots,r\}$. We set $\mathfrak{a}_i := r\tau^{-i}w$. We use the convention $\mathfrak{a}_i = \mathfrak{a}_{r+i}$. Let $\mu_r := \{\kappa \in \mathbb{C} \mid \kappa^r = 1\}$. It is regarded as the Galois group of the ramified covering φ with the action $(\kappa, w) \longmapsto \kappa w$. The meromorphic flat bundle $(\mathcal{V}^{\circ}, \nabla^{\circ})$ is naturally μ_r -equivariant. We have the ambiguity of the choice of \widehat{p} , i.e., the scalar multiplication of each \widehat{p}_i . We have $(\tau^{-1})^*\widehat{p}_j = \xi_j\widehat{p}_{j+1}$ for some $\xi_j \in \mathbb{C}^*$ with respect to the action. We have $\prod \xi_j = 1$.

4.2.3 Flat frames

We consider the following sectors:

$$S_1 := \{ w \mid |\arg(w) - (\pi/2 - \epsilon)| < \pi/2 \} \qquad S_2 := \{ w \mid |\arg(w) - (\pi/2 + 2\pi/r - \epsilon)| < \pi/2 \}$$
 (27)

Here, ϵ is a sufficiently small positive number. We have $\tau(S_1) = S_2$. We put $S = S_1 \cup S_2 \cup \{\arg(w) = 2\pi/r - \epsilon\}$. The last term is necessary only in the case r = 2.

It is easy to see that S_i (i = 1, 2) are strictly adapted to the connection ∇° . We have the following flat decompositions as in (23):

$$\mathcal{V}_{|S_i}^{\circ} = \bigoplus_{\mathfrak{a} \in \mathcal{I}(\nabla^{\circ})} \mathcal{V}_{S_i,\mathfrak{a}}^{\circ} \tag{28}$$

We set $\omega := e^{2\pi\sqrt{-1}b} \in R^*$. Let $C_1 = ((C_1)_{ij})$ be the $r \times r$ -matrix such that $(C_1)_{ij} = 1$ if i - j = 1, $(C_1)_{ij} = \omega$ if (i,j) = (r,1), and $(C_1)_{ij} = 0$ otherwise.

Lemma 4.19 There exist flat frames $\widetilde{\boldsymbol{p}}_{S_i} = (\widetilde{p}_{j,S_i} \mid j = 1, \dots, r)$ of $\mathcal{V}_{\mid S_i}^{\circ}$ such that (i) \widetilde{p}_{j,S_i} are flat sections of $\mathcal{V}_{S_i,\mathfrak{a}_j}^{\circ}$, (ii) $(\tau^{-1})^*\widetilde{\boldsymbol{p}}_{S_1} = \widetilde{\boldsymbol{p}}_{S_2}C_1$.

Proof We put $\omega_1 := e^{2\pi\sqrt{-1}b/r}$. Let $\widehat{\boldsymbol{p}}$ be a formal frame of \mathcal{V}° at ∞ satisfying (26). By adjusting the scalar multiplication, we may assume that $\widehat{p}_j = \omega_1^{j-1} \cdot (\tau^{1-j})^* \widehat{p}_1$. Then, we have $(\tau^{-1})^* \widehat{\boldsymbol{p}} = \widehat{\boldsymbol{p}} C_1 \omega_1^{-1}$.

On any strictly adapted sector S, we have a unique holomorphic frame \mathbf{p}_S of V such that \mathbf{p}_S is asymptotic to $\widehat{\mathbf{p}}$, and $\nabla \mathbf{p}_S = \mathbf{p}_S \left(\operatorname{diag}[\tau^{-1}, \dots, \tau^{-r}] \, r \, dw + \operatorname{diag}[b, \dots, b] dw/w \right)$ holds. We fix a branch of $\log w$ on $S_1 \cup S_2$. If r = 2, we impose that it is analytic on S. We define

$$\widetilde{\boldsymbol{p}}_{S_i} := \boldsymbol{p}_{S_i} \, \exp \left(\operatorname{diag}[-\tau^{-1}, \dots, -\tau^{-r}] r w \right) \exp(-b \log w),$$

which gives a flat frame on S. Then, we have the following equalities:

$$(\tau^{-1})^* \widetilde{\boldsymbol{p}}_{S_1} = (\tau^{-1})^* \boldsymbol{p}_{S_1} (\tau^{-1})^* \left(\exp\left(\operatorname{diag}\left[-\tau^{-1}rw, \dots, -\tau^{-r}rw\right]\right) \exp\left(-b\log w\right) \right)$$

$$= \boldsymbol{p}_{S_2} C_1 \omega_1^{-1} \exp\left(\operatorname{diag}\left[-\tau^{-2}rw, \dots, -\tau^{-r-1}rw\right]\right) \exp\left(-b\log w\right) \omega_1$$

$$= \widetilde{\boldsymbol{p}}_{S_2} \exp\left(\operatorname{diag}\left[\tau^{-1}rw, \dots, \tau^{-r}rw\right]\right) C_1 \exp\left(\operatorname{diag}\left[-\tau^{-2}rw, \dots, -\tau^{-r-1}rw\right]\right) = \widetilde{\boldsymbol{p}}_{S_2} C_1 \quad (29)$$

Thus, the proof of Lemma 4.19 is finished.

To adjust the signature in the later formulas, we set

$$y_{j,S_i} := \begin{cases} \omega \, \widetilde{p}_{j,S_i} & (j = 1, \dots, [(r-1)/2]) \\ \widetilde{p}_{j,S_i} & (j = [(r-1)/2] + 1, \dots, r). \end{cases}$$

We define y_{j,S_i} for any $j \in \mathbb{Z}$ by using the convention $y_{j+\ell r,S_i} = y_{j,S_i}$.

Lemma 4.20 If r = 2m, there exist complex numbers α_j (j = 1, ..., m - 1) and β_j (j = 1, ..., m) such that the following holds:

$$y_{j,S_1} = \begin{cases} y_{j,S_2} & (0 \le j \le m - 1) \\ y_{j,S_2} + \alpha_{-j} y_{-j,S_2} + \beta_{-j} y_{-j-1,S_2} & (-m < j < 0) \\ y_{-m,S_2} + \beta_m y_{m-1,S_2} & (j = -m) \end{cases}$$

If r = 2m + 1, there exist complex numbers α_j and β_j (j = 1, ..., m) such that the following holds:

$$y_{j,S_1} = \begin{cases} y_{j,S_2} & (0 \le j \le m) \\ y_{j,S_2} + \alpha_{-j}y_{-j,S_2} + \beta_{-j}y_{-j-1,S_2} & (-m \le j < 0) \end{cases}$$

Proof On the half line arg(w) = 0, we have

$$Re(-\tau^{-\ell}w) = Re(-\tau^{\ell}w) \quad (\ell = 1, \dots, [(r-1)/2])$$

On the half line $arg(w) = \pi/r$, we have

$$\operatorname{Re}(-\tau^{-\ell}w) = \operatorname{Re}(-\tau^{\ell+1}w) \quad (\ell = 0, 1, \dots, \lceil r/2 \rceil - 1)$$

On $S_1 \cap S_2$, we have $\text{Re}(-\tau^{-\ell}w) < \text{Re}(-\tau^{\ell}w)$ and $\text{Re}(-\tau^{-\ell}w) < \text{Re}(-\tau^{\ell+1}w)$ for ℓ as above. Then, the claim of the lemma follows.

4.2.4 Stokes factor

We set $x_{j,S_i} := y_{-\lceil r/2 \rceil + j - 1,S_i}$ and $\boldsymbol{x}_{S_i} = (x_{1,S_i}, \dots, x_{r,S_i})$. By construction, we have

$$(\tau^{-1})^* \boldsymbol{x}_{S_1} = \boldsymbol{x}_{S_2} C_1. \tag{30}$$

We can naturally regard x_{S_i} as tuples of flat sections on S. Let A be the matrix determined by $x_{S_1} = x_{S_2}A$. If r = 2m, we have the following:

$$A_{k\ell} = \begin{cases} 1 & (k = \ell) \\ \beta_j & (k, \ell) = (m + j, m - j + 1), \quad (1 \le j \le m) \\ \alpha_j & (k, \ell) = (m + j + 1, m - j + 1), \quad (1 \le j \le m - 1) \\ 0 & (\text{otherwise}) \end{cases}$$
(31)

If r = 2m + 1, we have the following:

$$A_{k\ell} = \begin{cases} 1 & (k = \ell) \\ \beta_j & (k,\ell) = (m+j, m-j+1), \quad (1 \le j \le m) \\ \alpha_j & (k,\ell) = (m+j+1, m-j+1), \quad (1 \le j \le m) \\ 0 & (\text{otherwise}) \end{cases}$$
(32)

For example, if r = 4, we have the following relations:

$$m{x}_{S_1} = m{x}_{S_2} \left(egin{array}{cccc} 1 & 0 & 0 & 0 \ 0 & 1 & 0 & 0 \ 0 & eta_1 & 1 & 0 \ eta_2 & lpha_1 & 0 & 1 \end{array}
ight) \qquad (au^{-1})^*m{x}_{S_1} = m{x}_{S_2} \left(egin{array}{cccc} 0 & 0 & 0 & \omega \ 1 & 0 & 0 & 0 \ 0 & 1 & 0 & 0 \ 0 & 0 & 1 & 0 \end{array}
ight)$$

If r = 5, we have the following relation:

$$m{x}_{S_1} = m{x}_{S_2} \left(egin{array}{ccccc} 1 & 0 & 0 & 0 & 0 & 0 \ 0 & 1 & 0 & 0 & 0 & 0 \ 0 & eta_1 & 1 & 0 & 0 & 0 \ eta_2 & lpha_1 & 0 & 1 & 0 & 0 \ lpha_2 & 0 & 0 & 0 & 1 \end{array}
ight) \qquad (au^{-1})^*m{x}_{S_1} = m{x}_{S_2} \left(egin{array}{ccccccc} 0 & 0 & 0 & 0 & \omega & \omega & \omega & 0 & 0 \ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{array}
ight)$$

4.2.5 The Stokes factor and the monodromy

The following proposition means that the non-zero entries of the Stokes factor A is described by the coefficients of the polynomial $P_a(T)$.

Proposition 4.21 We have the following formula:

$$P_{a}(T) = T^{r} - \sum_{j=1}^{[(r-1)/2]} \alpha_{j} T^{r-2j} - \sum_{j=1}^{[r/2]} \beta_{j} T^{r-2j+1} - \omega$$

Proof We naturally regard \boldsymbol{x}_{S_1} and $(\tau^{-1})^*\boldsymbol{x}_{S_1}$ as tuples of flat sections of $(\mathcal{V}^{\circ}, \nabla^{\circ})_{|S}$. The monodromy of (\mathcal{V}_r, ∇) along the loop in \mathbb{C}_q^* around 0 in the counter-clockwise direction is the relation between \boldsymbol{x}_{S_1} and $(\tau^{-1})^*\boldsymbol{x}_{S_1}$, and it is expressed by the matrix M determined by $\boldsymbol{x}_{S_1} = (\tau^{-1})^*\boldsymbol{x}_{S_1}M$. The characteristic polynomial of M is $P_{\boldsymbol{a}}(T)$. By the definition of A and the relation (30), we obtain $M = C_1^{-1}A$. In particular, the characteristic polynomial $\mathcal{P}_{C_1^{-1}A}(T)$ of $C_1^{-1}A$ is equal to $P_{\boldsymbol{a}}(T)$. It is elementary to prove the following formulas:

$$\frac{\partial}{\partial \beta_j} \mathcal{P}_{C_1^{-1}A}(T) = -T^{r+1-2j} \qquad \frac{\partial}{\partial \alpha_j} \mathcal{P}_{C_1^{-1}A}(T) = -T^{r-2j}$$

If $\alpha_j = 0$ and $\beta_j = 0$ for any j, we have $\mathcal{P}_{C_1^{-1}A}(T) = T^r - \omega$. Then, we obtain the claim of Proposition 4.21.

4.2.6 End of the proof of Theorem 4.14

Suppose $P_{\boldsymbol{a}}(T) \in R[T]$. We obtain that the entries of A are elements of R by Proposition 4.21. We also have $\omega \in R^*$. We set $S_{3/2} := \{w \mid |\arg(w) - (\pi/2 + \pi/r - \epsilon)| < \pi/2\}$. It is standard to construct an adapted flat frame $\boldsymbol{x}_{S_{3/2}}$ by using the factorization of the matrix A. Let $A^{(1)}$ and $A^{(2)}$ be the matrix as follows:

$$A_{k\ell}^{(1)} = \begin{cases} 1 & (k = \ell) \\ \beta_j & (k, \ell) = ([r/2] + j, [r/2] - j + 1), \quad (1 \le j \le [r/2]) \\ 0 & \text{otherwise} \end{cases}$$

$$A_{k\ell}^{(2)} = \left\{ \begin{array}{ll} 1 & (k=\ell) \\ \alpha_j & (k,\ell) = ([r/2]+j+1,[r/2]-j+1), & (1 \leq j \leq [(r-1)/2]) \end{array} \right.$$

We set $\boldsymbol{x}_{S_{3/2}} := \boldsymbol{x}_{S_2}A^{(2)}$. Then, we have $\boldsymbol{x}_{S_1} = \boldsymbol{x}_{S_{3/2}}A^{(1)}$. The entries of $A^{(i)}$ and $(A^{(i)})^{-1}$ are integers. For $\ell = j+1$ or j+3/2 $(j=0,\ldots,r-1)$, we set $S_\ell := \tau^j(S_{\ell-j})$. They are strictly adapted to $(\mathcal{V}^\circ,\nabla^\circ)$. We have adapted frames $\boldsymbol{x}_{S_\ell} := (\tau^{-j})^*\boldsymbol{x}_{S_{\ell-j}}C_1^{-j}$ of $(\mathcal{V}^\circ,\nabla^\circ)_{|S_\ell}$. The Stokes factors with respect to the frames $\boldsymbol{x}_{S_{j+3/2}}$ and $\boldsymbol{x}_{S_{j+1}}$ is $C_1^jA^{(1)}C_1^{-j}$. The Stokes factors with respect to the frames $\boldsymbol{x}_{S_{j+2}}$ and $\boldsymbol{x}_{S_{j+3/2}}$ is $C_1^jA^{(2)}C_1^{-j}$. We remark that $(\tau^{-r})^*\boldsymbol{x}_{S_\ell}C_1^{-r} = \boldsymbol{x}_{S_\ell}$, where we care the analytic continuation of $\log w$. Any strictly adapted sector can be deformed to one of S_ℓ preserving the strictly adaptedness. Therefore, we obtain Theorem 4.14, according to Lemma 4.8.

4.3 Integral structure on the pull back

4.3.1 Statement

We continue to use the notation in §4.2. Suppose that $e^{\pm 2\pi\sqrt{-1}a_i}$ are algebraic integers. Let m be a positive integer. We consider the following meromorphic connection $\nabla_a^{(m)}$ of \mathcal{V}_r :

$$\nabla_{\mathbf{a}}^{(m)} \mathbf{e} = \mathbf{e} \left(\mathcal{K}(r, m) - \operatorname{diag}[a_1, \dots, a_r] \right) m \frac{dq}{q}$$
(33)

Let $K \subset \mathbb{C}$ be an algebraic number field. Let R be the ring of integers in K.

Theorem 4.22 If $(\mathcal{V}_r, \nabla_{\mathbf{a}}^{(m)})$ is irreducible, the following conditions are equivalent.

- $(\mathcal{V}_r, \nabla_{\mathbf{a}}^{(m)})$ has an R-structure.
- $(\mathcal{V}_r, \nabla_{\mathbf{a}}^{(m)})$ has a K-structure.
- There exists $\gamma \in \mathbb{C}^*$ such that (i) $\gamma^m \in K$, (ii) $\prod_{i=1}^r (T \gamma e^{2\pi\sqrt{-1}a_i}) \in K[T]$.

Proof The first condition clearly implies the second. Let us show that the second condition implies the first. Let $\mathcal{K} \subset \mathbb{C}$ be a sufficiently large algebraic number field which contains K and $e^{2\pi\sqrt{-1}a_i}$ $(i=1,\ldots,r)$. Let \mathcal{R} denote the ring of integers in \mathcal{K} . Because $\prod_{i=1}^r (T-e^{2\pi\sqrt{-1}a_i}) \in \mathcal{R}[T]$, according to Theorem 4.14, $(\mathcal{V}_r, \nabla_{\boldsymbol{a}})$ has an \mathcal{R} -structure. Because $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$ is isomorphic to the pull back of $(\mathcal{V}_r, \nabla_{\boldsymbol{a}})$ by $\varphi_m(q) = q^m$, $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$ also has an \mathcal{R} -structure, denoted by $\mathcal{L}_{\mathcal{R}}$. Let \mathcal{L}_K be a K-structure of $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$ in the second condition. By Proposition 4.12, we may assume $\mathcal{L}_{\mathcal{R}} \otimes_{\mathcal{R}} \mathcal{K} = \mathcal{L}_K \otimes_K \mathcal{K}$ by adjusting \mathcal{L}_K . Then, we obtain an R-structure \mathcal{L}_R of $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$ by using the following lemma.

Lemma 4.23 Let V_K be a finite dimensional K-vector space. Suppose that we are given an \mathcal{R} -lattice $V_{\mathcal{R}}$ of $V_K := V_K \otimes_K \mathcal{K}$. Let $\iota : V_K \longrightarrow V_K$ denote the natural inclusion. Then, $V_R := \iota^{-1}(V_{\mathcal{R}})$ is an R-lattice of V_K . Moreover, if we are given decompositions $V_K = \bigoplus_{i \in \mathcal{I}} V_{Ki}$ and $V_{\mathcal{R}} = \bigoplus_{i \in \mathcal{I}} V_{\mathcal{R}i}$ such that $V_{Ki} \otimes_K \mathcal{K} = V_{\mathcal{R}i} \otimes_{\mathcal{R}} \mathcal{K}$, then we have an induced decomposition $V_R = \bigoplus_{i \in \mathcal{I}} V_{Ri}$ such that $V_{Ri} \otimes_R \mathcal{K} = V_{Ki}$ and $V_{Ri} \otimes_R \mathcal{R} = V_{\mathcal{R}i}$.

Proof If we regard $V_{\mathcal{R}}$ as an R-module by $R \subset \mathcal{R}$, then it is finitely generated over the Noetherian ring R. Because V_R is naturally an R-submodule of $V_{\mathcal{R}}$, it is finitely generated over R. It is torsion-free. Because R is a one dimensional regular ring, V_R is projective. Let $j:V_R\longrightarrow V_K$ be the natural inclusion. The induced morphism $j_K:V_R\otimes_R K\longrightarrow V_K$ is injective. For any $f\in V_K$, there exists an integer n_f such that $n_f\iota(f)\in V_{\mathcal{R}}$. Hence, j_K is surjective. Thus, we obtain the first claim. We can easily obtain the second claim by using the first claim. Thus, Lemma 4.23 is proved.

Let us show the equivalence of the second and third conditions in Theorem 4.22. We put $\kappa := \exp(2\pi\sqrt{-1}/m)$ and $\mu_m := \{\kappa^j \mid j = 0, 1, \dots, m-1\}$. We consider the μ_m -action on \mathbb{P}^1 given by $(\kappa^j, q) \longmapsto \kappa^j q$. We have a μ_m -action on \mathcal{V}_r given by $\kappa^* e_i \longleftrightarrow e_i$. The connection $\nabla_a^{(m)}$ is equivariant with respect to the action.

For any $\nu \in \mathbb{C}$, let $L(\nu)$ be the meromorphic flat bundle $\mathcal{O}_{\mathbb{P}^1}(*D) y_{\nu}$ with $\nabla y_{\nu} = y_{\nu} \nu dq/q$. The multi-valued flat section is given by $\rho_{\nu} := y_{\nu} \exp(-\nu \log q)$. We consider a μ_m -action on $L(\nu)$ given by $\kappa^* y_{\nu} = y_{\nu}$. The tensor product $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)}) \otimes L(\nu)$ is equipped with an induced μ_m -action. Its descent with respect to the μ_m -action is isomorphic to (\mathcal{V}, ∇') , where ∇' is given as follows:

$$\nabla' \mathbf{e} = \mathbf{e} \left(\mathcal{K}(r,1) - \operatorname{diag}[a_1 - \nu/m, \dots, a_r - \nu/m] \right) \frac{dq}{q}$$
(34)

Suppose the second condition. Let \mathcal{L}_K be a K-structure of $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$. By Proposition 4.12, there exists a $\gamma \in \mathbb{C}^*$ such that $(\kappa^{-1})^*\mathcal{L}_K = \gamma\mathcal{L}_K$. We have $\gamma^m \in K^*$. We take ν such that $\exp(-2\pi\sqrt{-1}\nu/m) = \gamma$. Then, $L(\nu)$ has a K-structure \mathcal{L}_{ν} given by $K\rho_{\nu} \subset \mathcal{L}_{\nu}$, and we have $(\kappa^{-1})^*\mathcal{L}_{\nu} = \gamma^{-1}\mathcal{L}_{\nu}$. Because the induced K-structure $\mathcal{L}_K \otimes \mathcal{L}_{\mu}$ is μ_m -equivariant, we obtain a K-structure of (\mathcal{V}_r, ∇') . Applying Theorem 4.14, we obtain $\prod_{i=1}^r (T - \gamma e^{2\pi\sqrt{-1}a_i}) \in K[T]$, i.e., the third condition holds.

Suppose the third condition. We take $\nu \in \mathbb{C}$ such that $\exp(-2\pi\sqrt{-1}\nu/m) = \gamma$. By Theorem 4.14, (\mathcal{V}_r, ∇') has a K-structure. It induces a K-structure on $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)}) \otimes L(\nu)$, and hence a K-structure on $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$. Thus, Theorem 4.22 is proved.

We have a special case of Theorem 4.22.

Corollary 4.24 Suppose that $a \in \mathbb{Q}^r$. Then, the following conditions are equivalent.

- $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(m)})$ has a \mathbb{Z} -structure.
- $(\mathcal{V}_r, \nabla_{\mathbf{a}}^{(m)})$ has a \mathbb{Q} -structure.
- There exists $\gamma \in \mathbb{C}^*$ such that (i) $\gamma^m \in \mathbb{Q}$, (ii) $\prod_{i=1}^r (T \gamma e^{2\pi\sqrt{-1}a_i}) \in \mathbb{Q}[T]$.

4.3.2 Examples

Let us consider the conditions in the case that r=m is a prime number, and that $K=\mathbb{Q}$ and $R=\mathbb{Z}$. We begin with a preliminary. Take $\gamma\in\mathbb{R}_{>0}\setminus\mathbb{Q}$ such that $\gamma^r\in\mathbb{Q}$. Let P be the minimal polynomial of $\eta:=\gamma e^{2\pi\sqrt{-1}v/(r^mu)}$, where u and v are integers such that g.c.d.(u,v)=g.c.d.(r,u)=g.c.d.(r,v)=1.

Lemma 4.25 Suppose $deg(P) \le r$.

- If r is an odd prime, $P = T^r \pm \gamma^r$.
- If r = 2, we have $P = T^2 \pm \gamma^2$ unless (u, m) = (1, 3), (3, 2).
- If r = 2 and (u, m) = (1, 3), we have $\gamma \in \sqrt{2}\mathbb{Q}_{>0}$ and $P(T) \in \{T^2 + 2bT + 2b^2 \mid b \in \mathbb{Q}\}$.
- If r = 2 and (u, m) = (3, 2), we have $\gamma \in \sqrt{3}\mathbb{Q}_{>0}$ and $P(T) \in \{T^2 + 3bT + 3b^2 \mid b \in \mathbb{Q}\}$.

Proof We set $K := \mathbb{Q}(\eta)$. By the hypothesis, we have $[K : \mathbb{Q}] \leq r$, where $[K_1 : K_2]$ denotes the degree for a finite extension K_1 over K_2 . Because $\eta^{r^{m+1}} = \gamma^{r^{m+1}} e^{2\pi\sqrt{-1}(rv)/u} \in K$, we have $e^{2\pi\sqrt{-1}/u} \in K$.

Let us consider the case that r is an odd prime. Suppose $m \geq 2$, and we shall derive a contradiction. We have $\eta^{r^{m-1}u} = \gamma^{r^{m-1}u}e^{2\pi\sqrt{-1}v/r} \in K$, and hence $e^{2\pi\sqrt{-1}/r} \in K$. Because $[\mathbb{Q}(e^{2\pi\sqrt{-1}/r}):\mathbb{Q}] = r-1$ and $[K:\mathbb{Q}] \leq r$, we obtain $K = \mathbb{Q}(e^{2\pi\sqrt{-1}/r})$ and $\deg P = r-1$. Because $\prod_{i=0}^{r^m u-1} (T - \gamma e^{2\pi\sqrt{-1}i/r^m u}) = T^{r^m u} - \gamma^{r^m u} \in \mathbb{Q}[T]$, a number conjugate to η over \mathbb{Q} is of the form $\gamma e^{2\pi\sqrt{-1}c}$ for $c \in \mathbb{Q}$. We have $P = \prod_{j=1}^{r-1} (T - \gamma e^{2\pi\sqrt{-1}c_j})$ for some $c_j \in \mathbb{Q}$ $(j=1,\ldots,r-1)$, and hence $P(0) = (-1)^{r-1}\gamma^{r-1}e^{2\pi\sqrt{-1}\sum c_j}$. If $e^{2\pi\sqrt{-1}\sum c_j} \notin \mathbb{R}$, we have $P(0) \notin \mathbb{Q}$ by considering the polar part. If $e^{2\pi\sqrt{-1}\sum c_j} \in \mathbb{R}$, then $e^{2\pi\sqrt{-1}\sum c_j} = \pm 1$, and hence $P(0) \notin \mathbb{Q}$. It contradicts with the hypothesis $P(T) \in \mathbb{Q}[T]$. Thus, we obtain m < 1.

the hypothesis $P(T) \in \mathbb{Q}[T]$. Thus, we obtain $m \leq 1$. If $m \leq 1$, we have $\gamma e^{2\pi \sqrt{-1}t/r} \in K$ for some integer t. Recall that $T^r - \gamma^r$ is irreducible in $\mathbb{Q}[T]$. Let $\overline{\mathbb{Q}}$ be an algebraic closure of \mathbb{Q} , and let $f : \overline{\mathbb{Q}} \longrightarrow \overline{\mathbb{Q}}$ be an automorphism of the field over \mathbb{Q} such that $f(\gamma e^{2\pi \sqrt{-1}t/r}) = \gamma$. Then, $f(e^{2\pi \sqrt{-1}/u}) \in K(\gamma) \subset \mathbb{R}$. We obtain u = 1, 2, and $\eta = \pm \gamma e^{2\pi \sqrt{-1}v/r}$. Then, its minimal polynomial is $T^r \pm \gamma^r$.

Let us consider the case r=2. If m=0, we have $\gamma\in K$ and $e^{2\pi\sqrt{-1}v/u}\in K$. Hence, we have $K=\mathbb{Q}(\gamma)$, and $e^{2\pi\sqrt{-1}v/u}\in\mathbb{Q}(\gamma)\subset\mathbb{R}$. We obtain $\eta=\pm\gamma$, and hence $P(T)=T^2-\gamma^2$. Suppose $m\geq 1$. We have $e^{2\pi\sqrt{-1}v/(2^{m-1}u)}\in K$. If $e^{2\pi\sqrt{-1}v/(2^{m-1}u)}\in\mathbb{Q}$, then we have $e^{2\pi\sqrt{-1}v/(2^mu)}\in\{\pm 1,\pm\sqrt{-1}\}$. Hence, $P=T^2\pm\gamma^2$. Suppose $e^{2\pi\sqrt{-1}v/(2^{m-1}u)}\notin\mathbb{Q}$. We have $K=\mathbb{Q}(e^{2\pi\sqrt{-1}v/(2^{m-1}u)})$. Because $[K:\mathbb{Q}]=2$, we have u=1 or 3.

In the case u=1, we have m=3, because $e^{2\pi\sqrt{-1}v/2^{m-1}} \notin \mathbb{Q}$ with degree 2. If (u,m)=(1,3), we have $K=\mathbb{Q}[\sqrt{-1}]$ and $\eta=\gamma e^{2\pi\sqrt{-1}v/8}\in\mathbb{Q}[-1]$, where v=1,3,5,7. Then, we obtain $\gamma=\sqrt{2}a$ for some $a\in\mathbb{Q}_{>0}$. The minimal polynomial of η is either one of the following $(\zeta_8:=e^{2\pi\sqrt{-1}/8})$:

$$\left(T - \sqrt{2}a\zeta_8\right)\left(T - \sqrt{2}a\zeta_8^7\right) = T^2 - 2aT + 2a^2, \qquad \left(T - \sqrt{2}a\zeta_8^3\right)\left(T - \sqrt{2}a\zeta_8^5\right) = T^2 + 2aT + 2a^2$$

Similarly, if u=3, we have m=1 or 2. If (u,m)=(3,1), we have $K=\mathbb{Q}(e^{2\pi\sqrt{-1}/3})=\mathbb{Q}(\sqrt{-3})$. Because $\gamma e^{2\pi\sqrt{-1}v/6}\in\mathbb{Q}(\sqrt{-3})$, we obtain $\gamma\in\mathbb{Q}_{>0}$, which contradicts with our assumption $\gamma\notin\mathbb{Q}$. If (u,m)=(3,2),

we have $\gamma e^{2\pi\sqrt{-1}v/12} \in \mathbb{Q}(\sqrt{-3})$, where v = 1, 5, 7, 11. Hence, we obtain $\gamma \in \sqrt{3}\mathbb{Q}_{>0}$. The minimal polynomial of η is either one of the following $(\zeta_{12} = e^{2\pi\sqrt{-1}/12})$:

$$\left(T - \sqrt{3}a\zeta_{12}\right)\left(T - \sqrt{3}a\zeta_{12}^{11}\right) = T^2 - 3aT + 3a^2, \qquad \left(T - \sqrt{3}a\zeta_{12}^5\right)\left(T - \sqrt{3}a\zeta_{12}^7\right) = T^2 + 3aT + 3a^2,$$

Thus, we obtain Lemma 4.25

If r is an odd prime, we have no exceptional cases in the following sense.

Proposition 4.26 Suppose (i) r is an odd prime, (ii) $\mathbf{a} \in \mathbb{Q}^r$, (iii) $(\mathcal{V}_r, \nabla_{\mathbf{a}}^{(r)})$ is irreducible. Then, $(\mathcal{V}_r, \nabla_{\mathbf{a}}^{(r)})$ has a \mathbb{Z} -structure, if and only if there exists a half-integer ℓ such that $\prod_{i=1}^r (T - e^{2\pi\sqrt{-1}(a_i + \ell/r)}) \in \mathbb{Q}[T]$.

Proof By Theorem 4.22, $(\mathcal{V}_r, \nabla_{\boldsymbol{a}}^{(r)})$ has a \mathbb{Z} -structure, if and only if there exist $\gamma \in \mathbb{R}_{>0}$ and a half-integer ℓ such that (i) $\gamma^r \in \mathbb{Q}$, (ii) $\prod_{i=1}^r (T - \gamma e^{2\pi\sqrt{-1}(a_i + \ell/r)}) \in \mathbb{Q}[T]$. Suppose $\gamma \notin \mathbb{Q}$. By using Lemma 4.25, we obtain $\prod_{i=1}^r (T - \gamma e^{2\pi\sqrt{-1}(a_i + \ell/r)}) = T^r \pm \gamma^r$. It implies that $\prod_{i=1}^r (T - e^{2\pi\sqrt{-1}(a_i + \ell/r)}) \in \mathbb{Q}[T]$. Hence, we obtain the proposition.

If r=2, we obtain the following proposition from Lemma 4.25. We have the exceptional cases.

Proposition 4.27 Suppose (i) $\mathbf{a} \in \mathbb{Q}^2$, (ii) $(\mathcal{V}_r, \nabla^{(2)})$ is irreducible. Then, $(\mathcal{V}_r, \nabla^{(2)})$ has a \mathbb{Z} -structure if and only if there exists an integer n such that one of the following holds:

- $\prod_{i=1}^{2} (T e^{2\pi\sqrt{-1}(a_i + n/4)}) \in \mathbb{Q}[T].$
- $\{a_1, a_2\} = \{-1/8 + n/4, -7/8 + n/4\}$
- $\{a_1, a_2\} = \{-1/12 + n/4, -11/12 + n/4\}$ or $\{-5/12 + n/4, -7/12 + n/4\}$.

Let us compute Stokes matrices in a case. The other cases can be computed similarly. Let $(a_1, a_2) = (-1/8, -7/8)$. We have $\gamma = \sqrt{2}$ and

$$(T - \gamma e^{2\pi\sqrt{-1}a_1})(T - \gamma e^{2\pi\sqrt{-1}a_2}) = T^2 - 2T + 2.$$

We have $b = -(a_1 + a_2) - (r - 1)/2 = 1/2$. We take a formal frame $\mathbf{p} = (\widehat{p}_1, \widehat{p}_2)$ compatible with the Hukuhara-Levelt-Turrittin decomposition such that

$$(-1)^* \widehat{\boldsymbol{p}} = \widehat{\boldsymbol{p}} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -2 \\ 1 & 0 \end{pmatrix}$$

We put $\widehat{\boldsymbol{p}}' := (\widehat{p}_2, \widehat{p}_1).$

Let $S_1 := \{-\epsilon < \arg q < \pi + \epsilon\}$ and $S_2 := \{-\pi - \epsilon < \arg q < \epsilon\}$. We put $T_1 := \{\pi - \epsilon < \arg q < \pi + \epsilon\}$ and $T_2 := \{-\epsilon < \arg q < \epsilon\}$. We have $S_1 \cap S_2 = T_1 \sqcup T_2$. We take flat frames $\widetilde{\boldsymbol{p}}_{S_i}$, which are lifts of $\widehat{\boldsymbol{p}}$. On T_1 , we have

$$\widetilde{\boldsymbol{p}}_{S_1} = \widetilde{\boldsymbol{p}}_{S_2} A, \quad A = \left(\begin{array}{cc} 1 & 0 \\ \alpha & 1 \end{array} \right)$$

We have

$$(-1)^*\widetilde{\boldsymbol{p}}_{S_1} = \widetilde{\boldsymbol{p}}_{S_2} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & -2 \\ 1 & 0 \end{pmatrix}$$

Let M be the monodromy matrix given by $\tilde{p}_{S_1} = (-1)^* \tilde{p}_{S_1}$. Then, we have

$$\sqrt{2}M = \begin{pmatrix} 0 & 2 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \alpha & 1 \end{pmatrix} = \begin{pmatrix} 2\alpha & 2 \\ -1 & 0 \end{pmatrix}$$

We obtain $T^2 - 2T + 2 = (T - 2\alpha)T + 2$ and hence $\alpha = 1$.

Let $\widetilde{\boldsymbol{p}}_{S_i}'$ be the lift of $\widehat{\boldsymbol{p}}'$. On T_2 , we have

$$\widetilde{\boldsymbol{p}}_{S_2}' = \widetilde{\boldsymbol{p}}_{S_1}' A', \quad A' = \left(\begin{array}{cc} 1 & 0 \\ \beta & 1 \end{array} \right)$$

We have

$$(-1)^*\widetilde{\boldsymbol{p}}_{S_2}' = \widetilde{\boldsymbol{p}}_{S_1} \frac{1}{\sqrt{2}} \begin{pmatrix} 0 & 1 \\ -2 & 0 \end{pmatrix}$$

Let M' be the monodromy matrix given by $\widetilde{p}'_{S_2} = (-1)^* \widetilde{p}'_{S_2}$. Then, we have

$$\sqrt{2}M' = \left(\begin{array}{cc} 0 & -1 \\ 2 & 0 \end{array}\right) \left(\begin{array}{cc} 1 & 0 \\ \beta & 1 \end{array}\right) = \left(\begin{array}{cc} -\beta & -1 \\ 2 & 0 \end{array}\right)$$

We obtain $T^2 - 2T + 2 = (T + \beta)T + 2$ and hence $\beta = -2$.

5 Harmonic bundle with homogeneity

5.1 Preliminary

5.1.1 Holomorphic vector fields

Let X be a complex manifold. The tangent bundle TX is equipped with a complex structure, i.e., an automorphism J with $J^2=-1$. We have the canonical decomposition $TX\otimes_{\mathbb{R}}\mathbb{C}=T^{1,0}X\oplus T^{0,1}X$, i.e., $T^{1,0}X$ and $T^{0,1}X$ denote the eigen spaces of J corresponding to $\sqrt{-1}$ and $-\sqrt{-1}$. The bundle $T^{1,0}X$ is naturally a holomorphic vector bundle on X, and $T^{0,1}X$ is naturally a holomorphic vector bundle on the conjugate X^{\dagger} . For the natural conjugation on $TX\otimes\mathbb{C}$, we have $T^{0,1}X=\overline{T^{1,0}X}$.

A section \mathfrak{v} of TX has the decomposition $\mathfrak{v} = \mathfrak{v}^{1,0} + \mathfrak{v}^{0,1}$, and we have $\overline{\mathfrak{v}^{1,0}} = \mathfrak{v}^{0,1}$. The section \mathfrak{v} is called holomorphic, if $\mathfrak{v}^{1,0}$ is a holomorphic section of $T^{1,0}X$, which is equivalent to that $\mathfrak{v}^{0,1}$ is a holomorphic section of $T^{1,0}X^{\dagger} = T^{0,1}X$ on X^{\dagger} . If we are given a local coordinate (z_1, \ldots, z_n) , we have the expression $\mathfrak{v} = \sum (f_i \partial_{z_i} + \overline{f_i} \partial_{\overline{z_i}})$. It is holomorphic if and only if f_i are holomorphic.

Lemma 5.1 Let $L_{\mathfrak{v}}$ denote the Lie derivative on $\Omega^{\bullet}_{\mathbf{X}}$. The following conditions are equivalent.

- ullet The vector field $oldsymbol{v}$ is holomorphic.
- $L_{\mathfrak{p}}\Omega_{\mathbf{Y}}^{p,q} \subset \Omega_{\mathbf{Y}}^{p,q}$ for any p,q.
- $[L_{\mathfrak{p}}, \overline{\partial}_X] = 0.$
- $[L_n, \partial_X] = 0.$

Proof If $\mathfrak{v} = \sum (f_i \partial_{z_i} + \overline{f}_i \partial_{\overline{z}})$, we have $L_{\mathfrak{v}}(dz_i) = df_i$, from which we can deduce the equivalence of the first and second conditions. Because $[L_{\mathfrak{v}}, d_X] = 0$, the second implies the third and fourth. If the third condition holds, we have $L_{\mathfrak{v}}(dz_i) = L_{\mathfrak{v}}\overline{\partial}_X z_i = \overline{\partial}_X (L_{\mathfrak{v}}z_i)$. Hence, we have $L_{\mathfrak{v}}\Omega_X^{1,0} \subset \Omega_X^{1,0}$. Similarly, we have $L_{\mathfrak{v}}\Omega_X^{0,1} \subset \Omega_X^{0,1}$. Then, we obtain the second. Clearly, the third and the fourth are equivalent.

5.1.2 An action on vector bundle

Let E be a C^{∞} -vector bundle on X. We identify it with the sheaf of C^{∞} -sections. Let \mathfrak{v} be a vector field on X. An action of \mathfrak{v} on E is a \mathbb{C} -linear map of sheaves $L^E_{\mathfrak{v}}: E \longrightarrow E$ such that $L^E_{\mathfrak{v}}(fs) = \mathfrak{v}(f) \, s + f L^E_{\mathfrak{v}}(s)$. It is standard that, an action of \mathfrak{v} on E uniquely induces an action of \mathfrak{v} on $\Omega^{\bullet}_X \otimes E$ satisfying $L^E_{\mathfrak{v}}(\omega \otimes s) = L_{\mathfrak{v}}(\omega) \otimes s + \omega \otimes L^E_{\mathfrak{v}}(s)$. A section f of E is called homogeneous of degree $\alpha \in \mathbb{C}$ with respect to \mathfrak{v} , if $L^E_{\mathfrak{v}}f = \alpha f$. An action of \mathfrak{v} on E_i (i = 1, 2) naturally induces an action on $E_1 \oplus E_2$, $E_1 \otimes E_2$ and $\operatorname{Hom}(E_1, E_2)$.

Suppose that E is a holomorphic vector bundle, and that \mathfrak{v} is holomorphic. An action $L^E_{\mathfrak{v}}$ of \mathfrak{v} on E is called holomorphic, if $[L^E_{\mathfrak{v}}, \overline{\partial}_E] = 0$. The following lemma is clear.

Lemma 5.2 Let E^{hol} denote the sheaf of holomorphic sections. A holomorphic action of \mathfrak{v} on E induces a \mathbb{C} -linear map $L^E_{\mathfrak{v}}: E^{\text{hol}} \longrightarrow E^{\text{hol}}$ such that $L^E_{\mathfrak{v}}(fs) = L_{\mathfrak{v}}(f)s + f L^E_{\mathfrak{v}}(s)$ for local sections $f \in \mathcal{O}_X$ and $s \in E^{\text{hol}}$. Conversely, such a \mathbb{C} -linear map on E^{hol} induces a holomorphic action of \mathfrak{v} on E.

5.2 Homogeneous harmonic bundle and integrable variation of twistor structure

Let X be a complex manifold with a holomorphic vector field \mathfrak{v} . A harmonic bundle $(E, \overline{\partial}_E, \theta, h)$ is called homogeneous of degree $m \in \mathbb{R}$, with respect to \mathfrak{v} , if E is equipped with a \mathfrak{v} -action $L^E_{\mathfrak{v}}$ such that $[L^E_{\mathfrak{v}}, \overline{\partial}_E] = 0$, $L^E_{\mathfrak{v}}(h) = 0$ and $[L^E_{\mathfrak{v}}, \theta] = \sqrt{-1}m\theta$. Note that we have $[L^E_{\mathfrak{v}}, \partial_{E,h}] = 0$ and $[L^E_{\mathfrak{v}}, \theta_h^{\dagger}] = -\sqrt{-1}m\theta^{\dagger}$. In the following, we impose that $m \neq 0$, and we shall observe that the associated variation of twistor structure is naturally integrable, if $m \neq 0$.

Remark 5.3 Suppose $[L_{\mathfrak{v}}^E, \theta] = \alpha \theta$ for some complex number $\alpha \notin \sqrt{-1}\mathbb{R}$, instead of $[L_{\mathfrak{v}}^E, \theta] = \sqrt{-1}m\theta$. Then, the local structure of the harmonic bundle is easy. See Proposition 5.8 below. Hence, we exclude the case.

5.2.1 The associated meromorphic flat connection

We set $\mathcal{X} := \mathbb{C}_{\lambda} \times X$ and $\mathcal{X}^{\lambda} := \{\lambda\} \times X$. Let $p_{\lambda} : \mathcal{X} \longrightarrow X$ denote the projection. We have the holomorphic vector bundle $\mathcal{E} = p_{\lambda}^{-1} E$ with the holomorphic structure $\overline{\partial}_E + \lambda \theta^{\dagger} + \overline{\partial}_{\lambda}$. It is equipped with the relative flat connections

$$abla^{\lambda}: \mathcal{E} \longrightarrow \mathcal{E} \otimes \Omega^{1}_{\mathcal{X}/\mathbb{C}_{\lambda}} \otimes \mathcal{O}_{\mathcal{X}}(\mathcal{X}^{0})$$

given by $\nabla^{\lambda} := (\overline{\partial}_E + \lambda \theta^{\dagger}) + (\partial_E + \lambda^{-1}\theta).$

Let $\widetilde{\mathfrak{v}} := \mathfrak{v} + m\sqrt{-1}(\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}})$. We have natural actions of $m\sqrt{-1}(\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}})$ and \mathfrak{v} on $p^{-1}(E)$. They give an action of $\widetilde{\mathfrak{v}}$ on $p^{-1}(E)$, denoted by $L_{\widetilde{\mathfrak{v}}}^{\mathcal{E}}$. We have $[L_{\widetilde{\mathfrak{v}}}^{\mathcal{E}}, \nabla^{\lambda}] = [L_{\widetilde{\mathfrak{v}}}^{\mathcal{E}}, \overline{\partial}_{\mathcal{E}}] = 0$. We can check by an elementary computation that \mathcal{E} has a unique meromorphic flat connection

$$\widetilde{\nabla}: \mathcal{E} \longrightarrow \mathcal{E} \otimes \Omega^1_{\mathcal{X}}(\log \mathcal{X}^0) \otimes \mathcal{O}(\mathcal{X}^0),$$

such that (i) $\widetilde{\nabla}_v = \nabla_v^{\lambda}$ for any vector fields v in the X-direction, (ii) $\widetilde{\nabla}_{\widetilde{\mathfrak{v}}} = L_{\widetilde{\mathfrak{v}}}^{\mathcal{E}}$.

5.2.2 Conjugate

Let X^{\dagger} denote the conjugate of X. We set $\mathcal{X}^{\dagger} := \mathbb{C}_{\mu} \times X^{\dagger}$. The projection $\mathcal{X}^{\dagger} \longrightarrow X^{\dagger}$ is denoted by p_{μ} . We obtain a holomorphic vector bundle $\mathcal{E}^{\dagger} := \left(p_{\mu}^{-1}E, p_{\mu}^{*}\partial_{E} + \mu\theta\right)$. It is equipped with the relative flat connection $\nabla^{\dagger\mu} = \overline{\partial}_{E} + \partial_{E} + \mu\theta + \mu^{-1}\theta^{\dagger}$.

Note that \mathfrak{v} naturally gives a holomorphic vector field on X^{\dagger} . The harmonic bundle $(E, \partial_E, \theta^{\dagger}, h)$ is homogeneous of degree -m with respect to \mathfrak{v} . By the same procedure, we obtain a meromorphic connection

$$\widetilde{\nabla}^{\dagger}: \mathcal{E}^{\dagger} \longrightarrow \mathcal{E}^{\dagger} \otimes \Omega^{1}_{\mathcal{X}^{\dagger}}(\log \mathcal{X}^{0\dagger}) \otimes \mathcal{O}(\mathcal{X}^{0\dagger}).$$

We identify \mathbb{C}_{λ}^* and \mathbb{C}_{μ}^* by $\lambda = \mu^{-1}$. We have a natural identifications $\mathcal{E}_{|\mathbb{C}_{\lambda}^* \times X} = \mathcal{E}_{|\mathbb{C}_{\mu}^* \times X^{\dagger}}$.

Lemma 5.4 On $\mathbb{C}^*_{\lambda} \times X$, we have $\widetilde{\nabla} = \widetilde{\nabla}^{\dagger}$.

Proof We have only to prove that $\widetilde{\nabla}_{\partial_{\lambda}} = \widetilde{\nabla}_{\partial_{\lambda}}^{\dagger}$. We remark $\sqrt{-1}m(\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}}) = -\sqrt{-1}m(\mu\partial_{\mu} - \overline{\mu}\partial_{\overline{\mu}})$. By the constructions, we have the equalities $\widetilde{\nabla}_{V} = \widetilde{\nabla}_{V}^{\dagger}$, if V is $\mathfrak{v} + \sqrt{-1}m(\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}})$, or if V is a vector field in the X-direction. We obtain $\widetilde{\nabla}_{\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}}} = \widetilde{\nabla}_{\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}}}^{\dagger}$. Because the $d\overline{\lambda}$ -parts are the same, we obtain $\widetilde{\nabla}_{\partial_{\lambda}} = \widetilde{\nabla}_{\partial_{\lambda}}^{\dagger}$.

5.2.3 Polarized integrable variation of pure twistor structure

Let $p: \mathbb{P}^1 \times X \longrightarrow X$ be the projection. Due to Simpson [45], we have the polarized variation of pure twistor structure $(p^{-1}(E), \mathbb{D}^{\triangle})$ of weight 0, associated to $(E, \overline{\partial}_E, \theta, h)$. (See [33] for a review.) Lemma 5.4 implies that the operators $\widetilde{\nabla}$ and $\widetilde{\nabla}^{\dagger}$ gives a $T\widetilde{T}E$ -structure $\widetilde{\mathbb{D}}^{\triangle}$ of $p^{-1}(E)$. ([17]. See §2.1 of [33] for a review of $T\widetilde{T}E$ -structure.) Hence, we obtain an integrable variation of pure twistor structure $(p^{-1}(E), \widetilde{\mathbb{D}}^{\triangle})$, which is an enrichment of $(p^{-1}(E), \mathbb{D}^{\triangle})$.

Let $\sigma: \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ be given by $\sigma(\lambda) = (-\overline{\lambda})^{-1}$. The induced map $\mathbb{P}^1 \times X \longrightarrow \mathbb{P}^1 \times X$ is also denoted by σ . A polarization S_h of $(p^{-1}(E), \mathbb{D}^{\triangle})$ is a pairing of $(p^{-1}(E), \mathbb{D}^{\triangle})$ and $\sigma^*(p^{-1}(E), \mathbb{D}^{\triangle})$ taking values in the trivial line bundle on $\mathbb{P}^1 \times X$, satisfying some conditions. In this case, it is given by $S_h(u, \sigma^*v) := h(u, \sigma^*v)$ for sections u and v of $p^{-1}(E)$ on \mathcal{U} and $\sigma(\mathcal{U})$, respectively, where \mathcal{U} is any open subset of $\mathbb{P}^1 \times X$.

Lemma 5.5 The polarization is compatible with the integrability, i.e., for local sections u, v of $p^{-1}(E)$, we have

$$\lambda \partial_{\lambda} S_h(u, \sigma^* v) = S_h(\widetilde{\nabla}_{\lambda \partial_{\lambda}} u, \sigma^* v) + S_h(u, \sigma^*(\widetilde{\nabla}_{\lambda \partial_{\lambda}} v)). \tag{35}$$

Proof We have $d_X h(u, v) = h(\nabla^{\lambda} u, \sigma^* v) + h(u, \sigma^* (\nabla^{\lambda} v))$. Hence, for a vector field V in the X-direction, we have

$$Vh(u, \sigma^*v) = h(\widetilde{\nabla}_V u, \sigma^*v) + h(u, \sigma^*(\widetilde{\nabla}_{\overline{V}}v)).$$

We have $\overline{\partial}_{\lambda}h(u,\sigma^*v)d\overline{\lambda} = h(\overline{\partial}_{\lambda}u\,d\overline{\lambda},\,\sigma^*v) + h(u,\sigma^*(\overline{\partial}_{\mu}v\,d\overline{\mu}))$, and hence

$$\overline{\lambda}\overline{\partial}_{\lambda}h(u,\sigma^*v) = h(\overline{\lambda}\widetilde{\nabla}_{\overline{\lambda}}u,\sigma^*v) + h(u,\sigma^*(\overline{\mu}\widetilde{\nabla}_{\overline{\mu}}v)).$$

Note $\sigma^*(\widetilde{\mathfrak{v}}) = \widetilde{\mathfrak{v}}$. We have $\widetilde{\mathfrak{v}}h(u, \sigma^*v) = h(L_{\widetilde{\mathfrak{v}}}^{\mathcal{E}}u, \sigma^*v) + h(u, \sigma^*(L_{\widetilde{\mathfrak{v}}}^{\mathcal{E}}v))$. We deduce

$$\lambda \partial_{\lambda} h(u, \sigma^* v) = h(\lambda \widetilde{\nabla}_{\lambda} u, \sigma^* v) + h(u, \sigma^* (\mu \widetilde{\nabla}_{\mu} v)),$$

i.e., we obtain (35)

5.2.4 Equivalence

Let $(V^{\triangle},\widetilde{\mathbb{D}}^{\triangle})$ be an integrable variation of twistor structure on X. If $\widetilde{\mathbb{D}}^{\triangle}_{\widetilde{\mathfrak{v}}}(s)$ is C^{∞} for any C^{∞} -section s of V^{\triangle} , then $(V^{\triangle},\widetilde{\mathbb{D}}^{\triangle})$ is called homogeneous of degree m with respect to \mathfrak{v} . If we describe $(V^{\triangle},\widetilde{\mathbb{D}}^{\triangle})$ as the gluing of TE-structure $(\mathcal{V},\widetilde{\nabla})$ and $\widetilde{T}E$ -structure $(\mathcal{V}^{\dagger},\widetilde{\nabla}^{\dagger})$, the condition is equivalent to $\widetilde{\nabla}_{\widetilde{\mathfrak{v}}}\mathcal{V}\subset\mathcal{V}$ and $\widetilde{\nabla}^{\dagger}_{\widetilde{\mathfrak{v}}}\mathcal{V}^{\dagger}\subset\mathcal{V}^{\dagger}$.

In §5.2.1–5.2.3, we explained that a harmonic bundle with homogeneity of degree m with respect to \mathfrak{v} induces a polarized integrable variation of pure twistor structure of weight 0 which is homogeneous of degree m with respect to \mathfrak{v} . The construction gives an equivalence, which is a variant of the fundamental result due to C. Simpson [45]. (See [17] and [41] for the TERP case and the integrable case.)

Proposition 5.6 Suppose that $(V^{\triangle}, \widetilde{\mathbb{D}}^{\triangle})$ is pure of weight 0 and equipped with a polarization S. Then, the corresponding harmonic bundle is naturally homogeneous of degree m with respect to \mathfrak{v} . The constructions are mutually converse.

Proof Let $p: \mathbb{P}^1 \times X = \mathfrak{X} \longrightarrow X$ denote the projection. Let $(E, \overline{\partial}_E, \theta, h)$ be the harmonic bundle corresponding to the polarized variation of pure twistor structure of weight 0. Let \mathbb{D}^{\triangle} and $\widetilde{\mathbb{D}}_{\mathbb{P}^1}^{\triangle}$ denote the restriction of $\widetilde{\mathbb{D}}^{\triangle}$ to the X-direction and \mathbb{P}^1 -direction, respectively. Let $d_{\mathbb{P}^1}''$ denote the (0,1)-part of $\widetilde{\mathbb{D}}_{\mathbb{P}^1}^{\triangle}$. We have $V^{\triangle} = p^{-1}(E)$ and $\mathbb{D}^{\triangle} = \overline{\partial}_E + \partial_E + \lambda \theta^{\dagger} + \lambda^{-1}\theta$. Note that $[d_{\mathbb{P}^1}'', \widetilde{\mathbb{D}}_{\widetilde{\mathfrak{p}}}^{\triangle}] = 0$. Hence, for any C^{∞} -section s of E, we can define a C^{∞} -section $L_{\mathfrak{p}}^E s$ of E by the condition $\widetilde{\mathbb{D}}_{\widetilde{\mathfrak{p}}}^{\triangle} p^{-1}(s) = p^{-1}(L_{\mathfrak{p}}^E s)$. It is easy to check that $L_{\mathfrak{p}}^E$ gives an action of \mathfrak{p} on E

We have another description of $L^E_{\mathfrak{v}}$. Let $\mathfrak{X}^0 := \{0\} \times X \subset \mathfrak{X}$. For any C^{∞} -section s of $E = V_{|\mathfrak{X}^0}^{\triangle}$, we take any C^{∞} -section \widetilde{s} of V^{\triangle} such that $\widetilde{s}_{|\mathfrak{X}^0} = s$. Then, we have $L^E_{\mathfrak{v}}(s) = \widetilde{\mathbb{D}}^{\triangle}_{\widetilde{\mathfrak{v}}}(\widetilde{s})_{|\mathfrak{X}^0}$.

Let $\mathbb{D}^{\triangle} = \mathbb{D}^{1,0} + \mathbb{D}^{0,1}$ denote the decomposition into the (1,0)-part and (0,1)-part. We have $\left[\widetilde{\mathbb{D}}_{\widetilde{\mathfrak{v}}}^{\triangle}, \mathbb{D}^{0,1}\right] = 0$ and $\left[\widetilde{\mathbb{D}}_{\widetilde{\mathfrak{v}}}^{\triangle}, \lambda \mathbb{D}^{1,0}\right] = \sqrt{-1}m\lambda \mathbb{D}^{1,0}$. By using the above description, we obtain $[L_{\mathfrak{v}}^{E}, \overline{\partial}_{E}] = 0$ and $[L_{\mathfrak{v}}^{E}, \theta] = \sqrt{-1}m\theta$. For C^{∞} -sections u and v of E, we have $S(p^{-1}u, \sigma^{*}p^{-1}v) = h(u, v)$. Because

$$\widetilde{\mathfrak{v}}S(u,\sigma^*v) = S\big(\widetilde{\mathbb{D}}_{\widetilde{\mathfrak{v}}}^{\triangle}u,\sigma^*v\big) + S\big(u,\sigma^*(\widetilde{\mathbb{D}}_{\widetilde{\mathfrak{v}}}^{\triangle}v)\big), \qquad (\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}})h(u,v) = 0,$$

we obtain $L_{\mathfrak{p}}^E h = 0$. It is clear that the constructions are mutually converse.

5.2.5 Appendix

Let $(E, \overline{\partial}_E, \theta, h)$ be a harmonic bundle on X. Suppose that E is equipped with an action of a holomorphic vector field \mathfrak{v} such that $[L^E_{\mathfrak{v}}, \overline{\partial}_E] = 0$, $L^E_{\mathfrak{v}}(h) = 0$ and $[L^E_{\mathfrak{v}}, \theta] = \alpha \theta$ for a complex number $\alpha \notin \sqrt{-1}\mathbb{R}$.

Lemma 5.7 We have $[\overline{\partial}_E, \partial_E] = 0$ and $[\theta, \theta^{\dagger}] = 0$.

Proof For each point of $P \in X$, there exists $\epsilon > 0$, such that $(E, \overline{\partial}_E, e^{s\alpha}\theta, h)$ are harmonic bundles for any $-\epsilon < s < \epsilon$ on a neighbourhood X_P of P. We obtain $[\overline{\partial}_E, \partial_E] + e^{2s \operatorname{Re}(\alpha)}[\theta, \theta^{\dagger}] = 0$. Then, the claim of the lemma follows.

Proposition 5.8 Let P be any point of X. There exist a simply connected neighbourhood X_P of P, holomorphic functions \mathfrak{a}_i $(i = 1, ..., \ell)$ and an orthogonal decomposition $(E, \theta)_{|X_P} = \bigoplus (E_i, d\mathfrak{a}_i \operatorname{id}_{E_i})$.

Proof The Higgs bundle (E, θ) naturally induces a coherent \mathcal{O}_{T^*X} -module. Its support with the reduced structure is called the spectral variety of the Higgs bundle, and denoted by $\Sigma_{E,\theta}$. We say that a point $Q \in X$ is generic with respect to (E, θ) , if the projection $\Sigma_{E,\theta} \longrightarrow X$ is unramified over Q.

Suppose that P is generic with respect to (E,θ) . We take a holomorphic coordinate (z_1,\ldots,z_n) around P. We have the description $\theta = \sum_{i=1}^n f_i dz_i$. Let f_i^{\dagger} denote the adjoint of f_i with respect to h. Because $[f_i,f_j^{\dagger}]=0$ for any i and j according to Lemma 5.7, we have an orthogonal decomposition $E_{|P}=\bigoplus_{\alpha\in\mathbb{C}^n}E_{\alpha}$ such that $f_{i|P}=\bigoplus_{\alpha}\alpha_i\operatorname{id}_{E_{\alpha}}$ and $f_{i|P}^{\dagger}=\bigoplus_{\alpha}\overline{\alpha}_i\operatorname{id}_{E_{\alpha}}$, where α_i denotes the i-th component of α . On a small neighbourhood $X_P\ni P$, we have a unique holomorphic decomposition $E_{|X_P}=\bigoplus_{\alpha\in\mathbb{C}^n}E'_{\alpha}$ such that $f_i(E'_{\alpha})\subset E'_{\alpha}$ and $E'_{\alpha|P}=E_{\alpha}$.

Because P is supposed to be generic with respect to (E,θ) , there exists a holomorphic function $\beta_{\boldsymbol{\alpha},i}$ such that $f_{i|E'_{\boldsymbol{\alpha}}} = \beta_{\boldsymbol{\alpha},i} \operatorname{id}_{E'_{\boldsymbol{\alpha}}}$. Because we have $\partial_E \theta = 0$, we obtain that $\sum_{i=1}^n \beta_{\boldsymbol{\alpha},i} dz_i$ is closed. Hence, there exist holomorphic functions $\mathfrak{b}(\boldsymbol{\alpha})$ such that $f = \bigoplus d\mathfrak{b}(\boldsymbol{\alpha}) \operatorname{id}_{E'_{\boldsymbol{\alpha}}}$. Hence, we are done in the case that P is generic. Note that the decomposition $E_{|X_P} = \bigoplus E'_{\boldsymbol{\alpha}}$ is orthogonal, and hence flat with respect to the Chern connection $\overline{\partial}_E + \partial_E$.

Let us consider the case that P is not necessarily generic. Let X_P be a simply connected small neighbourhood of P. There exists a point $Q \in X_P$ which is generic with respect to (E, θ) . On a small neighbourhood $X_Q \ni Q$ in X_P , we have holomorphic functions \mathfrak{a}_i and an orthogonal decomposition $(E, \theta)_{|X_Q} = \bigoplus (E_i, d\mathfrak{a}_i \operatorname{id}_{E_i})$. Because the decomposition is flat with respect to $\overline{\partial}_E + \partial_E$, we can uniquely extend it to a flat decomposition $E_{|X_P|} = \bigoplus E'_i$, which is orthogonal with respect to h. It is easy to deduce that $\theta(E'_i) \subset E'_i$, and that $\theta_{|E'_i|}$ has a unique eigenvalue $d\mathfrak{a}'_i$.

5.3 Semi-infinite variation of Hodge structure

5.3.1 Real structure and semi-infinite variation of Hodge structure

Let X be a complex manifold. Let $(E, \overline{\partial}_E, \theta, h)$ be a harmonic bundle on X. The dual vector bundle E^{\vee} is naturally equipped with a Higgs field θ^{\vee} and a hermitian metric $h_{E^{\vee}}$, so that $(E^{\vee}, \overline{\partial}_{E^{\vee}}, \theta^{\vee}, h^{\vee})$ is a harmonic bundle. Let C be a non-degenerate symmetric pairing of $(E, \overline{\partial}_E)$, such that the induced map $\Phi_C : E \longrightarrow E^{\vee}$ gives an isomorphism of harmonic bundles, i.e., it is an isometric with respect to h and h^{\vee} , and h is symmetric with respect to h. We have

$$\overline{\partial}C(u,v) = C(\overline{\partial}_E u,v) + C(u,\overline{\partial}_E v), \qquad C(\theta u,v) - C(u,\theta v) = 0.$$

Because Φ_C is an isometry, we have

$$\partial C(u,v) = C(\partial_E u, v) + C(u, \partial_E v), \qquad C(\theta^{\dagger} u, v) - C(u, \theta^{\dagger} v) = 0.$$

Hence, we obtain

$$\overline{\partial}C(u,v) = C((\overline{\partial}_E + \lambda \theta^{\dagger})u, v) + C(u, (\overline{\partial}_E - \lambda \theta^{\dagger})v) = 0$$
(36)

$$\partial C(u,v) = C((\partial_E + \lambda^{-1}\theta)u, v) + C(u, (\partial_E - \lambda^{-1}\theta)v) = 0$$
(37)

Let $j: \mathcal{X} \longrightarrow \mathcal{X}$ be given by $j(\lambda, x) = (-\lambda, x)$. According to (36) and (37), C naturally gives a ∇^{λ} -flat pairing $\mathcal{E} \otimes j_* \mathcal{E} \longrightarrow \mathcal{O}_{\mathcal{X}^*}$.

Let n be an integer. Suppose moreover that C is homogeneous of degree $\sqrt{-1}nm$ with respect to \mathfrak{v} , i.e., $L^E_{\mathfrak{v}}C = \sqrt{-1}nmC$. We have the induced pairing $C: \mathcal{E} \otimes j^*\mathcal{E} \longrightarrow \mathcal{O}_{\mathcal{X}}$. We set $P:=\lambda^{-n}C$. The following lemma is clear by construction.

Lemma 5.9 We have $L_{\widetilde{\mathfrak{p}}}^{\mathcal{E}}P=0$, and $\widetilde{\nabla}P=0$.

The pairing P and the polarization S_h induce a real structure of the local system on $\mathbb{C}^* \times X$ corresponding to $(\mathcal{E}, \widetilde{\nabla})$. (See §5.3.2 below.) Thus, it gives a variation of pure polarized TERP-structure. Because we are given an "Euler field" \mathfrak{v} such that $\mathfrak{v}\mathcal{E} \subset \mathcal{E}$, it is graded $\frac{\infty}{2}$ -VHS with real structure which is pure and polarized. (See [23]. The formulation here is slightly changed.) If the base space is a punctured disc, it is related with a Sabbah orbit or Hertling-Sevenheck orbit, ("nilpotent orbit" in the sense of Hertling and Sevenheck) studied in [18].

5.3.2 Appendix

Symmetric pairing Let V be a finite dimensional \mathbb{C} -vector space. The natural pairing $V^{\vee} \times V \longrightarrow \mathbb{C}$ is denoted by $\langle \cdot, \cdot \rangle$. Let $C: V \times V \longrightarrow \mathbb{C}$ be a bilinear map. It is equivalent to a linear map $\Phi_C: V \longrightarrow V^{\vee}$ given by $\langle \Phi_C(u), v \rangle = C(u, v)$. It is symmetric if and only if the dual $\Phi_C^{\vee}: V \longrightarrow V^{\vee}$ is equal to Φ_C . If C is non-degenerate, a pairing $C^{\vee}: V^{\vee} \times V^{\vee} \longrightarrow \mathbb{C}$ is induced by $C^{\vee}(f,g) = C(\Phi_C^{-1}(f), \Phi_C^{-1}(g))$. Note that we have $\Phi_{C^{\vee}} = \Phi_C^{-1}$, if C is non-degenerate and symmetric. Indeed,

$$\langle \Phi_{C^\vee}(f),\,g\rangle = C^\vee(f,g) = C\left(\Phi_C^{-1}(f),\Phi_C^{-1}(g)\right) = \langle f,\Phi_C^{-1}(g)\rangle = \langle (\Phi_C^{-1})^\vee(f),\,g\rangle.$$

Conjugate Let \overline{V} be the vector space such that $V = \overline{V}$ as a \mathbb{Z} -module, and that the scalar multiplication of $\alpha \in \mathbb{C}$ is given by $\alpha \bullet u = \overline{\alpha} u$. Let \overline{v} denote the element of \overline{V} corresponding to an element $v \in V$. Then, $\alpha \overline{v} = \overline{(\alpha v)}$. We may regard $\overline{(\cdot)}$ as a \mathbb{C} -linear isomorphism $V \simeq \overline{V}$.

We have a natural anti- \mathbb{C} -linear map $\Psi: \mathrm{Hom}_{\mathbb{C}}(\overline{V}, \mathbb{C}) \longrightarrow \mathrm{Hom}_{\mathbb{C}}(\overline{V}, \mathbb{C})$ given by $\Psi(f)(\overline{v}) = \overline{f(v)}$. It naturally gives a \mathbb{C} -isomorphism $\overline{\mathrm{Hom}_{\mathbb{C}}(V, \mathbb{C})} \simeq \mathrm{Hom}_{\mathbb{C}}(\overline{V}, \mathbb{C})$, expressed as $\overline{(V^{\vee})} \longrightarrow \overline{V}^{\vee}$.

We have the natural pairing $\overline{V}^{\vee} \times \overline{V} \longrightarrow \mathbb{C}$. Under the identifications, we have $\langle \overline{f}, \overline{v} \rangle = \overline{\langle f, v \rangle}$.

Hermitian pairing Let $h: V \times \overline{V} \longrightarrow \mathbb{C}$ be a bilinear map, which is equivalent to a linear map $\Phi_h: V \longrightarrow \overline{V}^{\vee}$ given by $\langle \Phi_h(u), \overline{v} \rangle = h(u, v)$. It is hermitian, if and only if $(\overline{\Phi_h})^{\vee} = \Phi_h$. Indeed, we have

$$h(u,v) = \langle \Phi_h(u), \overline{v} \rangle = \langle u, \Phi_h^{\vee}(\overline{v}) \rangle = \overline{\langle \overline{u}, \overline{\Phi}_h^{\vee}(v) \rangle}.$$

Because $\langle \overline{u}, \Phi_h(v) \rangle = h(v, u)$, we are done.

If h is non-degenerate and hermitian, we have the induced hermitian pairing h^{\vee} and \overline{h}^{\vee} on V^{\vee} and \overline{V}^{\vee} , respectively. We have $\Phi_{\overline{h}^{\vee}} = \Phi_h^{-1}$. Indeed,

$$\langle \Phi_{\overline{h}^\vee}(\overline{f}), g \rangle = \overline{h}^\vee(\overline{f}, \overline{g}) = h \left(\Phi_h^{-1}(\overline{f}), \Phi_h^{-1}(\overline{g}) \right) = \langle \overline{f}, \overline{\Phi_h^{-1}(\overline{g})} \rangle = \langle \overline{f}, \overline{\Phi}_h^{-1}(g) \rangle = \langle \Phi_h^{-1}(\overline{f}), g \rangle$$

We also have $\overline{\Phi}_{h^{\vee}} = \Phi_{\overline{h}^{\vee}}$.

Compatibility Let C be a non-degenerate symmetric pairing of V. Let h be a non-degenerate hermitian pairing of V. We say that C and h are compatible, if $h^{\vee}(\Phi_C(u), \Phi_C(v)) = h(u, v)$, i.e., Φ_C gives an isometry $(V, h) \simeq (V^{\vee}, h^{\vee})$. In that case, the following diagram is commutative:

$$\begin{array}{ccc} V & \xrightarrow{\Phi_C} & V^{\vee} \\ & & & \downarrow^{\Phi_h \vee} \\ & \overline{V}^{\vee} & \xrightarrow{\overline{\Phi}_{C^{\vee}}} & \overline{V} \end{array}$$

By the compatibility, we have $h^{\vee}(\Phi_C(u), f) = h(u, \Phi_{C^{\vee}}(f))$. The left hand side is equal to $\langle \Phi_h \vee \Phi_C(u), \overline{f} \rangle$, and the right hand side is equal to $\langle \Phi_h(u), \Phi_{C^{\vee}}(f) \rangle = \langle \overline{\Phi}_{C^{\vee}} \Phi_h(u), \overline{f} \rangle$. Hence, we have $\overline{\Phi}_{C^{\vee}} \circ \Phi_h = \Phi_{h^{\vee}} \circ \Phi_C =: \kappa$, which is a linear map $V \longrightarrow \overline{V}$. We have $\overline{\kappa} \circ \kappa = \mathrm{id}_V$. Indeed, $\overline{\kappa} \circ \kappa = \overline{\Phi}_{h^{\vee}} \circ \overline{\Phi}_C \circ \overline{\Phi}_{C^{\vee}} \circ \Phi_h = \overline{\Phi}_{h^{\vee}} \circ \underline{\Phi}_h = 1$. Hence, κ gives a real structure of the vector space V. In this case, we have the relation $h(u, v) = C(u, \overline{\kappa(v)})$.

Induced vector bundle with real structure on \mathbb{C}^* Let V be a \mathbb{C} -vector space with a non-degenerate hermitian pairing h and a non-degenerate symmetric pairing C. Assume that they are compatible.

Let σ , γ and j be the involutions of \mathbb{C}^*_{λ} , given by $\sigma(\lambda) = -\overline{\lambda}^{-1}$, $\gamma(\lambda) = \overline{\lambda}^{-1}$ and $j(\lambda) = -\lambda$, respectively. We set $\mathcal{V} := V \otimes \mathcal{O}_{\mathbb{C}^*_{\lambda}}$. Let S_h be the pairing $\mathcal{V} \otimes \sigma^* \mathcal{V} \longrightarrow \mathcal{O}_{\mathbb{C}^*}$ given by $S_h(u, \sigma^* v) = h(u, \sigma^* v)$. Let S_C be the pairing $\mathcal{V} \otimes j^* \mathcal{V} \longrightarrow \mathcal{O}_{\mathbb{C}^*}$ given by $S_C(u, j^* v) = C(u, j^* v)$. The pairing S_h satisfies the symmetry $\sigma^* S_h(\sigma^* u, v) = S_h(v, \sigma^* u)$. The pairing S_C satisfies the symmetry $j^* S_C(j^* u, v) = S_C(v, j^* u)$. Let $\Phi_{S_h} : \mathcal{V} \longrightarrow \sigma^* \mathcal{V}^{\vee}$ and $\Phi_{S_C} : \mathcal{V} \longrightarrow j^* \mathcal{V}^{\vee}$ denote the induced morphisms. We have the following commutativity by the compatibility of h and C:

$$\begin{array}{ccc} \mathcal{V} & \stackrel{\Phi_{SC}}{\longrightarrow} & j^* \mathcal{V}^{\vee} \\ & & \downarrow j^* \Phi_{S_h^{\vee}} \\ & & \sigma^* \mathcal{V}^{\vee} & \stackrel{\sigma^* \Phi_{S_{C}^{\vee}}}{\longrightarrow} & \gamma^* \mathcal{V} \end{array}$$

We set $\kappa := j^* \Phi_{S_h^{\vee}} \circ \Phi_{S_C}$ which is a morphism $\mathcal{V} \longrightarrow \gamma^* \mathcal{V}$ such that $\gamma^* \kappa \circ \kappa = \mathrm{id}$.

If we set $P = \lambda^{-n} S_C$, then we have the following $(-1)^n$ -symmetry:

$$j^*(P)(j^*u, v) = (-1)^n P(v, j^*v)$$

We have $P^{\vee} = (-1)^n \lambda^n S_C^{\vee}$. We have $\Phi_P = \lambda^{-n} \Phi_{S_C}$ and $\sigma^* \Phi_{P^{\vee}} = \sigma^* \left((-1)^n \lambda^n \Phi_{S_C^{\vee}} \right) = \lambda^{-n} \sigma^* \Phi_{S_C^{\vee}}$. Hence, the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{V} & \stackrel{\Phi_P}{\longrightarrow} & j^* \mathcal{V}^{\vee} \\ & & \downarrow j^* \Phi_{S_h^{\vee}} \\ & & \sigma^* \mathcal{V}^{\vee} & \stackrel{\sigma^* \Phi_{P^{\vee}}}{\longrightarrow} & \gamma^* \mathcal{V} \end{array}$$

We set $\kappa_n := j^* \Phi_{S_h^{\vee}} \circ \Phi_P = \lambda^{-n} \kappa$. We have $\gamma^* \kappa_n \circ \kappa_n = \mathrm{id}$. Hence, it also gives a real structure.

Assume that \mathcal{V} is equipped with a connection ∇ such that the pairings S_h and P are ∇ -flat. Then, κ is also flat. As in [17], it determines a real structure of the corresponding local system. See §2 of [33] for a review.

5.4 The prolongation of the action

Let X be a complex manifold with a simply normal crossing hypersurface D. Let $D = \bigcup_{i \in \Lambda} D_i$ denote the irreducible decomposition. Let \mathfrak{v} be a holomorphic vector field on $X \setminus D$ such that $\mathfrak{v}^{1,0} \in \Theta_X(\log D)$. Let $(E, \overline{\partial}_E, \theta, h)$ be a good wild harmonic bundle on $X \setminus D$ which is homogeneous of degree m with respect to \mathfrak{v} .

We set $\mathcal{X} := \mathbb{C}_{\lambda} \times X$ and $\mathcal{D} := \mathbb{C}_{\lambda} \times D$. As explained in §5.2, we have the holomorphic vector bundle \mathcal{E} on \mathcal{X} with a meromorphic flat connection $\widetilde{\nabla}$. According to [31] and [36] (see [33] for a review), \mathcal{E} is naturally extended to a filtered vector bundle $\mathcal{Q}_*\mathcal{E} = (\mathcal{Q}_a\mathcal{E} \mid a \in \mathbb{R}^{\Lambda})$ on $(\mathcal{X}, \mathcal{D})$. Let $\mathcal{Q}\mathcal{E}$ denote the locally free $\mathcal{O}_{\mathcal{X}}(*\mathcal{D})$ -module $\bigcup_a \mathcal{Q}_a \mathcal{E}$. According to [19] in the tame case and [33] in the wild case, $\widetilde{\nabla}$ gives a meromorphic flat connection of $\mathcal{Q}\mathcal{E}$:

$$\widetilde{\nabla}: \mathcal{QE} \longrightarrow \mathcal{QE} \otimes \Omega^1_{\mathcal{X}}(\log \mathcal{X}^0) \otimes \mathcal{O}_{\mathcal{X}}(\mathcal{X}^0).$$

Here, $\mathcal{X}^0 := \{0\} \times X$.

We have a \mathbb{C} -endomorphism $L^{\mathcal{Q}\mathcal{E}}_{\widetilde{\mathfrak{v}}}$ of $\mathcal{Q}\mathcal{E}$ given by $L^{\mathcal{Q}\mathcal{E}}_{\widetilde{\mathfrak{v}}}:=\widetilde{\nabla}_{\widetilde{\mathfrak{v}}}$. It is an extension of $L^{\mathcal{E}}_{\widetilde{\mathfrak{v}}}$ on \mathcal{E} . It satisfies $L^{\mathcal{Q}\mathcal{E}}_{\widetilde{\mathfrak{v}}}(fs)=\widetilde{\mathfrak{v}}(f)\,s+f\,L^{\mathcal{Q}\mathcal{E}}_{\widetilde{\mathfrak{v}}}(s)$ for local sections $f\in\mathcal{O}_{\mathcal{X}}(*\mathcal{D})$ and $s\in\mathcal{Q}\mathcal{E}$.

Proposition 5.10 For any $a \in \mathbb{R}^{\Lambda}$, we have $L_{\widetilde{\mathfrak{p}}}^{\mathcal{QE}}(\mathcal{Q}_a \mathcal{E}) \subset \mathcal{Q}_a \mathcal{E}$.

Proof We have only to consider the case $X = \{(z_1, \ldots, z_n) \mid |z_i| < 1\}$ and $D = \bigcup_{i=1}^{\ell} D_i$, where $D_i = \{z_i = 0\}$. Let us consider the case that the harmonic bundle $(E, \overline{\partial}_E, \theta, h)$ is unramifiedly good wild. Let O denote the origin $(0, \ldots, 0)$. We have a good set $\mathcal{I} \subset \mathcal{O}_X(*D)/\mathcal{O}_X$ and a formal decomposition

$$\left(\mathcal{Q}_*\mathcal{E},\widetilde{\nabla}\right)_{|\widehat{\mathbb{C}_\lambda\times\{O\}}} = \bigoplus_{\mathfrak{a}\in\mathcal{I}} \left(\mathcal{Q}_*\widehat{\mathcal{E}}_{\mathfrak{a}},\widehat{\nabla}_{\mathfrak{a}}\right)$$

such that $\widehat{\nabla}_{\mathfrak{a}}^{\mathrm{reg}} := \widehat{\nabla}_{\mathfrak{a}} - d(\widetilde{\mathfrak{a}}/\lambda) \operatorname{id}_{\mathcal{Q}\widehat{\mathcal{E}}_{\mathfrak{a}}}$ are logarithmic along \mathcal{D} , where we choose lifts $\widetilde{\mathfrak{a}} \in \mathcal{O}_X(*D)$ of \mathfrak{a} . Around any point (λ_0, O) with $\lambda_0 \neq 0$, for any $a \in \mathbb{R}^\ell$, we have $\widehat{\nabla}_{\mathfrak{a}}^{\mathrm{reg}} \mathcal{Q}_a \widehat{\mathcal{E}}_{\mathfrak{a}} \subset \mathcal{Q}_a \widehat{\mathcal{E}}_{\mathfrak{a}} \otimes \Omega^1_{\mathcal{X}}(\log \mathcal{D})$, and hence $\widehat{\nabla}_{\mathfrak{a},\widetilde{\mathfrak{p}}}^{\mathrm{reg}} (\mathcal{Q}_a \widehat{\mathcal{E}}_{\mathfrak{a}}) \subset \mathcal{Q}_a \widehat{\mathcal{E}}_{\mathfrak{a}}$. Because $[L_{\mathfrak{v}}^E, \theta] = \sqrt{-1}m\theta$, we obtain that $\mathfrak{v}(\widetilde{\mathfrak{a}}) - \sqrt{-1}m\widetilde{\mathfrak{a}}$ is holomorphic. We obtain that $d(\widetilde{\mathfrak{a}}/\lambda)(\widetilde{\mathfrak{v}})$ is holomorphic around (λ_0, O) . Hence, we obtain that $\widehat{\nabla}_{\mathfrak{a},\widetilde{\mathfrak{v}}}(\mathcal{Q}_a\widehat{\mathcal{E}}_{\mathfrak{a}}) \subset \mathcal{Q}_a\widehat{\mathcal{E}}_{\mathfrak{a}}$ around (λ_0, O) . We can conclude that $L_{\widetilde{\mathfrak{v}}}^{\mathcal{Q}\mathcal{E}}(\mathcal{Q}_a\mathcal{E}) \subset \mathcal{Q}_a\mathcal{E}$ outside \mathcal{D}^0 . By using the Hartogs property, we obtain $L_{\widetilde{\mathfrak{v}}}^{\mathcal{Q}\mathcal{E}}(\mathcal{Q}_a\mathcal{E}) \subset \mathcal{Q}_a\mathcal{E}$ on \mathcal{X} .

Let us consider the case that $(E, \overline{\partial}_E, \theta, h)$ is not necessarily unramified. We set $X' = \{(\zeta_1, \dots, \zeta_n) \mid |\zeta_i| < 1\}$ and $D' = \bigcup_{i=1}^{\ell} \{\zeta_i = 0\}$. Let $\varphi: (X', D') \longrightarrow (X, D)$ be the ramified covering given by $\varphi(\zeta_1, \dots, \zeta_n) = (\zeta_1^e, \dots, \zeta_\ell^e, \zeta_{\ell+1}, \dots, \zeta_n)$ such that $(E', \overline{\partial}_{E'}, \theta', h') := \varphi^{-1}(E, \overline{\partial}_E, \theta, h)$ is unramified. Let \mathfrak{v}' be the holomorphic vector field of $X' \setminus D'$ which is the lift of \mathfrak{v} . The harmonic bundle $(E', \overline{\partial}_{E'}, \theta', h')$ is homogeneous of degree $\sqrt{-1}m$ with respect to \mathfrak{v}' . We have $L_{\widetilde{\mathfrak{v}}'}^{\mathcal{Q}\mathcal{E}'}(\mathcal{Q}_{\mathbf{a}'}\mathcal{E}') \subset \mathcal{Q}_{\mathbf{a}'}\mathcal{E}'$ for any $\mathbf{a}' \in \mathbb{R}^{\ell}$. Note that a section f of $\mathcal{Q}\mathcal{E}$ is contained in $\mathcal{Q}_{\mathbf{a}}\mathcal{E}$ if and only if $\varphi^*f \in \mathcal{Q}_{ea}\mathcal{E}'$. Then, we obtain $L_{\widetilde{\mathfrak{v}}}^{\mathcal{Q}\mathcal{E}}(\mathcal{Q}_{\mathbf{a}}\mathcal{E}) \subset \mathcal{Q}_{\mathbf{a}}\mathcal{E}$.

5.5 tr-TLE structure

5.5.1 Limit mixed twistor structure

Let $X := \{(z_1, \ldots, z_n) \mid |z_i| < 1\}$ and $D := \bigcup_{i=1}^{\ell} \{z_i = 0\}$. Let \mathfrak{v} be a holomorphic vector field of $X \setminus D$ such that $\mathfrak{v}^{1,0} \in \Theta_X(\log D)$. Let $(E, \overline{\partial}_E, \theta, h)$ be a *tame* harmonic bundle on $X \setminus D$ which is homogeneous of degree $\sqrt{-1m}$ with respect to \mathfrak{v} .

For any $a \in \mathbb{R}^{\ell}$, we have the limit mixed twistor structure $(S_a^{\text{can}}(E), \mathbf{N})$. (See [31]. See [33] for a review.) Let us briefly recall the construction. We have the following coherent \mathcal{O} -module on $\mathbb{C}_{\lambda} \times \bigcap_{i=1}^{\ell} \{z_i = 0\}$:

$$\operatorname{Gr}_{\boldsymbol{a}}^{\mathcal{Q}}(\mathcal{E}) := \frac{\mathcal{Q}_{\boldsymbol{a}}(\mathcal{E})}{\sum_{\boldsymbol{b} \leq \boldsymbol{a}} \mathcal{Q}_{\boldsymbol{b}} \mathcal{E}}$$

Here, $\mathbf{b} \leq \mathbf{a}$ for $\mathbf{b} = (b_i), \mathbf{a} = (a_i) \in \mathbb{R}^{\ell}$ means $b_i \leq a_i$ for any i and $\mathbf{b} \neq \mathbf{a}$. According to [31], $\operatorname{Gr}_{\mathbf{a}}^{\mathcal{Q}}(\mathcal{E})$ is locally free. Hence, we have the vector bundle $\mathcal{G}_{\mathbf{a}}(E) := \operatorname{Gr}_{\mathbf{a}}^{\mathcal{Q}}(\mathcal{E})|_{\mathbb{C}_{\lambda} \times \{O\}}$ on \mathbb{C}_{λ} . They are equipped with the morphisms $\mathcal{N}_i : \mathcal{G}_{\mathbf{a}}(E) \longrightarrow \lambda^{-1}\mathcal{G}_{\mathbf{a}}(E)$ obtained as the nilpotent part of the residues $\operatorname{Res}_{z_i}(\widetilde{\nabla})$. We also obtain the vector bundle $\mathcal{G}_{-\mathbf{a}}^{\dagger}(E)$ on \mathbb{C}_{μ} with the morphisms $\mathcal{N}_i^{\dagger} : \mathcal{G}_{-\mathbf{a}}^{\dagger}(E) \longrightarrow \mu^{-1}\mathcal{G}_{-\mathbf{a}}^{\dagger}(E)$ from $(\mathcal{Q}_*\mathcal{E}^{\dagger}, \widetilde{\nabla}^{\dagger})$. We have an isomorphism $\Phi_{\mathbf{a}} : \mathcal{G}_{\mathbf{a}}(E)|_{\mathbb{C}_{\lambda}^*} \simeq \mathcal{G}_{-\mathbf{a}}^{\dagger}(E)|_{\mathbb{C}_{\mu}^*}$, such that $\Phi_{\mathbf{a}} \circ \mathcal{N}_i = -\mathcal{N}_i^{\dagger} \circ \Phi_{\mathbf{a}}$. (See [33] for a review of the construction.) Thus, we obtain a vector bundle $S_{\mathbf{a}}^{\operatorname{can}}(E)$ on \mathbb{P}^1 with morphisms $\mathcal{N}_i : S_{\mathbf{a}}^{\operatorname{can}}(E) \longrightarrow S_{\mathbf{a}}^{\operatorname{can}}(E) \otimes \mathcal{O}_{\mathbb{P}^1}(2)$. Let W denote the weight filtration of $\sum_{i=1}^{\ell} \mathcal{N}_i$. According to [31], $(S_{\mathbf{a}}^{\operatorname{can}}(E), W)$ is a mixed twistor structure, i.e., $\operatorname{Gr}_n^W S_{\mathbf{a}}^{\operatorname{can}}(E)$ are isomorphic to a direct sum of $\mathcal{O}_{\mathbb{P}^1}(n)$.

Because the harmonic bundle is homogeneous, the associated polarized variation of pure twistor structure is integrable (§5.2). As explained in [19] (see [33] for a review), $S_{\boldsymbol{a}}^{\operatorname{can}}(E)$ is equipped with the induced meromorphic connection ${}^{\boldsymbol{a}}\!\nabla$ in the case of polarized integrable variation of twistor structure. Let $\mathfrak{u} := \sqrt{-1}(\lambda\partial_{\lambda} - \overline{\lambda}\partial_{\overline{\lambda}})$. We also have an action $L_{m\mathfrak{u}}^{S^{\operatorname{can}}}$ on $S_{\boldsymbol{a}}^{\operatorname{can}}(E)$ induced by $L_{\overline{\mathfrak{v}}}^{\mathcal{QE}}$ (§5.4). It is equal to ${}^{\boldsymbol{a}}\!\nabla_{m\mathfrak{u}} + \sum f_i(a_i + \mathcal{N}_i)$, where $\mathfrak{v}_{|O} = \sum f_i \left(z_i \partial_{z_i}\right)_{|O}$. The morphisms \mathcal{N}_i are compatible with ${}^{\boldsymbol{a}}\!\nabla$ and $L_{m\mathfrak{u}}^{S^{\operatorname{can}}}$.

5.5.2 Mixed Hodge structure

Note that \mathfrak{u} is the fundamental vector field of the S^1 -action on \mathbb{P}^1 given by $(t,\lambda) \longmapsto t\lambda$. For our purpose in §6, it is reasonable to suppose the following.

Assumption For any $a \in \mathbb{R}^{\ell}$, the \mathfrak{u} -action on $S_a^{\text{can}}(E)$ is lifted to an S^1 -action.

It is naturally extended to a \mathbb{C}^* -action. (See Lemma 5.17 below.) The connection ${}^a\nabla$ and the morphisms \mathcal{N}_i are \mathbb{C}^* -invariant. In particular, the weight filtration is also \mathbb{C}^* -equivariant. Hence, $(S_a^{\text{can}}(E), W)$ comes from a complex mixed Hodge structure (H, F, G, W) with morphisms N_i , by the Rees construction. ([45] . See also [31].)

Recall that, by using the canonical decomposition of mixed Hodge structure by Deligne, we obtain a decreasing filtration F^{op} such that (i) it is 0-opposite to F, (ii) $N_i(F^{\text{op}})^j \subset (F^{\text{op}})^{j+1}$ for any $i = 1, \ldots, \ell$ and $j \in \mathbb{Z}$. Let

us briefly review it. We have the Deligne decomposition $H = \bigoplus_{p,q \in \mathbb{Z}} I^{p,q}$ which satisfies $F^p = \bigoplus_{r \geq p} I^{r,s}$ and $W_k = \bigoplus_{r+s \leq k} I^{r,s}$. Moreover, for any morphism of mixed Hodge structures $f: (H,F,G,W) \longrightarrow (H',F,G,W)$, we have $f(I^{r,s}(H)) \subset I^{r,s}(H')$. (See [37], for example.) We set $(F^{\text{op}})^q := \bigoplus_{q+r \leq 0} I^{r,s}$. Then, the filtrations F and F^{op} are 0-opposed, i.e., $F^p \cap (F^{\text{op}})^q = 0$ for p+q>0 and $H = \bigoplus_{p+q=0} F^p \cap F^q$. Because N_i $(i=1,\ldots,\ell)$ give morphisms from (H,F,G,W) to its (-1)-Tate twist, we obtain that $N_i((F^{\text{op}})^q) \subset (F^{\text{op}})^{q+1}$ by using the functoriality.

Applying the Rees construction to two filtrations F and F^{op} , we obtain a vector bundle $\mathcal{R}_{\boldsymbol{a}}$ on \mathbb{P}^1 , which is pure twistor structure of weight 0. It satisfies $\mathcal{R}_{\boldsymbol{a}|\mathbb{C}_{\lambda}} = S_{\boldsymbol{a}|\mathbb{C}_{\lambda}}^{\text{can}}$. We have induced morphisms

$$\mathcal{N}_i: \mathcal{R}_{\boldsymbol{a}} \longrightarrow \mathcal{R}_{\boldsymbol{a}} \otimes \mathcal{O}_{\mathbb{P}^1}(\{0\} - \{\infty\}), \quad (i = 1, \dots, \ell).$$

The bundle \mathcal{R}_a is equipped with a \mathbb{C}^* -action, which induces an action $L_{m\mathfrak{u}}^{\mathcal{R}_a}$ of $m\mathfrak{u}$. Because we have ${}^a\nabla_{m\mathfrak{u}} = L_{m\mathfrak{u}}^{\mathcal{R}_a} - \sum f_i(a_i + \mathcal{N}_i)$ on \mathbb{C}_{λ} , we have the induced connections

$${}^{\boldsymbol{a}}\nabla: \mathcal{R}_{\boldsymbol{a}} \longrightarrow \mathcal{R}_{\boldsymbol{a}} \otimes \Omega^1_{\mathbb{P}^1}(2\{0\} + \{\infty\}).$$

The residue of ${}^{a}\nabla$ at ∞ is semisimple.

Remark 5.11 Hertling and Sevenheck [18] obtained a mixed Hodge structure from their Sabbah orbit based on the above theorem that $(S_a^{can}(E), \mathbf{N})$ are mixed twistor structure in [31]. It seems related with the above mixed Hodge structure in the case that $n = \ell = m = 1$ and $\mathfrak{v} = \sqrt{-1}(z\partial_z - \overline{z}\partial_{\overline{z}})$. But, the construction in this paper is slightly different from theirs, at least apparently. Namely, we construct a mixed Hodge structure on the $L_{\mathfrak{u}}$ -equivariant sections, and they constructed a mixed Hodge structure on the space of multi-valued sections which are flat with respect to ${}^{\alpha}\nabla$. Their construction seems to have its origin in Singularity theory. Our construction might be natural from the view point of "Simpson's Meta theorem" [45]. We postpone to clarify the precise relation.

5.5.3 tr-TLE structure

We set $\mathfrak{X} := \mathbb{P}^1 \times X$ and $\mathfrak{D} := \mathbb{P}^1 \times D$. We set $\mathfrak{X}^{\infty} := \{\infty\} \times X$. From $(\mathcal{Q}_0 \mathcal{E}, \widetilde{\nabla})$, we obtain an $\mathcal{O}_{\mathfrak{X}}(*\mathfrak{X}^{\infty})$ -module $\widetilde{\mathcal{Q}}_0 \mathcal{E}$ such that $\widetilde{\nabla}$ is regular singular along \mathfrak{X}^{∞} . We continue to suppose the assumption in §5.5.2. We regard \mathbb{P}^1 as $\mathbb{C}_{\lambda} \cup \mathbb{C}_{\mu}$ by $\lambda \mu = 1$. We have the natural isomorphism $\widetilde{\mathcal{Q}}_0 \mathcal{E}_{|\mathbb{C}_{\mu}^* \times \{O\}} \simeq \bigoplus_{\boldsymbol{a} \in]-1,0]^{\ell}} \mathcal{R}_{\boldsymbol{a}|\mathbb{C}_{\mu}^*}$ which is the simultaneous generalized eigen decomposition of $\mathrm{Res}_{z_i}(\widetilde{\nabla})$ $(i=1,\ldots,\ell)$. We naturally regard $\bigoplus_{\boldsymbol{a} \in]-1,0]^{\ell}} \mathcal{R}_{\boldsymbol{a}|\mathbb{C}_{\mu}}$ as an $\mathcal{O}_{\mathbb{C}_{\mu}}$ -submodule $\widetilde{\mathcal{Q}}_0 \mathcal{E}_{|\mathbb{C}_{\mu} \times \{O\}}$. It determines a locally free $\mathcal{O}_{\mathbb{P}^1}$ -submodule $\mathcal{R} \subset \widetilde{\mathcal{Q}}_0 \mathcal{E}_{|\mathbb{P}^1 \times \{O\}}$.

Lemma 5.12 \mathcal{R} is a pure twistor structure of weight 0.

Proof For any $-1 < b \le 0$, let $\mathbf{b}_i \in \mathbb{R}^{\ell}$ denote the element whose j-th entry is b if j = i, or 0 otherwise. We have the naturally defined map $\mathcal{Q}_{\mathbf{b}_i}\mathcal{E}_{|\mathbb{C}_{\lambda}\times\{O\}} \longrightarrow \mathcal{Q}_0\mathcal{E}_{|\mathbb{C}_{\lambda}\times\{O\}}$. The image is denoted by ${}^iF_b(\mathcal{Q}_0\mathcal{E}_{|\mathbb{C}_{\lambda}\times\{O\}})$. Thus, we obtain the filtrations iF $(i = 1, \ldots, \ell)$. We set ${}^\ell\!F_a := \bigcap_{i=1}^{\ell} ({}^iF_{a_i})$. According to [31], we have an isomorphism

$$\mathcal{G}_{a}(E) \simeq \frac{\frac{\ell}{F_{a}}(\mathcal{Q}_{0}\mathcal{E}_{|\mathbb{C}_{\lambda}\times\{O\}})}{\sum_{b \leq a} \ell F_{b}(\mathcal{Q}_{0}\mathcal{E}_{|\mathbb{C}_{\lambda}\times\{O\}})},$$

and we have a splitting ${}^{\ell}\!F_{a} \simeq \bigoplus_{b \leq a} \mathcal{G}_{a}(E)$. The filtrations ${}^{i}\!F$ naturally induces filtrations ${}^{i}\!F$ of \mathcal{R} , and we have an isomorphism

$$\mathcal{R}_{a} \simeq \frac{\ell F_{a} \mathcal{R}}{\sum_{b \leq a} \ell F_{b} \mathcal{R}}.$$

In other words, for some N > 0, we can order a_1, \ldots, a_N and we can take a filtration \mathcal{F}_j $(j = 1, \ldots, N)$ of \mathcal{R} such that $\mathcal{F}_j/\mathcal{F}_{j-1} \simeq \mathcal{R}_{a_j}$. Because \mathcal{R}_a are pure twistor structure of weight 0, by construction, we obtain that \mathcal{R} is also a pure twistor structure of weight 0.

Proposition 5.13 We have a locally free $\mathcal{O}_{\mathfrak{X}}$ -submodule $\mathfrak{Q}_0\mathcal{E} \subset \widetilde{\mathcal{Q}}_0\mathcal{E}$ such that (i) $\mathfrak{Q}_0\mathcal{E}_{|\mathbb{P}^1\times\{O\}} = \mathcal{R}$, (ii) $\widetilde{\nabla}$ is logarithmic along \mathfrak{X}^{∞} with respect to $\mathfrak{Q}_0\mathcal{E}$.

Proof It follows from the correspondence between regular filtered flat bundles and filtered local systems ([44], [32]). It is also obtained by successive use of the elementary transforms. We shall give an outline of the latter argument. We have a meromorphic flat bundle $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})$ such that (i) $\widetilde{\mathcal{Q}}\mathcal{E}_{|\mathbb{C}_{\lambda}\times X}=\mathcal{Q}\mathcal{E}$, (ii) $\widetilde{\nabla}$ is regular singular along \mathfrak{X}^{∞} . We have $\mathfrak{X}=(\mathbb{C}_{\mu}\times X)\cup(\mathbb{C}_{\lambda}\times X)$ by $\mu=\lambda^{-1}$. We have only to construct a submodule of $\mathcal{V}:=\widetilde{\mathcal{Q}}\mathcal{E}_{|\mathbb{C}_{\mu}\times X}$ with the desired property. Let M be the Deligne lattice of $(\mathcal{V},\widetilde{\nabla})$. We have $M_{|\mathbb{C}_{\mu}^*\times X}=\mathcal{Q}_0\mathcal{E}_{|\mathbb{C}_{\mu}^*\times X}$ and $M_{|\mathbb{C}_{\mu}\times \{O\}}\otimes\mathcal{O}_{\mathbb{C}_{\mu}}(*0)=\widetilde{\mathcal{Q}}_0\mathcal{E}_{|\mathbb{C}_{\mu}\times \{O\}}=:\mathcal{U}$. The induced connection of \mathcal{U} is denoted by $\nabla_{\mathcal{U}}$. It is equal to $\bigoplus({}^{\alpha}\nabla)$.

We recall a general procedure called elementary transform. Let $Y:=\{(w_1,\ldots,w_n)\,|\,|w_i|<1\},\ D_Y:=\bigcup_{i=1}^\ell\{w_i=0\}$ and $Y_1:=\{w_1=0\}$. Let V be a holomorphic vector bundle on Y with a logarithmic flat connection ∇ . The holomorphic vector bundle $V_1:=V/w_1V$ on Y_1 is equipped with the endomorphism $\mathrm{Res}_{w_1}(\nabla)$ obtained as the residue with respect to w_1 . It is also equipped with an induced logarithmic flat connection ∇_1 , i.e., for a holomorphic local section s of V_1 , take a holomorphic local section s of V such that $s_{|Y_1}=s$, and $\nabla_1 s=\sum_{j=2}^n \widetilde{s}_{j|Y_1}\,dw_j$, where $\nabla \widetilde{s}=\sum_{j=1}^n \widetilde{s}_jdw_j$. Note that $V\otimes_{\mathcal{O}_Y}\mathcal{O}_Y(Y_1)$ is equipped with the induced logarithmic connection. Let $H\subset V_1$ be a subbundle, which is invariant with respect to $\mathrm{Res}_{w_1}(\nabla)$ and ∇_1 . We obtain a holomorphic vector bundles V_{Hi} (i=1,2) by the following exact sequences of \mathcal{O}_Y -modules:

$$0 \longrightarrow V_{H1} \longrightarrow V \longrightarrow V_1/H \longrightarrow 0 \tag{38}$$

$$0 \longrightarrow V_{H2} \longrightarrow V \otimes \mathcal{O}_Y(Y_1) \longrightarrow (V_1/H) \otimes \mathcal{O}_Y(Y_1) \longrightarrow 0$$
(39)

It is easy to see that V_{Hi} have induced logarithmic flat connections ∇ . They are called the elementary transforms of V along H of type (38) and (39), respectively. The kernel of the surjective morphisms $V_{H1}/w_1V_{H1} \longrightarrow H$ and $V_{H2}/w_1V_{H2} \longrightarrow H \otimes_{\mathcal{O}_Y} \mathcal{O}_Y(Y_1)$ are denoted by K_1 and K_2 . Then, V is the elementary transform of V_{H1} (resp. V_{H2}) along K_1 (resp. K_2) of type (39) (resp. (38)).

Let us return to the proof of Proposition 5.13. We have locally free $\mathcal{O}_{\mathbb{C}_{\mu}}$ -submodules $\mathcal{R}_i \subset \mathcal{U}$ (i = 0, ..., N) such that (i) $\mathcal{R}_0 = \mathcal{R}$, (ii) $\mathcal{R}_N = M_{|\mathbb{C}_{\mu} \times \{O\}}$, (iii) \mathcal{R}_i is the elementary transform of \mathcal{R}_{i-1} along $\operatorname{Res}(\nabla_{\mathcal{U}})$ -invariant subspaces $H_i \subset \mathcal{R}_{i-1}/\mu\mathcal{R}_{i-1}$. More precisely, H_i are direct sums of some generalized eigen spaces of $\operatorname{Res}(\nabla_{\mathcal{U}})$. We have the endomorphisms F_i of \mathcal{R} induced by $\operatorname{Res}_i(\widetilde{\nabla})$. By an induction, \mathcal{R}_i are equipped with induced endomorphisms F_i , and H_i are F_i -invariant.

We have either one of the morphisms $\mathcal{R}_i/\mu\mathcal{R}_i \longrightarrow H_i$ in the case (38) or $\mathcal{R}_i/\mu\mathcal{R}_i \longrightarrow H_i \otimes_{\mathcal{O}_{\mathbb{C}_{\mu}}} \mathcal{O}_{\mathbb{C}_{\mu}}(0)$ in the case (39). The kernel is denoted by K_i . They are invariant with respect to F_i and $\operatorname{Res}(\nabla_{\mathcal{U}})$, and we obtain \mathcal{R}_{i-1} from \mathcal{R}_i by applying an elementary transform along K_i of type (39) or (38), respectively.

We set $M_N := M$. We have a logarithmic connection $\widetilde{\nabla}'$ on $M_N' := M_N/\mu M_N$. Its residues at $(0,O) \in \mathbb{C}_{\mu} \times X$ along z_i are given by F_i . Note that M_N' is the Deligne lattice. We have the subspace $K_N \subset \mathcal{R}_N \simeq M_N'/(z_1,\ldots,z_n)M_N'$. Because it is invariant with respect to F_i , we have $\widetilde{\nabla}'$ -invariant subbundle $\widetilde{K}_N' \subset M_N'$. It is $\operatorname{Res}_{\mu}(\widetilde{\nabla})$ -invariant. If \mathcal{R}_N is the elementary transform of \mathcal{R}_{N-1} of type (38) (resp. (39)), let M_{N-1} be the elementary transform of M_N along \widetilde{K}_N' of type (39) (resp. (38)). By a descending induction with a similar procedure, we construct M_i ($i=0,\ldots,N$). Then, M_0 has the desired property.

By Lemma 5.12, there exists a neighbourhood $U \subset X$ of O such that, for each $P \in U$, the restriction $\mathfrak{Q}_0\mathcal{E}_{|\mathbb{P}^1_{\lambda}\times\{P\}}$ is pure twistor structure of weight 0. We state it as a proposition.

Proposition 5.14 There exists a neighbourhood $U \subset X$ of O such that $(\mathcal{Q}_0 \mathcal{E}, \widetilde{\nabla})_{|\mathbb{C}_{\lambda} \times (U \setminus D)}$ is extended to a tr-TLE structure $(\mathfrak{Q}_0 \mathcal{E}, \widetilde{\nabla})_{|\mathbb{C}_{\lambda} \times (U \setminus D)}$ in the sense of Hertling [17].

Lemma 5.15 The action $L_{\widetilde{\mathfrak{v}}}^{\mathcal{Q}_0\mathcal{E}}$ on $\mathcal{Q}\widetilde{\mathcal{E}}$ is extended to an action $L_{\widetilde{\mathfrak{v}}}^{\mathcal{Q}_0\mathcal{E}}$ on $\mathfrak{Q}_0\widetilde{\mathcal{E}}$.

Proof We define $L^{\mathfrak{Q}_0\mathcal{E}}_{\widetilde{\mathfrak{p}}} := \widetilde{\nabla}_{\widetilde{\mathfrak{p}}}$. Because $\widetilde{\nabla}$ is logarithmic along \mathfrak{X}^{∞} , we obtain that $L^{\mathfrak{Q}_0\mathcal{E}}_{\widetilde{\mathfrak{p}}}\mathfrak{Q}_0\mathcal{E} \subset \mathfrak{Q}_0\mathcal{E}$.

Let $\beta \in \mathbb{C}$, and let $\mathfrak{v}_1 := \beta \mathfrak{v}^{1,0} + \overline{\beta} \mathfrak{v}^{0,1}$. We set $\widetilde{\mathfrak{v}}_1 := \beta \widetilde{\mathfrak{v}}^{1,0} + \overline{\beta} \widetilde{\mathfrak{v}}^{0,1}$. Let P be any point of $U \setminus D$, and let γ_t be an integral curve of \mathfrak{v}_1 starting from P such that $\gamma_0 = P$. The holomorphic vector field induces an isomorphism $\mathbb{P}^1_{\lambda} \times \{P\} \simeq \mathbb{P}^1_{\lambda} \times \{\gamma_t\}$. We have an induced isomorphism $(\mathfrak{Q}\mathcal{E}, \widetilde{\nabla})_{|\mathbb{P}^1_{\lambda} \times \{P\}} \simeq (\mathfrak{Q}\mathcal{E}, \widetilde{\nabla})_{|\mathbb{P}^1_{\lambda} \times \{\gamma_t\}}$. (See the proof of Lemma 5.17 below.) Hence, $\mathfrak{Q}\mathcal{E}_{|\mathbb{P}^1_{\lambda} \times \{\gamma_t\}}$ is also a pure twistor structure of weight 0.

Corollary 5.16 In Proposition 5.14, U can be an open subset in X which is preserved by the flow by $\beta v^{1,0} + \overline{\beta} v^{0,1}$ for any $\beta \in \mathbb{C}$.

5.5.4 S^1 -action and \mathbb{C}^* -action (Appendix)

We will be mainly interested in the case that the vector field \mathfrak{v} is obtained as the fundamental vector field of an S^1 -action. So, as an appendix, we give a remark that an S^1 -action on a holomorphic vector bundle is extended to a \mathbb{C}^* -action.

Let Y be a complex manifold equipped with a holomorphic \mathbb{C}^* -action ρ . The restriction of ρ to S^1 is denoted by ρ_1 . Let E be a holomorphic vector bundle on Y with an S^1 -action ρ_{E1} such that (i) (E, ρ_{E1}) is an S^1 -equivariant vector bundle over (Y, ρ_1) , (ii) the fundamental vector field of the action ρ_{E1} is holomorphic. In this subsection, we shall identify TY and $T^{1,0}Y$.

Lemma 5.17 The S^1 -action ρ_{E1} is uniquely extended to a holomorphic \mathbb{C}^* -action ρ_E on E such that (E, ρ_E) is a \mathbb{C}^* -vector bundle on (Y, ρ) .

Proof Let \mathbb{P}_E denote the projective completion of E. The S^1 -action ρ_{E1} on E is uniquely extended to an S^1 -action of ρ_{E2} on \mathbb{P}_E , whose fundamental vector field V is also holomorphic. We obtain the holomorphic vector field $\sqrt{-1}V$, and we have $[V, \sqrt{-1}V] = 0$.

Let ν be the \mathbb{R} -action on Y given by $\nu(t,Q) = \rho(e^{-t},Q)$. Let V_1 denote the fundamental vector field of ν . Let $\pi: \mathbb{P}_E \longrightarrow Y$ denote the projection. Let $T_P\pi: T_P\mathbb{P}_E \longrightarrow T_{\pi(P)}Y$ denote its derivative. Then, $T_P\pi(\sqrt{-1}V_{|P}) = V_{1|Q}$ for any $P \in \pi^{-1}(Q)$.

Assume that $V_{1|Q} \neq 0$. By using the properness of π , there exist $\epsilon > 0$ and C^{∞} -map $F : (-\epsilon, \epsilon) \times \pi^{-1}(Q) \longrightarrow \mathbb{P}_E$ such that F(t, P) is an integral curve $\gamma(t)$ of $\sqrt{-1}V$ such that $\gamma(0) = P$. For each $t \in (-\epsilon, \epsilon)$, $F(t, \bullet)$ gives a diffeomorphism of $\pi^{-1}(Q) \simeq \pi^{-1}(\nu(t, Q))$. Then, it is easy to prove that an integral curve of $\sqrt{-1}V$ through any point of \mathbb{P}_E can be defined over \mathbb{R} . Hence, we have an \mathbb{R} -action $\nu_{\mathbb{P}_E}$ on \mathbb{P}_E whose fundamental vector field is $\sqrt{-1}V$. Because it preserves $\mathbb{P}_E \setminus E$, we obtain an \mathbb{R} -action ν_E on E. It commutes with ρ_{E1} , and satisfies $\phi \circ \nu_E(t, P) = \nu(t, \pi_1(P))$, where $\pi_1 : E \longrightarrow Y$. Hence, we obtain a holomorphic \mathbb{C}^* -action ρ_E such that π_1 is equivariant with respect to ρ_E and ρ . Let $\alpha_1, \alpha_2 \in \mathbb{C}$ and $\nu_1, \nu_2 \in E_{|Q}$ for $Q \in Y$. We have $\rho_E(t, \alpha_1 \nu_1 + \alpha_2 \nu_2) = \alpha_1 \rho_E(t, \nu_1) + \alpha_2 \rho_E(t, \nu_2)$ for $t \in S^1$. Hence, it holds for $t \in \mathbb{C}^*$, i.e., (E, ρ_E) is a \mathbb{C}^* -equivariant bundle over (Y, ρ) .

6 $\frac{\infty}{2}$ -VHS associated to Toda like harmonic bundle

6.1 Explicit description of the associated meromorphic flat bundle

6.1.1 Refinement for meromorphic prolongment

We set $X := \mathbb{P}^1_q$, $D_0 := \{0\}$, $D_\infty := \{\infty\}$ and $D := D_0 \cup D_\infty$. We set $\mathfrak{v} := \sqrt{-1}r(q\partial_q - \overline{q}\partial_{\overline{q}})$, which is the fundamental vector field of the S^1 -action on \mathbb{P}^1 given by $(t,q) \longmapsto t^r q$. We set $\mathcal{X} := \mathbb{C}_\lambda \times X$ and $\mathfrak{X} := \mathbb{P}^1 \times X$. We use the notation \mathcal{D} , \mathfrak{D} , etc., in similar meanings. We set $\mathfrak{X}^\lambda := \{\lambda\} \times X \subset \mathfrak{X}$ for $\lambda \in \mathbb{P}^1$.

Because $t^*\theta = t^m\theta$ and $t^*h = h$, the harmonic bundle $(E, \theta, h) := (E_r, \theta_{r,m}, h_a)$ on $X \setminus D$ is homogeneous of degree m with respect to \mathfrak{v} . As explained in §5.2 and §5.4, we have the locally free $\mathcal{O}_{\mathcal{X}}(*\mathcal{D})$ -module $\mathcal{Q}\mathcal{E}$ with the meromorphic flat connection $\widetilde{\nabla}$. We have the $\mathcal{O}_{\mathcal{X}}(*\mathcal{D}_{\infty})$ -submodule $\mathcal{Q}_0\mathcal{E} \subset \mathcal{Q}\mathcal{E}$. The connection $\widetilde{\nabla}$ is logarithmic along \mathcal{D}_0 .

Lemma 6.1 $Q\mathcal{E}$ is uniquely extended to a locally free $\mathcal{O}_{\mathfrak{X}}(*(\mathfrak{D} \cup \mathfrak{X}^{\infty}))$ -module $\widetilde{Q}\mathcal{E}$ such that $\widetilde{\nabla}$ gives a meromorphic flat connection of $\widetilde{Q}\mathcal{E}$ which is regular singular at any point of $\mathfrak{X}^{\infty} \setminus \mathfrak{D}_{\infty}$.

Proof Note that the harmonic bundle is not tame at ∞ . We have only to consider the issue locally around $(\lambda, q) = (\infty, \infty)$. We use the Riemann-Hilbert correspondence between meromorphic flat bundles and local systems with Stokes structure, studied in §4 of [36]. See [35] for a review.

By using the descent and pull back, we may assume m = r. Then, the eigenvalues of $\theta_{r,r}$ are of the form αdq ($\alpha \in \mathbb{C}$). Hence, the irregular values of $(\mathcal{QE}, \widetilde{\nabla})$ along $q = \infty$ are of the form $\alpha \lambda^{-1} q$ ($\alpha \in \mathbb{C}$).

It is clear that \mathcal{QE} is extended along $\mathfrak{X}^{\infty}\setminus\{(\infty,\infty)\}$ in a regular singular way. We take a blow up $\varphi:Z\longrightarrow\mathfrak{X}$ at (∞,∞) . Then, the pole and the zero of the function $\varphi^{-1}(\lambda^{-1}q)$ are separated. The pole of $\varphi^{-1}(\lambda^{-1}q)$ is the strict transform of $\mathbb{P}^1_{\lambda}\times\{\infty\}$.

We may restrict our interest to a neighbourhood of $\varphi^{-1}(\infty,\infty)$. Let D_1' be the exceptional divisor. Let $\pi: \widetilde{Z} \longrightarrow Z$ be the real blow up along $\varphi^{-1}(\mathbb{P}^1_{\lambda} \times \{\infty\}) \cup \varphi^{-1}(\mathfrak{X}^{\infty})$. Let \mathcal{L} be the local system corresponding to $(\mathcal{QE},\widetilde{\nabla})$. It is equipped with the Stokes structure outside $\pi^{-1}(D_1')$. (See [35] for a review of the Stokes structure. We shall use the notation there.)

Let D_2' be the strict transform of $\mathbb{P}^1_{\lambda} \times \{\infty\}$. Let P be the intersection of D_1' and D_2' . Let $Q \in \pi^{-1}(P)$. For a local coordinate (u, v) around P such that q = uv and $\mu = v$, the irregular values are of the form αu^{-1} ($\alpha \in \mathbb{C}$). We can take a point $Q' \in \pi^{-1}(D_2' \setminus P)$ such that (i) Q' is sufficiently close to Q, (ii) the orders $\leq_{Q'}$ and \leq_{Q} are the same on the set of the irregular values of $(Q\mathcal{E}, \widetilde{\nabla})$. We have a natural isomorphism $\mathcal{L}_Q \simeq \mathcal{L}_{Q'}$. Hence, we obtain a filtration of \mathcal{L}_Q induced by the Stokes filtration of $\mathcal{L}_{Q'}$. Thus, we obtain the Stokes structure of \mathcal{L} . By using the Riemann-Hilbert correspondence, we obtain a good meromorphic flat bundle on Z. Its push-forward to \mathfrak{X} gives the desired extension of $(Q\mathcal{E}, \widetilde{\nabla})$.

Remark 6.2 The first claim of the previous lemma holds in a more general situation, which will be argued elsewhere.

We have a locally free $\mathcal{O}_{\mathcal{X}}(*\mathcal{D}_{\infty})$ -module $\mathcal{Q}_0\mathcal{E}\subset\mathcal{Q}\mathcal{E}$.

Lemma 6.3 It is extended to a locally free $\mathcal{O}_{\mathfrak{X}}(*(\mathfrak{D}_{\infty} \cup \mathfrak{X}^{\infty}))$ -module $\widetilde{\mathcal{Q}}_0\mathcal{E}$.

Proof We have only to check the claim around $(\lambda, q) = (\infty, 0)$. We have the Deligne lattice M of $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})$ around $(\infty, 0)$. Then, it is easy to see that $\widetilde{\mathcal{Q}}_0\mathcal{E} = M(*\mathfrak{X}^\infty)$ around $(\infty, 0)$.

Let $\mathfrak{v}_m := m^{-1}\mathfrak{v}$. The harmonic bundle (E, θ, h) is homogeneous of degree 1 with respect to \mathfrak{v}_m . Even if we replace \mathfrak{v} with \mathfrak{v}_m , we clearly obtain the same meromorphic flat connection $\widetilde{\nabla}$.

For a positive integer ℓ , let $\varphi_{\ell}: \mathbb{P}^1 \longrightarrow \mathbb{P}^1$ be given by $\varphi(q) = q^{\ell}$. It gives $\varphi_{\ell*}(\mathfrak{v}_{\ell m}) = \mathfrak{v}_m$. The harmonic bundle $\varphi_{\ell}^{-1}(E,\theta,h)$ is homogeneous of degree ℓm with respect to \mathfrak{v} . Then, it is easy to observe that the meromorphic flat bundle associated to $\varphi_{\ell}^{-1}(E,\theta,h)$ is naturally isomorphic to $\varphi_{\ell}^*(\mathcal{QE},\widetilde{\nabla})$, where the induced map $\mathcal{X} \longrightarrow \mathcal{X}$, $(\lambda,q) \longmapsto (\lambda,q^{\ell})$ is also denoted by φ_{ℓ} .

6.1.2 Refinement for tr-TLE structure

Proposition 6.4 We have a locally free $\mathcal{O}_{\mathfrak{X}}(*\mathfrak{D}_{\infty})$ -submodule $\mathfrak{Q}_0\mathcal{E}\subset\widetilde{\mathcal{Q}}\mathcal{E}$ such that $(\mathfrak{Q}_0\mathcal{E},\widetilde{\nabla})$ is tr-TLE structure in the sense of Hertling [17], i.e., (i) $\widetilde{\nabla}$ is logarithmic with respect to $\mathfrak{Q}_0\mathcal{E}$ along \mathfrak{X}^{∞} , (ii) $\mathfrak{Q}_0\mathcal{E}_{|\mathbb{P}^1\times\{Q\}}$ is a pure twistor structure of weight 0 for each $Q\in X\setminus D_{\infty}$.

Proof We have already studied the issue locally around q = 0 in Proposition 5.13, by using the Deligne lattice. For the construction of $\mathfrak{Q}_0\mathcal{E}$ in this proposition, we have only to replace the Deligne lattice with $M \otimes \mathcal{O}_{\mathfrak{X}}(*\mathfrak{D}_{\infty})$, where M is the Deligne-Malgrange lattice of $\widetilde{\mathcal{Q}}\mathcal{E}$. (See [28] for Deligne-Malgrange lattice, or canonical lattice. See [34] for a review.) As explained in Proposition 5.14 and Corollary 5.16, it is a tr-TLE structure.

We set $\mathfrak{Q}\mathcal{E} := \mathfrak{Q}\mathcal{E} \otimes \mathcal{O}_{\mathfrak{X}}(*\mathfrak{D}_0)$. Note that we have a natural isomorphism $\mathfrak{Q}\mathcal{E}_{|\mathfrak{X}^0} \simeq \mathcal{V}_r$ by construction. We shall identify them. Let $p:\mathfrak{X} \longrightarrow X$ denote the projection.

Proposition 6.5 We have a unique holomorphic isomorphism $\Phi: p^*\mathcal{V}_r \simeq \mathfrak{Q}\mathcal{E}$ such that $\Phi_{|\lambda=0}$ is the identity.

Proof We have a natural isomorphism $\mathfrak{QE} \simeq p^*p_*\mathfrak{QE}$. Because $\mathfrak{QE}_{|\{0\}\times X} = \mathcal{V}_r$, we have a natural morphism $p_*\mathfrak{QE} \longrightarrow \mathcal{V}_r$. Because its restriction to X^* is an isomorphism, it is an isomorphism on X. Hence, we obtain a natural isomorphism $p^*\mathcal{V}_r \simeq \mathfrak{QE}$.

Under the isomorphism Φ , we have $\widetilde{\nabla} = p^{-1}\nabla_1 + \lambda^{-1}\theta + \widetilde{\nabla}_{\lambda}d\lambda$ for some connection ∇_1 of \mathcal{V}_r . Recall the decomposition $\mathcal{V}_r = \bigoplus_{i=1}^r \mathcal{V}^{(i)}$, where $\mathcal{V}^{(i)} = \mathcal{O}_X(*D)e_i$.

Proposition 6.6 ∇_1 is isomorphic to $\nabla_0 - \bigoplus a_i \operatorname{id}_{\mathcal{V}^{(i)}} dq/q$, where $\nabla_0 e_i = 0$ for $i = 1, \ldots, r$.

Proof We have only to consider the case m=1 by the remark given in the last of §6.1.1. Let σ_0 be an automorphism of \mathfrak{X} given by $\sigma_0(\lambda,q)=(\tau\lambda,q)$, where τ is the primitive r-the root. We have the \mathbb{C}^* -action on \mathcal{X} given by $t(\lambda,q)=(t\lambda,t^rq)$. The automorphism σ of \mathcal{V}_r induces an automorphism σ of $p^*\mathcal{V}_r$ over σ_0 . The \mathbb{C}^* -action on \mathcal{V}_r induces the action on $p^*\mathcal{V}_r$. Then, $\lambda^{-1}\theta$ and $\widetilde{\nabla}$ are σ -equivariant and \mathbb{C}^* -equivariant. Hence, ∇_1 is also σ -equivariant and \mathbb{C}^* -equivariant. Let $\mathcal{V}_{r,i}:=\mathcal{O}_{\mathbb{P}^1}(*D)e_i$. Then, ∇_1 preserves the decomposition $\mathcal{V}_r=\bigoplus \mathcal{V}_{r,i}$, and it is expressed as $\nabla_0-\bigoplus b_i\operatorname{id}_{\mathcal{V}_{r,i}}dq/q$, where $b_i\in\mathbb{C}$. The numbers b_i must be equal to the parabolic weights. (Recall the dependence of the parabolic weights and the eigenvalues of the residues $\operatorname{Res}(\nabla^\lambda)$ on λ in [18], [31], [44].)

6.1.3 Explicit description

Let us give an explicit description of $\widetilde{\nabla}$ under the isomorphism $\mathfrak{QE} \simeq p^* \mathcal{V}_r$ in Proposition 6.5. For given complex numbers $\gamma_1, \ldots, \gamma_r$, let $\operatorname{diag}[\gamma_1, \ldots, \gamma_r]$ denote the diagonal matrix whose (j, j)-entries are γ_j . By the construction of \mathbb{C}^* -action and $\widetilde{\nabla}$, we have $\widetilde{\nabla}_{m\lambda\partial_{\lambda}+rq\partial_q}\mathbf{e} = \mathbf{e} \operatorname{diag}[m, 2m, \ldots, rm]$. According to Proposition 6.6, we have

$$q\widetilde{\nabla}_{q}e = e\left(-\operatorname{diag}[a_{1},\ldots,a_{r}] + \frac{m}{\lambda}\mathcal{K}(r,m)\right).$$
 (40)

Hence, we obtain

$$\lambda \widetilde{\nabla}_{\lambda} \boldsymbol{e} = \boldsymbol{e} \Big(\operatorname{diag}[1, 2, \dots, r] + \frac{r}{m} \operatorname{diag}[a_1, \dots, a_r] - \frac{r}{\lambda} \mathcal{K}(r, m) \Big). \tag{41}$$

The formulas (40) and (41) describe $\widetilde{\nabla}$.

In particular, if m = r, we have the following:

$$q\widetilde{\nabla}_q e = e\left(-\operatorname{diag}[a_1,\ldots,a_r] + \frac{r}{\lambda}\mathcal{K}(r,r)\right)$$
 (42)

$$\lambda \widetilde{\nabla}_{\lambda} e = e \left(\operatorname{diag}[1, 2, \dots, r] + \operatorname{diag}[a_1, \dots, a_r] - \frac{r}{\lambda} \mathcal{K}(r, r) \right)$$
(43)

Let C_r be the matrix whose (i, j)-th entry is 1 if $i \equiv j + 1$ modulo r, or 0 otherwise. Then, by specializing (43) to q = 1, we obtain

$$\lambda \widetilde{\nabla}_{\lambda} \boldsymbol{e}_{|q=1} = \boldsymbol{e}_{|q=1} \left(\operatorname{diag}[1, 2, \dots, r] + \operatorname{diag}[a_1, \dots, a_r] - \frac{r}{\lambda} C_r \right).$$

By specializing (42) to $\lambda = 1$, we obtain

$$q\widetilde{\nabla}_q e_{|\lambda=1} = e_{|\lambda=1} \Big(-\operatorname{diag}[a_1, \dots, a_r] + r\mathcal{K}(r, r) \Big).$$

We set $v_i := q^{-i}e_i$. Then, we obtain

$$q\widetilde{\nabla}_q \boldsymbol{v} = \boldsymbol{v} \Big(-\operatorname{diag}[1,\ldots,r] - \operatorname{diag}[a_1,\ldots,a_r] + rC_r q \Big)$$

If we set $\zeta = q^{-1}$, we obtain the following:

$$\zeta \widetilde{\nabla}_{\zeta} \boldsymbol{v} = \boldsymbol{v} \Big(\operatorname{diag}[1, \dots, r] + \operatorname{diag}[a_1, \dots, a_r] - \frac{r}{\zeta} C_r \Big)$$

In particular, we obtain the following:

Corollary 6.7 We have an isomorphism
$$(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$$
 and $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathbb{P}^1_{\lambda}\times\{1\}}$.

Remark 6.8 We can also obtain the isomorphism in Corollary 6.7 by using the homogeneity without explicit computations.

6.2 Integral structure of Toda-like harmonic bundle

6.2.1 Special case

Let $(E_r, \theta_{r,1}, h)$ be the Toda-like harmonic bundle, corresponding to $\mathbf{a} = (a_i) \in \mathfrak{R}_{r,1}(a)$ as in Theorem 3.32. We study when $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathbb{P}^1_{\lambda} \times \{1\}}$ and $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathfrak{X}^1}$ have \mathbb{Z} -structures.

Theorem 6.9 $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure (resp. \mathbb{Q} -structure) if and only if $P_{\boldsymbol{a}}(T) \in \mathbb{Z}[T]$ (resp. \mathbb{Q} -structure). (See Theorem 6.15 below for general m.)

Proof It follows from Theorem 4.14, Corollary 4.17 and the formula (40).

If $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure, then we have $\boldsymbol{a}\in\mathbb{Q}^r$. Indeed, because we have $\prod_{i=1}^r(T-e^{2\pi\sqrt{-1}a_i})\in\mathbb{Z}[T]$, the number $e^{2\pi\sqrt{-1}a_i}$ and its conjugates over \mathbb{Q} are algebraic integers such that the absolute values are 1. It is a classical theorem of Kronecker that $e^{2\pi\sqrt{-1}a_i}$ are roots of unity, i.e., $a_i\in\mathbb{Q}$. Hence, when we are interested in \mathbb{Z} -structure, it is natural to impose the condition $\boldsymbol{a}\in\mathbb{Q}^r$. In that case, we have $P_{\boldsymbol{a}}(T)\in\mathbb{Z}[T]$ if and only if $P_{\boldsymbol{a}}(T)\in\mathbb{Q}[T]$, and therefore $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure if and only if it has \mathbb{Q} -structure.

Remark 6.10 The condition $P_a(T) \in \mathbb{Z}[T]$ implies an additional symmetry for the sequence (a_1, \ldots, a_n) . Hence, the harmonic bundle is equipped with a symmetric pairing. (See §3.1.6.) It induces a real structure by the procedure in §5.3. By Proposition 4.12, we can adjust that a \mathbb{Z} -structure is compatible with the \mathbb{R} -structure.

Remark 6.11 If the condition in Theorem 6.9 is satisfied, the meromorphic flat bundle $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})$ has a \mathbb{Z} -structure, although we do not discuss a \mathbb{Z} -structure of a meromorphic flat bundle on higher dimensional varieties in this paper. In particular, $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathbb{P}^1_{\mathbb{T}}\times\{1\}}$ has a \mathbb{Z} -structure, which also follows from Theorem 6.12 below.

6.2.2 Criterion

We shall freely use the notation in §3.1.5. Let h be the Toda-like harmonic metric of $(E, \theta) := (E_r, \theta_{r,r})$ corresponding to $\mathbf{a} \in \mathfrak{R}_{r,r} \cap \mathbb{Q}^r$. We have the associated meromorphic flat bundle $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})$ on \mathfrak{X} . Note that $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure (resp. \mathbb{Q} -structure) if and only if $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure (resp. \mathbb{Q} -

structure) by Corollary 6.7. We set $\delta_d = (1, \ldots, 1)$ for a positive integer d.

Theorem 6.12 The following conditions are equivalent.

- (P1) $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathbb{P}^1_{\lambda} \times \{1\}}$ has a \mathbb{Z} -structure.
- (P2) $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathbb{P}^1_{\lambda} \times \{1\}}$ has a \mathbb{Q} -structure.
- **(P3)** There exists $\gamma \in \mathbb{C}^*$ such that (i) $\gamma^r \in \mathbb{Q}$, (ii) $\prod_{i=1}^r (T \gamma e^{2\pi \sqrt{-1}a_i/r}) \in \mathbb{Q}[T]$.

Proof We shall use the description of $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})$ given in §6.1.3. We have only to study the existence of \mathbb{Q} -structure of $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$.

Let us consider the case $\mathcal{S}(\boldsymbol{a},r) = \{r\}$. By Proposition 3.18 and the Kobayashi-Hitchin correspondence, the filtered flat bundle $(\mathcal{Q}_*\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ is stable. Because the eigenvalues of the residue of θ are 0 at both $\{0\}$ and $\{\infty\}$, we obtain that $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ is irreducible as a meromorphic flat bundle. (See [40]. See also Lemma 19.4.3 of [36]. Note that $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ is deformed as in §11 [36].) Then, the equivalence of the three conditions follow from Theorem 4.22.

Let us consider the case $\mathcal{S}(\boldsymbol{a},r) \neq \{r\}$. We have the decomposition $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla}) = \bigoplus_s (\widetilde{\mathcal{Q}}\mathcal{E}^{(s)},\widetilde{\nabla}^{(s)})$ induced by (9). As remarked in the argument for the case $\mathcal{S}(\boldsymbol{a},r) = \{r\}$, each $(\widetilde{\mathcal{Q}}\mathcal{E}^{(s)},\widetilde{\nabla}^{(s)})_{|\mathfrak{X}^1}$ is irreducible. Because there is no common eigenvalue of $\theta^{(s)}$ and $\theta^{(s')}$ ($s \neq s'$), we have $\operatorname{Irr}(\widetilde{\nabla}^{(s)}) \cap \operatorname{Irr}(\widetilde{\nabla}^{(s')}) = \emptyset$ if $s \neq s'$. Let us show the equivalence of the conditions (**P1**) and (**P2**). Clearly (**P1**) implies (**P2**). Suppose (**P2**). By applying

Proposition 4.11 to the irreducible decomposition $(\widetilde{\mathcal{Q}}\mathcal{E}, \widetilde{\nabla})_{|\mathfrak{X}^1} = \bigoplus_s (\widetilde{\mathcal{Q}}\mathcal{E}^{(s)}, \widetilde{\nabla}^{(s)})_{|\mathfrak{X}^1}$, each $(\widetilde{\mathcal{Q}}\mathcal{E}^{(s)}, \widetilde{\nabla}^{(s)})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure. Hence, **(P1)** holds. We consider the following condition:

(P3') There exists $\gamma \in \mathbb{C}^*$ such that (i) $\gamma^{r_0} \in \mathbb{Q}$, (ii) $\prod_{i=1}^{r_0} (T - \gamma e^{2\pi\sqrt{-1}a_i/r_0}) \in \mathbb{Q}[T]$.

If (P3') holds, by applying the result in the case $\mathcal{S}(\boldsymbol{a},r) = \{r\}$ to each $(\widetilde{\mathcal{Q}}\mathcal{E}^{(s)},\widetilde{\nabla}^{(s)})_{|\mathfrak{X}^1}$, we obtain that (P2) holds. Conversely, if (P2) holds, by Proposition 4.11, each $(\widetilde{\mathcal{Q}}\mathcal{E}^{(s)},\widetilde{\nabla}^{(s)})_{|\mathfrak{X}^1}$ has a \mathbb{Q} -structure. By the result in the case $\mathcal{S}(\boldsymbol{a},r) = \{r\}$, we obtain that (P3') holds. Finally, the claim of the theorem follows from the equivalence of the conditions (P3) and (P3'). (See Lemma 6.16 below.)

For example, we obtain the following from Theorem 6.12 and Lemma 4.25 (or Proposition 4.26).

Corollary 6.13 Suppose that r is an odd prime. Then, $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathbb{P}^1_{\lambda}\times\{1\}}$ has a \mathbb{Z} -structure, if and only if there exists a half integer ℓ such that $\prod_{i=1}^r (T - e^{2\pi\sqrt{-1}(a_i+\ell)/r}) \in \mathbb{Z}[T]$.

Remark 6.14 According to the correspondence between Toda-like harmonic bundles and solutions of Toda lattice with opposite sign (see $\S 3.3$), we obtain a criterion when the meromorphic flat bundle associated to a solution of the Toda equation has a \mathbb{Z} -structure.

Variant Let h be the Toda-like harmonic metric of $(E_r, \theta_{r,m})$ corresponding to $\mathbf{a} \in \mathfrak{R}_{r,m} \cap \mathbb{Q}^r$. The following proposition can be proved by the argument in the proof of Theorem 6.12.

Theorem 6.15 The associated meromorphic flat bundle $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Q} -structure, if and only if there exists $\gamma \in \mathbb{C}^*$ such that (i) $\gamma^m \in \mathbb{Q}$, (ii) $\prod_{i=1}^r (T - \gamma e^{2\pi\sqrt{-1}a_i/m}) \in \mathbb{Q}[T]$. In that case, $(\widetilde{\mathcal{Q}}\mathcal{E},\widetilde{\nabla})_{|\mathfrak{X}^1}$ has a \mathbb{Z} -structure, more strongly.

Appendix Let r_0, j_0 be positive integers, and set $r_1 = r_0 j_0$. Let $\mu_p := \{ \kappa \in \mathbb{C} \mid \kappa^p = 1 \}$ for a positive integer p. We have the homomorphism $\Psi : \mu_{r_1} \longrightarrow \mu_{r_0}$ given by $\Psi(\kappa) = \kappa^{j_0}$. Let S_0 be a subset μ_{r_0} with a function $f_0 : S_0 \longrightarrow \mathbb{Z}_{>0}$. We put $S_1 := \Psi^{-1}(S_0)$ and $f_1 := f_0 \circ \Psi$. We take $\gamma_1 \in \mathbb{C}^*$ and put $\gamma_0 := \gamma_1^{j_0}$. For i = 0, 1, we set

$$P_i(T) := \prod_{b \in S_i} (T - \gamma_i b)^{f_i(b)}.$$

Because $P_1(T) = P_0(T^{j_0})$, the following lemma is clear.

Lemma 6.16 We have $P_0(T) \in \mathbb{Q}[T]$ if and only if $P_1(T) \in \mathbb{Q}[T]$.

References

- [1] E. Aldrovandi and G. Falqui, Geometry of Higgs and Toda fields on Riemann surfaces, J. Geom. Phys. 17 (1995), 25–48.
- [2] D. Baraglia, Cyclic Higgs bundles and the affine Toda equations, arXiv:1011.6421
- [3] S. Barannikov, Quantum periods. I. Semi-infinite variations of Hodge structures, Internat. Math. Res. Notices, 23, (2001), 1243–1264.
- [4] O. Biquard, Fibrés de Higgs et connexions intégrables: le cas logarithmique (diviseur lisse), Ann. Sci. École Norm. Sup. **30** (1997), 41–96.
- [5] O. Biquard and P. Boalch, Wild non-abelian Hodge theory on curves, Compos. Math. 140 (2004), 179–204.
- [6] S. Cecotti and C. Vafa, Topological-anti-topological fusion, Nuclear Phys. B, 367 (1991), 359-461.

- [7] S. Cecotti and C. Vafa, On classification of N=2 supersymmetric theories, Comm. Math. Phys. 158 (1993), 569-644
- [8] J. Dorfmeister, M. Guest, W. Rossman, The tt* structure of the quantum cohomology of CP1 from the viewpoint of differential geometry, Asian J. Math. 14 (2010), 417–437.
- [9] A. Douai, Quantum differential systems and some applications to mirror symmetry, arXiv:1203.5920
- [10] A. Douai, C. Sabbah, Gauss-Manin systems, Brieskorn lattices and Frobenius structures. I. Ann. Inst. Fourier (Grenoble) 53 (2003), 1055–1116
- [11] B. Dubrovin, Geometry and integrability of topological-antitopological fusion, Comm. Math. Phys., 152 (1993), 539–564.
- [12] G. Fujisaki, Fields and Galois theory, (in Japanese) Iwanami, Tokyo, 1983
- [13] M. A. Guest and C.-S. Lin, Nonlinear PDE aspects of the tt* equations of Cecotti and Vafa, arXiv:1010.1889.
- [14] M. A. Guest, C.-S. Lin, Some tt* structures and their integral Stokes data, arXiv:1209.2318
- [15] M. A. Guest, A. R. Its, C.-S. Lin. Isomonodromy aspects of the tt* equations of Cecotti and Vafa I. Stokes data, arXiv:1209.2045
- [16] D. Guzzetti, Stokes matrices and monodromy of the quantum cohomology of projective spaces, Comm. Math. Phys. **207** (1999), 341–383.
- [17] C. Hertling, tt* geometry, Frobenius manifolds, their connections, and the construction for singularities, J. Reine Angew. Math., **555** (2003), 77–161.
- [18] C. Hertling and C. Sevenheck, Nilpotent orbits of a generalization of Hodge structures, J. Reine Angew. Math., 609 (2007), 23–80
- [19] C. Hertling and C. Sevenheck *Limits of families of Brieskorn lattices and compactified classifying spaces*, Adv. Math. **223** (2010), 1155–1224.
- [20] N. Hitchin, The self-duality equations on a Riemann surface, Proc. London Math. Soc. (3) **55** (1987), 59–126.
- [21] N. Hitchin, *Monopoles, minimal surfaces and algebraic curves*. Presses de l'Université de Montréal, Montreal, QC, 1987.
- [22] H. Iritani, An integral structure in quantum cohomology and mirror symmetry for toric orbifolds, Adv. Math. 222 (2009), 1016–1079.
- [23] H. Iritani, tt*-geometry in quantum cohomology, arXiv:0906.1307
- [24] A. R. Its and V. Y. Novokshenov, The isomonodromic deformation method in the theory of Painlevé equations, Lecture Notes in Mathematics, 1191, (Springer, Berlin 1986)
- [25] B. M. McCoy, C. A. Tracy, and T. T. Wu, Painlevé functions of the third kind, J. Mathematical Phys. 18 (1977), 1058–1092.
- [26] L. Katzarkov, M. Kontsevich, and T. Pantev, Hodge theoretic aspects of mirror symmetry, IN From Hodge Theory to Integrability and TQFT: tt*-geometry eds. R. Y. Donagi and K. Wendland, Proc. of Symp. Pure Math. 78, Amer. Math. Soc. (2007), 87–174
- [27] A. Levelt, Jordan decomposition for a class of singular differential operators, Ark. Math. 13, (1975), 1–27
- [28] B. Malgrange, Connexions méromorphies 2, Le réseau canonique, Invent. Math. 124, (1996) 367–387.

- [29] E. Mann, Orbifold quantum cohomology of weighted projective spaces, J. Algebraic Geom. 17 (2008), 137–166.
- [30] T. Mochizuki, Kobayashi-Hitchin correspondence for tame harmonic bundles and an application Astérisque 309 (2006)
- [31] T. Mochizuki, Asymptotic behaviour of tame harmonic bundles and an application to pure twistor D-modules I, II, Mem. AMS. 185 (2007)
- [32] T. Mochizuki, Kobayashi-Hitchin correspondence for tame harmonic bundles II, Geometry&Topology 13, (2009), 359–455.
- [33] T. Mochizuki, Asymptotic behaviour of variation of pure polarized TERP structures, Publ. Res. Inst. Math. Sci. 47 (2011), 419–534.
- [34] T. Mochizuki, On Deligne-Malgrange lattices, resolution of turning points and harmonic bundles, Ann. Inst. Fourier (Grenoble) 59, (2009), 2819–2837
- [35] T. Mochizuki, The Stokes structure of a good meromorphic flat bundle, J. Inst. Math. Jussieu 10 (2011), 675–712.
- [36] T. Mochizuki, Wild harmonic bundles and wild pure twistor D-modules, Astérisque 340, 2011
- [37] C. Peters, and J. Steenbrink, Mixed Hodge structures, Springer-Verlag, Berlin, 2008.
- [38] T. Reichelt, A construction of Frobenius manifolds with logarithmic poles and applications, Comm. Math. Phys. 287 (2009), 1145–1187.
- [39] T. Reichelt and C. Sevenheck. Logarithmic Frobenius manifolds, hypergeometric systems and quantum D-modules, arXiv:1010.2118
- [40] C. Sabbah, Harmonic metrics and connections with irregular singularities, Ann. Inst. Fourier (Grenoble) 49 (1999), 1265–1291
- [41] C. Sabbah, Polarizable twistor D-modules Astérisque, 300, Société Mathématique de France, Paris, (2005).
- [42] M. Saito, On the structure of Brieskorn lattice, Ann. Inst. Fourier (Grenoble) 39 (1989), 27–72.
- [43] C. T. Simpson, Constructing variations of Hodge structure using Yang-Mills theory and applications to uniformization, J. Amer. Math. Soc. 1 (1988), 867–918.
- [44] C. T. Simpson, Harmonic bundles on noncompact curves, J. Amer. Math. Soc. 3 (1990), 713–770.
- [45] C. T. Simpson, Mixed twistor structures, math.AG/9705006.
- [46] C. T. Simpson, Katz's middle convolution algorithm, Pure Appl. Math. Q. 5 (2009), 781–852.
- [47] C. A. Tracy and H. Widom, Asymptotics of a class of solutions to the cylindrical Toda equations, Comm. Math. Phys. 190 (1998), 697–721.
- [48] K. Takasaki, The world of Integrable Systems (in Japanese), (Kyoritsu Shuppan, Tokyo) 2001
- [49] W. Wasow, Asymptotic expansions for ordinary equations, Reprint of 1976 edition. Dover Publications, Inc., New York, (1987)
- [50] H. Widom, Some classes of solutions to the Toda lattice hierarchy, Comm. Math. Phys. 184 (1997), 653–667.

Address

Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606-8502, Japan, takuro@kurims.kyoto-u.ac.jp