

Lévy path integral approach to the solution of the fractional Schrödinger equation with infinite square well

Jianping Dong¹

Department of Mathematics, College of Science, Nanjing University of Aeronautics and Astronautics, Nanjing 210016, China

The solution to the fractional Schrödinger equation with infinite square well is obtained in this paper, by use of the Lévy path integral approach. We obtain the even and odd parity wave functions of this problem, which are in accordance with those given by Laskin in [Chaos 10 (2000), 780–790].

1 Introduction

The phrase “fractional quantum mechanics” was first introduced by Nick Laskin [1, 2]. The fractional quantum system is described by the fractional Schrödinger equation (FSE) [3] containing the fractional Riesz operator [4]. It is well-known that Feynman and Hibbs [5] reformulated the standard Schrödinger equation [6] by use of the path integral approach considering the Gaussian probability distribution. The Lévy stochastic process is a natural generalization of the Gaussian process. The possibility of developing the path integral over the paths of the Lévy motion was discussed by Kac [7], who pointed out that the Lévy path integral generates the functional measure in the space of left (or right) continued functions having only discontinuities of the first kind. Recently, Laskin [1] generalized Feynman path integral to Lévy one, and developed the space-fractional Schrödinger equation containing the Riesz fractional derivative [3, 4]. Then, he constructed the fractional quantum mechanics and showed some properties of the space fractional quantum system [2, 8, 9]. Afterwards, the time-fractional Schrödinger equation containing the Caputo fractional derivative [4], and the space-time-fractional Schrödinger equation [10–12] were also given. Because of the double complexities of the fractional calculus and quantum mechanics, not too much progress has been made in this field. The solutions of the FSE with the linear potential, the delta potential, the infinite square well, and so on, have been obtained [8, 9, 13–16]. The fractional scattering problem for the one- and three-dimensional cases were also studied [17, 18], and the Thomas-Fermi theory for the multi-particle system was also generalized to the fractional quantum system [19].

This paper focuses on the solution to the fractional Schrödinger equation with infinite square well. This solution was firstly obtained by Laskin in the piece-wise solution approach, but then in Ref. [20], Jeng, et al. pointed out that this solution and some other obtained solutions to the fractional Schrödinger equation with specific potentials were incorrect, because of the nonlocal property of the fractional Riesz operator. Then, in Ref. [21], Bayin argued that the proof given by Jeng, et al., that the solution gotten by Laskin did not satisfy the FSE, was not true, and proved Laskin’s solution to be correct. But the controversy on this point still do not stop yet, see, for example, Ref. [22]. In this paper, we attempt to get the solution of the FSE with infinite square well in an indirect way: the Lévy path integral approach.

2 Lévy path integral and Fractional Schrödinger Equation

The Lévy path integral is generalized from the Feynman one by Laskin in Refs. [1, 9]. In this section, we give the introduction to the Lévy path integral and fractional Schrödinger equation, and show some properties useful in the following sections. If a particle at an initial time t_a starts from the point x_a and goes to a final point x_b at time t_b , we will say simply that the particle goes from a and b and its trajectory (path) $x(t)$ will have the property that $x(t_a) = x_a$ and $x(t_b) = x_b$. In quantum mechanics, then, we will have an quantum-mechanical amplitude, often called a kernel, which we may write $K(x_b t_b | x_a t_a)$, to get from the point a to the point b . This will be the sum over all of the paths that go between that end points and of a contribution from each. If we have the quantum particle moving in the potential then the quantum-mechanical amplitude $K_L(x_b t_b | x_a t_a)$ can be written as

$$K_L(x_b t_b | x_a t_a) = \int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}x(\tau) \cdot \exp \left\{ -\frac{i}{\hbar} \int_{t_a}^{t_b} d\tau V(x(\tau)) \right\}, \quad (1)$$

¹Email: Dongjp.sdu@gmail.com

where $V(x(\tau))$ is the potential energy as a functional of the Lévy particle path and the fractional path integral measure is defined as

$$\int_{x(t_a)=x_a}^{x(t_b)=x_b} \mathcal{D}x(\tau) = \lim_{N \rightarrow \infty} \int dx_1 \cdots dx_{N-1} \hbar^{-N} \left(\frac{iD_\alpha \varepsilon}{\hbar} \right)^{-N/\alpha} \cdot \prod_{j=1}^N L_\alpha \left\{ \frac{1}{\hbar} \left(\frac{\hbar}{iD_\alpha \varepsilon} \right)^{1/\alpha} |x_j - x_{j-1}| \right\} \cdots, \quad (2)$$

where D_α is the generalized 'fractional diffusion coefficient', the physical dimension of which is $[D_\alpha] = \text{erg}^{1-\alpha} \times \text{cm}^\alpha \times \text{sec}^{-\alpha}$ ($D_\alpha = 1/2m$ for $\alpha = 2$, m denotes the mass of a particle), $x_0 = x_a$, $x_N = x_b$, $\varepsilon = (t_b - t_a)/N$, and the Lévy probability distribution function L_α is expressed in term of Fox's H function (see [1]). The free particle kernel of the Lévy type can be expressed in the momentum representation by

$$K_L^{(0)}(x_b t_b | x_a t_a) = \frac{1}{2\pi\hbar} \int_{-\infty}^{+\infty} dp \cdot \exp \left\{ i \frac{p(x_b - x_a)}{\hbar} - i \frac{D_\alpha (t_b - t_a) |p|^\alpha}{\hbar} \right\}. \quad (3)$$

The kernel $K_L(x_b t_b | x_a t_a)$ which is defined by Eq. (12), describes the evolution of the fractional quantum-mechanical system,

$$\psi_f(x_b, t_b) = \int_{-\infty}^{+\infty} dx_a K_L(x_b t_b | x_a t_a) \psi_i(x_b, t_b), \quad (4)$$

where $\psi_i(x_b, t_b)$ is the fractional wave function of initial (at the $t = t_a$) state and $\psi_f(x_b, t_b)$ is the fractional wave function of the final (at the $t = t_b$) state. With the help of this equation, for the fractional quantum system, Laskin derived the fractional Schrödinger equation given as follows (in one-dimensional case),

$$i\hbar \frac{\partial \psi(x, t)}{\partial t} = H_\alpha \psi(x, t), \quad (5)$$

where $\psi(x, t)$ is the time-dependent wave function, and H_α ($1 < \alpha \leq 2$) is the fractional Hamiltonian operator given by

$$H_\alpha = -D_\alpha (\hbar \nabla)^\alpha + V(x, t). \quad (6)$$

Here, $(\hbar \nabla)^\alpha$ is the quantum Riesz fractional operator defined by

$$(\hbar \nabla)^\alpha \psi(x, t) = -\frac{1}{2\pi\hbar} \int_{-\infty}^{+\infty} dp e^{ipx/\hbar} |p|^\alpha \int_{-\infty}^{+\infty} e^{-ipx'/\hbar} \psi(x', t) dx'. \quad (7)$$

In Ref., the fractional Hamiltonian operator was proved to be a Hermitian or self-adjoint operator. When H_α is not explicitly dependent on t , that is, $V(x, t) = V(x)$, after separation of variables, Eq. (5) goes over to a form of steady state,

$$-D_\alpha (\hbar \nabla)^\alpha \phi(x) + V(x) \phi(x) = E \phi(x), \quad (8)$$

where $\psi(x, t)$ and $\phi(x)$ are related to each other by $\psi(x, t) = \phi(x) e^{-iEt/\hbar}$, in which E denotes the energy of the quantum system. Suppose that the solutions $\phi_n(x)$ (time-independent wave functions) to Eq. (8) corresponding to the set of energy levels E_n are not only orthogonal but also normalized, then following Feynman's method, we can obtain the relationship between the fractional quantum-mechanical kernel and the time-independent wave functions $\phi_n(x)$,

$$K_L(x_b t_b | x_a t_a) = \sum_{n=1}^{\infty} \phi_n(x_b) \phi_n^*(x_a) e^{-(i/\hbar) E_n (t_b - t_a)}, \quad \text{for } t_b > t_a \quad (9)$$

and $K_L(x_b t_b | x_a t_a) = 0$, for $t_b < t_a$; $K_L(x_b t_b | x_a t_a) = \delta(x_b - x_a)$, for $t_b = t_a$. Eq. (9) will be used in the following section to solve the fractional Schrödinger equation with infinite square well.

3 Infinite square well: wave functions and energy eigenvalues

In this section, we consider a particle in the infinite square well with the potential $V(x)$ defined as

$$V(x) = \begin{cases} 0, & |x| \leq l, \\ \infty, & |x| > l. \end{cases} \quad (10)$$

The solution to the fractional Schrödinger equation with this potential was firstly studied by Laskin in Ref. [8], in which this equation is solved in a piecewise approach. In this paper, to avoid treating the nonlocal property of the fractional Riesz operator, we get the solutions to this equation in the Lévy path integral approach using Eq. (9). That is, we turn to get the fractional quantum-mechanical kernel of this quantum system. Now we use the path cancellation method to get the transfer kernel.

An interpretation of the infinite potential barrier problem was the observation that there are two classical paths connecting the points (x_a, t_a) and (x_b, t_b) [23]. This is because the classical particle may go directly between the two points, or it may bounce off the wall once on the way. With the infinite well, the particle may bounce back and forth an arbitrary number of times on its way from (x_a, t_a) to (x_b, t_b) . This leads to an infinite number of classical paths connecting the two points. These paths correspond to an infinite sequence of images of the final point having coordinates

$$x_r = \begin{cases} 2lr + x_b, & \text{when } r = 2m, \\ 2lr - x_b, & \text{when } r = 2m + 1, \end{cases} \quad (11)$$

in which $m = 0, \pm 1, \pm 2, \dots$. Here, the subscript r is used to denote the number of reflections in the classical path corresponding to the image point, and to distinguish from the subscript n used for the energy eigenstates.

A generalization of the above arguments for the potential barrier would suggest that the propagator $K_L(x_b t_b | x_a t_a)$ should be the sum of contributions from each of these classical paths. The contribution from each classical path should be the free-particle propagator for its corresponding image point, multiplied by -1 for each time it is reflected. This leads to the kernel,

$$K_L(x_b t_b | x_a t_a) = \sum_{r=-\infty}^{+\infty} (-1)^r K_L^{(0)}(x_r t_b | x_a t_a) = \sum_{m=-\infty}^{+\infty} [K_L^{(0)}(x_{2m} t_b | x_a t_a) - K_L^{(0)}(x_{2m+1} t_b | x_a t_a)]. \quad (12)$$

Recalling Eqs. (3) and (11), the kernel can be calculated as follows,

$$\begin{aligned} K_L(x_b t_b | x_a t_a) &= \sum_{m=-\infty}^{+\infty} \frac{1}{2\pi\hbar} \int_{-\infty}^{+\infty} \left\{ \exp \left[i \frac{p(4ml + x_b - x_a)}{\hbar} - i \frac{D_\alpha(t_b - t_a)|p|^\alpha}{\hbar} \right] - \exp \left[i \frac{p(4ml + 2l - x_b - x_a)}{\hbar} - i \frac{D_\alpha(t_b - t_a)|p|^\alpha}{\hbar} \right] \right\} dp \\ &= \frac{1}{2\pi\hbar} \int_{-\infty}^{+\infty} \exp \left[-i \frac{D_\alpha(t_b - t_a)|p|^\alpha}{\hbar} \right] \left\{ \exp \left[i \frac{p[(x_b - l) - (x_a - l)]}{\hbar} \right] - \exp \left[i \frac{p[-(x_b - l) - (x_a - l)]}{\hbar} \right] \right\} \sum_{m=-\infty}^{+\infty} \exp \left[i \frac{4mlp}{\hbar} \right] dp \\ &= \frac{i}{\pi\hbar} \int_{-\infty}^{+\infty} \exp \left[-i \frac{D_\alpha(t_b - t_a)|p|^\alpha}{\hbar} \right] \exp \left[i \frac{-p(x_a - l)}{\hbar} \right] \sin \left(\frac{p(x_a - l)}{\hbar} \right) \sum_{m=-\infty}^{+\infty} \exp \left[i \frac{4mlp}{\hbar} \right] dp. \end{aligned} \quad (13)$$

Furthermore, using the Poisson summation rule

$$\sum_{r=-\infty}^{+\infty} \exp(2\pi i m \mu) = \sum_{n=-\infty}^{+\infty} \delta(\mu - n), \quad (14)$$

Then, we have

$$\begin{aligned}
K_L(x_b t_b | x_a t_a) &= \frac{i}{\pi \hbar} \int_{-\infty}^{+\infty} \exp \left[-i \frac{D_\alpha (t_b - t_a) |p|^\alpha}{\hbar} \right] \exp \left[i \frac{-p(x_a - l)}{\hbar} \right] \sin \left(\frac{p(x_a - l)}{\hbar} \right) \sum_{n=-\infty}^{+\infty} \delta \left(\frac{2lp}{\pi \hbar} - n \right) dp \\
&= \frac{i}{2l} \sum_{n=-\infty}^{+\infty} \int_{-\infty}^{+\infty} \exp \left[-i \frac{D_\alpha (t_b - t_a) |p|^\alpha}{\hbar} \right] \exp \left[i \frac{-p(x_a - l)}{\hbar} \right] \sin \left(\frac{p(x_a - l)}{\hbar} \right) \delta \left(p - \frac{n\pi \hbar}{2l} \right) dp \\
&= \frac{i}{2l} \sum_{n=-\infty}^{+\infty} \exp [-i E_n (t_b - t_a)] \exp [-i k_n (x_a - l)] \sin [k_n (x_a - l)],
\end{aligned} \tag{15}$$

in which

$$k_n = \frac{n\pi}{2l}, E_n = D_\alpha \hbar^\alpha |k_n|^\alpha.$$

Noting that the term with $n = 0$ vanishes, we can combine the positive and negative terms to get

$$K_L(x_b t_b | x_a t_a) = \frac{1}{l} \sum_{n=1}^{\infty} \exp [-i E_n (t_b - t_a)] \sin [k_n (x_a - l)] \sin [k_n (x_a - l)]. \tag{16}$$

Comparing Eq. (16) with Eq. (9) yields the wave functions and energy levels,

$$\phi_n(x) = \frac{1}{\sqrt{l}} \sin [k_n (x - l)] = \frac{1}{\sqrt{l}} \sin \left[\frac{n\pi}{2l} (x - l) \right], E_n = D_\alpha \left(\frac{n\pi \hbar}{2l} \right)^\alpha, n = 1, 2, 3, \dots \tag{17}$$

These results seem different from those given by Laskin. In fact, we have

$$\phi_n(x) = \begin{cases} \frac{1}{\sqrt{l}} \sin \left(\frac{j\pi}{l} x - j\pi \right) = \frac{(-1)^j}{\sqrt{l}} \sin \left(\frac{j\pi}{l} x \right), & \text{when } n = 2j, j = 1, 2, 3, \dots \\ \frac{1}{\sqrt{l}} \sin \left(\frac{j\pi}{l} x - j\pi - \frac{\pi}{2} \right) = \frac{(-1)^{j+1}}{\sqrt{l}} \cos \left(\frac{j\pi}{l} x \right). & \text{when } n = 2j + 1, j = 0, 1, 2, 3, \dots \end{cases} \tag{18}$$

Omitting the product factor, we can get the even and odd parity wave functions and the corresponding energy levels,

$$\begin{aligned}
\phi_n^{\text{odd}} &= \frac{1}{\sqrt{l}} \sin \left(\frac{n\pi}{l} x \right), E_n = D_\alpha \left(\frac{n\pi \hbar}{l} \right)^\alpha; \\
\phi_n^{\text{even}} &= \frac{1}{\sqrt{l}} \cos \left(\frac{n\pi}{l} x \right), E_n = D_\alpha \left(\left(n + \frac{1}{2} \right) \frac{\pi \hbar}{l} \right)^\alpha,
\end{aligned} \tag{19}$$

which are in accordance with the results given by Laskin.

4 Conclusions

In this paper, we solved the fractional Schrödinger equation with infinite square well in the Lévy path integral approach. Because of the nonlocal property of the fractional Riesz operator, the usual method for solving the standard Schrödinger equation can not be effective here. The solution to the equation gotten by the usual method has been proved to be true, but whether the usual method is right or not, or when the usual method is effective, still need more investigations. The method given in this paper, can be a way to solve the fractional Schrödinger equation indirectly.

Acknowledgements

This work was supported by the National Natural Science Foundation of China (Grant No. 11147109), the Specialized Research Fund for the Doctoral Program of Higher Education of China (Grant No. 20113218120030), and the Fundamental Research Funds for the Central Universities (Grant No. NS2012119).

References

- [1] N. Laskin, Fractional quantum mechanics and Lévy path integrals, *Phys. Lett. A* 268 (2000) 298–305.
- [2] N. Laskin, Fractional quantum mechanics, *Phys. Rev. E* 62 (2000) 3135–3145.
- [3] N. Laskin, Fractional Schrödinger equation, *Phys. Rev. E* 66 (2002) 056108.
- [4] A. A. Kilbas, H. M. Srivastava and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, Amsterdam, 2006.
- [5] R. P. Feynman and A. R. Hibbs, *Quantum Mechanics and Path Integrals*, McGraw-Hill, New York, 1965.
- [6] F. S. Levin, *An Introduction to Quantum Theory*, Cambridge University Press, 2002.
- [7] M. Kac, On Some Connections between Probability Theory and Differential and Integral Equations, in *Second Berkeley Symposium on Mathematical Statistics and Probability* (edited by Jerzy Neyman), University of California Press, Berkeley, California, 1951.
- [8] N. Laskin, Fractals and quantum mechanics, *Chaos* 10 (2000) 780–790.
- [9] N. Laskin, Levy flights over quantum paths, *Comm. Nonlin. Sci. Num. Sim.* 12(2), 2007.
- [10] M. Naber, Time fractional Schrödinger equation, *J. Math. Phys.* 45(8), 2004. 3339-3352.
- [11] S. W. Wang and M. Y. Xu, Generalized fractional Schrödinger equation with space-time fractional derivatives, *J. Math. Phys.* 48 (2007) 043502.
- [12] J. P. Dong, M. Y. Xu, Space-time fractional Schrödinger equation with time-independent potentials, *J. Math. Anal. Appl.*, 344, 1005-1017 (2008).
- [13] J. P. Dong, M. Y. Xu, Applications of continuity and discontinuity of a fractional derivative of the wave functions to fractional quantum mechanics, *J. Math. Phys.* 49, 052105 (2008).
- [14] J. P. Dong, M. Y. Xu, Some solutions to the space fractional Schrödinger equation using momentum representation method, *J. Math. Phys.* 48 (2007) 072105.
- [15] X. Y. Guo, M. Y. Xu, Some physical applications of fractional Schrödinger equation, *J. Math. Phys.* 47(2006) 082104.
- [16] E. C. Oliveira, F. S. Costa and J. J. Vaz, The fractional Schrödinger equation for delta potentials, *J. Math. Phys.* 51, 123517 (2010).
- [17] J. P. Dong, Generalized Lippmann-Schwinger equation in the fractional quantum mechanics, *Journal of Physics A: Mathematical and Theoretical*, 44 (2011) 215204;
- [18] J. P. Dong, Greens function for the time-dependent scattering problem in the fractional quantum mechanics, *Journal of Mathematical physics*, 52 (2011) 042103 ;
- [19] J. P. Dong, Applications of density matrix in the fractional quantum mechanics: Thomas-Fermi model and Hohenberg-Kohn theorems revisited, *Physics Letters A*, 375 (2011) 2787-2792.
- [20] M. Jeng, S.-L.-Y. Xu, E. Hawkins, and J. M. Schwarz, *J. Math. Phys.* 51, 062102 (2010).
- [21] S. S. Bayin, On the consistency of the solutions of the space fractional Schrödinger equation, *J. Math. Phys.* 53, 042105 (2012).
- [22] E. Hawkins and J. M. Schwarz, Comment on: “the consistency of solutions of the space fractional Schrödinger equation”, arXiv: 1210.1447.
- [23] M. Goodman, Path integral solution to the infinite square well, *American Journal of Physics*, 49(1981), 843.