HOMOLOGY CLASSES OF NEGATIVE SQUARE AND EMBEDDED SURFACES IN 4-MANIFOLDS

M. J. D. HAMILTON

ABSTRACT. Let X be a simply-connected closed oriented 4-manifold and A an embedded surface of genus g and negative self-intersection -N. We show that for fixed genus g there is an upper bound on N if the homology class of A is divisible or characteristic. In particular, for genus zero, there is a lower bound on the self-intersection of embedded spheres in these kinds of homology classes. This question is related to a problem from the Kirby list.

1. INTRODUCTION

A well-known problem in the smooth topology of simply-connected closed oriented 4-manifolds is to determine for a given second integral homology class the minimal genus of a smoothly embedded closed connected oriented surface representing that class. This problem dates back to the work of Kervaire and Milnor [6] and has been studied a lot since the introduction of the Donaldson and Seiberg-Witten invariants. The problem is harder and less is known in the case of homology classes of negative self-intersection (if the manifold does not admit an orientation reversing self-diffeomorphism). This is partly due to the fact that the Seiberg-Witten invariants are sensitive to the orientation of the 4-manifold. If the self-intersection number gets very negative the adjunction inequality [13] from Seiberg-Witten theory reduces to the fact that the genus of the surface is at least zero. For example, the minimal genus problem for the K3 surface for classes of non-negative square has been solved completely [10]. On the other hand, not much is known for classes of negative square. Finashin and Mikhalkin [2] constructed for any negative even number $n \geq -86$ an embedded sphere in the K3 surface with this self-intersection number; see also [1]. In general, Problem 4.105 from the Kirby list [7] asks if there is a (negative) lower bound on the self-intersection of embedded spheres in a given 4-manifold X (a 4-manifold with non-vanishing Seiberg-Witten invariants does not contain embedded spheres of positive self-intersection). One can also ask the following question: Given a genus g, is there a lower bound on the self-intersection number of embedded surfaces of genus g in the 4-manifold X?

In this note, we will show that there is a lower bound on the self-intersection number if we fix the type of the homology class A in following sense: We assume

Date: June 8, 2018.

²⁰¹⁰ Mathematics Subject Classification. Primary 57R95; Secondary 57N13, 57R57.

Key words and phrases. 4-manifold, homology class, branched covering, minimal genus.

M. J. D. HAMILTON

that the integral homology class A is divisible by an integer greater than one or that it is characteristic. The proof in both cases uses branched coverings in a similar way to the paper [8] due to Kotschick and Matić together with the solution of the $\frac{5}{4}$ -conjecture due to Furuta [3]. There is also another approach using the Gsignature theorem [5, 14] that leads to the best results for divisible classes in our situation. The result for divisible classes therefore is implicit in these papers, but it is a worthwhile preparation for the case of a characteristic class. In this case the other ingredient is a lemma that allows to represent a multiple of a given homology class by a surface of a certain genus. The argument here seems to work only with the $\frac{5}{4}$ -theorem.

Note that Kervaire and Milnor have shown that $A^2 \equiv \sigma(X) \mod 16$ if A is a characteristic homology class represented by an embedded sphere, where $\sigma(X)$ denotes the signature of X. The following lower bound on A^2 is a consequence of Theorem 4.2.

Corollary 1.1. Let X be a simply-connected closed oriented 4-manifold and A a characteristic class represented by an embedded sphere with $A^2 < 0$. Then

$$A^{2} \ge -(4b_{2}(X) - 5\sigma(X) - 8).$$

The spheres in the example of Finashin and Mikhalkin represent indivisible homology classes in the K3 surface. Unfortunately, the method presented here does not give a lower bound on the self-intersection number in this case. In the final section we indicate a strategy how to obtain a bound in the general situation, that depends on an open, yet not completely unrealistic, conjecture.

2. Branched coverings and the $\frac{5}{4}$ -theorem

We first need some preparations. From now on, X will denote a closed oriented simply-connected smooth 4-manifold whose second Betti number is nonzero, so that X is not a homotopy sphere (the results in this paper remain valid if $H_1(X;\mathbb{Z}) = 0$). All embedded surfaces in the following will be connected, closed, oriented and smoothly embedded. Suppose that $B \subset X$ is such an embedded surface in X of genus g_B , whose homology class $[B] \in H_2(X;\mathbb{Z})$ is divisible by a prime power $q = p^r$, where p is a prime and $r \ge 1$ an integer. We can then consider the q-fold cyclic branched covering $\phi: Y \to X$, branched over B. The second Betti number and the signature Y are known from [4, 5, 8, 14]. They are given by:

$$b_2(Y) = qb_2(X) + 2(q-1)g_B$$

$$\sigma(Y) = q\sigma(X) - \frac{q^2 - 1}{3q}B^2.$$

The formula for the second Betti number follows from the well-known formula for the Euler characteristic of a branched covering and the fact that the first Betti number of Y is zero. In addition it is known [12] that Y is spin if and only if the class $(q-1)\frac{1}{q}[B]$ is characteristic, i.e. the mod 2 reduction of its Poincaré dual is equal to the second Stiefel-Whitney class $w_2(X)$. Hence if q is odd, then Y is

spin if and only if X is spin. If q is even, then Y is spin if and only if $\frac{1}{q}[B]$ is characteristic.

In any case, we have the inequality

$$b_2(Y) \ge |\sigma(Y)|.$$

Suppose that Y is a spin manifold. Under our assumptions the second Betti number of Y is non-zero. In this situation Furuta's theorem [3] on the $\frac{5}{4}$ -conjecture applies and shows that

$$b_2(Y) \ge \frac{5}{4}|\sigma(Y)| + 2.$$

3. The case of divisible A

The following theorem can be deduced from results in [5, 14] due to Hsiang-Szczarba and Rohlin.

Theorem 3.1. Let X be a simply-connected 4-manifold and A an embedded surface in X of genus g_A and self-intersection $A^2 = -N < 0$. Suppose that the integral homology class [A] is divisible by a prime power q > 1. Then

(3.1)
$$N \le 2(b_2(X) - \sigma(X)) + 4g_A$$

if q = 2 and

(3.2)
$$N \le \frac{2q^2}{q^2 - 1}(b_2(X) - \sigma(X)) + \frac{4q^2}{q^2 - 1}g_A$$

if q is an odd prime power. Hence for fixed genus g_A there is an upper bound on N, depending only on X and q. For an arbitrary odd prime power $q \ge 3$ we have

(3.3)
$$N \le \frac{9}{4}(b_2(X) - \sigma(X)) + \frac{9}{2}g_A,$$

independent of q.

Proof. In [14] it is shown using branched coverings over X, branched along A, that

$$g_A \ge \left|\frac{\sigma(X)}{2} - \frac{A^2}{4}\right| - \frac{b_2(X)}{2}$$

if q = 2 and

$$g_A \ge \left|\frac{\sigma(X)}{2} - \frac{(q^2 - 1)A^2}{4q^2}\right| - \frac{b_2(X)}{2}$$

if q is odd. This implies inequalities (3.1) and (3.2). Suppose $q \ge 3$. With $b_2(X) \ge \sigma(X)$ we have

$$N \le \frac{2}{1 - \frac{1}{q^2}} (b_2(X) - \sigma(X)) + \frac{4}{1 - \frac{1}{q^2}} g_A$$
$$\le \frac{9}{4} (b_2(X) - \sigma(X)) + \frac{9}{2} g_A.$$

This is inequality (3.3).

M. J. D. HAMILTON

Note that the bound for q odd is greater than the one for q = 2 and tends to this one for large q. In any case q^2 has to divide N. Hence for a given prime power q and genus g_A there is only a small finite set of possible values for N. For completeness we also derive a bound using the $\frac{5}{4}$ -conjecture. The inequality coming from this conjecture is more crucial in the case of a characteristic class in Section 4.

Theorem 3.2. Let X be a simply-connected 4-manifold and A an embedded surface in X of genus g_A and self-intersection $A^2 = -N < 0$. Suppose that the integral homology class [A] is divisible by a prime power q > 1 and let $[C] = \frac{1}{q}[A]$. Assume also one of the following:

- If q is even, then the class [C] is characteristic.
- If q is odd, then X is spin.

Then

(3.4)
$$N \leq \frac{3q^2}{q^2 - 1} \left(\frac{4}{5} b_2(X) - \sigma(X) - \frac{8}{5q} \right) + \frac{24}{5} \frac{q}{q+1} g_A.$$

Hence for fixed genus g_A there is an upper bound on N, depending only on X and q. If X is spin and q odd, then

(3.5)
$$N \le \frac{27}{40} (4b_2(X) - 5\sigma(X)) + \frac{24}{5}g_A,$$

independent of q.

Proof. We consider the branched covering $Y \to X$ as in Section 2 with B = A. Note that under our assumptions the manifold Y is spin. Hence Furuta's theorem applies, which leads to inequality (3.4). Suppose that X is spin and $q \ge 3$. An argument similar to the one above leads to inequality (3.5). Here we have to use that $4b_2(X) - 5\sigma(X)$ is non-negative according to the $\frac{5}{4}$ -inequality applied to X. \Box

Example 3.3. Let X be the K3 surface and A a class divisible by two. With $b_2(X) = 22$ and $\sigma(X) = -16$ we get from inequality (3.1)

$$N \leq 76 + 4g_A$$
.

In particular, for $g_A = 0$, we see that there is an upper bound on N of 76 for homology classes divisible by 2 that are represented by an embedded sphere. We can use inequality (3.3) to derive the following upper bound on N if the class A is divisible by an odd prime:

$$N \le 85.5 + 4.5g_A.$$

4. The case of characteristic A

We begin with the following lemma:

Lemma 4.1. Let A be an embedded surface in X of genus g_A with self-intersection $A^2 = -N < 0$. Then the class [B] = 2[A] can be represented by a surface B of genus $g_B = N - 1 + 2g_A$.

4

Proof. The proof is elementary: Let A' be a section of the normal disk bundle of A, considered as a tubular neighbourhood, having precisely N transverse zeroes. Smoothing the intersections between A' and A results in the surface B. The claim also follows from the more general Lemma 7.7 in [9] under switching the orientation of X.

We can now prove the main theorem in this section.

Theorem 4.2. Let X be a simply-connected 4-manifold and A an embedded surface in X of genus g_A with self-intersection $A^2 = -N < 0$ such that the class [A] is characteristic. Then

(4.1)
$$N \le 4b_2(X) - 5\sigma(X) - 8 + 8g_A$$

Hence for fixed genus g_A there is an upper bound on N, depending only on X.

Proof. Let [B] = 2[A]. According to Lemma 4.1, the class [B] is represented by a surface B of genus $g_B = N - 1 + 2g_A$. Let $Y \to X$ be the 2-fold branched covering, branched over B. Then Y is spin and Furuta's theorem applies. We get:

$$2b_2(X) + 2g_B \ge \frac{5}{4}|2\sigma(X) + 2N| + 2 \ge \frac{5}{4}(2\sigma(X) + 2N) + 2.$$

This implies inequality (4.1).

Remark 4.3. Note that this idea only works for the 2-fold branched covering and not for higher degrees with the generalization of Lemma 4.1 due to the *q*-coefficients in the $\frac{5}{4}$ -inequality. This is still the case if we replace the $\frac{5}{4}$ -theorem by the $\frac{11}{8}$ -conjecture. The argument does not work with the inequality $b_2(Y) \ge |\sigma(Y)|$ nor with the *G*-signature theorem.

According to Kervaire and Milnor [6] a characteristic homology class A represented by a sphere has to satisfy

$$N = -A^2 \equiv -\sigma(X) \mod 16.$$

Together with the inequality in Theorem 4.2 this can be used to restrict the possible self-intersection numbers of characteristic homology classes represented by spheres.

Example 4.4. Let $X = \mathbb{CP}^2 \# k \overline{\mathbb{CP}}^2$ with $k \ge 1$ and A a characteristic homology class of self-intersection $A^2 = -N < 0$ represented by a sphere. By Theorem 4.2

$$N \le 9(k-1)$$

and according to Kervaire and Milnor

$$N \equiv k - 1 \bmod 16.$$

Hence if k = 2 then N = 1, if k = 3 then N = 2 or 18, and so on; compare with [11].

M. J. D. HAMILTON

5. The case of indivisible A

We consider the following conjecture, motivated by the results in Section 3.

Conjecture 5.1. Let X be a simply-connected 4-manifold and B an embedded surface in X of genus g_B with self-intersection $B^2 = -M < 0$ such that the class [B] is divisible by 2. Then

$$M \le c(X) + \kappa g_B,$$

where c(X) is a number that depends only on X and κ is a number less than 4.

Assuming that this is true we can derive a bound on $A^2 = -N$ without any assumptions on the class [A].

Theorem 5.2. Assume that Conjecture 5.1 holds. Let X be a simply-connected 4-manifold and A an embedded surface in X of genus g_A with self-intersection number $A^2 = -N < 0$. Then

(5.1)
$$N \le \frac{1}{4-\kappa} (c(X) + \kappa (2g_A - 1)).$$

Proof. The argument is a generalization of the proof of Theorem 4.2. Let [B] = 2[A]. According to Lemma 4.1, the class [B] is represented by a surface B of genus $g_B = N - 1 + 2g_A$. We have $M = -B^2 = 4N$. Hence Conjecture 5.1 implies

$$4N \le c(X) + \kappa(N - 1 + 2g_A)$$

This leads to inequality (5.1).

Remark 5.3. The inequality $b_2(Y) \ge |\sigma(Y)|$ and the inequality from the *G*-signature theorem have $\kappa = 4$. The $\frac{5}{4}$ -conjecture has $\kappa = \frac{16}{5} < 4$, but only if $\frac{1}{2}[B]$ is characteristic. This leads to the same bound as in Theorem 4.2. We could also formulate a conjecture if the class [B] is divisible by some other positive integer *d*. If [B] = d[A] with $A^2 = -N < 0$ and *A* is a surface of genus g_A , then [B] is represented according to Lemma 7.7 in [9] by a surface of genus

$$g_B = \frac{1}{2}d(d-1)N + 1 - d + dg_A.$$

In the conjecture we have to replace $\kappa < 4$ by $\kappa < \frac{2d}{d-1}$. However, for d > 2, this conjecture tends to get unrealistic.

REFERENCES

- 1. S. Akbulut, A fake compact contractible 4-manifold, J. Differential Geom. 33, 335–356 (1991).
- 2. S. Finashin, G. Mikhalkin, A (-86)-sphere in the K3 surface, Turkish J. Math. **21**, 129–131 (1997).
- 3. M. Furuta, Monopole equation and the $\frac{11}{8}$ -conjecture, Math. Res. Lett. 8, no. 3, 279–291 (2001).
- F. Hirzebruch, *The signature of ramified coverings*, in: Global Analysis (Papers in Honor of K. Kodaira), 253–265, Univ. Tokyo Press, Tokyo 1969.
- W. C. Hsiang, R. H. Szczarba, On embedding surfaces in four-manifolds, Proc. Symp. Pure Math. 22, 97–103 (1971).
- M. A. Kervaire, J. W. Milnor, On 2-spheres in 4-manifolds, Proc. Nat. Acad. Sci. U.S.A. 47, 1651–1657 (1961).

- 7. R. Kirby (Ed.), *Problems in Low-Dimensional Topology*, Dec. 22, 1995. http://www.math.berkeley.edu/~kirby/
- D. Kotschick, G. Matić, *Embedded surfaces in four-manifolds, branched covers, and* SO(3)invariants, Math. Proc. Cambridge Philos. Soc. 117, 275–286 (1995).
- 9. P. B. Kronheimer, T. S. Mrowka, *Embedded surfaces and the structure of Donaldson's polynomial invariants*, J. Differential Geom. **41**, 573–734 (1995).
- 10. T. Lawson, The minimal genus problem, Exposition. Math. 15, 385-431 (1997).
- 11. B.-H. Li, T.-J. Li, Smooth minimal genera for small negative classes in $\mathbb{C}P^2 \# n\overline{\mathbb{C}P^2}$ with $n \leq 9$, Topology Appl. 132, 1–15 (2003).
- S. Nagami, Existence of Spin structures on cyclic branched covering spaces over fourmanifolds, in: Perspectives of complex analysis, differential geometry and mathematical physics (St. Konstantin, 2000), 86–92, World Sci. Publ., River Edge, NJ, 2001.
- 13. P. Ozsváth, Z. Szabó, The symplectic Thom conjecture, Ann. of Math. (2) 151, 93–124 (2000).
- 14. V. A. Rohlin, *Two-dimensional submanifolds of four-dimensional manifolds*, Func. Analysis Appl. **5**, 39–48 (1971).

INSTITUTE FOR GEOMETRY AND TOPOLOGY, UNIVERSITY OF STUTTGART, PFAFFENWALDRING 57, 70569 STUTTGART, GERMANY

E-mail address: mark.hamilton@math.lmu.de