Comment on a recent attempt to explain the 'GSI anomaly' by initial-state e.m. spin-rotation coupling*

A. Gal¹

¹Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

A recently proposed solution [1, 2] of the 'GSI anomaly' by spin precession of the decaying heavy ions in the magnetic field that controls their circular motion at the GSI storage ring is dubious: the uncertainty in the computed electron-capture decay-rate modulation frequency is at least of order 10^7 Hz, by far exceeding the 1 Hz modulation frequency reported in the GSI experiment [3].

PACS numbers: 13.40.Em, 21.10.Ky, 23.40.-s

Two-body electron-capture decay rates of ¹⁴⁰Pr and ¹⁴²Pm hydrogen-like heavy ions coasting at the GSI Experimental Storage Ring (ESR) were reported to be modulated with period $\sim 7 \text{ s}$ [3]. This modulation is known as the 'GSI anomaly' [4]. Preliminary data exist also for ¹²²I [5]. It was concluded in Refs. [6, 7] that spin precession of the decaying ions in the ESR magnetic field $B \approx 1.2 \text{ T}$ could not give rise to such modulation since the precession frequency associated with the electron is of order $|\mu_e|B \approx 3.3 \cdot 10^{10}$ Hz and that associated with the nucleus is of order $\mu_p B \approx 2.5 \cdot 10^7$ Hz, where $\mu_{e(p)}$ is the electron (proton) magnetic moment, both these frequencies exceeding by many orders of magnitude the reported modulation frequency $P_{\rm GSI} \sim 1$ Hz. We note that electron-capture decay rates of implanted neutral atoms, not exposed to external magnetic fields, do not show evidence for GSI-like modulation [8, 9].




FIG. 1: $P \equiv P_{\rm LPS} - P_{\rm GSI}$ (Hz units) in $^{140}{\rm Pr}$ as a function of the relativistic Lorentz factor $\gamma_{e|n}$. Here, $P_{\rm GSI} \sim 1$ Hz and $P_{\rm LPS} = a(\gamma_{e|n})\mu_e B + b(\gamma_{e|n}, g_{\rm nucl})\mu_p B$, a and b depending on $\gamma_{e|n}$ and $g_{\rm nucl}$. Note that a 0.1% variation of $\gamma_{e|n}$ induces $\approx 4\cdot 10^7$ variation of P. Figure adapted from Ref. [1].

Recently, however, Lambiase et al. [1, 2] argued that a full treatment of spin-rotation coupling results in cancelations among components of the spin precession frequency $P_{\rm LPS}$, bringing $P_{\rm LPS}$ down to the level of $P_{\rm GSI} \sim 1$ Hz. Their P_{LPS} is responsible for splitting the two approximately $F = \frac{1}{2}$, $m = \pm \frac{1}{2}$ magnetic states of the rotating ion. Fig. 1 adapted from Ref. [1] shows the dependence of $P \equiv P_{\text{LPS}} - P_{\text{GSI}}$ on the relativistic Lorentz factor $\gamma_{e|n}$ in ¹⁴⁰Pr. To sustain a zero value for P, $(\gamma_{e|n}-1)$ must be known accurately to at least three digits in order to trust the reduction of the μ_e term in $P_{\rm LPS}$ (see caption) from order $10^{10}~\mathrm{Hz}$ down to order $10^7~\mathrm{Hz}$ at which it might get canceled with the μ_p term; and moreover, the nuclear gyromagnetic ratio g_{nucl} must be known accurately to at least seven digits so as to reduce the order of magnitude of P_{LPS} from 10⁷ Hz further down to $P_{\rm GSI} \sim 1$ Hz. In practice, $(\gamma_{e|n} - 1)$ is estimated semiclassically in Ref. [1] to a single digit only, and g_{nucl} is either experimentally unknown or known to a single digit only: $g_{\text{nucl}}(^{122}\text{I})=0.94\pm0.03$ [10]. Similar arguments were given by Faestermann [11] in response to earlier attempts by Lambiasse et al. to publish this kind of wishful thinking.

In summary, atomic and nuclear magnetic moments are much too large to resolve the 'GSI anomaly' as attempted by Lambiasse *et al.* [1, 2]. Magnetic moments of order $10^{-11}\mu_B$, many orders of magnitude below those provided by atoms and nuclei but less than one order of magnitude below the laboratory upper limit for neutrinos, are required to explain the 'GSI anomaly' by spin precession [12].

*In memoriam Paul Kienle (1931-2013) who pursued the 'GSI anomaly' with boundless imagination [3–6].

G. Lambiase, G. Papini, G. Scarpetta, Phys. Lett. B 718 (2013) 998 [arXiv:1205.0684(nucl-th)].

^[2] G. Lambiase, et al., Ann. Phys. 332 (2013) 143.

Yu.A. Litvinov, et al., Phys. Lett. B 664 (2008) 162;
P. Kienle, et al., Phys. Lett. B 726 (2013) 638.

^[4] P. Kienle, Found. Phys. 40 (2010) 733.

^[5] P. Kienle, Prog. Part. Nucl. Phys. 64 (2010) 439.

^[6] M. Faber, et al., J. Phys. G 37 (2010) 015102.

^[7] I.M. Pavlichenkov, Phys. Rev. C 81 (2010) 051602.

^[8] P.A. Vetter, et al., Phys. Lett. B 670 (2008) 196.

^[9] T. Faestermann, et al., Phys. Lett. B 672 (2009) 227.

^[10] V.R. Green, et al., Phys. Lett. B 173 (1986) 115.

^[11] T. Faestermann, arXiv:0907.1557(nucl-th).

^[12] A. Gal, Nucl. Phys. A 842 (2010) 102.