

On varieties of Hilbert type

Lior Bary-Soroker, Arno Fehm and Sebastian Petersen

September 25, 2018

Abstract

A variety X over a field K is of Hilbert type if $X(K)$ is not thin. We prove that if $f: X \rightarrow S$ is a dominant morphism of K -varieties and both S and all fibers $f^{-1}(s)$, $s \in S(K)$, are of Hilbert type, then so is X . We apply this to answer a question of Serre on products of varieties and to generalize a result of Colliot-Thélène and Sansuc on algebraic groups.

1 Introduction

In the terminology of thin sets (we recall this notion in Section 2), Hilbert's irreducibility theorem asserts that $\mathbb{A}_K^n(K)$ is not thin, for any number field K and any $n \geq 1$. As a natural generalization a K -variety X is called of Hilbert type if $X(K)$ is not thin. The importance of this definition stems from the observation of Colliot-Thélène and Sansuc [2] that the inverse Galois problem would be settled if every unirational variety over \mathbb{Q} was of Hilbert type.

In this direction, Colliot-Thélène and Sansuc [2, Corollary 7.15] prove that any connected reductive algebraic group over a number field is of Hilbert type. This immediately raises the question whether the same holds for all linear algebraic groups (note that these are unirational). Another question, asked by Serre [15, p. 21], is whether a product of two varieties of Hilbert type is again of Hilbert

type. The main result of this paper gives a sufficient condition for a variety to be of Hilbert type:

Theorem 1.1. *Let K be a field and $f: X \rightarrow S$ a dominant morphism of K -varieties. Assume that the set of $s \in S(K)$ for which the fiber $f^{-1}(s)$ is a K -variety of Hilbert type is not thin. Then X is of Hilbert type.*

As an immediate consequence we get the following result for a family of varieties over a variety of Hilbert type:

Corollary 1.2. *Let K be a field and $f: X \rightarrow S$ a dominant morphism of K -varieties. Assume that S is of Hilbert type and that for every $s \in S(K)$ the fiber $f^{-1}(s)$ is of Hilbert type. Then X is of Hilbert type.*

Using this result we resolve both questions discussed above affirmatively, see Corollary 3.4 and Proposition 4.2.

2 Background

Let K be a field. A K -variety is a separated scheme of finite type over K which is geometrically reduced and geometrically irreducible. Thus, an open subscheme of a K -variety is again a K -variety. If $f: X \rightarrow S$ is a morphism of K -varieties and $s \in S(K)$, then $f^{-1}(s) := X \times_S \text{Spec}(\kappa(s))$, where $\kappa(s)$ is the residue field of s , denotes the scheme theoretic fiber of f at s . This fiber is a separated scheme of finite type over K , which needs not be reduced or connected in general. We identify the set $f^{-1}(s)(K)$ of K -rational points of the fiber with the set theoretic fiber $\{x \in X(K) \mid f(x) = s\}$.

Let X be a K -variety. A subset T of $X(K)$ is called *thin* if there exists a proper Zariski-closed subset C of X , a finite set I , and for each $i \in I$ a K -variety Y_i with $\dim Y_i = \dim X$ and a dominant separable morphism $p_i: Y_i \rightarrow X$ of degree ≥ 2 such that

$$T \subseteq \bigcup_{i \in I} p_i(Y_i(K)) \cup C(K).$$

A K -variety X is of *Hilbert type* if $X(K)$ is not thin, cf. [15, Definition 3.1.2]. Note that X is of Hilbert type if and only if some (or every) open subscheme of X is of Hilbert type, cf. [15, p. 20]. A field K is *Hilbertian* if \mathbb{A}_K^1 is of Hilbert type. We note that if there exists a K -variety X of positive dimension such that X is of Hilbert type, then K is Hilbertian [5, Proposition 13.5.3].

3 Proof of Theorem 1.1

A key tool in the proof of Theorem 1.1 is the following consequence of Stein factorization.

Lemma 3.1. *Let K be a field and $\psi: Y \rightarrow S$ a dominant morphism of normal K -varieties. Then there exists a nonempty open subscheme $U \subset S$, a K -variety T and a factorization*

$$\psi^{-1}(U) \xrightarrow{g} T \xrightarrow{r} U$$

of ψ such that the fibers of g are geometrically irreducible and r is finite and étale.

Proof. See [10, Lemma 9]. □

Lemma 3.2. *Let K be a field and $f: X \rightarrow S$ a dominant morphism of normal K -varieties. Assume that the set Σ of $s \in S(K)$ for which $f^{-1}(s)$ is a K -variety of Hilbert type is not thin. Let I be a finite set and let $p_i: Y_i \rightarrow X$, $i \in I$, be finite étale morphisms of degree ≥ 2 . Then $X(K) \not\subseteq \bigcup_{i \in I} p_i(Y_i(K))$.*

Proof. For $i \in I$ consider the composite morphism $\psi_i := f \circ p_i: Y_i \rightarrow S$. By Lemma 3.1 there is a nonempty open subscheme U_i of S and a factorization

$$\psi_i^{-1}(U_i) \xrightarrow{g_i} T_i \xrightarrow{r_i} U_i$$

of ψ_i such that the morphism g_i has geometrically irreducible fibers, r_i is finite and étale, and such that T_i is a K -variety. We now replace successively S by $\bigcap_{i \in I} U_i$, X by $f^{-1}(S)$, T_i by $r_i^{-1}(S)$ and Y_i by $p_i^{-1}(X)$, to assume in addition that $r_i: T_i \rightarrow S$ is finite étale for every $i \in I$.

For $s \in S(K)$ denote by $X_s := f^{-1}(s)$ the fiber of f over s . Then X_s is a K -variety of Hilbert type for each $s \in \Sigma$. Furthermore we define $Y_{i,s} := \psi_i^{-1}(s)$ and let $p_{i,s}: Y_{i,s} \rightarrow X_s$ be the corresponding projection morphism. Then $p_{i,s}$ is a finite étale morphism of the same degree as p_i . In particular, the K -scheme $Y_{i,s}$ is geometrically reduced. For every $s \in S(K)$ and every $i \in I$ we have constructed a commutative diagram

$$\begin{array}{ccccc} Y_{i,s} & \longrightarrow & Y_i & \xrightarrow{g_i} & T_i \\ p_{i,s} \downarrow & & p_i \downarrow & \searrow \psi_i & \downarrow r_i \\ X_s & \longrightarrow & X & \xrightarrow{f} & S \end{array}$$

in which the left hand rectangle is cartesian. Set $J := \{i \in I : \deg(r_i) \geq 2\}$. Then $\bigcup_{i \in J} r_i(T_i(K)) \subseteq S(K)$ is thin, so by assumption there exists

$$s \in \Sigma \setminus \bigcup_{i \in J} r_i(T_i(K)).$$

For $i \in J$ there is no K -rational point of T_i over s , hence $Y_{i,s}(K) = \emptyset$ for every $i \in J$. For $i \in I \setminus J$, the finite étale morphism r_i is of degree 1, hence an isomorphism, and therefore $Y_{i,s}$ is geometrically irreducible. Thus, $Y_{i,s}$ is a K -variety. So since X_s is of Hilbert type, there exists $x \in X_s(K)$ such that $x \notin \bigcup_{i \in I \setminus J} p_{i,s}(Y_{i,s}(K))$. Thus

$$x \notin \bigcup_{i \in J \setminus I} p_{i,s}(Y_{i,s}(K)) = \bigcup_{i \in I} p_{i,s}(Y_{i,s}(K)),$$

hence $x \notin \bigcup_{i \in I} p_i(Y_i(K))$, as needed. \square

We shall use the following well-known fact.

Lemma 3.3. *Let K be a field, let X, Y be K -varieties with $\dim X = \dim Y$, and let $p: Y \rightarrow X$ be a dominant separable morphism. Then there exists a nonempty open subscheme U of X such that the restriction of p to a morphism $p^{-1}(U) \rightarrow U$ is finite and étale.*

Proof of Theorem 1.1. Let K be a field, and $f: X \rightarrow S$ a dominant morphism of K -varieties. Assume that the set Σ of those $s \in S(K)$ for which $f^{-1}(s)$ is of Hilbert type is not thin. Let $C \subseteq X$ be a proper Zariski-closed subset. Let I be a finite set and suppose that Y_i is a K -variety with $\dim(Y_i) = \dim(X)$ and

$p_i: Y_i \rightarrow X$ is a dominant separable morphism of degree ≥ 2 , for every $i \in I$. We have to show that $X(K) \not\subseteq C(K) \cup \bigcup_{i \in I} p_i(Y_i(K))$.

By Lemma 3.3 and [8, 6.12.6, 6.13.5] there exists a normal nonempty open subscheme $X' \subset X \setminus C$ such that the restriction of each p_i to a morphism $p_i^{-1}(X') \rightarrow X'$ is finite and étale. The image $f(X')$ contains a nonempty open subscheme S' of S (cf. [7, 1.8.4], [6, 9.2.2]). Furthermore, S' contains a nonempty normal open subscheme S'' . Let us define $X'' := f^{-1}(S'') \cap X'$ and $Y_i'' := p_i^{-1}(X'')$. Then the restriction of f to a morphism $f'': X'' \rightarrow S''$ is a surjective morphism of normal K -varieties, $\Sigma \cap S''(K)$ is not thin, and $f''^{-1}(s)$ is of Hilbert type for every $s \in \Sigma \cap S''(K)$ because it is an open subscheme of $f^{-1}(s)$. The restriction p_i'' of p_i to a morphism $Y_i'' \rightarrow X''$ is finite and étale for every $i \in I$. By Lemma 3.2 applied to f'' and the p_i'' we have

$$\begin{aligned} \emptyset &\neq X''(K) \setminus \bigcup_{i \in I} p_i''(Y_i''(K)) \\ &= X''(K) \setminus \bigcup_{i \in I} p_i(Y_i(K)) \\ &\subseteq X(K) \setminus \left(C(K) \cup \bigcup_{i \in I} p_i(Y_i(K)) \right), \end{aligned}$$

so $X(K) \not\subseteq C(K) \cup \bigcup_{i \in I} p_i(Y_i(K))$, as needed. \square

As an immediate consequence we get an affirmative solution of Serre's question mentioned in the introduction.

Corollary 3.4. *Let K be a field. If X, Y are K -varieties of Hilbert type, then $X \times Y$ is of Hilbert type.*

Proof. Denote by $f: X \times Y \rightarrow X$ the projection. Then $f^{-1}(x)$ is isomorphic to Y and hence of Hilbert type for every $x \in X(K)$. Thus Corollary 1.2 gives that $X \times Y$ is of Hilbert type. \square

4 Algebraic groups of Hilbert type

By an *algebraic group* over a field K we shall mean a connected smooth group scheme over K . Recall that such an algebraic group is a K -variety, see [9, Exp VI_A, 0.3, 2.1.2, 2.4]. If G is an algebraic group over K , then $G(K_s)$ is a $\text{Gal}(K)$ -group, where K_s denotes a separable closure of K and $\text{Gal}(K) = \text{Gal}(K_s/K)$ is the absolute Galois group of K , and there is an associated Galois cohomology pointed set $H^1(K, G) = H^1(\text{Gal}(K), G(K_s))$, which classifies isomorphism classes of $G(K_s)$ -torsors, cf. [12, Prop. 1.2.4].

Proposition 4.1. *Let K be a field and let*

$$1 \rightarrow N \rightarrow G \xrightarrow{p} Q \rightarrow 1$$

be a short exact sequence of algebraic groups over K . If $H^1(K, N) = 1$ and both N and Q are of Hilbert type, then G is of Hilbert type.

Proof. It suffices to show that $p^{-1}(x)$ is of Hilbert type for every $x \in Q(K)$, because then Corollary 1.2 implies the assertion. Let $x \in Q(K)$ and $F = p^{-1}(x)$. There is an exact sequence of $\text{Gal}(K)$ -groups

$$1 \rightarrow N(K_s) \rightarrow G(K_s) \rightarrow Q(K_s) \rightarrow 1,$$

where the right hand map is surjective, because for every point $x \in Q(K_s)$ the fiber over x is a non-empty K_s -variety and thus has a K_s -rational point. Since the $\text{Gal}(K)$ -set $F(K_s)$ is a coset of $N(K_s)$, it is a $N(K_s)$ -torsor. Our hypothesis $H^1(K, N) = 1$ implies that $F(K_s)$ is isomorphic to the trivial $N(K_s)$ -torsor $N(K_s)$. It follows that F is isomorphic to N as a K -variety, hence F is of Hilbert type. \square

Using this, we generalize the result of Colliot-Thélène and Sansuc [2, Corollary 7.15] from reductive groups to arbitrary linear groups.

Theorem 4.2. *Every linear algebraic group G over a perfect Hilbertian field K is of Hilbert type.*

Proof. We denote by G_u the unipotent radical of G (cf. [11, Proposition XVII.1.2]). We have a short exact sequence of algebraic groups over K

$$1 \rightarrow G_u \rightarrow G \rightarrow Q \rightarrow 1 \quad (*)$$

with Q reductive, cf. [11, Proposition XVII.2.2]. By [2, Corollary 7.15], Q is of Hilbert type. Since K is perfect, G_u is split, i.e. there exists a series of normal algebraic subgroups

$$1 = U_0 \subseteq \cdots \subseteq U_n = G_u$$

such that $U_{i+1}/U_i \cong \mathbb{G}_a$ for each i , cf. [1, 15.5(ii)]. The groups U_i are unipotent, hence $H^1(K, U_i) = 1$ by [14, Ch. III.2.1, Prop. 6], and \mathbb{G}_a is of Hilbert type since K is Hilbertian. Thus, an inductive application of Proposition 4.1 implies that G_u is of Hilbert type. Finally we apply Proposition 4.1 to the exact sequence $(*)$ to conclude that G is of Hilbert type. \square

Remark 4.3. The special case of Theorem 4.2 where G is simply connected and K is finitely generated is also a consequence of a result of Corvaja, see [3, Corollary 1.7].

Remark 4.4. We point out that Theorem 4.2 could be deduced also from Corollary 3.4 (instead of 1.1) via [13, Corollary 1] and the fact that a unipotent group over a perfect field is rational, cf. [9, XIV, 6.3].

As a consequence of Theorem 4.2, we get a more general statement for homogeneous spaces, which was pointed out to us by Borovoi:

Corollary 4.5. *If G is a linear algebraic group over a perfect Hilbertian field K , and H is algebraic subgroup of G , then the quotient G/H is of Hilbert type.*

Proof. For the existence of the quotient $Q := G/H$ see for example [1, Chapter II Theorem 6.8]. If \mathcal{H} denotes the generic fiber of $G \rightarrow Q$ and \bar{F} is an algebraic closure of the function field $K(Q)$ of Q , then $\mathcal{H}_{\bar{F}} \cong H_{\bar{F}}$ by translation on G . Thus, \mathcal{H} is geometrically irreducible since H is, so [2, Proposition 7.13] implies that Q is of Hilbert type. \square

We also get a complete classification of the algebraic groups that are of Hilbert type over a number field:

Corollary 4.6. *An algebraic group G over a number field K is of Hilbert type if and only if it is linear.*

Proof. If G is linear, then it is of Hilbert type by Theorem 4.2. Conversely, assume that G is of Hilbert type. Chevalley's theorem [4, Theorem 1.1] gives a short exact sequence of algebraic groups over K ,

$$1 \rightarrow H \rightarrow G \rightarrow A \rightarrow 1$$

with H linear and A an abelian variety. As in the proof of Corollary 4.5 we conclude that the generic fiber of $G \rightarrow A$ is geometrically irreducible, and therefore A is of Hilbert type. Since no nontrivial abelian variety over a number field is of Hilbert type, cf. [5, Remark 13.5.4], A is trivial and $G \cong H$ is linear. \square

Acknowledgements

The authors would like to thank Mikhail Borovoi for pointing out to them Corollary 4.5, Jean-Louis Colliot-Thélène and Moshe Jarden for helpful comments on a previous version, and Daniel Krashen for some references concerning algebraic groups. This research was supported by the Lion Foundation Konstanz – Tel Aviv, the Alexander von Humboldt Foundation, the Hermann Minkowski Minerva Center for Geometry at Tel Aviv University, and by a grant from the GIF, the German-Israeli Foundation for Scientific Research and Development.

References

- [1] Armand Borel. *Linear algebraic groups*. Springer, 1991.
- [2] Jean-Louis Colliot-Thélène and Jean-Jacques Sansuc. Principal homogeneous spaces under flasque tori: applications. *Journal of Algebra*, 106:148–205, 1987.
- [3] Pietro Corvaja. Rational fixed points for linear groups. *Ann. Sc. Norm. Super. Pisa, Cl. Sci. (5)*, 6(4):561–597, 2007.

- [4] Brian Conrad. A modern proof of Chevalley's theorem. *J. Ramanujan Math. Soc.*, (17):1–18, 2002.
- [5] M. Fried and M. Jarden. *Field Arithmetic*. Ergebnisse der Mathematik III **11**. Springer, 2008. 3rd edition, revised by M. Jarden.
- [6] Alexander Grothendieck. Éléments de géométrie algébrique (rédigé avec la coopération de Jean Dieudonné): II. Étude cohomologique des faisceaux cohérents. *Publ. Math. IHES*, (11):5–167, 1961.
- [7] Alexander Grothendieck. Éléments de géométrie algébrique (rédigé avec la coopération de Jean Dieudonné): IV. Étude locale des schémas et des morphismes des schémas, Premier partie. *Publ. Math. IHES*, (20):5–259, 1964.
- [8] Alexander Grothendieck. Éléments de géométrie algébrique (rédigé avec la coopération de Jean Dieudonné): IV. Étude locale des schémas et des morphismes des schémas, Seconde partie. *Publ. Math. IHES*, (24):5–231, 1966.
- [9] Alexander Grothendieck. Séminaire de Géométrie Algébrique 3. *Springer*, 1970.
- [10] János Kollár. Rationally connected varieties and fundamental groups. In *Higher dimensional varieties and rational points*, p. 69–92. Springer, 2003.
- [11] James Milne. *Basic Theory of Affine Group Schemes*. Available at WWW.JMILNE.ORG, 2012
- [12] Jürgen Neukirch, Alexander Schmidt and Kay Wingberg. *Cohomology of Number Fields*. Springer, 2000.
- [13] Maxwell Rosenlicht. Questions of rationality for solvable algebraic groups over nonperfect fields. *Annali di matematica pura ed applicata*, 62:97-120, 1963.
- [14] Jean-Pierre Serre. *Galois Cohomology*. Second edition. Springer, 2002.
- [15] Jean-Pierre Serre. *Topics in Galois theory*. Second edition. Springer, 2008.