The corona algebra of the stabilized Jiang-Su algebra

Huaxin Lin and Ping Wong Ng

Abstract

Let \mathcal{Z} be the Jiang-Su algebra and \mathcal{K} the C^* -algebra of compact operators on an infinite dimensional separable Hilbert space. We prove that the corona algebra $M(\mathcal{Z} \otimes \mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$ has real rank zero. We actually prove a more general result.

1 Introduction

The Jiang-Su algebra \mathcal{Z} is a projectionless unital simple separable infinite dimensional amenable C^* -algebra with the same K-theory as that of the complex field \mathbb{C} (see [17]). It had been an interesting problem to find projectionless unital infinite dimensional simple C^* -algebras ([19], [2], [3], [8]). However, the Jiang-Su algebra is the nicest that one can get. In fact \mathcal{Z} is an inductive limit of sub-homogeneous C^* -algebras. It is one of many products of the decade of 1990's in the Elliott program, the program of classification of amenable C^* -algebras. However, more recently, \mathcal{Z} has played a much important role in the study of C^* -algebras. (See, for example, [54], [53], [33].). It becomes increasingly important to fully understand the structure of the Jiang-Su algebra.

In many ways, \mathcal{Z} behaves like \mathbb{C} , as its K-theory suggests. One indirect, but important feature, of the unital simple C^* -algebra \mathbb{C} , is the simplicity (as well as complexity) of the Calkin algebra $M(\mathbb{C} \otimes \mathcal{K})/\mathbb{C} \otimes \mathcal{K}$. Let A be a non-unital but σ -unital C^* -algebra. The quotient M(A)/A is called the corona algebra of A. The Calkin algebra is the corona algebra of $\mathbb{C}\otimes\mathcal{K}$, the stablization of \mathbb{C} . One of the important consequences of the simplicity of the Calkin algebra is that it has real rank zero. As we know, many important results in operator algebras (as well as operator theory) are related to the Calkin algebra, such as Fredholm index theory, the BDF-theory, C^* -algebra extension theory and the KK-theory. The fact that the Calkin algebra has real rank zero plays a crucial role in all these even though the term real rank zero was invented much later ([6]). From that point of view, the corona algebra $M(\mathcal{Z} \otimes \mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$, inevitably, is also important, given the central role \mathcal{Z} is playing in the study of classification of amenable C^* -algebras. By now, we know that the corona algebra of $\mathcal{Z} \otimes \mathcal{K}$ is not simple. In fact it has a unique proper closed ideal. Nevertheless, we would like to know whether the corona algebra of $\mathcal{Z} \otimes \mathcal{K}$ has some other similar structure to that of $\mathbb{C} \otimes \mathcal{K}$. The problem whether the corona algebra $M(\mathcal{Z} \otimes \mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$ has real rank zero has been around for many years and was specifically raised at the American Institute of Mathematics in 2009, as well as other places ([21]). Moreover, there is a long history of similar questions (e.g., ([6], [61], [15], [25], [27], [26], [39], [59], [60], [42]). The purpose of this paper is to establish that the corona algebra of $\mathcal{Z} \otimes \mathcal{K}$ does have real rank zero, despite the fact it is not simple and the fact that \mathcal{Z} has no proper projection.

One of the earliest applications of K-theory in operator algebras is the proof of the following theorem: an extension of AF-algebras is again an AF-algebra ([5] and [12]). This extends to a more general form: Let $0 \to B \to E \to A \to 0$ be a short exact sequence of C^* -algebras, where the ideal and quotient have real rank zero. Then E has real rank zero if and only if $K_1(B)$ is trivial (see [6]; also see [61]). Denote by $Q(\mathcal{Z})$ the corona algebra of $\mathcal{Z} \otimes \mathcal{K}$. Let J be the closed ideal of $M(\mathcal{Z} \otimes \mathcal{K})$ such that $\pi(J)$ is the unique proper nontrivial ideal of the corona algebra,

where $\pi: M(\mathcal{Z} \otimes \mathcal{K}) \to Q(\mathcal{Z})$ is the quotient map. We have that $Q(\mathcal{Z})/\pi(J)$ is purely infinite and simple (this is well-known; an explicit reference is [21]; it also follows immediately from [30] Theorem 3.5; see also [55] Theorem 2.2 and its proof). By a result of S. Zhang ([56]), it has real rank zero. It is also known that $\pi(J)$ is also purely infinite and simple (this is also well-known; an explicit reference is [21]; it also follows immediately from the definition of J in [24] 2.2, Remark 2.9 and Lemma 2.1; see also [55] Theorem 2.2 and its proof). Therefore, from the above mentioned result, if $K_1(J)$ is trivial, then $Q(\mathcal{Z})$ has real rank zero. Much of the work of this study is to show just that.

The general strategy to show that $K_1(J)$ is trivial is taken from an original idea of Elliott ([11]). However, unlike the case that Elliott considered, $A \otimes \mathcal{K}$ (in particular, in the case that $A = \mathcal{Z}$) lacks sufficiently many projections. Instead, we need to use positive elements and Cuntz comparison. The method to adapt Cuntz's relation to compare positive elements in the multiplier algebra of a non-unital simple C^* -algebra was initially used in [24] (and [28]). When A has real rank zero, one has the uniform bound for the length of path of unitaries in A which connects to the identity. This fact plays an important role in many of the earlier studies of multiplier algebras. In general, however, for a unital simple C^* -algebra A, the exponential length could be infinite. In other words, there will not be any control of the length. The idea to controlling the exponential length of unitaries in the multiplier algebras via the closure of the commutator subgroup instead of controlling the exponential length in $U_0(A)$ directly is new. We need results of exponential length of unitaries in the unitization of a non-unital simple C^* -algebra. The method to actually controlling the exponential length of unitaries in the closure of commutators for unital simple C^* -algebras was taken from [36] which is based on the results in the connection to the Elliott program of classification of amenable simple C^* -algebras.

The paper is organized as follows. Section 2 serves the purpose of setting up notation and terminology that will be used in subsequent sections. Section 3 contributes to the understanding of exponential length of unitaries in the closure of commutator subgroup of non-unital simple C^* -algebras. The computation of the exponential length plays critical role in dealing with unitary group of some ideals in the multiplier algebras and corona algebras of some simple C^* -algebras, in particular, those of $\mathcal{Z} \otimes \mathcal{K}$. Section 4 contains a number of technical lemmas that use the Cuntz relation to produce and deal with projections in the multiplier algebra of certain non-unital simple C^* -algebras. The main technical lemma is 4.15 which allows us to connect a unitary to one which is nontrivial only in a small corner. Section 5 contains the main result. We show that the corona algebra of $A \otimes \mathcal{K}$ has real rank zero for a class of unital simple C^* -algebras A which includes the projectionless simple C^* -algebra \mathcal{Z} . We expect a number of direct applications of the main result that $M(\mathcal{Z} \otimes \mathcal{K})/\mathcal{Z} \otimes \mathcal{K}$ has real rank zero. However, these would be the subject of future projects.

Acknowldgments This project began in 2009 when both authors visited American Institute of Mathematics during the Workshop on the Cuntz semigroup. It restarted in May 2012 when both authors were in the Research Center for Operator Algebras in East China Normal University and partially supported by the Center. We thank Leonel Robert for pointing out that the proof of Proposition 4.4 can be shortened.

2 Notations

Definition 2.1. For any C^* -algebra C, denote by T(C) the tracial state space of C. Let $\tau \in T(C)$ and let $n \geq 1$ be an integer. We will also use τ for the trace $\tau \otimes Tr$, where Tr is the standard (non-normalized) trace on M_n .

Definition 2.2. Let A be a unital C^* -algebra. Denote by U(A) the unitary group of A, by $U_0(A)$ the path connected component of U(A) containing the identity. Denote by CU(A) the

closure of the commutator group of $U_0(A)$.

Definition 2.3. Let A be a unital C^* -algebra with $T(A) \neq \emptyset$. Let $u \in U_0(A)$. Suppose that $\{u(t): t \in [0,1]\}$ is a continuous path of unitaries which is also piecewise smooth such that u(0) = u and u(1) = 1. Define the de la Harpe-Skandalis determinant as follows:

$$Det(u)(\tau) := Det(u(t))(\tau) := \int_{[0,1]} \tau(\frac{du(t)}{dt}u(t)^*) dt \text{ for all } \tau \in T(A).$$
 (e 2.1)

Note that, if $u_1(t)$ is another continuous path which is piecewise smooth with $u_1(0) = u$ and $u_1(1) = 1$, then $\text{Det}((u(t)) - \text{Det}(u_1(t)) \in \rho_A(K_0(A))$. Suppose that $u, v \in U(A)$ and $uv^* \in U_0(A)$. Let $\{w(t) : t \in [0,1]\} \subset U(A)$ be a piecewise smooth and continuous path such that w(0) = u and w(1) = v. Define

$$R_{u,v}(\tau) = \operatorname{Det}(w(t))(\tau) = \int_{[0,1]} \tau(\frac{dw(t)}{dt}w(t)^*)dt \text{ for all } \tau \in T(A).$$

Note that $R_{u,v}$ is well-defined (independent of the choices of the path) up to elements in $\rho_A(K_0(A))$.

Definition 2.4. Let \mathfrak{p} be a supernatural number. Denote by $M_{\mathfrak{p}}$ the UHF-algebra associated with \mathfrak{p} . Denote by $\mathbb{Q}_{\mathfrak{p}}$ the $K_0(M_{\mathfrak{p}})$ which is identified with a subgroup of \mathbb{Q} . Denote by Q the UHF-algebra with $K_0(Q) = \mathbb{Q}$ and $[1_Q] = 1$.

Definition 2.5. Let A be a unital simple C^* -algebra. We write TR(A) = 0 if the tracial rank of A is zero (see [29]).

Denote by \mathcal{A}_0 the class of unital simple C^* -algebras such that $TR(A \otimes U) = 0$ for some infinite dimensional UHF-algebra U. It follows from [40] that if $A \in \mathcal{A}_0$ then $TR(A \otimes U) = 0$ for all infinite dimensional simple AF-algebras U.

Denote by \mathcal{Z} the Jiang-Su algebra. It is a projectionless simple ASH-algebra with $K_0(\mathcal{Z}) = \mathbb{Z}$ and $K_1(\mathcal{Z}) = 0$. Note that \mathcal{Z} has a unique tracial state and $\mathcal{Z} \in \mathcal{A}_0$.

A unital separable C^* -algebra A is said to be \mathcal{Z} -stable if $A \otimes \mathcal{Z} \cong A$.

Let \mathfrak{p} and \mathfrak{q} be two relative prime supernatural numbers. Define

$$\mathcal{Z}_{\mathfrak{p},\mathfrak{q}} = \{ f \in C([0,1], M_{\mathfrak{p}\mathfrak{q}}) : f(0) \in M_{\mathfrak{p}} \otimes 1_{M_{\mathfrak{q}}} \text{ and } f(1) \in 1_{M_{\mathfrak{p}}} \otimes M_{\mathfrak{q}} \}.$$

By Theorem 3.4 of [49], there is a trace-collapsing unital *-embedding $\varphi : \mathcal{Z}_{\mathfrak{p},\mathfrak{q}} \to \mathcal{Z}_{\mathfrak{p},\mathfrak{q}}$. Moreover, \mathcal{Z} is the stationary C*-inductive limit $\mathcal{Z} = \lim(\mathcal{Z}_{\mathfrak{p},\mathfrak{q}},\varphi)$ (each building block is isomorphic to $\mathcal{Z}_{\mathfrak{p},\mathfrak{q}}$ and each connecting map is the same as φ).

2.6. Let A be a unital C*-algebra and let τ be a tracial state of A. Let $a \in M_n(A)_+$. Then $d_{\tau}(a) =_{df} \lim_{n \to \infty} \tau(a^{1/n})$.

For a C*-algebra A and $b, c \in A_+$, $b \leq c$ if b is Cuntz subequivalent to c, i.e., there exists a sequence $\{x_n\}$ in A such that $x_n c x_n^* \to b$.

A simple C^* -algebra A with $T(A) \neq \emptyset$ is said to have strict comparison for positive element (by traces), if, for any two positive elements $a, b \in M_n(A)$, $a \lesssim b$ whenever $d_{\tau}(a) < d_{\tau}(b)$ for all $\tau \in T(A)$.

2.7. Let C be a non-unital but σ -unital nonelementary simple C*-algebra. In [24], it is proven that there exists a unique smallest ideal J of M(C) which properly contains C. J has the form $J = \overline{J_0}$ where

$$J_0 = \{x \in M(C) : \forall a \in C_+ - \{0\}, \exists n_0 \ni (e_m - e_n) x^* x (e_m - e_n) \preceq a, \forall m > n \ge n_0\}$$

In the above, $\{e_n\}$ is an approximate identity for C. J is independent of the choice of the approximate identity $\{e_n\}$. (See [24] 2.2, Lemma 2.1 and Remark 2.9. See also [28].)

If, in addition, C has the form $C = A \otimes \mathcal{K}$ where A is unital C*-algebra with unique tracial state τ and if also C has strict comparison of positive elements, then J (as defined above) is the unique ideal of M(C) which sits properly between C and M(C). Moreover, in this case, J can be characterized by

$$J = \overline{\{x \in M(C) : \tau(x^*x) < \infty\}}.$$

(See [48]; see also [11], [23], [13], and [57].)

Definition 2.8. Retaining the notation from 2.7, recall that a C*-algebra C is said to have continuous scale if for all $x \in C_+ \setminus \{0\}$, there exists n_0 such that for all $m > n \ge n_0$, $(e_m - e_n) \le x$. ([24] Definition 2.5.) (Note that this implies that $M(C) = J_0 = J$.) If C has continuous scale and $T(C) \ne \emptyset$, then the map $\tau \to \sup\{\tau(e_n) : n \in \mathbb{N}\}$ is a continuous function. Moreover, if, in addition, C is not stable, $d_{\mu}(a) < \infty$ for all $\mu \in T(C)$.

Let A be a unital simple C^* -algebra and let $C \subset A \otimes \mathcal{K}$ be a non-unital hereditary C^* -subalgebra. Suppose that C has continuous scale. Suppose also that $T(A) \neq \emptyset$. Then

$$f(t) = \sup\{t(a) : a \in C \text{ and } 0 \le a \le 1\}$$

(for $t \in T(A)$) is a continuous affine function in Aff(T(A)). Since A is simple,

$$\inf\{f(t): t \in T(A)\} > 0.$$

For each $t \in T(A)$, define

$$\tau(c) = t(c)/f(t)$$
 for all $c \in C$.

Then τ is a normalized trace on C. Note that every $\tau \in T(C)$ has this form. This also implies that T(C) is compact.

Let C be a non-unital and σ -unital nonelementary simple C^* -algebra. Let $J \subseteq M(C)$ be the ideal as defined in 2.7 and let $a \in J_+ \setminus \{0\}$. Then for all $\delta > 0$, $C_1 = \overline{(a-\delta)_+ C(a-\delta)_+}$ has continuous scale.

3 Exponential length

The following is known.

Lemma 3.1. (cf. 10.8 of [34]) Let $\epsilon > 0$ and let $\Delta : (0,1) \to (0,1)$ be a non-decreasing map. There exist $\eta > 0$, $\delta > 0$ and a finite subset $\mathcal{G} \in C(\mathbb{T})_{s.a.}$ satisfying the following: Suppose that A is a unital separable simple C^* -algebra with TR(A) = 0, and suppose that $u, v \in U(A)$ are two unitaries such that $\operatorname{sp}(u) = \operatorname{sp}(v) = \mathbb{T}$,

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(A)$$
 (e 3.2)

and for all arcs I_a with length at least $a \ge \eta$, where $\varphi : C(\mathbb{T}) \to A$ is defined by $\varphi(f) = f(u)$ for all $f \in C(\mathbb{T})$ and

$$|\tau(g(u)) - \tau(g(v))| < \delta \text{ for all } g \in \mathcal{G}, \text{ for all } \tau \in T(A),$$

$$[u] = [v] \text{ in } K_1(A),$$

$$(e 3.3)$$

Then there exists a unitary $w \in U(A)$ such that

$$||w^*uw - v|| < \epsilon. \tag{e 3.4}$$

The following is a non-unital version of the above.

Lemma 3.2. Let $\epsilon > 0$ and let $\Delta : (0,1) \to (0,1)$ be a non-decreasing map. There exist $\eta > 0$, $\delta > 0$ and a finite subset $\mathcal{G} \in C(\mathbb{T})_{s.a.}$ satisfying the following: Suppose that A is a unital separable simple C^* -algebra with TR(A) = 0, C is a non-unital hereditary C^* -subalgebra of $A \otimes \mathcal{K}$ with continuous scale, $B = \tilde{C}$ and suppose that $u, v \in U(B)$ are two unitaries such that $\operatorname{sp}(u) = \operatorname{sp}(v) = \mathbb{T}$,

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.5)

and for all arcs I_a with length at least $a \ge \eta$, where $\varphi : C(\mathbb{T}) \to B$ is defined by $\varphi(f) = f(u)$ for all $f \in C(\mathbb{T})$ and

$$|\tau(g(u)) - \tau(g(v))| < \delta \text{ for all } g \in \mathcal{G} \text{ and } \text{ for all } \tau \in T(C),$$
 (e 3.6)

$$[u] = [v] \text{ in } K_1(B) \text{ and } \pi(u) = \pi(v),$$
 (e 3.7)

where $\pi: B \to B/C = \mathbb{C}$ is the quotient map. Then there exists a unitary $w \in U(B)$ such that

$$||w^*uw - v|| < \epsilon. \tag{e 3.8}$$

Proof. Without loss of generality, we may assume that $\pi(u) = \pi(v) = 1$. Let $\epsilon > 0$ and let $\{e_n\}$ be an approximate identity for C consisting of projections. Let $\Delta_1(a) = (1/3)\Delta(3a/4)$ for $a \in (0,1)$ and $\Delta_2 = \Delta_1/2$.

Let $\delta > 0$ and $\eta_1 > 0$ (in place of η) and a finite subset $\mathcal{G} \subset C(\mathbb{T})$ be required by Lemma 3.1 for $\epsilon/2$ and Δ_2 . Put $\eta = \eta_1/4$. Suppose that u and v are two unitaries in B satisfy the assumption for the above δ and η .

Let $\epsilon/4 > \epsilon_0 > 0$. Since C has continuous scale, there exists $N \ge 1$ and unitaries $u_1, v_1 \in e_N A e_N = e_N C e_N$ such that

$$||u - ((1 - e_N) + u_1)|| < \epsilon_0$$
 and $||v - ((1 - e_N) + v_1)|| < \epsilon_0$,

and

$$\tau(1 - e_N) < \min\{\delta/16, \Delta(\eta)/16\}$$
 (e 3.9)

for all $\tau \in T(C)$.

By choosing sufficiently small ϵ_0 , we may assume that

$$|\tau(g(u_1)) - \tau(g(v_1))| < \delta \text{ for all } \tau \in T(e_N A e_N)$$
(e 3.10)

and for all $q \in \mathcal{G}$. Moreover (by Lemma 3.4 of [35]),

$$\mu_{\tau \circ \psi}(I_a) \ge \Delta_1(a) \text{ for all } \tau \in T(C)$$
 (e 3.11)

and for all arcs I_a with length $a \ge \eta$, where $\psi(f) = f((1 - e_N) + u_1)$ for all $f \in C(\mathbb{T})$. It follows from (e 3.9) and (e 3.11) that

$$\mu_{\tau \circ v_1}(I_a) > \Delta_2(a)$$
 for all $\tau \in T(e_N A e_N)$

and for all $a \geq \eta$, where $\psi_1(f) = f(u_1)$ for all $f \in C(\mathbb{T})$. It follows from 3.1 that there exists a unitary $w_1 \in e_N A e_N$ such that

$$||u_1 - w_1^* v_1 w_1|| < \epsilon/2.$$

Let $w = (1 - e_N) + w_1$. Then

$$||u - w^*vw|| < \epsilon.$$

Proposition 3.3. Let A be a unital simple C^* -algebra, $C \subset A \otimes K$ be a non-unital hereditary C^* -subalgebra with continuous scale and let $B = \tilde{C}$. Let $u \in U(B)$ with $\operatorname{sp}(u) = \mathbb{T}$. Then there exists a nondecreasing function $\Delta : (0,1) \to (0,1)$ such that

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.12)

and for all arcs I_a with length $a \in (0,1)$, where $\varphi : C(\mathbb{T}) \to B$ defined by $\varphi(f) = f(u)$ for all $f \in C(\mathbb{T})$. Moreover, one may also require that $\lim_{a\to 0+} \Delta(a) = 0$.

Proof. Let $\pi: B \to B/C = \mathbb{C}$ be the quotient map. Let $\lambda = \pi(u)$. So $\lambda \in \mathbb{T}$. Fix $a \in (0,1)$. Consider finitely many open arcs $I_{a,1}, I_{a,2}, ..., I_{a,m} \subset \mathbb{T}$ all with radius a/2 such that $\lambda \in I_{a,1}$ and every arc I_a with length at least a contains one of $I_{a,j}$ and $\bigcup_{j=1}^m I_{a,j} = \mathbb{T}$. Choose a non-zero positive function $f_{a,j} \in C(\mathbb{T})$ such that the support of $f_{a,j}$ contained in $I_{a,j}$ and $1 \geq f_{a,j}(t) > 0$ for all $t \in I_{a,j}, j = 1, 2, 3, ..., m$. Since $f_{a,j} \in C$ for j = 2, 3, ..., m, and T(C) is compact (since C has continuous scale),

$$d_{a,j} = \inf\{\tau(f_{a,j}) : \tau \in T(C)\} > 0.$$

For $f_{a,1}$, it dominates a non-zero positive element in C. It follows that

$$d_{a,1} = \inf\{\tau(f_{a,1}) : \tau \in T(C)\} > 0.$$

Define $D(a) = \min\{a, d_{a,j} : j = 1, 2, ..., m\}$. Then define

$$\Delta(a) = \sup\{D(\eta) : 0 < \eta \le a\}.$$

Note that, for each $a \in (0,1), I_a \supset I_{b,j}$ for some j and any $b \leq a$. Therefore

$$\mu_{\tau \circ \varphi}(I_a) \ge \mu_{\tau \circ \varphi}(I_{b,j})$$

for all $\tau \in T(C)$. It follows that

$$\mu_{\tau \circ \varphi}(I_a) \geq D(b)$$
 for all $\tau \in T(C)$ and for all $a \geq b$.

Consequently

$$\mu_{\tau \circ \varphi}(I_a) \geq \Delta(a)$$
 for all $\tau \in T(C)$.

Lemma 3.4. Let A be a unital simple separable C^* -algebra with TR(A) = 0, $C \subset A \otimes K$ be a non-unital hereditary C^* -subalgebra with continuous scale and let $B = \tilde{C}$. Let $u \in U_0(B)$ with $\operatorname{sp}(u) = \mathbb{T}$ and $\pi(u) = 1$, where $\pi : B \to \mathbb{C}$ is the quotient map. Then, for any $\epsilon > 0$, there exists a selfadjoint element $h \in C$ with $\operatorname{sp}(h) = [-2\pi, 2\pi]$ such that

$$||u - \exp(ih)|| < \epsilon \text{ and } \tau(h) = 0 \text{ for all } \tau \in T(C).$$
 (e 3.13)

Moreover, we may assume that

$$\sup_{\tau \in T(C)} \lim_{n \to \infty} \tau(|h|^{1/n}) < 1.$$
 (e 3.14)

Proof. First we note that the assumption that $sp(u) = \mathbb{T}$ implies that A is infinite dimensional. Let $\epsilon > 0$. Let $\varphi : C(\mathbb{T}) \to B$ be the homomorphism defined by $\varphi(f) = f(u)$ for all $f \in C(\mathbb{T})$. It follows from 3.3 that there exists a non-decreasing function $\Delta : (0,1) \to (0,1)$ such that

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta(a)$$
 for all $\tau \in T(C)$

and for all arcs I_a with length $a \in (0,1)$. Let $\eta > 0$, $\delta > 0$ and $\mathcal{G} \subset C(\mathbb{T})_{s.a.}$ a finite subset be required by 3.2 for $\epsilon/4$ and Δ . To simplify the notation, without loss of generality, we may assume that $||g|| \leq 1$ for all $g \in \mathcal{G}$.

Let $\epsilon/4 > \epsilon_0 > 0$. Let $\{e_n\}$ be an approximate identity for C consisting of projections. We choose an integer $N \ge 1$ and unitary $u_1 \in e_N C e_N$ such that $u_1 \in U_0(e_N C e_N)$ and

$$||u - ((1 - e_N) + u_1)|| < \epsilon_0/2 \text{ and } \tau(1 - e_N) < \min\{\epsilon_0, \Delta(\eta/2)/4\}$$
 (e 3.15)

for all $\tau \in T(C)$. Since $e_N C e_N$ has real rank zero and is infinite dimensional, $u_1 \in CU(e_N C e_N)$. It follows from Theorem 4.5 of [36] that there exists a selfadjoint element $b_1 \in e_N C e_N$ with $||b_1|| \leq 2\pi$ such that

$$||u_1 - e_N \exp(ib_1)|| < \epsilon_0/2 \text{ and } \tau(b_1) = 0 \text{ for all } \tau \in T(e_N C e_N).$$
 (e 3.16)

(This can be derived directly from the fact that C has real rank zero). We assume that $e_{N+1} - e_N \neq 0$. Let $q_1, q_2 \in (e_{N+1} - e_N)C(e_{N+1} - e_N)$ be mutually orthogonal and mutually equivalent projections such that

$$|\tau(q_2)| = |\tau(q_1)| < \min\{\epsilon_0/32\pi, \Delta_1(\eta/2)/32\pi\} \text{ for all } \tau \in T(C).$$
 (e 3.17)

We choose ϵ_0 sufficiently small such that $\epsilon_0 < \delta/16\pi$ and

$$|\tau(g(u)) - \tau(g(u_1'))| < \delta/2 \text{ for all } g \in \mathcal{G}$$
 (e 3.18)

where $u_1' = (1 - e_N) + u_1$. Let $b_2 \in q_1Cq_1$ with $\operatorname{sp}(b_2) = [-2\pi, 2\pi]$. Let $z \in U(B)$ such that $z^*q_1z = q_2$. Let $b_3 = -z^*b_2z$. Note that $\tau(b_2 + b_3) = 0$ for all $\tau \in T(C)$. Define $u_2 = (1 - e_{N+1}) + (e_{N+1} - e_N - q_1 - q_2) + (q_1 + q_2) \exp(i(b_2 + b_3)) + u_1$. By (e 3.18), (e 3.17) and the fact that $||g|| \leq 1$ for all $g \in \mathcal{G}$, we estimate that

$$|\tau(g(u)) - \tau(g(u_2))| < \delta \text{ for all } g \in \mathcal{G}.$$
 (e.3.19)

It follows from 3.2 that there exists a unitary $w \in B$ such that

$$||u - w^* u_2 w|| < \epsilon/2. \tag{e 3.20}$$

Now let $h = w^*(b_1 + b_2 + b_3)w$. Then $sp(h) = [-2\pi, 2\pi]$,

$$||u - \exp(ih)|| < \epsilon$$
 and $\tau(h) = 0$ for all $\tau \in T(C)$.

Moreover (e 3.14) also holds since $h \in (e_N + q_1 + q_2)C(e_N + q_1 + q_2)$.

Corollary 3.5. Let A be a unital simple infinite dimensional separable C^* -algebra of tracial rank zero and let $C \subset A \otimes K$ be a non-unital hereditary C^* -subalgebra with continuous scale. Suppose that $u \in U(\tilde{C})$ with $\operatorname{sp}(u) = \mathbb{T}$ and suppose that $\{e_n\} \subset C$ is an approximate identity consisting of projections. Then, for any $\epsilon > 0$ and any $\sigma > 0$, there exists $k \geq 1$ and a unitary $w \in e_k Ce_k$ with $\operatorname{sp}(w) = \mathbb{T}$ such that

$$||u - \lambda(1 - e_k + w)|| < \epsilon \text{ and } \tau(1 - e_k) < \sigma \text{ for all } \tau \in T(C),$$
 (e 3.21)

where $\lambda = \pi(u)$ and $\pi : \tilde{C} \to \tilde{C}/C$ is the quotient map.

Proof. Without loss of generality, we may assume that $\pi(u) = 1$. In the above proof, let

$$w = (e_{N+1} - e_N - q_1 - q_2) + (q_1 + q_2) \exp(i(b_2 + b_3) + u_1)$$

Then, since $sp(b_2) = [-2\pi, 2\pi]$, $sp(w) = \mathbb{T}$. Let k = N + 1. The corollary then follows.

The following is a non-unital version of Corollary 3.9 of [31].

Lemma 3.6. Let $\epsilon > 0$. There exists $\delta > 0$ satisfying the following: For any unital separable simple C^* -algebra A with real rank zero and stable rank one, any hereditary C^* -subalgebra $C \subset A \otimes K$ with continuous scale, and any unitary $u \in \tilde{C}$ with $\operatorname{sp}(u) = \mathbb{T}$, if $v \in \tilde{C}$ is another unitary with [v] = 0 in $K_1(\tilde{C})$ such that

$$||[u, v]|| < \delta \text{ and } bott_1(u, v) = 0,$$
 (e 3.22)

there exists a continuous path of unitaries $\{V(t): t \in [0,1]\}$ in \tilde{C} with V(0) = v and V(1) = 1 such that

$$||[u, V(t)]|| < \epsilon \text{ for all } t \in [0, 1] \text{ and } \operatorname{length}(V(t)) \le \pi + \epsilon.$$
 (e 3.23)

Moreover, if $\pi(v) = 1$, one can require that $\pi(V(t)) = 1$ for all $t \in [0, 1]$.

Proof. Without loss of generality, we may assume that $\pi(u)=1$. Put $\tilde{C}=B$. Let $1/2>\epsilon>0$. From Corollary 3.9 of [31], one has the following statement: There exists $\delta>0$ and $\sigma>0$ satisfying the following: for any unital separable simple C^* -algebra A_0 with real rank zero and stable rank one, and unitary $u'\in A_0$ with $\mathrm{sp}(u')$ being σ -dense in $\mathbb T$ and any unitary $v'\in A_0$ with [v']=0 in $K_1(A_0)$ such that $\|[u,v']\|<\delta$ and $\mathrm{bott}_1(u',v')=0$, there exists a continuous path of unitaries $\{v'(t):t\in[0,1]\}\subset A_0$ such that v'(0)=v',v'(1)=1 and $\|[u',v'(t)]\|<\epsilon/4$ for all $t\in[0,1]$ and $\mathrm{length}(\{v'(t)\}:t\in[0,1]\}\leq\pi+\epsilon/2$. Choose such δ and σ .

Choose $\delta_1 > 0$ satisfying the following: if u' and v' are two unitaries with $\operatorname{sp}(u') = \mathbb{T}$, then $\operatorname{sp}(v')$ is $\sigma/2$ -dense in \mathbb{T} , provided that $||u'-v'|| < \delta_1$.

Let $\{e_n\}$ be an approximate identity for C consisting of projections. Choose $\theta > 0$ satisfying the following: for any unitaries w_1, w_2, w_3 , bott₁ (w_1, w_3) and bott₁ (w_2, w_3) are well defined and

$$bott_1(w_1, w_3) = bott_1(w_2, w_3)$$

provided that $||[w_i, w_3]|| < \theta$ and $||w_1 - w_2|| < \theta$.

Choose $\epsilon_1 = \min\{\epsilon/4, \delta/4, \delta_1/2, \theta\}$. Choose an integer $N \ge 1$ and unitaries $u_1, v_1 \in e_N A e_N = e_N C e_N$ such that

$$||u - ((1 - e_N) + u_1)|| < \epsilon_1, ||v - (\lambda(1 - e_N) + v_1)|| < \epsilon_1 \text{ and } ||[u_1, v_1]|| < \epsilon_1$$
 (e.3.24)

where $\lambda = \pi(v)$. Moreover, by the choice of θ ,

$$bott_1(u, v) = bott_1(u'_1, v'_1),$$
 (e 3.25)

where

$$u'_1 = (1 - e_N) + u_1$$
 and $v'_1 = \lambda(1 - e_N) + v_1$.

Note that

$$bott_1(u'_1, v'_1) = bott_1(u_1, v_1).$$
 (e 3.26)

It follows from (e 3.25), and (e 3.26)

$$bott_1(u_1, v_1) = 0.$$
 (e 3.27)

By the choice of δ_1 , we conclude that $\operatorname{sp}(u_1')$ is $\sigma/2$ -dense in \mathbb{T} . Thus $\operatorname{sp}(u_1)$ is σ -dense in \mathbb{T} . By applying the statement at the beginning of this proof (from Cor.3.9 of [31]), we obtain a continuous path of unitaries $\{w(t): t \in [1/2, 1]\} \subset e_N A e_N$ such that

$$w(1/2) = v_1, \ w(1) = e_N \text{ and } \|[u_1, w(t)]\| < \epsilon/4 \text{ for all } t \in [1/2, 1].$$
 (e 3.28)

Moreover,

$$length\{w(t): t \in [1/2, 1]\} \le \pi + \epsilon/2. \tag{e 3.29}$$

By (e 3.24), there exists $h \in B_{s.a.}$ such that

$$||h|| < 2\arcsin(\epsilon/4) \text{ and } v\exp(ih) = v_1'.$$
 (e 3.30)

Define $w_0(t) = v \exp(i2th)$ for $t \in [0, 1/2]$. Then

$$w_0(0) = v$$
, $w_0(1/2) = v'_1$ and length $\{w_0(t) : t \in [0, 1/2]\} < \epsilon/2$.

Now define $V(t) = w_0(t)$ if $t \in [0, 1/2)$ and $V(t) = \lambda(1 - e_N) + w(t)$ for $t \in [1/2, 1]$. Then V(t) is continuous and

length
$$\{V(t)\} \le \pi + \epsilon/2 + \epsilon/2 = \pi + \epsilon$$
.

One also verifies that

$$||[u, V(t)]|| < \epsilon \text{ for all } t \in [0, 1].$$
 (e 3.31)

For the very last part of the lemma, assume that $\pi(v) = 1$. Then $\lambda = 1$. By (e 3.30), $\pi(h) = 0$. It follows that $\pi(w_0(t)) = 1$ for all $t \in [0, 1/2]$. One then checks that

$$\pi(V(t)) = 1 \text{ for all } t \in [0, 1].$$

Lemma 3.7. Let C be a non-unital hereditary C^* -subalgebra of a separable C^* -algebra with stable rank one, and let $B = \tilde{C}$. Suppose that u and v are two unitaries in B such that $uv^* \in U_0(B)$ and $\pi(u) = \pi(v)$, where $\pi: B \to B/C = \mathbb{C}$ be the quotient map. Then, we can always assume that

$$R_{u,v}(t_0) = 0$$

where t_0 is the tracial state of B such that $(t_0)|_C = 0$.

Proof. We may write

$$uv^* = \prod_{j=1}^k \exp(\pi i a_j)$$

where $a_j \in B_{s.a.}$. Since $\pi(uv^*) = 1$,

$$\sum_{j=1}^{k} \pi(a_j) = 2m$$

for some integer m. Define $h_1 = a_1 - 2m$, $h_j = a_j$, j = 2, 3, ..., k. Define

$$U(t) = \prod_{j=1}^{k} \exp(\pi i (1-t)h_j) v \ t \in [0,1].$$

Then U(t) is a continuous piecewise smooth path with U(0) = u and U(1) = v. Moreover,

$$\sum_{j=1}^{k} \pi(h_j) = 0.$$

It follows that

$$\frac{1}{2\pi i} \int_0^1 t_0(\frac{dU(t)}{dt}U(t)^*)dt = 0.$$

The following is a non-unital version for a special case of Lemma 3.5 of [36].

Lemma 3.8. Let $\epsilon > 0$ and let $\Delta : (0,1) \to (0,1)$ be a non-decreasing function. There is $\delta > 0$, $\eta > 0$, $\sigma > 0$ and there is a finite subset $\mathcal{G} \subset C(\mathbb{T})_{s.a.}$ satisfying the following: For any unital separable simple C^* -algebra A with TR(A) = 0, any non-unital hereditary C^* -subalgebra $C \subset A \otimes \mathcal{K}$ with continuous scale, any pair of unitaries $u, v \in \tilde{C}$ such that $sp(u) = \mathbb{T}$ and [u] = [v] in $K_1(\tilde{C})$, $\pi(u) = \pi(v)$,

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$

for all intervals I_a with length at least η , where $\varphi: C(\mathbb{T}) \to \tilde{C}$ is the homomorphism defined by $\varphi(f) = f(u)$ for all $f \in C(\mathbb{T})$,

$$|\tau(g(u)) - \tau(g(v))| < \delta \text{ for all } \tau \in T(C)$$
 (e 3.32)

and for all $g \in \mathcal{G}$, for any $a \in Aff(T(C))$ with $a - R_{u,v}|_{T(C)} \in \rho_C(K_0(C))$ and $||a|| < \sigma$ and $y \in K_1(\tilde{C})$, there is a unitary $w \in \tilde{C}$ such that

$$[w] = y, ||u - w^*vw|| < \epsilon \text{ and}$$
 (e 3.33)

$$\frac{1}{2\pi i}\tau(\log(u^*w^*vw)) = a(\tau) \text{ for all } \tau \in T(C).$$
 (e 3.34)

Proof. Note that, since $\operatorname{sp}(u) = \mathbb{T}$, A has infinite dimension. Since TR(A) = 0, $CU(A) = U_0(A)$ and $\overline{\rho_A(K_0(A))} = \operatorname{Aff}(T(A))$. We will use these facts in the following proof. Without loss of generality, we may assume that $\pi(u) = \pi(v) = 1$. By 3.7, we may assume that

$$R_{u,v}(t_0) = 0, (e 3.35)$$

where t_0 is the tracial state of B that vanishes on C.

Let $\epsilon > 0$ and Δ be given. Choose $\epsilon/2 > \theta > 0$ such that, $\log(u_1)$, $\log(u_2)$ and $\log(u_1u_2)$ are well defined and

$$\tau(\log(u_1 u_2)) = \tau(\log(u_1)) + \tau(\log(u_2)) \tag{e 3.36}$$

for every tracial state τ and for any unitaries u_1, u_2 such that

$$||u_i - 1|| < \theta, \quad i = 1, 2.$$

We may choose even smaller θ such that

$$bott_1(u_1, v_1) = bott_1(u_2, v_1)$$
 (e 3.37)

provided that $||[u_1, v_1]|| < \theta$ and $||u_1 - u_2|| < \theta$. Let $\delta' > 0$ (in place of δ) be required by Lemma 3.1 of [36] for $\theta/2$ (in place of ϵ). Put $\sigma = \delta'/2$. Let $1/2 > \delta > 0$ and η be required by 3.2 for $\min\{\sigma, \theta/2, 1\}$ (in place of ϵ) and Δ . Suppose u and v satisfy the assumption for the above δ , η and σ . Then, by 3.2, there exists a unitary $z \in U(\tilde{C})$ such that

$$||u - z^*vz|| < \min\{\sigma, \theta/2, 1\}.$$
 (e 3.38)

Let $b = \frac{1}{2\pi i} \log(u^*z^*vz)$. Then $||b|| < \min\{\theta, \sigma, 1\}$. By Lemma 3.2 of [36], $\hat{b} - R_{u,v} \in \rho_C(K_0(C))$. Note that, by $(e \ 3.38)$, $||\log(u^*z^*vz)|| < \pi$. Since $\pi(u) = \pi(v) = 1$, $\pi(u^*z^*vz) = 1$. It follows that $\pi(\log(u^*z^*vz)) = 0$. In particular, $t_0(b) = 0$.

Let $a \in Aff(T(C))$ be such that $||a|| < \sigma$ and $a - R_{u,v}|_{T(C)} \in \rho_C(K_0(C))$ as given by the lemma. It follows that $a - \hat{b} \in \rho_C(K_0(C))$. Moreover, $||a - \hat{b}|| < 2\sigma < \delta'$. Let $p, q \in M_k(C)$ be projections such that

$$\tau(p-q) = a(\tau) - \tau(b) \text{ for all } \tau \in T(C).$$
 (e 3.39)

Let $\{e_n\}$ be an approximate identity for C consisting of projections. Choose an integer $N \ge 1$ and unitaries $u_1, v_1 \in e_N A e_N = e_N C e_N$ such that

$$||u - ((1 - e_N) + u_1)|| < \min\{\delta/4, \theta/2\}, ||v - ((1 - e_N) + v_1)|| < \min\{\delta/4, \theta/2\}$$
 (e 3.40)

and
$$\tau(1 - e_N) < \eta$$
 for all $\tau \in T(C)$. (e 3.41)

By 3.5, we may also assume that $\operatorname{sp}(u_1) = \mathbb{T}$. By (e 3.39) and (e 3.41),

$$|\tau(p-q)| < \sigma \text{ for all } \tau \in T(e_N A e_N).$$
 (e 3.42)

Note that $K_1(\tilde{C}) = K_1(C) = K_1(eNAe_N)$. It follows from Lemma 3.1 of [36] that there exists a unitary $z_1 \in e_N Ae_N$ such that

$$[z_1] = y - [z], ||[u_1, z_1]|| < \theta/2 \text{ and } bott_1(u_1, z_1)(\tau) = \tau(p - q)$$
 (e 3.43)

for all $\tau \in T(e_N A e_N)$. Put $z_2 = (1 - e_N) + z_1$, $w = z z_2$ and $u'_1 = (1 - e_N) + u_1$. It follows that

$$[w] = y$$
 and $||u - w^*vw|| < \theta < \epsilon$ (e 3.44)

$$bott_1(u'_1, z_2)(\tau) = a(\tau) - \tau(b) \text{ for all } \tau \in T(C).$$
 (e 3.45)

Put $u_0 = u((1 - e_N) + u_1^*)$ and $v_0 = v(\bar{\lambda}(1 - e_N) + v_1^*)$. So

$$||u_0 - 1|| < \min\{\delta/4, \theta/2\} \text{ and } ||v_0 - 1|| < \min\{\delta/4, \theta/2\}.$$
 (e 3.46)

By the choice of θ , we have

$$bott_1(u, z_2) = bott_1(u'_1, z_2).$$
 (e 3.47)

We compute that (using (e 3.36), (e 3.37), (e 3.47) and the Exel formula (see Lemma 3.5 of [33]))

$$\frac{1}{2\pi i}\tau(\log(u^*w^*vw)) = \frac{1}{2\pi i}\tau(\log(u^*z_2^*z^*vzz_2))$$
 (e 3.48)

$$= \frac{1}{2\pi i} \tau(\log(u^* z_2^* (uu^*) z^* vzz_2)$$
 (e 3.49)

$$= \frac{1}{2\pi i} \tau(\log((z_2 u^* z_2^* u)(u^* z^* v z))$$
 (e 3.50)

$$= \frac{1}{2\pi i} (\tau(\log(z_2 u^* z_2^* u)) + \tau(\log(u^* z^* v z)))$$
 (e 3.51)

$$= \frac{1}{2\pi i} (\tau(\log(u^* z_2^* u z_2)) + \tau(b)$$
 (e 3.52)

$$= bott_1(u, z_2)(\tau) + \tau(b)$$
 (e 3.53)

$$= a(\tau) - \tau(b) + \tau(b) = a(\tau) \quad \text{for all } \tau \in T(C).$$
 (e 3.54)

The above proof also contains the following

Lemma 3.9. (see 3.1 of [36]) Let $\epsilon > 0$. There exists $\delta > 0$ satisfying the following: Suppose that A is a unital separable simple C^* -algebra with TR(A) = 0, $C \subset A \otimes K$ is a non-unital hereditary C^* -subalgebra with continuous scale and suppose that $u \in U(\tilde{C})$ with $\operatorname{sp}(u) = \mathbb{T}$. Then, for any $x \in K_0(C)$ with $\|\rho_C(x)\| < \delta$ and any $y \in K_1(C)$, there exists a unitary $v \in A$ such that

$$[v] = y, ||[u, v]|| < \epsilon \text{ and } bott_1(u, v) = x.$$
 (e 3.55)

The next result is a special case of [51] Corollary 3.1. We provide a short proof for the convenience of the reader.

Lemma 3.10. Let A be a unital separable simple C^* -algebra which is \mathcal{Z} -stable and let $C \subset A \otimes \mathcal{K}$ be a hereditary C^* -subalgebra with continuous scale. Then $C \otimes \mathcal{Z} \cong C$.

Proof. Let $C = \overline{a(A \otimes \mathcal{K})a}$ for some positive element $a \in A \otimes \mathcal{K}$. By the assumption, we may assume that

$$a \lesssim E_N$$
,

where $E_N = \mathrm{id}_{M_N(A)}$. Since A is \mathcal{Z} -stable, A has stable rank one. We may assume that $a \in M_N(A)$. So $C \otimes \mathcal{Z}$ is a hereditary C^* -subalgebra of $M_N(A) \otimes \mathcal{Z} \cong M_N(A)$. There is an isomorphism $\varphi : M_n(A) \otimes \mathcal{Z} \to M_n(A)$ such that $\varphi(c \otimes 1_{\mathcal{Z}})$ is approximately unitarily equivalent to c for all $c \in M_n(A)$. Let $B_1 = \overline{b(A \otimes M_N)b} = C \otimes \mathcal{Z}$, where $b = a \otimes 1_{\mathcal{Z}}$. Then

$$[a] = [\varphi(b)]$$

in the Cuntz semigroup. Therefore $C \otimes \mathcal{Z} \cong C$.

Lemma 3.11. Let $\epsilon > 0$ and let $\Delta : (0,1) \to (0,1)$ be a non-decreasing map. There exist $\eta > 0$, $\delta > 0$ and a finite subset $\mathcal{G} \in C(\mathbb{T})_{s.a.}$ satisfying the following: Suppose that A is a \mathcal{Z} -stable unital separable simple C^* -algebra such that $TR(A \otimes Q) = 0$, C is a non-unital hereditary C^* -subalgebra of $A \otimes \mathcal{K}$ with continuous scale, $B = \tilde{C}$, and suppose that $u, v \in U(B)$ are two unitaries such that $\operatorname{sp}(u) = \mathbb{T}$,

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.56)

and for all arcs I_a with length at least $a \ge \eta$, where $\varphi : C(\mathbb{T}) \to B$ is defined by $\varphi(f) = f(u)$ for all $f \in C(\mathbb{T})$ and

$$|\tau(g(u)) - \tau(g(v))| < \delta \text{ for all } g \in \mathcal{G}, \text{ for all } \tau \in T(C),$$

$$[u] = [v] \text{ in } K_1(C), uv^* \in CU(B) \text{ and } \pi(u) = \pi(v),$$

$$(e 3.57)$$

where $\pi: B \to B/C = \mathbb{C}$ is the quotient map. Then there exists a unitary $w \in U(B \otimes \mathcal{Z})$ such that

$$||w^*(u \otimes 1)w - (v \otimes 1)|| < \epsilon.$$
 (e 3.58)

Proof. Without loss of generality, we may assume that $\pi(u) = \pi(v) = 1$. We first note, by [37], that $TR(A \otimes M_{\mathfrak{r}}) = 0$ for any supernatural number \mathfrak{r} . Let $\varphi : C(\mathbb{T}) \to B$ be the monomorphism defined by $\varphi(f) = f(u)$.

It follows from 3.3 that there is a non-decreasing function $\Delta:(0,1)\to(0,1)$ such that

$$\mu_{\tau \circ \varphi}(O_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.59)

for all open balls O_a of \mathbb{T} with radius $a \in (0,1)$.

Let $\epsilon > 0$. As in 3.10, we may assume that $C \subset M_n(A)$ for integer $n \geq 1$.

Let \mathfrak{p} and \mathfrak{q} be a pair of relatively prime supernatural numbers of infinite type with $\mathbb{Q}_{\mathfrak{p}} + \mathbb{Q}_{\mathfrak{q}} = \mathbb{Q}$. Denote by $M_{\mathfrak{p}}$ and $M_{\mathfrak{q}}$ the UHF-algebras associated to \mathfrak{p} and \mathfrak{q} respectively. Let $i_{\mathfrak{r}}: C \to C \otimes M_{\mathfrak{r}}$ be the embedding defined by $i_{\mathfrak{r}}(c) = c \otimes 1$ for all $a \in C$, where \mathfrak{r} is a supernatural number. Note $Tr(M_n(A) \otimes M_{\mathfrak{r}}) = 0$. Define $u_{\mathfrak{r}} = i_{\mathfrak{r}}(u)$ and $v_{\mathfrak{r}} = i_{\mathfrak{r}}(v)$. Denote $B_{\mathfrak{r}} = C \otimes M_{\mathfrak{r}}$, $\mathfrak{r} = \mathfrak{p}$, \mathfrak{q} . Denote by $\varphi_{\mathfrak{r}}: C(\mathbb{T}) \to B_{\mathfrak{r}}$ the homomorphisms defined by $\varphi_{\mathfrak{r}}(f) = f(u_{\mathfrak{r}})$ for all $f \in C(\mathbb{T})$, $\mathfrak{r} = \mathfrak{p}$, \mathfrak{q} .

Let $\delta_1 > 0$ (in place of δ) be required by 3.6 for $\epsilon/6$. Without loss of generality, we may assume that $\delta_1 < \epsilon/12$ and is small enough such that $\text{bott}_1(u_1, z_j)$ and $\text{bott}_1(u_1, w_j)$ are well defined and

$$bott_1(u_1, w_j) = bott_1(u_1, z_1) + \dots + bott_1(u_1, z_j)$$
 (e 3.60)

if u_1 is a unitary and z_j is any unitaries with $||[u_1, u_j]|| < \delta_1$, where $w_j = z_1 \cdots z_j$, j = 1, 2, 3, 4. Let $\delta_2 > 0$ (in place δ) be require by 3.9 for $\delta_1/8$ (in place of ϵ).

Furthermore, one may assume that δ_2 is sufficiently small such that for any unitaries z_1, z_2 in a C*-algebra with tracial states, $\tau(\frac{1}{2\pi i}\log(z_iz_j^*))$ (i, j = 1, 2, 3) is well defined and

$$\tau(\frac{1}{2\pi i}\log(z_1 z_2^*)) = \tau(\frac{1}{2\pi i}\log(z_1 z_3^*)) + \tau(\frac{1}{2\pi i}\log(z_3 z_2^*))$$

for any tracial state τ , whenever $||z_1 - z_3|| < \delta_2$ and $||z_2 - z_3|| < \delta_2$. We may further assume that $\delta_2 < \min\{\delta_1, \epsilon/6, 1\}$.

Let $\delta > 0$, $\eta > 0$ and $\delta_3 > 0$ (in place of σ) required by 3.8 for δ_2 (in place of ϵ).

Now assume that u and v are two unitaries which satisfy the assumption of the lemma with above δ and η .

Since $uv^* \in CU(B)$, $R_{u,v} \in \overline{\rho_B(K_0(B))}$. Since $\pi(u) = \pi(v) = 1$, we may assume that $R_{u,v}(t_0) = 0$ (see 3.7). It follows that there is $a \in \text{Aff}(T(C))$ with $||a|| < \delta_3/2$ such that $a - R_{u,v}|_{T(C)} \in \rho_C(K_0(C))$. Then the image of $a_{\mathfrak{p}} - R_{u_{\mathfrak{p}},v_{\mathfrak{p}}}$ is in $\rho_{C\otimes M_{\mathfrak{p}}}(K_0(C\otimes M_{\mathfrak{p}}))$, where $a_{\mathfrak{p}}$ is the image of a under the map induced by $\iota_{\mathfrak{p}}$. The same holds for \mathfrak{q} . Note that

$$\mu_{\tau \circ \varphi_{\tau}}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.61)

and for all a > 0 (and certainly holds for all $a \ge \eta$). By Lemma 3.8 there exist unitaries $z_{\mathfrak{p}} \in B_{\mathfrak{p}}$ and $z_{\mathfrak{q}} \in B_{\mathfrak{q}}$ such that

$$[z_{\mathfrak{r}}] = 0 \text{ in } K_1(B_{\mathfrak{r}}), \ \mathfrak{r} = \mathfrak{p}, \mathfrak{q}, \ \|u_{\mathfrak{p}} - z_{\mathfrak{p}}^* v_{\mathfrak{p}} z_{\mathfrak{p}}\| < \delta_2 \text{ and } \|u_{\mathfrak{q}} - z_{\mathfrak{q}}^* v_{\mathfrak{q}} z_{\mathfrak{q}}\| < \delta_2.$$

Moreover,

$$\tau(\frac{1}{2\pi i}\log(u_{\mathfrak{p}}^*z_{\mathfrak{p}}^*v_{\mathfrak{p}}z_{\mathfrak{p}})) = a_{\mathfrak{p}}(\tau) \text{ for all } \tau \in \mathcal{T}(C_{\mathfrak{p}}) \text{ and}$$
 (e 3.62)

$$\tau(\frac{1}{2\pi i}\log(u_{\mathfrak{q}}^*z_{\mathfrak{q}}^*v_{\mathfrak{q}}z_{\mathfrak{q}})) = a_{\mathfrak{q}}(\tau) \text{ for all } \tau \in \mathcal{T}(C_{\mathfrak{q}}).$$
 (e 3.63)

We then identify $u_{\mathfrak{p}}, u_{\mathfrak{q}}$ with $u \otimes 1$ and $z_{\mathfrak{p}}$ and $z_{\mathfrak{q}}$ with the elements in the unitization of $C \otimes M_{\mathfrak{p}} \otimes M_{\mathfrak{q}}$ which is also identified with the unitization of $C \otimes Q$. In the following computation, we also identify T(C) with $T(C_{\mathfrak{p}}), T(C_{\mathfrak{q}})$, and $T(C_{\mathfrak{p}}), \text{ or } T(C_{\mathfrak{q}})$ with $T(C \otimes Q)$ by identifying τ with $\tau \otimes t$, where t is the unique tracial state on $M_{\mathfrak{p}}$, or $M_{\mathfrak{q}}$, or Q. In particular,

$$a_{\mathfrak{p}}(\tau \otimes t) = \tau(a) \text{ for all } \tau \in T(C) \text{ and}$$
 (e 3.64)

$$a_{\mathfrak{q}}(\tau \otimes t) = \tau(a) \text{ for all } \tau \in T(C)$$
 (e 3.65)

We compute that by the Exel formula (see 3.5 of [33]),

$$(\tau \otimes t)(\operatorname{bott}_{1}(u \otimes 1, z_{\mathfrak{p}}^{*}z_{\mathfrak{q}})) = (\tau \otimes t)(\frac{1}{2\pi i}\log(z_{\mathfrak{p}}^{*}z_{\mathfrak{q}}(u^{*} \otimes 1)z_{\mathfrak{q}}^{*}z_{\mathfrak{p}}(u \otimes 1))) \qquad (e \, 3.66)$$

$$= (\tau \otimes t)(\frac{1}{2\pi i}\log(z_{\mathfrak{q}}(u^* \otimes 1)z_{\mathfrak{q}}^*z_{\mathfrak{p}}(u \otimes 1)z_{\mathfrak{p}}^*) \qquad (e 3.67)$$

$$= (\tau \otimes t)(\frac{1}{2\pi i}\log(z_{\mathfrak{q}}(u^* \otimes 1)z_{\mathfrak{q}}^*(v \otimes 1))$$
 (e 3.68)

$$+(\tau \otimes t)(\frac{1}{2\pi i}\log((v^* \otimes 1)z_{\mathfrak{p}}(u \otimes 1)z_{\mathfrak{p}}^*)$$
 (e 3.69)

$$= (\tau \otimes t)(\frac{1}{2\pi i}\log(u_{\mathfrak{q}}^* z_{\mathfrak{q}}^* v_{\mathfrak{q}} z_{\mathfrak{q}})$$
 (e 3.70)

$$+(\tau \otimes t)(\frac{1}{2\pi i}\log(v_{\mathfrak{p}}^*z_{\mathfrak{p}}u_{\mathfrak{p}}z_{\mathfrak{p}}^*)) \tag{e 3.71}$$

$$= \tau(a) - \tau(a) = 0 (e 3.72)$$

for all $\tau \in T(C)$. It follows that

$$\tau(\text{bott}_1(u \otimes 1, z_{\mathfrak{p}}^* z_{\mathfrak{q}})) = 0. \tag{e 3.73}$$

for all $\tau \in T(C \otimes Q)$.

Since the UHF-algebras D := Q, $M_{\mathfrak{p}}$ or $M_{\mathfrak{q}}$ have unique trace, the map $\rho_C \otimes \mathrm{id}_{K_0(D)}$ is the same as the map $\rho_{C \otimes D}$ if $K_0(C \otimes D)$ is identified as $K_0(C) \otimes K_0(D)$ respectively.

Let $y = \text{bott}_1(u \otimes 1, z_{\mathfrak{p}}^* z_{\mathfrak{q}}) \in \ker \rho_{C \otimes Q}$.

Since \mathbb{Q} , \mathbb{Q}_p and \mathbb{Q}_q are flat \mathbb{Z} -modules, as in the proof of 5.3 of [38],

$$\ker \rho_{C \otimes Q} = \ker \rho_C \otimes \mathbb{Q} \tag{e 3.74}$$

$$\ker \rho_{C \otimes M_{\mathfrak{r}}} = \ker \rho_C \otimes \mathbb{Q}_{\mathfrak{r}}, \quad \mathfrak{r} = \mathfrak{p} \text{ and } \mathfrak{r} = \mathfrak{q}.$$
 (e 3.75)

It follows that there are $x_1, x_2, ..., x_l \in \rho_C(K_0(C))$ and $r_1, r_2, ..., r_l \in \mathbb{Q}$ such that

$$y = \sum_{i=1}^{l} x_j \otimes r_j.$$

Since $\mathbb{Q} = \mathbb{Q}_{\mathfrak{p}} + \mathbb{Q}_{\mathfrak{q}}$, one has $r_{j,\mathfrak{p}} \in \mathbb{Q}_{\mathfrak{p}}$ and $r_{j,\mathfrak{q}} \in \mathbb{Q}_{\mathfrak{q}}$ such that $r_j = r_{j,\mathfrak{p}} - r_{j,\mathfrak{q}}$. So

$$y = \sum_{j=1}^{l} x_j \otimes r_{j,\mathfrak{p}} - \sum_{j=1}^{l} x_j \otimes r_{j,\mathfrak{q}}.$$

Put $y_{\mathfrak{p}} = \sum_{j=1}^{l} x_{j} \otimes r_{j,\mathfrak{p}} \in \ker \rho_{C \otimes M_{\mathfrak{p}}}$ and $y_{\mathfrak{q}} = \sum_{j=1}^{l} x_{j} \otimes r_{j,\mathfrak{q}} \in \ker \rho_{C \otimes M_{\mathfrak{q}}}$. It follows from 3.9 that there are unitaries $w_{\mathfrak{p}} \in B_{\mathfrak{p}}$ and $w_{\mathfrak{q}} \in B_{\mathfrak{q}}$ such that

$$[w_{\mathfrak{r}}] = 0 \text{ in } K_1(B_{\mathfrak{r}}), \ \mathfrak{r} = \mathfrak{p}, \mathfrak{q}, \ \|[u_{\mathfrak{p}}, w_{\mathfrak{p}}]\| < \delta_1/8, \ \|[u_{\mathfrak{q}}, w_{\mathfrak{q}}]\| < \delta_1/8 \text{ and}$$
 (e 3.76)

$$\operatorname{bott}_1(u_{\mathfrak{p}}, w_{\mathfrak{p}}) = y_{\mathfrak{p}} \text{ and } \operatorname{bott}_1(u_{\mathfrak{q}}, w_{\mathfrak{q}}) = y_{\mathfrak{q}}.$$
 (e 3.77)

Put $W_{\mathfrak{p}} = z_{\mathfrak{p}}^* w_{\mathfrak{p}} \in \widetilde{C \otimes M_{\mathfrak{p}}}$ and $W_{\mathfrak{q}} = z_{\mathfrak{q}}^* w_{\mathfrak{q}} \in \widetilde{C \otimes M_{\mathfrak{q}}}$. Then

$$\|u_{\mathfrak{p}} - W_{\mathfrak{p}}^* v_{\mathfrak{p}} W_{\mathfrak{p}}\| < \delta_2 + \delta_1/8 < \epsilon/6 \text{ and } \|u_{\mathfrak{q}} - W_{\mathfrak{q}}^* v_{\mathfrak{q}} W_{\mathfrak{q}}\| < \delta_2 + \delta_1/8 < \epsilon/6.$$
 (e 3.78)

Suppose that $\pi(W_{\mathfrak{p}}) = \lambda_{\mathfrak{p}}$ and $\pi(W_{\mathfrak{q}}) = \lambda_{q}$. Replacing $W_{\mathfrak{p}}$ by $\bar{\lambda}_{\mathfrak{p}}W_{\mathfrak{p}}$ and $W_{\mathfrak{q}}$ by $\bar{\lambda}_{q}W_{\mathfrak{q}}$, we may assume that $\pi(W_{\mathfrak{p}}) = \pi(W_{\mathfrak{q}}) = 1$.

Note, again, that $u_{\mathfrak{r}} = u \otimes 1$ and $v_{\mathfrak{r}} = v \otimes 1$, $\mathfrak{r} = \mathfrak{p}$, \mathfrak{q} . With identification of $W_{\mathfrak{r}}, w_{\mathfrak{r}}, z_{\mathfrak{r}}$ with unitaries in the unitization $C \otimes Q$, we also have

$$||[u \otimes 1, W_{\mathfrak{p}}^* W_{\mathfrak{q}}]|| < \delta_1/4 \text{ and}$$
 (e 3.79)

$$bott_{1}(u \otimes 1, W_{\mathfrak{p}}^{*}W_{\mathfrak{q}}) = bott_{1}(u \otimes 1, w_{\mathfrak{p}}^{*}z_{\mathfrak{p}}z_{\mathfrak{q}}^{*}w_{\mathfrak{q}})$$

$$= bott_{1}(u \otimes 1, w_{\mathfrak{p}}^{*}) + bott_{1}(u \otimes 1, z_{\mathfrak{p}}^{*}z_{\mathfrak{q}}) + bott_{1}(u \otimes 1, w_{\mathfrak{q}})(e 3.81)$$

$$= -y_{\mathfrak{p}} + (y_{\mathfrak{p}} - y_{\mathfrak{q}}) + y_{\mathfrak{q}} = 0.$$

$$(e 3.82)$$

Let $Z_0 = W_{\mathfrak{p}}^* W_{\mathfrak{q}}$. Then $Z_0 \in U_0(C \otimes Q)$ since $C \otimes Q$ has stable rank one. Note $\pi(Z_0) = \pi(W_{\mathfrak{p}}^*)\pi(W_{\mathfrak{q}}) = 1$. Then it follows from the choice of δ_1 , (e 3.61), (e 3.82) and 3.6 that there is a continuous path of unitaries $\{Z(t) : t \in [0,1]\} \subset \widetilde{C \otimes Q}$ such that $Z(0) = Z_0$ and Z(1) = 1 and

$$||[u \otimes 1, Z(t)]|| < \epsilon/6 \text{ for all } t \in [0, 1].$$
 (e 3.83)

Moreover, by 3.6, we may assume that

$$\pi(Z(t)) = 1 \text{ for all } t \in [0, 1].$$
 (e 3.84)

Define $U(t) = W_{\mathfrak{q}}Z(t)$. Then

$$U(0) = W_{\mathfrak{p}}, \ U(1) = W_{\mathfrak{q}} \text{ and } \pi(U(t)) = 1 \text{ for all } t \in [0, 1].$$
 (e 3.85)

So, in particular, $U(0) \in B_{\mathfrak{p}}$ and $U(1) \in B_{\mathfrak{q}}$. Therefore, by (e 3.85), $U \in \widetilde{C \otimes \mathcal{Z}_{\mathfrak{p},\mathfrak{q}}} \subset \widetilde{C \otimes \mathcal{Z}}$ is a unitary and, by (e 3.78) and (e 3.83),

$$||u \otimes 1 - U^*(v \otimes 1)U|| < \epsilon/3. \tag{e 3.86}$$

The following is a non-unital version of Theorem 4.6 of [36].

Theorem 3.12. Let A be a unital separable simple \mathcal{Z} -stable C^* -algebra in \mathcal{A}_0 . Let C be a non-unital hereditary C^* -subalgebra of $A \otimes \mathcal{K}$ with continuous scale and let $B = \tilde{C}$. Suppose that $u \in CU(B)$. Then, for any $\epsilon > 0$, there exists a self-adjoint element $h \in B$ with $||h|| \leq 1$ such that

$$||u - \exp(i2\pi h)|| < \epsilon. \tag{e 3.87}$$

Proof. Note that the assumption that $u \in CU(B)$ implies that $\pi(u) = 1$, where $\pi : B \to \mathbb{C}$ is the quotient map. Since C has continuous scale, without loss of generality, we may assume that $C \subset M_k(A)$ for some $k \geq 1$. To simplify notation, we may further assume, without loss of generality, that $C \subset A$. We assume that $\operatorname{sp}(u) = \mathbb{T}$, otherwise $u = \exp(ig(u))$ for some (real-valued) continuous branch of logarithm with $||g(u)|| \leq 2\pi$. Let $\epsilon > 0$. Let $\varphi : C(\mathbb{T}) \to A$ be defined by $\varphi(f) = f(u)$. It is a unital monomorphism. It follows from 3.3 that there is a non-decreasing function $\Delta_1 : (0,1) \to (0,1)$ such that

$$\mu_{\tau \circ \varphi}(I_a) \ge \Delta_1(a) \text{ for all } \tau \in T(C)$$
 (e 3.88)

for all arcs I_a of \mathbb{T} with length $a \in (0,1)$. Define $\Delta(a) = (1/3)\Delta_1(3a/4)$ for all $a \in (0,1)$.

Choose $\delta_1 > 0$ satisfying the following: If h_1, h_2 are two selfadjoint elements in any unital C^* -algebra with $||h_j|| \leq 3\pi$, j = 1, 2, such that

$$||h_1 - h_2|| < \delta_1,$$

then

$$\|\exp(ih_1) - \exp(ih_2)\| < \epsilon/4.$$

We may assume that $\delta_1 < \epsilon/4$. Note, by [37], for any supernatural number \mathfrak{p} of infinite type, $TR(A \otimes M_{\mathfrak{p}}) = 0$. Let \mathfrak{p} and \mathfrak{q} be two relatively prime supernatural numbers of infinite type. Consider $u \otimes 1$. Denote by $u_{\mathfrak{p}}$ for $u \otimes 1$ in $A \otimes M_{\mathfrak{p}}$. For any $\delta_1/2 > \epsilon_0 > 0$, by 3.4, there is a selfadjoint element $h_{\mathfrak{p}} \in C \otimes M_{\mathfrak{p}}$ with $\operatorname{sp}(h_{\mathfrak{p}}) = [-2\pi, 2\pi]$ such that

$$||u_{\mathfrak{p}} - \exp(ih_{\mathfrak{p}})|| < \epsilon_0 \text{ and } \tau(h_{\mathfrak{p}}) = 0 \text{ for all } \tau \in T(C \otimes M_{\mathfrak{p}}).$$
 (e 3.89)

Moreover, for some 1 > r > 0,

$$\lim_{n \to \infty} \tau(|h_{\mathfrak{p}}|^{1/n}) < 1 - r \text{ for all } \tau \in T(C).$$
 (e 3.90)

Let $\Gamma: C([-2\pi, 2\pi])_{s.a.} \to \text{Aff}(T(C))$ be defined by

$$\Gamma(f)(\tau) = (\tau \otimes t)(f(h_p)) \text{ for all } f \in C([-2\pi, 2\pi])_{s.a.}$$
 (e 3.91)

and for all $\tau \in T(C)$, where t is the unique tracial state on $M_{\mathfrak{p}}$.

Let (note we now assume that $C \subset A$)

$$d(\tau) = \sup\{\tau(c) : 0 \le c \le 1 \text{ and } c \in C\} \text{ for all } \tau \in T(A).$$
 (e 3.92)

Note that $\inf_{\tau \in T(A)}(d(\tau)) > 0$ since A is simple. Since C has continuous trace, $d \in Aff(T(A))$. Define $\Gamma_1 : C([-2\pi, 2\pi])_{s.a.} \to Aff(T(A))$ by

$$\Gamma_1(f)(\tau) = d(\tau)\Gamma(f)(\tau/d(\tau))$$
 for all $\tau \in T(A)$. (e 3.93)

Let $\eta > 0$, $\delta > 0$ and let \mathcal{G} be a finite subset as required by 3.11 for $\delta_1/4$ (in place of ϵ) and Δ . Note $\Delta(a) = (1/3)\Delta_1(3a/4)$ for all $a \in (0,1)$. Choose ϵ_0 sufficiently small, so the following holds (see Lemma 3.4 of [33]): For any unitary $v \in C \otimes M_{\mathfrak{p}}$, if $||u_{\mathfrak{p}} - v|| < \epsilon_0$, then

$$\mu_{\tau \circ \psi}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.94)

and for all arcs I_a with length $a \geq \eta$, where $\psi : C(\mathbb{T}) \to A \otimes M_{\mathfrak{p}}$ is the homomorphism defined by $\psi(g) = g(v)$ for all $g \in C(\mathbb{T})$, and

$$|\tau(g(u_{\mathfrak{p}})) - \tau(g(v))| < \delta \text{ for all } \tau \in T(C)$$
 (e 3.95)

and for all $g \in \mathcal{G}$. Note each $\tau \in C \otimes M_{\mathfrak{p}}$ may be written as $s \otimes t$, where $s \in T(C)$ is any tracial state and $t \in T(M_{\mathfrak{p}})$ is the unique tracial state. It follows from 3.8 of [36] that there exists a selfadjoint element $h_0 \in A$ with $\operatorname{sp}(h) = [-2\pi, 2\pi]$ such that

$$\tau(f(h_0)) = \Gamma_1(f)(\tau) = \left(\frac{\tau \otimes t}{d(\tau)}\right)(f(h_{\mathfrak{p}}))d(\tau) \text{ for all } f \in C([-2\pi, 2\pi])$$
 (e 3.96)

and for all $\tau \in T(A)$. It follows that

$$d_{\tau}(|h_0|) < d(\tau)(1-r).$$

for all $\tau \in T(A)$ (see (e 3.90)). Let $\{e_n\}$ be an approximate identity for C (e_n may not be projections). Since C has continuous scale, we may assume

$$d_{\tau}(|h_0|) < \tau(e_n)$$
 for all $\tau \in T(A)$

and for some $n \geq 1$. Note that $0 \leq e_n \leq 1$. So, in particular,

$$d_{\tau}(|h_0|) < d_{\tau}(e_n)$$
 for all $\tau \in T(C)$.

By strict comparison, we may assume that $h_0 \in e_n Ce_n$. Note that

$$\tau(h_0) = 0 \text{ for all } \tau \in T(C). \tag{e 3.97}$$

Define $v_1 = \exp(ih_0) \in A$ and denote by $\psi_1 : C(\mathbb{T}) \to A$ the homomorphism given by $\psi_1(f) = f(v_1)$ for all $f \in C(\mathbb{T})$. Note that, by (e 3.96),

$$\tau(g(v_1)) = (\tau \otimes t)g(\exp(ih_{\mathfrak{p}})) \text{ for all } \tau \in T(C)$$
 (e 3.98)

and for all $f \in C(\mathbb{T})$. By the choice of ϵ_0 , as in (e 3.94) and (e 3.95),

$$\mu_{\tau \circ \psi_1}(I_a) = \mu_{(\tau \circ \otimes t) \circ \psi_1}(I_a) \ge \Delta(a) \text{ for all } \tau \in T(C)$$
 (e 3.99)

and for all arcs I_a with length $a \geq \eta$. Moreover,

$$|\tau(g(u)) - \tau(g(v_1))| = |(\tau \otimes t)(g(u_{\mathfrak{p}})) - (\tau \otimes t)(g(\exp(ih_{\mathfrak{p}})))| < \delta$$
 (e 3.100)

for all $\tau \in T(A)$ and for all $g \in \mathcal{G}$. Note since $\tau(h_0) = 0$ for all $\tau \in T(C)$, $v_1 \in CU(C)$. It follows from 3.11 that there exists a unitary $W \in \widetilde{C} \otimes \mathcal{Z}$ such that

$$||(u \otimes 1) - W^*(v_1 \otimes 1)W|| < \delta_1/4.$$
 (e 3.101)

Denote by $i: A \to A \otimes \mathcal{Z}$ defined by $i(a) = a \otimes 1$ and define $j: A \otimes \mathcal{Z} \to A$ such that $j \circ i$ is approximately inner. Since A is separable, there is $e \in C_+$ with $0 \leq e \leq 1$ which is strictly positive in C. Let $f_c(t) = 1$ if $t \in [c, 1]$, $f_c(t) = 0$ if $t \in [0, c/2]$ and linear in [c/2, c]. So $\{f_{1/n}(e)\}$ and $\{f_{1/n}(e) \otimes 1\}$ form approximate identities for C and for $C \otimes \mathcal{Z}$, respectively. Choose 1 > c > 0 such that

$$||(f_c(e) \otimes 1)W^*(v_1 \otimes 1)W - W^*(v_1 \otimes 1)W|| < \delta_1/16\pi \text{ and}$$
 (e 3.102)

$$||W^*(v_1 \otimes 1)W(f_c(e) \otimes 1) - W^*(v_1 \otimes 1)W|| < \delta_1/16\pi$$
 (e 3.103)

Let $z \in U(A)$ such that

$$||z^*j \circ \imath(a)z - a|| < \delta_1/16\pi$$
 (e 3.104)

for $a \in \{u, v_1, h, f_c(e), e, f_{c/2}(e)\}$. Then,

$$||u - z^* j(W^*(v_1 \otimes 1)W)z|| < \delta_1/16\pi.$$
 (e 3.105)

Moreover,

$$||f_{c/2}(e)z^*j(f_c(e)\otimes 1)z - z^*j(f_c(e)\otimes 1)z|| < \delta_1/8\pi$$
 (e 3.106)

Put $H = z^*j(W^*)j(h_0 \otimes 1)j(W)z$, $H_1 = z^*j(f_c(e) \otimes 1)j(W^*)j(h_0 \otimes 1)j(W)j(f_c(e) \otimes 1)z$ and $h = f_{c/2}(e)z^*j(W^*)j(h_0 \otimes 1)j(W)zf_{c/2}(e)$. Then $||H|| \leq 2\pi$, $||H_1|| \leq 2\pi$ and $||h|| \leq 2\pi$. We note that $z^*j(W^*)j(h_0 \otimes 1)j(W)z \in A$ and C is hereditary, $h \in C$ (recall that we assumed that $C \subset A$ at the beginning of this proof). Since $v_1 = \exp(ih_0)$, by $(e \cdot 3.105)$,

$$||u - \exp(iH)|| < \delta_1/16\pi.$$
 (e 3.107)

It follows from (e 3.102) and (e 3.103) that

$$||H - H_1|| < 4\delta_1/16 = \delta_1/4 \tag{e 3.108}$$

and by (e 3.106),

$$||H_1 - h|| < 4\delta_1/8 = \delta_1/2.$$
 (e 3.109)

Therefore

$$||H - h|| < \delta_1.$$
 (e 3.110)

By the choice of δ_1 , we have

$$\|\exp(iH) - \exp(ih)\| < \epsilon/4. \tag{e 3.111}$$

It follows from (e 3.107) and (e 3.111) that

$$||u - \exp(ih)|| < \epsilon. \tag{e 3.112}$$

The point is that now $h \in C$.

4 Unitaries in a small corner

The main goal of this section is to prove a technical lemma of 4.15 which is used to compute K_1 groups of J (as in 2.7).

Key steps in the proof of Lemma 4.15 are Lemmas 4.11 and 4.12, which are attempts to generalize a well-known argument of Elliott ([11], proof of Theorem 2.4). Elliott's argument roughly says the following: If B is a nonunital UHF-algebra and u is a unitary in M(B), then u can be approximately factored into the product of two appropriate block diagonal unitaries. Each block diagonal unitary will turn out to be homotopic to a block diagonal unitary with a trivial block (i.e., a block which is a projection).

The argument of Elliott has been generalized by a number of authors (e.g., [41] Part 2, [27] Lemma 2.6 and [58] Lemma 1.6). However, unlike the case [41], J is very different from the full multiplier algebra $M(B \otimes \mathcal{K})$. Moreover, B does not have real rank zero which were cases in [11], [27] and in [58]. In fact, our main case is $B = \mathcal{Z}$ which completely lacks of projections. Nevertheless we will try to carry out Elliott's idea without projections in hand.

Let A be a C*-algebra. Recall that $A \otimes \mathcal{K}$ has has strict comparison of positive elements by traces if for all $a, b \in (A \otimes \mathcal{K})_+$, the condition that $d_{\tau}(a) < d_{\tau}(b)$ whenever $d_{\tau}(b) < \infty$ for every nonzero densely defined lower semicontinuous trace τ on $A \otimes \mathcal{K}$, implies that $a \leq b$.

Often, we will abbreviate the above by just saying that A has strict comparison of positive elements or A has strict comparison.

Proposition 4.1. Let A be a unital simple C^* -algebra which has the strict comparison (in the sense of 2.6). Suppose that A has only finitely many extremal tracial states. Then $A \otimes \mathcal{K}$ has the strict comparison in the sense of above.

Proof. Suppose that $a, b \in A \otimes \mathcal{K}_+$ such that $d_{\tau}(a) < d_{\tau}(b)$ for all tracial states $\tau \in T(A)$ for which $d_{\tau}(b) < \infty$. Without loss of generality, we may assume that $0 \le a, b \le 1$. Let $\{\tau_1, \tau_2, ..., \tau_k\}$ be the set of extremal tracial states of A. We may assume that $d_{\tau_i}(b) < \infty$ for all $i \le k_1$ and $d_{\tau_i}(b) = \infty$ if $k \ge i > k_1$. Set

$$\delta = \min\{d_{\tau_i}(b) - d_{\tau_i}(a) : 1 \le i \le k_1\}.$$
 (e 4.113)

Let $\{e_{i,j}\}$ be a system of matrix unit for \mathcal{K} . Set $e_n = \sum_{i=1}^n e_{i,i}, n = 1, 2, ...$ For any $1/2 > \epsilon > 0$, there exists an integer $n_1 \ge 1$ such that

$$||e_{n_1}ae_{n_1} - a|| < \epsilon/4. \tag{e 4.114}$$

Put $a_1 = e_{n_1} a e_{n_1}$. Then $d_{\tau}(a_1) < \infty$ for all $\tau \in T(A)$. Moreover, since $a_1 \lesssim a$,

$$d_{\tau}(a_1) < d_{\tau}(a) \text{ for all } \tau \in T(A). \tag{e 4.115}$$

There exists an integer $m_0 \ge 1$ such that

$$\tau_i(b^{1/m_0}) > d_{\tau_i}(b) - \delta/2 \text{ for all } 1 \le i \le k_1 \text{ and}$$
 (e 4.116)

$$\tau_i(b^{1/m_0}) > d_{\tau_i}(a_1) + \delta/2 \text{ for all } k \ge i > k_1.$$
 (e 4.117)

Choose an integer $n_2 \geq n_1$ such that

$$||e_{n_2}b^{1/m_0}e_{n_2} - b^{1/m_0}|| < \min\{\epsilon/2, \delta/4\}.$$
 (e 4.118)

Put $b_1 = e_{n_2} b^{1/m_0} e_{n_2}$. Then we compute that

$$d_{\tau}(b_1) \ge \tau(b_1) > d_{\tau}(a_1) \text{ for all } \tau \in T(A).$$
 (e 4.119)

By viewing both b_1 and a_1 as elements in $M_{n_2}(A)$, we obtain an element $x \in e_{n_2}(A \otimes \mathcal{K})e_{n_2}$ such that

$$||x^*e_{n_2}b^{1/m_0}e_{n_2}x - a_1|| < \epsilon/4$$
 (e 4.120)

One then finds an element $y \in A \otimes \mathcal{K}_+$ such that

$$||x^*e_{n_1}ybye_{n_1}x - a_1|| < \epsilon/2.$$
 (e 4.121)

It follows that

$$||x^*e_{n_2}ybye_{n_2}x - a|| < \epsilon.$$
 (e 4.122)

Since this holds for every $\epsilon > 0$, we conclude that $a \leq b$.

Proposition 4.2. Suppose that A is a simple, separable, stably finite C^* -algebra such that $A \otimes \mathcal{K}$ has strict comparison of positive elements by traces. Suppose, in addition, that A has stable rank one.

Suppose that $p, q \in M(A \otimes K) \setminus A \otimes K$ are two projections such that $\tau(p) = \tau(q)$ for every densely defined, lower semicontinuous trace τ .

Then $p \sim q$, i.e., there exists a partial isometry $v \in M(A \otimes K)$ such that

$$a = v^*v$$
 and $p = vv^*$.

Proof. Set $C = A \otimes \mathcal{K}$. Let $a \in pCp$ and $b \in qCq$ be two strictly positive elements (in the respective C*-algebras). Then neither are projections. By the assumption,

$$d_{\tau}(a) = d_{\tau}(b)$$

for every densely defined, lower semicontinuous trace τ .

Since C has strict comparison for positive elements and A has stable rank one, one obtains a partial isometry $v \in C^{**}$ such that $vv^* = p$, $v^*v = q$, $xv \in C$ for all $x \in \overline{Ca}$ and $vy \in C$ for all $y \in \overline{bC}$. Moreover,

$$v^* \overline{aCa} v = \overline{bCb}.$$

(See, for example, [32] Theorem 1.7.) For any $c \in C$, since $q \in M(C)$, $qc \in C$. It follows that $qc \in \overline{bC}$. Therefore

$$vc = vqc \in C$$
.

Similarly, $cv = cpv \in C$. Therefore $v \in M(C)$.

The next result is standard (see, for example, [16] Lemma 1.3.2). We sketch the argument for the convenience of the reader:

Lemma 4.3. Suppose that B is a stable C^* -algebra. Then

$$H_B \cong B$$

where H_B is the standard Hilbert B-module, and the isomorphism is (unitary) isomorphism of Hilbert B-modules.

Sketch of proof: Since B is a stable C*-algebra, let $\{W_n\}_{n=1}^{\infty}$ be a sequence of elements of M(B) such that the following hold:

- (1) $W_n^*W_n = 1$ (i.e., W_n is an isometry), for all $n \ge 1$.
- (2) $W_m^* W_n = 0$ for all $m \neq n$.
- (3) $1_{M(B)} = \sum_{n=1}^{\infty} W_n W_n^*$, where the sum converges strictly in M(B).

Let $\Phi: H_B \to B$ be given by

$$\Phi(b_1, b_2, b_3,) =_{df} \sum_{n=1}^{\infty} W_n b_n$$

for all $(b_1, b_2, b_3, ...) \in H_B$.

One can check that Φ is an (unitary) isomorphism of Hilbert B-modules.

We now fix some notation that will be used in the next proposition and its corollary. For a simple C*-algebra A, we let Ped(A) denote the Pedersen ideal of A (see [44]). Recall that Ped(A) is hereditarily closed, and Ped(A) can be characterized as the minimal dense two-sided (algebraic) ideal of A ([44], [22]). For a nonzero positive element $e \in Ped(A)$, let $T_e(A)$ denote the set of all densely defined, lower semicontinuous traces τ on A such that $\tau(e) = 1$. When endowed with the topology of pointwise convergence on Ped(A), $T_e(A)$ is a Choquet simplex. (See [50] Proposition 3.4. Note that their notation for $T_e(A)$ is different from ours, and their terminology is also slightly different.)

Of course, if A is unital then $T_1(A) = T(A)$, where T(A) is as defined in Definition 2.1.

Proposition 4.4. Suppose that A is a simple, separable, stably finite C^* -algebra, and let e be a nonzero positive element of Ped(A).

Suppose, in addition, that for every bounded strictly positive affine lower semicontinuous function $f: T_e(A) \to (0, \infty)$, there exists a non-zero $a \in (A \otimes \mathcal{K})_+$ which is not Cuntz equivalent to a projection such that $d_{\tau}(a) = f(\tau)$ for all $\tau \in T_e(A)$.

Then for every bounded, strictly positive, affine, lower semicontinuous function $f: T_e(A) \to (0, \infty)$, there exists a projection $p \in M(A \otimes \mathcal{K}) \setminus (A \otimes \mathcal{K})$ such that

$$\tau(p) = f(\tau) \text{ for all } \tau \in T_e(A).$$

Proof. Let $a \in C_+$ (= $(A \otimes \mathcal{K})_+$) be such that a is not Cuntz equivalent to a projection in C and

$$d_{\tau}(a) = f(\tau) \tag{e 4.123}$$

for all $\tau \in T_e(A)$. By the Kasparov Absorption Theorem ([20] Theorem 2) and by Lemma 4.3, let $p \in M(C) \setminus C \cong L_C(C) \setminus K_C(C)$ be a projection such that

$$pC \cong \overline{aC}$$

where the isomorphism is (unitary) isomorphism of Hilbert C-modules. (In the above, $L_C(C)$ is the C*-algebra of all adjointable operators on the Hilbert module C, and $K_C(C)$ is the C*-subalgebra of all compact adjointable operators. See, for example, [52] Chapter 15.)

Since C is separable, we can find a strictly positive element $b \in pCp$. Hence,

$$\overline{bC} \cong \overline{aC}$$
.

Hence, by [43] Proposition 4.3 (see also [32]),

$$d_{\tau}(b) = d_{\tau}(a)$$

for all $\tau \in T_e(A)$.

But we have that for all $\tau \in T_e(A)$, $f(\tau) = d_{\tau}(a)$ and $\tau(p) = d_{\tau}(b)$. Hence, for all $\tau \in T_e(A)$,

$$f(\tau) = \tau(p)$$

as required. \Box

Remark 4.5. We note that by [7] Theorem 5.5, if A is a unital simple exact finite and \mathbb{Z} -stable C*-algebra and if we take $e =_{df} 1$ then A satisfies the hypotheses of Proposition 4.4.

We can generalize Proposition 4.4 to the case where the lower semicontinuous function is unbounded or takes the value ∞ .

Corollary 4.6. Suppose that A is a simple, separable, stably finite C^* -algebra, and let e be a nonzero positive element of Ped(A).

Suppose, in addition, that A has the property that, for every bounded strictly positive affine lower semicontinuous function $f: T_e(A) \to (0, \infty)$, there exists a nonzero $a \in (A \otimes \mathcal{K})_+$ which is not Cuntz equivalent to a projection such that $d_{\tau}(a) = f(\tau)$ for all $\tau \in T_e(A)$.

(E.g., A can be unital simple exact finite and \mathcal{Z} -stable and $e = 1_A$.)

Then for every strictly positive, affine, lower semicontinuous function $f: T_e(A) \to (0, \infty]$, there exists a projection $p \in M(A \otimes \mathcal{K}) \setminus (A \otimes \mathcal{K})$ such that

$$\tau(p) = f(\tau)$$
 for all $\tau \in T_e(A)$.

Proof. Let $\{q_n\}_{n=1}^{\infty}$ be a sequence of pairwise orthogonal projections in $M(A \otimes \mathcal{K})$ such that $q_n \sim 1$ for all $n \geq 1$ and $\sum_{n=1}^{\infty} q_n = 1$, where the sum converges in the strict topology on $M(A \otimes \mathcal{K})$.

Next, f is the pointwise limit of a strictly increasing sequence of strictly positive affine continuous functions on $T_e(A)$ (see [1] Lemma 5.3; see also [14] Theorem 11.12 or [10] (Edwards' Separation Theorem); see also [50] Proposition 4.1 and Lemma 4.2). In other words, let $\{f_n\}_{n=1}^{\infty}$ be a sequence of strictly positive affine continuous functions on $T_e(A)$ with $f_n < f_{n+1}$ for all n such that $f_n \to f$ pointwise (where ∞ is allowed to be the limit at a point).

For all $n \ge 1$, let g_n be the strictly positive affine continuous function on $T_e(A)$ that is given by $g_n =_{df} f_n - f_{n-1}$ (where $f_0 =_{df} 0$). For each $n \ge 1$, apply Proposition 4.4 to get a projection $p_n \in q_n M(A \otimes \mathcal{K})q_n$ such that

$$\tau(p_n) = g_n(\tau)$$
 for all $\tau \in T_e(A)$.

Let $p =_{df} \sum_{n=1}^{\infty} p_n$. Then the sum converges strictly and p is a projection in $M(A \otimes \mathcal{K}) - (A \otimes \mathcal{K})$ such that

$$\tau(p) = f(\tau)$$
 for all $\tau \in T_e(A)$.

The next statement follows from [18]:

Theorem 4.7. Let A be a unital simple stably finite C^* -algebra with strict comparison of positive elements by traces such that T(A) has finitely many extreme points.

Suppose that $a, b \in M(A \otimes \mathcal{K})_+$ with $a \in Ideal(b)$ and $d_{\tau}(a) < d_{\tau}(b)$ for all $\tau \in T(A)$ for which $d_{\tau}(b) < \infty$.

Then $a \leq b$.

Proof. This follows immediately from [18] Theorem 5.4.

The next lemma is standard (e.g., see Lemma 2.1 in [11]).

Lemma 4.8. For every $\epsilon > 0$, there exists $\delta > 0$ such that the following holds: If C is a unital C*-algebra and $p, q \in C$ projections such that

$$pq \approx_{\delta} p$$

then there exists a unitary $u \in C$ such that

$$upu^* \le q \text{ and } ||u-1|| < \epsilon. \tag{e 4.124}$$

The next lemma is an observation of Rørdam ([47] section 4).

Lemma 4.9. Let C be a unital C^* -algebra and let $x \in C$ be a nilpotent element (i.e., $x^2 = 0$). Then

$$x \in \overline{GL(C)}$$

where the closure is in the norm topology.

NOTE: For the rest of this section, unless otherwise stated, A is a unital separable simple \mathcal{Z} -stable C*-algebra with unique tracial state τ which is the only normalized quasitrace and J is the unique proper C*-ideal of $M(A \otimes \mathcal{K})$ which properly contains $A \otimes \mathcal{K}$ (see 2.7 and 2.8).

Lemma 4.10. Let $p \in M(A \otimes K)$ be a projection, and suppose that $a \in pJp$ is a nonzero positive element such that

$$2d_{\tau}(a) < \tau(p)$$

and there exists positive $c \in pJp \setminus A \otimes \mathcal{K}$ such that $c \perp a$.

Then for every $\epsilon > 0$, there exists a projection $q \in pJp \setminus p(A \otimes \mathcal{K})p$ such that

$$d_{\tau}(a) < \tau(q) < d_{\tau}(a) + \epsilon$$
 and $qa \approx_{\epsilon} a$.

Proof. For simplicity, let us assume that $||a|| \le 1$ and $\epsilon < \min\{1, \frac{\tau(p) - 2d_{\tau}(a)}{2}\}$. Let $h: [0,1] \to [0,1]$ be the unique continuous function such that

$$h(t) = \begin{cases} 1 & t \in [\epsilon/64, 1] \\ 0 & t \in [0, \epsilon/128] \\ \text{linear on } [\epsilon/128, \epsilon/64] \end{cases}$$

Hence, $p - h(a) \in p(A \otimes \mathcal{K})p$ is a positive element such that

$$\tau(p - h(a)) = \tau(p) - \tau(h(a)) \ge \tau(p) - d_{\tau}(a) > d_{\tau}(a) + \epsilon.$$

Also, since $c \in Her(p - h(a)), p - h(a) \notin A \otimes \mathcal{K}$.

Hence, by Proposition 4.4, let $r_1 \in J \setminus A \otimes \mathcal{K}$ be a projection such that

$$d_{\tau}(a) < \tau(r_1) < d_{\tau}(a) + \epsilon < \tau(p - h(a)).$$

By Theorem 4.7,

$$r_1 \lesssim p - h(a)$$
.

It follows that there is a projection $r \in Her(p - h(a))$ such that $[r] = [r_1]$. In particular, $\tau(r) = \tau(r_1)$. Hence,

$$r \perp (a - \epsilon/32)_+$$
.

By Theorem 4.7 again, $a \leq r$. Hence, let $x \in pJp$ be an element such that

$$x^*x = (a - \epsilon/8)_+$$
 and $xx^* \le r$.

Let

$$x = v|x|$$

be the polar decomposition of x (in A^{**}). Then

$$(a - \epsilon/8)_+ = |x|^2$$
 and $v|x|^2 v^* \le r$.

Note that since $(a - \epsilon/8)_+ \perp r$, $x^2 = 0$. Hence, by Lemma 4.9,

$$x \in \overline{GL(pJp)}.$$

Hence, by Theorem 5 in [45], let $u \in pJp$ be a unitary such that

$$u(a - 3\epsilon/4)_+ u^* = v(a - 3\epsilon/4)_+ v^* \le r.$$

Hence,

$$(a - 3\epsilon/4)_{+} \le u^* r u.$$

In particular,

$$u^*rua \approx_{\epsilon} a.$$

Taking $q =_{df} u^* r u$, we are done.

Lemma 4.11. Let $p \in J \setminus (A \otimes K)$ be a projection and let $u \in pJp$ be a unitary. Suppose that $p_1 \in pJp \setminus p(A \otimes K)p$ is a projection such that

$$100\tau(p_1) < \tau(p).$$

Then for every $\epsilon > 0$ and for every projection $q \in J \setminus A \otimes K$ with $q \perp p$, there exist projections $e_1, e_2, e_3 \in J$ such that

$$p_1 = e_1 < e_2 < e_3 < p + q$$

where the inequalities are strict, and there exists a unitary $w \in (p+q)J(p+q)$ such that

$$||w - (u+q)|| < \epsilon$$
 and $we_1 w^* \le e_2 \le we_3 w^* < p+q$.

Moreover,

$$e_2 \leq p + q - e_2$$
.

Proof. By the virtue of Propositions 4.2 and 4.4, replacing q by a subprojection if necessary, we may assume that

$$\tau(q) < (\tau(p) - \tau(p_1))/10^{10000}$$
.

By Theorem 4.7, Proposition 4.4 and Proposition 4.2, we can decompose q into a direct sum of non-zero projections

$$q = q' + q'' + q'''$$

such that $q', q'', q''' \in J \setminus (A \otimes \mathcal{K})$.

Let $\delta_5 > 0$ (in place of δ) associated with $\epsilon/100$ (in place of ϵ) be given by Lemma 4.8. Let $\delta_{j-1} > 0$ (in place of δ) associated with $\delta_j/100$ (in place of ϵ) given by Lemma 4.8, j = 5, 4, 3, 2, 1. We may assume that $\delta_5 < \epsilon/100$, and that for all j, $\delta_{j-1} < \delta_j/100$.

Take $e_1 =_{df} p_1$.

By hypothesis,

$$\tau(e_1) < \tau(p)/100.$$

Hence,

$$e_1 + ue_1u^* \in pJp \text{ and } d_{\tau}(e_1 + ue_1u^*) < \tau(p)/50.$$

It follows from Lemma 4.10 that there exists a projection $q_1 \in (p+q')J(p+q')$ such that

$$d_{\tau}(e_1 + ue_1u^*) < \tau(q_1) < d_{\tau}(e_1 + ue_1u^*) + \min\{\delta_1, \tau(p)\}/10^{100},$$
 (e 4.125)

$$q_1e_1 \approx_{\delta_1} e_1$$
 and $q_1ue_1u^* \approx_{\delta_1} ue_1u^*$.

By Lemma 4.8 and the definition of δ_1 , we can find a projection $e_2 \in (p+q')J(p+q')$ such that

$$||e_2 - q_1|| < \delta_2/50$$

(and hence $e_2 \sim q_1$) and

$$e_1 \leq e_2$$
.

As a consequence, (since $\delta_1 < \delta_2/100$),

$$e_2(ue_1u^*) \approx_{\delta_2} ue_1u^*$$
.

It follows from Lemma 4.8 and the definition of δ_2 that there is a unitary $w_1 \in (p+q')J(p+q')$ such that

$$||w_1 - (u + q')|| < \delta_3/50$$

and

$$w_1 e_1 w_1^* \le e_2.$$

Since

$$d_{\tau}(e_2 + w_1^* e_2 w_1) \le 2(\tau(p)/50 + \min\{\delta_1, \tau(p)\}/10^{100}) \le \tau(p)/25 + 2\tau(p)/10^{100},$$

and by Lemma 4.10, let $q_2 \in (p + q' + q'')J(p + q' + q'')$ be a projection such that

$$d_{\tau}(e_2 + w_1^* e_2 w_1) < \tau(q_2) < d_{\tau}(e_2 + w_1^* e_2 w_1) + \min\{\delta_2, \tau(p)\}/10^{100},$$

$$q_2e_2 \approx_{\delta_2/100} e_2$$

and

$$q_2 w_1^* e_2 w_1 \approx_{\delta_2/100} w_1^* e_2 w_1.$$

By Lemma 4.8 and the definition of δ_2 , there exists a projection $e_3 \in (p+q'+q'')J(p+q'+q'')$ such that

$$||e_3 - q_2|| < \delta_3/50$$

(and hence $e_3 \sim q_2$), and

$$e_2 \leq e_3$$
.

Hence,

$$e_3(w_1^*e_2w_1) \approx_{\delta_3} w_1^*e_2w_1.$$

Hence,

$$(w_1 + q'')e_3(w_1 + q'')^*e_2 \approx_{\delta_3} e_2.$$

Note that since

$$e_1 \le e_2 \le e_3,$$

$$(w_1 + q'')e_1(w_1 + q'')^* \le (w_1 + q'')e_3(w_1 + q'')^*.$$

Also, by our construction of e_2 ,

$$(w_1 + q'')e_1(w_1 + q'')^* \le e_2.$$

Hence,

$$(w_1+q'')(e_3-e_1)(w_1+q'')^*(e_2-(w_1+q'')e_1(w_1+q'')^*) \approx_{\delta_3} (e_2-(w_1+q'')e_1(w_1+q'')^*).$$

Hence, by Lemma 4.8 and by the definition of δ_3 , there exists a unitary $w_2' \in (p + q' + q'' - w_1e_1w_1^*)J(p + q' + q'' - w_1e_1w_1^*)$ with

$$||w_2' - (p + q' + q'' - w_1 e_1 w_1^*)|| < \delta_4/100$$

and

$$e_2 - w_1 e_1 w_1^* \le w_2' (w_1 + q'') (e_3 - e_1) (w_1 + q'')^* w_2'^*.$$

Hence, if we take

$$w =_{df} (w_2' + w_1 e_1 w_1^*)(w_1 + q'') + q'''$$

then $w \in (p+q)J(p+q)$ is a unitary,

$$||w - (u+q)|| < \epsilon, \ p_1 = e_1 < e_2 < e_3 < p,$$

and $we_1 w^* < e_2 < we_3 w^* < p$ (e 4.126)

as required. Finally, since $q_1 \sim e_2$, by (e 4.125) and by applying Theorem 4.7, we also have

$$e_2 \leq p + q - e_2$$
.

Lemma 4.12. Let $p \in J \setminus (A \otimes K)$ be a projection and $u \in pJp$ be a unitary. Let $1/2^{10} > r > 0$. Then for every projection $q \in J \setminus A \otimes K$ with $q \perp p$, there is a (norm-) continuous path of unitaries $\{u(t): t \in [0,1]\}$ in U((p+q)J(p+q)) such that

$$u(0) = u + q$$
, $u(1) = ((p + q - f_2) + u_1)((p + q - f_1) + v_1)$

where $v_1 \in U(f_1Jf_1)$, $u_1 \in U(f_2Jf_2)$, $\tau(p+q-f_j) > r\tau(p)$ for and $f_j \in (p+q)J(p+q) \setminus A \otimes K$ is a projection, j = 1, 2.

Proof. By Proposition 4.4 and Theorem 4.7, there is a projection $p_1 \in pJp \setminus p(A \otimes \mathcal{K})p$ such that $\tau(p_1) > r\tau(p)$ and $100\tau(p_1) < \tau(p)$ (equivalently, $\beta < \tau(p_1) < \tau(p)/100$ for some $\beta < \tau(p)/2^{10}$.) Since $q \in J \setminus (A \otimes \mathcal{K})$ is a projection, by Propositions 4.4 and 4.2, we can decompose q into orthogonal projections

$$q = q' \oplus q''$$

where $q', q'' \in J \setminus (A \otimes \mathcal{K})$.

It follows from Lemma 4.11 that there exists a unitary $w_0 \in (p+q')J(p+q')$ with

$$||w_0 - (u + q')|| < \epsilon/10000$$

(which implies that w_0 is path-connected to u+q' in (p+q')J(p+q')) and there are projections $e_1, e_2, e_3 \in (p+q')J(p+q') \setminus (A \otimes \mathcal{K})$ such that

$$p_1 = e_1 < e_2 < e_3 < p + q' < p + q$$

(where the inequalities are strict), and

$$w_0 e_1 w_0^* < e_2 < w_0 e_3 w_0^* < p + q' < p + q.$$

Moreover,

$$e_2 \leq p + q - e_2.$$
 (e 4.127)

It follows that

$$e_1 \leq p + q - e_1.$$
 (e 4.128)

Define $w = w_0 + q''$. Hence, we have that

$$||w - (u+q)|| < \epsilon/10000,$$
 (e 4.129)

$$p_1 = e_1 < e_2 < e_3 < e_4 =_{df} p + q,$$
 (e 4.130)

$$we_1w^* < e_2 < we_3w^* < p+q,$$
 (e4.131)

and there exists a positive $c \in (p+q)J(p+q) - (A \otimes \mathcal{K})$ such that

$$c \perp e_3 + we_3w^*$$
.

From the above and by Proposition 4.2,

$$e_1 \sim we_1 w^*$$
, $e_2 - e_1 \sim e_2 - we_1 w^*$,
 $e_3 - e_2 \sim we_3 w^* - e_2$ and $p + q - e_3 \sim p + q - we_3 w^*$.

Recall that $e_4 =_{df} p + q$. Put $e_0 =_{df} 0$. Hence, for j = 1, 2, there is a unitary $w_{2j} \in (e_{2j} - e_{2j-2})J(e_{2j} - e_{2j-2})$ such that

$$w_{2j}(e_{2j-1}-e_{2j-2})w_{2j}^*=we_{2j-1}w^*-e_{2j-2}.$$

Define

$$z =_{df} w_2 \oplus w_4$$
 and $y =_{df} z^* w$,

two unitaries in (p+q)J(p+q). Then w=zy. Clearly, by definition,

$$(e_{2j} - e_{2j-2})z = w_{2j} = z(e_{2j} - e_{2j-2})$$
 and
 $\tau(e_2 - e_0) = \tau(e_2) \ge \tau(e_1) > r\tau(p).$

Also, since

$$ze_1z^* = we_1w^*,$$

 $e_1 = z^*we_1w^*z.$

In other words,

$$e_1 = ye_1y^*$$
 or $e_1y = ye_1$.

Hence,

$$y = y_1 \oplus y_3$$

where $y_1 \in e_1 J e_1$ is a unitary and $y_3 \in (p+q-e_1)J(p+q-e_1)$ is a unitary. Moreover,

$$\tau(e_2) > \tau(e_1) = \tau(p_1) > r\tau(p).$$
 (e 4.132)

By (e 4.127) and (e 4.128), there are partial isometries $Z_1, Z_2 \in (p+q)J(p+q)$ such that

$$Z_i^* Z_i = e_i, \ Z_i Z_i^* =_{df} e_i' \le p + q - e_i, \ i = 1, 2.$$

Define a unitary $U_i \in (p+q)J(p+q)$ by

$$U_i = Z_i + Z_i^* + (p + q - e_i - e_i'), i = 1, 2.$$

We may write

$$w = zy = (w_2 \oplus U_2 w_2^* U_2^* U_2 w_2 U_2^* w_4)(y_1 \oplus U_1 y_1^* U_1^* U_1 y_1 U_1^* y_3).$$
 (e 4.133)

(In the above, we are simplifying notation by using " $U_2w_2U_2^*$ " to denote $U_2w_2U_2^* + (p+q-e_2-e_2)$, a unitary in $(p+q-e_2)J(p+q-e_2)$. Similarly, we are using " $U_1y_1U_1^*$ " to denote $U_1y_1U_1^* + (p+q-e_1-e_1')$.)

Put $f_i = e_i$, i = 1, 2. So $f_i \in (p+q)J(p+q) \setminus A \otimes \mathcal{K}$ are projections.

$$u_1 = U_2 w_2 U_2^* w_4 \in (p+q-f_2) J(p+q-f_2)$$
 and $v_1 = U_1 y_1 U_1^* y_3 \in (p+q-f_1) J(p+q-f_1)$.

Then, by (e 4.133) and (e 4.129), u + q is connected to

$$(e_2+u_1)(e_1+v_1)$$

via a norm continuous path in U(p+q)J(p+q)). Moreover, we compute that, by (e 4.132), (e 4.127) and (e 4.128),

$$\tau(p+q-f_i) \ge \tau(e_i) > r\tau(p), \ i = 1, 2.$$

Lemma 4.13. Let $p \in J \setminus (A \otimes K)$ be a projection and $u \in pJp$ be a unitary. Let 1 > r > 0. Then for every projection $q \in J \setminus A \otimes K$ with $q \perp p$, there is an integer $m \geq 1$ and a (norm-) continuous path of unitaries $\{u(t) : t \in [0,1]\}$ in U((p+q)J(p+q)) such that

$$u(0) = u + q, \ u(1) = ((p + q - f_1) + u_1)((p + q - f_2) + u_2)...((p + q - f_m) + u_m)$$

where $u_j \in U(f_jJf_j)$, $\tau(p+q-f_j) > r\tau(p)$, and $f_j \in (p+q)J(p+q) \setminus A \otimes \mathcal{K}$ is a projection, j = 1, 2, ..., m.

Proof. There is an integer $n \ge 1$ such that $1 - (1 - s)^n > r$, where $s = (3/4)2^{-10}$. By Theorem 4.7, Proposition 4.2 and Proposition 4.4, there are mutually orthogonal projections $q_l \in qJq \setminus A \otimes \mathcal{K}$, l = 1, 2, ..., n, such that $q = \sum_{l=1}^{n} q_l$.

It suffices to prove the following statement:

There are projections

$$f_1, f_2, ..., f_{2^k} \in (p + \sum_{j=1}^k q_j)J(p + \sum_{j=1}^k q_j) \setminus A \otimes \mathcal{K},$$
 (e 4.134)

unitaries
$$u_i \in U(f_j J f_j)$$
 such that $\tau(f_j) < \tau(\sum_{j=1}^k q_j) + (1-s)^k \tau(p)$, (e 4.135)

j=1,2,..,k, and u is connected to $\prod_{j=1}^{2^k}((p+q-f_j)+u_j)$ by a norm continuous path of unitaries in (p+q)J(p+q) for all $1 \le k \le n$.

We prove this by induction on k.

The case k=1 follows immediately from Lemma 4.12 with q_1 in place of q and s in place of r.

Suppose that we have proven the statement for k. We now prove the statement for k+1.

Now for $1 \leq j \leq k$, apply Lemma 4.12 to u_j (in place of u), f_j (in place of p), s (in place of r) and q_{k+1} (in place of q) to get projections $f_{j,l} \in (f_j + q_{k+1})J(f_j + q_{k+1}) \setminus A \otimes K$ (l = 1, 2) and unitaries $u_{j,l} \in f_{j,l}Jf_{j,l}$ such that $u_j + q_{k+1}$ is connected to $(f_j + q_{k+1} - f_{j,1} + u_{j,1})(f_j + q_{k+1} - f_{j,2} + u_{j,2})$ by a norm continuous path of unitaries in $(f_j + q_{k+1})J(f_j + q_{k+1})$ and

$$\tau(f_j + q_{k+1} - f_{j,l}) > s\tau(f_j).$$
 (e 4.136)

From this and the induction hypothesis, it follows that

$$\tau(f_{j,l}) < \tau(f_j) + \tau(q_{k+1}) - s\tau(f_j)$$
(e 4.137)

$$= \tau(q_{k+1}) + (1-s)\tau(f_j)$$
 (e 4.138)

$$<\sum_{j=1}^{k+1} \tau(q_j) + (1-s)^{k+1} \tau(p).$$
 (e 4.139)

The statement above holds for k+1 by renaming the $f_{j,l}s$ and $u_{j,l}s$ which completes the induction.

Lemma 4.14. Let $p, q, r, s \in J \setminus A \otimes K$ be projections such that

$$q, r, s \le p$$

$$\{q, r\} \perp s \tag{e 4.140}$$

and
$$\tau(q) \le \tau(s) \le \tau(r)$$
. (e 4.141)

Suppose that $u \in qJq$ is a unitary. Then there is a continuous path of unitaries $\{u(t) : t \in [0,1]\} \subset pJp$ such that

$$u(0) = u + (p - q)$$
 and $u(1) = v + (p - r)$,

where where $v \in rJr$ is a unitary.

Proof. Since $s \perp q$ and $q \leq s$, we may view u + (p - q) as $u + (p - q) = diag(u, s, p - q - s) = diag(u, u^*u, p - q - s)$ (with coordinates in (q, s, p - q - s)) which is norm path connected to diag(q, u, p - q - s), which has the form w + (p - s) for some unitary $w \in sJs$.

But since $s \perp r$ and $s \leq r$, w + (p - s) can be viewed as $w + (p - s) = diag(w, r, p - s - r) = diag(w, w^*w, p - s - r)$ (with coordinates in (s, r, p - s - r)) which is norm path connected to diag(s, w, p - s - r), which has the form v + (p - r) for some unitary $v \in rJr$.

Lemma 4.15. Let $p \in J \setminus A \otimes K$ be a projection and let $q \in J \setminus A \otimes K$ be another projection with $p \perp q$ and $q \leq p$.

Let $u \in pJp$ be a unitary. Then there exists a continuous path of unitaries $\{u(t) : t \in [0,1]\} \subset U((p+q)J(p+q))$ such that

$$u(0) = u + q \text{ and } u(1) = p + v,$$
 (e 4.142)

where $v \in U(qJq)$.

Proof. The result of the lemma follows immediately from Lemma 4.13 and Lemma 4.14. By Proposition 4.4, Proposition 4.2 and Theorem 4.7, we can decompose q into pairwise orthogonal projections

$$q = q' \oplus q'' \oplus q'''$$

where $q', q'', q''' \in J \setminus (A \otimes \mathcal{K})$ and $\tau(q') < \tau(q'') < \tau(q''')$, which implies that $q' \leq q'' \leq q'''$.

By applying Lemma 4.13 and by passing to a unitary which connects with u + q' by a continuous path of unitaries in U((p+q)J(p+q)), without loss of generality, we may write

$$u + q' = \prod_{j=1}^{m} ((p + q' - f_j) + u_j)$$
 (e 4.143)

such that $u_j \in U(f_j J f_j)$, where $f_j \leq p + q'$ and $f_j \leq q''$, j = 1, 2, ..., m. Note that

$$f_i, q'', q''' \le p + q$$
, $f_i \perp q'', f_j \perp q''', q''' \perp q''$ and $f_i \le q'' \le q'''$, $j = 1, 2, ..., m$.

Thus, by Lemma 4.14, for each j, there is a continuous path of unitaries $\{w_j(t): t \in [0,1]\} \subset (p+q)J(p+q)$ such that

$$w_j(0) = (p + q' + q'' + q''' - f_j) + u_j \text{ and } w_j(1) = v_j + (p + q' + q''),$$
 (e 4.144)

where $v_j \in U(q'''Jq'''), j = 1, 2, ..., m$. Define

$$u(t) = \prod_{j=1}^{m} w_j(t)$$
 for all $t \in [0, 1]$.

Then $\{u(t): t \in [0,1]\}$ is a continuous path of unitaries in (p+q)J(p+q). Moreover,

$$u(0) = u + q$$
 and $u(1) = v + p$,

where $v = \prod_{i=1}^{m} (v_i + q' + q'') \in U(qJq)$.

5 The main results

Lemma 5.1. Let A be a unital C^* -algebra and $B = A \otimes K$. Then M(B) contains a sequence of mutually orthogonal projections $\{p_n\}$ each of which is equivalent to the identity $1_{M(B)}$, $p_n \notin B$, n = 1, 2, ..., and $\sum_{n=1}^{\infty} p_n$ converges strictly to $1_{M(B)}$.

Proof. This is known. One can have a partition of \mathbb{N} into a sequence $\{I_n\}$ of mutually disjoint infinite subsets such that $1 \in I_n$ and $I_n \cap \{1, 2, ..., n-1\} = \emptyset$. Let $\{e_{i,j}\}$ be a system of matrix unit for \mathcal{K} . We identify e_{ii} with $1_A \otimes e_{ii}$. Define

$$p_n = \sum_{j \in I_n} e_{jj}.$$

The convergence is in the strict topology. Let $e_n = \sum_{i=1}^n e_{ii}$. Then $\{e_n\}$ forms an approximate identity for B. Note that, by the construction,

$$p_m e_n = 0$$
 for all $m > n$.

It follows that $\sum_{k=1}^{n} p_k$ converges strictly to $1_{M(B)}$.

The following holds in a much more general situation.

Lemma 5.2. Let A be a unital separable simple \mathcal{Z} -stable C^* -algebra with unique tracial state τ , and let $B = A \otimes \mathcal{K}$. Let J be the smallest ideal of M(B) which properly contains B.

Then, for any sequence of positive numbers $\{\alpha_n\}$ with $\alpha_n \geq \alpha_{n+1}$ and $\sum_{n=1}^{\infty} \alpha_n < \infty$, there exists a sequence of mutually orthogonal projections $\{q_n\} \in M(B) - B$ such that $q_{n+1} \lesssim q_n$, $\tau(q_n) = \alpha_n$, n = 1, 2, ..., such that $\sum_{n=1}^{\infty} q_n$ converges in the strict topology and $p = \sum_{n=1}^{\infty} q_n \in J$.

Proof. Let $\{p_n\}$ be as in 5.1.

By Proposition 4.4 and since $p_n \sim 1_{M(B)}$, let $q_n \in J - B$ be such that $q_n \leq p_n$ and $\tau(q_n) = \alpha_n$ for all $n \geq 1$. Then $\sum_n q_n$ converges strictly to an element of J.

Corollary 5.3. Let A be a unital separable simple \mathcal{Z} -stable C^* -algebra with unique tracial state τ , let $B = A \otimes \mathcal{K}$, let J be the smallest ideal of M(B) which properly contains B, and let $p \in M(B)$ be a projection such that $\tau(p) = \infty$. Then, for any sequence of positive numbers $\{\alpha_n\}$ with $\alpha_n \geq \alpha_{n+1}$ and $\sum_{n=1}^{\infty} \alpha_n < \infty$, there exists a sequence of mutually orthogonal projections $\{q_n\} \in M(B) - B$ such that $q_{n+1} \lesssim q_n$, $\tau(q_n) = \alpha_n$, n = 1, 2, ..., such that $\sum_{n=1}^{\infty} q_n$ converges in the strict topology to a projection $q \in J$ such that $q \leq p$.

Proof. This follows from 5.2 and the fact that $p \sim 1_{M(B)}$.

Lemma 5.4. Let B be a non-unital and σ -unital simple C^* -algebra, $J \subset M(B)$ be an ideal containing B such that J/B is purely infinite and simple and let $p \in J$ be a non-zero projection. Suppose that $u \in pM(B)p$ is a unitary such that [u] = 0 in $K_1(pM(B)p)$. Then, for any $\epsilon > 0$, there are two selfadjoint elements $H_1, H_2, \in pM(B)p$ and a unitary $w \in p + pBp$ such that $||H_1|| \leq \pi$, $||H_2|| < \epsilon$ and

$$u = w \exp(iH_1) \exp(iH_2).$$
 (e 5.145)

In particular, $w^*u \in U_0(pM(B)p)$.

Proof. Note that pM(B)p/pBp is purely infinite and simple. Let $\pi: pM(B)p \to pM(B)p/pBp$ be the quotient map. Suppose that u is as in the lemma. Then $\pi(u) \in U_0(pJp/pBp)$. Note that pJp/pBp is a unital purely infinite simple C^* -algebra. It follows from a result of Chris Phillips ([46]) that, for any $\epsilon > 0$, there are selfadjoint elements $h_1, h_2 \in pM(B)p/pBp$ such that

$$\pi(u) = \exp(ih_1) \exp(ih_2)$$
 and $||h_1|| \le \pi$ and $||h_2|| < \epsilon$. (e 5.146)

There are selfadjoint elements $H_1, H_2 \in pM(B)p$ such that $||H_1|| \le \pi$, $||H_2|| \le \epsilon$ and $\pi(H_j) = h_j$, j = 1, 2. Let $u_j = \exp(iH_j)$, j = 1, 2. Then

$$w = uu_2^* u_1^* \in p + pBp. \tag{e 5.147}$$

It follows that $u = w \exp(iH_1) \exp(H_2)$ and $w^*u \in U_0(pM(B)p)$.

Lemma 5.5. Let $A \in \mathcal{A}_0$ be a unital separable simple \mathcal{Z} -stable C^* -algebra with unique tracial state τ , let $B = A \otimes \mathcal{K}$ and let $J \subset M(B)$ be an ideal containing B such that J/B is purely infinite and simple. Suppose that $p \in J$ and $u \in pM(B)p \setminus B$ be a unitary. Then, for any projection $q \in J \setminus B$ and pq = 0 = qp, any $\epsilon > 0$, there is a unitary $v \in qM(B)q$ and selfadjoint elements $h_1, h_2 \in (p+q)M(B)(p+q)$ and $h_3 \in \mathbb{C}(p+q) + (p+q)B(p+q)$ such that

$$||h_1|| < \epsilon/2, ||h_2|| \le \pi, ||h_3|| \le 2\pi,$$
 (e 5.148)

$$||(u+v) - (p+q)\exp(ih_1)\exp(ih_2)\exp(ih_3)|| < \epsilon \text{ and}$$
 (e 5.149)

$$(u+v) - (p+q)\exp(ih_1)\exp(ih_2)\exp(ih_3) \in (p+q)B(p+q).$$
 (e 5.150)

Proof. There is, by 4.6 and 4.7, a projection $q_1 \in J \setminus B$ such that $q_1 \leq q$ and there is a unitary $z \in (p+q)M(B)(p+q)$ such that

$$z^*q_1z < p$$
.

It follows from Lemma 4.15 that there is a unitary $v_1 \in q_1Jq_1$ such that

$$u + v_1 + (q - q_1) \in U_0((p+q)J(p+q)).$$
 (e 5.151)

Put P = p + q. It follows from 5.4 that there are selfadjoint elements $h_1, h_2 \in PJP$ with $||h_1|| < \epsilon/2$ and $||h_2|| \le \pi$ such that

$$u + v_1 + (q - q_1) = \exp(ih_1)\exp(ih_2)w,$$

where $w \in P + PBP$. Since B is simple and has stable rank one, there exists $w_1 \in q + qBq$ such that $[w_1] = -[w]$. Thus $w(w_1 + p) \in U_0(P + PBP)$. Let

$$w(w_1 + p) = \prod_{k=1}^{m} \exp(ia_k),$$

where $a_k \in (\mathbb{C}P + PBP)_{s.a.}$. Let $\pi(a_k) = \alpha_k$, k = 1, 2, ..., m. Since $w(w_1 + p) \in P + PBP$, $\sum_{k=1}^{m} \alpha_k = 2N\pi$ for some integer N. By replacing a_k by $a_k - \alpha_k P$, we may assume that $\pi(a_k) = 0$, k = 1, 2, ..., m. In other words, $a_k \in PBP$, k = 1, 2, ..., m. Choose $b \in (qBq)_{s.a.}$ such that

$$\tau(b) = \sum_{k=1}^{m} \tau(a_k).$$
 (e 5.152)

Set

$$v_2 = q \exp(-ib). \tag{e 5.153}$$

Now let $v = (v_1 + (q - q_1))w_1v_2$ Then

$$u + v = \exp(ih_1) \exp(ih_2) w(w_1 + p)(v_2 + p).$$

Note that

$$w(w_1 + p)(v_2 + p) = P \prod_{k=1}^{m} \exp(ia_k) \exp(-ib).$$

But

$$\tau(-b) + \sum_{k=1}^{m} \tau(a_k) = 0.$$

It follows that

$$w(w_1 + p)(v_2 + p) \in CU(\mathbb{C}P + PBP).$$

By 3.12, there is $h_3 \in PBP_{s.a.}$ with $||h_3|| \leq 2\pi$ such that

$$||w(w_1+p)(v_2+p) - P\exp(ih_3)|| < \epsilon.$$

It follows that

$$||(u+v) - P\exp(ih_1)\exp(ih_2)\exp(ih_3)|| < \epsilon.$$

Lemma 5.6. Let A be a unital separable simple \mathcal{Z} -stable C^* -algebra with unique tracial state τ and let J be the smallest ideal in $M(A \otimes \mathcal{K})$ that properly contains $A \otimes \mathcal{K}$. Then, for any finite subset $\mathcal{F} \subset J$, any projection $e \in J$ and any $\epsilon > 0$, there exists a projection $p \in J$ such that

$$e \le p \text{ and } ||xp - x|| < \epsilon \text{ for all } x \in \mathcal{F}.$$
 (e 5.154)

Proof. Let $1/2 > \epsilon > 0$, finite subset $\mathcal{F} \subset J$ and a projection $e \in J$ be given. We may assume, without loss of generality, that $||x|| \le 1$ for all $x \in \mathcal{F}$. Let $\{e_{\alpha}\}$ be an approximate identity for J. There exists α such that

$$||xe_{\alpha} - x|| < \epsilon/64 \text{ and } ||ee_{\alpha} - e|| < \epsilon/64.$$
 (e 5.155)

We may assume that $||e_{\alpha}|| > 1/2$. Put $b = (e_{\alpha} - \epsilon/64)_{+}$. Let

$$f_{\epsilon/65,\epsilon/64}(t) =_{df} \begin{cases} 1 & t \in [\epsilon/64,1] \\ 0 & t \in [0,\epsilon/65] \\ \text{linear on } [\epsilon/65,\epsilon/64]. \end{cases}$$

Since $e_{\alpha} \in J$, $1_{M(A \otimes \mathcal{K})} - f_{\epsilon/65,\epsilon/64}(e_{\alpha})$ is a full element of $M(A \otimes \mathcal{K})$, and $d_{\tau}(1_{M(A \otimes \mathcal{K})} - f_{\epsilon/65,\epsilon/64}(e_{\alpha})) = \infty$. Hence, by Theorem 4.7, there is a positive element $c \notin A \otimes \mathcal{K}$, $c \perp b$. Recall also, by 2.8, that $d_{\tau}(b) < \infty$. It follows from Lemma 4.10 (with $p = 1_{M(A \otimes \mathcal{K})}$) that there is a projection $q \in J$ such that

$$||bq - b|| < \epsilon/64. \tag{e 5.156}$$

Therefore

$$||e_{\alpha}q - e_{\alpha}|| < \epsilon/64 + ||bq - b|| + \epsilon/64 < 3\epsilon/64.$$
 (e 5.157)

We estimate that

$$||xq - x|| < ||xe_{\alpha}q - xq|| + ||xe_{\alpha}q - xe_{\alpha}|| + ||xe_{\alpha} - x||$$
 (e 5.158)

$$<\epsilon/64 + 3\epsilon/64 + \epsilon/64 = 5\epsilon/64$$
 and (e 5.159)

$$||eq - e|| < 5\epsilon/64.$$
 (e 5.160)

There exists a unitary $u \in M(A \otimes \mathcal{K})$ such that $||u-1|| < 10\epsilon/64$ such that $u^*qu \ge e$. Let $p = u^*qu$.

Theorem 5.7. Let $A \in \mathcal{A}_0$ be a unital separable simple \mathcal{Z} -stable C^* -algebra with unique tracial state τ and let J be the smallest ideal in $M(A \otimes \mathcal{K})$ that properly contains $A \otimes \mathcal{K}$.

Then $K_1(J) = 0$. In fact, for any $u \in \tilde{J}$,

$$cel(u) \leq 7\pi$$
.

Proof. Let $u \in M_n(\tilde{J})$. Let $\Pi: M_n(\tilde{J}) \to M_n$ be the quotient map. Let $z \in M_n$ such that $\Pi(u) = z$. Identify z with the scalar matrix in $M_n(\tilde{J})$ and put $v = uz^*$. Then $v \in \widetilde{M}_n(J)$. Moreover, $\Pi(v) = 1$. Since $M_n(J) \cong J$, without loss of generality, we may assume that $u \in 1 + J$ but we need to show that $\operatorname{cel}(u) \leq 6\pi$.

Let $\epsilon > 0$. By applying 5.6, we may assume that there is a projection $P \in J$ such that

$$u = (1 - P) + u_0,$$

where $u_0 \in PJP$ is a unitary.

Note that $\tau(1-P)=\infty$. Then, by applying 5.3, one obtains a sequence of mutually orthogonal nonzero projections $\{q_n\}$ in $J-A\otimes \mathcal{K}$ such that $\sum_{n=1}^{\infty}q_n$ converges in the strict topology and $\sum_{n=1}^{\infty}q_n\in(1-P)J(1-P)$. By 5.5, putting $P=q_0$, there are a sequence of unitaries, $u_n\in q_nJq_n$ and sequences of selfadjoint elements $\{h_j^{2n-1}\}\subset (q_{2n-1}+q_{2n})J(q_{2n-1}+q_{2n})$ and $\{h_i^{(2n)}\}\subset (q_{2n}+q_{2n+1})J(q_{2n}+q_{2n+1})$ such that

$$||h_1^{(n)}|| \le \pi, ||h_2^{(n)}|| \le 2\pi,$$
 (e 5.161)

$$\|(u_{2n-1} + u_{2n}) - \exp(ih_1^{(n)}) \exp(ih_2^{(n)})\| < \epsilon/2^n \text{ and } (e 5.162)$$

$$\|(u_{2n} + u_{2n+1}) - \exp(ih_1^{2n}) \exp(ih_2^{(2n)})\| < \epsilon/2^n,$$
 (e 5.163)

 $n = 0, 1, 2, \dots$ Since $\sum_{n=1}^{\infty} q_n$ converges in the strict topology and by (e 5.161), so do $\sum_{n=0}^{\infty} u_n$, $\sum_{n=1}^{\infty} u_n$, $\sum_{n=0}^{\infty} h_1^{(n)}$, $\sum_{n=1}^{\infty} h_1^{(n)}$, $\sum_{n=0}^{\infty} h_2^{(n)}$ and $\sum_{n=1}^{\infty} h_2^{(n)}$. Define

$$U_0 = \sum_{n=0}^{\infty} u_n, \ U_1 = \sum_{n=1}^{\infty} u_n \tag{e 5.164}$$

$$H_{0,1} = \sum_{n=0}^{\infty} h_1^{(n)}, \ H_{0,2} = \sum_{n=0}^{\infty} h_2^{(n)}$$
 (e 5.165)

$$H_{1,1} = \sum_{n=1}^{\infty} h_1^{(n)} \text{ and } H_{1,2} = \sum_{n=1}^{\infty} h_2^{(n)}.$$
 (e 5.166)

Then

$$U_0 = u_0 + U_1, (e 5.167)$$

$$||U_1 - \exp(iH_{1,1})\exp(iH_{1,2})|| < \epsilon$$
 (e 5.168)

$$||U_0 - \exp(iH_{0,1})\exp(iH_{02})|| < \epsilon$$
 (e 5.169)

$$||H_{0,1}|| \le \pi \text{ and } ||H_{0,2}|| \le 2\pi.$$
 (e 5.170)

It follows from (e 5.168) that $[U_1] = 0$ in $K_1(\tilde{J})$. Then, by (e 5.169) that $[u_0] = 0$ in $K_1(\tilde{J})$. It follows that [u] = 0 in $K_1(\tilde{J})$. Since u is arbitrarily chosen, we conclude that

$$K_1(\tilde{J}) = 0.$$

Moreover, by (e 5.168), (e 5.169) and (e 5.170), one computes that

$$cel(u) \le 3\pi + 3\pi = 6\pi.$$

So in general,

$$cel(u) \leq 7\pi$$
.

Theorem 5.8. Let $A \in \mathcal{A}_0$ be a unital separable simple \mathcal{Z} -stable C^* -algebra with unique tracial state. Then $M(A \otimes \mathcal{K})/(A \otimes \mathcal{K})$ has real rank zero.

Proof. Let $B =_{df} A \otimes K$, and let $\pi : M(B) \to M(B)/B$ be the natural quotient map. Let J be the smallest ideal in M(B) which properly contains B. Consider the six-term exact sequence:

(e 5.171)

$$K_0(J) \rightarrow K_0(M(B)) \rightarrow K_0(M(B)/J)$$
 $\uparrow \qquad \qquad \downarrow$
 $K_1(M(B)/J) \leftarrow K_1(M(B)) \leftarrow K_1(J)$

From Theorem 5.7, $K_1(J) = 0$. The above six-term exact sequence implies that the map $K_0(M(B)) \to K_0(M(B)/J)$ is surjective. But it is also known that $K_0(M(B)) = 0$ ([4] Proposition 12.2.1; see [41]) Hence, $K_0(M(B)/J) = 0$. Since $M(B)/J \cong \pi(M(B))/\pi(J)$, $K_0(\pi(M(B))/\pi(J)) = 0$. Therefore the map $K_0(\pi(M(B))) \to K_0(\pi(M(B))/\pi(J))$ is surjective.

Since both $\pi(J)$ and $\pi(M(B))/\pi(J)$ are simple purely infinite (this is well-known; an explicit reference can be found in [21]; it also follows immediately from [30] Theorem 3.5 and the definition of J in [24] 2.2, Remark 2.9 and Lemma 2.1; see also [55] Theorem 2.2 and its proof), both $\pi(J)$ and $\pi(M(B))/\pi(J)$ have real rank zero. It follows from [6] Theorem 3.14 and Proposition 3.15 that $\pi(M(B))$ has real rank zero.

Theorem 5.9. $M(\mathcal{Z} \otimes \mathcal{K})/(\mathcal{Z} \otimes \mathcal{K})$ has real rank zero.

Remark 5.10. We note that Theorem 5.8 includes many other C*-algebras (see [37])—including crossed products coming from uniquely ergodic minimal homeomorphisms on a compact metric space with finite topological dimension (e.g., see [53], [8].)

Finally, we end the paper with some K-theory computations that follow immediately from our work.

Corollary 5.11. Let $\pi: M(\mathcal{Z} \otimes \mathcal{K}) \to M(\mathcal{Z} \otimes \mathcal{K})/(\mathcal{Z} \otimes \mathcal{K}) = Q(\mathcal{Z})$ be the natural quotient map. Let J be the unique smallest ideal of $M(\mathcal{Z} \otimes \mathcal{K})$ which properly contains $\mathcal{Z} \otimes \mathcal{K}$.

Then we have the following: $K_0(J) = \mathbb{R}$, $K_1(J) = 0$, $K_0(\pi(J)) = \mathbb{R}/\mathbb{Z} \cong \mathbb{T}$, $K_1(\pi(J)) = 0$, $K_0(Q(\mathcal{Z})/\pi(J)) = 0$ and $K_1(Q(\mathcal{Z})/\pi(J)) = \mathbb{R}$.

Proof. Let $B =_{df} \mathcal{Z} \otimes \mathcal{K}$.

That $K_0(J) = \mathbb{R}$ follows immediately from Proposition 4.2 and Proposition 4.4.

That $K_1(J) = 0$ follows immediately from Theorem 5.7.

That $K_0(Q(\mathcal{Z})/\pi(J)) = 0$ was computed in the proof of Theorem 5.8.

Hence, the six-term exact sequence (e 5.171) from Theorem 5.8 thus becomes the following six-term exact sequence:

Hence, $K_1(Q(\mathcal{Z})/\pi(J)) = K_1(M(B)/J) = \mathbb{R}$.

Next, the exact sequence

$$0 \to B \to J \to J/B \to 0$$

results in the six term exact sequence

$$K_0(B) \rightarrow K_0(J) \rightarrow K_0(J/B)$$
 $\uparrow \qquad \qquad \downarrow$
 $K_1(J/B) \leftarrow K_1(J) \leftarrow K_1(B)$

which, by the above results, becomes

Since the map $\mathbb{Z} \to \mathbb{R}$ is the natural inclusion, $K_1(\pi(J)) = 0$ and $K_0(\pi(J)) = \mathbb{R}/\mathbb{Z} \cong \mathbb{T}$. \square

References

- [1] R. Antoine, J. Bosa and F. Perera, Completions of monoids with applications to the Cuntz semigroup, Internat. J. Math., 22 (2011), no. 6, 837–861. A copy is available at http://arxiv.org/pdf/1003.2874.
- [2] B. E. Blackadar, A simple C^* -algebra with no nontrivial projections. Proc. Amer. Math. Soc. 78 (1980), no. 4, 504–508.
- [3] B. E. Blackadar, A simple unital projectionless C*-algebra. J. Operator Theory, 5 (1981), no. 1, 63–71.
- [4] B. E. Blackadar, K-Theory for operator algebras. Second edition. Mathematical Sciences Research Institute Publications, 5. Cambridge University Press, Cambridge, 1998.
- [5] L. G. Brown, *Extensions of AF-algebras*, Operator Algebras and Applications, (Kingston, Ont. 1980) R. V. Kadison (ed.), Proc. Symp. Pure Math., **38** (1982) 175–176.
- [6] L. G. Brown and G. K. Pedersen, C*-algebras of real rank zero, Journal of Functional Analysis, 99 (1991), 131–149.
- [7] N. P. Brown, F. Perera and A. S. Toms, *The Cuntz semigroup, the Elliott conjecture, and dimension functions on C*-algebras*, J. Reine Angew. Math. **621** (2008), 191–211. A copy is available at http://arxiv.org/pdf/0609182v2.

- [8] A. Connes, An analogue of the Thom isomorphism for crossed products of a C*-algebra by an action of \mathbb{R} . Adv. in Math. 39 (1981), no. 1, 31–55.
- [9] K. T. Coward, G. A. Elliott and C. Ivanescu, The Cuntz semigroup as an invariant for C*-algebras, J. Reine Angew. Math. 623 (2008), 161–193. A copy is available at http://arxiv.org/pdf/0705.0341.
- [10] D. A. Edwards, Separation des fonctions reelles definies sur un simplexe de Choquet, C. R. Acad. Sci. Paris, 261 (1965), 2798-2800.
- [11] G. A. Elliott, *Derivations of matroid C*-algebras, II*, Annals of Mathematics, Second Series **100** (1974), no. 2, 407–422.
- [12] G. A. Elliott, Automorphisms determined by multipliers on ideals of a C*-algebra, J. Functional Analysis, 23 (1976) no. 1, 1–10.
- [13] G. A. Elliott, The ideal structure of the multiplier algebra of an AF-algebra, C. R. Math. Rep. Acad. Sci. Canada, 9 (1987), 225–230.
- [14] K. Goodearl, Partially ordered abelian groups with interpolation. Math. Surveys and Monographs, 20, Amer. Math. Soc., Providence, 1986.
- [15] N. Higson and M. Rordam, *The Weyl-von Neumann Theorem for multipliers*, Canad. J. Math., **43** (1991) no. 2, 322–330.
- [16] K. K. Jensen and K. Thomsen, Elements of KK-theory. Mathematics: Theory and Applications. Birkhauser Boston, Inc., Boston, MA, 1991.
- [17] X. Jiang and H. Su, On a simple unital projectionless C^* -algebra, Amer. J. Math., 121 (1999), 359–413.
- [18] V. Kaftal, P. W. Ng and S. Zhang, Strict comparison of positive elements in multiplier algebras. Preprint. A copy is available at http://arxiv.org/pdf/1501.05463.
- [19] I. Kaplansky, Functional analysis, some aspects of analysis and probability, Wiley, New York, 1958, pp. 1–34.
- [20] G. G. Kasparov, Hilbert C*-modules: Theorems of Stinespring and Voiculescu, J. Operator Theory, 4 (1980), 133–150.
- [21] D. Kucerovsky, P. W. Ng and F. Perera, Purely infinite corona algebras of simple C*-algebras, Math. Ann. **346** (2010), no. 1, 23–40.
- [22] K. B. Laursen and A. M. Sinclair, Lifting matrix units in C*-algebras. II. Math. Scand. 37 (1975), no. 1, 167–172.
- [23] H. Lin, *Ideals of multiplier algebras of simple AF C*-algebras*, Proc. Amer. Math. Soc., **104** (1988), 239–244.
- [24] H. Lin, Simple C*-algebras with continuous scales and simple corona algebras, Proc. Amer. Math. Soc., 112 (1991), no. 3, 871–880.
- [25] H. Lin, Generalized Weyl-von Neumann Theorems, Internat. J. Math. 2 (1991) no. 6, 725–739.

- [26] H. Lin, Exponential rank of C*-algebras with real rank zero and the Brown-Pedersen Conjectures, Journal of Functional Analysis, 114 (1993), 1-11.
- [27] H. Lin, Generalized Weyl-von Neumann Theorems. II. Math. Scand., 77 (1995) no. 1, 129–147.
- [28] H. Lin, Simple corona C*-algebras, Proc. Amer. Math. Soc. **132** (2004), 3215–3224.
- [29] H. Lin, The tracial topological rank of C*-algebras, Proc. London Math. Soc., 83 (2001), 199–234.
- [30] H. Lin, Full extensions and approximate unitary equivalence, Pacific J. Math., 229 (2007) no. 2, 389–428.
- [31] H. Lin, Approximate homotopy of homomorphisms from C(X) into a simple C^* -algebra, Mem. Amer. Math. Soc., **205** (2010), no. 963, vi+131 pp. ISBN: 978-0-8218-5194-4.
- [32] H. Lin, Cuntz semigroups of C*-algebras of stable rank one and projective Hilbert modules, preprint. A copy is available at http://arxiv.org/pdf/1001.4558.
- [33] H. Lin, Asymptotic unitary equivalence and classification of simple amenable C*-algebras, Invent. Math. **183** (2011), 385–450.
- [34] H. Lin, Approximate unitary equivalence in simple C*-algebras of tracial rank one, Trans. Amer. Math. Soc. **364** (2012), no. 4, 2021–2086.
- [35] H. Lin, Homomorphisms from AH-algebras, preprint, arXiv:1102.4631.
- [36] H. Lin, Exponentials in simple Z-stable simple C*-algebras, J. Funct. Anal. 266 (2014), no. 2, 754–791.
- [37] H. Lin and Z. Niu, The range of a class of classifiable separable simple amenable C*-algebras, J. Funct. Anal., **260**, (2011), 1–29.
- [38] H. Lin and Z. Niu, Homomorphisms into a simple \mathbb{Z} -stable C^* -algebras, preprint (arXiv:1003.1760).
- [39] H. Lin and H. Osaka, Real rank of multiplier algebras of C*-algebras of real rank zero. Operator algebras and their applications (Waterloo, ON, 1994/1995), 235–241. Fields Inst. Commun., 13, Amer. Math. Soc., Providence, RI, 1997.
- [40] H. Lin and W. Sun, Tensor products of classifiable C*-algebras, preprint (arXiv:1203.3737).
- [41] J. A. Mingo, K-theory and multipliers of stable C*-algebras, Transactions of the American Mathematical Society, **299** (1987), no. 1, 397–411.
- [42] G. J. Murphy, Diagonality in C*-algebras, Math. Z., 199 (1988) no. 2, 279–284.
- [43] E. Ortega, M. Rordam and H. Thiel, The Cuntz semigroup and comparison of open projections, J. Funct. Anal., 260 (2011), 3474–3493.
- [44] G. K. Pedersen, Measure Theory for C*-algebras. Math. Scand. 19 (1966), 131–145.
- [45] G. K. Pedersen, Unitary extensions and polar decompositions in a C*-algebra, J. Operator Theory 17 (1987), 357–364.

- [46] N.C. Phillips, Approximation by unitaries with finite spectrum in purely infinite C*-algebras,
 J. Funct. Anal. 120 (1994), 98–106.
- [47] M. Rordam, On the structure of simple C*-algebras tensored with a UHF-algebra, Journal of Functional Analysis, 100 (1991), 1–17.
- [48] M. Rordam, *Ideals in the multiplier algebra of a stable C*-algebra*, J. Operator Theory, **25** (1991), 283–298.
- [49] M. Rordam and W. Winter, *The Jiang-Su algebra revisited*, J. Reine Angew. Math., **642** (2010), 129–144.
- [50] A. Tikuisis and A. Toms, On the structure of the Cuntz semigroup in (possibly) nonunital C*-aglebras. To appear in Canad. Math. Bull.
- [51] A. S. Toms and W. Winter, Strongly self-absorbing C*-algebras. Trans. Amer. Math. Soc. **359** (2007), no. 8, 3999–4029.
- [52] N. E. Wegge-Olsen, K-theory and C*-algebras. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1993.
- [53] W. Winter, Decomposition rank and Z-stability. Invent. Math. 179 (2010), no. 2, 229–301.
- [54] W. Winter, Localizing the Elliott conjecture at strongly self-absorbing C*-algebras. J. Reine Angew. Math. **692** (2014), 193–231.
- [55] S. Zhang, On the structure of projections and ideals of corona algebras, Canadian J. Math. 41 (1989), 721–742.
- [56] S. Zhang, A property of purely infinite simple C*-algebras. Proc. Amer. Math. Soc. 109 (1990), no. 3, 717–720.
- [57] S. Zhang, A Riesz decomposition property and ideal structure of multiplier algebras, J. Operator Theory, 24 (1990), 209–226.
- [58] S. Zhang, Trivial K_1 -flow of AF algebras and finite von Neumann algebras. Journal of Functional Analysis **92** (1990), 77–91.
- [59] S. Zhang, K_1 -groups, quasidiagonality, and interpolation by multiplier projections, Trans. Amer. Math. Soc. **325** (1991) no. 2, 792–818.
- [60] S. Zhang, C*-algebras with real rank zero and their corona and multiplier algebras. IV. Internat. J. Math. 3 (1992) no. 2, 309–330.
- [61] S. Zhang, Certain C*-algebras with real rank zero and their corona and multiplier algebras. I. Pacific J. Math. 155 (1992), no. 1, 169–197.