

On the synectic metric in the tangent bundle of a Riemannian manifold

Melek ARAS*

September 25, 2018

Abstract

The purpose of this paper is to investigate applications the covariant derivatives of the covector fields and killing vector fields with respect to the synectic lift ${}^S g = {}^C g + {}^V a$ in a the Riemannian manifold to its tangent bundle $T(M_n)$, where ${}^C g$ -complete lift of the Riemannian metric, ${}^V a$ -vertical lift of the symmetric tensor field of type $(0, 2)$ in M_n .

Keywords: Tensor bundle ; Metric connection; Covector field; Levi-Civita connections; Killing vector field

1. Introduction

Suppose that there is given the following Riemannian metric

$${}^S \tilde{g}_{CB} dx^C dx^B = a_{ji} dx^j dx^i + 2g_{ji} dx^j \delta y^i \quad (1)$$

in tangent bundle in $T(M_n)$ over a Riemannian manifold M_n with metric g , where a_{ji} are components of a symmetric tensor field of type $(0, 2)$ in M_n and $\delta y^h = dy^h + \Gamma_i^h dx^i$, $\Gamma_i^h = y^j \Gamma_{ji}^h$ with respect to the induced coordinates (x^h, y^h) in $\pi^{-1}(U) \subset T(M_n)$. We call this metric the synectic metric. The synectic metric ${}^S g = {}^C g + {}^V a$ has respectively components[2] and its contravariant components

$${}^S g = ({}^S \tilde{g}_{CB}) = \begin{pmatrix} a_{ji} + \partial g_{ji} & g_{ji} \\ g_{ji} & 0 \end{pmatrix}, {}^S \tilde{g}^{CB} = \begin{pmatrix} 0 & g^{ji} \\ g^{ji} & x^{\bar{s}} \partial_s g^{ji} - a_{\bar{s}}^{ji} \end{pmatrix} \quad (2)$$

where $\partial g_{ji} = x^{\bar{s}} \partial_s g_{ji}$ and $a_{\bar{s}}^{ti} = g^{jt} a_{js} g^{si}$.

Components of the Riemannian connection determined by the synectic metric ${}^S g$ are [1]

$$\begin{cases} {}^S \Gamma_{ji}^k = \Gamma_{ji}^k, & {}^S \Gamma_{\bar{j}\bar{i}}^k = \Gamma_{\bar{j}\bar{i}}^k, & {}^S \Gamma_{\bar{j}\bar{i}}^k = \Gamma_{\bar{j}\bar{i}}^k, & {}^S \Gamma_{\bar{j}\bar{i}}^k = 0 \\ {}^S \Gamma_{\bar{j}\bar{i}}^k = {}^S \Gamma_{\bar{j}\bar{i}}^k = {}^S \Gamma_{\bar{j}\bar{i}}^k = 0, & {}^S \Gamma_{\bar{j}\bar{i}}^k = x^{\bar{t}} \partial_t \Gamma_{\bar{j}\bar{i}}^k + H_{\bar{j}\bar{i}}^k \end{cases} \quad (3)$$

*Department of Mathematics, Faculty of Arts and Sciences, Giresun University, 28049, Turkey e-mail: melek.aras@iresun.edu.tr; melek.aras25@hotmail.com

with respect to the induced coordinates in $T(M_n)$, Γ_{ji}^k being Christoffel symbols constructed with g_{ji} . Where $H_{ji}^k = \frac{1}{2}g^{ks}(\nabla_j a_{si} + \nabla_i a_{js} - \nabla_s a_{ji})$ is a tensor of type (1, 2) and $\nabla_s a_{ji} = \partial_s a_{ji} - \Gamma_{kj}^l a_{li} - \Gamma_{ki}^l a_{jl}$.

The metric connection $\tilde{\nabla}$ of the synectic metric satisfies $\tilde{\nabla}_C {}^S g_{BA} = 0$ and has non-trivial torsion tensor \tilde{T}_{CB}^A , which is skew-symmetric in the indices C and B . Then the metric connection $\tilde{\nabla}$ of the synectic metric has components[1]

$$\begin{cases} \tilde{\Gamma}_{ji}^h = \tilde{\Gamma}_{ji}^{\tilde{h}} = \tilde{\Gamma}_{ji}^{\tilde{h}} = \Gamma_{ji}^h, \\ \tilde{\Gamma}_{ji}^{\tilde{h}} = \tilde{\Gamma}_{ji}^{\tilde{h}} = \tilde{\Gamma}_{ji}^{\tilde{h}} = \Gamma_{ji}^h, \\ \tilde{\Gamma}_{ji}^{\tilde{h}} = x^t \partial_t \Gamma_{ji}^h + H_{ji}^h - y^k R_{kjih} \end{cases} \quad (4)$$

with respect to the induced coordinates, Γ_{ji}^k being Christoffel symbols formed with g_{ji} , where $H_{ji}^k = \frac{1}{2}g^{ks}(\nabla_j a_{si} + \nabla_i a_{js} - \nabla_s a_{ji})$.

Given a vector field \tilde{X} in $T(M_n)$, the 1-form, $\tilde{\omega}$ defined by $\tilde{\omega}(\tilde{Y}) = \tilde{g}(\tilde{X}, \tilde{Y})$, \tilde{Y} being an arbitrary element of $T_0^1(M_n)$, is called the covector field associated with \tilde{X} and denoted by \tilde{X}^* . If \tilde{X} has local components \tilde{X}^A , then the associated covector field \tilde{X}^* of \tilde{X} has local components $\tilde{X}_C = \tilde{g}_{CA} \tilde{X}^A$.

Let ω be a 1-form in M_n with components ω_i . Then the vertical, complete and horizontal lifts of ω to $T(M_n)$ have respectively components[4]

$$({}^V \omega_B) = (\omega_i, 0), \quad ({}^C \omega_B) = (\partial \omega_i, \omega_i), \quad ({}^H \omega_B) = (-\Gamma_i^k \omega_k, \omega_i)$$

with respect to the induced coordinates in $T(M_n)$.

The associated covector fields of the vertical, complete and horizontal lifts to $T(M_n)$, with the synectic metric, of a vector field X with components X^h in M_n are respectively

$$(X_i, 0), \quad (y^s \partial_s X_i + a_{ij} X^j, X_i), \quad (\Gamma_i^h X_h + a_{ij} X^j, X_i) \quad (5)$$

with respect to the induced coordinates, where $X_j = g_{ji} X^i$ are components of the covector field X^* associated with X .

A vector field $X \in \mathfrak{S}_0^1(M_n)$ is said to be a Killing vector field of a Riemannian manifold with metric g , if $L_X g = 0$ [3]. In terms of components g_{ji} of g , X is a Killing vector field if and only if

$$L_X g = X^\alpha \nabla_\alpha g_{ji} + g_{\alpha i} \nabla_j X^\alpha + g_{j\alpha} \nabla_i X^\alpha = \nabla_j X_i + \nabla_i X_j = 0,$$

X^α being components of X , where ∇ is the Riemannian connection of the metric g .

2. Main Results

We now take a vector field X in M_n with components X^h . Then, since the associated covector fields of the lifts of X have respectively components give by (5), we have by (3) and (5)

$$\left\{ \begin{array}{l} {}^S\nabla_B^V X_A = \begin{pmatrix} \nabla_j X_i & 0 \\ 0 & 0 \end{pmatrix}, \\ {}^S\nabla_B^C X_A = \begin{pmatrix} \partial(\nabla_j X_i) + \nabla_j(a_{il}X^l) - H_{ji}^m X_m & \nabla_j X_i \\ \nabla_j X_i & 0 \end{pmatrix} \end{array} \right. \quad (6)$$

and consequently

$$\left\{ \begin{array}{l} {}^S\nabla_B^V X_A + {}^S\nabla_A^V X_B = \begin{pmatrix} \nabla_j X_i + \nabla_i X_j & 0 \\ 0 & 0 \end{pmatrix}, \\ {}^S\nabla_B^C X_A + {}^S\nabla_A^C X_B = \begin{pmatrix} {}^S\nabla_j^C X_i + {}^S\nabla_i^C X_j & {}^S\nabla_j^C X_{\bar{i}} + {}^S\nabla_{\bar{i}}^C X_j \\ {}^S\nabla_j^C X_i + {}^S\nabla_i^C X_{\bar{j}} & {}^S\nabla_j^C X_{\bar{i}} + {}^S\nabla_{\bar{i}}^C X_{\bar{j}} \end{pmatrix} \\ {}^S\nabla_j^C X_i + {}^S\nabla_i^C X_j = \partial(\nabla_j X_i + \nabla_i X_j) \\ \quad + \nabla_j(a_{il}X^l) + \nabla_i(a_{jl}X^l) - H_{ji}^m X_m - H_{ij}^m X_m \\ {}^S\nabla_j^C X_{\bar{i}} + {}^S\nabla_{\bar{i}}^C X_j = \nabla_j X_i + \nabla_i X_j \\ {}^S\nabla_j^C X_i + {}^S\nabla_i^C X_{\bar{j}} = \nabla_j X_i + \nabla_i X_j^S \\ \nabla_j^C X_{\bar{i}} + {}^S\nabla_{\bar{i}}^C X_{\bar{j}} = 0 \end{array} \right. \quad (7)$$

with respect to the induced coordinates ,where $X_j = g_{jk}X^k$. From (7) we have

Theorem 1 *Necessary and sufficient conditions in order that (a) the vertical, (b) complete lifts to $T(M_n)$, with the symplectic metric, of a vector field X in M_n be a Killing vector field in $T(M_n)$ are that, respectively, (a) X is a Killing vector field in M_n and (b) X is Killing vector field with vanishing covariant derivative in M_n and the covariant derivative of symmetric tensor field a of type $(0,2)$ vanishes.*

We also have by (2) and (6)

$$\left\{ \begin{array}{l} {}^S\nabla_B^V X_A - {}^S\nabla_A^V X_B = \begin{pmatrix} \nabla_j X_i - \nabla_i X_j & 0 \\ 0 & 0 \end{pmatrix}, \\ {}^S\nabla_B^C X_A - {}^S\nabla_A^C X_B = \begin{pmatrix} {}^S\nabla_j^C X_i - {}^S\nabla_i^C X_j & {}^S\nabla_j^C X_{\bar{i}} - {}^S\nabla_{\bar{i}}^C X_j \\ {}^S\nabla_j^C X_i - {}^S\nabla_i^C X_{\bar{j}} & {}^S\nabla_j^C X_{\bar{i}} - {}^S\nabla_{\bar{i}}^C X_{\bar{j}} \end{pmatrix} \\ {}^S\nabla_j^C X_i - {}^S\nabla_i^C X_j = \partial(\nabla_j X_i - \nabla_i X_j) \\ \quad + \nabla_j(a_{il}X^l) - \nabla_i(a_{jl}X^l) - H_{ji}^m X_m + H_{ij}^m X_m \\ {}^S\nabla_j^C X_{\bar{i}} + {}^S\nabla_{\bar{i}}^C X_j = \nabla_j X_i - \nabla_i X_j \\ {}^S\nabla_j^C X_i + {}^S\nabla_i^C X_{\bar{j}} = \nabla_j X_i - \nabla_i X_j^S \\ \nabla_j^C X_{\bar{i}} - {}^S\nabla_{\bar{i}}^C X_{\bar{j}} = 0 \end{array} \right. \quad (8)$$

with respect to the induced coordinates ,where ${}^S g^{BA} {}^S\nabla_B^V X_A = 0$ and ${}^S g^{BA} {}^S\nabla_B^C X_A = 2g^{ji}\nabla_j X_i$. Thus we have, from (8), respectively,

Theorem 2 *The vertical lift of a vector field in M_n to $T(M_n)$ with the symplectic metric Sg is harmonic if and only if the vector field in M_n is closed.*

Theorem 3 *The complete lift of a vector field in M_n to $T(M_n)$ with the symplectic metric Sg is harmonic if and only if the vector field in M_n is harmonic and the covariant derivative of symmetric tensor field a of type $(0, 2)$ vanishes.*

We consider a vector field $X \in \mathfrak{S}_0^1(M_n)$. Then its vertical, complete and horizontal lifts have components of the form

$${}^V X = \begin{pmatrix} 0 \\ X^h \end{pmatrix}, \quad {}^C X = \begin{pmatrix} X^h \\ \partial X^h \end{pmatrix}, \quad {}^H X = \begin{pmatrix} X^h \\ -\Gamma_i^h X^i \end{pmatrix} \quad (9)$$

with respect to the induced coordinates in $T(M_n)$, where $\Gamma_i^h X^i = y^s \Gamma_{si}^h X^i$.

Let X be a vector field in M_n with local components X^k . Then, from (9) and (4), we see that, the covariant derivatives of the vertical, complete and horizontal lifts of $X \in \mathfrak{S}_0^1(M_n)$ with the metric connection $\tilde{\nabla}$ have respectively components

$$\left\{ \begin{array}{l} \tilde{\nabla}_B {}^V X^A = \begin{pmatrix} 0 & 0 \\ \nabla_j X^h & 0 \end{pmatrix}, \\ \tilde{\nabla}_B {}^C X^A = \begin{pmatrix} \nabla_j X^h & 0 \\ \partial(\nabla_j X^h) + H_{jm}^h X^m - y^k R_{kjm}^h X^m & \nabla_j X^h \end{pmatrix}, \\ \tilde{\nabla}_B {}^H X^A = \begin{pmatrix} \nabla_j X^h & 0 \\ -\Gamma_i^h (\nabla_j X^i) + H_{jm}^h X^m & 0 \end{pmatrix} \end{array} \right. \quad (10)$$

with respect to the induced coordinates $T(M_n)$.

Remark 4 $\tilde{\nabla} = \bar{\nabla} + {}^V H$, where $\bar{\nabla}$ is the metric connection with the metric ${}^C g$ [1]

Remark 5 *The metric connection $\bar{\nabla}$ coincides with the horizontal lift ${}^H \nabla$ of Levi Civita connection ∇ of g in M_n [4]. Thus we have*

Proposition 6 *Necessary and sufficient conditions in order that (a) the vertical, (b) complete and horizontal lifts of a vector field in M_n to $T(M_n)$ with the metric connection $\tilde{\nabla}$ be parallel in $T(M_n)$ are that, respectively, (a) the vector field given in M_n is parallel (b) the vector field given in M_n is parallel and the covariant derivative of symmetric tensor field a of type $(0, 2)$ vanishes.*

Since $\nabla_j X^h = t\delta_j^h$ with constant t implies $R_{kji}^h X^i = 0$, we have also

Proposition 7 *The complete lift of a vector in M_n to $T(M_n)$ with the metric connection $\tilde{\nabla}$ is concurrent if and only if the vector field given in M_n is concurrent and the covariant derivative of symmetric tensor field a of type $(0, 2)$ vanishes.*

A vector field $X \in \mathfrak{S}_0^1(M_n)$ is said to be an infinitesimal isometry or a Killing vector field of a Riemannian manifold with metric g , if $L_X g = 0$. In terms of components g_{ji} of g , X is an infinitesimal isometry if and only if

$$X^\gamma \partial_\gamma g_{\alpha\beta} + g_{\alpha\gamma} \partial_\beta X^\gamma + g_{\gamma\beta} \partial_\alpha X^\gamma = 0, \quad (11)$$

X^α being components of X [3], where the indices α, β and γ run over the range $\{1, 2, \dots, m\}$.

Let there be given in M_n a Riemannian metric g with components g_{ji} . Let \tilde{X} be a vector field with components $\begin{pmatrix} \tilde{X}^k \\ \tilde{X}^{\bar{k}} \end{pmatrix}$ with respect to the induced coordinates in $T(M_n)$.

with respect to the induced coordinates in $T(M_n)$. Then, taking account of (2), we see by virtue of (11) that \tilde{X} is an infinitesimal isometry in $T(M_n)$ with metric ${}^S g$ if and only if

$$\begin{cases} \left(\tilde{X}^h \partial_h \partial g_{ji} + \tilde{X}^{\bar{h}} \partial_h g_{ji} \right) + \left(\partial g_{jh} \partial_i \tilde{X}^h + g_{jh} \partial_i \tilde{X}^{\bar{h}} \right) \\ + \left(\partial g_{hi} \partial_j \tilde{X}^h + g_{hi} \partial_j \tilde{X}^{\bar{h}} \right) + \left(\tilde{X}^h \partial_h a_{ji} + a_{jh} \partial_i \tilde{X}^h + a_{hi} \partial_j \tilde{X}^h \right) = 0 \end{cases} \quad (12)$$

$$\tilde{X}^h \partial_h g_{ji} + g_{jh} \partial_i \tilde{X}^h + \left(\partial g_{hi} \partial_j \tilde{X}^h + g_{hi} \partial_j \tilde{X}^{\bar{h}} \right) + a_{hi} \partial_j \tilde{X}^h = 0 \quad (13)$$

$$\tilde{X}^h \partial_h g_{ji} + g_{hi} \partial_j \tilde{X}^h + \left(\partial g_{jh} \partial_i \tilde{X}^h + g_{jh} \partial_i \tilde{X}^{\bar{h}} \right) + a_{jh} \partial_i \tilde{X}^h = 0 \quad (14)$$

$$g_{jh} \partial_i \tilde{X}^h + g_{hi} \partial_j \tilde{X}^h = 0. \quad (15)$$

We shall now prove

Lemma 8 *Let C be an element of $\mathfrak{S}_1^1(M_n)$. Then $L_{\iota C} {}^S g = 0$ holds if and only if $C = 0$.*

Proof. Denote by C_i^k the local components of C . Then ιC has components $\begin{pmatrix} 0 \\ y^i C_i^k \end{pmatrix}$ with respect to the induced coordinates in $T(M_n)$. Thus, substituting $\tilde{X}^k = 0$ and $\tilde{X}^{\bar{k}} = y^i C_i^k$ in (13), we have $g_{ki} C_j^k = 0$, which implies $C_j^k = 0$, i.e., $C = 0$. ■

References

- [1] Aras, M. The Metric Connection with Respect to the Synectic Metric. Hacett. J. Math. Stat, 2012, 41(2), 169-173.
- [2] Vishnevskii V V, Shirokov A and Shurygin V V, Spaces over algebras (Kazan University Press) (1985)

- [3] Yano, K. The Theory of Lie Derivatives and Applications, Elsevier, Amsterdam, 1957.
- [4] Yano, K. and Ishihara, S. Tangent and Cotangent Bundles, Marcel Dekker Inc. New York, 1973.