

ON CHARACTERISTIC INTEGRALS OF TODA FIELD THEORIES

ZHAOHU NIE

ABSTRACT. Characteristic integrals of Toda field theories associated to general simple Lie algebras are constructed using systematic techniques, and complete mathematical proofs are provided. Plenty of examples illustrating the results are presented in explicit forms.

1. INTRODUCTION

First consider the famous Liouville equation, for which we take the following version:

$$(1.1) \quad u_{xy} = -e^{2u},$$

where x and y are the independent variables, and $u = u(x, y)$ is an unknown function. Let

$$(1.2) \quad I = u_{xx} - u_x^2.$$

Then it is easily checked that $I_y = \frac{\partial}{\partial y} I = 0$ for a solution $u(x, y)$ to (1.1). I is thus called a *characteristic integral* of (1.1).

Toda field theories are generalizations of the Liouville equation (1.1). Let \mathfrak{g} be a simple Lie algebra of rank n with Cartan matrix $A = (a_{ij})_{i,j=1}^n$. For $1 \leq i \leq n$, let $u^i = u^i(x, y)$ be n unknown functions of the independent variables x and y . The Toda field theory associated to \mathfrak{g} is the system of nonlinear PDEs:

$$(1.3) \quad u_{xy}^i = -e^{\rho_i} := -\exp\left(\sum_{j=1}^n a_{ij}u^j\right), \quad 1 \leq i \leq n.$$

The Liouville equation (1.1) is the Toda field theory associated to $A_1 = \mathfrak{sl}_2$.

Toda field theories are fundamental integrable systems with rich properties and important applications in mathematics and physics. They have been extensively studied, and we refer to the two books [BBT03, LS92] for surveys on them. In this paper, we are concerned with the explicit forms of their characteristic integrals.

Definition 1.4. A differential polynomial in the $u^i(x, y)$ for $1 \leq i \leq n$ is a polynomial in the u^i and their partial derivatives with respect to x of various orders u_x^i, u_{xx}^i, \dots .

A *characteristic integral* of the Toda field theory (1.3) is a differential polynomial I in the $u^i(x, y)$ for $1 \leq i \leq n$ such that $I_y = 0$ for the solutions $u^i(x, y)$ to (1.3).

By the symmetry between x and y in the Toda field theory (1.3), there are differential polynomials \tilde{I} in the u^i and their y -partial derivatives such that $\tilde{I}_x = 0$ but they clearly follow the same patterns as those in the above definition.

Since a differential polynomial in characteristic integrals is another such integral, the characteristic integrals form a differential algebra. Shabat and Yamilov [SY81] showed that the characteristic integrals of the Toda field theory (1.3) form a *polynomial* differential algebra generated by n *primitive characteristic integrals*, and that the system (1.3) has such a complete set of characteristic integrals if and only if the matrix $A = (a_{ij})$ is equivalent to the Cartan matrix of a simple Lie algebra.

For a differential monomial in the u^i , we call by its *degree* the sum of the orders of differentiation multiplied by the algebraic degrees of the corresponding factors. Therefore the I in (1.2) has a homogeneous degree 2. The primitive characteristic integrals can be chosen to be homogeneous and their degrees are equal to the degrees of the Lie algebra \mathfrak{g} . Recall that the algebra of adjoint-invariant functions on \mathfrak{g} is a polynomial algebra on n homogenous generators, whose degrees we call the degrees of \mathfrak{g} [Kos59]. For a proof of this structure theorem, see [Sha95] and [FF96, Theorem 2.4.10 and Proposition 2.4.7]. This paper is concerned with constructing these primitive characteristic integrals.

Many works [LS92, BFO⁺90, FOR⁺92, Dem10] have been devoted to the characteristic integrals, very often under different names such as *local conservation laws*, *chiral currents*, *intermediate integrals* or *W-algebras* and from different viewpoints. In particular, the works of Guryeva and Zhiber [GZ04, GZ05] have shown that the Toda field theories (1.3) for classical and exceptional Lie algebras are of Liouville type, and have obtained explicit formulas for the generalized Laplace invariants for these systems, which can be used to construct the characteristic integrals and higher symmetries.

However to this author, the construction results about characteristic integrals for the Toda field theories (1.3) are not explicit enough, in terms of both the formulas and the proofs. Therefore we would like to present here systematic and concrete techniques for constructing the primitive characteristic integrals together with new and self-contained proofs.

Furthermore, many works [LS92, HO96, ZS01] utilize the characteristic integrals to obtain explicit solutions to the original Toda field theories (1.3) by the method of Darboux. Recently, Anderson, Fels and Vassiliou [AFV09] developed a very general approach to the study of Darboux integrable systems.

By constructing enough characteristic integrals, this paper explicitly shows that the Toda field theories (1.3) are Darboux integrable. In sequels to this work, Anderson and the author will apply the characteristic integrals found in this paper and the structure theorem in [AFV09] to realize the Toda field theory (1.3) as the quotient of two standard differential systems [Yam93] depending on x and y separately. We will also apply the method and result of this paper to the setting of differential invariants for the standard differential systems, generalizing work by Mari Beffa [MB08].

This paper is organized as follows. In Section 2, we present our main results. Our main theorem is Theorem 2.6, which is applicable to all simple Lie algebras. It employs the zero curvature representation [LS79] of the Toda field theories (1.3) under a Drinfeld-Sokolov gauge [DS84]. When the Lie algebra \mathfrak{g} has a non-branching representation (see (2.10)), which is the case for Lie algebras of types A, B, C and \mathfrak{g}_2 , we present a more concrete formula in Theorem 2.11, which is easier to use. We stress that in both cases, we provide complete and novel mathematical proofs.

In Sections 3, we demonstrate the more direct method in Theorem 2.11 in all the applicable cases. In Section 4, we illustrate the general method in Theorem 2.6 by the example of the characteristic integral related to the Pfaffian of D_4 . We stress that our formulas are easily implemented using mathematical softwares such as Maple, which this author and Anderson have done. Maple programs and formulas for the characteristic integrals can be found at the DifferentialGeometry Software Project website at the Digital Commons of Utah State Univeristy (<http://digitalcommons.usu.edu/dg/>). This makes our results readily usable for various purposes.

Acknowledgment. The author thanks Ian Anderson for introducing this topic to him. He thanks Anderson and Pawel Nurowski for helps with Maple. The author also thanks László Fehér and Luen-Chau Li for their interests and useful correspondences. He thanks the referee for detailed comments and some references.

2. MAIN RESULTS

Let us first introduce some terminology. Let $\mathfrak{h} \subset \mathfrak{g}$ be a Cartan subalgebra, and we denote the corresponding set of roots of \mathfrak{g} by Δ , the sets of positive/negative roots by Δ_{\pm} , and the set of positive simple roots by $\pi = \{\alpha_i\}_{i=1}^n$. Let $\mathfrak{g} = \mathfrak{h} \oplus \bigoplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha}$ be the root space decomposition.

For a root α , define its height by $\text{ht}(\alpha) = \sum_{i=1}^n c_i$ if $\alpha = \sum_{i=1}^n c_i \alpha_i$. Also define the principal height gradation

$$(2.1) \quad \mathfrak{g} = \bigoplus_{k=-m}^m \mathfrak{g}_k, \quad \mathfrak{g}_k = \bigoplus_{\text{ht}(\alpha)=k} \mathfrak{g}_{\alpha}, \quad \mathfrak{g}_0 = \mathfrak{h},$$

where m is the maximal height of the roots. We also denote by $\mathfrak{n} = \bigoplus_{\alpha \in \Delta_+} \mathfrak{g}_{\alpha} = \bigoplus_{k>0} \mathfrak{g}_k$ the maximal nilpotent subalgebra, and by N the corresponding unipotent group.

For $\alpha \in \Delta_+$, let e_{α} and $e_{-\alpha}$ be root vectors in the root spaces \mathfrak{g}_{α} and $\mathfrak{g}_{-\alpha}$ such that for $H_{\alpha} = [e_{\alpha}, e_{-\alpha}] \in \mathfrak{h}$, we have $\alpha(H_{\alpha}) = 2$. Then the Cartan matrix $A = (a_{ij})_{i,j=1}^n$ of \mathfrak{g} is defined by $a_{ij} = \alpha_i(H_{\alpha_j})$.

Let us recall the zero curvature representation of (1.3) following [LS79]. Let

$$(2.2) \quad \mathbf{u} = \sum_{i=1}^n u_x^i H_{\alpha_i}, \quad \epsilon = \sum_{i=1}^n e_{-\alpha_i}, \quad Y = \sum_{i=1}^n e^{\rho_i} e_{\alpha_i},$$

where as in (1.3) $\rho_i = \sum_{j=1}^n a_{ij} u^j$. Then the Toda field theory (1.3) is equivalent to the following zero curvature equation

$$(2.3) \quad [-\partial_x + \epsilon + \mathbf{u}, \partial_y + Y] = 0.$$

Now let us recall the definition of a Kostant slice $\mathfrak{s} \subset \mathfrak{g}$ [Kos63], which is used in a Drinfeld-Sokolov gauge [DS84]. Let \mathfrak{s} be a complement of $[\epsilon, \mathfrak{g}]$ in \mathfrak{g} , that is,

$$(2.4) \quad \mathfrak{g} \cong \mathfrak{s} \oplus [\epsilon, \mathfrak{g}].$$

Then by [Kos63], $\mathfrak{s} \subset \mathfrak{n}$, and $\dim(\mathfrak{s}) = n$ is equal to the rank. We call \mathfrak{s} a Kostant slice, and let $\{s_j\}_{j=1}^n$ be a homogeneous basis of \mathfrak{s} with respect to the height gradation (2.1).

By [DS84], we can bring the first element in (2.3) into its Drinfeld-Sokolov gauge. More precisely, there exists an element $g \in N$ (whose components are differential

polynomials of the u^i) such that

$$(2.5) \quad g(-\partial_x + \epsilon + \mathbf{u})g^{-1} = -\partial_x + \epsilon + \mathbf{I}, \quad \mathbf{I} = \sum_{j=1}^n I_j s_j \in \mathfrak{s},$$

where the components I_j are differential polynomials of the u^i .

Theorem 2.6. *For the solutions u^i to (1.3), the differential polynomials I_j for $1 \leq j \leq n$ defined in (2.5) through a Drinfeld-Sokolov gauge are primitive characteristic integrals, that is, $\partial_y I_j = 0$.*

Proof. Suppose that for the $g \in N$ in (2.5), we have

$$g^{-1}(\partial_y + Y)g = \partial_y + \tilde{Y}.$$

Since $g \in N$ and $Y \in \mathfrak{g}_1$ by (2.2), we have $\tilde{Y} \in \mathfrak{n}$. Suppose $\tilde{Y} = \sum_{i=1}^p Y_i$ by the height decomposition (2.1), where p is the biggest height of \mathfrak{g} .

By the invariance of the zero-curvature equation (2.3) under the adjoint action by g , from (2.5) we get

$$(2.7) \quad \left[-\partial_x + \epsilon + \sum_{j=1}^n I_j s_j, \partial_y + \sum_{i=1}^p Y_i \right] = 0.$$

We will prove by induction that all the $Y_i = 0$. Clearly this then implies that $\partial_y I_j = 0$ for $1 \leq j \leq n$.

First recall the basic result of Kostant [Kos59] that $\ker(\text{ad}_\epsilon) \cap (\mathfrak{h} \oplus \mathfrak{n}) = 0$. The term on the left of (2.7) with height zero is $[\epsilon, Y_1] = 0$, which then implies that $Y_1 = 0$.

Now assume $i \geq 2$ and that $Y_j = 0$ for $j \leq i-1$. Then since the $s_j \in \mathfrak{n}$, the term on the left of (2.7) with height $i-1$ is

$$[\epsilon, Y_i] - \sum_{\text{ht}(s_j)=i-1} (\partial_y I_j) s_j = 0.$$

By the decomposition (2.4), we get $[\epsilon, Y_i] = 0$ which then implies that $Y_i = 0$.

These I_j are primitive since without further differentiation, we are obtaining the primitive invariant functions on the Lie algebra by [Kos63]. \square

Remark 2.8. The calculations of $g \in N$ and the I_j in (2.5) can be done inductively using the decomposition (2.4) as in [DS84, FOR⁺92]. For the reader's convenience, we provide some details. Consider

$$g = e^{a_1} \cdots e^{a_m}, \quad a_i \in \mathfrak{g}_i, \quad i = 1, \dots, m.$$

The formula (2.5) becomes

$$\partial_x g \cdot g^{-1} + g(\epsilon + \mathbf{u})g^{-1} = \epsilon + \mathbf{I}.$$

Inductively for $i \geq 1$, the $i-1$ th component of the left hand side can be uniquely written as $[\epsilon, a_i] + J_{i-1}$ with $J_{i-1} \in \mathfrak{s} \cap \mathfrak{g}_{i-1}$. Then exactly this a_i will do the job of the Drinfeld-Sokolov gauging at grade $i-1$. The algorithm is easily implemented on a mathematical software such as Maple, as long as we know the structure equations of the Lie algebra. This author has worked out the example of E_6 Toda field theory by this method. In Section 4, we will see the example of D_4 .

Furthermore very often there are more direct formulas for the characteristic integrals as we explain now.

Take an irreducible representation $\phi : \mathfrak{g} \rightarrow \text{End } V$. Let the $\beta_k \in \mathfrak{h}^*$ for $1 \leq k \leq m$ be the weights of ϕ , and

$$(2.9) \quad V = \bigoplus_{k=1}^m V_{\beta_k},$$

the weight space decomposition. We assume that $\dim V_{\beta_k} = 1$ and also that the representation ϕ does not branch. That is, for each weight β_k there is one unique negative simple root $-\alpha_{i_k}$ such that $\beta_k - \alpha_{i_k}$ is another weight of ϕ . Order our weights such that $\beta_{k+1} = \beta_k - \alpha_{i_k}$ for $1 \leq k \leq m-1$ and we draw the following weight diagram

$$(2.10) \quad \beta_m \xleftarrow{-\alpha_{i_{m-1}}} \dots \xleftarrow{-\alpha_{i_2}} \beta_2 \xleftarrow{-\alpha_{i_1}} \beta_1.$$

Such non-branching representations occur for the first fundamental representations of the Lie algebras A_n, B_n, C_n and \mathfrak{g}_2 .

Theorem 2.11. *If a non-branching representation ϕ as above exists with weight diagram (2.10), then we have*

$$(2.12) \quad [(\partial_x - \beta_1(\mathbf{u}))(\partial_x - \beta_2(\mathbf{u})) \cdots (\partial_x - \beta_m(\mathbf{u})), \partial_y] = 0,$$

for a \mathbf{u} (2.2) satisfying (2.3) or equivalently the Toda field theory (1.3). The product is in the sense of composition for operators on functions of x and y , and we strictly follow the order. If by the Leibniz rule, we expand

$$(\partial_x - \beta_1(\mathbf{u}))(\partial_x - \beta_2(\mathbf{u})) \cdots (\partial_x - \beta_m(\mathbf{u})) = \partial_x^m + \sum_{j=1}^m I_j \partial_x^{m-j},$$

then (2.12) implies that

$$(2.13) \quad \partial_y I_j = 0, \quad 1 \leq j \leq m.$$

Proof. In the weight decomposition (2.9), we let $v_{\beta_1} \in V_{\beta_1}$ be a weight vector for the highest weight, and inductively we define the other weight vectors by $v_{\beta_k} = \phi(e_{-\alpha_{i_{k-1}}})v_{\beta_{k-1}}$ for $2 \leq k \leq m$ by (2.10). Therefore by (2.2)

$$(2.14) \quad \phi(\epsilon)v_{\beta_{k-1}} = v_{\beta_k}, \quad 2 \leq k \leq m.$$

The zero curvature equation (2.3) is the compatibility condition for the following system of equations. Let $\psi(x, y) = \sum_{k=1}^m \psi_k(x, y)v_{\beta_k}$ be a function of x and y with values in V . Then (2.3) implies that the following system of equations has solutions

$$(2.15) \quad \begin{cases} (-\partial_x + \phi(\mathbf{u} + \epsilon))\psi = 0 \\ (\partial_y + \phi(Y))\psi = 0. \end{cases}$$

Then by (2.14) and $\mathbf{u} \in \mathfrak{h}$ (2.2), the first equation, at the weight vector v_{β_k} , means that

$$(2.16) \quad (\partial_x - \beta_k(\mathbf{u}))\psi_k = \psi_{k-1}, \quad 2 \leq k \leq m.$$

When $k = 1$, we actually have

$$(\partial_x - \beta_1(\mathbf{u}))\psi_1 = 0,$$

since β_1 is the highest weight. Therefore combining them, we have

$$(\partial_x - \beta_1(\mathbf{u}))(\partial_x - \beta_2(\mathbf{u})) \cdots (\partial_x - \beta_m(\mathbf{u}))\psi_m = 0.$$

On the other hand, the second equation in (2.15), at the lowest weight vector v_{β_m} , quickly gives that

$$\partial_y \psi_m = 0,$$

since $Y \in \mathfrak{n}$ (2.2). Therefore the above two equations have a solution $\psi_m = \psi_m(x, y)$, which is general. The implied compatibility condition is exactly (2.12). The rest of the theorem is clear. \square

We note that again the summands of the I_j with only derivatives of order 1 again are just the algebraic invariant functions. Therefore if we know the degrees of the Lie algebras, by the structure theorem we know the primitive characteristic integrals.

3. EXAMPLES FOR THEOREM 2.11

The first fundamental representations of the Lie algebras A_n, B_n, C_n and \mathfrak{g}_2 are non-branching representations (2.10). Keeping our spirit of being as explicit as possible, we present the formulas in these cases. In this section we follow [FH91] for notation and choices of root vectors. The Cartan subalgebras \mathfrak{h} always consist of diagonal matrices. We let $L_i \in \mathfrak{h}^*$ denote the linear function of taking the i th element on the diagonal. We also let E_{ij} denote the matrix with a 1 at the (i, j) -position and zero everywhere else. For simplicity, we write ∂ for ∂_x .

Example 3.1 (A_n). The simple roots for A_n are $\alpha_i = L_i - L_{i+1}$ and the $H_{\alpha_i} = E_{i,i} - E_{i+1,i+1}$, $1 \leq i \leq n$. The Cartan matrix is

$$\begin{pmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{pmatrix}$$

The degrees of A_n (that is, the degrees of primitive adjoint-invariant functions) are $2, 3, \dots, n+1$. The weight diagram for the first fundamental representation is

$$L_{n+1} \xleftarrow{\alpha_n} \cdots \xleftarrow{\alpha_2} L_2 \xleftarrow{\alpha_1} L_1.$$

The \mathbf{u} in (2.2) is

$$\mathbf{u} = \text{Diag}(u_x^1, u_x^2 - u_x^1, \dots, u_x^n - u_x^{n-1}, -u_x^n)$$

By Theorem 2.11, consider the expansion

$$\begin{aligned} & (\partial - u_x^1)(\partial + u_x^1 - u_x^2) \cdots (\partial + u_x^{n-1} - u_x^n)(\partial + u_x^n) \\ (3.2) \quad & = \partial^{n+1} + \sum_{j=1}^n I_j \partial^{n-j}. \end{aligned}$$

Then the I_j for $1 \leq j \leq n$ are primitive characteristic integrals of the A_n Toda field theory (1.3).

Example 3.3 ($C_n, n \geq 2$). The simple roots for C_n are $\alpha_i = L_i - L_{i+1}$ for $1 \leq i \leq n-1$ and $\alpha_n = 2L_n$. Also $H_{\alpha_i} = E_{i,i} - E_{i+1,i+1} - E_{n+i,n+i} + E_{n+i+1,n+i+1}$ for $1 \leq i \leq n-1$ and $H_{\alpha_n} = E_{n,n} - E_{2n,2n}$. The Cartan matrix is

$$\begin{pmatrix} 2 & -1 & & & & \\ -1 & 2 & -1 & & & \\ & & \ddots & \ddots & \ddots & \\ & & & -1 & 2 & -1 \\ & & & & -2 & 2 \end{pmatrix}$$

The degrees of C_n are $2, 4, \dots, 2n$. The weight diagram for the first fundamental representation is

$$-L_1 \xleftarrow{-\alpha_1} \dots \xleftarrow{-\alpha_{n-1}} -L_n \xleftarrow{-\alpha_n} L_n \xleftarrow{-\alpha_{n-1}} \dots \xleftarrow{-\alpha_2} L_2 \xleftarrow{-\alpha_1} L_1.$$

The \mathbf{u} in (2.2) is

$$\mathbf{u} = \text{Diag}(u_x^1, u_x^2 - u_x^1, \dots, u_x^n - u_x^{n-1}, -u_x^1, -u_x^2 + u_x^1, \dots, -u_x^n + u_x^{n-1}).$$

By Theorem 2.11, consider the expansion

$$\begin{aligned} & (\partial - u_x^1)(\partial + u_x^1 - u_x^2) \cdots (\partial + u_x^{n-1} - u_x^n) \\ & (\partial + u_x^n - u_x^{n-1}) \cdots (\partial + u_x^2 - u_x^1)(\partial + u_x^1) \\ (3.4) \quad & = \partial^{2n} + \sum_{j=1}^n I_j \partial^{2n-2j} + \sum_{j=1}^{n-1} J_j \partial^{2n-2j-1}. \end{aligned}$$

Then the I_j for $1 \leq j \leq n$ are primitive characteristic integrals of the C_n Toda field theory, and the J_j are some differential polynomials in them.

Example 3.5 ($B_n, n \geq 2$). The simple roots for B_n are $\alpha_i = L_i - L_{i+1}$ for $1 \leq i \leq n-1$ and $\alpha_n = L_n$. Also $H_{\alpha_i} = E_{i,i} - E_{i+1,i+1} - E_{n+i,n+i} + E_{n+i+1,n+i+1}$ for $1 \leq i \leq n-1$ and $H_{\alpha_n} = 2E_{n,n} - 2E_{2n,2n}$. The Cartan matrix is

$$\begin{pmatrix} 2 & -1 & & & & \\ -1 & 2 & -1 & & & \\ & & \ddots & \ddots & \ddots & \\ & & & -1 & 2 & -2 \\ & & & & -1 & 2 \end{pmatrix}$$

The degrees of B_n are $2, 4, \dots, 2n$. The weight diagram for the first fundamental representation is

$$(3.6) \quad -L_1 \xleftarrow{-\alpha_1} \dots \xleftarrow{-\alpha_{n-1}} -L_n \xleftarrow{-\alpha_n} 0 \xleftarrow{-\alpha_n} L_n \xleftarrow{-\alpha_{n-1}} \dots \xleftarrow{-\alpha_2} L_2 \xleftarrow{-\alpha_1} L_1.$$

The \mathbf{u} in (2.2) is

$$\mathbf{u} = \text{Diag}(u_x^1, u_x^2 - u_x^1, \dots, 2u_x^n - u_x^{n-1}, -u_x^1, -u_x^2 + u_x^1, \dots, -2u_x^n + u_x^{n-1}, 0).$$

By Theorem 2.11, consider the expansion

$$\begin{aligned} & (\partial - u_x^1)(\partial + u_x^1 - u_x^2) \cdots (\partial + u_x^{n-1} - 2u_x^n) \\ & \partial(\partial + 2u_x^n - u_x^{n-1}) \cdots (\partial + u_x^2 - u_x^1)(\partial + u_x^1) \\ (3.7) \quad & = \partial^{2n+1} + \sum_{j=1}^n I_j \partial^{2n-2j+1} + \sum_{j=1}^n J_j \partial^{2n-2j}. \end{aligned}$$

Then the I_j for $1 \leq j \leq n$ are primitive characteristic integrals of the B_n Toda field theory, and the J_j are some differential polynomials in them.

Example 3.8 (\mathfrak{g}_2). The Cartan matrix is

$$\begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$$

For simplicity we call the unknown functions u^1, u^2 by u and v . The Toda system is

$$\begin{cases} u_{xy} = -e^{2u-v} \\ v_{xy} = -e^{-3u+2v} \end{cases}$$

Although we can work abstractly with just the above Cartan matrix, we choose to use the embedding of $\mathfrak{g}_2 \subset \mathfrak{so}_7 = B_3$ such that the two roots are

$$\alpha_1 = L_1 - L_2, \quad \alpha_2 := L_2 - L_3,$$

and

$$\begin{aligned} H_{\alpha_1} &= \text{Diag}(1, -1, 2, -1, 1, -2, 0) \\ H_{\alpha_2} &= \text{Diag}(0, 1, -1, 0, -1, 1, 0). \end{aligned}$$

The \mathbf{u} in (2.2) is

$$\mathbf{u} = \text{Diag}(u_x, -u_x + v_x, 2u_x - v_x, -u_x, u_x - v_x, -2u_x + v_x, 0).$$

The first fundamental representation of \mathfrak{g}_2 is the restriction of that of B_3 . Therefore we follow the weight diagram (3.6). The degrees of \mathfrak{g}_2 are 2 and 6.

By Theorem 2.11, consider the expansion

$$\begin{aligned} & (\partial - u_x)(\partial + u_x - v_x)(\partial - 2u_x + v_x) \\ & \quad \partial(\partial + 2u_x - v_x)(\partial - u_x + v_x)(\partial + u_x) \\ (3.9) \quad & = \partial^7 + I_1 \partial^5 + I_2 \partial + \sum_{j=1}^3 J_j \partial^{5-j} + J_4. \end{aligned}$$

Then the I_1 and I_2 are primitive characteristic integrals of the \mathfrak{g}_2 Toda field theory, and the J_j are some differential polynomials in them.

4. AN EXAMPLE OF THEOREM 2.6

It can be said that Theorem 2.11 is a quick way to compute a Drinfeld-Sokolov gauge (2.5) when there is a non-branching representation, and this is the viewpoint in [BFO⁺90]. When such a simple representation does not exist, a Drinfeld-Sokolov gauge can still be computed through an inductive procedure (see [DS84, FOR⁺92]). In the particular case of D_n , where an explicit matrix presentation of the Lie algebra is easy to write down, one can use a computer algebra system like Maple to solve a Drinfeld-Sokolov gauge easily and therefore obtain the characteristic integrals according to Theorem 2.6.

As stated in Remark 2.8, the Drinfeld-Sokolov gauging can be achieved without using a representation. Intrinsically one can use the adjoint representation constructed from the structure equations of the Lie algebra. This author has done this for the E_6 Toda field theory.

REFERENCES

- [AFV09] Ian M. Anderson, Mark E. Fels, and Peter J. Vassiliou, *Superposition formulas for exterior differential systems*, Adv. Math. **221** (2009), no. 6, 1910–1963.
- [BBT03] Olivier Babelon, Denis Bernard, and Michel Talon, *Introduction to classical integrable systems*, Cambridge Monographs on Mathematical Physics, Cambridge University Press, Cambridge, 2003.
- [BFO⁺90] J. Balog, L. Fehér, L. O’Raifeartaigh, P. Forgács, and A. Wipf, *Toda theory and W-algebra from a gauged WZNW point of view*, Ann. Physics **203** (1990), no. 1, 76–136.
- [Dem10] D. K. Demskoi, *Integrals of open two-dimensional lattices*, Theoret. and Math. Phys. **163** (2010), no. 1, 466–471.
- [DS84] V. G. Drinfel’d and V. V. Sokolov, *Lie algebras and equations of Korteweg-de Vries type*, Current problems in mathematics, Vol. 24, Itogi Nauki i Tekhniki, Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1984, pp. 81–180 (Russian).
- [FOR⁺92] L. Fehér, L. O’Raifeartaigh, P. Ruelle, I. Tsutsui, and A. Wipf, *Generalized Toda theories and W-algebras associated with integral gradings*, Ann. Physics **213** (1992), no. 1, 1–20.
- [FF96] Boris Feigin and Edward Frenkel, *Integrals of motion and quantum groups*, Integrable systems and quantum groups (Montecatini Terme, 1993), Lecture Notes in Math., vol. 1620, Springer, Berlin, 1996, pp. 349–418.
- [FH91] William Fulton and Joe Harris, *Representation theory*, Graduate Texts in Mathematics, vol. 129, Springer-Verlag, New York, 1991. A first course; Readings in Mathematics.
- [GZ04] A. M. Gur’eva and A. V. Zhiber, *Laplace invariants of two-dimensionalized open Toda chains*, Teoret. Mat. Fiz. **138** (2004), no. 3, 401–421 (Russian, with Russian summary); English transl., Theoret. and Math. Phys. **138** (2004), no. 3, 338–355.
- [GZ05] ———, *Laplace invariants of Toda lattices with exceptional Cartan matrices* (2005), available at <http://arxiv.org/abs/nlin/0512001>.
- [Kos59] Bertram Kostant, *The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group*, Amer. J. Math. **81** (1959), 973–1032.
- [Kos63] ———, *Lie group representations on polynomial rings*, Amer. J. Math. **85** (1963), 327–404.
- [HO96] Erling G. B. Hohler and Kåre Olaussen, *Using conservation laws to solve Toda field theories*, Internat. J. Modern Phys. A **11** (1996), no. 10, 1831–1853.
- [LS79] A. N. Leznov and M. V. Saveliev, *Representation of zero curvature for the system of nonlinear partial differential equations $x_{\alpha, z\bar{z}} = \exp(kx)_{\alpha}$ and its integrability*, Lett. Math. Phys. **3** (1979), no. 6, 489–494.
- [LS92] ———, *Group-theoretical methods for integration of nonlinear dynamical systems*, Progress in Physics, vol. 15, Birkhäuser Verlag, Basel, 1992. Translated and revised from the Russian; Translated by D. A. Leuities.
- [MB08] G. Mari Beffa, *Geometric Hamiltonian structures on flat semisimple homogeneous manifolds*, Asian J. Math. **12** (2008), no. 1, 1–33.
- [Sha95] A. B. Shabat, *Higher symmetries of two-dimensional lattices*, Phys. Lett. A **200** (1995), no. 2, 121–133.
- [SY81] A. B. Shabat and R. L. Yamilov, *Exponential systems of type II and Cartan matrices* (1981), Preprint No. 1 Bashkir Branch USSR Acad. Sci. Ufa.
- [Yam93] Keizo Yamaguchi, *Differential systems associated with simple graded Lie algebras*, Progress in differential geometry, Adv. Stud. Pure Math., vol. 22, Math. Soc. Japan, Tokyo, 1993, pp. 413–494.
- [ZS01] A. V. Zhiber and V. V. Sokolov, *Exactly integrable hyperbolic equations of Liouville type*, Uspekhi Mat. Nauk **56** (2001), no. 1(337), 63–106 (Russian, with Russian summary); English transl., Russian Math. Surveys **56** (2001), no. 1, 61–101.

E-mail address: zhaohu.nie@usu.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, UTAH STATE UNIVERSITY, LOGAN, UT 84322-3900