

# Hamilton's Harnack inequality and the $W$ -entropy formula on complete Riemannian manifolds

Xiang-Dong Li \*

October 3, 2018

**Abstract.** In this paper, we prove Hamilton's Harnack inequality and the gradient estimates of the logarithmic heat kernel for the Witten Laplacian on complete Riemannian manifolds. As applications, we prove the  $W$ -entropy formula for the Witten Laplacian on complete Riemannian manifolds, and prove a family of logarithmic Sobolev inequalities on complete Riemannian manifolds with natural geometric condition.

## 1 Introduction

Differential Harnack inequality is an important tool in the study of heat equation on complete Riemannian manifolds [19, 10, 20, 39, 8, 9]. In [16], Hamilton proved a Harnack inequality for positive and bounded solutions to the heat equation on compact Riemannian manifolds. More precisely, let  $M$  be a complete Riemannian manifold, and suppose that there exists a constant  $K \geq 0$  such that

$$Ric \geq -K.$$

Then, for any positive and bounded solution  $u$  to the heat equation

$$\partial_t u = \Delta u,$$

it holds

$$|\nabla \log u|^2 \leq \left( \frac{1}{t} + 2K \right) \log(A/u), \quad \forall x \in M, t > 0,$$

where

$$A := \sup\{u(t, x) : x \in M, t \geq 0\}.$$

In [18], Kotschwar extended Hamilton's Harnack inequality to complete Riemannian manifolds with non-negative Ricci curvature. As application, he proved the following gradient estimates of the logarithmic of the heat kernel  $p_t(x, y)$  of the Laplace-Beltrami operator on complete Riemannian manifolds with non-negative Ricci curvature. More precisely, for all  $\delta > 0$ , there exists a constant  $C = C(n, \delta) > 0$ , such that

$$|\nabla \log p_t(x, y)|^2 \leq \frac{2}{t} \left( C + \frac{d^2(x, y)}{(4 - \delta)t} \right), \quad \forall x, y \in M, t > 0.$$

The purpose of this paper is to prove Hamilton's Harnack inequality and gradient estimates of the logarithmic of the heat kernel of the Witten Laplacian on complete Riemannian

---

\*Research supported by NSFC No. 11371351, Key Laboratory RCDSDS, CAS, No. 2008DP173182, and a Hundred Talents Project of AMSS, CAS.

manifolds with weighted volume measure. As application, we derive the  $W$ -entropy formula and prove the rigidity theorem of the  $W$ -entropy for the Witten Laplacian on complete Riemannian manifolds with natural Bakry-Emery curvature condition. Indeed, the relation between the  $W$ -entropy functional and the differential Harnack inequality has been recently investigated by many authors [35, 33, 34, 9, 12, 26].

Let  $M$  be a complete Riemannian manifold with a fixed Riemannian metric  $g$ ,  $\phi \in C^2(M)$  and  $d\mu = e^{-\phi}dv$ , where  $v$  is the Riemannian volume measure on  $(M, g)$ , and let

$$L = \Delta - \nabla\phi \cdot \nabla$$

be the Witten Laplacian on  $(M, g)$  with respect to the weighted volume measure  $\mu$ . For all  $u, v \in C_0^\infty(M)$ , the following integration by parts formula holds

$$\int_M \langle \nabla u, \nabla v \rangle d\mu = - \int_M Luv d\mu = - \int_M uLv d\mu.$$

In [5], Bakry and Emery proved that for all  $u \in C_0^\infty(M)$ ,

$$L|\nabla u|^2 - 2\langle \nabla u, \nabla Lu \rangle = 2|\nabla^2 u|^2 + 2Ric(L)(\nabla u, \nabla u), \quad (1)$$

where

$$Ric(L) = Ric + \nabla^2\phi.$$

The formula (1) can be viewed as a natural extension of the Bochner-Weitzenböck formula. The quantity  $Ric(L) = Ric + \nabla^2\phi$ , called the Bakry-Emery Ricci curvature on the weighted Riemannian manifolds  $(M, g, \phi)$ , plays as a good substitute of the Ricci curvature in many problems in comparison geometry and analysis on complete Riemannian manifolds with smooth weighted volume measures. See [5, 14, 15, 20, 23, 24, 32, 42, 30, 31] and reference therein.

Following [5, 27, 20], we introduce the  $m$ -dimensional Bakry-Emery Ricci curvature on  $(M, g, \phi)$  by

$$Ric_{m,n}(L) := Ric + \nabla^2\phi - \frac{\nabla\phi \otimes \nabla\phi}{m-n},$$

where  $m \geq n$  is a constant, and  $m = n$  if and only if  $\phi$  is a constant. When  $m = \infty$ , we have  $Ric_{\infty,n}(L) = Ric(L)$ . When  $m \in \mathbb{N}$ , the  $m$ -dimensional Bakry-Emery Ricci curvature  $Ric_{m,n}(L)$  has a very natural geometric interpretation. Indeed, consider the warped product metric on  $M^n \times S^{m-n}$  defined by

$$\tilde{g} = g_M \oplus e^{-\frac{2\phi}{m-n}} g_{S^{m-n}}.$$

where  $S^{m-n}$  is the unit sphere in  $\mathbb{R}^{m-n+1}$  with the standard metric  $g_{S^{m-n}}$ . By a classical result in Riemannian geometry, the quantity  $Ric_{m,n}(L)$  is equal to the Ricci curvature of the above warped product metric  $\tilde{g}$  on  $M^n \times S^{m-n}$  along the horizontal vector fields. See [27, 20, 42].

We now state the main results of this paper. The first result of this paper is the following improved version of the Harnack inequality for the heat equation of the Witten Laplacian on complete Riemannian manifolds.

**Theorem 1.1** *Let  $M$  be a complete Riemannian manifold, and  $\phi \in C^2(M)$ . Suppose that there exists a constant  $K \geq 0$  such that*

$$Ric(L) \geq -K.$$

Let  $u$  be a positive and bounded solution to the heat equation

$$\partial_t u = Lu.$$

Let  $A := \sup\{u(t, x) : x \in M, t \geq 0\}$ . Then

$$|\nabla \log u|^2 \leq \frac{2K}{1 - e^{-2Kt}} \log(A/u). \quad (2)$$

The following result extends the Hamilton Harnack inequality to the Witten Laplacian on complete Riemannian manifolds.

**Theorem 1.2** *Under the same condition as in Theorem 1.1, the Halmilton Harnack inequality holds*

$$|\nabla \log u|^2 \leq \left(\frac{1}{t} + 2K\right) \log(A/u). \quad (3)$$

Letting  $t \rightarrow \infty$  in Theorem 1.2, we can derive the following gradient estimate and the Liouville theorem due to Brighton [7]. See also Remark 2.3 below.

**Corollary 1.3** *Let  $u$  be a bounded and positive  $L$ -harmonic function  $u$  on  $M$ . Then, under the same condition as in Theorem 1.1, we have*

$$|\nabla \log u| \leq 2K \log(A/(u - \inf u)),$$

where  $A = \sup u - \inf u$ . In particular, if  $\text{Ric} + \nabla^2 \phi \geq 0$ , then every bounded  $L$ -harmonic function must be constant.

Let  $p_t(x, y)$  be the heat kernel of  $L$  with respect to the reference measure  $\mu$ , i.e., the fundamental solution of the heat equation  $\partial_t u = Lu$ . The following result extends Hamilton and Kotschwar's gradient estimate to the logarithmic heat kernel of the Witten Laplacian on complete Riemannian manifolds with natural geometric condition.

**Theorem 1.4** *Let  $M$  be a complete Riemannian manifolds, and  $\phi \in C^2(M)$ . Suppose that there exist some constant  $m \geq n$  and  $K \geq 0$  such that*

$$\text{Ric}_{m,n}(L) \geq -K.$$

Then, for all  $T > 0$ , there exists a constant  $C(K, m, n, T) > 0$  such that

$$|\nabla \log p_t(x, y)| \leq C(K, m, n, T) \left( \frac{d(x, y)}{t} + \frac{1}{\sqrt{t}} \right)$$

holds for all  $x, y \in M$  and  $t \in (0, T]$ .

The following result gives the higher order gradient estimates for the logarithmic heat kernel of the Witten Laplacian on complete Riemannian manifolds with natural geometric condition.

**Theorem 1.5** *Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold. Suppose that  $\|\nabla^k \text{Riem}\| \leq C_k$  for some constant  $C_k > 0$ ,  $0 \leq k \leq N$ ,  $\phi \in C^{N+1}(M)$  with  $\nabla \phi \in C_b^N(M)$ , and there exists some  $m \geq n$ ,  $K \geq 0$  such that  $\text{Ric}_{m,n}(L) \geq -K$ . Then, for all  $T > 0$ , there exists a constant  $C_N(K, m, n, T) > 0$  such that*

$$|\nabla^N \log p_t(x, y)| \leq C_N(K, m, n, T) \left( \frac{d(x, y)}{t} + \frac{1}{\sqrt{t}} \right)^N$$

holds for all  $x, y \in M$  and  $t \in (0, T]$ .

As an application of Theorem 1.4 and Theorem 1.5, we prove the following  $W$ -entropy formula for the Witten Laplacian on complete Riemannian manifolds, which extends a previous result due to Ni [33, 34] for the case  $m = n$  and  $L = \Delta$ .

**Theorem 1.6** *Let  $(M, g)$  be a complete Riemannian manifold. Suppose that  $\|\nabla^k \text{Riem}\| \leq C_k$  for some constant  $C_k > 0$ ,  $0 \leq k \leq 2$ ,  $\phi \in C^4(M)$  with  $\nabla \phi \in C_b^3(M)$ , and there exists some  $m \geq n$ ,  $K \geq 0$  such that  $\text{Ric}_{m,n}(L) \geq -K$ . Let  $u(t, x) = \frac{e^{-f}}{(4\pi t)^{m/2}}$  be the heat kernel of the heat equation  $\frac{\partial u}{\partial t} = Lu$ . Let*

$$W_m(u, t) = \int_M (t|\nabla f|^2 + f - m) \frac{e^{-f}}{(4\pi t)^{m/2}} d\mu$$

be the  $W$ -entropy of the Witten Laplacian  $L$  on  $(M, g, \mu)$ . Then, for all  $t > 0$ , we have

$$\begin{aligned} \frac{dW_m(u, t)}{dt} &= -2 \int_M t \left( \left| \nabla^2 f - \frac{g}{2t} \right|^2 + \text{Ric}_{m,n}(L)(\nabla f, \nabla f) \right) u d\mu \\ &\quad - \frac{2}{m-n} \int_M t \left( \nabla \phi \cdot \nabla f + \frac{m-n}{2t} \right)^2 u d\mu. \end{aligned} \quad (4)$$

In particular, if  $\text{Ric}_{m,n}(L) \geq 0$ , then  $W_m(u, t)$  is non-increasing in  $t$ , and  $W_m(u, t)$  attains its minimum at some point  $t_0 > 0$ , i.e.,  $\frac{dW_m(u, t)}{dt} = 0$  holds at  $t = t_0$ , if and only if  $(M, g)$  is isometric to Euclidean space  $\mathbb{R}^n$ ,  $m = n$ ,  $\phi \equiv C$  for a constant  $C \in \mathbb{R}$ , and

$$u(t, x) = \frac{e^{-\frac{\|x\|^2}{4t}}}{(4\pi t)^{n/2}}, \quad \forall x \in \mathbb{R}^n, t > 0.$$

In our previous paper [23], we proved the  $W$ -entropy formula (4) for the Witten Laplacian on complete Riemannian manifolds satisfying the condition in Theorem 1.6 and the following uniform volume lower bound condition: there exists a constant  $r_0 > 0$  such that

$$\mu_0 = \inf\{\mu(B(x, r_0)) : x \in M\} > 0. \quad (5)$$

By Theorem 2.8 in [22], on complete Riemannian manifolds with  $\text{Ric}_{m,n}(L) \geq -K$  and satisfying the uniformly lower bound condition (5), the following Sobolev inequality holds

$$\|f\|_{\frac{2m}{m-2}}^2 \leq C \left( \int_M |\nabla f|^2 d\mu + \int_M |f|^2 d\mu \right), \quad \forall f \in C_0^\infty(M).$$

Thus, at least from a geometric and analytic point of view, the uniformly volume lower bound condition (5) is too restrictive for the validity of the  $W$ -entropy formula (4) for the Witten Laplacian on complete Riemannian manifolds. Our new observation in Theorem 1.6 here is that we can remove the uniform volume lower bound condition (5) to establish the  $W$ -entropy formula (4). This is essentially based on the fact that Theorem 1.4 and Theorem 1.5 hold on complete Riemannian manifolds on which we need not to assume the uniform volume lower bound condition (5). Indeed, just after the paper [23] was accepted, we observed that the uniform volume lower bound condition (5) can be removed. See the Note added in proof of [23].

As an application of Theorem 1.6, we prove the following result, which indicates that the uniform volume lower bound condition (5) allows us to derive a family of logarithmic Sobolev inequalities on complete Riemannian manifolds.

**Theorem 1.7** *Let  $(M, g, \phi)$  be a weighted complete Riemannian manifold as in Theorem 1.6 and satisfying the uniform volume lower bound condition (5). Then, for any  $\tau > 0$ , the*

following logarithmic Sobolev inequality holds: for all  $v \in W^{1,2}(M, \mu)$  with  $\int_M v^2 d\mu = 1$ , we have

$$\int_M v^2 \log v^2 d\mu \leq 4\tau \int_M |\nabla v|^2 - m \left( 1 + \frac{1}{2} \log(4\pi\tau) \right) - \mu(\tau),$$

where

$$\mu(\tau) := \inf \left\{ W_m(u, \tau) : u = \frac{e^{-f}}{(4\pi\tau)^{m/2}}, \int_M u d\mu = 1 \right\}$$

is a finite number.

**Remark 1.8** Theorem 1.4 and Theorem 1.5 are improved versions of Theorem 4.1 and Theorem 4.2 obtained in [23], where the uniform volume lower bound condition (5) is assumed. When  $M$  is a compact Riemannian manifold and  $\phi$  is a constant, Theorem 1.4 and Theorem 1.5 are due to Hamilton [16], Sheu [36], Hsu [17] and Stroock-Turesky [38]. In the case where  $L = \Delta$  is the Laplace-Beltrami operator on complete Riemannian manifolds with bounded geometry condition (in particular, the uniform volume lower bound condition (5) holds for  $\mu = \nu$ ) and for small time  $t > 0$ , Theorem 1.4 is due to Engoulatov [13]. In [18, 11], the authors proved the gradient estimates in Theorem 1.4 for  $L = \Delta$  on complete Riemannian manifolds with non-negative Ricci curvature.

**Remark 1.9** In [25], Songzi Li and the author gave a new proof of the  $W$ -entropy formula on Riemannian manifolds by using the warped metric product

$$\tilde{g} = g_M \oplus e^{-\frac{2\phi}{m-n}} g_N.$$

on  $M \times N$ , where  $(N, g_N)$  is any fixed compact Riemannian manifold of dimension  $m - n$ . This gives a natural geometric interpretation of the third term appeared in (4). See Remark 2.2 in [25]. Moreover, we extended the  $W$ -entropy formula (4) to the heat equation of the time dependent Witten Laplacian on compact Riemannian manifolds on which the metrics  $\{g(t), t \in [0, T]\}$  and the potential functions  $\{\phi(t), t \in [0, T]\}$  evolve along the  $m$ -dimensional Perelman Ricci flow and the conjugate heat equation

$$\begin{aligned} \frac{\partial g}{\partial t} &= -2 \left( Ric + \nabla^2 \phi - \frac{\nabla \phi \otimes \nabla \phi}{m-n} \right), \\ \frac{\partial \phi}{\partial t} &= -R - \Delta \phi + \frac{|\nabla \phi|^2}{m-n}. \end{aligned}$$

In this case, without assuming  $Ric_{m,n}(L) \geq 0$ , we prove in [25] that the  $W$ -entropy for the positive solution of the forward heat equation

$$\partial_t u = Lu$$

of the time dependent Witten Laplacian  $L = \Delta - \nabla \phi \cdot \nabla$ , is non-increasing in time  $t \in [0, T]$ .

The rest of this paper is organized as follows. In Section 2, we prove the Hamilton type Harnack inequalities (Theorem 1.1 and Theorem 1.2) for the Witten Laplacian on complete Riemannian manifolds. In Section 3, we use Theorem 1.2 to prove the first order gradient estimate of the logarithmic heat kernel of the Witten Laplacian, i.e., Theorem 1.4. In Section 4, we use a probabilistic approach to prove the gradient estimates of the logarithmic heat kernel of the Witten Laplacian, i.e., Theorem 1.4 and Theorem 1.5. In Section 4, we prove the  $W$ -entropy formula and derive the monotonicity and rigidity theorem of the  $W$ -entropy for the Witten Laplacian on complete Riemannian manifolds, i.e., Theorem 1.6. In Section 5, we prove a family of logarithmic Sobolev inequalities on complete Riemannian manifolds with the uniform volume lower bound condition, i.e., Theorem 1.7.

## 2 Hamilton's Harnack inequality

In this section we give two proofs of Theorem 1.1 and Theorem 1.2.

**Direct Proof of Theorem 1.1.** Similarly to [16], a direct calculation and the generalized Bochner formula yield

$$\begin{aligned} (\partial_t - L) \frac{|\nabla u|^2}{u} &= -\frac{2}{u} |\nabla^2 u - u^{-1} \nabla u \otimes \nabla u|^2 - 2u^{-1} (\text{Ric} + \nabla^2 \phi)(\nabla u, \nabla u) \\ &\leq 2Ku^{-1} |\nabla u|^2. \end{aligned}$$

Moreover

$$(\partial_t - L)(u \log(A/u)) = \frac{|\nabla u|^2}{u}.$$

Let  $\psi(t) = \frac{1-e^{-2Kt}}{2K}$ . Then

$$\psi'(t) + 2K\psi(t) = 1.$$

Define

$$h(x, t) := \psi(t) \frac{|\nabla u|^2}{u} - u \log(A/u).$$

We have

$$\begin{aligned} (\partial_t - L)h &= \psi'(t) \frac{|\nabla u|^2}{u} + \psi(t) (\partial_t - L) \frac{|\nabla u|^2}{u} - (\partial_t - L)u \log(A/u) \\ &\leq \psi'(t) \frac{|\nabla u|^2}{u} + 2K\psi(t) \frac{|\nabla u|^2}{u} - \frac{|\nabla u|^2}{u} \\ &= 0. \end{aligned}$$

Note that

$$h(x, 0) = -u \log(A/u)(x, 0) \leq 0.$$

In the case  $M$  is compact, the maximum principle yields that  $h(x, t) \leq 0$  for all time  $t > 0$  and  $x \in M$ . In the case  $M$  is a complete non-compact Riemannian manifold with  $\text{Ric}(L) \geq -K$ , we can give a probabilistic proof to (2) as follows. Indeed, on any complete Riemannian manifold  $M$  with  $\text{Ric}(L) \geq -K$ , a previous result due to Bakry [3] says that the  $L$ -diffusion process on  $M$  has infinity lifetime. Let  $X_t$  be the  $L$ -diffusion process on  $M$  starting from  $X_0 = x$ . Applying Itô's formula to  $h(X_t, T-t)$ ,  $t \in [0, T]$ , we have

$$h(X_t, T-t) = h(X_0, T) + \int_0^t \nabla h(X_s, T-s) \cdot dW_s + \int_0^t \left( L - \frac{\partial}{\partial t} \right) h(X_s, T-s) ds,$$

where the second term in the right hand side is the Itô's stochastic integral with respect to a  $M$ -valued Brownian motion  $\{W_s, s \in [0, t]\}$ . In particular, taking  $t = T$ , we obtain

$$h(X_T, 0) = h(X_0, T) + \int_0^T \nabla h(X_s, T-s) \cdot dW_s + \int_0^T \left( L - \frac{\partial}{\partial t} \right) h(X_s, T-s) ds.$$

Taking the expectation on both sides, the martingale property of Itô's integral implies that

$$E[h(X_T, 0)] = h(x, T) + E \left[ \int_0^T \left( L - \frac{\partial}{\partial t} \right) h(X_s, T-s) ds \right] \geq h(x, T).$$

As  $h(y, 0) \leq 0$  for all  $y \in M$ , we derive that  $h(x, T) \leq 0$  for all  $T > 0$  and  $x \in M$ .  $\square$

**Alternative proof of Theorem 1.1.** By [4, 6], we know that  $Ric(L) \geq -K$  if and only the following version of logarithmic Sobolev inequality holds: for all  $T > 0$ ,  $f \in C_b(M)$  with  $f > 0$ ,

$$\frac{|\nabla P_T f|^2}{P_T f} \leq \frac{2K}{1 - e^{-2KT}} (P_T(f \log f) - P_T f \log P_T f). \quad (6)$$

Replacing  $f$  by  $P_t f$  and using the fact that  $0 < P_t f(x) \leq A$  for all  $x \in M$ , one has

$$\frac{|\nabla P_{T+t} f|^2}{P_{T+t} f} \leq \frac{2K}{1 - e^{-2KT}} (P_T(P_t f \log A) - P_{T+t} f \log P_{T+t} f).$$

Thus

$$|\nabla \log P_{T+t} f|^2 \leq \frac{2K}{1 - e^{-2KT}} \log(A/P_{T+t} f), \quad \forall t > 0.$$

Taking  $t \rightarrow 0^+$  we have

$$|\nabla \log P_T f|^2 \leq \frac{2K}{1 - e^{-2KT}} \log(A/P_T f).$$

This finishes the proof of Theorem 1.1.  $\square$

**Proof of Theorem 1.2.** We can derive the Hamilton Harnack inequality (3) from the Harnack inequality (2) in Theorem 1.1. More precisely, using the inequality  $\frac{2K}{1 - e^{-2KT}} \leq 2K + \frac{1}{T}$ , we have

$$|\nabla \log P_T f|^2 \leq \left(2K + \frac{1}{T}\right) \log(A/P_T f).$$

Note that, if we follow Hamilton [16] to use the function  $\psi(t) = \frac{t}{2Kt+1}$  (instead of  $\psi(t) = \frac{1 - e^{-2Kt}}{2K}$ ), then  $\psi'(t) + 2K\psi(t) \leq 1$ . By the same argument as in the direct proof of Theorem 1.1, we can give a direct proof of the Harnack inequality (3).  $\square$

**Remark 2.1** As was pointed out in Section 1, Theorem 1.2 was first proved by Hamilton [16] for the heat equation of the Laplace-Beltrami operator on compact Riemannian manifolds and was extended to complete Riemannian manifolds with Ricci curvature bounded from below by Kotschwar [18]. For a probabilistic proof of Hamilton's Harnack inequality for  $L = \Delta$  on compact or complete Riemannian manifolds, see [1]. The local version of Theorem 1.1 for the Laplace-Beltrami operator was proved by Souplet-Zhang [37]. In [2] (Theorem 5.1), Arnaudon, Tahlmaier and Wang extended the local estimate of Souplet-Zhang [37] to the heat equation of the Laplacian with general drift  $L = \Delta + Z$ , where  $Z$  is not necessary of gradient form  $Z = \nabla\phi$  and the Laplacian comparison theorem for  $L = \Delta + Z$  is essentially used. Taking  $R \rightarrow \infty$  in [2], we can derive the following estimate for the bounded and positive solution of the heat equation  $\partial_t u = Lu$ : there exists a constant  $C > 0$ , which depends only on  $n = \dim M$ , such that for all  $x \in M$  and  $t > 0$ , it holds

$$|\nabla \log u|^2 \leq C \left(\frac{1}{t} + K\right) [1 + \log(A/u)]. \quad (7)$$

where  $A := \sup\{u(t, x) : x \in M, t \geq 0\}$ .

As a corollary of the above theorem, we can derive the following gradient estimate and the Liouville theorem due to Brighton [7] for bounded  $L$ -harmonic functions.

**Corollary 2.2** (i.e., Corollary 1.3)) Suppose that  $\text{Ric}(L) = \text{Ric} + \nabla^2\phi \geq -K$  for some constant  $K \geq 0$ . Then every bounded  $L$ -harmonic function  $u$  satisfies

$$|\nabla \log u| \leq 2K \log(A/(u - \inf u)), \quad (8)$$

where  $A = \sup u - \inf u$ . In particular, if  $\text{Ric}(L) = \text{Ric} + \nabla^2\phi \geq 0$ , then every bounded  $L$ -harmonic function must be constant.

*Proof.* Applying (3) to  $u(t, x) = u(x) - \inf u$ ,  $x \in M$ ,  $t > 0$ , we have

$$|\nabla \log u(x)|^2 \leq \left(2K + \frac{1}{T}\right) \log(A/(u(x) - \inf u)), \quad \forall T > 0.$$

Taking  $T \rightarrow \infty$ , we derive (8). In particular, if  $K = 0$ , we have  $|\nabla \log u| = 0$ . This finishes the proof of Corollary 2.2.  $\square$

**Remark 2.3** In the case  $L = \Delta$ , Yau [40] proved that, on complete Riemannian manifolds with Ricci curvature bounded below by  $-K$ , i.e.,  $\text{Ric} \geq -K$ , every harmonic function which is bounded below by a constant satisfies the gradient estimate

$$|\nabla \log u| \leq \sqrt{(n-1)K}(u - \inf u).$$

In particular, Yau [40] proved the Strong Liouville theorem holds: On complete Riemannian manifolds with non-negative Ricci curvature, every positive (and hence bounded) harmonic function must be constant. See also [39]. In [20], the author extended Yau's gradient estimate and Strong Liouville theorem to the  $L$ -harmonic functions on complete Riemannian manifolds via the Bakry-Emery Ricci curvature: Suppose that  $\text{Ric}_{m,n}(L) \geq -K$  for some constant  $K \geq 0$ . Then every  $L$ -harmonic function  $u$ , which is bounded from below, satisfies the following gradient estimate

$$|\nabla \log u| \leq \sqrt{(m-1)K}(u - \inf u).$$

In particular, if  $\text{Ric}_{m,n}(L) \geq 0$ , then all positive (and hence bounded)  $L$ -harmonic functions must be constant. In [21], the local gradient estimates was proved for  $L$ -harmonic functions: If  $\text{Ric}_{m,n}(L) \geq -K$ , then every positive  $L$ -harmonic function  $u$  on  $M$  satisfies: for all  $o \in M$ , and  $R > 0$ , we have

$$|\nabla \log u(x)| \leq \sqrt{(m-1)K}u(x) + C(K, m)(R^{-1/2} + R^{-3/2}), \quad \forall x \in B(o, R).$$

In [7], Brighton proved that, on complete Riemannian manifolds with  $\text{Ric} + \nabla^2\phi \geq 0$ , every bounded  $L$ -harmonic function must be constant, i.e., Corollary 1.3. Our method for the proof of Corollary 1.3 is different from [7]. For more recent results, see [30, 31].

### 3 Proof of Theorem 1.4

In this section, we use Hamilton's Harnack inequality to prove Theorem 1.4. This proof is similar to the one given in [18, 11] for the gradient estimate of the logarithmic heat kernel of the Laplace-Beltrami operator on complete Riemannian manifolds with non-negative Ricci curvature.

To prove Theorem 1.4, we need the following estimates on the heat kernel of the Witten Laplacian on complete Riemannian manifolds with weighted volume measure.



**Proposition 3.1** *Suppose that there exist some constants  $m \geq n$ ,  $m \in \mathbb{N}$  and  $K \geq 0$  such that  $\text{Ric}_{m,n}(L) \geq -K$ . Then, for any small  $\varepsilon > 0$ , there exist some constants  $C_i = C_i(m, n, K, \varepsilon) > 0$ ,  $i = 1, 2$ , such that for all  $x, y \in M$  and  $t > 0$ ,*

$$p_t(x, y) \leq \frac{C_1}{\mu(B_y(\sqrt{t}))} \exp\left(-\frac{d^2(x, y)}{4(1+\varepsilon)t} + \alpha\varepsilon Kt\right) \times \left(\frac{d(x, y) + \sqrt{t}}{\sqrt{t}}\right)^{m/2} \exp\left(\frac{\sqrt{(m-1)K}d(x, y)}{2}\right), \quad (9)$$

where  $\alpha$  is a constant depending only on  $m$ , and

$$p_t(x, y) \geq C_2 e^{-(1+\varepsilon)\lambda_{K,m}t} \mu^{-1}(B_y(\sqrt{t})) \exp\left(-\frac{d^2(x, y)}{4(1-\varepsilon)t}\right) \left[\frac{\sqrt{K}d(x, y)}{\sinh \sqrt{K}d(x, y)}\right]^{\frac{m-1}{2}}, \quad (10)$$

where

$$\lambda_{K,m} = \frac{(m-1)^2 K}{8}.$$

*Proof.* The lower bound estimate (10) has proved in [23]. By Theorem 5.4 in [20], for all  $\varepsilon > 0$ , there exists a constant  $C_3 = C_3(m, n, K, \varepsilon) > 0$

$$p_t(x, y) \leq \frac{C_3}{\sqrt{\mu(B_x(\sqrt{t}))\mu(B_y(\sqrt{t}))}} \exp\left(-\frac{d^2(x, y)}{4(1+\varepsilon)t} + \alpha\varepsilon Kt\right). \quad (11)$$

Similarly to [20] (Step 2 of p. 1324), using the Bishop-Gromov relative volume comparison theorem [32, 20], we have

$$\begin{aligned} \mu(B_y(\sqrt{t})) &\leq \mu\left(B_x(d(x, y) + \sqrt{t}) \setminus B_x(d(x, y) - \sqrt{t})\right) \\ &\leq \mu(B_x(\sqrt{t})) \frac{V_{m,K}(d(x, y) + \sqrt{t}) - V_{m,K}(d(x, y) - \sqrt{t})}{V_{m,K}(\sqrt{t})} \\ &\leq \mu(B_x(\sqrt{t})) \frac{V_{m,K}(d(x, y) + \sqrt{t})}{V_{m,K}(\sqrt{t})}, \end{aligned}$$

where  $V_{m,K}(r)$  denotes the volume of geodesic balls of radius  $r$  in the  $m$ -dimensional hyperbolic space form  $\mathbb{H}^m(K)$  of constant sectional curvature  $K/m - 1$ . Using again the Bishop-Gromov volume comparison theorem on  $\mathbb{H}^m(K)$ , we have

$$\begin{aligned} \frac{\mu(B_y(\sqrt{t}))}{\mu(B_x(\sqrt{t}))} &\leq \frac{V_{m,K}(d(x, y) + \sqrt{t})}{V_{m,K}(\sqrt{t})} \\ &\leq \left(\frac{d(x, y) + \sqrt{t}}{\sqrt{t}}\right)^m \exp[\sqrt{(m-1)K}d(x, y)]. \end{aligned} \quad (12)$$

By (11) and (12), we can obtain the upper bound estimate (9).  $\square$

**Proof of Theorem 1.4.** Fix  $T > 0$ , and let  $u(t, x)$  be a positive and bounded solution to the heat equation  $\partial_t u = Lu$ ,  $t \in (0, t_1)$ . Let

$$A := \sup\{u(t, x) : 0 \leq t \leq t_1, x \in M\}.$$

By Hamilton's Harnack inequality (3), we have

$$t|\nabla \log u(t, x)|^2 \leq (1 + 2Kt) \log(A/u(t, x)), \quad \forall (t, x) \in [0, t_1] \times M. \quad (13)$$

Let  $s \in (0, T]$ ,  $y \in M$ ,  $t_1 = s/2$  and  $u(t, x) = p_{s/2+t}(x, y)$ . By (13), (9) and (10), we have

$$t|\nabla_x \log p_{s/2+t}(x, y)|^2 \leq C_{K,m,T}(1 + Kt) \times \left[ 1 + \frac{d(x, y)}{\sqrt{t}} + \log \left( \frac{C_1 \mu(B(y, \sqrt{s/2+t}))}{C_2 \mu(B(y, \sqrt{s/2}))} \exp \left( C_3 \frac{d^2(x, y)}{s/2+t} + C_4 d(x, y) \right) \right) \right].$$

In particular, taking  $t = s/2$  and changing  $s$  by  $t$  we get

$$\frac{t}{2}|\nabla_x \log p_t(x, y)|^2 \leq C_{K,m,T}(1 + K\frac{t}{2}) \times \left[ 1 + \frac{d(x, y)}{\sqrt{t}} + \log \left( \frac{C_1 \mu(B(y, \sqrt{t}))}{C_2 \mu(B(y, \sqrt{t/2}))} \exp \left( C_3 \frac{d^2(x, y)}{t} + C_4 d(x, y) \right) \right) \right].$$

By the Bishop-Gromov volume comparison (23), we derive

$$t|\nabla_x \log p_t(x, y)|^2 \leq C_{K,m,T} \left( 1 + \frac{d^2(x, y)}{t} + \frac{d(x, y)}{\sqrt{t}} + d(x, y) \right),$$

which yields

$$|\nabla_x \log p_t(x, y)| \leq C_{K,m,T} \left( \frac{d(x, y)}{t} + \frac{1}{\sqrt{t}} \right).$$

This finishes the analytic proof of Theorem 1.4. □

## 4 Gradient estimates of the heat kernel

In this section we use a probabilistic approach to prove Theorem 1.4 and Theorem 1.5. Our proof is inspired by previous works initiated by Sheu [36], and developed by Hsu [17] and Engoulatov [13].

Let  $O(M)$  be the orthogonal frame bundle over  $M$ ,  $\pi : O(M) \rightarrow M$  the canonical projection map. Let  $H_1, \dots, H_n$  be the canonical horizontal vector fields on  $O(M)$ . Let  $B_t$  be the standard Brownian motion on  $\mathbb{R}^n$ . Following Malliavin [28, 29], we define the horizontal  $L$ -diffusion process  $U_t$  on  $O(M)$  by the following Stratonovich SDE on  $O(M)$ :

$$dU_t = \sum_{i=1}^n H_i(U_t) \circ dB_t^i - \nabla^H \phi(U_t) dt, \quad (14)$$

where  $\nabla^H \phi$  denotes the horizontal lift of the gradient vector field  $\nabla \phi$  on  $O(M)$ , which is the unique horizontal vector field on  $O(M)$  such that

$$\pi_*(\nabla^H \phi) = \nabla \phi.$$

Let

$$X_t = \pi(U_t), \quad \forall t > 0.$$

Then  $X_t$  is a diffusion process on  $M$  with infinitesimal generator  $L$ . Moreover, we have

$$dX_t = U_t \circ dB_t - \nabla \phi(X_t) dt. \quad (15)$$

Let  $\Delta_{O(M)} = \sum_{i=1}^n H_i^2$  be the horizontal Laplace-Beltrami on  $O(M)$ . The horizontal Witten Laplacian is defined by

$$L_{O(M)} = \Delta_{O(M)} - \nabla^H \phi \cdot \nabla^H.$$

Fix  $T > 0$ . Let  $\mathbb{P}_{x,y,T}$  be the conditional  $L$ -diffusion processes starting from  $x$  and ending at  $y$  at time  $T$ , i.e.,

$$\mathbb{P}_{x,y,T}(\cdot) = \mathbb{P}_x(\cdot | X_T = y).$$

Let

$$J(t, u) = \log p_{T-t}(\pi u, y), \quad \forall u \in O(M), t \in [0, T].$$

By the Girsanov transformation, we have

$$\left. \frac{d\mathbb{P}_{x,y,T}}{d\mathbb{P}_x} \right|_{\mathcal{F}_t} = \frac{p_{T-t}(X_t, y)}{p_T(x, y)} = \exp\{J(t, U_t) - J(0, U_0)\}, \quad \forall t < T.$$

where  $\mathcal{F}_t = \sigma(X_s, s \leq t)$  is the natural  $\sigma$ -field generated by  $X_t$ . By Itô's formula, we have

$$dJ(t, U_t) = \frac{\partial J}{\partial t}(t, U_t)dt + \sqrt{2}\langle \nabla^H J(t, U_t), dB_t \rangle + L_{O(M)}J(t, U_t)dt,$$

Note that  $J(t, u)$  satisfies the Hamilton-Jacobi equation

$$\frac{\partial J}{\partial t} + L_{O(M)}J + |\nabla^H J|^2 = 0, \quad (16)$$

which yields

$$\left. \frac{d\mathbb{P}_{x,y,T}}{d\mathbb{P}_x} \right|_{\mathcal{F}_t} = \exp \left[ \int_0^t \langle \sqrt{2}\nabla^H J(s, U_s), dB_s \rangle - \int_0^t |\nabla^H J(s, U_s)|^2 ds \right], \quad \forall t < T.$$

It follows, by Girsanov's theorem, that the process

$$b_t = B_t - \sqrt{2} \int_0^t \nabla^H J(s, U_s) ds, \quad s \in [0, T], \quad (17)$$

is a Brownian motion under  $\mathbb{P}_{x,y,T}$ . Substituting (17) into (14) and (15), we have the following lemma.

**Lemma 4.1** *The conditional  $L$ -diffusion process  $X_t$  on  $M$  starting from  $x$  with terminal end  $y$  at the time  $T$ , and its horizontal lift  $U_t$ , are the solutions of the following Stratonovich SDEs:*

$$\begin{aligned} dU_t &= \sum_{i=1}^n H_i(U_t) \circ \left[ \sqrt{2} db_t^i + 2H_i \log p_{T-t}(U_t, y) dt \right] - \nabla^H \phi(U_t) dt, \\ dX_t &= U_t \circ \left[ \sqrt{2} db_t + 2\nabla^H \log p_{T-t}(U_t, y) dt \right] - \nabla \phi(X_t) dt, \end{aligned}$$

where  $b_t$  is a standard Brownian motion on  $\mathbb{R}^n$  under  $\mathbb{P}_{x,y,T}$ .

**Proposition 4.2** *For all  $0 < t < T$ , we have*

$$\mathbb{E}[d|\nabla^H J(t, U_t)|^2] = 2\mathbb{E} \left[ |\nabla^H \nabla^H J|^2 + (\text{Ric} + \nabla^2 \phi)(\nabla^H J, \nabla^H J) \right]. \quad (18)$$

*Proof.* By Lemma 4.1 and Itô's formula, we have

$$d|\nabla^H J(t, U_t)|^2 = \langle \nabla^H |\nabla^H J|^2, \sqrt{2}db_t + 2\nabla^H J dt \rangle + (\partial_t + L_{O(M)})|\nabla^H J|^2 dt.$$

By the generalized Bochner-Weitzenböck formula, the action of the horizontal Witten Laplacian  $L_{O(M)}$  on  $|\nabla^H J|^2$  is given by

$$L_{O(M)}|\nabla^H J|^2 = 2\langle \nabla^H L_{O(M)}J, \nabla^H J \rangle + 2|\nabla^H \nabla^H J|^2 + 2(\text{Ric} + \nabla^2\phi)(\nabla^H J, \nabla^H J).$$

On the other hand, using the Hamilton-Jacobi equation (16), we have

$$\partial_t |\nabla^H J|^2 = 2\langle \partial_t \nabla^H J, \nabla^H J \rangle = -2\langle L_{O(M)} \nabla^H J, \nabla^H J \rangle - 2\langle \nabla^H |\nabla^H J|^2, \nabla^H J \rangle.$$

Combining this with the previous calculation, we have

$$d|\nabla^H J(t, U_t)|^2 = \langle \nabla^H |\nabla^H J|^2, \sqrt{2}db_t \rangle + 2[|\nabla^H \nabla^H J|^2 + (\text{Ric} + \nabla^2\phi)(\nabla^H J, \nabla^H J)] dt.$$

Taking expectation with respect to  $\mathbb{E}_{x,y,T}$ , we complete the proof of Proposition 4.2.  $\square$

**Proposition 4.3** *Suppose that  $\text{Ric} + \nabla^2\phi \geq -K$ . Then for all  $T > 0$ , we have*

$$|\nabla \log p_T(x, y)|^2 \leq 2 \left( \frac{1}{T} + K \right) \mathbb{E} \left[ \log \frac{p_{T/2}(X_{T/2}, y)}{p_T(x, y)} \right]. \quad (19)$$

*Proof.* Based on Proposition 4.2, the proof of Proposition 4.3 is similar to the argument used in [17, 13]. For the convenience of the reader, we give the detail here. Under the condition  $\text{Ric} + \nabla^2\phi \geq -K$ , integrating (18) from 0 to  $t$ , we have

$$\mathbb{E}[|\nabla^H J(t, U_t)|^2] - |\nabla^H J(0, U_0)|^2 \geq -2KE \left[ \int_0^t |\nabla^H J(s, U_s)|^2 ds \right].$$

Integrating again with respect to  $t$  we can obtain

$$\frac{T}{2} |\nabla^H J(0, U(0))|^2 \leq (1 + KT) \mathbb{E} \left[ \int_0^{T/2} |\nabla^H J(s, U_s)|^2 ds \right]. \quad (20)$$

By Itô's formula and the Hamilton-Jacobi equation (16), we have

$$\begin{aligned} dJ(t, U_t) &= (\partial_t + L_{O(M)})J dt + \langle \sqrt{2}\nabla^H J, db_t + \sqrt{2}\nabla^H J dt \rangle \\ &= \langle \sqrt{2}\nabla^H J, db_t \rangle + |\nabla^H J|^2 dt. \end{aligned}$$

Taking expectation with respect to  $\mathbb{E}_{x,y,T}$  and integrating from  $t = 0$  to  $t = T/2$ , we get

$$\mathbb{E}[J(T/2, U_{T/2})] - J(0, U_0) = \mathbb{E} \left[ \int_0^{T/2} |\nabla^H J|^2 dt \right]. \quad (21)$$

Combining (20) with (21), we finish the proof of (19).  $\square$

**Probabilistic proof of Theorem 1.4.** By Proposition 3.1, we have

$$\begin{aligned} \frac{p_{t/2}(X_{t/2}, y)}{p_t(x, y)} &\leq C_1(K, m, T) \frac{\mu(B_y(\sqrt{t}))}{\mu(B_y(\sqrt{t/2}))} \exp \left( \frac{d^2(x, y)}{4(1-\varepsilon)t} + \frac{\sqrt{(m-1)K}d(X_{t/2}, y)}{2} \right) \\ &\quad \times \left( \frac{d(X_{t/2}, y) + \sqrt{t}}{\sqrt{t}} \right)^{m/2} \left[ \frac{\sqrt{K}d(x, y)}{\sinh \sqrt{K}d(x, y)} \right]^{-\frac{m-1}{2}}. \end{aligned} \quad (22)$$

The Bishop-Gromov volume comparison theorem ([32, 20]) implies

$$\frac{\mu(B_y(\sqrt{t}))}{\mu(B_y(\sqrt{t/2}))} \leq 2^{m/2} \exp(\sqrt{m-1}Kt) \leq 2^{m/2} \exp(\sqrt{m-1}KT). \quad (23)$$

Using (22), (23) and the elementary inequality  $x/\sinh x \geq e^{-x}$ , we obtain

$$\begin{aligned} \log \left[ \frac{p_{t/2}(X_{t/2}, y)}{p_t(x, y)} \right] &\leq C_2(K, m, T) \left[ 1 + \frac{d(X_{t/2}, y)}{\sqrt{t}} + \frac{d^2(x, y)}{t} + d(X_{t/2}, y) + d(x, y) \right] \\ &\leq C_3(K, m, T) \left[ 1 + \frac{d^2(X_{t/2}, y)}{t} + \frac{d^2(x, y)}{t} + d(X_{t/2}, y) + d(x, y) \right]. \end{aligned}$$

By Itô's formula, we can prove that (cf. [23])

$$\mathbb{E}_{x, y, T}[d^k(x_{t/2}, y)] \leq (d^k(x, y) + 1)e^{C_4(K, m)t}, \quad \forall t > 0, k = 1, 2.$$

Taking expectation with respect to  $\mathbb{E}_{x, y, T}$  in (24), we have

$$\mathbb{E}_{x, y, T} \left[ \log \frac{p_{t/2}(X_{t/2}, y)}{p_t(x, y)} \right] \leq C_5(K, m, T) \left[ 1 + \frac{d^2(x, y) + 1}{t} + d(x, y) \right].$$

By Proposition 4.3, we have

$$\begin{aligned} |\nabla \log p_t(x, y)|^2 &\leq C_6(K, m, T) \left[ \frac{1}{t} + \frac{d^2(x, y) + 1}{t^2} + \frac{d(x, y)}{t} \right] \\ &\leq C_7(K, m, T) \left[ \frac{d(x, y)}{t} + \frac{1}{\sqrt{t}} \right]^2. \end{aligned}$$

The proof of Theorem 1.4 is completed.  $\square$

**Proof of Theorem 1.5.** The proof is similarly to Sheu [36] and Hsu [17]. Let  $I = \{i_1, \dots, i_N\}$ , and denote  $H_I = H_{i_1} \dots H_{i_N}$ , then by Lemma 4.1 and repeatedly using Itô's formula, we have

$$dH_I J = \langle \nabla^H H_I J, \sqrt{2}db_t \rangle + [\partial_t H_I J + L_{O(M)} H_I J + 2\langle \nabla^H H_I J, \nabla^H J \rangle] dt. \quad (24)$$

Substituting the Hamilton-Jacobi equation (16) into (24), we have

$$dH_I J = \langle \nabla^H H_I J, \sqrt{2}db_t \rangle + (F_I + G_I)dt. \quad (25)$$

where

$$F_I = [L_{O(M)}, H_I]J = [\Delta_{O(M)}, H_I]J - [\nabla^H \phi, H_I]J,$$

and

$$G_I = 2\langle \nabla^H H_I J, \nabla^H J \rangle - H_I |\nabla^H J|^2.$$

By (21) and (22), we have

$$\mathbb{E} \left[ \int_0^{T/2} |\nabla^H J(t, U_t)|^2 dt \right] \leq CT \left[ \frac{d(x, y)}{T} + \frac{1}{\sqrt{T}} \right]^2, \quad (26)$$

and by Theorem 1.4, we have

$$|H_I J(0, U_0)| \leq C \left[ \frac{d(x, y)}{T} + \frac{1}{\sqrt{T}} \right]. \quad (27)$$

Note that

$$[\nabla^H \phi, H_I] J = \sum_{k=1}^n [(H_k \phi) H_k, H_I] J = \sum_{k=1}^n (H_k \phi [H_k, H_I] J - H_I H_k \phi H_k J),$$

and by Cartan's structure theorem, we have

$$[H_i, H_j] = V_{ij}, \quad [H_i, V_{jk}] = \Omega_{jk}^{il} H_l, \quad [V_{ij}, V_{kl}] = c_{ij,kl}^{ab} V_{ab},$$

where  $V_{ij}$  are the vertical fields on  $O(M)$ ,  $\Omega$  is the  $o(n)$ -valued curvature form on  $O(M)$ , and  $c_{ij,kl}^{ab}$  are the structure constants of  $o(n)$ . The assumption of Theorem 1.5 says that the  $L^\infty$ -norms  $\|\nabla^k \Omega\|_\infty$  are bounded for all  $k = 0, 1, \dots, N$ , and the  $L^\infty$ -norms  $\|\nabla^k \phi\|_\infty$  are bounded for  $k = 1, \dots, N + 1$ . This yields

$$|F_I| \leq C_1 |\nabla^H J| + \dots + C_1 |(\nabla^H)^{|I|} J|, \quad (28)$$

$$|G_I| \leq C_1 |\nabla^H J| + \dots + C_1 |(\nabla^H)^{|I|} J| + C_1 |(\nabla^H)^{|I|} J|^2, \quad (29)$$

where  $C_1 > 0$  is a constant depending only on  $\|\nabla^i \Omega\|_\infty$  and  $\|\nabla^i \phi\|_\infty$ ,  $i = 1, \dots, |I|$ . By induction, and using (25), (26), (27), (28) and (29), we can prove that for all  $I$  with  $2 \leq |I| \leq N$ ,

$$\mathbb{E} \left[ \int_0^{T/2} |H_I J(t, U_t)|^2 dt \right] \leq C_2 \left[ \frac{d(x, y)}{T} + \frac{1}{\sqrt{T}} \right]^{2(|I|-1)}, \quad (30)$$

and for all  $I$  with  $|I| \leq N$ ,

$$|H_I J(0, U_0)| \leq C_3 \left[ \frac{d(x, y)}{T} + \frac{1}{\sqrt{T}} \right]^{|I|}. \quad (31)$$

where  $C_2$  and  $C_3$  are two constants depending on  $K, m, n, T$  and  $C_1$ . The proof of Theorem 1.5 is completed. For the detail of the above induction argument for  $L = \Delta$  on compact Riemannian manifolds, see Hsu [17].  $\square$

## 5 Proofs of Theorem 1.6

Let

$$H(u) = - \int_M u \log u d\mu.$$

be the Boltzmann-Shannon-Nash entropy of the heat equation of the Witten Laplacian on  $(M, g, \mu)$ .

To prove Theorem 1.6, we need the following entropy dissipation formulas for the Witten Laplacian on complete Riemannian manifolds.

**Theorem 5.1** ([23]) *Let  $M$  be a complete Riemannian manifold, and  $\phi \in C^2(M)$ . Suppose that there exists a constant  $K \geq 0$  such that*

$$\text{Ric}(L) \geq -K.$$

Then, for any positive solution to the heat equation

$$\partial_t u = Lu$$

with  $u(\cdot, 0) \in L^1(\mu) \cap L^4(d(\cdot, o)d\mu)$ , where  $o \in M$  is any fixed point, we have

$$\frac{d}{dt}H(u) = - \int_M Lu \log u d\mu. \quad (32)$$

**Theorem 5.2** *Let  $(M, g)$  be an  $n$ -dimensional complete Riemannian manifold. Suppose that  $\|\nabla^k \text{Riem}\| \leq C_k$  for some constant  $C_k > 0$ ,  $0 \leq k \leq 3$ ,  $\phi \in C^4(M)$  with  $\nabla \phi \in C_b^3(M)$ , and there exists some  $m \geq n$ ,  $K \geq 0$  such that  $\text{Ric}_{m,n}(L) \geq -K$ . Then, for the fundamental solution of the heat equation  $\partial_t u = Lu$ , we have*

$$\frac{d^2}{dt^2}H(u) = - \int_M \left( \frac{|Lu|^2}{u} - \left\langle \nabla Lu, \frac{\nabla u}{u} \right\rangle \right) d\mu, \quad (33)$$

and

$$\frac{d^2}{dt^2}H(u) = -2 \int_M (|\nabla^2 \log u|^2 + \text{Ric}(L)(\nabla \log u, \nabla \log u)) u d\mu. \quad (34)$$

*Proof.* Based on Theorem 5.1, we can prove (33) by the same argument as used in the proof of Theorem 4.3 in [23]. Integrating by part yields

$$\begin{aligned} \int_M \frac{|Lu|^2}{u} d\mu &= - \int_M \left\langle \nabla \left( \frac{Lu}{u} \right), \frac{\nabla u}{u} \right\rangle u d\mu \\ &= - \int_M \langle \nabla Lu, \nabla \log u \rangle d\mu + \int_M Lu |\nabla \log u|^2 d\mu. \end{aligned}$$

Hence

$$\begin{aligned} \frac{d^2}{dt^2}H(u) &= 2 \int_M \left\langle \nabla Lu, \frac{\nabla u}{u} \right\rangle d\mu - \int_M Lu |\nabla \log u|^2 d\mu \\ &= 2 \int_M \left\langle \nabla Lu, \frac{\nabla u}{u} \right\rangle d\mu - \int_M uL |\nabla \log u|^2 d\mu. \end{aligned}$$

By the Bochner formula for the Witten Laplacian, we have (see [23])

$$L|\nabla \log u|^2 = 2\langle \nabla L \log u, \nabla \log u \rangle + 2|\nabla^2 \log u|^2 + 2\text{Ric}(L)(\nabla \log u, \nabla \log u)$$

Therefore

$$\begin{aligned} \frac{d^2}{dt^2}H(u) &= 2 \int_M \left\langle \nabla Lu, \frac{\nabla u}{u} \right\rangle d\mu - 2 \int_M \langle \nabla L \log u, \nabla u \rangle d\mu \\ &\quad - 2 \int_M (|\nabla^2 \log u|^2 + \text{Ric}(L)(\nabla \log u, \nabla \log u)) u d\mu. \end{aligned}$$

Again, integrating by parts yields

$$\int_M \langle \nabla L \log u, \nabla u \rangle d\mu = - \int_M L \log u \cdot Lu d\mu = \int_M \langle \nabla \log u, \nabla Lu \rangle d\mu.$$

Thus

$$\frac{d^2}{dt^2}H(u) = -2 \int_M (|\nabla^2 \log u|^2 + \text{Ric}(L)(\nabla \log u, \nabla \log u)) u d\mu.$$

The proof of Theorem 5.2 is completed.  $\square$

**Remark 5.3** Theorem 5.2 is an improvement of Theorem 4.3 in [23]. The entropy dissipation formulas (32) and (33) play an important role in the study of the convergence of the heat kernel measure  $p_t(\cdot, y)d\mu(y)$  towards the equilibrium measure  $\mu$ . In the case where  $M$  is compact and  $\phi \in C^3(M)$ , it is well-known that the entropy dissipation formulas (32) and (33) hold. However, as far as we know, even in the non-weighted case, it seems that there is no explicit result in the literature (except our previous paper [23]) which ensures the entropy dissipation formulas (32) and (33) on complete non-compact Riemannian manifolds. Theorem 5.1 and Theorem 5.2 give us natural geometric conditions on  $(M, g, \phi)$  to ensure (32) and (33).

**Proof of Theorem 1.6.** We simplify the proof of Theorem 2.3 in [23] as follows. Let

$$H_m(u, t) = - \int_M u \log u d\mu - \frac{m}{2} (1 + \log(4\pi t)).$$

By Theorem 5.1 and Theorem 5.2, we have

$$\frac{d}{dt}H_m(u, t) = - \int_M uL \log u d\mu - \frac{m}{2t} = \int_M \left( |\nabla \log u|^2 - \frac{m}{2t} \right) u d\mu, \quad (35)$$

and

$$\frac{d^2}{dt^2}H_m(u, t) = -2 \int_M (|\nabla^2 \log u|^2 + Ric(L)(\nabla \log u, \nabla \log u)) u d\mu + \frac{m}{2t^2}. \quad (36)$$

By the definition formula of the  $W$ -entropy, we have

$$W_m(u, t) = \frac{d}{dt}(tH_m(u, t)) = H_m(u, t) + t \frac{d}{dt}H_m(u, t), \quad (37)$$

and

$$\frac{d}{dt}W_m(u, t) = 2 \frac{d}{dt}H_m(u, t) + t \frac{d^2}{dt^2}H_m(u, t). \quad (38)$$

Substituting (32) and (35) into (37), and using  $u = \frac{e^{-f}}{(4\pi t)^{m/2}}$ , we obtain

$$\begin{aligned} W_m(u, t) &= - \int_M u \log u d\mu - \frac{m}{2} (1 + \log(4\pi t)) - t \int_M uL \log u d\mu - \frac{m}{2} \\ &= \int_M \left[ t|\nabla \log u|^2 - \log u - m - \frac{m}{2} \log(4\pi t) \right] u d\mu \\ &= \int_M [t|\nabla f|^2 + f - m] \frac{e^{-f}}{(4\pi t)^{m/2}} d\mu. \end{aligned}$$

This proves (4) in Theorem 1.6. Substituting (35) and (36) into (38), we obtain

$$\frac{d}{dt}W_m(u, t) = -2 \int_M uL \log u d\mu - 2 \int_M t(|\nabla^2 \log u|^2 + Ric(L)(\nabla \log u, \nabla \log u)) u d\mu - \frac{m}{2t}.$$

Noting that

$$2t|\nabla^2 \log u|^2 + \frac{m}{2t} = 2t \left| \nabla^2 \log u - \frac{g}{2t} \right|^2 + \frac{m-n}{2t} + 2\Delta \log u,$$

and by the integration by part formula

$$\begin{aligned} \int_M \Delta \log u u d\mu &= \int_M (L \log u + \nabla \phi \cdot \nabla \log u) u d\mu \\ &= - \int_M |\nabla \log u|^2 u d\mu + \int_M \nabla \phi \cdot \nabla \log u u d\mu, \end{aligned}$$



we have

$$\begin{aligned} \frac{d}{dt}W_m(u, t) &= -2t \int_M \left| \nabla^2 \log u - \frac{g}{2t} \right|^2 u d\mu + 2 \int_M \nabla \phi \cdot \nabla \log u u d\mu \\ &\quad - 2 \int_M t \text{Ric}(L)(\nabla \log u, \nabla \log u) u d\mu - \frac{m-n}{2t}. \end{aligned}$$

By the same argument as in [23], the above identity yields (4). The monotonicity of  $W$  follows and the rigidity part of Theorem 1.6 can be proved by the same argument used in Theorem 2.4 in [23].  $\square$

## 6 LSI and lower boundedness of the $\mu$ -entropy

Following [35], we introduce the best logarithmic Sobolev constant (called also the  $\mu$ -entropy)

$$\mu(\tau) := \inf \left\{ W_m(u, \tau) : u = \frac{e^{-f}}{(4\pi\tau)^{m/2}}, \int_M u d\mu = 1 \right\}.$$

In [23], the author proved that  $\mu(\tau)$  is always bounded from below on compact Riemannian manifolds  $M$  with  $\phi \in C^2(M)$ . In this section, we prove a family of logarithmic Sobolev inequalities (LSI) on complete Riemannian manifolds with the uniform volume lower bound condition. The lower boundedness of  $\mu(\tau)$  is also proved.

**Proof of Theorem 1.7.** Let  $v = \sqrt{u}$ . Then we can rewrite  $W_m(u, \tau)$  as follows

$$W_m(u, \tau) = 4\tau \int_M |\nabla v|^2 d\mu - \int_M v^2 \log v^2 d\mu - \left( m + \frac{m}{2} \log(4\pi\tau) \right).$$

We need to prove, for all  $v \in C_0^\infty(M)$  with  $\int_M v^2 d\mu = 1$ , we have

$$4\tau \int_M |\nabla v|^2 d\mu - \int_M v^2 \log v^2 d\mu - m \left( 1 + \frac{1}{2} \log(4\pi\tau) \right) \geq \mu(\tau) > -\infty. \quad (39)$$

Under the conditions of theorem, by Theorem 2.8 in [22], the following  $L^2$ -Sobolev inequality holds: there exists a constant  $C_1(m, K, \mu_0) > 0$ , depending only on  $m, K, \mu_0$ , such that

$$\|f\|_{\frac{2m}{m-2}}^2 \leq C_1(m, K, \mu_0) (\|\nabla f\|_2^2 + \|f\|_2^2), \quad \forall f \in C_0^\infty(M).$$

By [10], the above  $L^2$ -Sobolev inequality implies that, for any  $\varepsilon > 0$ , there exists a constant  $\beta(\varepsilon) > 0$  such that the following logarithmic Sobolev inequality holds

$$\int_M f^2 \log f^2 d\mu \leq \varepsilon \int_M |\nabla f|^2 + \beta(\varepsilon) \|f\|_2^2 + \|f\|_2^2 \log \|f\|_2^2, \quad \forall f \in C_0^\infty(M), \quad (40)$$

where for some constant  $C_2(m, K, \mu_0) > 0$  depending only on  $m, K, \mu_0$ , we have

$$\beta(\varepsilon) \leq C_2(m, K, \mu_0) - \frac{m}{2} \log \varepsilon.$$

Applying (40) to  $f = v \in C_0^\infty(M)$  with  $\int_M v^2 d\mu = 1$ , and taking  $\varepsilon = 4\tau$ , we have

$$\int_M v^2 \log v^2 d\mu \leq 4\tau \int_M |\nabla v|^2 - m \left( 1 + \frac{1}{2} \log(4\pi\tau) \right) - \mu(\tau),$$

where

$$-\mu(\tau) := \beta(4\tau) + m \left( 1 + \frac{1}{2} \log(4\pi\tau) \right),$$

This proves (39) with

$$\mu(\tau) \geq -\left(C_2(m, K, \mu_0) + m + \frac{m}{2} \log(4\pi)\right), \quad \forall \tau > 0.$$

This finishes the proof of Theorem 1.7.  $\square$

**Acknowledgement.** The author would like to thank F. Baudoin for helpful discussion on the alternative proof of Theorem 1.1 in September 2011. He would like also to thank D. Bakry, E. Carlen, E. Hsu and A. Thalmaier for their interests on this work.

## References

- [1] M. Arnaudon, A. Thalmaier, Li-Yau type gradient estimates and Harnack inequalities by stochastic analysis, *Advanced Studies in Pure Mathematics*, 57 (2010) 29-48.
- [2] M. Arnaudon, A. Thalmaier, F.-Y. Wang, Gradient estimate and Harnack inequality on non-compact Riemannian manifolds, *Stochastic Processes Appl.* 119 (2009) 3653-3670.
- [3] D. Bakry, Un critère de non-explosion pour certaines diffusions sur une variété riemannienne complète, *C. R. Acad. Sci. Paris, Sér. I* 303 (1986) 23-26.
- [4] D. Bakry, Functional inequalities for Markov semigroups, Notes de cours du Tata Institute, Bombay, 2002. In *Probability measures on groups: recent directions and trends*, 91-147. Proceedings of the CIMPA-TIFR School (2002), Tata Institute of Fundamental Research. Narosha Publishing House, 2006.
- [5] D. Bakry, M. Emery, Diffusion hypercontractives, *Sém. Prob. XIX, Lect. Notes in Maths.* 1123 (1985), 177-206.
- [6] D. Bakry, M. Ledoux, A logarithmic Sobolev form of the Li-Yau parabolic inequality, *Rev. Mat. Iberoam.* 22 (2006), No. 2, 683-702.
- [7] K. Brighton, A Liouville-type theorem for smooth metric measure spaces, arXiv:1006.0751
- [8] B. Chow, P. Lu, L. Ni, Hamilton's Ricci Flow, *Lectures in Contemporary Maths.*, Sciences Press, Beijing, Amer. Math. Soc. 2006.
- [9] B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, P. Lu, L. Ni, *The Ricci Flow, Part I: Techniques and Applications-Geometric Aspects*, Mathematical Surveys and Monographs Ser. 2007.
- [10] E.B. Davies, *Heat Kernel and spectral Theory*, Cambridge University Press, Cambridge, New York, 1990.
- [11] B. Driver, M. Gordia, Integrated Harnack inequalities on Lie groups, *J. Diff. Geom.* 83 (2009) 501-550
- [12] K. Ecker, A formula relating entropy monotonicity to Harnack inequalities, *Comm. Anal. Geom.* 15 (2007), no. 5, 1025-1061.
- [13] A. Engoulatov, A universal bound on the gradient of logarithm of the heat kernel for manifolds with bounded Ricci curvature. *J. Funct. Anal.* 238, 518-529 (2006)
- [14] F. Fang, X.-D. Li, Z.-L. Zhang, Two generalizations of Cheeger-Gromoll splitting theorem via Bakry-Emery Ricci curvature, *Ann. Inst. Fourier (Grenoble)* 59 (2009), no. 2, 563-573.

- [15] A. Futaki, H.-Z. Li, X.-D. Li, On the first eigenvalue of the Witten Laplacian and the diameter of compact shrinking Ricci solitons, [arxiv.org/abs/1111.6364v3](https://arxiv.org/abs/1111.6364v3)
- [16] R.S. Hamilton, A matrix Harnack estimate for the heat equation, *Comm. Anal. Geom.* 1 (1993), no. 1, 113-126.
- [17] E. Hsu, Estimates of derivatives of the heat kernel on a compact Riemannian manifold. *Proc. Am. Math. Soc.* 127 (1999), 37393744.
- [18] B. Kotschwar, Hamiltons gradient estimate for the heat kernel on complete manifolds, *Proc. of Amer. Math. Soc.* 135 (2007), no. 9, 3013-3019.
- [19] P. Li, S.-T. Yau, On the parabolic kernel of the Schrödinger operator, *Acta. Math.* 156 (1986), 153-201.
- [20] X.-D. Li, Liouville theorems for symmetric diffusion operators on complete Riemannian manifolds, *J. Math. Pures Appl.* 84 (2005), 1295-1361.
- [21] X.-D. Li, On the Strong Liouville theorem for weighted Laplacian on complete Riemannian manifolds, preprint, 2007.
- [22] X.-D. Li, Sobolev inequalities on forms and  $L^{p,q}$ -cohomology on complete Riemannian manifolds, *J. Geom. Anal.* 20 (2010), 354-387.
- [23] X.-D. Li, Perelman’s entropy formula for the Witten Laplacian on Riemannian manifolds via Bakry-Emery Ricci curvature, *Math. Ann.* 353 (2012), 403-437.
- [24] X.-D. Li, Perelman’s W-entropy for the Fokker-Planck equation over complete Riemannian manifolds, *Bull. Sci. Math.* 135 (2011) 871-882.
- [25] S. Li, X.-D. Li, Perelmans entropy formula for the Witten Laplacian on manifolds with time dependent metrics and potentials, [arXiv:1303.6019v2](https://arxiv.org/abs/1303.6019v2)
- [26] P. Lu, L. Ni, J.-J. Vazquez, C. Villani, Local Aronson-Benilan estimates and entropy formulae for porous medium and fast diffusion equations on manifolds, *Jour. Math. Pures Appl.*, Vol 91 (2009), no. 1, 1-19.
- [27] J. Lott, Some geometric properties of the Bakry-Emery Ricci tensor. *Comment. Math. Helv.* 78 (2003), 865-883.
- [28] P. Malliavin, Formule de la moyenne, calcul des perturbations et thorie d’annulation pour les formes harmoniques, *J. Funct. Anal.* 17 (1974), 274-291.
- [29] P. Malliavin, *Stochastic Analysis*. Springer, Berlin, 1997.
- [30] O. Munteanu, J. Wang, Smooth metric measure spaces with non-negative curvature, [arxiv.1103.0746v2](https://arxiv.org/abs/1103.0746v2).
- [31] O. Munteanu, J. Wang, Analysis of weighted Laplacian and applications to Ricci solitons, [arxiv.org/abs/1112.3027v1](https://arxiv.org/abs/1112.3027v1)
- [32] Z.-M. Qian, Estimates for weight volumes and applications, *Quart. J. Math. Oxford Ser.* 48 (1987), 235-242.
- [33] L. Ni, The entropy formula for linear equation, *J. Geom. Anal.* 14 (1), 87-100, (2004).
- [34] L. Ni, Addenda to “The entropy formula for linear equation”, *J. Geom. Anal.* 14 (2), 329-334, (2004).

- [35] G. Perelman, The entropy formula for the Ricci flow and its geometric applications, <http://arXiv.org/abs/math/0211159>.
- [36] S.-J. Sheu, Some estimates of the transition density of a nondegenerate diffusion Markov process. *Ann. Probab.* 19(2), 538-561 (1991)
- [37] P. Souplet, Q. S. Zhang, Sharp gradient estimate and Yau-Liouville theorem for the heat equation on noncompact manifolds. *Bull London Math Soc*, 38 (2006), 1045-1053.
- [38] D. Stroock, J. Turesky, Upper bounds on derivatives of the logarithm of the heat kernel. *Commun. Anal. Geom.* 6, 669-685 (1998).
- [39] R. Schoen, S.-T. Yau, *Lectures on Differential Geometry*, International Press, 1994.
- [40] S.T. Yau, Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry, *Indiana Univ. Math. J.*, (1976), 659-670.
- [41] C. Villani, *Optimal Transport, Old and New*, Springer, 2008.
- [42] G. Wei, W. Wylie, Comparison geometry for the Bakry-Emery Ricci tensor, *J. Diff. Geom.* (2009), 377-405.

Xiang-Dong Li, ACADEMY OF MATHEMATICS AND SYSTEM SCIENCE, CHINESE ACADEMY OF SCIENCES, 55, ZHONGGUANCUN EAST ROAD, BEIJING, 100190, P. R. CHINA,  
E-mail: xdli@amt.ac.cn