A SUFFICIENT CONDITION FOR *p*-VALENTLY HARMONIC FUNCTIONS

TOSHIO HAYAMI

ABSTRACT. For normalized harmonic functions $f(z) = h(z) + \overline{g(z)}$ in the open unit disk \mathbb{U} , a sufficient condition on h(z) for f(z) to be *p*-valent in \mathbb{U} is discussed. Moreover, some interesting examples and images of f(z) satisfying the obtained condition are enumerated.

1. INTRODUCTION AND DEFINITIONS

For a fixed p $(p = 1, 2, 3, \dots)$, a meromorphic function f(z) in a domain \mathbb{D} is said to be *p*-valent (or multivalent of order *p*) in \mathbb{D} if for each w_0 (infinity included) the equation $f(z) = w_0$ has at most *p* roots in \mathbb{D} where the roots are counted in accordance with their multiplicity and if there is some w_1 such that the equation $f(z) = w_1$ has exactly *p* roots in \mathbb{D} . In particular, f(z) is said to be univalent (one-to-one) in \mathbb{D} when p = 1.

A complex-valued harmonic function f(z) in \mathbb{D} is given by $f(z) = h(z) + \overline{g(z)}$ where h(z) and g(z) are analytic in \mathbb{D} . We call h(z) and g(z) the analytic part and co-analytic part of f(z), respectively. A necessary and sufficient condition for f(z) to be locally univalent and sense-preserving in \mathbb{D} is |h'(z)| > |g'(z)| for all $z \in \mathbb{D}$ (see [2] or [6]). The theory and applications of harmonic functions are stated in a book due to Duren [3]. Let $\mathcal{H}(p)$ denote the class of functions f(z) of the from

$$f(z) = h(z) + \overline{g(z)} = z^p + \sum_{n=p+1}^{\infty} a_n z^n + \sum_{n=p}^{\infty} b_n z^n$$
(1.1)

which are harmonic in the open unit disk $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$. We next denote by $\mathcal{S}_{\mathcal{H}}(p)$ the class of functions $f(z) \in \mathcal{H}(p)$ which are *p*-valent and sense-preserving in \mathbb{U} . Then, we say that $f(z) \in \mathcal{S}_{\mathcal{H}}(p)$ is a *p*-valently harmonic function in \mathbb{U} .

In the present paper, we discuss a sufficient condition about h(z) for $f(z) \in \mathcal{H}(p)$ given by (1.1), satisfying

$$g'(z) = z^{m-1}h'(z) \tag{1.2}$$

for some m $(m = 2, 3, 4, \dots)$, to be in the class $S_{\mathcal{H}}(p)$. In this case, we can write

$$f(z) = h(z) + \overline{z^{m-1}h(z) - (m-1)\int_0^z \zeta^{m-2}h(\zeta)d\zeta}$$
(1.3)

which means that f(z) is well defined if h(z) is given.

²⁰¹⁰ Mathematics Subject Classification. Primary 30C45, Secondary 58E20.

Key words and phrases. Harmonic function, multivalent function, univalent function.

TOSHIO HAYAMI

2. Main Result

Our result is contained in

Theorem 2.1. Let $h(z) = z^p + \sum_{n=p+1}^{\infty} a_n z^n$ be analytic in the closed unit disk $\overline{\mathbb{U}} = \{z \in \mathbb{C} : |z| \leq 1\}$ with $H(z) = h'(z)/z^{p-1} \neq 0$ $(z \in \overline{\mathbb{U}})$ and let

$$F(t) = (2p + m - 1)t + 2\arg(H(e^{it})) \qquad (-\pi \le t < \pi)$$
(2.1)

for some $m \ (m = 2, 3, 4, \cdots)$. If for each $k \in K = \{0, \pm 1, \pm 2, \cdots, \pm \lfloor \frac{2p+m+1}{2} \rfloor\}$ where $[\]$ is the Gauss symbol, the equation

$$F(t) = 2k\pi \tag{2.2}$$

has at most a single root in $[-\pi,\pi)$ and for all $k \in K$ there exist exactly 2p+m-1 such roots in $[-\pi,\pi)$, then the harmonic function $f(z) = h(z) + \overline{g(z)}$ with $g'(z) = z^{m-1}h'(z)$ belongs to the class $S_{\mathcal{H}}(p)$ and maps \mathbb{U} onto a domain surrounded by 2p+m-1 concave curves with 2p+m-1 cusps.

Proof. We first consider the function $\varphi(t)$ defined as

$$\varphi(t) = f(e^{it}) = h(e^{it}) + \overline{g(e^{it})} \qquad (-\pi \le t < \pi).$$
(2.3)

Supposing that

$$\varphi'(t) = i\left(zh'(z) - \overline{z}\overline{g'(z)}\right) = iz\left(h'(z) - \overline{z}^{m+1}\overline{h'(z)}\right) = 0 \qquad (z = e^{it}), \tag{2.4}$$

we need the following equation

$$z^{m+1}\frac{h'(z)}{h'(z)} = 1$$
 $\left(\overline{z} = e^{-it} = \frac{1}{z}\right).$ (2.5)

This means that

$$\arg\left(e^{i(m+1)t}\frac{h'(e^{it})}{h'(e^{it})}\right) = (m+1)t + 2\arg\left(e^{i(p-1)t}H(e^{it})\right)$$
$$= (2p+m-1)t + 2\arg\left(H(e^{it})\right) = 2k\pi$$
(2.6)

for some $k \in K$ which gives us the equation (2.2). In consideration of the assumption of the theorem, there are 2p + m - 1 distinct roots on the unit circle $\partial \mathbb{U} = \{z \in \mathbb{C} : |z| = 1\}$ and they divide $\partial \mathbb{U}$ onto 2p + m - 1 arcs. Moreover, since $g''(z) = (m - 1)z^{m-2}h'(z) + z^{m-1}h''(z)$, we obtain that

$$\varphi''(t) = -\left(zh'(z) + z^2h''(z) + \overline{z}\overline{g'(z)} + \overline{z}^2\overline{g''(z)}\right)$$
$$= -\left(zh'(z) + z^2h''(z) + m\overline{z}^m\overline{h'(z)} + \overline{z}^{m+1}\overline{h''(z)}\right)$$
(2.7)

and

$$\varphi''(t)\overline{\varphi'(t)} = -\left(zh'(z) + z^{2}h''(z) + m\overline{z}^{m}\overline{h'(z)} + \overline{z}^{m+1}\overline{h''(z)}\right)(-i)\left(\overline{z}\overline{h'(z)} - z^{m}h'(z)\right)$$

$$= i\left(-(m-1)|h'(z)|^{2} + z\overline{h'(z)}h''(z) - \overline{z}\overline{h'(z)}h''(z) - z^{m+1}h'(z)^{2} + \overline{z}^{m+1}\overline{h'(z)}^{2} + (m-1)\overline{z}^{m+1}\overline{h'(z)}^{2} - z^{m+2}h'(z)h''(z) + \overline{z}^{m+2}\overline{h'(z)}h''(z)\right)$$

$$= i\left\{(m-1)\left(\overline{z}^{m+1}\overline{h'(z)}^{2} - |h'(z)|^{2}\right) + 2i\mathrm{Im}\left(z\overline{h'(z)}h''(z) - z^{m+1}h'(z)^{2} - z^{m+2}h'(z)h''(z)\right)\right\}.$$
(2.8)

This leads us that

$$\operatorname{Im}\left(\varphi''(t)\overline{\varphi'(t)}\right) = (m-1)|h'(z)|^{2}\operatorname{Re}\left\{\overline{z}^{m+1}\left(\frac{\overline{h'(z)}}{|h'(z)|}\right)^{2} - 1\right\} \leq 0$$

$$(2.9)$$

for all $z = e^{it}$. Therefore, it follows that

$$\operatorname{Im}\left(\frac{\varphi''(t)}{\varphi'(t)}\right) = \frac{1}{|\varphi'(t)|^2} \operatorname{Im}\left(\varphi''(t)\overline{\varphi'(t)}\right) \leq 0$$
(2.10)

which shows that $\varphi(t)$ maps $[-\pi, \pi)$ onto a union of 2p + m - 1 concave curves. By the help of a simple geometrical observation, we know that the image of $\partial \mathbb{U}$ as a union of 2p + m - 1 concave arcs is a simple curve. Thus, f(z) is *p*-valent in \mathbb{U} and the image $f(\mathbb{U})$ is a domain surrounded by 2p + m - 1 concave curves with 2p + m - 1 cusps. \Box

Remark 2.1. If we take p = 1 in Theorem 2.1, then we readily arrive at the univalence criterion for harmonic functions due to Hayami and Owa [5, Theorem 2.1] (see also [7]).

3. Some Illustrative Examples and Image Domains

In this section, we discuss functions $f(z) = h(z) + \overline{g(z)}$ satisfying the conditions of Theorem 2.1 and their image domains.

Example 3.1. Let $h(z) = z^p$. Then we easily see that the equation (2.2) becomes

$$(2p+m-1)t = 2k\pi$$
 $\left(k = 0, \pm 1, \pm 2, \cdots, \pm \left[\frac{2p+m+1}{2}\right]\right)$ (3.1)

which satisfies the conditions of Theorem 2.1. Hence, the function

$$f(z) = h(z) + \overline{g(z)} = z^p + \frac{p}{p+m-1} z^{p+m-1} \qquad (g'(z) = z^{m-1} h'(z))$$
(3.2)

belongs to the class $S_{\mathcal{H}}(p)$ and it maps \mathbb{U} onto a domain surrounded by 2p + m - 1 concave curves with 2p + m - 1 cusps. Taking p = 2 and m = 4 for example, we know that the function

$$f(z) = z^2 + \frac{2}{5}\overline{z}^5$$
(3.3)

is a 2-valently harmonic function in \mathbb{U} and it maps \mathbb{U} onto the domain surrounded by 7 concave curves with 7 cusps as shown in Figure 1.




FIGURE 1. The image of $f(z) = z^2 + \frac{2}{5}\overline{z}^5$.

Remark 3.1. Since we can rewrite F(t) given by (2.1) as follows:

$$F(t) = (2p + m - 1)t + 2\text{Im}\left(\log H(e^{it})\right), \qquad (3.4)$$

we have that

$$F'(t) = 2p + m - 1 + 2\operatorname{Re}\left(\frac{e^{it}h''(e^{it})}{h'(e^{it})} - (p - 1)\right)$$

= $m + 1 + 2\operatorname{Re}\left(\frac{e^{it}h''(e^{it})}{h'(e^{it})}\right)$ (3.5)

which implies that F(t) is increasing if

$$\operatorname{Re}\left(1+\frac{zh''(z)}{h'(z)}\right) > -\frac{m-1}{2} \qquad (z \in \mathbb{U}).$$

$$(3.6)$$

By the above remark, we derive the following exapmle.

Example 3.2. Let
$$h(z) = z^p + \frac{c}{p+1} z^{p+1} \left(|c| \leq p - \frac{2p}{2p+m+1} \right)$$
. Then the equation (2.2) becomes
 $F(t) = (2p+m-1)t + 2\arg\left(p + ce^{it}\right)$. (3.7)

Noting

$$\operatorname{Re}\left(1+\frac{zh''(z)}{h'(z)}\right) = p+1-\operatorname{Re}\left(\frac{p}{p+cz}\right)$$
$$> p+1-\frac{p}{p-|c|} \ge -\frac{m-1}{2} \qquad (z \in \mathbb{U}),$$
(3.8)

$$F(-\pi) = -(2p+m-1)\pi - 2\arctan\left(\frac{|c|\sin\theta}{p-|c|\cos\theta}\right)$$
(3.9)

and

$$F(\pi) = (2p + m - 1)\pi - 2\arctan\left(\frac{|c|\sin\theta}{p - |c|\cos\theta}\right)$$
(3.10)

where $0 \leq \theta = \arg(c) < 2\pi$, we see that F(t) satisfies the conditions of Theorem 2.1. Hence, the function

$$f(z) = h(z) + \overline{g(z)} = z^p + \frac{c}{p+1} z^{p+1} + \overline{\frac{p}{p+m-1} z^{p+m-1} + \frac{c}{p+m} z^{p+m}}$$
(3.11)

belongs to the class $S_{\mathcal{H}}(p)$ and it maps \mathbb{U} onto a domain surrounded by 2p + m - 1 concave curves with 2p + m - 1 cusps. Putting p = 3, m = 2 and $c = i\left(|c| = 1 < \frac{7}{3}\right)$, we know that the function

$$f(z) = z^3 + \frac{i}{4}z^4 + \frac{3}{4}\overline{z}^4 - \frac{i}{5}\overline{z}^5$$
(3.12)

is a 3-valently harmonic function in \mathbb{U} and it maps \mathbb{U} onto the domain surrounded by 7 concave curves with 7 cusps as shown in Figure 2. We check only the boundary of $f(\mathbb{U})$, for the sake of simplicity.




FIGURE 2. The image of $f(z) = z^3 + \frac{i}{4}z^4 + \frac{3}{4}\overline{z}^4 - \frac{i}{5}\overline{z}^5$.

4. Appendix

The next result was conjectured by Mocanu [8] and proved by Bshouty and Lyzzaik [1].

Theorem 4.1. If h(z) and g(z) are analytic in \mathbb{U} , with $h'(0) \neq 0$, which satisfy

$$g'(z) = zh'(z) \tag{4.1}$$

5

and

$$\operatorname{Re}\left(1 + \frac{zh''(z)}{h'(z)}\right) > -\frac{1}{2} \tag{4.2}$$

for all $z \in \mathbb{U}$, then the harmonic function f(z) = h(z) + g(z) is univalent close-to-convex in \mathbb{U} .

We immediately notice that the above theorem is closely related to Theorem 2.1 and Remark 3.1 with p = 1 and m = 2. This motivates us to state

Conjecture 4.1. If the function f(z) given by (1.1) is harmonic in \mathbb{U} which satisfies

$$g'(z) = z^{m-1}h'(z) (4.3)$$

and

$$\operatorname{Re}\left(1+\frac{zh''(z)}{h'(z)}\right) > -\frac{m-1}{2} \qquad (z \in \mathbb{U})$$

$$(4.4)$$

for some m $(m = 2, 3, 4, \cdots)$, then f(z) is p-valent in \mathbb{U} .

Finally, in view of the process of proving Theorem 2.1, we obtain the following interesting example. Example 4.1. If we consider special functions h(z) and g(z) given by

$$h'(z) = \frac{pz^{p-1}}{1+z^{2p+m-1}} \quad and \quad g'(z) = \frac{pz^{p+m-2}}{1+z^{2p+m-1}} \qquad \left(g'(z) = z^{m-1}h'(z)\right), \tag{4.5}$$

then the function $\varphi(t)$ given by (2.3) satisfies

$$\varphi'(t) = iz \left(h'(z) - \overline{z^{m+1}} h'(z) \right)$$

= $iz \left(\frac{pz^{p-1}}{1 + z^{2p+m-1}} - \frac{p\overline{z}^{p+m}}{1 + \overline{z}^{2p+m-1}} \right)$
= $iz \left(\frac{pz^{p-1}}{1 + z^{2p+m-1}} - \frac{pz^{p-1}}{1 + z^{2p+m-1}} \right) = 0 \qquad (z = e^{it})$ (4.6)

for any $z^{2p+m-1} = e^{i(2p+m-1)t} \neq -1$. Thus, the function

$$f(z) = h(z) + \overline{g(z)} = \int_0^z \frac{p\zeta^{p-1}}{1 + \zeta^{2p+m-1}} d\zeta + \overline{\int_0^z \frac{p\zeta^{p+m-2}}{1 + \zeta^{2p+m-1}}} d\zeta$$
(4.7)

is a member of the class $S_{\mathcal{H}}(p)$ and it maps \mathbb{U} onto a domain surrounded by 2p + m - 1 straight lines with 2p + m - 1 cusps. Indeed, setting p = 2 and m = 2, we know that

$$f(z) = \int_0^z \frac{2\zeta}{1+\zeta^5} d\zeta + \overline{\int_0^z \frac{2\zeta^2}{1+\zeta^5} d\zeta}$$
(4.8)

is a 2-valently harmonic function and it maps \mathbb{U} onto a star as shown in Figure 3. Furthermore, if we take p = 1 in (4.7), then we see that the function

$$f_{m+1}(z) = h(z) + \overline{g(z)} = \int_0^z \frac{1}{1+\zeta^{m+1}} d\zeta + \overline{\int_0^z \frac{\zeta^{m-1}}{1+\zeta^{m+1}} d\zeta}$$
(4.9)

is univalent in \mathbb{U} and it maps \mathbb{U} onto a (m+1)-sided polygon. For example, the function

$$f_8(z) = \int_0^z \frac{1}{1+\zeta^8} d\zeta + \overline{\int_0^z \frac{\zeta^6}{1+\zeta^8} d\zeta} \qquad (m=7)$$
(4.10)

maps \mathbb{U} onto an octagon (schlicht domain) as shown in Figure 4.

 $\mathbf{6}$




FIGURE 4. The image of $f_8(z) = \int_0^z \frac{1}{1+\zeta^8} d\zeta + \overline{\int_0^z \frac{\zeta^6}{1+\zeta^8} d\zeta}.$

TOSHIO HAYAMI

References

- [1] D. Bshouty and A. Lyzzaik, *Close-to-convexity criteria for planar harmonic mappings*, Complex Anal. Oper. Theory **5**(2011), 767–774.
- [2] J. Clunie and T. Sheil-Small, Harmonic univalent functions, Ann. Acad. Sci. Fenn. Ser. A I Math. 9(1984), 3–25.

[3] P. L. Duren, *Harmonic Mappings in the Plane*, Cambridge University Press, Cambridge, 2004.

[4] A. W. Goodman, Univalent Functions, Vol. I and Vol. II, Mariner Publishing Company, Tampa, Florida (1983).

- [5] T. Hayami and S. Owa, Hypocycloid of n + 1 cusps harmonic function, Bull. Math. Anal. Appl. 3(2011), 239–246.
- [6] H. Lewy, On the non-vanishing of the Jacobian in certain one-to-one mappings, Bull. Amer. Math. Soc. 42(1936), 689–692.

[7] P. T. Mocanu, Three-cornered hat harmonic functions, Complex Var. Elliptic Equ. 54(2009), 1079–1084.

[8] P. T. Mocanu, Injectively conditions in the complex plane, Complex Anal. Oper. Theory 5(2011), 759–766.

Toshio Hayami Department of Mathematics, Kinki University Higashi-Osaka, Osaka 577-8502, Japan *E-mail address*: ha_ya_to112@hotmail.com