# COEFFICIENT CONDITIONS FOR HARMONIC CLOSE-TO-CONVEX FUNCTIONS

TOSHIO HAYAMI

ABSTRACT. New sufficient conditions, concerned with the coefficients of harmonic functions  $f(z) = h(z) + \overline{g(z)}$  in the open unit disk U normalized by f(0) = h(0) = h'(0) - 1 = 0, for f(z) to be harmonic close-to-convex functions are discussed. Furthermore, several illustrative examples and the image domains of harmonic close-to-convex functions satisfying the obtained conditions are enumerated.

#### 1. INTRODUCTION

For a continuous complex-valued function f(z) = u(x, y) + iv(x, y) (z = x + iy), we say that f(z) is harmonic in the open unit disk  $\mathbb{U} = \{z \in \mathbb{C} : |z| < 1\}$  if both u(x, y) and v(x, y) are real harmonic in  $\mathbb{U}$ , that is, u(x, y) and v(x, y) satisfy the Laplace's equations

$$\Delta u = u_{xx} + u_{yy} = 0 \qquad \text{and} \qquad \Delta v = v_{xx} + v_{yy} = 0$$

A complex-valued harmonic function f(z) in  $\mathbb{U}$  is given by f(z) = h(z) + g(z) where h(z) and g(z) are analytic in  $\mathbb{U}$ . We call h(z) and g(z) the analytic part and the co-analytic part of f(z), respectively. A necessary and sufficient condition for f(z) to be locally univalent and sense-preserving in  $\mathbb{U}$  is |h'(z)| >|g'(z)| in  $\mathbb{U}$  (see, [2] or [8]). Let  $\mathcal{H}$  denote the class of harmonic functions f(z) in  $\mathbb{U}$  with f(0) = h(0) = 0and h'(0) = 1. Thus, every normalized harmonic function f(z) can be written by

$$f(z) = h(z) + \overline{g(z)} = z + \sum_{n=2}^{\infty} a_n z^n + \sum_{n=1}^{\infty} b_n z^n \in \mathcal{H}$$

where  $a_1 = 1$  and  $b_0 = 0$ , for convenience.

We next denote by  $S_{\mathcal{H}}$  the class of functions  $f(z) \in \mathcal{H}$  which are univalent and sense-preserving in  $\mathbb{U}$ . Since the sense-preserving property of f(z), we see that  $|b_1| = |g'(0)| < |h'(0)| = 1$ . If  $g(z) \equiv 0$ , then  $S_{\mathcal{H}}$  reduces to the class S consisting of normalized analytic univalent functions. Furthermore, for every function  $f(z) \in S_{\mathcal{H}}$ , the function

$$F(z) = \frac{f(z) - \overline{b_1 f(z)}}{1 - |b_1|^2} = z + \sum_{n=2}^{\infty} \frac{a_n - \overline{b_1} b_n}{1 - |b_1|^2} z^n + \sum_{n=2}^{\infty} \frac{b_n - b_1 a_n}{1 - |b_1|^2} z^n$$

<sup>2010</sup> Mathematics Subject Classification. Primary 30C45, Secondary 58E20.

Key words and phrases. Coefficient condition, harmonic function, univalent function, close-to-convex function.

## TOSHIO HAYAMI

is also a member of  $\mathcal{S}_{\mathcal{H}}$ . Therefore, we consider the subclass  $\mathcal{S}_{\mathcal{H}}^0$  of  $\mathcal{S}_{\mathcal{H}}$  defined as

$$S^0_{\mathcal{H}} = \{f(z) \in S_{\mathcal{H}} : b_1 = g'(0) = 0\}.$$

Conversely, if  $F(z) \in S^0_{\mathcal{H}}$ , then  $f(z) = F(z) + \overline{b_1 F(z)} \in S_{\mathcal{H}}$  for any  $b_1$  ( $|b_1| < 1$ ).

We say that a domain  $\mathbb{D}$  is a close-to-convex domain if the complement of  $\mathbb{D}$  can be written as a union of non-intersecting half-lines (except that the origin of one half-line may lie on one of the other half-lines). Let  $\mathcal{C}$ ,  $\mathcal{C}_{\mathcal{H}}$  and  $\mathcal{C}^{0}_{\mathcal{H}}$  be the respective subclasses of  $\mathcal{S}$ ,  $\mathcal{S}_{\mathcal{H}}$  and  $\mathcal{S}^{0}_{\mathcal{H}}$  consisting of all functions f(z) which map  $\mathbb{U}$  onto a certain close-to-convex domain.

Bshouty and Lyzzaik [1] have stated the following result.

**Theorem 1.1.** If  $f(z) = h(z) + \overline{g(z)} \in \mathcal{H}$  satisfies

$$g'(z) = zh'(z)$$
 and  $\operatorname{Re}\left(1 + \frac{zh''(z)}{h'(z)}\right) > -\frac{1}{2}$ 

for all  $z \in \mathbb{U}$ , then  $f(z) \in \mathcal{C}^0_{\mathcal{H}} \subset \mathcal{S}^0_{\mathcal{H}}$ .

A simple and interesting example is below.

Example 1.1. The function

$$f(z) = \frac{1 - (1 - z)^2}{2(1 - z)^2} + \frac{z^2}{2(1 - z)^2} = z + \sum_{n=2}^{\infty} \frac{n + 1}{2} z^n + \sum_{n=2}^{\infty} \frac{n - 1}{2} \overline{z}^n$$

satisfies the conditions of Theorem 1.1, and therefore f(z) belongs to the class  $\mathcal{C}^{0}_{\mathcal{H}}$ . We now show that  $f(\mathbb{U})$  is actually close-to-convex domain. It follows that

$$f(z) = \left(\frac{z}{2(1-z)^2} + \frac{z}{2(1-z)}\right) + \overline{\left(\frac{z}{2(1-z)^2} - \frac{z}{2(1-z)}\right)} \\ = \operatorname{Re}\left(\frac{z}{(1-z)^2}\right) + i\operatorname{Im}\left(\frac{z}{1-z}\right).$$

Setting

$$f(re^{i\theta}) = \frac{-2r^2 + r(1+r^2)\cos\theta}{(1+r^2 - 2r\cos\theta)^2} + \frac{r\sin\theta}{1+r^2 - 2r\cos\theta}i = u + iv$$

for any  $z = re^{i\theta} \in \mathbb{U}$   $(0 \leq r < 1, 0 \leq \theta < 2\pi)$ , we see that

$$-4(u+v^2) = \frac{4r(r-\cos\theta)(1-r\cos\theta)}{(1+r^2-2r\cos\theta)^2} = \frac{4r(r-t)(1-rt)}{(1+r^2-2rt)^2} \equiv \phi(t) \qquad (-1 \le t = \cos\theta \le 1).$$

Since

$$\phi'(t) = \frac{-4r(1-r^2)^2}{(1+r^2-2rt)^3} \le 0,$$

we obtain that

$$\phi(t) \leq \phi(-1) = \frac{4r}{(1+r)^2} \equiv \psi(r)$$

Also, noting that

$$\psi'(r) = \frac{4(1-r)}{(1+r)^3} > 0,$$

we know that

$$\psi(r) < \psi(1) = 1$$

which implies that

$$u > -v^2 - \frac{1}{4}.$$

Thus, f(z) maps  $\mathbb{U}$  onto the following close-to-convex domain.



Remark 1.1. Let  $\mathcal{M}$  be the class of all functions satisfying the conditions of Theorem 1.1. Then, it was earlier conjectured by Mocanu [9, 10] that  $\mathcal{M} \subset S^0_{\mathcal{H}}$ . Furthermore, we can immediately see that the function f(z) in Example 1.1 is a member of the class  $\mathcal{M}$  and it shows that  $f(z) \in \mathcal{M}$  is not necessarily starlike with respect to the origin in  $\mathbb{U}$  (f(z) is starlike with respect to the origin in  $\mathbb{U}$  if and only if  $tw \in f(\mathbb{U})$  for all  $w \in f(\mathbb{U})$  and t ( $0 \leq t \leq 1$ )).

Remark 1.2. For the function  $f(z) = h(z) + \overline{g(z)} \in \mathcal{H}$  given by

$$g'(z) = z^{n-1}h'(z)$$
  $(n = 2, 3, 4, \cdots),$ 

letting  $w(t) = f(e^{it}) = h(e^{it}) + \overline{g(e^{it})}$   $(-\pi \leq t < \pi)$ , we know that

$$\operatorname{Im}\left(\frac{w''(t)}{w'(t)}\right) \leq 0 \qquad (-\pi \leq t < \pi)$$

which means that f(z) maps the unit circle  $\partial \mathbb{U} = \{z \in \mathbb{C} : |z| = 1\}$  onto a union of several concave curves (see, [6, Theorem 2.1]).

Jahangiri and Silverman [7] have given the following coefficient inequality for  $f(z) \in \mathcal{H}$  to be in the class  $\mathcal{C}_{\mathcal{H}}$ .

**Theorem 1.2.** If  $f(z) \in \mathcal{H}$  satisfies

$$\sum_{n=2}^{\infty} n|a_n| + \sum_{n=1}^{\infty} n|b_n| \le 1,$$

then  $f(z) \in \mathcal{C}_{\mathcal{H}}$ .

Example 1.2. The function

$$f(z) = z + \frac{1}{5}\overline{z}^5$$

belongs to the class  $\mathcal{C}^0_{\mathcal{H}} \subset \mathcal{C}_{\mathcal{H}}$  and satisfies the condition of Theorem 1.2. Indeed, f(z) maps  $\mathbb{U}$  onto the following hypocycloid of six cusps (cf. [3] or [6]).



FIGURE 2. The image of  $f(z) = z + \frac{1}{5}\overline{z}^5$ .

The object of this paper is to find some sufficient conditions for functions  $f(z) \in \mathcal{H}$  to be in the class  $\mathcal{C}_{\mathcal{H}}$ . In order to establish our results, we have to recall here the following lemmas due to Clunie and Sheil-small [2].

**Lemma 1.1.** If h(z) and g(z) are analytic in  $\mathbb{U}$  with |h'(0)| > |g'(0)| and  $h(z) + \varepsilon g(z)$  is close-to-convex for each  $\varepsilon$  ( $|\varepsilon| = 1$ ), then  $f(z) = h(z) + \overline{g(z)}$  is harmonic close-to-convex.

**Lemma 1.2.** If  $f(z) = h(z) + \overline{g(z)}$  is locally univalent in  $\mathbb{U}$  and  $h(z) + \varepsilon g(z)$  is convex for some  $\varepsilon$   $(|\varepsilon| \leq 1)$ , then f(z) is univalent close-to-convex.

We also need the following result due to Hayami, Owa and Srivastava [5].

Lemma 1.3. If a function 
$$F(z) = z + \sum_{n=2}^{\infty} A_n z^n$$
 is analytic in  $\mathbb{U}$  and satisfies  

$$\sum_{n=2}^{\infty} \left[ \left| \sum_{k=1}^n \left\{ \sum_{j=1}^k (-1)^{k-j} j(j+1) \begin{pmatrix} \alpha \\ k-j \end{pmatrix} A_j \right\} \begin{pmatrix} \beta \\ n-k \end{pmatrix} \right| + \left| \sum_{k=1}^n \left\{ \sum_{j=1}^k (-1)^{k-j} j(j-1) \begin{pmatrix} \alpha \\ k-j \end{pmatrix} A_j \right\} \begin{pmatrix} \beta \\ n-k \end{pmatrix} \right| \right] \leq 2$$

for some real numbers  $\alpha$  and  $\beta$ , then F(z) is convex in  $\mathbb{U}$ .

#### 2. Main results

Our first result is contained in

**Theorem 2.1.** If  $f(z) \in \mathcal{H}$  satisfies the following condition

$$\sum_{n=2}^{\infty} \left| na_n - e^{i\varphi}(n-1)a_{n-1} \right| + \sum_{n=1}^{\infty} \left| nb_n - e^{i\varphi}(n-1)b_{n-1} \right| \le 1$$

for some real number  $\varphi$   $(0 \leq \varphi < 2\pi)$ , then  $f(z) \in C_{\mathcal{H}}$ .

*Proof.* Let 
$$F(z) = z + \sum_{n=2}^{\infty} A_n z^n$$
 be analytic in U. If  $F(z)$  satisfies  
$$\sum_{n=2}^{\infty} \left| nA_n - e^{i\varphi}(n-1)A_{n-1} \right| \leq 1$$

then it follows that

$$\begin{aligned} \left| (1 - e^{i\varphi}z)F'(z) - 1 \right| &= \left| \sum_{n=2}^{\infty} \left( nA_n - e^{i\varphi}(n-1)A_{n-1} \right) z^{n-1} \right| \\ &\leq \sum_{n=2}^{\infty} \left| nA_n - e^{i\varphi}(n-1)A_{n-1} \right| \cdot |z|^{n-1} \\ &< \sum_{n=2}^{\infty} \left| nA_n - e^{i\varphi}(n-1)A_{n-1} \right| \leq 1 \quad (z \in \mathbb{U}). \end{aligned}$$

This gives us that

$$\operatorname{Re}\left((1-e^{i\varphi}z)F'(z)\right) > 0 \qquad (z \in \mathbb{U}).$$

that is, that  $F(z) \in \mathcal{C}$ . Then, it is sufficient to prove that

$$F(z) = \frac{h(z) + \varepsilon g(z)}{1 + \varepsilon b_1} = z + \sum_{n=2}^{\infty} \frac{a_n + \varepsilon b_n}{1 + \varepsilon b_1} z^n \in \mathcal{C}$$

#### TOSHIO HAYAMI

for each  $\varepsilon$  ( $|\varepsilon| = 1$ ) by Lemma 1.1. From the assumption of the theorem, we obtain that

$$\begin{split} \sum_{n=2}^{\infty} \left| n \frac{a_n + \varepsilon b_n}{1 + \varepsilon b_1} - e^{i\varphi} (n-1) \frac{a_{n-1} + \varepsilon b_{n-1}}{1 + \varepsilon b_1} \right| \\ & \leq \frac{1}{1 - |b_1|} \sum_{n=2}^{\infty} \left[ \left| na_n - e^{i\varphi} (n-1)a_{n-1} \right| + \left| nb_n - e^{i\varphi} (n-1)b_{n-1} \right| \right] \leq \frac{1 - |b_1|}{1 - |b_1|} = 1. \end{split}$$
This completes the proof of the theorem.

This completes the proof of the theorem.

## Example 2.1. The function

$$f(z) = -\log(1-z) + \overline{\left(-mz - \log(1-z)\right)} = z + \sum_{n=2}^{\infty} \frac{1}{n} z^n + (1-m)\overline{z} + \sum_{n=2}^{\infty} \frac{1}{n} \overline{z}^n \quad (0 < m \le 1)$$

satisfies the condition of Theorem 2.1 with  $\varphi = 0$  and belongs to the class  $C_{\mathcal{H}}$ . In particular, putting m = 1, we obtain the following.



FIGURE 3. The image of  $f(z) = -\overline{z} - 2\log|1-z|$ .

By making use of Lemma 1.2 with  $\varepsilon = 0$  and applying Lemma 1.3, we readily obtain the next theorem. **Theorem 2.2.** If  $f(z) \in \mathcal{H}$  is locally univalent in  $\mathbb{U}$  and satisfies

$$\sum_{n=2}^{\infty} \left[ \left| \sum_{k=1}^{n} \left\{ \sum_{j=1}^{k} (-1)^{k-j} j(j+1) \begin{pmatrix} \alpha \\ k-j \end{pmatrix} a_j \right\} \begin{pmatrix} \beta \\ n-k \end{pmatrix} \right| + \left| \sum_{k=1}^{n} \left\{ \sum_{j=1}^{k} (-1)^{k-j} j(j-1) \begin{pmatrix} \alpha \\ k-j \end{pmatrix} a_j \right\} \begin{pmatrix} \beta \\ n-k \end{pmatrix} \right| \right] \leq 2$$

for some real numbers  $\alpha$  and  $\beta$ , then  $f(z) \in C_{\mathcal{H}}$ .

Putting  $\alpha = \beta = 0$  in the above theorem, we arrive at the following result due to Jahangiri and Silverman [7].

**Theorem 2.3.** If  $f(z) \in \mathcal{H}$  is locally univalent in  $\mathbb{U}$  with

$$\sum_{n=2}^{\infty} n^2 |a_n| \le 1$$

then  $f(z) \in \mathcal{C}_{\mathcal{H}}$ .

Furthermore, taking  $\alpha = 1$  and  $\beta = 0$  in the theorem, we have

**Corollary 2.1.** If  $f(z) \in \mathcal{H}$  is locally univalent in  $\mathbb{U}$  and satisfies

$$\sum_{n=2}^{\infty} \left\{ n \left| (n+1)a_n - (n-1)a_{n-1} \right| + (n-1) \left| na_n - (n-2)a_{n-1} \right| \right\} \le 2,$$

then  $f(z) \in \mathcal{C}_{\mathcal{H}}$ .

# Example 2.2. The function

$$f(z) = -\int_0^z \frac{\log(1-t)}{t} dt + \overline{\left(z + (1-z)\log(1-z)\right)} = z + \sum_{n=2}^\infty \frac{1}{n^2} z^n + \sum_{n=2}^\infty \frac{1}{n(n-1)} \overline{z}^n$$

satisfies the conditions of Corollary 2.1 and belongs to the class  $C_{\mathcal{H}}$ .



FIGURE 4. The image of  $f(z) = -\int_0^z \frac{\log(1-t)}{t} dt + \overline{\left(z + (1-z)\log(1-z)\right)}.$ 

#### TOSHIO HAYAMI

## 3. Appendix

A sequence  $\{c_n\}_{n=0}^{\infty}$  of non-negative real numbers is called a convex null sequence if  $c_n \to 0$  as  $n \to \infty$ and

$$c_n - c_{n+1} \geqq c_{n+1} - c_{n+2} \geqq 0$$

for all  $n \ (n = 0, 1, 2, \cdots)$ .

The next lemma was obtained by Fejér [4].

**Lemma 3.1.** Let  $\{c_n\}_{k=0}^{\infty}$  be a convex null sequence. Then, the function

$$p(z) = \frac{c_0}{2} + \sum_{n=1}^{\infty} c_n z^n$$

is analytic and satisfies  $\operatorname{Re}(p(z)) > 0$  in  $\mathbb{U}$ .

Applying the above lemma, we deduce

**Theorem 3.1.** For some b (|b| < 1) and some convex null sequence  $\{c_n\}_{n=0}^{\infty}$  with  $c_0 = 2$ , the function

$$f(z) = h(z) + \overline{g(z)} = z + \sum_{n=2}^{\infty} \frac{c_{n-1}}{n} z^n + b\left(z + \sum_{n=2}^{\infty} \frac{c_{n-1}}{n} z^n\right)$$

belongs to the class  $C_{\mathcal{H}}$ .

*Proof.* Let us define F(z) by

$$F(z) = \frac{h(z) + \varepsilon g(z)}{1 + \varepsilon b} = z + \sum_{n=2}^{\infty} \frac{c_{n-1}}{n} z^n$$

for each  $\varepsilon$  ( $|\varepsilon| = 1$ ). Then, we know that

$$F'(z) = \frac{c_0}{2} + \sum_{n=1}^{\infty} c_n z^n \qquad (c_0 = 2).$$

By virtue of Lemma 1.1 and Lemma 3.1, it follows that  $\operatorname{Re}(F'(z)) > 0$   $(z \in \mathbb{U})$ , that is,  $F(z) \in \mathcal{C}$ . Thus, we conclude that  $f(z) = h(z) + \overline{g(z)} \in \mathcal{C}_{\mathcal{H}}$ .

In the same manner, we also have

**Theorem 3.2.** For some b (|b| < 1) and some convex null sequence  $\{c_n\}_{n=0}^{\infty}$  with  $c_0 = 2$ , the function

$$f(z) = h(z) + \overline{g(z)} = z + \sum_{n=2}^{\infty} \frac{1}{n} \left( 1 + \sum_{j=1}^{n-1} c_j \right) z^n + b \left( z + \sum_{n=2}^{\infty} \frac{1}{n} \left( 1 + \sum_{j=1}^{n-1} c_j \right) z^n \right)$$

belongs to the class  $C_{\mathcal{H}}$ .

*Proof.* Let us define F(z) by

$$F(z) = \frac{h(z) + \varepsilon g(z)}{1 + \varepsilon b} = z + \sum_{n=2}^{\infty} \frac{1}{n} \left( 1 + \sum_{j=1}^{n-1} c_j \right) z^n$$

for each  $\varepsilon$  ( $|\varepsilon| = 1$ ). Then, we know that

$$(1-z)F'(z) = \frac{c_0}{2} + \sum_{n=1}^{\infty} c_n z^n \qquad (c_0 = 2).$$

Therefore, by the help of Lemma 1.1 and Lemma 3.1, we obtain that  $\operatorname{Re}\left((1-z)F'(z)\right) > 0$   $(z \in \mathbb{U})$ , that is,  $F(z) \in \mathcal{C}$  which implies that  $f(z) = h(z) + \overline{g(z)} \in \mathcal{C}_{\mathcal{H}}$ .

*Remark* 3.1. The sequence

$$\{c_n\}_{n=0}^{\infty} = \left\{2, 1, \frac{2}{3}, \cdots, \frac{2}{n+1}, \cdots\right\}$$

is a convex null sequence because

$$\lim_{n \to \infty} c_n = \lim_{n \to \infty} \left( \frac{2}{n+1} \right) = 0, \qquad c_n - c_{n+1} = \frac{2}{(n+1)(n+2)} \ge 0$$

and

$$(c_n - c_{n+1}) - (c_{n+1} - c_{n+2}) = \frac{4}{(n+1)(n+2)(n+3)} \ge 0 \qquad (n = 0, 1, 2, \cdots).$$

Setting  $b = \frac{1}{4}$  in Theorem 3.1 with the above sequence  $\{c_n\}_{n=0}^{\infty}$ , we derive

Example 3.1. The function

$$f(z) = -z - 2\int_0^z \frac{\log(1-t)}{t} dt - \frac{1}{4}\left(z + 2\int_0^z \frac{\log(1-t)}{t} dt\right) = z + \sum_{n=2}^\infty \frac{2}{n^2} z^n + \frac{1}{4}\left(z + \sum_{n=2}^\infty \frac{2}{n^2} z^n\right)$$

is in the class  $C_{\mathcal{H}}$ .



FIGURE 5. The image of f(z) in Example 3.1.

Moreover, we know

*Remark* 3.2. The sequence

$$\{c_n\}_{n=0}^{\infty} = \left\{2, 1, \frac{1}{2}, \cdots, 2^{1-n}, \cdots\right\}$$

is a convex null sequence because

$$\lim_{n \to \infty} c_n = \lim_{n \to \infty} 2^{1-n} = 0, \qquad c_n - c_{n+1} = 2^{-n} \ge 0$$

 $\quad \text{and} \quad$ 

$$(c_n - c_{n+1}) - (c_{n+1} - c_{n+2}) = 2^{-(n+1)} \ge 0$$
  $(n = 0, 1, 2, \cdots).$ 

Hence, letting  $b = \frac{1}{4}$  in Theorem 3.2 with the sequence  $\{c_n\}_{n=0}^{\infty} = \{2^{1-n}\}_{n=0}^{\infty}$ , we have **Example 3.2.** The function

$$f(z) = -3\log(1-z) + 4\log\left(1-\frac{z}{2}\right) + \overline{\left(-\frac{3}{4}\log(1-z) + \log\left(1-\frac{z}{2}\right)\right)}$$
$$= z + \sum_{n=2}^{\infty} \frac{1}{n} \left(1 + \sum_{j=1}^{n-1} 2^{1-j}\right) z^n + \overline{\frac{1}{4} \left(z + \sum_{n=2}^{\infty} \frac{1}{n} \left(1 + \sum_{j=1}^{n-1} 2^{1-j}\right) z^n\right)}$$

is in the class  $\mathcal{C}_{\mathcal{H}}$ .



FIGURE 6. The image of f(z) in Example 3.2.

10

#### References

[1] D. Bshouty and A. Lyzzaik, *Close-to-convexity criteria for planar harmonic mappings*, Complex Anal. Oper. Theory **5**(2011), 767–774.

[2] J. Clunie and T. Sheil-Small, Harmonic univalent functions, Ann. Acad. Sci. Fenn. Ser. A I Math. 9(1984), 3–25.

[3] P. L. Duren, Harmonic Mappings in the Plane, Cambridge University Press, Cambridge, 2004.

 [4] L. Fejér, Über die positivität von summen, die nach trigonometrischen oder Legendreschen, funktionen fortschreiten. I, Acta Szeged 2(1925), 75-86(German).

 [5] T. Hayami, S. Owa and H. M. Srivastava, Coefficient inequalities for certain classes of analytic and univalent functions, J. Inequal. Pure Appl. Math. 8(2007), Article 95, 1-10.

[6] T. Hayami and S. Owa, Hypocycloid of n + 1 cusps harmonic function, Bull. Math. Anal. Appl. 3(2011), 239-246.

[7] J. M. Jahangiri and H. Silverman, Harmonic close-to-convex mappings, J. Appl. Math. Stoch. Anal. 15(2002), 23-28.

[8] H. Lewy, On the non-vanishing of the Jacobian in certain one-to-one mappings, Bull. Amer. Math. Soc. 42(1936), 689–692.

[9] P. T. Mocanu, Three-cornered hat harmonic functions, Complex Var. Elliptic Equ. 54(2009), 1079–1084.

[10] P. T. Mocanu, Injectively conditions in the complex plane, Complex Anal. Oper. Theory 5(2011), 759–766.

Toshio Hayami Department of Mathematics, Kinki University Higashi-Osaka, Osaka 577-8502, Japan

E-mail address: ha\_ya\_to112@hotmail.com