

ON THE PROBLEM OF CLASSIFYING SIMPLE COMPACT QUANTUM GROUPS

SHUZHOU WANG

Dedicated to Professor S. L. Woronowicz on the Occasion of his 70th Birthday

ABSTRACT. We review the notion of simple compact quantum groups and examples, and discuss the problem of construction and classification of simple compact quantum groups. Several new quantum groups constructed by Banica, Curran and Speicher since the author's first paper on simple quantum groups are shown to be simple using results of Raum and Weber.

1. INTRODUCTION

There are two main areas in the operator algebraic approach to quantum groups: compact quantum groups and locally compact quantum groups. The former has a satisfactory axiomatic theory due to Woronowicz [83, 85]. While the latter has witnessed great progress due to concerted efforts of several generations of mathematicians (cf. an incomplete list including [47, 48, 41, 42, 1, 50, 54]), it still does not have an axiomatic framework that contains the non-compact Drinfeld-Jimbo quantum groups [40] as examples except for special cases such as $SL_q(2, \mathbb{C})$ (cf. e.g. [55]), nor has the existence of Haar weight been established in general. This is in stark contrast with the fact that compact real forms of the Drinfeld-Jimbo quantum groups are special examples in the theory of compact quantum groups (see [59, 60, 62, 61, 51]), and Haar measure, Peter-Weyl theorem and Tannaka-Krein duality can be established from a very simple set of axioms [83, 84, 85]. Within this well established framework of compact quantum groups, recent work on compact quantum groups has been primarily on the construction, classification, structure and other (operator) algebraic properties of specific classes of compact quantum groups.

The modern impetus in the theory of quantum groups came as a result of the discovery of new examples of Hopf algebras by Drinfeld-Jimbo on the algebraic side [40] and

2010 *Mathematics Subject Classification*. Primary 46L65; Secondary 17B37, 20G42, 58B32, 46L87, 46L55, 46L60, 46L89, 16T05, 81R50, 81R60.

Key words and phrases. simple quantum groups, quantum permutation groups, quantum automorphism groups, compact quantum groups, Woronowicz C^* -algebras, Hopf Algebras.

Woronowicz on the analytic side [82]. These are deformation quantizations of the classical Lie algebras and Lie groups, and much of the literature on quantum groups had been devoted to this approach to quantum groups.

Starting in his thesis [70], the author took a different direction than the traditional deformation quantization method by viewing quantum groups as intrinsic objects and found in a series of papers (including [69] in collaboration with Van Daele) several classes of universal compact quantum groups that *can not be obtained* as deformations of Lie groups or Lie algebras, the most important of these are the universal compact quantum groups of Kac type $A_u(n)$ and their orthogonal counterpart $A_o(n)$ [71], the more general universal compact quantum groups $A_u(Q)$ and their self-conjugate counterpart $B_u(Q)$ [69, 74], where $Q \in GL(n, \mathbb{C})$, and the quantum automorphism groups $A_{aut}(B, tr)$ [75] of finite dimensional C^* -algebras B endowed with a tracial functional tr , including the quantum permutation groups $A_{aut}(X_n)$ on the space X_n of n points. These objects have been an international focus of study in the subject of compact quantum groups and interest in them continues unabated (cf. [2]-[35], [44], [49], [63]-[68]).

The quantum groups $A_u(Q)$ have the remarkable universal property that can be used to give following alternative and *concrete definition* of compact matrix quantum groups that was originally defined by Woronowicz [83] more abstractly: a compact matrix quantum group is a quotient $A_u(Q)/I$, where I is a Woronowicz C^* -ideal in $A_u(Q)$. This means in geometric language that every compact matrix quantum group (including compact Lie group) is a quantum subgroup of $A_u(Q)$ for an appropriate choice of Q . In contrast, Drinfeld-Jimbo quantum groups and other deformations of Lie groups do not enjoy this property, but are quantum subgroups of $A_u(Q)$. Similarly, the quantum groups $B_u(Q)$ have the universal property that every compact matrix quantum group with self-conjugate fundamental representation is of the form $B_u(Q)/I$, where I is a Woronowicz C^* -ideal in $B_u(Q)$.

Without universal compact quantum groups, the Drinfeld-Jimbo quantum groups and other quantum groups obtained by deformation would be the end of the story. However, the outpouring of papers on the universal quantum groups in the last few years (e.g. [2]-[35], [44], [49], [63]-[68]) demonstrates depth of the subject: despite much work achieved so far, we have only seen the tip of the iceberg and the story is far from the end.

Although compact quantum groups have a satisfactory axiomatic framework and Drinfeld-Jimbo quantum groups are their special examples, after the discovery of these new classes of universal compact quantum groups, it is a natural program to classify simple compact

quantum groups. This program was initiated in [79], where it was shown that all compact quantum groups mentioned above are simple in generic cases.

The *main goals* of the program on simple compact quantum groups are: (1) construct and classify simple compact quantum groups and their irreducible representations, (2) understand the structure of simple compact quantum groups and structure of compact quantum groups in terms of the simple ones, and (3) develop new applications of simple compact quantum groups in other areas of mathematics and physics, such as quantum symmetries in noncommutative geometry and algebraic quantum field theory. For goal (1), one would like to develop a theory of simple compact quantum groups that parallels the Killing-Cartan theory and the Cartan-Weyl theory for simple compact Lie groups. For this purpose, one must first construct all simple compact quantum groups. Though the work so far provides several infinite classes of examples of these, it should be pointed out that the construction of simple compact quantum groups is only at the beginning stage for this task at the moment, as all the simple compact quantum groups known so far are almost classical in the sense that their representation rings are isomorphic to those of ordinary compact groups and in particular are commutative. The first examples of simple compact quantum groups that are not almost classical should be directly related to the universal quantum groups $A_u(Q)$ (see the footnote after Problem 4.1), where $Q \in GL(n, \mathbb{C})$ are positive, $n \geq 2$, though these quantum groups are not simple themselves. The representation ring of $A_u(Q)$ is highly noncommutative, being roughly the free product of two copies of the ring of integers, according to Banica [3]. To construct other simple compact quantum groups, the most natural idea is to study quantum automorphism groups of appropriate quantum spaces, such as those in the author's papers [75, 76], the papers of Banica, Bichon, Goswami and their collaborators [5]-[11], [32, 33], [44], [24]-[31]. In retrospect, both simple Lie groups and finite simple groups are automorphism groups. This suggests viability of this approach to the program. Natural mathematical and physical structures that have compact quantum automorphic group symmetries are compact commutative and noncommutative Riemannian manifolds in the sense of Connes [36, 38]. Such symmetries should be investigated first.

The following heuristic may indicate the depth of the problem on classification problem of simple compact quantum groups. The finite dimensional factors are classified by the *discrete set* of natural numbers while the classification of infinite dimensional von Neumann factors involves *continuous parameters*. Similarly, simple compact Lie groups are classified by a *discrete set* of Cartan matrices, but the classification of simple compact quantum groups involves *continuous parameters*. However, since the algebraic structures

of compact quantum groups are richer and more rigid than those of von Neumann algebras, the classification of simple compact quantum groups might be more accessible than the classification of infinite dimensional factors. Even if a classification of simple compact quantum groups up to isomorphism is unattainable, just as von Neumann factors are far from being classified up to isomorphism, experience has demonstrated that the study of universal quantum groups and quantum automorphism groups is fruitful (cf. [71, 69, 74, 76, 78, 79], [2]-[35], [44], [49], [63]-[68] and references therein), and other types of classification theories may also be considered, such as the classification of easy quantum groups [21, 18, 19, 81] and the classification of restricted classes of quantum automorphism groups [8]-[12] by Banica *et al.*

An outline of the paper is as follows. In §2, we review the notion of compact quantum groups and simple compact quantum groups. In §3, we review examples of simple compact quantum groups constructed so far in [79]. In §4, we discuss problem of the fine structure of $A_u(Q)$ and simple quantum quotient groups from $A_u(Q)$, as well as quantum subgroups from free products of compact quantum groups. In §5, we give a list of problems related to almost classical compact quantum groups and compact quantum groups with property F . In §6 and §7, we discuss the problem of constructing simple compact quantum groups from quantum automorphisms of finite graphs and other quantum subgroups of the quantum permutation groups, easy quantum groups and quantum isometry groups. In §6, several new quantum groups constructed by Banica, Curran and Speicher [17] since [79] are shown to be simple using results in Raum [56] and Weber [81] along with results in [79].

2. COMPACT QUANTUM GROUPS AND SIMPLE COMPACT QUANTUM GROUPS

We first recall the definition of compact (matrix) quantum groups and then the notion of simple compact quantum groups.

There are several equivalent definitions of compact (matrix) quantum groups, each has its own advantages over the others. We briefly describe below another equivalent definition, which has the advantage of letting the reader to “visualize” all compact quantum groups more concretely. This equivalent definition is essentially in the literature, but not in the explicit form we describe below.

Notation: For elements u_{ij} ($i, j = 1, \dots, n$) of a C^* -algebra A , we define the following elements in the $n \times n$ matrix algebra $M_n(A)$ over A : $u := (u_{ij})_{i,j=1}^n$, $\bar{u} := (u_{ij}^*)_{i,j=1}^n$, $u^t := (u_{ji})_{i,j=1}^n$ and $u^* := \bar{u}^t$, i.e. $u^* = (u_{ji}^*)_{i,j=1}^n$.

Definition 2.1. (cf. [70, 71, 69, 74, 78]) The **universal compact matrix quantum groups** are defined to be the family of pairs $(A_u(Q), \Delta_u)$, where $Q \in GL(n, \mathbb{C})$, $Q > 0$, and u_{ij} ($i, j = 1, \dots, n$) are generators of the universal C^* -algebra $A_u(Q)$ that satisfies the following sets of relations:

$$u^*u = I_n = uu^*, \quad u^t Q \bar{u} Q^{-1} = I_n = Q \bar{u} Q^{-1} u^t,$$

and $\Delta_u : A_u(Q) \rightarrow A_u(Q) \otimes A_u(Q)$ is the uniquely defined morphism such that

$$\Delta_u(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

Note that instead of restricting Q to positive matrices, one can still define $(A_u(Q), \Delta_u)$ for any invertible Q . For such Q , one has the following free product decomposition [78],

$$A_u(Q) \cong A_u(P_1) * A_u(P_2) * \cdots * A_u(P_k)$$

for appropriate positive matrices P_1, P_2, \dots, P_k with compatible coproducts as in [71].

Definition 2.2. (cf. [83]) A **compact matrix quantum group** is a triple (A, Δ, π) , where $\pi : A_u(Q) \rightarrow A$ and $\Delta : A \rightarrow A \otimes A$ are C^* -morphisms such that

- (1) π is surjective, and
- (2) $\Delta\pi = (\pi \otimes \pi)\Delta_u$.

Note that using [71], conditions (1) and (2) above is equivalent to (1) plus the following condition:

$$(2)' \quad \Delta_u(\ker(\pi)) \subset \ker(\pi \otimes \pi).$$

Hence a compact matrix quantum group can be defined even more simply as a pair (A, π) satisfying (1) and (2)'.

The C^* -algebra A in the definition of a compact matrix quantum group (A, Δ, π) is called a **finitely generated Woronowicz C^* -algebra**. The morphism Δ is called the **coproduct** of A .

It can be shown that there is a Hopf $*$ -algebra structure $(\mathcal{A}_u(Q), \Delta, \varepsilon, S)$ on the dense $*$ -subalgebra $\mathcal{A}_u(Q)$ of $A_u(Q)$ such that

$$S(u_{ij}) = u_{ji}^*, \quad \varepsilon(u_{ij}) = \delta_{ij}, \quad i, j = 1, 2, \dots, n.$$

This $*$ -Hopf algebra structure induces Hopf $*$ -algebra structure on the dense $*$ -subalgebra \mathcal{A} of A in the above definition of compact matrix quantum group, and on the dense $*$ -subalgebra \mathcal{A} of A in the definition the compact quantum group below.

As in 2.3 of [71], one can define morphism between compact matrix quantum groups as opposite of morphisms of finitely generated Woronowicz C^* -algebras. Under these morphisms, compact matrix quantum groups form a category, though this category is not closed under inverse limits, which leads to the following notion of compact quantum groups using 3.1 of [71],

Definition 2.3. (cf. [1, 85]) A **compact quantum group** (A, Δ) is an inverse limit of compact matrix quantum groups $(A_\lambda, \Delta_\lambda, \pi_{\lambda\lambda'})$.

We will see in the next few paragraphs that the above notion of compact quantum groups is equivalent to the elegant and abstract one in [85].

As an inductive limit (instead of inverse limit) of finitely generated Woronowicz C^* -algebras A_λ , the C^* -algebra A in the definition above is called a **Woronowicz C^* -algebra**. Kernels of morphisms between Woronowicz C^* -algebras are called **Woronowicz C^* -ideals**, which can also be intrinsically defined as in 2.3 of [71].

Remark: Intuitively, we think of $A = C(G)$, where G is a compact quantum group, even there might be no points in “ G ” other than the identity. We also use the notations $A_G = C(G)$ and $C(G_A) = A$. As in the literature, by abuse of terminology, A is also called a compact (matrix) quantum group besides being called a (finitely generated) Woronowicz C^* -algebra.

As usual, one defines the Haar state/measure. One then establishes the existence and uniqueness of the Haar state/measure on any compact matrix quantum group just as in [83]. Using this and 3.3 of [71], one establishes the existence and uniqueness of the Haar state/measure on any compact quantum group G . In the following, the Haar state on $A = C(G)$ is denoted by h_G or simply h if no confusion arises.

Using the Haar state/measure h , the Peter-Weyl theory for all compact quantum groups can be developed as in [85] or [53]. As a result, we see that the above definition of compact quantum groups is equivalent to the one in [85].

We use \mathcal{A}_G to denote the dense $*$ -subalgebra of A_G consisting of coefficients of finite dimensional representations of G . As a consequence of the Peter-Weyl theory for compact quantum groups, \mathcal{A}_G is a Hopf $*$ -algebra.

We need to recall the notion of normal quantum subgroups [71, 80] to define simple compact quantum groups.

Let (N, π) be a **quantum subgroup** of a compact quantum group G , which, as defined in [70, 71], means that $\pi : C(G) \rightarrow C(N)$ is a surjection of C^* -algebras such that $(\pi \otimes \pi)\Delta_G = \Delta_N\pi$, where Δ_G, Δ_N are coproducts of $C(G)$ and $C(N)$, respectively.

Define

$$C(G/N) := \{a \in C(G) \mid (id \otimes \pi)\Delta(a) = a \otimes 1_N\},$$

$$C(N \setminus G) := \{a \in C(G) \mid (\pi \otimes id)\Delta(a) = 1_N \otimes a\},$$

Δ being coproduct on $C(G)$, 1_N the unit of $C(N)$.

Definition 2.4. (cf. [71, 79]) We say N is **normal** in G if it satisfies one of the equivalent conditions in the proposition below.

Proposition 2.5. (cf. [79]) *Let N be a quantum subgroup of a compact quantum group G . The following conditions are equivalent:*

- (1) $C(N \setminus G)$ is a Woronowicz C^* -subalgebra of $C(G)$.
- (2) $C(G/N)$ is a Woronowicz C^* -subalgebra of $C(G)$.
- (3) $C(G/N) = C(N \setminus G)$.
- (4) For every irreducible representation u^λ of G , either $h_N \pi(u^\lambda) = I_{d_\lambda}$ or $h_N \pi(u^\lambda) = 0$, where h_N is the Haar measure on N , $d_\lambda = \dim(u^\lambda)$ and I_{d_λ} the $d_\lambda \times d_\lambda$ identity matrix.

Among the above four equivalent formulations of the notion of normal quantum subgroups, condition (4) is the most convenient for our purposes. If G is a compact group, then the above definition coincides with the usual notion of closed normal subgroups.

To avoid complications with classification of finite quantum groups, we want to restrict the notion of simple quantum groups to quantum groups that are connected.

Definition 2.6. (cf. [79]) A compact quantum group G is called **connected** if for each non-trivial irreducible representation $u^\alpha \in \hat{G}$, the C^* -algebra $C^*(u_{ij}^\alpha)$ generated by the coefficients of u^α is of infinite dimension.

A compact matrix quantum group G is called **simple** (resp. **absolutely simple**) if it is connected and has no non-trivial connected normal quantum subgroups (resp. non-trivial normal quantum subgroups) and no non-trivial representations of dimension one.

Note that compact Lie groups are (commutative) examples of compact quantum groups. It is easy to show that a compact Lie group is simple (resp. connected) in the usual sense if and only if it is simple (resp. connected) in the sense above.

Problem 2.7. *In the definition of simple compact quantum groups, if one replaces the condition “it has no non-trivial representations of dimension one” with the apparently less stringent condition “ $C(G) \neq C^*(\Gamma)$ where Γ is a discrete group”, do we get an equivalent definition?*

Note according to [82], the condition $C(G) \neq C^*(\Gamma)$ above means that G is a nonabelian compact quantum group. For a compact Lie group G , this condition simply means G is a nonabelian Lie group, and the answer to the above question is affirmative by Weyl's dimension formula.

Just as the notion of simple compact Lie groups excludes the torus groups, the definition of simple quantum groups above (including the alternative one formulated in Problem 2.7) excludes the compact quantum groups coming from group C^* -algebras $C^*(\Gamma)$ of discrete groups Γ (i.e., abelian quantum groups). This is important because the classification of discrete groups is out of reach.

3. EXAMPLES OF SIMPLE COMPACT QUANTUM GROUPS THAT ARE ALMOST CLASSICAL AND HAVE PROPERTY F

The simple compact quantum groups that have been constructed so far share many properties common to compact Lie groups. We recall two of these properties [79].

Just as for compact groups, the **representation ring** (also called the **fusion ring**) $R(G)$ of a compact quantum group G is a partially ordered algebra over the integers \mathbb{Z} generated by the irreducible characters of G . The set of characters of G is a semi-ring that defines the order of $R(G)$.

Definition 3.1. (cf. [79]) A compact quantum group G is said to have **property F** if each Woronowicz C^* -subalgebra of $C(G)$ is of the form $C(G/N)$ for some normal quantum subgroup N of G .

A compact quantum group is called **almost classical** if its representation ring $R(G)$ is order isomorphic to the representation ring of a compact group.

In plain language, a compact quantum G is said to have property F if its quantum function algebra $C(G)$ behaves exactly as the function algebras of compact groups with respect to normal subgroups. Note that compact quantum groups $C^*(\Gamma)$ from dual Γ of a discrete group Γ do not have this property unless the discrete group Γ is abelian, in which case the group C^* -algebra $C^*(\Gamma)$ is a genuine function algebra over the Pontryagin dual $\hat{\Gamma}$ of the discrete abelian group Γ .

The notion dual to property F is given by following definition, which captures the property that compact quantum group $C^*(\Gamma)$ has with respect to normal quantum subgroups:

Definition 3.2. (cf. [79]) A compact quantum G is said to have **property FD** if each of quantum subgroup of G is normal.

Proofs of assertions in [79] concerning properties F and FD , along with other related properties of compact quantum groups, can be found in [80].

Quantum groups $B_u(Q)$ (cf. [70, 71, 69, 74, 78])

Keeping the notation for definition of $A_u(Q)$ in §2. Let $Q \in GL(n, \mathbb{C})$ be such that $Q\bar{Q} = \pm I_n$, $n \geq 2$. $B_u(Q)$ is defined to be the universal C^* -algebra with generators u_{ij} ($i, j = 1, 2, \dots, n$) that satisfy that following sets of relations:

$$u^*u = I_n = uu^*, \quad u^tQuQ^{-1} = I_n = QuQ^{-1}u^t.$$

It can be shown that there is a well-defined morphism

$$\Delta : B_u(Q) \rightarrow B_u(Q) \otimes B_u(Q)$$

such that

$$\Delta(u_{ij}) = \sum_{k=1}^n u_{ik} \otimes u_{kj}.$$

and that $(B_u(Q), \Delta)$ is a compact matrix quantum group. The quantum groups $B_u(Q)$ have the universal property that every compact matrix quantum group with self-conjugate fundamental representation is of the form $B_u(Q)/I$, where I is a Woronowicz C^* -ideal in $B_u(Q)$.

Note: $B_u(Q)$ is also denoted by $A_o(Q^*)$ by Banica *et al.*

When $Q = I_n$, $B_u(Q)$ is just $A_o(n)$, the universal orthogonal quantum group of Kac type introduced in 4.5 of [71]. Banica *et al.* also invented the notation $O^+(n)$ to signify $A_o(n) = C(O^+(n))$.

In addition, when $Q = \begin{bmatrix} 0 & -I_n \\ I_n & 0 \end{bmatrix}$, $B_u(Q)$ is a quantum symplectic group. This is one of the reasons that we use the notation $B_u(Q)$ instead of the notation $A_o(Q)$, as the latter only captures the special case $Q = I_n$. Following the notation $O^+(n)$ of Banica *et al.*, we denote the above universal symplectic quantum group by $Sp^+(n)$. The quantum symplectic group has also appeared in recent work of Bhowmick, D'Andrea, Das and Dąbrowski on quantum gauge symmetries (see [26]).

For any invertible $Q \in GL(n, \mathbb{C})$ that does not satisfy $Q\bar{Q} = \pm I_n$, $B_u(Q)$ can be defined by the same relations as above, but

$$\begin{aligned} B_u(Q) \cong & A_u(P_1) * A_u(P_2) * \cdots * A_u(P_k) * \\ & * B_u(Q_1) * B_u(Q_2) * \cdots * B_u(Q_l). \end{aligned}$$

for certain $P_i > 0$ and Q_j such that $Q_j \bar{Q}_j$'s are scalars (cf. [78]).

We note that both $A_u(Q)$ and $B_u(Q)$ can be alternatively described as quantum automorphism groups of appropriate spaces (cf. [76, 26]).

Quantum automorphism groups $A_{aut}(B, \tau)$ (cf. [75])

Let B be a finite dimensional C^* -algebra and ϕ a functional on B . In general the quantum automorphism group $A_{aut}(B, \phi)$ that preserves the system (B, ϕ) exists, which is the universal object in the category of compact quantum groups acting on the system (B, ϕ) . However, as shown in Banica [4], only when ϕ is the *canonical trace* τ does the quantum group $A_{aut}(B, \tau)$ have a representation theory that is relatively easy to describe, where the trace τ on B is called *canonical* if it coincides with the restriction to B of the unique tracial state on the algebra $L(B)$ of operators with B acting by the GNS representation associated with the trace τ . If one identifies B with $\bigoplus_{k=1}^m M_{n_k}(\mathbb{C})$, then

$$\tau\left(\sum_{k=1}^m b_k\right) = \sum_{k=1}^m \frac{n_k^2}{n} Tr(b_k),$$

where $b_k \in M_{n_k}$, n is the dimension of B , and Tr is ordinary trace on M_{n_k} , i.e. $Tr(b_k)$ is the sum of the diagonal entries of the matrix b_k .

In either case, explicit description of $A_{aut}(B, \phi)$ or $A_{aut}(B, \tau)$ in terms of generators and relation is complicated for a general finite dimensional C^* -algebra B . However, when $B = C(X_n)$ is the commutative C^* -algebra of functions on the space where X_n is the space of n points, the **quantum permutation group** $A_{aut}(X_n) := A_{aut}(C(X_n))$ has a surprisingly simple description in terms of generators and relations: The C^* -algebra $A_{aut}(X_n)$ is generated by self-adjoint projections a_{ij} such that each row and column of the matrix $(a_{ij})_{i,j=1}^n$ adds up to 1, i.e.,

$$\begin{aligned} a_{ij}^2 &= a_{ij} = a_{ij}^*, \quad i, j = 1, \dots, n, \\ \sum_{j=1}^n a_{ij} &= 1, \quad i = 1, \dots, n, \\ \sum_{i=1}^n a_{ij} &= 1, \quad j = 1, \dots, n. \end{aligned}$$

Banica *et al.* invented the very convenient geometric notation S_n^+ so that $A_{aut}(X_n) = C(S_n^+)$. In [34], Bichon generalizes the quantum permutation groups to purely algebraic context and establishes the universal property of the quantum permutation groups in complete generality.

Theorem 3.3. (cf. 4.1 and 4.7 in [79])

(a) For $Q \in GL(n, \mathbb{C})$ such that $Q\bar{Q} = \pm I_n$ and $n \geq 2$, $B_u(Q)$ is almost classical simple compact quantum group with property F .

(b) Let B be a finite dimensional C^* -algebra endowed with its canonical trace τ and $\dim(B) \geq 4$, then quantum group $A_{aut}(B, \tau)$ is an almost classical absolutely simple compact quantum groups with property F .

As special cases of $B_u(Q)$ and $A_{aut}(B, \tau)$, we have

Corollary 3.4. (a) The universal orthogonal quantum groups $O^+(n)$ (for $n \geq 2$) and the universal symplectic quantum groups $Sp^+(n)$ (for $n \geq 1$) are simple.

(b) The quantum permutation groups S_n^+ are simple for $n \geq 4$.

The main ideas used in the proof of Theorem 3.3 include

- (1) Banica's fundamental work of on the structure of fusion rings of these quantum groups (cf. Théorème 1 and Theorem 4.1 in [2, 4] respectively);
- (2) Correspondence between Hopf $*$ -ideals and Woronowicz C^* -ideals (cf. 4.2 - 4.3 in [79]);
- (3) Reconstruction of a normal quantum group from the identity in the quotient quantum group (cf. 4.4 in [79]).

In addition to the fusion rings of compact quantum groups such as those considered in [2, 4], the matters in (2) and (3) above are of interest in their own right and worth further investigation. The correspondence between Hopf $*$ -ideals and Woronowicz C^* -ideals and related matters in (2) relate algebraic and analytical aspects of compact quantum groups. In purely Hopf algebras context, reconstruction of a normal quantum group from the quotient quantum group in (3) has been an issue since 1970's and is related to several other important and old open questions [80].

Quantum groups K_q , K_q^u and K_J (cf. [62, 61, 52, 58, 73] and [59, 60])

A unified study of the compact quantum groups K_q , K_q^u is due to Soibelman & Vaksman, Levendorskii [62, 61, 52]. The $*$ -Hopf algebras \mathcal{A}_{K_q} are algebras of “representative functions” of Drinfeld-Jimbo quantum groups $U_q(\mathfrak{g})$ and define in a sense their “compact real form”. The quantum group K_q is a deformation of the Poisson Lie group $K(1, 0)$ (cf. [52]). The $*$ -Hopf algebras $\mathcal{A}_{K_q^u}$ are twisting of the $*$ -Hopf algebras \mathcal{A}_{K_q} by an element $u \in \wedge^2 \mathfrak{h}_{\mathbb{R}}$. The quantum group K_q^u is a deformation of the Poisson Lie group $K(1, u)$ (cf.

[52]). As shown in [73], K_q^u is an example of Rieffel's deformation from action of finite dimensional vector space as conjectured by Rieffel (cf. [57, 58] for the background).

Rieffel's quantum group deformation K_J [58] depends on $J = S \oplus (-S)$, where S is a skew symmetric operator on the Lie algebra (viewed as \mathbb{R}^n) of a torus subgroup of the compact Lie group K . For appropriate choice of S , K_J is a deformation of Poisson Lie group $K(0, u)$ [52, 58]. An action of $\mathbb{R}^d := \mathbb{R}^n \times \mathbb{R}^n$ on $A = C(K)$ can be constructed and Rieffel's theory of deformation for action of \mathbb{R}^d [57] can be applied to obtain A_J [58], also denoted $C(K_J)$.

Precise description of K_q , K_q^u and K_J require more space than appropriate in this paper. For our purposes, these quantum groups can be roughly described as follows:

- (1) The associated dense Hopf $*$ -algebras \mathcal{A}_{K_q} , $\mathcal{A}_{K_q^u}$ and \mathcal{A}_{K_J} are the same vector space as the un-deformed/un-twisted ones \mathcal{A}_K , \mathcal{A}_{K_q} and \mathcal{A}_K respectively, but the algebras \mathcal{A}_{K_q} , $\mathcal{A}_{K_q^u}$ and \mathcal{A}_{K_J} have deformed products;
- (2) The Hopf $*$ -algebras \mathcal{A}_{K_q} , $\mathcal{A}_{K_q^u}$ and \mathcal{A}_{K_J} have the same coproduct as the un-deformed/un-twisted ones \mathcal{A}_K , \mathcal{A}_{K_q} and \mathcal{A}_K respectively;
- (3) Representation theories of the deformed/twisted quantum groups K_q , K_q^u and K_J are the same as the un-deformed/un-twisted ones K , K_q and K respectively.

Theorem 3.5. cf. (5.1, 5.4 and 5.6 in [79]) *If K is a simple compact Lie group, then K_q , K_q^u , K_J are almost classical simple compact quantum groups with property F .*

The main ideas used in the proof of Theorem 3.5 include

- (1) Representation theory of these quantum groups;
- (2) Correspondence between Hopf $*$ -ideals and Woronowicz C^* -ideals, as in the proof of Theorem 3.3;
- (3) Reconstruction of a normal quantum group from the identity in the quotient quantum group, as in the proof of Theorem 3.3;
- (4) The normal subgroups of the undeformed Lie group remain to be normal subgroups of the deformed quantum groups and explicit identification of normal quantum subgroups of the deformed quantum groups.

Other deformations of compact Lie groups, though constructed not as systematic as the ones considered by Drinfeld-Jimbo, Soibelman *et al.* and Rieffel, are scattered in the literature. We believe the general ideas used in the proof of Theorem 3.5 can also be applied to such deformations.

4. QUOTIENT QUANTUM GROUPS FROM $A_u(Q)$ AND FREE PRODUCTS

In [78], the quantum groups $A_u(Q)$ are classified up to isomorphism for positive matrices $Q > 0$ and the quantum groups $B_u(Q)$ are classified up to isomorphism for matrices Q with $Q\bar{Q} = \pm I_n$. It is shown that the corresponding $A_u(Q)$ (resp. $B_u(Q)$) is not a free product, or a tensor product, or a crossed product. However, for general non-singular matrices Q , we have the decomposition theorem expressing $A_u(Q)$ and $B_u(Q)$ in terms of free product of the forgoing quantum groups (cf. Theorem 3.1 and 3.3 in [78]). In the light of these results, the following problem seems to be fundamental:

Problem 4.1. *Study further the fine structure of $A_u(Q)$ for positive matrices $Q \in GL(n, \mathbb{C})$ and $n \geq 2$; Determine their simple quotient quantum groups.*

A solution of this problem will also provide the first examples¹ of simple compact quantum groups that are not almost classical (see §3). Note that the $A_u(Q)$'s have the 1 dimensional diagonal torus \mathbb{T} as their (connected) normal quantum subgroup, as observed by Bichon (private communication, cf. 4.5 in [79]), so they are not simple. However, they are very close to being simple. For example, they have no non-trivial irreducible representations of dimension one [3, 78].

It is worth noting that in Problem 4.1, simple quotient quantum groups of $A_u(Q)$ should be easier to determine than simple quantum subgroups of $A_u(Q)$, since the latter is tantamount to finding all simple quantum groups due to the universal property of $A_u(Q)$, which include all simple compact quantum groups in [79] as reviewed in §3, simple compact Lie groups, as well as all the other unknown simple compact quantum groups.

By investigating the fine structure of concrete quantum groups such as $A_u(Q)$, one can expect to gain insights into the structure of general compact quantum groups and simple quantum groups. Sections §6 and §7 below contain more directions of research on this approach to quantum groups.

Suitable modifications of the method for the proofs of the main results in section 4 of the paper [79] should yield a solution to Problem 4.1. The extra work needed for this

¹After this paper was accepted for publication, Alexandru Chirvăsitu informed the author that in the preprint “Free unitary quantum groups are (almost) simple”, he showed that the quantum group generated by $u_{ij}u_{kl}^*$ and $u_{kl}^*u_{ij}$ is a simple compact quantum group with noncommutative representation ring and without property F , where u_{ij} are the generators of $A_u(Q)$ with $Q > 0$. In the same preprint, he also showed that the quotient quantum group of $A_u(Q)$ by its central subgroup \mathbb{T} in the proof of 4.5 in [79] has no normal quantum subgroups but is not finitely generated, and all normal subgroups are subgroups of \mathbb{T} , giving a complete classification of simple quotient groups of $A_u(Q)$. See also footnote before Problem 5.4 below.

problem that do not appear in [79] is that there are more Woronowicz subalgebras in $A_u(Q)$ to consider than therein. Some preliminary computations of these subalgebras give optimism to a positive solution of the problem. One of the main ingredients in this calculation is Banica's fundamental result on fusion rules of the irreducible representations of the quantum group $A_u(Q)$ [3].

The general $A_u(Q)$ (resp. $B_u(Q)$) for arbitrary $Q \in GL(n, \mathbb{C})$ is not simple if $C^*(\mathbb{Z})$ appears in its free product decompositions as described in [78]. This is because of the fact that $A_u(Q') = C^*(\mathbb{Z}) = C(\mathbb{T})$ (resp. $B_u(Q') = C^*(\mathbb{Z}/2\mathbb{Z})$) for $Q' \in GL(1, \mathbb{C})$ and the following result ([80]):

Proposition 4.2. *Let G_1, G_2 be compact quantum groups. Let $G = G_1 \hat{*} G_2$ be the free product compact quantum group [71] underlying $A_{G_1} * A_{G_2}$. Let π_1 be the natural embedding of G_1 into G defined by the surjection*

$$\pi_1 : A_{G_1} * A_{G_2} \rightarrow A_{G_1}, \quad \pi_1 = id_1 * \epsilon_2.$$

*If G_1 has at least one irreducible representation of dimension greater than one, then (G_1, π_1) is **not** a normal quantum subgroup of $G_1 \hat{*} G_2$. Otherwise, (G_1, π_1) is normal in $G_1 \hat{*} G_2$.*

The hat in the symbol $\hat{*}$ above signifies the ‘‘Fourier transform’’ of free product $*$ reminiscent of classical case in which $G_k = \hat{\Gamma}_k$, where Γ_k are discrete abelian groups.

Note that a compact Lie group being simple means roughly that it is not a direct product of proper connected subgroups. A similar result also holds for quantum groups: If G_A is a simple compact quantum group, then A_G is not a tensor product (i.e. G is not a direct product of its non-trivial quantum subgroups) [72, 80].

However, the proposition above says that the evident quantum subgroups (G_1, π_1) and (G_2, π_2) of $G_1 \hat{*} G_2$ are not normal in $G_1 \hat{*} G_2$ when G_1 and G_2 have no non-trivial representations of dimension one. Along with other results of the author, this may suggest that the following problem on the structure of simple quantum groups has a positive solution.

Problem 4.3. *Let G_1 and G_2 be simple compact quantum groups. Is $G_1 \hat{*} G_2$ also simple?*

The results in author's paper [71] should be useful for a solution of this problem. In particular, Theorem 1.1 there should play a role, as it did in [2, 3, 33, 78]. Note that the formula for the Haar measure on $G_1 \hat{*} G_2$ and the classification its irreducible representations are given in explicit formulas in Theorem 1.1 in [71]. According to the postulates in the definition of a normal quantum subgroup (Definition 2.4), these are important ingredients in determining whether $G_1 \hat{*} G_2$ has normal quantum subgroups.

The results and methods of the paper [79] (especially section 4 therein) should also be useful for this problem.

A positive solution to Problem 4.3 would have the following implication: $G_1 \hat{*} G_2$ would be a *simple compact quantum group* when both G_1 and G_2 are merely *simple compact Lie groups*.

The following easier variation of Problem 4.3 is also of interest and is related to the problem of determining whether several families of easy quantum groups are simple (cf. §6 below).

Problem 4.4. *Let G_1 be a simple compact quantum group and G_2 a finite quantum group. Is $G_1 \hat{*} G_2$ also simple?*

Note that in view of the problems in §6 below, it would be interesting to solve Problem 4.4 for G_2 a finite group. Also related problems can be formulated for Bichon's free wreath product of compact quantum groups [33] (cf. §6 below):

Problem 4.5. *Let G_1 be a simple compact quantum group and $G_2 = S_n^+$ the quantum permutation group. Is the free wreath product $G_1 \hat{*} S_n^+$ also simple?*

5. PROBLEMS RELATED TO ALMOST CLASSICAL COMPACT QUANTUM GROUPS AND PROPERTY F

As reviewed in §3, the simple compact quantum groups known so far are almost classical and have property F . Such quantum groups seems to be most accessible at the moment. The following problem evidently is less difficult than the general problem of classifying simple compact quantum groups and should be attempted first:

Problem 5.1. *Classify simple compact quantum groups that are almost classical and have property F .*

The following closely related problems should be considered also:

Problem 5.2. (a) *Classify almost classical simple compact quantum groups.*

(b) *Classify simple compact quantum groups with property F .*

Problem 5.3. *Does simple compact quantum groups with property FD exist? If so, construct and classify them.*

In Problem 5.1 and Problem 5.2, it would be interesting enough to restrict consideration to almost classical simple compact quantum groups that have the same representation rings as simple compact Lie groups.

In another direction, for the apparently more difficult problem of classifying simple compact quantum groups that are not almost classical or without property F , we do not have a single example of them. Therefore the following is a basic problem²:

Problem 5.4. (a) *Construct an example of simple compact quantum group that is not almost classical.*

(b) *Construct an example of simple compact quantum group that does not have property F .*

A more concrete problem than Problem 5.4 is the following

Problem 5.5. *Construct simple compact quantum groups with noncommutative representation ring.*

For the problems in this section, results on general structure of compact quantum groups such as those in the previous sections should also be useful. In this direction, we have the following result (cf. [80]).

Theorem 5.6. *Let G be a compact quantum group with property F . Then its quantum subgroups and quotient groups G/N by normal quantum subgroups N also have property F .*

It would be of interest to develop other general results on the structure of compact quantum groups.

6. QUANTUM AUTOMORPHISM GROUPS OF FINITE GRAPHS AND EASY QUANTUM GROUPS

How and where do we find quantum groups satisfying the properties in the problems in §5 above? The most natural approach, in our opinion, is by considering quantum automorphism groups of appropriate commutative and noncommutative spaces and their quantum subgroups. Much of the recent work on compact quantum groups falls into this category. We would like to mention three classes of these quantum groups: quantum automorphism groups of finite graphs and other quantum subgroups of the quantum

²Alexandru Chirvăsitu informed the author that he has since solved Problem 5.4 and Problem 5.5 - see footnote after Problem 4.1.

permutation groups, easy quantum groups and quantum isometry groups. The last of these three is discussed in the next section. We briefly look at the first two in this section.

In part to understand quantum subgroups of the quantum permutation groups [75], Bichon [32] constructed the *quantum automorphism groups of finite graphs*, which are quantum subgroups of the former preserving the edges of the graphs. To further understanding this new class of quantum groups, Bichon [33] also defined free wreath product of compact quantum groups using the quantum permutation groups and proved the beautiful formula stating that the free wreath product of the quantum automorphism group of a graph by the quantum permutation group $A_{\text{aut}}(X_n)$ is the quantum automorphism group of n disjoint copies of the graph. Banica also independently studied quantum subgroups of the quantum permutation groups [5, 6]. These works lead to a great deal of further studies of quantum automorphism of finite graphs and quantum subgroups of the quantum permutation groups, cf. [5] - [12] and references therein.

It is instructive to see an immediate application of the quantum automorphism groups of finite graphs to related problems in the last section. Clearly, a quantum quotient group G/N of an almost classical quantum group G is also almost classical. However a *crucial* observation is that a quantum subgroup of an almost classical quantum group needs not be almost classical. For example, the quantum permutation groups $A_{\text{aut}}(X_n)$ are almost classical (cf. [4, 75, 79] and §3), but according to of Bichon [33], their quantum subgroups $A_2(\mathbb{Z}/m\mathbb{Z})$, as the quantum automorphism group of certain graphs, are not almost classical if $m \geq 3$ (see Corollary 2.7 and the paragraph following Corollary 4.3 of [33]). Though the quantum groups $A_2(\mathbb{Z}/m\mathbb{Z})$ are not simple (see proposition 2.6 of [33]), and noting that the quantum automorphism group of the trivial graph (i.e. the quantum permutation group) is simple, it is natural to expect that it is possible to obtain simple compact quantum groups that are not almost classical by considering quantum automorphism groups of other appropriate finite graphs, including other free wreath products, thus solving Problem 5.4. Note that since the free wreath is constructed from the free product, such problems are related to see Problem 4.1 and the discussions following it.

In another important and related new direction that has origins in works on free wreath product and quantum subgroups of the quantum permutation group discussed above, Banica and Speicher [21] initiated the study of easy quantum groups and found several interesting families of new compact quantum groups. A compact matrix quantum group G with fundamental representation u is called **easy** if

- (1) G lies between S_n and $O^+(n)$ (cf. §3 on $B_u(Q)$ with $Q = I_n$); and

(2) For any $k, l \geq 0$, $\text{Hom}(u^{\otimes k}, u^{\otimes l})$ is linearly generated by operators T_p canonically associated with partitions p of $k + l$.

See [21] or [17] or [81] for a description of T_p .

In addition to the six families of free easy quantum groups (also called orthogonal quantum groups) in [21, 17], Weber also found another new family $B'_n{}^+$ of free easy quantum groups in [81], where we follow his different notation from [21]. In [56], Raum computed the fusion rings of several easy quantum groups in [21] using free products. In works of tour de force, Banica and Vergnioux computed the fusion rings of the quantum reflection group H_n^s in [22] and the half-librated orthogonal quantum group $O^*(n)$ in [23]. It would be interesting to see if any of these quantum groups provide solutions to some of the problems in this section. As the fusion rings of H_n^s and $O^*(n)$ are non-commutative, the following problems seem to be most appealing:

Problem 6.1. (a) *Is the quantum reflection group H_n^s simple?*

(b) *Is the half-librated orthogonal quantum group $O^*(n)$ simple?*

A positive answer to either (a) or (b) would provide the first simple compact quantum group that is not almost classical and has noncommutative representation ring.

In the light of Theorem 4.1 in Raum [56] and 3.1 and 3.2 in Weber [81] whose notation we follow, the relevant quantum groups considered in that theorem are either not simple or rely on solution of problems in §4. For instance, using their results above along with our Theorem 3.3, we see that

(1) B_n^+ is simple for $n \geq 3$;

(2) $B'_n{}^+$ ($n \geq 3$) and $S'_n{}^+$ ($n \geq 4$) are simple modulo two components (i.e. disconnected);

(3) $B_n^{\#+}$ is not simple because of Proposition 4.2, but it seems not far from being simple (a concept to be precised) because of 4.1.(3) in [56] and 3.2.(a) in [81].

It is not clear if the quantum groups H_n^+ , H_n^* , $H_n^{(s)}$ and $H_n^{[s]}$ are simple. Note that as a free wreath product, the quantum group H_n^+ is related to the general question Problem 4.5.

7. QUANTUM ISOMETRY GROUPS OF COMMUTATIVE AND NONCOMMUTATIVE RIEMANNIAN SPACES

After Banica's initial investigation of quantum isometry groups of finite spaces [5, 6], a recent conceptual breakthrough in compact quantum groups is Debashish Goswami's [44] theory of quantum isometry groups of spectral triples à la Connes [36], where the universal quantum groups $A_u(Q)$ plays an essential rôle in the proof of existence. Using

this notion he computed with Bhowmick [27, 24] several examples of quantum isometry groups and found that they are either isomorphic to classical isometry groups or are among the examples studied earlier by Rieffel and the author [58, 73]. Using the same circle of ideas they subsequently developed [28] an improved notion of quantum isometry group without relying on existence of a good Laplacian as required in [44].

The quantum isometry groups computed so far are either classical groups, or known quantum groups, or combinations of both based on free product or tensor product. Because of this, Goswami made a rigidity conjecture earlier stating to the effect that connected spaces do not admit non-trivial quantum symmetries. On the other hand, Huichi Huang [46] has shown that connected non-smooth metric spaces admit faithful action even by quantum permutation groups, disproving this earlier conjecture. Most striking of all is that Goswami, along with his collaborators, have recently shown in a series of papers [45, 39] a modified rigidity result stating that a connected and oriented Riemannian manifold does not have quantum symmetries other than the classical ones.

As a fundamental new concept, one would naturally wonder if fundamentally new compact quantum groups can be constructed using quantum isometry groups. In the light of results of Huang and Goswami *et al.* above, one should look at quantum isometry groups of non-smooth metric spaces or disconnected Riemannian manifolds that are non-classical (i.e. not compact groups). It would be interesting to find if any such quantum groups are simple:

Problem 7.1. *Construct examples of simple quantum isometry groups of non-smooth metric spaces or disconnected Riemannian manifolds.*

It is conceivable that quantum isometry groups of disconnected Riemannian manifolds will be related to quantum subgroups of the quantum permutation group as considered in §6, since quantum permutation group can permute the connected components in quantum manner just as it does on the finite space.

Another direction of research in the theory of quantum isometry group is the following. In the newly developed notion of quantum isometry group in [28], the quantum groups in the categories \mathbf{Q}'_R and \mathbf{Q}' that are used to define the quantum isometry group there does not carry a C^* -algebraic action on the spectral triple, and as a result, the universal object (i.e, the quantum isometry group) does not carry a C^* -algebraic action in general. See [30] for an example of this situation. (We refer the reader to the above cited papers for detailed description of \mathbf{Q}'_R and \mathbf{Q}' due to space limitation.)

To address this problem, we believe the categories \mathbf{Q}'_R and \mathbf{Q}' are too large, and propose the following alternative for the notion of quantum isometry groups that will always carry C^* -algebraic action.

First, by a **compact quantum transformation group** (A, α, u) of a compact type spectral triple $(\mathcal{B}, \mathcal{H}, D)$ we mean a compact quantum transformation group (cf. [75]) (A, α) of B that satisfies

(QT1) There is a unitary representation $u \in L(\mathcal{H} \otimes A_G)$ of G_A on \mathcal{H} such that (ρ, u) is α -covariant: $(\rho \otimes 1)\alpha(b) = u(\rho(b) \otimes 1)u^*$, $b \in B$;

(QT2) D is an intertwiner of u with itself: $(D \otimes 1)u = u(D \otimes 1)$.

A **morphism** from another quantum transformation group $(\tilde{A}, \tilde{\alpha}, \tilde{u})$ to (A, α, u) is defined to be a morphism π from $(\tilde{A}, \tilde{\alpha})$ to (A, α) (cf. [75]) that satisfies $\tilde{u} = (id_{\mathcal{H}} \otimes \pi)u$.

The above defines **category \mathcal{C} of compact quantum transformation groups** of $(\mathcal{B}, \mathcal{H}, D)$ with objects $\{(A, \alpha, u)\}$ and morphisms $\{\pi\}$. Similar to [75], the **quantum isometry group** of $(\mathcal{B}, \mathcal{H}, D)$ is defined to be a universal object of category \mathcal{C} if it exists.

As in [75], universal object does not always exist in \mathcal{C} in general. Therefore, as in [75] and [28], consider **measured spectral triple** $(\mathcal{B}, \mathcal{H}, D, \phi)$ with a (usually positive) functional ϕ on \mathcal{B} and consider the category \mathcal{C}_ϕ of compact quantum transformation groups that satisfies (QT1)-(QT2) and

(QT3) $(\phi \otimes id)(\alpha(b)) = \phi(b)1_A$ for $b \in \mathcal{B}$

The **quantum isometry group** of $(\mathcal{B}, \mathcal{H}, D, \phi)$ is defined to be the universal object of category \mathcal{C}_ϕ if it exists. The following can be proved and justifies in part the above notion quantum isometry group.

Proposition 7.2. (1) *Let $(\mathcal{B}, \mathcal{H}, D)$ be the spectral triple associated with a compact Riemannian manifold M . Then the universal object in the category of compact transformation groups of $(\mathcal{B}, \mathcal{H}, D)$ in the sense above is the **isometry group** of M .*

(2) *For an arbitrary spectral triple $(\mathcal{B}, \mathcal{H}, D)$, the universal object in the category of compact transformation groups of $(\mathcal{B}, \mathcal{H}, D)$ in the sense above is the **compact group of isometries** of $(\mathcal{B}, \mathcal{H}, D)$ in the sense of Connes (see p6200 of [37]).*

Since composition of two continuous maps is continuous, the quantum isometry group defined above is evidently contained in the quantum isometry groups defined by Goswami [44] and Bhowmick and Goswami [28]. Because of this, and using ideas in [75] and [28], the following seems to be quite plausible:

Problem 7.3. *For a measured spectral triple $(\mathcal{B}, \mathcal{H}, D, \phi)$, show that universal object exists in the category \mathcal{C}_ϕ .*

Many other problems naturally arise:

(i) Calculate the quantum isometry groups in the sense above for the classical spaces such as the flat spheres, tori, and other Riemannian spaces;

(ii) Calculate the quantum isometry groups in the sense above for the noncommutative tori;

(iii) Study the properties of quantum isometry groups in general and apply them to study other properties of noncommutative spaces;

(iv) Construct simple compact quantum groups through the study of quantum isometry groups.

As in the definitions of quantum isometry groups by Goswami and Bhowmick, and because a quantum isometry group in our sense is contained in theirs, examples of simple quantum isometry groups in our sense might need to be constructed out of non-smooth metric spaces or disconnected Riemannian manifolds, as in Problem 7.1.

Acknowledgments. The author would like to thank Hanfeng Li and Yi-Jun Yao for invitations to Chongqing University and Fudan University respectively during the summer of 2012, where the author had the opportunity to work on this paper. The author would also like to thank Hanfeng Li and Dechao Zheng at Chongqing University, and Yi-Jun Yao and Guoliang Yu at Fudan University for their hospitality and for creating a congenial atmosphere. He also wish to record his thanks to Huichi Huang, Pan Ma and Qinggang Ren for making his visit enjoyable. The author is indebted to Professor Marek Bozejko for drawing his attention to easy quantum groups during the conference in September 2011 celebrating Professor Woronowicz's seventieth birthday.

REFERENCES

- [1] Baaĵ, S. and Skandalis, G.: Unitaires multiplicatifs et dualité pour les produits croisés de C^* -algèbres, *Ann. Sci. Ec. Norm. Sup.* **26** (1993), 425-488.
- [2] Banica, T.: Théorie des représentations du groupe quantique compact libre $O(n)$, *C. R. Acad. Sci. Paris t.* **322**, Serie I (1996), 241-244.
- [3] Banica, T.: Le groupe quantique compact libre $U(n)$, *Commun. Math. Phys.* **190** (1997), 143-172.
- [4] Banica, T.: Symmetries of a generic coaction. *Math. Ann.* **314** (1999), 763-780.
- [5] Banica, T.: Quantum automorphism groups of homogeneous graphs. *J. Funct. Anal.* **224** (2005), no. 2, 243–280.
- [6] Banica, T.: Quantum automorphism groups of small metric spaces. *Pacific J. Math.* **219** (2005), no. 1, 27–51.
- [7] Banica, T., Bichon, J.: Free product formulae for quantum permutation groups. *J. Inst. Math. Jussieu* **6** (2007), no. 3, 381–414.

- [8] Banica, T., Bichon, J.: Quantum automorphism groups of vertex-transitive graphs of order ≤ 11 . *J. Algebraic Combin.* **26** (2007), no. 1, 83–105.
- [9] Banica, T., Bichon, J.: Quantum groups acting on 4 points, *J. Reine Angew. Math.*, **626** (2009), 74–114.
- [10] Banica, T., Bichon, J., Chenevier, G.: Graphs having no quantum symmetry. *Ann. Inst. Fourier (Grenoble)* **57** (2007), no. 3, 955–971.
- [11] Banica, T., Bichon, J., Collins, B.: The hyperoctahedral quantum group. *J. Ramanujan Math. Soc.* **22** (2007), no. 4, 345–384.
- [12] Banica, T., Bichon, J., Collins, B.: Quantum permutation groups: a survey, *Banach Center Publications* 78, 13–34, 2007.
- [13] Banica, T., Bichon, J., Natale, S.: Finite quantum groups and quantum permutation groups. *Adv. Math.* **229** (2012), 3320–3338
- [14] Banica, T., Collins, B.: Integration over quantum permutation groups. *J. Funct. Anal.* **242** (2007), no. 2, 641–657.
- [15] Banica, T., Collins, B.: Integration over compact quantum groups. *Publ. Res. Inst. Math. Sci.* **43** (2007), no. 2, 277–302.
- [16] Banica, T., Collins, B.: Integration over the Pauli quantum group. *J. Geom. Phys.* **58** (2008), 942–961.
- [17] Banica, T., Curran, S., Speicher, R.: Classification results for easy quantum groups. *Pacific J. Math.* **247** (2010), no. 1, 1–26.
- [18] Banica, T., Curran, S., Speicher, R.: Stochastic aspects of easy quantum groups. *Probab. Theory Related Fields* **149** (2011), 435–462.
- [19] Banica, T., Curran, S., Speicher, R.: De Finetti theorems for easy quantum groups. *Ann. Probab.* **40** (2012), 401–435.
- [20] Banica, T., Moroianu, S.: On the structure of quantum permutation groups. *Proc. Amer. Math. Soc.* **135** (2007), no. 1, 21–29.
- [21] Banica, T., Speicher, R.: Liberation of orthogonal Lie groups. *Adv. Math.* **222** (2009), 1461–1501.
- [22] Banica, T. and Vergnioux, R.: Fusion rules for quantum reflection groups. *J. Noncommut. Geom.* **3** (2009), 327–359.
- [23] Banica, T. and Vergnioux, R.: Invariants of the half-liberated orthogonal group. *Ann. Inst. Fourier* **60** (2010), 2137–2164.
- [24] Bhowmick, J.: Quantum Isometry Group of the n tori. *Proc. Amer. Math. Soc.* **137** (2009), no. 9, 3155–3161.
- [25] Bhowmick, J., D’Andrea, F., Dąbrowski, L.: Quantum isometries of the finite noncommutative geometry of the standard model. *Comm. Math. Phys.* **307** (2011), no. 1, 101–131
- [26] Bhowmick, J., D’Andrea, F., Das, B., Dąbrowski, L.: Quantum gauge symmetries in Noncommutative Geometry. arXiv:1112.3622
- [27] Bhowmick, J. and Goswami, D.: Quantum Isometry Groups: Examples and Computations, *Commun. Math. Phys.* **285** (2009), 421–444,
- [28] Bhowmick, J. and Goswami, D.: Quantum group of orientation-preserving Riemannian isometries. *J. Funct. Anal.* **257** (2009), no. 8, 2530–2572.

- [29] Bhowmick, J. and Goswami, D.: Quantum isometry groups of the Podleś spheres. *J. Funct. Anal.* **258** (2010), no. 9, 2937-2960.
- [30] Bhowmick, J., Goswami, D.: Some counterexamples in the theory of quantum isometry groups. *Lett. Math. Phys.* **93** (2010), no. 3, 279-293.
- [31] Bhowmick, J., Goswami, D., Skalski, A.: Quantum Isometry Groups of 0- Dimensional Manifolds. *Trans. Amer. Math. Soc.* **363** (2011), no. 2, 901-921.
- [32] Bichon, J.: Quantum automorphism groups of finite graphs. *Proc. Amer. Math. Soc.* **131** (2003), no. 3, 665–673
- [33] Bichon, J.: Free wreath product by the quantum permutation group. *Algebr. Represent. Theory* **7** (2004), 343–362.
- [34] Bichon, J.: Algebraic quantum permutation groups. *Asian-Eur. J. Math.* **1** (2008), no. 1, 1–13.
- [35] Bichon, J., De Rijdt, A., Vaes, S.: Ergodic coactions with large multiplicity and monoidal equivalence of quantum groups. *Comm. Math. Phys.* **262** (2006), 703–728.
- [36] Connes, A.: *Noncommutative Geometry*, Academic Press, 1994.
- [37] Connes, A.: Noncommutative geometry and reality, *J. Math. Phys.* **36**:11 (1995), 6194-6231
- [38] Connes, A.: Gravity coupled with matter and the foundation of non commutative geometry, *Commun. Math. Phys.* **182** (1996), 155-176.
- [39] Das, B. and Goswami, D and Joardar, S.: Rigidity of action of compact quantum groups II. arXiv:1206.1718
- [40] Drinfeld, V.G.: Quantum groups, *in* Proc. of the ICM-1986, Berkeley, Vol I, Amer. Math. Soc., Providence, R.I., 1987, pp798–820.
- [41] Enock, M. et Schwartz, J.-M.: Une dualité dans les algèbres de von Neumann, *Bull. Soc. Math. France. Suppl. Mémoire* **44** (1975), 1-144.
- [42] Enock, M. et Schwartz, J.-M.: *Kac algebras and duality of locally compact groups*. (English summary) With a preface by Alain Connes. With a postface by Adrian Ocneanu. Springer-Verlag, Berlin, 1992. x+257 pp. ISBN: 3-540-54745-2
- [43] Fredenhagen, K. and Rehren, K.H. and Schroer, B.: Superselection sectors with braid group statistics and exchange algebras I, *Comm. Math. Phys.* **125** (1989), 201-226.
- [44] Goswami, D.: Quantum Group of Isometries in Classical and Noncommutative Geometry, *Comm. Math. Phys.* **285** (2009), no. 1, 141-160.
- [45] Goswami, D.: Rigidity of action of compact quantum groups I, III arXiv:1106.5107, arXiv:1207.6470
- [46] Huang, Huichi: Faithful compact quantum group actions on connected compact metrizable spaces, arXiv:1202.1175
- [47] Kac, G.: Ring groups and the duality principle, I, II *Proc. Moscow Math. Soc.* **12** (1963), 259-303; *ibid.* **13** (1965), 84-113.
- [48] Kac, G. and Vainerman, L. I.: Nonunimodular ring groups and Hopf Von Neumann algebras, *Math. USSR Sb.* **23** (1974), 185-214.
- [49] Köstler, C., Speicher, R.: A noncommutative de Finetti theorem: Invariance under quantum permutations is equivalent to freeness with amalgamation. *Comm. Math. Phys.* **291** (2009), no. 2, 473-490.
- [50] Kustermans, Johan and Vaes, Stefaan Locally compact quantum groups. *Ann. Sci. École Norm. Sup.* (4) **33** (2000), no. 6, 837-934.

- [51] Levendorskii, S.: Twisted algebra of functions on compact quantum group and their representations, *Algebra i analiz* **3**:2 (1991), 180-198. *St. Petersburg Math. J.* **3**:2 (1992), 405-423.
- [52] Levendorskii, S. and Soibelman, Y.: Algebra of functions on compact quantum groups, Schubert cells, and quantum tori, *Commun. Math. Phys.* **139** (1991), 141-170.
- [53] Maes, Annand Van Daele, Alfons: Notes on compact quantum groups. *Nieuw Arch. Wisk.* (4) **16** (1998), no. 1-2, 73-112.
- [54] Masuda, T.; Nakagami, Y.; Woronowicz, S. L. A C^* -algebraic framework for quantum groups. *Internat. J. Math.* **14** (2003), no. 9, 903-1001
- [55] Podleś, P. and Woronowicz S. L.: Quantum deformation of Lorentz group, *Commun. Math. Phys.* **130** (1990), 381-431.
- [56] Raum, S.; Isomorphisms and fusion rules of orthogonal free quantum groups and their free complexifications. *Proc. Amer. Math. Soc.* **140** (2012), no. 9, 3207-3218.
- [57] Rieffel, M.: Deformation quantization for actions of \mathbf{R}^n , *Memoirs A.M.S.* no. **506**, 1993.
- [58] Rieffel, M.: Compact quantum groups associated with toral subgroups, *Contemp. Math.* **145** (1993), 465-491.
- [59] Rosso, M.: Comparaison des groupes $SU(2)$ quantiques de Drinfeld et Woronowicz, *C. R. Acad. Sci. Paris* **304** (1987), 323-326.
- [60] Rosso, M.: Algèbres enveloppantes quantifiées, groupes quantiques compacts de matrices et calcul différentiel non-commutatif, *Duke Math. J.* **61** (1990), 11-40.
- [61] Soibelman, Y.: Algebra of functions on compact quantum group and its representations, *Algebra i analiz* **2**:1 (1990), 190-212. *Leningrad Math. J.* **2**:1 (1991), 161-178.
- [62] Soibelman, Y. and Vaksman, L.: The algebra of functions on quantum $SU(2)$, *Funct. Anal. ego Pril.* **22**:3 (1988), 1-14. *Funct. Anal. Appl.* **22**:3 (1988), 170-181.
- [63] Soltan, P.M.: Quantum families of maps and quantum semigroups on finite quantum spaces. *J. Geom. Phys.* **59** (2009), no. 3, 354-368.
- [64] Vaes, S. and Vennet, N.V.: Identification of the Poisson and Martin boundaries of orthogonal discrete quantum groups. *Journal of the Institute of Mathematics of Jussieu* **7** (2008), 391-412.
- [65] Vaes, S. and Vennet, N.V.: Poisson boundary of the discrete quantum group $\widehat{A_u(F)}$. *Compos. Math.* **146** (2010), no. 4, 1073-1095.
- [66] Vaes, S. and Vergnioux, R.: The boundary of universal discrete quantum groups, exactness and factoriality. *Duke Mathematical Journal* **140** (2007), 35-84.
- [67] Vergnioux, R., Voigt, C.: The K-theory of free quantum groups. arXiv:1112.3291
- [68] Voigt, C.: The Baum-Connes conjecture for free orthogonal quantum groups. *Adv. Math.* **227** (2011), 1873 - 1913.
- [69] Van Daele, A. and Wang, S.Z.: Universal quantum groups, *International J. of Math.* **7** no2 (1996), 255-264.
- [70] Wang, S. Z.: *General Constructions of Compact Quantum Groups*, Ph.D. Thesis, University of California at Berkeley, March, 1993.
- [71] Wang, S.Z.: Free products of compact quantum groups, *Commun. Math. Phys.* **167** (1995), 671-692.
- [72] Wang, S.Z.: Tensor products and crossed products of compact quantum groups, *Proc. London Math. Soc.* **71** (1995), 695-720.

- [73] Wang, S.Z.: Deformations of compact quantum groups via Rieffel quantization, *Commun. Math. Phys.* **178**:3 (1996), 747-764.
- [74] Wang, S.Z.: Problems in the theory of quantum groups, in *Quantum Groups and Quantum Spaces*, Banach Center Publication 40 (1997), Inst. of Math., Polish Acad. Sci., Editors: R. Budzynski, W. Pusz, and S. Zakrzewski. pp67-78
- [75] Wang, S.Z.: Quantum symmetry groups of finite spaces, *Commun. Math. Phys.* **195**:1 (1998), 195-211.
- [76] Wang, S.Z.: Ergodic actions of universal quantum groups on operator algebras, *Commun. Math. Phys.* **203** (1999), 481-498.
- [77] Wang, S.Z.: Classification of quantum groups $SU_q(n)$, *J. London Math. Soc.* **59**:2 (1999), 669-680
- [78] Wang, S.Z.: Structure and isomorphism classification of quantum groups $A_u(Q)$ and $B_u(Q)$, *Journal of Operator Theory* **48**(3) (2002), 573-583
- [79] Wang, S.Z.: Simple compact quantum groups I, *J. Funct. Anal.* **256** (2009), no. 10, 3313-3341.
- [80] Wang, S.Z.: Equivalent notions of normal quantum subgroups, compact quantum groups with properties F and FD , and other applications, preprint.
- [81] Weber, M.: On the classification of easy quantum groups - The nonhyperoctahedral and the half-liberated case. arXiv:1201.4723
- [82] Woronowicz, S. L.: Twisted $SU(2)$ group. An example of noncommutative differential calculus, *Publ. RIMS, Kyoto Univ.* **23** (1987), 117-181.
- [83] Woronowicz, S. L.: Compact matrix pseudogroups, *Commun. Math. Phys.* **111** (1987), 613-665.
- [84] Woronowicz, S. L.: Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups, *Invent. Math.* **93** (1988), 35-76.
- [85] Woronowicz, S. L.: Compact quantum groups, in *Quantum Symmetries*, Les Houches Summer School-1995, Session LXIV, A. Connes, K. Gawedzki, J. Zinn-Justin, ed., Elsevier Science, Amsterdam, 1998; pp845-884.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GA 30602, USA

FAX: 706-542-5907, TEL: 706-542-0884

E-mail address: `szwang@math.uga.edu`