

A TALE OF TWO SURFACES

ARNAUD BEAUVILLE

ABSTRACT. We point out a link between two surfaces which have appeared recently in the literature: the surface of cuboids and the Schoen surface. Both give rise to a surface with $q = 4$, whose canonical map is 2-to-1 onto a complete intersection of 4 quadrics in \mathbb{P}^6 with 48 nodes.

1. INTRODUCTION

The aim of this note is to point out a link between two surfaces which have appeared recently in the literature: the *surface of cuboids* ([ST], [vL]) and the surface (actually a family of surfaces) discovered by Schoen [S]. We will show that both surfaces give rise to a surface X with $q = 4$, whose canonical map is 2-to-1 onto a complete intersection of 4 quadrics $\Sigma \subset \mathbb{P}^6$ with 48 nodes. In the first case (§2) X is a quotient $(C \times C')/(\mathbb{Z}/2)^2$, where C and C' are genus 5 curves with a free action of $(\mathbb{Z}/2)^2$. In the second case (§3), X is a double étale cover of the Schoen surface.

When the canonical map of a surface X of general type has degree > 1 onto a surface, that surface either has $p_g = 0$ or is itself canonically embedded ([B1], Th. 3.1). Our surfaces X provide one more example of the latter case, which is rather exceptional (see [CPT] for a list of the examples known so far).

2. THE SURFACE OF CUBOIDS AND ITS DEFORMATIONS

In \mathbb{P}^4 , with coordinates $(x, y; u, v, w)$, we consider the curve C given by

$$(1) \quad u^2 = a(x, y) \quad , \quad v^2 = b(x, y) \quad , \quad w^2 = c(x, y)$$

where a, b, c are quadratic forms in x, y . We assume that the zeros $\{p'_a, p''_a; p'_b, p''_b; p'_c, p''_c\}$ of a, b, c form a set $Z \subset \mathbb{P}^1$ of 6 distinct points. Then C is a smooth curve of genus 5, canonically embedded. It is preserved by the group $\Gamma_+ \cong (\mathbb{Z}/2)^3$ which acts on \mathbb{P}^4 by changing the signs of u, v, w . Let $\Gamma \subset \Gamma_+$ be the subgroup (isomorphic to $(\mathbb{Z}/2)^2$) which changes an even number of signs. It acts freely on C , so the quotient curve $B := C/\Gamma$ has genus 2. The subring of Γ -invariant elements in $\oplus H^0(C, K_C^n)$ is generated by x, y and $z := uvw$, with the relation $z^2 = abc$; thus B is the double cover of \mathbb{P}^1 branched along Z .

Let JB_2 be the group of 2-torsion line bundles on B (isomorphic to $(\mathbb{Z}/2)^4$). The Γ -covering $\pi : C \rightarrow B$ corresponds to a subgroup $\cong (\mathbb{Z}/2)^2$ of JB_2 , namely the kernel of $\pi^* : JB \rightarrow JC$. Since the divisor $\pi^*(p'_a + p''_a)$ is cut out on C by the canonical divisor $u = 0$, we have $\pi^*(p'_a - p''_a) \sim 0$, and similarly for b and c ; thus $\text{Ker } \pi^* = \{0, p'_a - p''_a, p'_b - p''_b, p'_c - p''_c\}$. This is a *Lagrangian* subgroup of JB_2 for the Weil pairing [M2]; conversely, any Lagrangian subgroup of JB_2 is of that form. Thus the

curves C we are considering are exactly the $(\mathbb{Z}/2)^2$ -étale covers of a curve B of genus 2 associated to a Lagrangian subgroup of JB_2 . In particular they form a 3-dimensional family.

The group $\Gamma_+/\Gamma \cong \mathbb{Z}/2$ acts on $B = C/\Gamma$ through the hyperelliptic involution, so C/Γ_+ is isomorphic to \mathbb{P}^1 .

Proposition 1. *Let C, C' be two genus 5 curves of type (1), and let X be the quotient of $C \times C'$ by the diagonal action of $\Gamma \cong (\mathbb{Z}/2)^2$.*

1) X is a minimal surface of general type with $q = 4$, $p_g = 7$, $K^2 = 32$.

2) The involution i_X of X defined by the action of $\Gamma_+/\Gamma \cong \mathbb{Z}/2$ has 48 fixed points. The canonical map $\varphi_X : X \rightarrow \mathbb{P}^6$ factors through i_X , and induces an isomorphism of X/i_X onto a complete intersection of 4 quadrics in \mathbb{P}^6 with 48 nodes.

Proof : The computation of the numerical invariants of X is straightforward.

Let us denote by $(x', y'; u', v', w')$ the coordinates on C' , and by a', b', c' the corresponding quadratic forms. A basis of the space $H^0(X, K_X) = (H^0(C, K_C) \otimes H^0(C', K_{C'}))^{\Gamma}$ is given by the elements

$$X = x \otimes x' \quad Y = x \otimes y' \quad Z = y \otimes x' \quad T = y \otimes y' \quad U = u \otimes u' \quad V = v \otimes v' \quad W = w \otimes w' .$$

They satisfy the relations

$$XT - YZ = 0 \quad , \quad U^2 = A(X, Y, Z, T) \quad , \quad V^2 = B(X, Y, Z, T) \quad , \quad W^2 = C(X, Y, Z, T) ,$$

where A, B, C are quadratic forms satisfying $A(X, Y, Z, T) = a(x, y) \otimes a(x', y')$ and the analogous relations for B and C .

Let Σ be the surface defined by these 4 quadratic forms, and let $\varphi : X \rightarrow \Sigma$ be the induced map. We have $\varphi \circ i_X = \varphi$, so φ induces a map $\bar{\varphi}$ from $X/i_X = (C \times C')/\Gamma_+$ into Σ . We consider the commutative diagram

$$\begin{array}{ccc} (C \times C')/\Gamma_+ & \xrightarrow{\bar{\varphi}} & \Sigma \\ & \searrow p & \swarrow q \\ & Q \cong \mathbb{P}^1 \times \mathbb{P}^1 & \end{array}$$

where $p : (C \times C')/\Gamma_+ \rightarrow (C/\Gamma_+) \times (C'/\Gamma_+)$ is the quotient map by Γ_+ , and q the projection $(X, Y, Z, T; U, V, W) \mapsto (X, Y, Z, T)$. The group $(\mathbb{Z}/2)^3$ acts on Σ by changing the signs of (U, V, W) ; then $\bar{\varphi}$ is an equivariant map of $(\mathbb{Z}/2)^3$ -coverings, hence an isomorphism.

It remains to show that i_X has 48 fixed points. These fixed points are the images (mod. Γ) of the points of $C \times C'$ fixed by one of the elements of $\Gamma_+ \setminus \Gamma$. Such an element changes the sign of one of the coordinates $\ell = u, v$ or w , hence fixes the 64 points (m, m') of $C \times C'$ with $\ell(m) = \ell(m') = 0$. This gives $(3 \times 64)/4 = 48$ fixed points in X . ■

Example. Let us take for C and C' the curve

$$u^2 = xy \quad , \quad v^2 = x^2 - y^2 \quad , \quad w^2 = x^2 + y^2 .$$

The set of zeros of a, b, c is $\{0, \infty, \pm 1, \pm i\}$, so the genus 2 curve B is given by $z^2 = x(x^4 - 1)$.

We get for Σ the following equations :

$$XT = YZ = U^2 \quad , \quad V^2 = X^2 - Y^2 - Z^2 + T^2 \quad , \quad W^2 = X^2 + Y^2 + Z^2 + T^2 \quad ;$$

or, after the linear change of variables $X = x + t$, $T = t - x$, $Y = y + iz$, $Z = y - iz$, $U = u$, $V = 2v$, $W = 2w$:

$$t^2 = x^2 + y^2 + z^2 \quad , \quad u^2 = y^2 + z^2 \quad , \quad v^2 = x^2 + z^2 \quad w^2 = x^2 + y^2 \quad .$$

These are the equations of the *surface of cuboids*, studied in [ST], [vL]. It encodes the relations in a cuboid (= rectangular box) between the sides x, y, z , the diagonals of the faces u, v, w , and the big diagonal t . Thus the surface of cuboids belongs to a 6-dimensional family of intersection of 4 quadrics in \mathbb{P}^6 with 48 nodes.

Remark 1. The surfaces X fit into a tower of $(\mathbb{Z}/2)^2$ -étale coverings :

$$C \times C' \longrightarrow X \xrightarrow{r} B \times B' \quad .$$

The abelian covering r is the pull back of a $(\mathbb{Z}/2)^2$ -étale covering of $JB \times JB'$:

$$\begin{array}{ccc} X & \xhookrightarrow{\alpha} & A \\ \downarrow r & & \downarrow \\ B \times B' & \xhookrightarrow{\quad} & JB \times JB' \end{array} \quad .$$

The abelian variety A is the Albanese variety of X , and α is the Albanese map. Since the quotient X/i_X is regular, i_X acts as (-1) on the space $H^0(X, \Omega_X^1)$; therefore if we choose α so that it maps a fixed point of i_X to 0, i_X is induced by (-1_A) .

3. THE SCHOEN SURFACE

The Schoen surfaces S have been defined in [S], and studied in [CMR]. A Schoen surface S is contained in its Albanese variety A ; it has the following properties:

- a) $K_S^2 = 16$, $p_g = 5$, $q = 4$ (hence $\chi(\mathcal{O}_S) = 2$);
- b) The canonical map $\varphi_S : S \rightarrow \mathbb{P}^4$ factors through an involution i_S with 40 fixed points, and induces an isomorphism of S/i_S onto the complete intersection of a quadric and a quartic in \mathbb{P}^4 with 40 nodes [CMR].

Since S/i_S is a regular surface, i_S acts as (-1) on the space $H^0(S, \Omega_S^1)$. Therefore if we choose the Albanese embedding $S \hookrightarrow A$ so that it maps a fixed point of i_S to 0, i_S is induced by the involution (-1_A) .

Let ℓ be a line bundle of order 2 on A ; we denote by $\pi : B \rightarrow A$ the corresponding étale double cover, and put $X := \pi^{-1}(S)$. The restriction of ℓ to S , which we will still denote by ℓ , is nontrivial (because the restriction map $\text{Pic}^0(A) \rightarrow \text{Pic}^0(S)$ is an isomorphism), hence X is connected.

Proposition 2. *X is a minimal surface of general type with $q = 4$, $p_g = 7$, $K_X^2 = 32$.*

Proof : The formulas $K_X^2 = 32$ and $\chi(\mathcal{O}_X) = 4$ are immediate; we must prove $q(X) = 4$, that is, $H^1(S, \ell) = 0$.

By construction [S] the Schoen surfaces fit into a flat family over the unit disk Δ :

$$\begin{array}{ccc} S & \hookrightarrow & \mathcal{A} \\ & \searrow & \downarrow \\ & & \Delta \end{array}$$

where:

- \mathcal{A}/Δ is a smooth family of abelian varieties;
- at a point $z \neq 0$ of Δ , \mathcal{S}_z is a Schoen surface, and $\mathcal{S}_z \hookrightarrow \mathcal{A}_z$ is the Albanese embedding;
- $\mathcal{A}_0 = JC \times JC$ for a genus 2 curve C ; \mathcal{S}_0 is the union of JC embedded diagonally in $JC \times JC$, and of $C \times C \subset JC \times JC$ (we choose an Abel-Jacobi embedding $C \subset JC$). These two components intersect transversally along the diagonal $C \subset C \times C$.

The line bundle ℓ extends to a line bundle \mathcal{L} of order 2 on \mathcal{A} . Let ℓ_0 be the restriction of \mathcal{L} to \mathcal{S}_0 ; we want to compute $H^1(\mathcal{S}_0, \ell_0)$. We have an exact sequence

$$(2) \quad 0 \rightarrow \ell_0 \rightarrow \ell_0|_{JC} \oplus \ell_0|_{C \times C} \rightarrow \ell_0|_C \rightarrow 0.$$

The line bundle \mathcal{L}_0 on $JC \times JC$ can be written $\alpha \boxtimes \beta$, where α and β are 2-torsion line bundles on JC , not both trivial; we use the same letters to denote their restriction to C . The cohomology exact sequence associated to (2) gives

$$\begin{aligned} H^0(JC, \alpha \otimes \beta) \oplus H^0(C \times C, \alpha \boxtimes \beta) &\longrightarrow H^0(C, \alpha \otimes \beta) \longrightarrow H^1(\mathcal{S}_0, \ell_0) \xrightarrow{u} \\ &H^1(JC, \alpha \otimes \beta) \oplus H^1(C \times C, \alpha \boxtimes \beta) \longrightarrow H^1(C, \alpha \otimes \beta). \end{aligned}$$

The restriction map $H^0(JC, \alpha \otimes \beta) \rightarrow H^0(C, \alpha \otimes \beta)$ is surjective, so u is injective. If α and β are nontrivial, $H^1(C \times C, \alpha \boxtimes \beta)$ is zero, and the restriction map $H^1(JC, \alpha \otimes \beta) \rightarrow H^1(C, \alpha \otimes \beta)$ is injective, so $H^1(\mathcal{S}_0, \ell_0) = 0$. If, say, β is trivial, $H^1(JC, \alpha)$ is zero and the map $H^1(C \times C, \text{pr}_1^* \alpha) \rightarrow H^1(C, \alpha)$ is bijective, hence $H^1(\mathcal{S}_0, \ell_0) = 0$ again.

By semi-continuity this implies $H^1(\mathcal{S}_z, \mathcal{L}_z) = 0$ for z general in Δ , or equivalently $q(\tilde{\mathcal{S}}_z) = q(\mathcal{S}_z) = 4$, where $\tilde{\mathcal{S}} \rightarrow \mathcal{S}$ is the étale double covering defined by \mathcal{L} . But q is a topological invariant, so this holds for all $z \neq 0$ in Δ , hence $H^1(S, \ell) = 0$. ■

The surface X has a natural action of $(\mathbb{Z}/2)^2$, given by the involution i_X induced by (-1_B) and the involution τ associated to the double covering $X \rightarrow S$, which is induced by a translation of B . We want to determine how these involutions act on $H^0(X, K_X)$. The decomposition of $H^0(X, K_X)$ into eigenspaces for τ is

$$H^0(X, K_X) \cong H^0(S, K_S) \oplus H^0(S, K_S \otimes \ell).$$

By property b) above, i_S acts trivially on $H^0(S, K_S)$. It remains to study how it acts on $H^0(S, K_S \otimes \ell)$, or equivalently on $H^2(S, \ell)$. To define this action we choose the isomorphism $u : (-1_A)^* \ell \xrightarrow{\sim} \ell$ over A such that $u(0) = 1$, and we consider the involutions $H^p(i_S, u) : H^p(S, \ell) \xrightarrow{i_S^*} H^p(S, i_S^* \ell) \xrightarrow{u|_S} H^p(S, \ell)$.

Proposition 3. *There exist line bundles ℓ of order 2 on A for which i_S acts trivially on $H^2(S, \ell)$. In that case i_X has 48 fixed points.*

Proof : We will denote by A_2 and \hat{A}_2 the 2-torsion subgroups of A and its dual abelian variety \hat{A} , and similarly for B . The fixed point set of i_S is $A_2 \cap S$, and that of i_X is $B_2 \cap X$.

We apply the holomorphic Lefschetz formula to the automorphism i_S of S and the i_S -linearization $u|_S : i_S^* \ell \rightarrow \ell$:

$$\sum_p (-1)^p \operatorname{Tr} H^p(i_S, u) = \frac{1}{4} \sum_{a \in A_2 \cap S} u(a).$$

(At a point a of A_2 , $u(a) : \ell_a \rightarrow \ell_a$ is the multiplication by a scalar, which we still denote $u(a)$.)

Let $a \in A_2$. By [M1], property iv) p. 304, we have $u(a) = (-1)^{\langle a, \ell \rangle}$, where $\langle \cdot, \cdot \rangle : A_2 \times \hat{A}_2 \rightarrow \mathbb{Z}/2$ is the canonical pairing. On the other hand, dualizing the exact sequence of $(\mathbb{Z}/2)$ -vector spaces

$$0 \rightarrow (\mathbb{Z}/2)\ell \rightarrow \hat{A}_2 \xrightarrow{\hat{\pi}} \hat{B}_2$$

and using the canonical pairings we get an exact sequence

$$B_2 \xrightarrow{\pi} A_2 \xrightarrow{\langle \cdot, \ell \rangle} \mathbb{Z}/2 \rightarrow 0.$$

Thus $u(a) = 1$ if $a \in \pi(B_2)$, and $u(a) = -1$ otherwise. For $i = 0$ or 1 , let f_i be the number of points $a \in A_2 \cap S$ with $\langle a, \ell \rangle = i$. The right hand side of the Lefschetz formula is $\frac{1}{4}(f_0 - f_1)$.

We have $H^0(S, \ell) = H^1(S, \ell) = 0$ (Proposition 2), hence $\dim H^2(S, \ell) = \chi(\mathcal{O}_S) = 2$. Thus the left hand side is $\operatorname{Tr} H^2(i_S, u) \in \{2, 0, -2\}$. Since $f_0 + f_1 = 40$ this gives $f_i \in \{16, 20, 24\}$; the case $f_0 = 24$ corresponds to $H^2(i_S, u) = 1$. Moreover the number of fixed points of i_X is $\#(B_2 \cap X) = 2f_0$. Thus the Proposition will follow if we can find ℓ in \hat{A}_2 with $f_0 = 24$.

Put $F := A_2 \cap S$. Consider the homomorphism $\hat{A}_2 \rightarrow (\mathbb{Z}/2)^F$ given by $\ell \mapsto (\langle a, \ell \rangle)_{a \in F}$. For $\ell \neq 0$, the weight of the element $j(\ell)$ of $(\mathbb{Z}/2)^F$ (that is, the number of its nonzero coordinates) is f_1 , which belongs to $\{16, 20, 24\}$. Therefore j is injective; its image is a 8-dimensional vector subspace of $(\mathbb{Z}/2)^F$, that is, a linear code, such that the weight of any nonzero vector belongs to $\{16, 20, 24\}$. A simple linear algebra lemma ([B2], lemma 1) shows that a code in $(\mathbb{Z}/2)^{40}$ of dimension ≥ 7 contains elements of weight < 20 ; thus there exist elements ℓ in \hat{A}_2 with $f_1 = 16$, hence $f_0 = 24$. \blacksquare

From now on we choose ℓ so that i_S acts as trivially on $H^2(S, \ell)$. Thus i_X acts trivially on $H^0(X, K_X)$ and has 48 fixed points.

Proposition 4. *The canonical map $\varphi_X : X \rightarrow \mathbb{P}^6$ factors through i_X , and induces an isomorphism of X/i_X onto a complete intersection of 4 quadrics in \mathbb{P}^6 with 48 nodes.*

Proof : Since i_X acts as trivially on $H^0(X, K_X)$, we have a commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{\varphi_X} & \Sigma & \hookrightarrow & \mathbb{P}^6 \\ \downarrow \pi & & \downarrow p_\Sigma & & \downarrow p \\ S & \xrightarrow{\varphi_S} & \Xi & \hookrightarrow & \mathbb{P}^4 \end{array}$$

where φ_X and φ_S are the canonical maps, Σ and Ξ their images, p the projection corresponding to the injection $H^0(S, K_S) \rightarrow H^0(X, K_X)$, p_Σ its restriction to Σ .

The map $\varphi_S \circ \pi : X \rightarrow \Xi$ gives the quotient of X by the action of $(\mathbb{Z}/2)^2$. Since τ acts non-trivially on $H^0(X, K_X)$, φ_X identifies Σ with the quotient X/i_X . Thus all the maps in the left hand square of the above diagram are double coverings, étale outside finitely many points. In particular, since $K_X^2 = 32$, we have $\deg \Sigma = 16$.

We choose bases (x_0, \dots, x_4) and (u, v) of the $(+1)$ and (-1) -eigenspaces in $H^0(X, K_X)$ with respect to τ . The elements u^2, uv, v^2 of $H^0(X, K_X^{\otimes 2})$ are invariant under τ and i_X , therefore they are pull-back of i_S -invariant forms in $H^0(S, K_S^{\otimes 2})$. Such a form comes from an element of $H^0(\Xi, \mathcal{O}_\Xi(2))$, hence from an element of $H^0(\mathbb{P}^4, \mathcal{O}_{\mathbb{P}^4}(2))$. Thus we have

$$u^2 = a(x) \quad uv = b(x) \quad v^2 = c(x)$$

where a, b, c are quadratic forms in x_0, \dots, x_4 . Moreover the irreducible quadric Q containing Ξ is defined by a quadratic form $q(x)$ which vanishes on Σ .

Thus Σ is contained in the subvariety V of \mathbb{P}^6 defined by these 4 quadratic forms. If V is a surface, it has degree 16 and therefore is equal to Σ . Thus it suffices to prove that the morphism $p_V : V \rightarrow Q$ induced by the projection p is not surjective.

Assume that p_V is surjective; it has degree 2, and we have a cartesian diagram

$$\begin{array}{ccc} \Sigma & \hookrightarrow & V \\ \downarrow p_\Sigma & & \downarrow p_V \\ \Xi & \hookrightarrow & Q \end{array}$$

The variety V is irreducible: otherwise Σ is contained in one of its component, which maps birationally to Q , and p_Σ has degree 1, a contradiction. Since $Q \setminus \text{Sing}(Q)$ is simply connected, p_V is branched along a surface $R \subset Q$. Since Ξ is an ample divisor in Q (cut out by a quartic equation), it meets R along a curve, and p_Σ is branched along that curve, a contradiction. ■

Remark 2. It follows that $\Xi = p(\Sigma)$ is defined by the equations $q(x) = b(x)^2 - a(x)c(x) = 0$. The 40 nodes of Ξ break into two sets: the 16 points in \mathbb{P}^4 defined by $a(x) = b(x) = c(x) = q(x) = 0$ are the images by p_Σ of smooth points of Σ fixed by the involution induced by τ ; p_Σ is étale over the other 24 nodes of Ξ , giving rise to the 48 nodes of Σ .

Remark 3. The two families of surfaces X that we have constructed are different; in fact, a surface X_1 of the first family is not even homeomorphic to a surface X_2 of the second one. Indeed X_1 admits an irrational genus 2 pencil $X \rightarrow B$, and this is a topological property [C]. But for a general member

X_2 of the second family, the Albanese variety of the corresponding Schoen surface is simple [S], so its double cover $\text{Alb}(X_2)$ is also simple; therefore X_2 cannot have an irrational pencil of genus 2.

It follows that the corresponding surfaces Σ belong to two different connected components of the moduli space of complete intersections of 4 quadrics in \mathbb{P}^6 with an even set of 48 nodes.

REFERENCES

- [B1] A. Beauville : *L'application canonique pour les surfaces de type général*. Invent. math. **55** (1979), 121-140.
- [B2] A. Beauville : *Sur le nombre maximum de points doubles d'une surface dans \mathbb{P}^3 ($\mu(5) = 31$)*. Journées de Géométrie algébrique d'Angers, 207-215; Sijthoff & Noordhoff (1981).
- [CMR] C. Ciliberto, M. Mendes Lopes, X. Roulleau : *On Schoen surfaces*. Preprint arXiv:1303.1750.
- [C] F. Catanese : *Moduli and classification of irregular Kaehler manifolds (and algebraic varieties) with Albanese general type fibrations*. Invent. Math. **104** (1991), no. 2, 263-289.
- [CPT] C. Ciliberto, R. Pardini, F. Tovena : *Regular canonical covers*. Math. Nachr. **251** (2003), 19-27.
- [vL] R. van Luijk : *On perfect cuboids*. Undergraduate thesis, Universiteit Utrecht (2000).
- [M1] D. Mumford : *On the equations defining abelian varieties I*. Invent. Math. **1** (1966), 287-354.
- [M2] D. Mumford : *Tata lectures on theta, II*. Progress in Mathematics, **43**. Birkhäuser Boston, Inc., Boston, MA, 1984.
- [S] C. Schoen : *A family of surfaces constructed from genus 2 curves*. Internat. J. Math. **18** (2007), no. 5, 585-612.
- [ST] D. Testa, M. Stoll : *The surface parametrizing cuboids*. Preprint arXiv:1009.0388.

LABORATOIRE J.-A. DIEUDONNÉ, UMR 7351 DU CNRS, UNIVERSITÉ DE NICE, PARC VALROSE, F-06108 NICE CEDEX 2, FRANCE

E-mail address: arnaud.beauville@unice.fr