

ABOUT THE ALGEBRAIC CLOSURE OF THE FIELD OF POWER SERIES IN SEVERAL VARIABLES IN CHARACTERISTIC ZERO

GUILLAUME ROND

ABSTRACT. We begin this paper by constructing different algebraically closed fields containing an algebraic closure of the field of power series in several variables over a characteristic zero field. Each of these fields depends on the choice of an Abhyankar valuation and is constructed via a generalization of the Newton-Puiseux method for this valuation.

Then we study the Galois group of a polynomial with power series coefficients. In particular by examining more carefully the case of monomial valuations we are able to give several results concerning the Galois group of a polynomial whose discriminant is a weighted homogeneous polynomial times a unit. One of our main results is a generalization of Abhyankar-Jung Theorem for such polynomials, classical Abhyankar-Jung Theorem being devoted to polynomials whose discriminant is a monomial times a unit.

CONTENTS

1. Introduction	1
2. Notations and Abhyankar valuations	6
3. Homogeneous elements with respect to an Abhyankar valuation	9
3.1. Graded ring of an Abhyankar valuation and support	9
3.2. Homogeneous elements	10
4. Newton method and algebraic closure of $\mathbb{k}[[\mathbf{x}]]$ with respect to an Abhyankar valuation	14
4.1. Newton method	14
4.2. Analytically irreducible polynomials	18
5. Monomial valuation case: Eisenstein Theorem	22
6. Approximation of monomial valuations by divisorial monomial valuations	29
7. A generalization of Abhyankar-Jung Theorem	38
8. Diophantine Approximation	47
Notations	48
References	49

1. INTRODUCTION

When \mathbb{k} is an algebraically closed field of characteristic zero, we can always express the roots of a polynomial with coefficients in the field of power series over \mathbb{k} , denoted by $\mathbb{k}((t))$, as formal Laurent series in $t^{\frac{1}{k}}$ for some positive integer k . This result was known by Newton (at least formally see [BK] p. 372) and had been rediscovered by Puiseux in the complex analytic case [Pu1], [Pu2] (see [BK] or [Cu] for a presentation of this result). A modern way to reformulate

2000 *Mathematics Subject Classification*. Primary: 13F25. Secondary: 11J25, 12J20, 12F99, 13J05, 14B05, 32B10.

This work has been partially supported by ANR projects STAAVF (ANR-2011 BS01 009) and SUSI (ANR-12-JS01-0002-01).

this fact is to say that an algebraic closure of $\mathbb{k}((t))$ is the field of Puiseux power series \mathbb{P} defined in the following way:

$$\mathbb{P} := \bigcup_{k \in \mathbb{N}} \mathbb{k} \left(\left(t^{\frac{1}{k}} \right) \right).$$

The proof of this result, called the Newton-Puiseux method, consists essentially in constructing the roots of a polynomial $P(Z) \in \mathbb{k}[[t]][Z]$ by successive approximations in a similar way to Newton method in numerical analysis. These approximations converge since $\mathbb{k} \left(\left(t^{\frac{1}{k}} \right) \right)$ is a complete field with respect to the Krull topology.

This result, applied to a polynomial with coefficients in $\mathbb{k}[[t]]$ defining a germ of algebraic curve $(X, 0)$, provides an uniformization of this germ, i.e. a parametrization of this germ.

On the other hand this description of the algebraic closure of $\mathbb{k}((t))$ describes very easily the Galois group of $\mathbb{k}((t)) \rightarrow \mathbb{P}$, since this one is generated by the multiplication of the k -th roots of unity by $t^{\frac{1}{k}}$ for any positive integer k . In particular if an irreducible monic polynomial $P(Z) \in \mathbb{C}[[t]][Z]$ has a root which is a convergent power series in $t^{\frac{1}{k}}$, i.e. an element of $\mathbb{C}\{t^{\frac{1}{k}}\}$, then its other roots are also in $\mathbb{C}\{t^{\frac{1}{k}}\}$ and the coefficients of $P(Z)$ are convergent power series.

When \mathbb{k} is a characteristic zero field (but not necessarily algebraically closed), we can prove in the same way that an algebraic closure of $\mathbb{k}((t))$ is

$$(1) \quad \mathbb{P} := \bigcup_{\mathbb{k}'} \bigcup_{k \in \mathbb{N}} \mathbb{k}' \left(\left(t^{\frac{1}{k}} \right) \right).$$

where the first union runs over all finite field extensions $\mathbb{k} \rightarrow \mathbb{k}'$.

The aim of this work is double: the first one consists in finding representations of the roots of a polynomial whose coefficients are power series in several variables over a characteristic zero field. Our main results regarding these representations are Theorem 4.2 for Abhyankar valuations and its stronger version for monomial valuations (see Theorem 5.12). The second goal is to describe the Galois group of such polynomials. In particular we concentrate our study to irreducible polynomials that remain irreducible as polynomials with coefficients in the completion of the valuation ring associated to a monomial valuation. Our main result regarding this problem is a generalization of Abhyankar-Jung Theorem to polynomials whose discriminant is weighted homogeneous (see Theorems 7.5 and 7.7).

But let us present in more details the situation, the problems and the results given in this paper. It is tempting to find such a similar expression to (1) for the algebraic closure of the field of power series in n variables, $\mathbb{k}((x_1, \dots, x_n))$, for $n \geq 2$. But it appears easily that the algebraic closure of this field admits a really more complicated description and considering only power series depending on $x_1^{\frac{1}{k}}, \dots, x_n^{\frac{1}{k}}$ is not sufficient. For instance it is easy to see that a square root of $x_1 + x_2$ can not be expressed as such a power series.

Nevertheless there exist positive results in some specific cases, the most famous one being the Abhyankar-Jung theorem:

Theorem (Abhyankar-Jung Theorem). *If \mathbb{k} is an algebraically closed field of characteristic zero, then any polynomial with coefficients in $\mathbb{k}[[x_1, \dots, x_n]]$, whose discriminant has the form $ux_1^{\alpha_1} \dots x_n^{\alpha_n}$ where $u \in \mathbb{k}[[x_1, \dots, x_n]]$ is a unit and $\alpha_1, \dots, \alpha_n \in \mathbb{Z}_{\geq 0}$, has its roots in $\mathbb{k}[[x_1^{\frac{1}{k}}, \dots, x_n^{\frac{1}{k}}]]$ for some positive integer k .*

Such a polynomial is called a quasi-ordinary polynomial and this theorem asserts that the roots of quasi-ordinary polynomials are Puiseux power series in several variables. It provides not only a

description of the roots of a quasi-ordinary polynomial but also a description of its Galois group. This result has first being proven by Jung in the complex analytic case, then by Abhyankar in the general case ([Ju], [Ab]).

In the general case, a naive approach involves the use of Newton-Puiseux theorem n times (i.e. the formula (1) for the algebraic closure of $\mathbb{k}((t))$). For example in the case where $n = 2$ and \mathbb{k} is an algebraically closed field of characteristic zero, this means that the algebraic closure of $\mathbb{k}((x_1, x_2))$ is included in

$$\mathbb{L} := \bigcup_{k_2 \in \mathbb{N}} \bigcup_{k_1 \in \mathbb{N}} \mathbb{k} \left(\left(x_1^{\frac{1}{k_1}} \right) \right) \left(\left(x_2^{\frac{1}{k_2}} \right) \right).$$

But this field, which is algebraically closed, is very much larger than the algebraic closure of $\mathbb{k}((x_1, x_2))$ (see [Sa] for some thoughts about this). Moreover the action of the k_1 -th and k_2 -th roots of unity are not sufficient to generate the Galois group of the algebraic closure since there exist elements of $\mathbb{k}((x_1))((x_2))$ which are algebraic over $\mathbb{k}((x_1, x_2))$ but are not in $\mathbb{k}((x_1, x_2))$. For instance consider

$$x_1 \sqrt{1 + \frac{x_2}{x_1}} = \sum_{i \in \mathbb{Z}_{\geq 0}} a_i \frac{1}{x_1^{i-1}} x_2^i \in \mathbb{Q}((x_1))((x_2)) \setminus \mathbb{Q}((x_1, x_2))$$

for some well chosen rational numbers $a_i \in \mathbb{Q}$, $i \in \mathbb{Z}_{\geq 0}$.

Nevertheless a deeper analysis of the Newton-Puiseux method leads to the fact that it is enough to consider the field of fractions of the ring of elements

$$f = \sum_{(l_1, l_2) \in \mathbb{Z}^2} a_{l_1, l_2} x_1^{\frac{l_1}{k_1}} x_2^{\frac{l_2}{k_2}} \in \mathbb{L}$$

for some $k_1, k_2 \in \mathbb{N}$ whose support is included in a rational strongly convex cone of \mathbb{R}^2 . Here the support of f is the set

$$\text{Supp}(f) := \{(l_1, l_2) \in \mathbb{Z}^2 / a_{l_1, l_2} \neq 0\}.$$

This result has been proven by MacDonald [McD] (see also [Go], [Aro], [AI], [SV]). But once more, for any rational strongly convex cone of \mathbb{R}^2 , denoted by σ , $\mathbb{R}_{\geq 0}^2 \subsetneq \sigma$, there exist elements whose support is in σ but that are not algebraic over $\mathbb{k}((x_1, x_2))$.

One of the main difficulties comes from the fact that $\mathbb{k}((x_1, \dots, x_n))$ is not a complete field with respect to the topology induced by the maximal ideal of $\mathbb{k}[[x_1, \dots, x_n]]$ (called the Krull topology; it is induced by the following norm $\left| \frac{f}{g} \right| := e^{\text{ord}(g) - \text{ord}(f)}$ for any $f, g \in \mathbb{k}[[x_1, \dots, x_n]]$, $g \neq 0$, where $\text{ord}(f)$ is the order of the series f in the usual sense). Indeed, in order to apply the Newton-Puiseux method we have to work with a complete field since the roots are constructed by successive approximations. A very natural idea is to replace $\mathbb{k}((x_1, \dots, x_n))$ by its completion. But the completion of $\mathbb{k}((x_1, \dots, x_n))$ is not algebraic over $\mathbb{k}((x_1, \dots, x_n))$, thus the fields we construct in this way are bigger than the algebraic closure of $\mathbb{k}((x_1, \dots, x_n))$. In fact we need to replace the completion of $\mathbb{k}((x_1, \dots, x_n))$ by its henselization in the completion. The problem is that there is no general criterion to distinguish elements of the henselization from other elements of the completion. In some sense this problem is analogous to the fact that there is no general criterion to determine if a real number is algebraic or not over the rationals. One more issue is that choosing the Krull topology is arbitrary and we may replace this one by any topology induced by an other norm (or valuation) on this field.

In this paper we first investigate the use of the Newton-Puiseux method with respect to "tame"

valuations (i.e. replace $\mathbb{k}((x_1, \dots, x_n))$ by its completion for this valuation). By a "tame" valuation we mean a rank one (or real valued) valuation that satisfies the equality in the Abhyankar inequality (see Definition 2.1). These valuations are called Abhyankar valuations (cf. [ELS]) or quasi-monomial valuations (cf. [FJ]) and, essentially, these are monomial valuations after some sequence of blowing-ups. This is the first part of this work. If ν is such a valuation, we denote by $\widehat{\mathbb{K}}_\nu$ the completion of $\mathbb{k}((x_1, \dots, x_n))$ for the topology induced by this valuation. This field will play the role of $\mathbb{k}((t))$ in the classical Newton-Puiseux method. Then we have to define the elements that will play the role of $t^{\frac{1}{k}}$. This is where the first difficulty appears, since instead of working over $\widehat{\mathbb{K}}_\nu$, we need to work over the graded ring associated to ν . Both are isomorphic but there is no canonical isomorphism between them. In the case of $\mathbb{k}((t))$ where t is a single variable, such an isomorphism is defined by identifying the \mathbb{k} -vector space of homogeneous elements of degree i of the graded ring with the \mathbb{k} -vector space of homogeneous polynomials of degree i , i.e. $\mathbb{k}t^i$. But this identification depends on the choice of a uniformizer of $\mathbb{k}[[t]]$. In the case of $\mathbb{k}((x_1, \dots, x_n))$ an isomorphism will be determined by the choice of "coordinates" such that the valuation ν is monomial in these coordinates since Abhyankar valuations are monomial valuations after a sequence of blow-ups (cf. Remark 3.6). This is the reason why we restrict our study to these valuations. Nevertheless when such an isomorphism is chosen, we are able to define the elements that will play the role of $t^{\frac{1}{k}}$, this the aim of Section 3. These elements are called *homogeneous elements with respect to ν* (cf. Definitions 3.15 and 3.17). These are defined as being the roots of weighted homogeneous polynomials with coefficients in the graded ring of $\mathbb{k}[[x_1, \dots, x_n]]$ for the valuation ν . If \mathbb{k} is the field of complex numbers and the weights of the monomial valuation are positive integers, we can think about these homogeneous elements as weighted homogeneous algebraic (multivalued) functions. In fact we can replace $\widehat{\mathbb{K}}_\nu$ by a smaller field, the subfield of $\widehat{\mathbb{K}}_\nu$ whose elements have support included in a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$. Let us remark that this field is similar to the field of generalized power series $\cup_{\Gamma} \mathbb{C}((t^\Gamma))$ where the sum runs over all finitely generated semigroups Γ of $\mathbb{R}_{\geq 0}$ (see [Ri] for instance). Our first result is that the inductive limit of the extensions of $\widehat{\mathbb{K}}_\nu$ by homogeneous elements with respect to ν is algebraically closed (see Theorem 4.2). This field is $\lim_{\substack{\longrightarrow \\ \gamma_1, \dots, \gamma_s}} \widehat{\mathbb{K}}_\nu[\gamma_1, \dots, \gamma_s]$ where the limit runs over

all subsets $\{\gamma_1, \dots, \gamma_s\}$ of homogeneous elements with respect to ν and is denoted by $\overline{\mathbb{K}}_\nu$. The field extension $\mathbb{k}((x_1, \dots, x_n)) \rightarrow \overline{\mathbb{K}}_\nu$ factors through the field extension $\mathbb{k}((x_1, \dots, x_n)) \rightarrow \widehat{\mathbb{K}}_\nu$. While the Galois group of the field extension $\widehat{\mathbb{K}}_\nu \rightarrow \overline{\mathbb{K}}_\nu$ is easily described by the Galois group of weighted homogeneous polynomials, the Galois group of the algebraic closure of $\mathbb{k}((x_1, \dots, x_n))$ in $\widehat{\mathbb{K}}_\nu$ is more complicated. So it is very natural to study irreducible polynomials over $\mathbb{k}((x_1, \dots, x_n))$ which remain irreducible over $\widehat{\mathbb{K}}_\nu$, since their Galois groups are described by the Galois groups of weighted homogeneous polynomials. Proposition 4.14 shows that this property is an open property with respect to the topology induced by the chosen valuation. Let us mention that these polynomials are called ν -analytically irreducible polynomials in [Te] and their study is motivated by the construction of key polynomials for Abhyankar valuations (not necessarily of rank 1) in order to prove local uniformization.

Then we investigate more deeply the particular case of monomial valuations. In Section 5, using an idea of Tougeron [To] based on a work of Gabrielov [Ga], for any monomial valuation ν we construct a field, smaller than the ones constructed previously using the Newton-Puiseux method, and containing an algebraic closure of $\mathbb{k}((x_1, \dots, x_n))$. The main result (see Theorem 5.12) is a non-archimedean version of Eisenstein Theorem (classical Eisenstein Theorem concerns algebraic power series over \mathbb{Q}). The tool we use here is an effective version of the Implicit Function

Theorem (see Proposition 5.10). The elements we need to consider are of the form

$$(2) \quad \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}}$$

where the a_i and δ are weighted homogeneous polynomials for the weights corresponding to the given monomial valuation, Λ is a finitely sub-semigroup of $\mathbb{R}_{\geq 0}$, $\nu\left(\frac{a_i}{\delta^{m(i)}}\right) = i$ for all $i \in \Lambda$ and $i \mapsto m(i)$ is bounded by an affine function. In the particular case where the weights are \mathbb{Q} -linearly independent this corresponds to the result of MacDonald (see Theorem 6.9).

In Section 7 we use this description of the roots of polynomials with coefficients in $\mathbb{C}\{x_1, \dots, x_n\}$ to make a topological and complex analytical study of such polynomials whose discriminant is a weighted homogeneous polynomial multiplied by a unit. This study has been inspired by the work of Tougeron in [To] and more particularly by Remarque 2.7 of [To] where it is noticed that the elements of the form (2) define analytic functions on an open domain of \mathbb{C}^n which is the complement of some hornshaped neighborhood of $\{\delta = 0\}$ (see Definition 7.1). This study is possible in the case of monomial valuations whose weights are positive integers. To obtain the same results in the case of general monomial valuations we need to approximate general monomial valuations by divisorial monomial valuations, i.e. monomial valuations whose weights are positive integers. This is the subject of Section 6.

One of the main results we obtain in Section 7 is the following theorem which gives a criterion for an irreducible polynomial over $\mathbb{k}((x_1, \dots, x_n))$ to remain irreducible over $\widehat{\mathbb{K}}_\nu$:

Theorem. 7.5 *Let \mathbb{k} be a field of characteristic zero and $\alpha \in \mathbb{R}_{>0}^n$. Let \mathbf{x} denotes the set of variables (x_1, \dots, x_n) and let ν_α be the monomial valuation given by the weights α_i . Let $P(Z) \in \mathbb{k}[[\mathbf{x}]][[Z]]$ be a monic polynomial whose discriminant is equal to δu where $\delta \in \mathbb{k}[[\mathbf{x}]]$ is a weighted homogeneous polynomial for the weights $\alpha_1, \dots, \alpha_n$ and $u \in \mathbb{k}[[\mathbf{x}]]$ is a unit. If $P(Z)$ factors as $P(Z) = P_1(Z) \dots P_s(Z)$ where $P_i(Z)$ is an irreducible monic polynomial of $\mathbb{k}[[\mathbf{x}]][[Z]]$, then $P_i(Z)$ is irreducible in $\widehat{V}_\alpha[Z]$ where \widehat{V}_α denotes the completion of the valuation ring of ν_α .*

Then we show that Abhyankar-Jung Theorem is in fact a generalization of this result when the α_i are \mathbb{Q} -linearly independent (see Corollary 7.9) and we give the following generalization of Abhyankar-Jung Theorem for polynomials whose discriminant is weighted homogeneous with respect to weights $\alpha_1, \dots, \alpha_n \in \mathbb{R}_{>0}$:

Theorem. 7.7 *We assume that the hypothesis of Theorem 7.5 are satisfied. Let us set $N := \dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \dots + \mathbb{Q}\alpha_n)$. Then there exist $\gamma_1, \dots, \gamma_N$ integral homogeneous elements with respect to ν_α and a weighted homogeneous polynomial for the weights $\alpha_1, \dots, \alpha_n$ denoted by $c(\mathbf{x}) \in \mathbb{k}[[\mathbf{x}]]$ such that the roots of $P(Z)$ are in $\frac{1}{c(\mathbf{x})}\mathbb{k}'[[\mathbf{x}]][[\gamma_1, \dots, \gamma_N]]$ where $\mathbb{k} \rightarrow \mathbb{k}'$ is a finite field extension.*

Indeed in the case $N = n$, i.e. $\alpha_1, \dots, \alpha_n$ are \mathbb{Q} -linearly independent, the only weighted homogeneous polynomials are the monomials and the integral homogeneous elements with respect to ν_α are of the form \mathbf{x}^β where $\beta \in \mathbb{Q}_{>0}^n$ (see Remark 3.18). Abhyankar-Jung Theorem simply asserts that we may choose $c(\mathbf{x}) = 1$, a fact that we are able to prove in this case (see Corollary 7.9).

We remark that this result (along with Theorem 7.5) shows that the Galois group of an irreducible monic polynomial with coefficients in $\mathbb{k}[[x_1, \dots, x_n]]$ whose discriminant is weighted homogeneous is generated by the Galois group of one weighted homogeneous polynomial (see Remark 7.8).

Finally in Section 8 we give a result of Diophantine approximation (it is just an direct generalization of [Ro1] and [II]) that gives a necessary condition for an element of $\widehat{\mathbb{K}}_\nu$ to be algebraic over $\mathbb{k}((x_1, \dots, x_n))$.

At the end we give a list of notations for the convenience of the reader.

Let us mention that this work has been motivated by the understanding of the paper [To] of Tougeron where the study we make for monomial valuations is made in the case of the (x_1, \dots, x_n) -adic valuation of $\mathbb{k}((x_1, \dots, x_n))$.

I would like to thank Guy Casale and Adam Parusiński for their answers to my questions regarding the proofs of Lemma 7.4 and Lemma 7.2 respectively. I also thank H. Mourtada for the valuable discussions we had on these problems and his comments that helped to improve the presentation of this paper. I also thank the referees for their valuable suggestions.

2. NOTATIONS AND ABHYANKAR VALUATIONS

Let \mathbb{N} denote the set of positive integers and $\mathbb{Z}_{\geq 0}$ the set of non-negative integers. Let \mathbf{x} denote the multi-variable (x_1, \dots, x_n) where $n \geq 2$. Let \mathbb{k} denote a characteristic zero field. Then $\mathbb{k}[[\mathbf{x}]] = \mathbb{k}[[x_1, \dots, x_n]]$ denotes the ring of formal power series in n variables over \mathbb{k} and we denote by \mathbb{K}_n its fraction field and by \mathfrak{m} its maximal ideal.

When (A, \mathfrak{m}) is a local domain, a *valuation* on A is a function $\nu : A \setminus \{0\} \rightarrow \Gamma^+$, where Γ is an ordered subgroup of \mathbb{R} and $\Gamma^+ := \Gamma \cap \mathbb{R}_{\geq 0}$, such that

$$\nu(fg) = \nu(f) + \nu(g) \text{ and } \nu(f + g) \geq \min\{\nu(f), \nu(g)\} \quad \forall f, g \in A.$$

We will also impose that $\nu(f) > 0$ if and only if $f \in \mathfrak{m}$. We set $\nu(0) = \infty$ where $\infty > i$ for any $i \in \Gamma$.

Such valuation ν extends to \mathbb{K}_A , the fraction field of A , by

$$\nu\left(\frac{f}{g}\right) := \nu(f) - \nu(g)$$

for any $f, g \in A, g \neq 0$. We will always assume that $\nu : \mathbb{K}_A \rightarrow \Gamma$ is surjective. In this case Γ is called the *value group* of ν . The image of $A \setminus \{0\}$ by ν is called the *semigroup of ν* and we denote it by Σ . Then Γ is the group generated by Σ . Let us denote by V_ν the valuation ring of ν :

$$V_\nu := \left\{ \frac{f}{g} / f, g \in A, \nu(f) \geq \nu(g) \right\}.$$

This is a local ring whose maximal ideal, denoted by \mathfrak{m}_V , is the set of elements f/g such that $\nu(f/g) > 0$. Its residue field $\frac{V_\nu}{\mathfrak{m}_V}$ is denoted by \mathbb{k}_ν .

Let us denote by \widehat{V}_ν the completion of V_ν which is defined as follows: For any $\lambda \in \Gamma$ let us set $I_\lambda := \{v \in V_\nu / \nu(v) \geq \lambda\}$. The family of ideals $\{I_\lambda\}_{\lambda \in \Gamma}$ as a system of neighbourhoods of 0 makes V_ν into a topological ring. Then \widehat{V}_ν is the completion of V_ν for this topology. We can also remark that the family $\{V_\nu/I_\lambda\}_\lambda$ is an inverse system and its inverse limit is exactly \widehat{V}_ν .

Then \widehat{V}_ν is an equicharacteristic complete valuation ring and its residue field is isomorphic to \mathbb{k}_ν .

In this paper we will only consider a particular case of valuations, called Abhyankar valuations:

Definition 2.1. A valuation ν is called an *Abhyankar valuation* if the following equality holds:

$$\text{tr. deg}_{\mathbb{k}} \mathbb{k}_\nu + \dim_{\mathbb{Q}} \Gamma \otimes_{\mathbb{Z}} \mathbb{Q} = n.$$

This equality is called the *Abhyankar's Equality*.

Remark 2.2. If $\dim_{\mathbb{Q}} \Gamma \otimes \mathbb{Q} = 1$, then $\Gamma \simeq \mathbb{Z}$. Otherwise Γ is a dense subgroup of \mathbb{R} .

Example 2.3. The first example is the \mathfrak{m} -adic valuation denoted by ord on the ring $A = \mathbb{k}[[x]]$, and defined by

$$\text{ord}(f) := \max\{n \in \mathbb{N} / f \in \mathfrak{m}^n\} \quad \forall f \in \mathbb{k}[[\mathbf{x}]] \setminus \{0\}.$$

In this case its value group Γ is equal to \mathbb{Z} and its semigroup Σ is equal to $\mathbb{Z}_{\geq 0}$.

Example 2.4. Let $\alpha := (\alpha_1, \dots, \alpha_n) \in (\mathbb{R}_{>0})^n$. Let us denote by ν_α the monomial valuation on $A = \mathbb{k}[[x]]$ defined by $\nu_\alpha(x_i) := \alpha_i$ for $1 \leq i \leq n$. For instance $\nu_{(1, \dots, 1)} = \text{ord}$. Here we have $\Gamma = \mathbb{Z}\alpha_1 \oplus \dots \oplus \mathbb{Z}\alpha_n$ and $\Sigma = \mathbb{Z}_{\geq 0}\alpha_1 \oplus \dots \oplus \mathbb{Z}_{\geq 0}\alpha_n$.

Example 2.5. If Γ is isomorphic to \mathbb{Z} and ν is an Abhyankar valuation, then ν is a *divisorial valuation*. For such valuation there exists a proper birational dominant map $\pi : X \rightarrow \text{Spec}(\mathbb{k}[[\mathbf{x}]])$ and E an irreducible component of the exceptional locus of π such that ν is the composition of π^* with the \mathfrak{m}_E -adic valuation of the ring $\mathcal{O}_{X,E}$.

Remark 2.6. Geometrically, an Abhyankar valuation is a monomial valuation at a point lying on the exceptional divisor E of some proper birational map $(Y, E) \rightarrow (\mathbb{k}^n, 0)$. More precisely we have the following:

The restriction of ν to $\mathbb{k}[\mathbf{x}]$ is an Abhyankar valuation with the same value group as ν . We denote it by $\tilde{\nu}$. By Proposition 2.8 [ELS] there exists a regular local domain (A, \mathfrak{m}_A) , an injective morphism

$$\pi : \mathbb{k}[\mathbf{x}] \rightarrow A$$

inducing an isomorphism between the fields of fractions and a regular system of parameter z_1, \dots, z_r of A such that $\tilde{\nu}(z_1), \dots, \tilde{\nu}(z_r)$ freely generate the value group of $\tilde{\nu}$ (or the value group of ν since both are equal). Let us denote by μ the restriction of $\tilde{\nu}$ to A . Then π induces an isomorphism between V_ν and V_μ . Thus it induces an isomorphism between $\widehat{V}_{\tilde{\nu}}$ and \widehat{V}_μ . Moreover the completion of A is isomorphic to $\mathbb{L}[[z_1, \dots, z_r]]$ where $\mathbb{k} \rightarrow \mathbb{L}$ is a field extension of transcendence degree $n - r$ (here $\mathbb{L} = \frac{A}{\mathfrak{m}_A}$) and μ extends to a valuation on \widehat{A} which is exactly the monomial valuation that sends z_i onto $\nu(z_i)$ for all i .

Remark 2.7. If $n = 2$, in fact any discrete valuation (i.e. $\Gamma = \mathbb{Z}$) is an Abhyankar valuation [HOV].

Definition 2.8. Let $\alpha \in \mathbb{R}_{>0}^n$. A polynomial $f \in \mathbb{k}[[\mathbf{x}]]$ is called (α) -homogeneous of degree i if every nonzero monomial $c\mathbf{x}^\beta$ of f satisfies

$$\sum_{k=0}^n \alpha_k \beta_k = i$$

or equivalently $\nu_\alpha(c\mathbf{x}^\beta) = i$. This means that f is weighted homogeneous of degree i where x_j has weight α_j for every j .

Example 2.9. Let ν_α be a monomial valuation as before. Any power series $g \in \mathbb{k}[[\mathbf{x}]]$ can be written $g = \sum_{i \in \Sigma} g_i$ where g_i is a (α) -homogeneous polynomial of degree $i \in \Sigma$. Let us denote by i_0 the least $i \in \Sigma$ such that $g_i \neq 0$. Then we can write formally

$$g = g_{i_0} \left(1 + \sum_{i > i_0} \frac{g_i}{g_{i_0}} \right)$$

and this equality is satisfied in \widehat{V}_{ν_α} . Now if $f \in \mathbb{k}[[\mathbf{x}]]$, $g \neq 0$ and $\nu(f) \geq \nu(g)$ we can write

$$\frac{f}{g} = \left(\sum_i \frac{f_i}{g_{i_0}} \right) \left(1 + \sum_{i > i_0} \frac{g_i}{g_{i_0}} \right)^{-1}$$

where $f = \sum_i f_i$ where f_i is (α) -homogeneous of degree $i \in \Sigma$.

Thus any element of V_{ν_α} is of the form $\sum_{i \geq 0, i+i_0 \in \Sigma} \frac{a_i(\mathbf{x})}{b_i(\mathbf{x})}$ for some $i_0 \in \Sigma$, where $a_i(\mathbf{x})$ and $b_i(\mathbf{x})$ are (α) -homogeneous and $\nu_\alpha \left(\frac{a_i(\mathbf{x})}{b_i(\mathbf{x})} \right) = i$ for any $i \in \mathbb{R}$.

On the other hand \widehat{V}_{ν_α} is the set of elements of the form $\sum_{i \in \Lambda} \frac{a_i(\mathbf{x})}{b_i(\mathbf{x})}$ where Λ is a finite or countable subset of Γ^+ with no accumulation point, where $a_i(\mathbf{x})$ and $b_i(\mathbf{x})$ are (α) -homogeneous and $\nu_\alpha \left(\frac{a_i(\mathbf{x})}{b_i(\mathbf{x})} \right) = i$ for any $i \in \mathbb{R}$.

Let us denote by $\widehat{\mathbb{K}}_\nu$ the fraction field of \widehat{V}_ν . The valuation ν defines an ultrametric norm on $\widehat{\mathbb{K}}_\nu$, denoted by $|\cdot|_\nu$, defined by

$$\left| \frac{f}{g} \right|_\nu = e^{\nu(g) - \nu(f)} \quad \forall f \in \mathbb{k}[\mathbf{x}], g \in \mathbb{k}[\mathbf{x}] \setminus \{0\}.$$

Then $\widehat{\mathbb{K}}_\nu$ is the completion of \mathbb{K}_n for the topology induced by this norm and this norm (thus the valuation ν) extends canonically on $\widehat{\mathbb{K}}_\nu$. We shall also denote by ν the extension of ν to \mathbb{K}_ν .

Let us denote by $\mathbb{K}_\nu^{\text{alg}}$ the algebraic closure of \mathbb{K}_n in $\widehat{\mathbb{K}}_\nu$. We also denote by V_ν^{alg} the ring of elements of \widehat{V}_ν which are algebraic over \mathbb{K}_n : $V_\nu^{\text{alg}} := \mathbb{K}_\nu^{\text{alg}} \cap \widehat{V}_\nu$. We have the following lemma:

Lemma 2.10. *The ring V_ν^{alg} is a valuation ring (associated to the valuation ν) and $\mathbb{K}_\nu^{\text{alg}}$ is its fraction field. Moreover $V_\nu \rightarrow V_\nu^{\text{alg}}$ is the henselization of V_ν in \widehat{V}_ν .*

Proof. If $f, g \in V_\nu^{\text{alg}}$ and $\nu(f) \geq \nu(g)$, then $\frac{f}{g} \in \mathbb{K}_\nu^{\text{alg}} \cap \widehat{V}_\nu = V_\nu^{\text{alg}}$ so V_ν^{alg} is a valuation ring. For $f \in \mathbb{K}_\nu^{\text{alg}}$ there exists $N \in \mathbb{N}$ such that $x_1^N f \in \mathbb{K}_\nu^{\text{alg}} \cap \widehat{V}_\nu = V_\nu^{\text{alg}}$ since $\nu(x_1^N) > 0$. Thus $\mathbb{K}_\nu^{\text{alg}}$ is the fraction field of V_ν^{alg} .

By construction the elements of the henselization of V_ν are algebraic over V_ν . On the other hand every element of \widehat{V}_ν which is algebraic over V_ν is in the Henselization of V_ν (see Corollary 1.2.1 [M-B]). \square

Thus we can summarize the situation with the following commutative diagram, where the bottom part corresponds to the quotient fields of the rings of the upper part:

$$\begin{array}{ccccc}
 \mathbb{k}[\mathbf{x}] & \longrightarrow & V_\nu & \longrightarrow & \widehat{V}_\nu \\
 & & \downarrow & \searrow & \downarrow \\
 & & & & V_\nu^{\text{alg}} \\
 & & \downarrow & \swarrow & \downarrow \\
 & & \mathbb{K}_n & \longrightarrow & \widehat{\mathbb{K}}_\nu \\
 & & \downarrow & \searrow & \downarrow \\
 & & & & \mathbb{K}_\nu^{\text{alg}}
 \end{array}$$

3. HOMOGENEOUS ELEMENTS WITH RESPECT TO AN ABHYANKAR VALUATION

3.1. Graded ring of an Abhyankar valuation and support. Let A be an integral domain and let $\nu : A \rightarrow \Gamma^+$ be a valuation where Γ is a subgroup of \mathbb{R} . We define $\text{Gr}_\nu A = \bigoplus_{i \in \Gamma^+} \frac{\mathfrak{p}_{\nu,i}}{\mathfrak{p}_{\nu,i}^+}$ where $\mathfrak{p}_{\nu,i} := \{f \in A / \nu(f) \geq i\}$ and $\mathfrak{p}_{\nu,i}^+ := \{f \in A / \nu(f) > i\}$.

Definition 3.1. Let Γ^+ be a sub-semigroup of $\mathbb{R}_{\geq 0}$. A Γ^+ -graded ring is a ring A that has a direct sum of abelian groups, $A = \bigoplus_{i \in \Gamma^+} A_i$, such that $A_i A_j \subset A_{i+j}$ for any $i, j \in \Gamma^+$.

For any $j \in \Gamma^+$, $\bigoplus_{i \in \Gamma^+, i \geq j} A_i$ is an ideal of A . This family of ideals as a system of neighborhoods of 0 makes A into a topological whose completion is denoted by \widehat{A} or $\widehat{\bigoplus_{i \in \Gamma^+} A_i}$. The completion of A is the set of elements that are written as a series $\sum_{i \in \Lambda} a_i$ where $\Lambda \subset \Gamma^+$ is either a finite set, either a countable subset of $\mathbb{R}_{>0}$ with no accumulation point, and $a_i \in A_i$ for any $i \in \Lambda$.

A *complete* (Γ^+ -)graded ring is the completion of a (Γ^+ -)graded ring.

Remark 3.2. Let A be a complete graded ring. If A_0 is a field then A is a local ring and its maximal ideal is $\mathfrak{m} := \widehat{\bigoplus_{i > 0} A_i}$.

For any $a \in A$ we can write $a = \sum_{i \in \Lambda} a_i$ where $a_i \in A_i$ for any i . If $a \neq 0$ let us set $\nu(a) := \min\{i \in \Gamma^+ / a_i \neq 0\}$. Set $\nu(0) = \infty$. Then ν is an order function, i.e. $\nu(ab) \geq \nu(a) + \nu(b)$ and $\nu(a+b) \geq \min\{\nu(a), \nu(b)\}$. Moreover ν is a valuation if and only if A is an integral domain. The order function ν is called the *order function of A* .

Example 3.3. For a given Abhyankar valuation ν on $\mathbb{k}[\mathbf{x}]$ the rings $\text{Gr}_\nu \mathbb{k}[\mathbf{x}]$ and $\text{Gr}_\nu V_\nu$ are Γ^+ -graded rings and $\widehat{\text{Gr}_\nu \mathbb{k}[\mathbf{x}]}$ and $\widehat{\text{Gr}_\nu V_\nu}$ are complete Γ^+ -graded rings.

Remark 3.4. The ring $\widehat{\text{Gr}_\nu V_\nu}$ is isomorphic to the ring of generalized power series $\mathbb{k}_\nu[[t^{\Gamma^+}]]$ where t is a single variable.

Remark 3.5. The elements of $\widehat{\text{Gr}_\nu V_\nu}$ are the elements of the form $\sum_{i \in \Lambda} a_i$ where $a_i \in \frac{\mathfrak{p}_{\nu,i}}{\mathfrak{p}_{\nu,i}^+}$ for all $i \in \Lambda$ where Λ is either a finite set, either a countable subset of $\mathbb{R}_{\geq 0}$ with no accumulation point.

Remark 3.6. Let us consider a monomial valuation ν on $\mathbb{k}[\mathbf{x}]$, let us say $\nu := \nu_\alpha$ where $\alpha \in \mathbb{R}_{>0}^n$. In this case $\frac{\mathfrak{p}_{\nu,i}}{\mathfrak{p}_{\nu,i}^+}$ is isomorphic to the \mathbb{k} -vector space of rational fractions $\frac{a(\mathbf{x})}{b(\mathbf{x})}$ where $a(\mathbf{x})$ and $b(\mathbf{x})$ are (α) -homogeneous polynomials and $\nu_\alpha \left(\frac{a(\mathbf{x})}{b(\mathbf{x})} \right) = i$. Thus, by Example 2.9 $\widehat{\text{Gr}_\nu V_\nu}$ and \widehat{V}_ν are \mathbb{k} -isomorphic.

Let us now consider a general Abhyankar valuation ν on $\mathbb{k}[\mathbf{x}]$. By Remark 2.6 there exist a regular local domain (A, \mathfrak{m}_A) , an injective morphism

$$\pi : \mathbb{k}[\mathbf{x}] \rightarrow A$$

inducing an isomorphism between the fields of fractions and such that, if we denote by μ the restriction of ν to A , the following properties hold:

The extension of μ to \widehat{A} is a monomial valuation (denoted by $\widehat{\mu}$) and π induces isomorphisms $V_\nu \simeq V_\mu$ and $\widehat{V}_\nu \simeq \widehat{V}_\mu$.

We have $\widehat{V}_\mu = \widehat{V}_{\widehat{\mu}}$ and $\text{Gr}_\nu V_\nu \simeq \text{Gr}_\mu V_\mu = \text{Gr}_\mu \widehat{V}_\mu$. Thus $\widehat{\text{Gr}_\nu V_\nu}$ and \widehat{V}_ν are \mathbb{k} -isomorphic by the monomial case.

We can summarize this in the following proposition:

Proposition 3.7. *The choice of a proper birational map π and parameters z_1, \dots, z_r as in Remark 2.6 yields an isomorphism*

$$\widehat{\text{Gr}_\nu V_\nu} \simeq \widehat{V}_\nu.$$

Remark 3.8. A different choice of π and z_1, \dots, z_r would give an other isomorphism between these two rings.

Definition 3.9. Let $A = \widehat{\bigoplus}_{i \in \Gamma^+} A_i$ be a complete Γ^+ -graded ring. Let $a \in A$, $a = \sum_{i \in \Gamma^+} a_i$, $a_i \in A_i$ for any i . The support of a is the subset I of Γ^+ defined by $i \in I$ if and only if $a_i \neq 0$. We denote this set I by $\text{Supp}(a)$.

Definition 3.10. Let ν be an Abhyankar valuation defined on $\mathbb{k}[[\mathbf{x}]]$. Let us fix a \mathbb{k} -isomorphism φ between $\widehat{\text{Gr}_\nu V_\nu}$ and \widehat{V}_ν as in Proposition 3.7. Let $a \in \widehat{V}_\nu$ and let us write $\varphi(a) = \sum_{i \in \Gamma^+} a_i$ with $a_i \in \frac{\mathfrak{p}_{\nu,i}^+}{\mathfrak{p}_{\nu,i}^+}$. The ν -support with respect to φ of a is the subset of Γ^+ defined as

$$\text{Supp}_{\nu,\varphi}(a) := \{i \in \Gamma^+ / a_i \neq 0\}.$$

When the isomorphism is clear from the context we will skip the mention of φ and denote the ν -support of a by $\text{Supp}_\nu(a)$.

Proposition 3.11. *Let ν be an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$ and let φ be a \mathbb{k} -isomorphism between $\widehat{\text{Gr}_\nu V_\nu}$ and \widehat{V}_ν as in Proposition 3.7. Then there exists a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$, denoted by Λ , such that the ν -support of any element of $\mathbb{k}[[\mathbf{x}]]$ with respect to φ is included in Λ .*

Proof. By Remark 2.6, we may assume that ν is a monomial valuation. Thus the proposition comes from the following lemma applied to $\Sigma = \mathbb{Z}_{\geq 0}^n$: □

Lemma 3.12. *Let Σ be a strongly convex rational cone of \mathbb{R}^n . Let $\alpha \in \mathbb{R}_{> 0}^n$ such that $\langle \alpha, \beta \rangle > 0$ for any $\beta \in \Sigma$, $\beta \neq 0$. Then there exists a finitely generated subgroup of $\mathbb{R}_{\geq 0}$, denoted by Λ , such that $\text{Supp}_{\nu_\alpha}(f) \subset \Lambda$ for any $f \in \mathbb{k}[[x^\beta, \beta \in \Sigma \cap \mathbb{Z}^n]]$ where $\mathbb{k}[[x^\beta, \beta \in \Sigma \cap \mathbb{Z}^n]]$ denotes the ring of formal Laurent series whose support is included in $\Sigma \cap \mathbb{Z}^n$.*

Proof. By Gordan Lemma, $\Sigma \cap \mathbb{Z}^n$ is a finitely generated semigroup, let us say $\Sigma \cap \mathbb{Z}^n$ is generated by u_1, \dots, u_k . Let us set $r_i := \langle \alpha, u_i \rangle$, $1 \leq i \leq k$. Since any element of $\Sigma \cap \mathbb{Z}^n$ is a $\mathbb{Z}_{\geq 0}$ -linear combination of u_1, \dots, u_k , then $\langle \alpha, \beta \rangle$ is a $\mathbb{Z}_{\geq 0}$ -linear combination of r_1, \dots, r_k for any $\beta \in \Sigma \cap \mathbb{Z}^n$. Let us denote by Λ the semigroup of $\mathbb{R}_{\geq 0}$ generated by r_1, \dots, r_k . Then $\text{Supp}_{\nu_\alpha}(f) \subset \Lambda$. □

Remark 3.13. Proposition 3.11 does not imply that the semigroup Σ of ν is finitely generated, which is not true in general for Abhyankar valuations which are not monomial valuations.

3.2. Homogeneous elements. From now on we fix an Abhyankar valuation ν on $\mathbb{k}[[\mathbf{x}]]$ and a \mathbb{k} -isomorphism φ between $\widehat{\text{Gr}_\nu V_\nu}$ and \widehat{V}_ν induced by an injective birational morphism π as in Remark 3.6 and we will skip to mention it in the following. There are several reasons for that. The first one is that we are interested in effective results on the algebraic elements over $\mathbb{k}[[\mathbf{x}]]$, thus we are interested by valuations which are given effectively and this will be the case essentially through a map π as in Remark 2.6. In particular we will investigate more deeply the case of monomial valuations and, in this case, the set of variables x_1, \dots, x_n will be fixed from the beginning, thus φ is quite natural in this case. The last reason is that we will give properties on the ν -support of algebraic elements, and Proposition 3.11 will allow us to consider only elements whose ν -support is included in a finitely generated sub-semigroup of $\mathbb{R}_{> 0}$, and this fact does not depend on φ .

Definition 3.14. Let ν be an Abhyankar valuation defined on $\mathbb{k}[[\mathbf{x}]]$. We will denote by V_ν^{fg} the subset of \widehat{V}_ν of elements whose ν -support is included in a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ (when we identify \widehat{V}_ν and $\widehat{\text{Gr}_\nu V_\nu}$ via φ). It is straightforward to check that V_ν^{fg} is a valuation ring. We denote by $\mathbb{K}_\nu^{\text{fg}}$ its fraction field.

Definition 3.15. Let A be a complete Γ^+ -graded domain and let ν be its order function (which is a valuation since A is a domain). A *homogeneous element with respect to ν* is an element γ of a finite extension of A such that its minimal polynomial $Q(Z)$ is irreducible in $A[Z]$ and has the following form:

$$Z^q + g_1 Z^{q-1} + \cdots + g_q$$

where $g_k \in A_{i(k)}$ with $i(k) \in \Gamma$ for $1 \leq k \leq q$ such that $k \cdot i(l) = l \cdot i(k)$ for all k and l . In this case $d := \frac{i(k)}{k} \in \frac{1}{q!} \Gamma$ is called the *order* of γ .

Example 3.16. Let $\alpha \in \mathbb{R}_{>0}^n$ such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n$ i.e. the α_i are \mathbb{Q} -linearly independent. Then the value group of ν_α is the following group:

$$\Gamma = \mathbb{Z}\alpha_1 + \cdots + \mathbb{Z}\alpha_n$$

and for any $i \in \Gamma$ there exists a unique $(\beta_{i,1}, \dots, \beta_{i,n}) \in \mathbb{Z}^n$ such that

$$i = \beta_{i,1}\alpha_1 + \cdots + \beta_{i,n}\alpha_n.$$

Thus if $i \in \Gamma^+$ this means that $\frac{\mathfrak{p}_{\nu_\alpha, i}}{\mathfrak{p}_{\nu_\alpha, i}^+}$ is isomorphic to the one dimensional \mathbb{k}_{ν_α} -vector space generated by $x_1^{\beta_{i,1}} \cdots x_n^{\beta_{i,n}}$. Let us remark here that \mathbb{k}_{ν_α} is equal to \mathbb{k} since the α_i are \mathbb{Q} -linearly independent. Thus if $g_k \in \frac{\mathfrak{p}_{\nu_\alpha, dk}}{\mathfrak{p}_{\nu_\alpha, dk}^+}$ for $1 \leq k \leq q$ we have that

$$Z^q + g_1 Z^{q-1} + \cdots + g_q = x_1^{\beta_{qd,1}} \cdots x_n^{\beta_{qd,n}} (T^q + g'_1 T^{q-1} + \cdots + g'_q)$$

where $Z = x_1^{\beta_{d,1}} \cdots x_n^{\beta_{d,n}} T$ and $g'_1, \dots, g'_q \in \mathbb{k}$. If $g_q \neq 0$ then $\beta_{qd,j} \in \mathbb{Z}$ for any j but $\beta_{d,j} = \frac{\beta_{qd,j}}{d}$ may not be an integer. Then the roots of $T^q + g'_1 T^{q-1} + \cdots + g'_q$ are algebraic over \mathbb{k} . Thus homogeneous elements with respect to ν_α are of the form $c\mathbf{x}^\beta$ where c is algebraic over \mathbb{k} and $\beta \in \mathbb{Q}^n$ with $\langle \alpha, \beta \rangle := \alpha_1 \beta_1 + \cdots + \alpha_n \beta_n \geq 0$.

Definition 3.17. Let ν be an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$. Let $A = \widehat{\text{Gr}_\nu V_\nu}$ and γ be a homogeneous element with respect to ν . Let $Q(Z)$ be its minimal polynomial: $Q(Z) = Z^q + g_1 Z^{q-1} + \cdots + g_q$ with $g_k \in \frac{\mathfrak{p}_{\nu, dk}}{\mathfrak{p}_{\nu, dk}^+}$ for $1 \leq k \leq q$. We say that γ is an *integral homogeneous element with respect to ν* if g_k is the image of an element of $\mathbb{k}[[\mathbf{x}]] \cap \mathfrak{p}_{\nu, dk}$ for all k .

Example 3.18. Let $\alpha \in \mathbb{R}_{>0}^n$ such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n$ and let us keep the notations of Example 3.16. Then γ is an integral homogeneous element with respect to ν_α if $g_k \in \frac{\mathbb{k}[[\mathbf{x}]] \cap \mathfrak{p}_{\nu_\alpha, dk}}{\mathbb{k}[[\mathbf{x}]] \cap \mathfrak{p}_{\nu_\alpha, dk}^+}$ for $1 \leq k \leq q$. Since $g_q \neq 0$ this means that $\beta_{qd,j} \in \mathbb{Z}_{\geq 0}$ for all j . Thus integral homogeneous elements with respect to ν_α are of the form $c\mathbf{x}^\beta$ where c is algebraic over \mathbb{k} and $\beta \in \mathbb{Q}_{\geq 0}^n$.

Example 3.19. Let ν be an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$ and let us assume that \mathbb{k} is not algebraically closed. Let c be in the algebraic closure of \mathbb{k} , $c \notin \mathbb{k}$. Then c is a root of a polynomial equation with coefficients in \mathbb{k} and since \mathbb{k} is a subfield of \mathbb{k}_ν , this shows that c is an integral homogeneous element of order 0 with respect to ν .

Remark 3.20. Let ν be an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$ and let γ be a homogeneous element of order d with respect to ν . Let us denote by $Q(Z)$ its minimal polynomial, say

$$Q(Z) = Z^q + g_1 Z^{q-1} + \cdots + g_q$$

where $g_k \in \frac{\mathfrak{p}_{\nu_\alpha, dk}}{\mathfrak{p}_{\nu_\alpha, dk}^+}$ for $1 \leq k \leq q$. Each g_k is the image in $\widehat{\text{Gr}_\nu V_\nu}$ of some fraction $\frac{f_k}{h_k}$ where $f_k, h_k \in \mathbb{k}[[\mathbf{x}]]$. Set $h := h_1 \cdots h_k$, let h_0 be the image of h in $\widehat{\text{Gr}_\nu V_\nu}$ and set $\gamma' := h_0 \gamma$. Then γ' is a

homogeneous element annihilating $Z^q + g'_1 Z^{q-1} + \dots + g'_q$ where g'_k is the image of $\frac{f_k}{h_k} h^{k-1} \in \mathbb{k}[[\mathbf{x}]]$ in $\widehat{\text{Gr}_\nu V_\nu}$, thus it is an integral homogeneous element with respect to ν . Moreover we have

$$\text{Frac}(\widehat{\text{Gr}_\nu V_\nu})[\gamma] = \text{Frac}(\widehat{\text{Gr}_\nu V_\nu})[\gamma'].$$

Definition 3.21. Let ν be an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$,

$$P(Z_1, \dots, Z_m) \in \widehat{V}_\nu[Z_1, \dots, Z_m]$$

and $\mathbf{d} := (d_1, \dots, d_m) \in \mathbb{R}_{>0}^m$. One says that $P(Z_1, \dots, Z_m)$ is (ν, \mathbf{d}) -homogeneous of degree $d \in \mathbb{R}$ if for every nonzero monomial $gZ_1^{\beta_1} \dots Z_m^{\beta_m}$ of $P(Z)$ one has $g \in \frac{\mathfrak{p}_{\nu, k}^{\nu, k}}{\mathfrak{p}_{\nu, k}^+}$ with $k + \beta_1 d_1 + \dots + \beta_m d_m = d$.

Remark 3.22. Let ν be an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$. Let γ be a homogeneous element of order d with respect to ν . Let us denote by $P(Z)$ its minimal monic polynomial. Then $P(Z)$ is (ν, d) -homogeneous.

Conversely if $P(Z) \in \widehat{V}_\nu[Z]$ satisfies $P(\gamma) = 0$ for some element γ algebraic over \widehat{V}_ν , and if $P(Z)$ is a nonzero (ν, d) -homogeneous, then the divisors of P in $\widehat{V}_\nu[Z]$ are also (ν, d) -homogeneous, thus the minimal polynomial of γ is (ν, d) -homogeneous. Hence γ is a homogeneous element of order d with respect to ν .

Lemma 3.23. Let γ_1 and γ_2 be two homogeneous elements of order d_1 and d_2 respectively with respect to the valuation ν and let $k \in \mathbb{Z}$. Then

- i) γ_1^k is homogeneous of order kd_1 ,
- ii) if $e_1 d_1 = e_2 d_2$ with $e_1, e_2 \in \mathbb{N}$, then $\gamma_1^{e_1} + \gamma_2^{e_2}$ is homogeneous of order $d_1 e_1$,
- iii) $\gamma_1 \gamma_2$ is homogeneous of order $d_1 + d_2$.

Proof. If γ is homogeneous of order $d \in \mathbb{Q}$, then γ^k , $k \in \mathbb{N}$, is homogeneous of order kd . Indeed a polynomial having γ^k as a root is $Q(Z) := \text{Res}_X(P(X), Z - X^k)$ where P is the minimal monic polynomial of γ over $\mathbb{k}(\mathbf{x})$. But $P(X)$ is (ν, d) -homogeneous and $Z - X^k$ is (ν, d, kd) -homogeneous. Thus $Q(Z)$ is (ν, d, kd) -homogeneous, hence (ν, kd) -homogeneous since it does not depend on X . This proves that γ^k is homogeneous of order kd .

In order to show ii) we may assume, by i), that γ_1 and γ_2 are homogeneous of same order $d = e_1 d_1 = e_2 d_2$. Let us denote by $P_1(Z)$ and $P_2(Z)$ the minimal monic polynomials of γ_1 and γ_2 respectively. Then $Q(Z) := \text{Res}_X(P_1(Z - X), P_2(X))$ is (ν, d, d) -homogeneous, thus (ν, d) -homogeneous since it does not depend on X . Since $Q(\gamma_1 + \gamma_2) = 0$, $\gamma_1 + \gamma_2$ is homogeneous of order d .

In order to show iii) let us denote by $P_1(X)$ the minimal monic polynomial of γ_1 (this is a (ν, d_1) -homogeneous polynomial) and $P_2(Z)$ the minimal monic polynomial of γ_2 ((ν, d_2) -homogeneous). Let us denote by k the degree in Z of $P_1(Z)$ and set $R(X, Y) := X^k P_1(Y/X)$. Then $\gamma_1 \gamma_2$ is a root of $Q(Z) := \text{Res}_X(R(X, Z), P_2(X))$. Moreover $R(X, Z)$ is $(\nu, d_2, d_1 + d_2)$ -homogeneous. Thus $Q(Z)$ is $(\nu, d_1 + d_2)$ -homogeneous, which proves that $\gamma_1 \gamma_2$ is homogeneous of order $d_1 + d_2$. \square

Lemma 3.24. Let $P(T, Z)$ be a nonzero (ν, d_1, d_2) -homogeneous polynomial of $\widehat{V}_\nu[T, Z]$ and let γ_1 be a homogeneous element of order d_1 with respect to ν . If an element γ_2 belonging to a finite extension of $\mathbb{k}(\mathbf{x})$ satisfies $P(\gamma_1, \gamma_2) = 0$, then γ_2 is a homogeneous element of order d_2 with respect to ν .

Proof. Let $Q(T) \in \widehat{V}_\nu[T]$ be a nonzero (ν, d_1) -homogeneous polynomial such that $Q(\gamma_1) = 0$. Let us denote $R(Z) = \text{Res}_T(P(T, Z), Q(T))$. Then $R(Z)$ is a (ν, d_2) -homogeneous polynomial such that $R(\gamma_2) = 0$. This proves the result. \square

Remark 3.25. Let A be a complete Γ^+ -graded integral domain, let say A is the completion of $A' := \bigoplus_{i \in \Gamma^+} A_i$, and let ν be its order valuation. Let $Q(Z)$ be an irreducible polynomial of $A[Z]$ having the following form:

$$Z^q + g_1 Z^{q-1} + \cdots + g_q$$

where $g_k \in A_{dk}$ for $1 \leq k \leq q$ and $d \in \frac{1}{q!}\Gamma^+$. The ring $B := \frac{A[Z]}{(Q(Z))}$ is an integral domain and ν extends to a valuation of this ring by defining $\nu(Z) := d$ and

$$\nu\left(\sum_{i=0}^{q-1} a_i Z^i\right) := \inf_i \{\nu(a_i) + di\}.$$

Let us set $B' := A'[Z]/(Q(Z))$. Then B is a complete $\frac{1}{q!}\Gamma$ -graded domain since B is the completion of

$$B' = \bigoplus_{i \in \Gamma^+} \bigoplus_{\substack{0 \leq j \leq \min\{\lfloor \frac{i}{d} \rfloor, q\} \\ j \in \frac{1}{q!}\Gamma^+}} A_{i-dj} Z^j.$$

Definition 3.26. Let γ be an algebraic element over A whose minimal polynomial is the polynomial $Q(Z)$ as in the previous remark. Then the integral domain B constructed in the previous remark is denoted by $A[\gamma]$.

By induction we can define $A[\gamma_1, \dots, \gamma_s]$ where γ_{i+1} is a homogeneous element over $A[\gamma_1, \dots, \gamma_i]$ for $1 \leq i < s$. When ν is an Abhyankar valuation on $\mathbb{k}[[\mathbf{x}]]$ and $A = \widehat{V}_\nu, V_\nu^{\text{alg}}$ or V_ν^{fg} , the valuation ν extends to $A[\gamma_1, \dots, \gamma_i]$ as in Remark 3.25. Then we denote by $A[\langle \gamma_1, \dots, \gamma_s \rangle]$ the valuation ring associated to the order valuation of $A[\gamma_1, \dots, \gamma_s]$. In this case the elements of $A[\langle \gamma_1, \dots, \gamma_s \rangle]$ are the elements which are finite sums of the form $b\gamma_1^{j_1} \dots \gamma_s^{j_s}$ where $b \in \text{Frac}(A)$ and $\nu(b) \geq -(j_1\nu(\gamma_1) + \dots + j_s\nu(\gamma_s))$.

Definition 3.27. If ν is an Abhyankar valuation we denote by

$$\overline{V}_\nu := \varinjlim_{\gamma_1, \dots, \gamma_s} \widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_s \rangle]$$

the direct limit over all subsets $\{\gamma_1, \dots, \gamma_s\}$ of homogeneous elements with respect to ν and by $\overline{\mathbb{K}}_\nu$ its fraction field. By Remark 3.20 we may restrict the limit over the subsets of integral homogeneous elements.

In the same way we define

$$\overline{V}_\nu^{\text{fg}} := \varinjlim_{\gamma_1, \dots, \gamma_s} V_\nu^{\text{fg}}[\langle \gamma_1, \dots, \gamma_s \rangle],$$

$$\overline{V}_\nu^{\text{alg}} := \varinjlim_{\gamma_1, \dots, \gamma_s} V_\nu^{\text{alg}}[\langle \gamma_1, \dots, \gamma_s \rangle],$$

the limits being taken over all subsets $\{\gamma_1, \dots, \gamma_s\}$ of (integral) homogeneous elements with respect to ν , and we denote by $\overline{\mathbb{K}}_\nu^{\text{fg}}$ and $\overline{\mathbb{K}}_\nu^{\text{alg}}$ their respective fraction fields.

The following result provides an upper bound on the number of homogeneous elements we need to consider:

Proposition 3.28. *Let ν be an Abhyankar valuation on $\mathbb{k}[\mathbf{x}]$ and let Γ denote its value group. Set $N := \dim_{\mathbb{Q}} \Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$ and let $\gamma_1, \dots, \gamma_s$ be homogeneous elements with respect to ν . Then there exist integral homogeneous elements $\gamma'_1, \dots, \gamma'_N$ with respect to ν such that $\widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_s \rangle] = \widehat{V}_\nu[\langle \gamma'_1, \dots, \gamma'_N \rangle]$.*

This equality remains true if we replace \widehat{V}_ν by V_ν^{alg} or V_ν^{fg} .

Proof. We will prove this proposition by induction on s . Let $\gamma_1, \dots, \gamma_{N+1}$ be nonzero homogeneous elements with respect to ν . Let d_i be the order of γ_i , for $1 \leq i \leq N+1$. By assumption on N the d_i are \mathbb{Q} -linearly dependent. Thus, after a permutation of the g_i , there exists an integer $1 \leq l \leq N$ and integers $p_i \in \mathbb{Z}_{\geq 0}$, $q_i \in \mathbb{N}$ for all $1 \leq i \leq N+1$, such that

$$(3) \quad \frac{p_1}{q_1} d_1 + \dots + \frac{p_l}{q_l} d_l = \frac{p_{l+1}}{q_{l+1}} d_{l+1} + \dots + \frac{p_{N+1}}{q_{N+1}} d_{N+1}.$$

Set $r_i := \frac{p_1 \cdots p_{N+1}}{p_i}$ for $1 \leq i \leq N+1$. Let us denote $\gamma'_i := \gamma_i^{\frac{1}{q_i r_i}}$. Then we have

$$\widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_{N+1} \rangle] \subset \widehat{V}_\nu[\langle \gamma'_1, \dots, \gamma'_{N+1} \rangle].$$

By (3) and Lemma 3.23, $\gamma'_1 \cdots \gamma'_l$ and $\gamma'_{l+1} \cdots \gamma'_{N+1}$ are homogeneous elements of same order. By the Primitive Element Theorem there exists $c \in \mathbb{k}$ such that

$$\mathbb{k}(\mathbf{x})[\gamma'_1 \cdots \gamma'_l, \gamma'_{l+1} \cdots \gamma'_{N+1}] = \mathbb{k}(\mathbf{x})[\gamma'_1 \cdots \gamma'_l + c \gamma'_{l+1} \cdots \gamma'_{N+1}].$$

Moreover $\gamma := \gamma'_1 \cdots \gamma'_l + c \gamma'_{l+1} \cdots \gamma'_{N+1}$ is a homogeneous element with respect to ν of same order as $\gamma'_1 \cdots \gamma'_l$ and $\gamma'_{l+1} \cdots \gamma'_{N+1}$ by Lemma 3.23. Since

$$\mathbb{k}(\mathbf{x})[\gamma'_1, \dots, \gamma'_l] = \mathbb{k}(\mathbf{x})[\gamma'_1, \dots, \gamma'_{l-1}, \gamma'_1 \cdots \gamma'_l]$$

and

$$\mathbb{k}(\mathbf{x})[\gamma'_{l+1}, \dots, \gamma'_{N+1}] = \mathbb{k}(\mathbf{x})[\gamma'_{l+1}, \dots, \gamma'_N, \gamma'_{l+1} \cdots \gamma'_{N+1}],$$

we have

$$\mathbb{k}(\mathbf{x})[\gamma'_1, \dots, \gamma'_{N+1}] = \mathbb{k}(\mathbf{x})[\gamma'_1, \dots, \gamma'_{l-1}, \gamma'_{l+1}, \dots, \gamma'_N, \gamma].$$

Thus γ'_l is a finite sum of products of elements $a_i(\mathbf{x}) \in \mathbb{k}(\mathbf{x})$ and powers of $\gamma'_1, \dots, \gamma'_{l-1}, \gamma'_{l+1}, \dots, \gamma'_N, \gamma$ and by homogeneity we may assume that $a_i(\mathbf{x})$ are (ν) -homogeneous. Thus

$$\widehat{V}_\nu[\langle \gamma'_1, \dots, \gamma'_{N+1} \rangle] = \widehat{V}_\nu[\langle \gamma'_1, \dots, \gamma'_{l-1}, \gamma'_{l+1}, \dots, \gamma'_N, \gamma \rangle].$$

By Remark 3.20 we may assume that the γ'_i are integral homogeneous elements.

The proof is the same if we replace \widehat{V}_ν by V_ν^{alg} or V_ν^{fg} . □

4. NEWTON METHOD AND ALGEBRAIC CLOSURE OF $\mathbb{k}[\mathbf{x}]$ WITH RESPECT TO AN ABHYANKAR VALUATION

4.1. Newton method.

Lemma 4.1. *Let (A, \mathfrak{m}) be a complete graded local ring. Let B be the set of the elements of A whose support is included in a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$. Then B is a Henselian local domain.*

Proof. Let us prove that B is a ring: let b_1 and b_2 be two elements of B whose supports are included in Λ_1 and Λ_2 respectively. Thus we can write $b_i = \sum_{j \in \Lambda_i} b_{i,j}$ where $b_{i,j}$ is a homogeneous element of degree j for any $i = 1, 2$ and $j \in \Lambda_1$ or Λ_2 . Let Λ be the finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ generated by Λ_1 and Λ_2 . Then $\text{Supp}(b_1 + b_2)$ and $\text{Supp}(b_1 b_2)$ are included in Λ . This proves that B is a ring. Since $B \subset A$, B is a domain.

It is clear that $\mathfrak{m} \cap B$ is an ideal of B . If $b \in B \setminus (\mathfrak{m} \cap B)$, then there exists $a \in A$ such that

$ab = 1$. Let us write $b = \sum_{i \in \Lambda} b_i$ where b_i is homogeneous of degree i and Λ is a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$. Since $b \notin \mathfrak{m}$, then $b_0 \neq 0$. In this case we have

$$a = b^{-1} = \frac{1}{b_0} \left(1 + \sum_{i \in \Lambda \setminus \{0\}} \frac{b_i}{b_0} \right)^{-1} = \frac{1}{b_0} \sum_{k=1}^{\infty} (-1)^k \left(\sum_{i \in \Lambda \setminus \{0\}} \frac{b_i}{b_0} \right)^k.$$

Thus $\text{Supp}(a) \subset \Lambda$. This proves that B is a local ring with maximal ideal $\mathfrak{m} \cap B$.

Now let $P(Z) \in B[Z]$, such that $P(0) \in \mathfrak{m} \cap B$ and $P'(0) \notin \mathfrak{m}$. We denote by ν the order function of A , i.e. if $a \in A$, $a \neq 0$, $a = \sum_i a_i$ where a_i is homogeneous of degree i , $\nu(a) := \inf\{i / a_i \neq 0\}$ and the initial term of a is $\text{in}(a) := a_{\nu(a)}$. Since A is a complete local ring it is a Henselian local ring and there exists $a \in \mathfrak{m}$ such that $P(a) = 0$. We can construct a by using the fact that

$$(4) \quad P(Z) = P(0) + P'(0)Z + Q(Z)Z^2$$

where $Q(Z) \in B[Z]$. Indeed, let Λ denote a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ containing the supports of all the coefficients of $P(Z)$. In this case $a_1 := \text{in}(a) = -\frac{\text{in}(P(0))}{\text{in}(P'(0))}$ is a homogeneous element of degree $d_1 \in \Lambda$, $d_1 > 0$. If we set $P_1(Z) := P(Z + a_1)$, we see that

$$\nu(P_1(0)) = \nu(P(a_1)) > d_1,$$

$P'_1(0) = P'(0) = 0$ and $a - a_1$ is the solution of $P_1(Z) = 0$ given by the Hensel Lemma. Then we replace P by P_1 in Equation (4) and repeat the same argument, using the fact that the coefficients of $P_1(Z)$ have support included in Λ . Thus we see that $\text{in}(a - a_1) = -\frac{\text{in}(P_1(0))}{\text{in}(P'_1(0))}$ is a homogeneous element of degree $d_2 \in \Lambda$, $d_2 > d_1$. We repeat this operation a countable number of times (since Λ is countable) in order to construct a and we see that $\text{Supp}(a) \subset \Lambda$. □

Now we can prove the following theorem:

Theorem 4.2. *Let \mathbb{k} be a field of characteristic zero and ν be an Abhyankar valuation of $\mathbb{k}[[\mathbf{x}]]$. Let $N := \dim_{\mathbb{Q}} \Gamma \otimes_{\mathbb{Z}} \mathbb{Q}$. Let*

$$P(Z) \in V_{\nu}^{\text{fg}}[\langle \gamma_1, \dots, \gamma_N \rangle][Z]$$

(resp. $\widehat{V}_{\nu}[\langle \gamma_1, \dots, \gamma_N \rangle][Z]$) be a monic polynomial of degree d where γ_i is a homogeneous element with respect to ν for $1 \leq i \leq N$. Then there exist integral homogeneous elements $\gamma'_1, \dots, \gamma'_N$ such that the roots of $P(Z)$ are in $V_{\nu}^{\text{fg}}[\langle \gamma'_1, \dots, \gamma'_N \rangle]$ (resp. $\widehat{V}_{\nu}[\langle \gamma'_1, \dots, \gamma'_N \rangle]$).

Proof. Let us prove the case $P(Z) \in V_{\nu}^{\text{fg}}[\langle \gamma_1, \dots, \gamma_N \rangle][Z]$. We write $P(Z) = Z^d + a_1 Z^{d-1} + \dots + a_d$. By replacing Z by $Z - \frac{1}{d} a_1$ we can assume that $a_1 = 0$. Let i_0 be an integer such that

$$\frac{\nu(a_{i_0})}{i_0} \leq \frac{\nu(a_i)}{i}, \quad \text{for every } 2 \leq i \leq d.$$

Let γ be a i_0 th root of $\text{in}_{\nu}(a_{i_0})$, i.e. γ is a homogeneous element such that $\gamma^{i_0} = \text{in}_{\nu}(a_{i_0})$. By the definition of i_0 , for every $2 \leq i \leq d$ we can write

$$a_i = \gamma^i a'_i$$

with $a'_i \in V_{\nu}^{\text{fg}}[\langle \gamma_1, \dots, \gamma_N, \gamma \rangle]$. Then we have

$$P(\gamma Z) = \gamma^d Z^d + \gamma^{d-2} a_2 Z^{d-2} + \dots + a_d = \gamma^d (Z^d + a'_2 Z^{d-2} + \dots + a'_d)$$

Let $S(Z) := Z^d + a'_2 Z^{d-2} + \dots + a'_d$ and let $\overline{S}(Z)$ be the image of $S(Z)$ in the residue field $\mathbb{L} = V_{\nu}^{\text{fg}}[\langle \gamma_1, \dots, \gamma_s, \gamma \rangle] / \mathfrak{m}$ where $\mathbb{k}_{\nu} \rightarrow \mathbb{L}$ is finite and \mathfrak{m} is the maximal ideal of $V_{\nu}^{\text{fg}}[\langle \gamma_1, \dots, \gamma_s, \gamma \rangle]$. If $\overline{S}(Z) = (Z + \overline{a})^d$ where $\overline{a} \in \mathbb{L}$, since $a_1 = 0$ and $\text{char}(\mathbb{L}) = 0$, this would imply $\overline{a} = 0$. But $\overline{S}(Z) \neq Z^d$ since its coefficient of Z^{d-i_0} is nonzero. Thus we can factor $\overline{S}(Z) = \overline{S}_1(Z) \overline{S}_2(Z)$ such

that $\overline{S}_1(Z)$ and $\overline{S}_2(Z)$ are coprime monic polynomials in $\mathbb{L}[\gamma'][[Z]]$ where γ' is algebraic over \mathbb{L} , i.e. γ' is a homogeneous element of degree 0 with respect to ν . Since $V_\nu^{\text{fg}}[\langle\gamma_1, \dots, \gamma_N, \gamma, \gamma'\rangle]$ is a Henselian local ring by Lemma 4.1, by Hensel Lemma the polynomial $S(Z)$ factors as $S(Z) = S_1(Z)S_2(Z)$ where the images of $S_1(Z)$ and $S_2(Z)$ in $V_\nu^{\text{fg}}[\langle\gamma_1, \dots, \gamma_N, \gamma, \gamma'\rangle]$ are $\overline{S}_1(Z)$ and $\overline{S}_2(Z)$ and the ν -support of the coefficients of $S_1(Z)$ and $S_2(Z)$ are contained in a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$.

Since $\deg_Z(S_1(Z)), \deg_Z(S_2(Z)) < d = \deg_Z(P(Z))$, the theorem is proven by induction on d by using Proposition 3.28 and Remark 3.20.

The case $P(Z) \in \widehat{V}_\nu[\langle\gamma_1, \dots, \gamma_N\rangle][Z]$ is proven in a similar way by using the fact that $\widehat{V}_\nu[\langle\gamma_1, \dots, \gamma_N, \gamma, \gamma'\rangle]$ is a complete local ring, thus a Henselian local ring. \square

Remark 4.3. The proof of this theorem is what we call the *Newton-Puiseux method*. Usually the term of Newton-Puiseux method is used when one compute the roots of a monic polynomial with coefficients in the ring of power series in one variable: one root is constructed by computing step by step its coefficients. The fact that the ring of formal power series is a complete local ring allows to conclude that this process converges. But when we want to find roots of a polynomial in a local ring that is not complete but only Henselian, it is more convenient to use the Hensel Lemma as we have done here. The proof we used here appeared for the first time in [BM] (to the knowledge of the author). Of course if ν is a divisorial valuation \widehat{V}_ν is isomorphic to the ring of formal power series in one variable over the residue field \mathbb{k}_ν and the previous theorem may be proven by using the classical Newton-Puiseux method.

Corollary 4.4. *The field $\overline{\mathbb{K}}_\nu^{\text{fg}}$ (resp. $\overline{\mathbb{K}}_\nu$) is algebraically closed and it is the algebraic closure of $\mathbb{K}_\nu^{\text{fg}}$ (resp. \mathbb{K}_ν).*

Proof. Let $P(Z) \in \overline{\mathbb{K}}_\nu^{\text{fg}}[Z]$ be an irreducible polynomial. By multiplying $P(Z)$ by an element of V_ν^{fg} , we may assume that

$$P(Z) \in V_\nu^{\text{fg}}[\langle\gamma_1, \dots, \gamma_N\rangle][Z]$$

for some homogeneous elements $\gamma_1, \dots, \gamma_N$ with respect to ν . We write $P(Z) = a_d Z^d + \dots + a_0$, $a_i \in V_\nu^{\text{fg}}[\langle\gamma_1, \dots, \gamma_N\rangle]$, $0 \leq i \leq d$. We set $Q(Z) := a_d^{d-1} P(Z/a_d)$. Then $Q(Z)$ is a monic polynomial of $V_\nu^{\text{fg}}[\langle\gamma_1, \dots, \gamma_N\rangle][Z]$ and if z is a root of $Q(Z)$, then $\frac{z}{a_d}$ is a root of $P(Z)$. Hence the result comes from Theorem 4.2.

The assertion concerning $\overline{\mathbb{K}}_\nu$ is proven similarly. \square

We have the similar result for $\overline{\mathbb{K}}^{\text{alg}}$:

Lemma 4.5. *The algebraic closure of \mathbb{K}_n in $\overline{\mathbb{K}}_\nu$ is equal to $\overline{\mathbb{K}}_\nu^{\text{alg}}$. In particular $\overline{\mathbb{K}}_\nu^{\text{alg}}$ is algebraically closed.*

Proof. Let $\gamma_1, \dots, \gamma_s$ be homogeneous elements with respect to ν . Let us denote by q_{i+1} the degree of the minimal polynomial of γ_{i+1} over $\mathbb{K}_n[\gamma_1, \dots, \gamma_i]$ for $0 \leq i \leq s-1$. Thus any element z of $\widehat{\mathbb{K}}_\nu[\gamma_1, \dots, \gamma_s]$ can be uniquely written as $z = \sum_{i \in I} A_{i_1, \dots, i_s} \gamma_1^{i_1} \dots \gamma_s^{i_s}$ where $A_{i_1, \dots, i_s} \in \widehat{\mathbb{K}}_\nu$ for all $i \in I$ and $I = \{0, \dots, q_1 - 1\} \times \dots \times \{0, \dots, q_s - 1\}$.

In order to prove the lemma we need to show that $A_{i_1, \dots, i_s} \in \mathbb{K}_\nu^{\text{alg}}$ for any i_1, \dots, i_s when z is algebraic over $\mathbb{k}[\mathbf{x}]$. In this case let $\mathbb{L} := \widehat{\mathbb{K}}_\nu[\gamma_1, \dots, \gamma_{s-1}]$ and let us write $z := \sum_{i=0}^{q_s-1} B_i \gamma_s^i$

where $B_i \in \mathbb{L}$ for all i . Let us set $\zeta_1 := \gamma_s$ and let $\zeta_2, \dots, \zeta_{q_s}$ be the conjugates of ζ_1 over $\overline{\mathbb{K}}_\nu[\gamma_1, \dots, \gamma_{s-1}]$. Let us define $z_j = \sum_{i=0}^{q_s-1} B_i \zeta_j^i$ for $1 \leq j \leq q_s$. Then we have

$$\begin{pmatrix} z_1 \\ z_2 \\ \vdots \\ z_{q_s} \end{pmatrix} = \begin{pmatrix} 1 & \zeta_1 & \cdots & \zeta_1^{q_s-1} \\ 1 & \zeta_2 & \cdots & \zeta_2^{q_s-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \zeta_{q_s} & \cdots & \zeta_{q_s}^{q_s-1} \end{pmatrix} \begin{pmatrix} B_0 \\ B_1 \\ \vdots \\ B_{q_s-1} \end{pmatrix}.$$

The matrix $\begin{pmatrix} 1 & \zeta_1 & \cdots & \zeta_1^{q_s-1} \\ 1 & \zeta_2 & \cdots & \zeta_2^{q_s-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \zeta_{q_s} & \cdots & \zeta_{q_s}^{q_s-1} \end{pmatrix}$ is invertible and its entries are algebraic over $\mathbb{k}(\mathbf{x})$, z_j is algebraic over $\mathbb{k}[\mathbf{x}]$ for all j , hence B_j is algebraic over $\mathbb{k}[\mathbf{x}]$ for all j . By induction on s we see that $A_{i_1, \dots, i_s} \in \mathbb{K}_\nu^{\text{alg}}$ for any i_1, \dots, i_s . \square

We can summarize the situation with the following commutative diagram where the bottom part corresponds to the quotient fields of the rings of the upper part and all the morphisms are injective:

$$\begin{array}{ccccccc} \mathbb{k}[\mathbf{x}] \subset V_\nu & \longrightarrow & V_\nu^{\text{alg}} & \longrightarrow & V_\nu^{\text{fg}} & \longrightarrow & \widehat{V}_\nu \\ \downarrow & & \searrow & & \searrow & & \searrow \\ \mathbb{K}_n & \longrightarrow & \mathbb{K}_\nu^{\text{alg}} & \longrightarrow & \mathbb{K}_\nu^{\text{fg}} & \longrightarrow & \widehat{\mathbb{K}}_\nu \\ & & \downarrow & & \downarrow & & \downarrow \\ & & \mathbb{K}_\nu^{\text{alg}} & \longrightarrow & \mathbb{K}_\nu^{\text{fg}} & \longrightarrow & \mathbb{K}_\nu \end{array}$$

Example 4.6. Let $g(T) = \sum_{i=1}^{\infty} c_i T^i \in \mathbb{Q}[[T]]$ be a formal power series which is not algebraic over $\mathbb{Q}[T]$. Let $\alpha := (\alpha_1, \alpha_2) \in \mathbb{N}^n$. Let us set

$$f := g\left(\frac{x_2^{\alpha_1}}{x_1^{\alpha_2}}\right) = \sum_{i=1}^{\infty} c_i \frac{x_2^{\alpha_1 i}}{x_1^{\alpha_2 i}} \in \mathbb{k}((x_1))((x_2)).$$

But $f \notin \overline{\mathbb{K}}_{\nu_\alpha}$: let $P(Z) = a_0(\mathbf{x})Z^d + \cdots + a_d(\mathbf{x}) \in \widehat{V}_{\nu_\alpha}[Z]$ be a polynomial such that $P(f) = 0$. Let us write $a_i(\mathbf{x}) = \sum_{k=0}^{\infty} a_{i,k}(\mathbf{x})$ where $a_{i,k}(\mathbf{x})$ is a (α_1, α_2) -homogeneous rational fraction of degree k . By homogeneity we have

$$a_{0,k} f^d + a_{1,k} f^{d-1} + \cdots + a_{d,k} = 0 \quad \forall k \in \mathbb{N}.$$

This implies that

$$a_{0,k}(1, T)g(T^{\alpha_1})^d + a_{1,k}(1, T)g(T^{\alpha_1})^{d-1} + \cdots + a_{d,k}(1, T) = 0 \quad \forall k \in \mathbb{N}.$$

Thus $a_{i,k}(\mathbf{x}) = 0$ for all $0 \leq i \leq d$ and $0 \leq k$. Hence $P(Z) = 0$ and $f \notin \overline{\mathbb{K}}_{\nu_\alpha}$.

On the other hand, $h := g\left(\frac{x_1^{2\alpha_2}}{x_2^{\alpha_1}}\right) = \sum_{i=1}^{\infty} c_i \frac{x_1^{2\alpha_2 i}}{x_2^{\alpha_1 i}} \in \widehat{\mathbb{K}}_{\nu_\alpha}$ but h is not algebraic over $\mathbb{k}((x_1))((x_2))$.

4.2. Analytically irreducible polynomials.

Proposition 4.7. *Let $P(Z) \in V_\nu^{\text{fg}}[Z]$ (resp. $V_\nu^{\text{alg}}[Z]$) be an irreducible monic polynomial. Then $P(Z)$ is irreducible in $\widehat{V}_\nu[Z]$.*

Proof. By Corollary 4.4, $P(Z)$ splits in $V_\nu^{\text{fg}}[\langle \gamma_1, \dots, \gamma_s \rangle]$ for some homogeneous elements $\gamma_1, \dots, \gamma_s$ with respect to ν . Since

$$V_\nu^{\text{fg}}[\langle \gamma_1, \dots, \gamma_s \rangle] \cap \widehat{V}_\nu = V_\nu^{\text{fg}}$$

the result follows.

The proof is the same for V_ν^{alg} . \square

Lemma 4.8. *Let σ be a $\widehat{\mathbb{K}}_\nu$ -automorphism of $\overline{\mathbb{K}}_\nu$. For any $z \in \overline{\mathbb{K}}_\nu$ we have $\nu(\sigma(z)) = \nu(z)$.*

Proof. Let $z \in \widehat{\mathbb{K}}_\nu[\gamma_1, \dots, \gamma_s]$ where $\gamma_1, \dots, \gamma_s$ are homogeneous elements with respect to ν . Let us write $z := \sum_{i \in \Lambda} z_i$ where z_i is homogeneous of degree i for every i and Λ is a countable subset of \mathbb{R} with no accumulation point (see Remark 3.5). If $i_0 = \nu(z)$, then $z_{i_0} \neq 0$ and $\nu(z_i) = 0$ for all $i < i_0$. Since σ acts only on the homogeneous elements $\gamma_1, \dots, \gamma_s$, we have $\sigma(z) = \sum_i \sigma(z_i)$. For all i , $\sigma(z_i)$ is homogeneous of degree i and $\sigma(z_i) = 0$ if and only if $z_i = 0$. This proves that $i_0 = \nu(\sigma(z))$. \square

Definition 4.9. Let $P(Z) \in A[Z]$ where A is an integral domain. We write

$$P(Z) = a_0 Z^d + a_1 Z^{d-1} + \dots + a_d.$$

Let $\nu : A \rightarrow \mathbb{R}_{\geq 0}$ be a valuation. The *Newton polygon* of P is the convex hull of the set

$$\{(\nu(a_i), d - i) \in \mathbb{R}_{\geq 0}^2 / i = 0, \dots, d\} + \mathbb{R}_{\geq 0}^2.$$

Corollary 4.10. *Let $P(Z) \in \widehat{V}_\nu[Z]$ be an irreducible monic polynomial. Then the Newton polygon of $P(Z)$ has only one edge. The result remains valid if we replace \widehat{V}_ν by V_ν^{alg} or V_ν^{fg} .*

Proof. Let z be a root of $P(Z)$ in \overline{V}_ν . Let σ be a $\widehat{\mathbb{K}}_\nu$ -automorphism of $\overline{\mathbb{K}}_\nu$. Then $\nu(\sigma(z)) = \nu(z)$ by Lemma 4.8. The finite product of the distinct linear forms $Z - \sigma(z)$ obtained in this way is a monic polynomial with coefficients in $\widehat{\mathbb{K}}_\nu$ and divides $P(Z)$. Since $P(Z)$ is irreducible, both polynomials are equal. This proves that all the roots of $P(Z)$ have same valuation, hence the Newton polygon of $P(Z)$ has only one edge.

The cases V_ν^{alg} and V_ν^{fg} are deduced from Lemma 4.7. \square

Example 4.11. Let $P(Z) := Z^3 + 3x_1x_2Z - 2x_1^4 \in \mathbb{k}[[x_1, x_2]][Z]$. We see that $P(Z)$ has one root of order 2 and two roots of order 1 in $\overline{V}_{\text{ord}}^{\text{fg}}$. By Corollary 4.10, $P(Z)$ has at least one root in $V_{\text{ord}}^{\text{alg}}$ of order 2.

Let $\sqrt{1+U} := 1 + \sum_{i \geq 1} a_i U^i$, $a_i \in \mathbb{Q}$ for all i , the formal powers series whose square is equal to $1+U$, and let $\sqrt[3]{1+U} := 1 + \sum_{i \geq 1} b_i U^i$, $b_i \in \mathbb{Q}$ for all i , the formal power series whose cube is equal to $1+U$. Then the roots of $P(Z)$ are

$$a \sqrt[3]{q + \sqrt{q^2 + p^3}} + b \sqrt[3]{q - \sqrt{q^2 + p^3}}$$


with $(a, b) = (1, 1)$, (j, j^2) or (j^2, j) and $p = x_1x_2$ and $q = x_1^4$. But

$$\sqrt[3]{q + \varepsilon \sqrt{q^2 + p^3}} = \sqrt[3]{x_1^4 + \varepsilon \sqrt{x_1^3 x_2^3 + x_1^8}} = \sqrt[3]{\varepsilon} \sqrt{x_1 x_2} + \eta$$

where $\varepsilon = 1$ or -1 and $\text{ord}(\eta) > 1$. Both order 1 roots of $P(Z)$ have initial term of the form $\alpha\sqrt{x_1x_2}$ where $\alpha \in \mathbb{C}^*$. Thus $P(Z)$ has only one root in $V_{\text{ord}}^{\text{alg}}$ and even in $\mathbb{K}_{\text{ord}}^{\text{fg}}$.

Let z be the only root of $P(Z)$ in $V_{\text{ord}}^{\text{alg}}$. If $z \in \mathbb{K}_n$, since $P(Z)$ is monic and $\mathbb{k}[[\mathbf{x}]]$ is an integral domain, then $z \in \mathbb{k}[[\mathbf{x}]]$. But $\text{in}(z) = \frac{2}{3}\frac{x_1^3}{x_2} \notin \mathbb{k}[[\mathbf{x}]]$. Thus $z \notin \mathbb{K}_n$, hence $P(Z)$ is irreducible in $\mathbb{K}_n[Z]$. This shows that $\mathbb{K}_n \rightarrow \mathbb{K}_{\text{ord}}^{\text{alg}}$ is not a normal extension in general.

Corollary 4.12. *Let $P(Z) := Z^d + a_1(\mathbf{x})Z^{d-1} + \dots + a_d(\mathbf{x}) \in \mathbb{k}[[\mathbf{x}]][[Z]]$ be an irreducible polynomial having its roots in $\mathbb{k}[[x_1^{\frac{1}{e}}, \dots, x_n^{\frac{1}{e}}]]$ for some positive integer e . Then the Newton polyhedron of $P(Z)$ is the convex hull of the cone of \mathbb{N}^{n+1} centered in $(0, \dots, 0, d)$ and generated by the convex hull of the Newton polyhedra of $a_d(\mathbf{x})$ in \mathbb{N}^n .*



Proof. Let $\alpha \in \mathbb{N}^n$. Let $z_1, \dots, z_d \in \mathbb{k}[[x_1^{\frac{1}{e}}, \dots, x_n^{\frac{1}{e}}]]$ be the roots of $P(Z)$. Then $z_i \in \widehat{V}_{\nu_\alpha}[x_1^{\frac{1}{e}}, \dots, x_n^{\frac{1}{e}}]$ for any i , the $x_i^{\frac{1}{e}}$ being homogeneous elements with respect to ν_α . Let $G \simeq (\mathbb{Z}/e\mathbb{Z})^n$ be the Galois group of the extension $\widehat{V}_{\nu_\alpha} \rightarrow \widehat{V}_{\nu_\alpha}[x_1^{\frac{1}{e}}, \dots, x_n^{\frac{1}{e}}]$. The z_i are conjugated under the action of G , thus $P(Z) := \prod_{i=1}^d (Z - z_i)$ is irreducible in $\widehat{V}_{\nu_\alpha}[Z]$. This being true for any $\alpha \in \mathbb{N}^n$, the result follows from Corollary 4.10. □

We finish this section by giving two results relating the roots of a polynomial $P(Z)$ to the roots of polynomials approximating $P(Z)$. First of all we give the following definition:

Definition 4.13. Let $P(Z) \in A[[Z]]$ where A is an integral domain and let ν be a valuation on A . We define

$$\nu(P(Z)) := \min_a \nu(a)$$

where a runs over all the coefficients of $P(Z)$.

The following proposition is the analogue of Proposition 2.6 of [To]:

Proposition 4.14. *Let $P(Z) \in V_\nu^{\text{fg}}[[Z]]$ be a monic polynomial with no multiple factor. Let us write $P(Z) = P_1(Z) \dots P_r(Z)$ where $P_i(Z) \in V_\nu^{\text{fg}}[[Z]]$, $1 \leq i \leq r$, are irreducible monic polynomials. Let $Q(Z) \in V_\nu^{\text{fg}}[[Z]]$ be a monic polynomial and let z_1, \dots, z_d be the roots of $P(Z)$. If*

$$\deg(Q(Z)) = \deg(P(Z))$$

and

$$\nu(Q(Z) - P(Z)) > d \max_{i \neq j} \{\nu(z_i - z_j)\}$$

then we may factor $Q(Z) = Q_1(Z)\dots Q_r(Z)$ such that $Q_i(Z) \in V_\nu^{\text{fg}}[Z]$ is an irreducible monic polynomial, $1 \leq i \leq r$, and

$$\nu(Q_i(Z) - P_i(Z)) \geq \frac{\nu(Q(Z) - P(Z))}{d}.$$

The result is still valid if we replace V_ν^{fg} by V_ν^{alg} or \widehat{V}_ν .

Proof. Since $P(Z)$ has no multiple factor and since $\text{char}(\mathbb{k}) = 0$, we have $z_i \neq z_j$ for all $i \neq j$. Let us set $r := \max_{i \neq j} \{\nu(z_i - z_j)\}$. Let z'_i , $1 \leq i \leq d$, be the roots of $Q(Z)$. Let z be a root of $P(Z)$ in $V_\nu^{\text{fg}}[\langle \gamma_1, \dots, \gamma_N \rangle]$. Let us write $P(Z) = Z^d + a_1 Z^{d-1} + \dots + a_d$ and $Q(Z) = Z^d + b_1 Z^{d-1} + \dots + b_d$. Then

$$\prod_{1 \leq i \leq d} (z - z'_i) = Q(z) = Q(z) - P(z) = \sum_{i=1}^d (b_i - a_i) z^{d-i}.$$

Thus there exists at least one i such that

$$\nu(z'_i - z) \geq \frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d} = \frac{\nu(Q(Z) - P(Z))}{d} > r.$$

Let t be another root of $P(Z)$. Then

$$\nu(z'_i - t) = \nu(z'_i - z + z - t) = \nu(z - t) \leq r$$

since $\nu(z'_i - z) \geq \frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d} > r \geq \nu(z - t)$. Thus for any root of $P(Z)$ denoted by z , there is only one i such that

$$\nu(z - z'_i) \geq \frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d}.$$

Let $\sigma_1(z), \dots, \sigma_e(z)$ be the conjugates of z over $\mathbb{K}_\nu^{\text{fg}}$. Set

$$R(Z) := (Z - z) \prod_{j=1}^e (Z - \sigma_j(z)) \in V_\nu^{\text{fg}}[Z].$$

Then $R(Z)$ is an irreducible factor of $P(Z)$. Moreover $\sigma_1(z'_i), \dots, \sigma_e(z'_i)$ are conjugates of z'_i over $\mathbb{K}_\nu^{\text{fg}}$. Let σ be a $\mathbb{K}_\nu^{\text{fg}}$ -automorphism of $\overline{\mathbb{K}_\nu^{\text{fg}}}$. Then $\sigma(z)$ is a conjugate of z thus there exists j such that $\sigma(z) = \sigma_j(z)$. Moreover $\sigma(z)$ is a root of $P(Z)$ and $\nu(\sigma(z'_i) - \sigma(z)) \geq \frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d}$ by Lemma 4.8. Thus we have

$$\begin{aligned} \nu(\sigma(z'_i) - \sigma_j(z)) &= \nu(\sigma(z'_i) - \sigma(z)) = \nu(z'_i - z) = \\ &= \nu(\sigma_j(z'_i) - \sigma_j(z)) \geq \frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d} \end{aligned}$$

and since there is only one root of $Q(Z)$ whose difference with $\sigma_j(z)$ has valuation greater than $\frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d}$, we necessarily have $\sigma(z'_i) = \sigma_j(z'_i)$. Thus $\sigma_1(z'_i), \dots, \sigma_e(z'_i)$ are the conjugates of z'_i over $\mathbb{K}_\nu^{\text{fg}}$. Thus the polynomial

$$S(Z) := (Z - z'_i) \prod_{j=1}^e (Z - \sigma_j(z'_i))$$

is irreducible in $V_\nu^{\text{fg}}[Z]$ and

$$\nu(S(Z) - R(Z)) \geq \frac{\min_{1 \leq i \leq d} \{\nu(a_i - b_i)\}}{d}.$$

The proof for \widehat{V}_ν is the same and the case V_ν^{alg} is proven with the help of Lemma 4.7. \square

Remark 4.15. Let us remark that $\nu(Q(Z) - P(Z)) > \frac{d}{2}\nu(\Delta_P)$, where Δ_P is the discriminant of $P(Z)$, implies that

$$\nu(Q(Z) - P(Z)) > d \max_{i \neq j} \{\nu(z_i - z_j)\}.$$

Remark 4.16. This result is not true if $P(Z)$ has multiple factors. For example, let ν be a divisorial valuation and let us consider $P(Z) = Z^2$ and let $Q(Z) = X^2 + a$ where $\nu(a) = 2k + 1$ and $k \in \mathbb{N}$. Since $\nu(a)$ is odd and since the value group of ν is \mathbb{Z} , then it is not a square in \widehat{V}_ν and $Q(Z)$ is irreducible but $P(Z)$ is not irreducible.

Proposition 4.17. *Let ν be an Abhyankar valuation and let*

$$N := \dim_{\mathbb{Q}} \Gamma \otimes_{\mathbb{Z}} \mathbb{Q}.$$

Let $P(Z) \in \widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_N \rangle][Z]$ be a monic polynomial where $\gamma_1, \dots, \gamma_N$ are homogeneous elements with respect to ν . Then there exist integral homogeneous elements with respect to ν , denoted by $\gamma'_1, \dots, \gamma'_N$, and $c \in \mathbb{R}_{>0}$ such that the roots of $P(Z)$ are in $\widehat{V}_\nu[\langle \gamma'_1, \dots, \gamma'_N \rangle]$ and for any monic polynomial $Q(Z) \in \widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_N \rangle][Z]$ such that $\deg(Q(Z)) = \deg(P(Z))$ and $\nu(P(Z) - Q(Z)) \geq c$, the roots of $Q(Z)$ are in $\widehat{V}_\nu[\langle \gamma'_1, \dots, \gamma'_N \rangle]$.

Proof. The proof of this proposition is based on the proof of Theorem 4.2. So let us use the notations of that proof. Let us write $Q(Z) = Z^d + b_1 Z^{d-1} + \dots + b_d$ and let us define $R(Z) := Z^d + b'_1 Z^{d-1} + \dots + b'_d$ where $b'_i := \frac{b_i}{\gamma^i}$ for $1 \leq i \leq d$. We have $Q(\gamma Z) = \gamma^d R(Z)$. Let us assume that $\nu(b'_i - a'_i) > 0$ for all $1 \leq i \leq d$ (i.e. if $\nu(b_i - a_i) > \nu(\gamma^i)$ for all i , thus we assume here that $c > d\nu(\gamma)$). Then $\overline{R}(Z) = \overline{S}(Z)$ ($\overline{R}(Z)$ denotes the image of $R(Z)$ in $\mathbb{L}[Z]$) and the factorization $\overline{R}(Z) = \overline{S}_1(Z)\overline{S}_2(Z)$ lifts to a factorization $R(Z) = R_1(Z)R_2(Z)$ of $R(Z)$ as the product of two monic polynomials as in the proof of Theorem 4.2.

Lemma 4.18. *In the previous situation there exist two constants $a > 0$, $b \geq 0$ depending only on $S_1(Z)$ and $S_2(Z)$ such that for any $c > \max\{b, \nu(\gamma^d)\}$, we have $\nu(R_i(Z) - S_i(Z)) > \frac{c-b}{a}$ for $i = 1, 2$.*

Proof of Lemma 4.18. Let us denote by $r_{i,k}$ the coefficient of Z^k of the polynomial $R_i(Z)$, for $i = 1, 2$ and $0 \leq k \leq \deg_Z(R_i(Z))$, and let us denote by r the vector whose coordinates are the $r_{i,k}$. The coefficient of Z^k of $R_1(Z)R_2(Z) - S_1(Z)S_2(Z)$, for $0 \leq k \leq d$, is a polynomial $f_k(r)$ whose coefficients are in $\widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_N, \gamma, \gamma' \rangle]$ and depend themselves on the coefficients of $S(Z)$. By Theorem 1.2 [M-B], there exist $a > 0$, $b \geq 0$ such that

$$\begin{aligned} \forall c > b, \forall r \in \widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_N, \gamma, \gamma' \rangle]^{d+2} \text{ such that } \nu(f_k(r)) \geq c \quad \forall k \\ \exists r' \in \widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_N, \gamma, \gamma' \rangle]^{d+2} \text{ such that } f_k(r') = 0 \quad \forall k \\ \text{and } \nu(r'_{i,j} - r_{i,j}) \geq \frac{c-b}{a} \quad \forall i, j. \end{aligned}$$

Let us denote by $R'_i(Z)$ the polynomial whose coefficients are the $r'_{i,j}$ where $0 \leq j \leq \deg(R_i)$. Then $R'_1(Z)R'_2(Z) = S_1(Z)S_2(Z)$. Moreover $\overline{R}'_i(Z) = \overline{R}_i(Z) = \overline{S}_i(Z)$ if $\frac{c-b}{a} > 0$. Since the roots of $\overline{S}_1(Z)$ and $\overline{S}_2(Z)$ are different, and since $\widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_N, \gamma, \gamma' \rangle][Z]$ is a GCD domain, then $R'_i(Z) = S_i(Z)$ for $i = 1, 2$. This proves the lemma. \square

Here we remark that, the constants a , b , $\nu(\gamma)$ depend only on $P(Z)$. Thus the result is proven by induction on the degree of $P(Z)$ (since $\deg(S_i(Z)) < \deg(P(Z))$ for $i = 1, 2$) and using Proposition 3.28 and Remark 3.20. \square

5. MONOMIAL VALUATION CASE: EISENSTEIN THEOREM

We will first construct a subring of V_ν^{fg} containing V_ν^{alg} when ν is a monomial valuation.

Definition 5.1. Let $\alpha \in \mathbb{R}_{>0}^n$ and let δ be a (α) -homogeneous polynomial of degree d . We define

$$\mathcal{V}_{\alpha,\delta} := \left\{ A \in \widehat{V}_{\nu_\alpha} / \exists \Lambda \text{ a finitely generated sub-semigroup of } \mathbb{R}_{\geq 0}, \right. \\ \left. \begin{aligned} &\forall i \in \Lambda \exists a_i \in \mathbb{k}[\mathbf{x}] \text{ } (\alpha)\text{-homogeneous,} \\ &\exists a \geq 0, b \in \mathbb{R} \forall i \in \Lambda \exists m(i) \in \mathbb{N} \text{ s.t. } m(i) \leq ai + b, \\ &\nu_\alpha \left(\frac{a_i}{\delta^{m(i)}} \right) = i \text{ and } A = \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}} \end{aligned} \right\}.$$

With this notation we say that $i \mapsto ai + b$ is a *bounding function* for $\sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}}$.

By Lemma 3.12 we have $\mathbb{k}[\mathbf{x}] \subset \mathcal{V}_{\alpha,\delta} \subset V_{\nu_\alpha}^{\text{fg}}$, by identifying a formal power series $\sum_{\beta \in \mathbb{Z}_{\geq 0}^n} c_\beta x^\beta$

to $\sum_{i \in \Lambda} \frac{a_i(x)}{\delta(x)^{m(i)}}$ with $a_i(x) := \sum_{\alpha_1 \beta_1 + \dots + \alpha_n \beta_n = i} c_\beta x^\beta$ et $m(i) = 0$ for all $i \in \Lambda$. We extend in an obvious way the addition and multiplication of $\mathbb{k}[[x]]$ to $\mathcal{V}_{\alpha,\delta}$: this defines a \mathbb{k} -algebra structure over $\mathcal{V}_{\alpha,\delta}$. We have easily the following lemma:

Lemma 5.2. *If $i \mapsto ai + b$ is a bounding function of A and $B \in \mathcal{V}_{\alpha,\delta}$ then it is also a bounding function of $A + B$ and the function $i \mapsto ai + 2b$ is a bounding function of AB .*

Proof. Let us write

$$A = \sum_{i \in \Lambda} \frac{a_i}{\delta^{ai+b}}, \quad B = \sum_{i \in \Lambda} \frac{b_i}{\delta^{ai+b}}$$

where Λ is a semigroup and the a_i and b_i are (α) -homogeneous polynomials and

$$\nu_\alpha \left(\frac{a_i}{\delta^{ai+b}} \right) = \nu_\alpha \left(\frac{b_i}{\delta^{ai+b}} \right) = i \quad \forall i \in \Lambda.$$

Then we have

$$A + B = \sum_{i \in \Lambda} \frac{a_i + b_i}{\delta^{ai+b}}$$

$$\text{and } AB = \sum_{i \in \Lambda} \sum_{j \in \Lambda, j \leq i} \frac{a_j b_{i-j}}{\delta^{aj+b} \delta^{a(i-j)+b}} = \sum_{i \in \Lambda} \sum_{j \in \Lambda, j \leq i} \frac{a_j b_{i-j}}{\delta^{ai+2b}}.$$

This proves the lemma. \square

Remark 5.3. If $A \in \mathcal{V}_{\alpha,\delta}$ satisfies $\nu_\alpha(A) > 0$ then A admits a bounding function which is linear. Indeed let $i \mapsto ai + b$ be a bounding function of A and let $i_0 := \nu_\alpha(A)$. Then $i \mapsto \left(a + \frac{b}{i_0}\right) i$ is a bounding function of A .

Definition 5.4. Let $A := \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}} \in \mathcal{V}_{\alpha,\delta}$, $A \neq 0$. Let i_0 be the least element of Λ such that $a_{i_0} \neq 0$. We say that $\frac{a_{i_0}}{\delta^{m(i_0)}}$ is the *initial term* of A with respect to ν_α or its (α) -*initial term*. We denote it by $\text{in}_\alpha(A)$.

Lemma 5.5. *Let δ and δ' be two (α) -homogeneous polynomials. We have the following properties:*

- i) The (α) -homogeneous irreducible divisors of δ divide δ' if and only if $\mathcal{V}_{\alpha,\delta'} \subset \mathcal{V}_{\alpha,\delta}$. We denote by \mathcal{V}_α the inductive limit of the $\mathcal{V}_{\alpha,\delta}$.
- ii) The valuation ν_α is well defined on $\mathcal{V}_{\alpha,\delta}$ and extends to \mathcal{V}_α . Its valuation ring is exactly \mathcal{V}_α .

Proof. It is clear that if the irreducible divisors of δ divide δ' then $\mathcal{V}_{\alpha,\delta} \subset \mathcal{V}_{\alpha,\delta'}$. On the other hand if $\mathcal{V}_{\alpha,\delta} \subset \mathcal{V}_{\alpha,\delta'}$, then $\frac{1}{\delta} \in \mathcal{V}_{\alpha,\delta'}$, thus there exist a (α) -homogeneous polynomial $a \in \mathbb{k}[x]$ and an integer $m \in \mathbb{N}$ such that $\frac{1}{\delta} = \frac{a}{\delta'^m}$, hence $a\delta = \delta'^m$. This proves i).

If $A \in \mathcal{V}_{\alpha,\delta}$ and $B \in \mathcal{V}_{\alpha,\delta'}$ satisfy $\nu_\alpha(B) \geq \nu_\alpha(A)$, let $\frac{a_k(x)}{\delta(x)^{m(k)}}$ denote the first nonzero term in the expansion of A . Then we can check easily that $\frac{B}{A} \in \mathcal{V}_{\alpha,\delta\delta'^{a_k}}$. This proves ii). \square

Definition 5.6. For any $\alpha \in \mathbb{R}_{>0}^n$ we denote by \mathcal{K}_α the fraction field of \mathcal{V}_α and

$$\overline{\mathcal{K}}_\alpha := \lim_{\substack{\longrightarrow \\ \gamma_1, \dots, \gamma_s}} \mathcal{K}_\alpha[\gamma_1, \dots, \gamma_s]$$

the limit being taken over all subsets $\{\gamma_1, \dots, \gamma_s\}$ of (integral) homogeneous elements with respect to ν .

If $\gamma_1, \dots, \gamma_s$ are homogeneous elements with respect to ν_α we denote by $\mathcal{V}_{\alpha,\delta}[\langle \gamma_1, \dots, \gamma_s \rangle]$ the ring of elements $\sum_{\underline{k}} A_{\underline{k}} \gamma^{\underline{k}}$ where the sum is finite, $\underline{k} := (k_1, \dots, k_s)$, $A_{\underline{k}} = \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}}$ where $a_i \in \mathbb{k}[x]$ is (α) -homogeneous, there exist two constants $a \geq 0$, $b \in \mathbb{R}$ such that $m(i) \leq ai + b$ for all i and there exists $i_0 \in \Lambda$ such that $\nu_\alpha\left(\frac{a_i}{\delta^{m(i)}}\right) = i - i_0$ and $\nu(\gamma^{\underline{k}}) \geq i_0$.

This means that $\mathcal{V}_{\alpha,\delta}[\langle \gamma_1, \dots, \gamma_s \rangle]$ is the subring of $\mathcal{K}_\alpha[\gamma_1, \dots, \gamma_s]$ whose elements have non negative valuation ν_α . In the same way we denote by $\mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_s \rangle]$ the ring of elements of $\mathcal{K}_\alpha[\gamma_1, \dots, \gamma_s]$ having a non negative valuation ν_α . The field of fractions of $\mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_s \rangle]$ is exactly $\mathcal{K}_\alpha[\gamma_1, \dots, \gamma_s]$.

Remark 5.7. We will see later (see Remark 6.10) that these fields $\overline{\mathcal{K}}_\alpha$ coincide with those introduced in [AI] when $\dim_{\mathbb{Q}}(\alpha_1\mathbb{Q} + \dots + \alpha_n\mathbb{Q}) = n$ where it is proven that they are algebraically closed.

Remark 5.8. For any $\alpha \in \mathbb{R}_{>0}^n$ it is clear that $\mathcal{V}_\alpha \subset V_{\nu_\alpha}^{\text{fg}}$ but both rings are never equal if $\dim_{\mathbb{Q}}(\alpha_1\mathbb{Q}_1 + \dots + \alpha_n\mathbb{Q}) < n$. For instance, let $n = 2$ and $\alpha = (1, 1)$ and set

$$z = \sum_{i \in \mathbb{N}} \frac{x_1^{(i+1)^2}}{x_2^{i^2}} \text{ or } \sum_{i \in \mathbb{N}} \frac{x_1^i}{x_1 + ix_2}.$$

Then obviously $z \in V_{\nu_\alpha}^{\text{fg}}$ but $z \notin \mathcal{V}_\alpha$.

Proposition 5.9. If $\alpha_1, \dots, \alpha_n$ are linearly independent over \mathbb{Q} then $\mathcal{V}_\alpha = V_{\nu_\alpha}^{\text{fg}}$.

Proof. Let us denote by $\alpha^* : \mathbb{Q}^n \rightarrow \mathbb{R}$ the \mathbb{Q} -linear map defined by $\alpha^*(u) = \langle \alpha, u \rangle$ for any $u \in \mathbb{Q}^n$. Since the α_i are \mathbb{Q} -linearly independent then α^* is injective.

If Λ is a finitely generated sub-semigroup of $\mathbb{Z}\alpha_1 + \dots + \mathbb{Z}\alpha_n$ let β_1, \dots, β_s be generators of Λ . Then $\alpha^{*-1}(\Lambda)$ is a finitely generated semigroup whose generators are $b_1 = \alpha^{*-1}(\beta_1), \dots, b_s = \alpha^{*-1}(\beta_s) \in \mathbb{Z}^n$.

If the support of $z \in V_{\nu_\alpha}^{\text{fg}}$ is in Λ , since α^* is injective z can be written as

$$z = \sum_{\underline{k} \in \mathbb{Z}_{\geq 0}^s} a_{\underline{k}} \mathbf{x}^{k_1 b_1 + \dots + k_s b_s}$$

where $a_{\underline{k}} \in \mathbb{k}$ for all $\underline{k} = (k_1, \dots, k_s)$. Let us remark that the monomial $a_{\underline{k}} \mathbf{x}^{k_1 b_1 + \dots + k_s b_s}$ is (α) -homogeneous of degree $k_1 \beta_1 + \dots + k_s \beta_s$.

Let us write $b_i = b_{1,i} - b_{2,i}$ where $b_{1,i}, b_{2,i} \in \mathbb{Z}_{\geq 0}^n$. Then we have

$$\begin{aligned} \mathbf{x}^{k_1 b_1 + \dots + k_s b_s} &= \frac{\mathbf{x}^{k_1 b_{1,1} + \dots + k_s b_{1,s}}}{\mathbf{x}^{k_1 b_{2,1} + \dots + k_s b_{2,s}}} \\ &= \frac{\mathbf{x}^{k_1 b_{1,1} + \dots + k_s b_{1,s} + (\max_i \{k_i\} - k_1) b_{2,1} + \dots + (\max_i \{k_i\} - k_s) b_{2,s}}}{\mathbf{x}^{(b_{2,1} + \dots + b_{2,s}) \max_i \{k_i\}}}. \end{aligned}$$

Moreover

$$\max_i \{k_i\} \leq \max_j \left\{ \frac{1}{\beta_j} \right\} (k_1 \beta_1 + \dots + k_s \beta_s).$$

This shows that $z \in \mathcal{V}_{\alpha, \mathbf{x}^{b_{2,1} + \dots + b_{2,s}}}$ and

$$i \mapsto \max_j \left\{ \frac{1}{\beta_j} \right\} i$$

is a bounding function of z . □

Then we give the following version of the Implicit Function Theorem inspired by Lemma 1.2 [Ga] (see also Lemma 2.2. [To]):

Proposition 5.10. *Let $\alpha \in \mathbb{R}_{>0}^n$ and let $P(Z) \in \mathcal{V}_{\alpha, \delta}[\langle \gamma_1, \dots, \gamma_s \rangle][Z]$, $P(Z) = \sum_{k=0}^d a_k Z^k$, where γ_i is homogeneous for all i with respect to ν_α and $d \geq 2$.*

Let $u \in \mathcal{V}_{\alpha, \delta}[\langle \gamma_1, \dots, \gamma_s \rangle]$ such that $\nu_\alpha(P(u)) > 2\nu_\alpha(P'(u))$. Let $\frac{\tilde{\delta}}{\delta^m}$ denote the initial term of $P'(u)$ with respect to ν_α .

Then there exists a unique solution \bar{u} in $\mathcal{V}_{\alpha, \delta \tilde{\delta}}[\langle \gamma_1, \dots, \gamma_s \rangle]$ of $P(Z) = 0$ such that

$$\nu_\alpha(\bar{u} - u) \geq \nu_\alpha(P(u)) - \nu_\alpha(P'(u)).$$

Proof. • By replacing $P(Z)$ by $P(u + Z)$ we can assume that $u = 0$. In this case we have that $P(u) = P(0) = a_0$ and $P'(u) = P'(0) = a_1$.

The valuation ν_α is defined on the ring $\mathcal{V}_{\alpha, \delta}[\langle \gamma_1, \dots, \gamma_s \rangle]$ and we denote by V its valuation ring. We denote by \widehat{V} the completion of V . Let V^{fg} be the subring of \widehat{V} of all elements of \widehat{V} whose ν_α -support is included in a finitely generated semigroup. Then V^{fg} is a Henselian local ring by Lemma 4.1. We set $Z = \frac{\tilde{\delta}}{\delta^m} Y$. Thus we are looking for solving the following equation:

$$\tilde{P}(Y) := \frac{\delta^{2m}}{\tilde{\delta}^2} P \left(\frac{\tilde{\delta}}{\delta^m} Y \right) = a_0 \frac{\delta^{2m}}{\tilde{\delta}^2} + a_1 \frac{\delta^m}{\tilde{\delta}} Y + a_2 Y^2 + \dots + a_d \frac{\tilde{\delta}^{d-2}}{\delta^{(d-2)m}} Y^d = 0.$$

From now on we denote by \tilde{a}_k the coefficients of $\tilde{P}(Y)$:

$$\tilde{a}_k := a_k \frac{\tilde{\delta}^{k-2}}{\delta^{(k-2)m}} \quad k = 0, \dots, d.$$

Since

$$\nu_\alpha(a_0) = \nu_\alpha(P(0)) > 2\nu_\alpha(P'(0)) = 2\nu_\alpha(a_1) = \nu_\alpha \left(\frac{\tilde{\delta}^2}{\delta^{2m}} \right)$$

we have that $\tilde{a}_0 \in V^{\text{fg}}$. By assumption we have that $\nu_\alpha(\tilde{a}_1) = 0$ thus $\tilde{a}_1 \in V^{\text{fg}}$. Since $\nu_\alpha \left(\frac{\tilde{\delta}}{\delta^m} \right) \geq 0$ we have that $\tilde{a}_k \in V^{\text{fg}}$ for all $k \geq 2$. Moreover we have

$$\nu_\alpha(\tilde{P}(0)) > 0 \text{ and } \nu_\alpha(\tilde{P}'(0)) = 0.$$

Thus by Hensel Lemma this equation has a unique solution $y \in V^{\text{fg}}$ such that

$$\nu_\alpha(y) = \nu_\alpha \left(a_0 \frac{\delta^{2m}}{\tilde{\delta}^2} \right) = \nu_\alpha(P(0)) - 2\nu_\alpha(P'(0)) > 0.$$

Hence there exists a unique solution $z := \frac{\tilde{\delta}}{\delta^m} y \in V^{\text{fg}}$ of the equation $P(Z) = 0$ such that

$$\nu_\alpha(z) \geq \nu_\alpha(P(u)) - \nu_\alpha(P'(u)).$$

Now we have to show that z or $y \in \mathcal{V}_{\alpha, \delta \tilde{\delta}}[\langle \gamma_1, \dots, \gamma_s \rangle]$.

- We can write $y = \tilde{a}_0 \tilde{y}$ where $\tilde{y} \in V^{\text{fg}}$ and $\nu_\alpha(\tilde{y}) = 0$. Then \tilde{y} is a root of the polynomial

$$\begin{aligned} \tilde{P}(\tilde{a}_0 Y) &= \tilde{a}_0 + \tilde{a}_1 \tilde{a}_0 Y + \tilde{a}_2 \tilde{a}_0^2 Y^2 + \dots + \tilde{a}_d \tilde{a}_0^d Y^d = \\ &= \tilde{a}_0 (1 + \tilde{a}_1 Y + \tilde{a}_2 \tilde{a}_0 Y^2 + \dots + \tilde{a}_d \tilde{a}_0^{d-1} Y^d) \end{aligned}$$

and $y \in \mathcal{V}_{\alpha, \delta \tilde{\delta}}[\langle \gamma_1, \dots, \gamma_s \rangle]$ if and only if $\tilde{y} \in \mathcal{V}_{\alpha, \delta \tilde{\delta}}[\langle \gamma_1, \dots, \gamma_s \rangle]$.

Since $\nu_\alpha(\tilde{a}_0) > 0$, by replacing \tilde{P} (resp. y) by $1 + \tilde{a}_1 Y + \tilde{a}_2 \tilde{a}_0 Y^2 + \dots + \tilde{a}_d \tilde{a}_0^{d-1} Y^d$ (resp. \tilde{y}), we may assume that

$$\nu_\alpha(\tilde{a}_i) > 0 \text{ for } i \geq 2.$$

In this case we have $\tilde{a}_0 = 1$, $\text{in}_\alpha(\tilde{a}_1) = 1$ and $\text{in}_\alpha(y) = -1$.

Let Λ be a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ containing the ν_α -supports of y and the \tilde{a}_k . We denote by λ_l , $l \in \mathbb{Z}_{\geq 0}$, its elements ordered as follows:

$$\lambda_0 := 0 < \lambda_1 < \lambda_2 < \dots < \lambda_l < \lambda_{l+1} < \dots.$$

Let us expand the coefficients of $\tilde{P}(Y)$ as

$$\tilde{a}_k = \sum_{l \in \mathbb{Z}_{\geq 0}} \tilde{a}_{k, \lambda_l}$$

where \tilde{a}_{k, λ_l} is homogeneous of degree λ_l with respect to ν_α . For every $l \in \mathbb{N}$ let Y_{λ_l} be a new variable and set $Y^* := \sum_{l \in \mathbb{N}} Y_{\lambda_l}$. We extend the valuation ν_α to $V^{\text{fg}}[Y_{\lambda_1}, \dots, Y_{\lambda_l}, \dots]$ by setting $\nu_\alpha(Y_{\lambda_l}) := \lambda_l$ for any $l \in \mathbb{N}$. We may write formally $\tilde{P}(Y^*) = \sum_l \tilde{P}_{\lambda_l}(Y^*)$ where $\tilde{P}_{\lambda_l}(Y^*) \in \mathbb{Z}[\tilde{a}_{k, \lambda_i}][Y_{\lambda_j}]$ is the homogeneous term of degree λ_l with respect to ν_α .

Since $\text{in}_\alpha(\tilde{a}_1) = 1$ the equation

$$(5) \quad \tilde{P}(Y) = \tilde{a}_0 + \tilde{a}_1 Y + \tilde{a}_2 Y^2 + \dots + \tilde{a}_d Y^d = 0,$$

where Y is replaced by Y^* , yields the following equation, for every $l \in \mathbb{Z}_{\geq 0}$:

$$(6) \quad \tilde{P}_{\lambda_l}(Y^*) = Y_{\lambda_l} + Q_{\lambda_l}(Y^*) = 0,$$

where $Q_{\lambda_l}(Y^*) \in \mathbb{Z}[\tilde{a}_{k, \lambda_i}][Y_{\lambda_j}]$ is a polynomial depending only on the variables \tilde{a}_{k, λ_i} ($\lambda_i \leq \lambda_l$) and Y_{λ_j} ($j < l$). Since y is a solution of Equation (5), by replacing Y^* by y we have $\tilde{P}_{\lambda_l}(y) = 0$, hence

$$y_{\lambda_l} = -Q_{\lambda_l}(y_{\lambda_j}, j < l) \quad \forall l \in \mathbb{N}.$$

So by induction on l we see that we may write

$$y_{\lambda_l} = \frac{c_l}{(\delta \tilde{\delta})^{m(\lambda_l)}}$$

for some $c_l \in \mathbb{k}[x][\gamma_1, \dots, \gamma_s]$ and $m(\lambda_l) \in \mathbb{N}$ for all l .

Let $i \mapsto ai + b$ be a common bounding functions of the coefficients of $\tilde{a}_0, \tilde{a}_1, \tilde{a}_2, \dots, \tilde{a}_d$ seen as elements of $\mathcal{V}_{\alpha, \delta \tilde{\delta}}[\langle \gamma_1, \dots, \gamma_s \rangle]$. By Remark 5.3 we may assume that $b = 0$ since $\nu_\alpha(\tilde{a}_k) > 0$ for

$k \geq 2$ and $\text{in}_\alpha(\tilde{a}_0) = \text{in}_\alpha(\tilde{a}_1) = 1$.

Thus we have

$$(\delta\tilde{\delta})^{a\lambda_i}\tilde{a}_{k,\lambda_i} \in \mathbb{k}[x][\gamma_1, \dots, \gamma_s] \quad \forall i.$$

Let $m(\lambda_l)$ be the least integer such that $(\delta\tilde{\delta})^{m(\lambda_l)}y_{\lambda_l} \in \mathbb{k}[x][\gamma_1, \dots, \gamma_s]$. We will show by induction on l that

$$(7) \quad m(\lambda_l) \leq a\lambda_l.$$

This inequality is satisfied for $l = 0$ since $\text{in}_\alpha(y) = -1$ implies that $m(\lambda_0) = 0$.

We fix an integer $l > 0$ and we assume that (7) is satisfied for any integer less than l .

Let Q be a monomial of $Q_i(Y^*)$. We may write

$$Q = \tilde{a}_{k,\lambda_i}y_{\lambda_{j_1}} \cdots y_{\lambda_{j_k}}$$

where $k \leq d$, $j_1 \leq \cdots \leq j_k < l$ and $\lambda_i + \lambda_{j_1} + \cdots + \lambda_{j_k} = \lambda_l$.

Then

$$(\delta\tilde{\delta})^{a\lambda_i + a(\lambda_{j_1} + \cdots + \lambda_{j_k})}Q = (\delta\tilde{\delta})^{a\lambda_l}Q \in \mathbb{k}[[x]][\gamma_1, \dots, \gamma_s].$$

This proves (7). So $y \in \mathcal{V}_{\alpha, \delta\tilde{\delta}}[\langle \gamma_1, \dots, \gamma_s \rangle]$. \square

We deduce from this proposition the main result of this part (Theorem 5.12) which is a general version of Eisenstein Theorem for algebraic power series over \mathbb{Q} . First we recall the classical Eisenstein Theorem:

Theorem 5.11. [Ei] *Let $\sum_{k \in \mathbb{Z}_{\geq 0}} a_k T^k \in \mathbb{Q}[[T]]$ be a power series algebraic over $\mathbb{Q}[[T]]$. Then there exists an integers $a \in \mathbb{N}$ such that*

$$a^{k+1}a_k \in \mathbb{Z}$$

for every integer k .

Theorem 5.12 (Eisenstein Theorem). *Let \mathbb{k} be a field of characteristic zero. Let $\alpha \in \mathbb{R}_{>0}^n$ and let us set $N = \dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n)$. Let*

$$P(Z) \in \mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_s \rangle][Z]$$

be a monic polynomial where $\gamma_1, \dots, \gamma_s$ are homogeneous elements with respect to ν_α . Then there exist integral homogeneous elements with respect to ν_α , denoted by $\gamma'_1, \dots, \gamma'_N$, such that $P(Z)$ has all its roots in $\mathcal{V}_\alpha[\langle \gamma'_1, \dots, \gamma'_N \rangle]$.

Proof. By replacing $P(Z)$ by one of its irreducible factors we may assume that $P(Z)$ is irreducible. Let

$$z \in V_{\nu_\alpha}^{\text{fg}}[\langle \gamma'_1, \dots, \gamma'_N \rangle]$$

be a root of $P(Z)$ where γ'_i is an integral homogeneous with respect to ν_α (by Theorem 4.2 such a z exists). Since $P(Z)$ is irreducible, then $P'(z) \neq 0$. Let us set $i_0 := \max\{\nu_\alpha(z - z')\}$ where the maximum is taken over all the roots z' of P different from z . Let us take $\tilde{z} \in \mathcal{V}_\alpha[\langle \gamma'_1, \dots, \gamma'_N \rangle]$ such that

$$(8) \quad \nu_\alpha(\tilde{z} - z) > \max\{2\nu_\alpha(P'(z)), i_0 + \nu_\alpha(P'(z))\}.$$

For instance if we expand $z = \sum_{i \in \Lambda} z_i$ where Λ is a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ and z_i is a homogeneous element of degree i with respect to ν_α we can choose

$$\tilde{z} := \sum_{i \leq \max\{2\nu_\alpha(P'(z)), i_0 + \nu_\alpha(P'(z))\}} z_i.$$

By replacing $P(Z)$ by $P(Z + \tilde{z})$ we may assume that $\tilde{z} = 0$. In this case $P'(\tilde{z}) = P'(0) = a_{d-1}$ if we write

$$P(Z) = Z^d + a_1 Z^{d-1} + \cdots + a_{d-1} Z + a_d.$$

Now $\text{in}_{\nu_\alpha}(a_{d-1}) \in \mathbb{k}(\mathbf{x})[\gamma'_1, \dots, \gamma'_N]$ so if we denote by a the product of the conjugates of $\text{in}_{\nu_\alpha}(a_{d-1})$ over $\mathbb{k}(\mathbf{x})$ different from $\text{in}_{\nu_\alpha}(a_{d-1})$ we have $\text{in}_{\nu_\alpha}(aa_{d-1}) \in \mathbb{k}(\mathbf{x})$ and a is a homogeneous element with respect to ν_α by Lemma 4.8. Let b be a homogeneous element such that $b^{d-1} = a$. By Proposition 3.28 we may assume that $b \in V_{\nu_\alpha}^{\text{fg}}[\langle \gamma'_1, \dots, \gamma'_N \rangle]$. We have that

$$\begin{aligned} b^d P\left(\frac{Z}{b}\right) &= b^d \left(\frac{1}{b^d} Z^d + \frac{a_1}{b^{d-1}} Z^{d-1} + \cdots + \frac{a_{d-1}}{b} Z + a_d \right) \\ &= Z^d + a_1 b Z^{d-1} + \cdots + b^{d-1} a_{d-1} Z + b^d a_d. \end{aligned}$$

By replacing $P(Z)$ by $b^d P\left(\frac{Z}{b}\right)$ we may assume that $\text{in}_{\nu_\alpha}(P'(\tilde{z})) = \text{in}_{\nu_\alpha}(a_{d-1}) \in \mathbb{k}(\mathbf{x})$.

Since $P(\tilde{z}) - P(z) \in (\tilde{z} - z)$ then by Inequality (8)

$$\nu_\alpha(P(\tilde{z})) > 2\nu_\alpha(P'(z))$$

and

$$\nu_\alpha(P(\tilde{z})) > i_0 + \nu_\alpha(P'(z)).$$

In the same way, since $P'(\tilde{z}) - P'(z) \in (\tilde{z} - z)$, Inequality (8) yields

$$\nu_\alpha(P'(\tilde{z})) = \nu_\alpha(P'(z)).$$

Then we apply Proposition 5.10 (with $u := \tilde{z} = 0$), and we get a root $\bar{z} \in \mathcal{V}_\alpha[\langle \gamma'_1, \dots, \gamma'_N \rangle]$ of $P(Z)$ such that

$$\nu_\alpha(\tilde{z} - \bar{z}) \geq \nu_\alpha(P(\tilde{z})) - \nu_\alpha(P'(\tilde{z})) > i_0.$$

Thus

$$\nu_\alpha(z - \bar{z}) = \nu_\alpha(z - \tilde{z} + \tilde{z} - \bar{z}) > i_0 = \max_{\substack{z' \neq z \\ P(z')=0}} \{\nu_\alpha(z - z')\}.$$

Hence $z = \bar{z} \in \mathcal{V}_\alpha[\langle \gamma'_1, \dots, \gamma'_N \rangle]$. □

Corollary 5.13. *The field $\mathbb{K}_{\nu_\alpha}^{\text{alg}}$ is a subfield of \mathcal{K}_α .*

Proof. Let $z \in \mathbb{K}_{\nu_\alpha}^{\text{alg}}$ and let $P(Z) = a_0 Z^d + \cdots + a_d \in \mathbb{k}[\mathbf{x}][Z]$ be a polynomial such that $P(z) = 0$. Then $a_0 z \in \mathbb{K}_{\nu_\alpha}^{\text{alg}}$ is a root of the polynomial $a_0^{d-1} P(Z/a_0) = Z^d + a_1 Z^{d-1} + a_2 a_0 Z^{d-2} + \cdots + a_d a_0^{d-1}$ which is a monic polynomial. Hence $a_0 z \in \mathcal{V}_\alpha$ by Theorem 5.12 and $z \in \mathcal{K}_\alpha$. □

Example 5.14. Let us assume that $\text{Disc}_Z(P(Z))$ is normal crossing after a formal change of coordinates and let us assume that \mathbb{k} is algebraically closed. This means that there exist power series $x_i(\mathbf{y}) \in (\mathbf{y})\mathbb{k}[[\mathbf{y}]]$ ($\mathbf{y} = (y_1, \dots, y_n)$), for $1 \leq i \leq n$, such that the morphism of \mathbb{k} -algebras $\varphi : \mathbb{k}[[\mathbf{x}]] \rightarrow \mathbb{k}[[\mathbf{y}]]$ defined by $\varphi(f(\mathbf{x})) = f(x_1(\mathbf{y}), \dots, x_n(\mathbf{y}))$ is an isomorphism, and such that

$$\varphi(\text{Disc}_Z(P(Z)))\mathbb{k}[[\mathbf{y}]] = y_1^{e_1} \cdots y_m^{e_m} \mathbb{k}[[\mathbf{y}]], \quad m \leq n.$$

By Abhyankar-Jung Theorem [Ab] (or [KV], [PR], [MS]), the roots of $P(Z)$ can be written as

$$t_k = \sum_{l=0}^d t_{k,l}(\mathbf{y}) \mathbf{w}^l$$

where $\mathbf{w} = \mathbf{y}^\beta$ for some $\beta \in \mathbb{Q}_{\geq 0}^m \times \{0\}^{n-m}$, $d \in \mathbb{Z}_{\geq 0}$ and the $t_{k,l}(\mathbf{y})$ are power series with coefficients in \mathbb{k} . Let us write:

$$\beta = \left(\frac{b_1}{e}, \dots, \frac{b_m}{e}, 0, \dots, 0 \right)$$

for some non negative integers b_1, \dots, b_m and $e \in \mathbb{N}$. Let us denote by $f_i(\mathbf{x})$, $1 \leq i \leq n$, the power series satisfying $\varphi(f_i(\mathbf{x})) = y_i$.

Let $\alpha \in \mathbb{N}^n$ and write $f_i(\mathbf{x}) = l_{i,\alpha}(\mathbf{x}) + \varepsilon_{i,\alpha}(\mathbf{x})$ where $l_{i,\alpha}(\mathbf{x})$ is (α) -homogeneous and $\nu_\alpha(\varepsilon_i(\mathbf{x})) > \nu_\alpha(l_{i,\alpha}(\mathbf{x}))$ for any i . Thus we have for $1 \leq i \leq m$:

$$y_i^{\frac{1}{e}} = l_{i,\alpha}(\mathbf{x})^{\frac{1}{e}} \left(1 + \frac{\varepsilon_{i,\alpha}(\mathbf{x})}{l_{i,\alpha}(\mathbf{x})} \right)^{\frac{1}{e}} = l_{i,\alpha}(\mathbf{x})^{\frac{1}{e}} \left(1 + \sum_{k \geq 1} c_k \frac{\varepsilon_{i,\alpha}(\mathbf{x})^k}{l_{i,\alpha}(\mathbf{x})^k} \right)$$

where $c_k \in \mathbb{Q}$ for all k - here

$$c_k = \frac{\frac{1}{e} \left(\frac{1}{e} - 1 \right) \cdots \left(\frac{1}{e} - k + 1 \right)}{k!}.$$

Hence

$$\begin{aligned} \mathbf{w} = y_1^{\frac{b_1}{e}} \cdots y_m^{\frac{b_m}{e}} = \\ l_{1,\alpha}(\mathbf{x})^{\frac{b_1}{e}} \cdots l_{m,\alpha}(\mathbf{x})^{\frac{b_m}{e}} \prod_{j=1}^m \left(1 + \sum_{k \geq 1} c_k \frac{\varepsilon_{j,\alpha}(\mathbf{x})^k \prod_{p \neq j} l_{p,\alpha}(\mathbf{x})^k}{\left(\prod_{p=1}^m l_p(\mathbf{x}) \right)^k} \right)^{b_j}. \end{aligned}$$

We remark that $\text{Disc}_Z(P(Z)) = \prod_{p=1}^m l_{p,\alpha}(\mathbf{x})^{e_p} + \varepsilon(\mathbf{x})$ with

$$\nu_\alpha(\varepsilon(\mathbf{x})) > \nu_\alpha \left(\prod_{p=1}^m l_{p,\alpha}(\mathbf{x})^{e_p} \right).$$

Let $\gamma := \prod_{j=1}^m l_{j,\alpha}(\mathbf{x})^{\frac{b_j}{e}}$ be a root of the polynomial

$$Z^e - \prod_{j=1}^m l_{j,\alpha}(\mathbf{x})^{b_j}$$

(in particular it is an integral homogeneous element with respect to ν_α), and set $\delta := \prod_{j=1}^m l_{j,\alpha}(\mathbf{x})^{e_p}$. Here δ is the (α) -initial term of the discriminant of $P(Z)$. Hence we obtain the following three cases:

- i) If φ is a linear change of coordinates (i.e. $\alpha = (1, \dots, 1)$ and $\varepsilon_{i,\alpha} = 0 \forall i$), then the roots of $P(Z)$ are in $\mathbb{k}[\mathbf{x}][\gamma]$ (since in this case $\mathbf{w} = \gamma$).
- ii) If φ is a quasi-linear change of variables (i.e. $\alpha \in \mathbb{N}^n$ and $\varepsilon_{i,\alpha} = 0 \forall i$), then the roots of $P(Z)$ are still in $\mathbb{k}[\mathbf{x}][\gamma]$ (since in this case we also have $\mathbf{w} = \gamma$).
- iii) If (at least) one of the $\varepsilon_{i,\alpha}$ is not zero, then the roots of $P(Z)$ are in $\mathcal{V}_{\alpha,\delta}[\langle \gamma \rangle]$.

This example will be generalized later (see Theorem 7.7).

Example 5.15. Let $P(Z) = Z^2 + 2aZ + b$ where a and b are power series over \mathbb{k} and let $\alpha \in \mathbb{Q}_{>0}^2$. Let δ denote the (α) -initial term of the discriminant of $P(Z)$, i.e. the (α) -initial term of $a^2 - b$. Then the roots of $P(Z)$ are of the form $-a + \sqrt{a^2 - b} \in \mathcal{V}_{\alpha,\delta}[\langle \gamma \rangle]$ where γ is a root square of δ .

Example 5.16. Let $P(Z) = Z^3 + 3x_2^2 Z - 2(x_1^3 + \varepsilon)$ where ε is a homogeneous polynomial of degree greater or equal to 4. Its discriminant is $D := x_1^6 + x_2^6 + 2x_1^3 \varepsilon + \varepsilon^2$ whose initial term is $x_1^6 + x_2^6$. The roots of P are

$$a\sqrt[3]{x_1^3 + \varepsilon + \sqrt{D}} + b\sqrt[3]{x_1^3 + \varepsilon - \sqrt{D}}$$

with $(a, b) = (1, 1)$, (j, j^2) or (j^2, j) . But we have

$$\sqrt[3]{x_1^3 + \varepsilon + \sqrt{D}} = \gamma_1 \sqrt[3]{1 + \varepsilon + \frac{\gamma_2}{x_1^3 + \gamma_2} \sqrt{1 + \frac{2x_1^3\varepsilon + \varepsilon^2}{\delta} - \frac{\gamma_2}{x_1^3 + \gamma_2}}}$$

with $\gamma_2^2 = x_1^6 + x_2^6$, $\gamma_1^3 = x_1^3 + \gamma_2$ and $\delta = x_1^6 + x_2^6$ is the initial term of D . Thus

$$\sqrt[3]{x_1^3 + \varepsilon + \sqrt{D}} \in \mathcal{V}_{(1,1),\delta} \left[\left\langle \gamma_1, \gamma_2, \frac{\gamma_2\varepsilon}{x_1^3 + \gamma_2} \right\rangle \right].$$

By doing the same remark for $\sqrt[3]{x_1^3 + \varepsilon - \sqrt{D}}$, we see that there exist $\gamma_1, \dots, \gamma_5$ homogeneous elements with respect to ord such that the roots of $P(Z)$ are in $\mathcal{V}_{(1,1),\delta}[\langle \gamma_1, \dots, \gamma_5 \rangle]$. But there is no reason that the roots of $P(Z)$ are in $\mathcal{V}_{\alpha,\delta}[\langle \gamma \rangle]$ where γ is one (integral) homogeneous element with respect to ν_α .

6. APPROXIMATION OF MONOMIAL VALUATIONS BY DIVISORIAL MONOMIAL VALUATIONS

In several cases, it will be easier to work with a monomial valuation ν_α which is divisorial, i.e. such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \dots + \mathbb{Q}\alpha_n) = 1$. In order to extend some results which are proven for divisorial monomial valuations to general monomial valuations, we will approximate monomial valuations by divisorial monomial valuations. The aim of this section is to explain how this can be done.

Definition 6.1. Let $\alpha \in \mathbb{R}_{>0}^n$. Let $\alpha^* : \mathbb{Q}^n \rightarrow \mathbb{R}$ be the \mathbb{Q} -linear morphism defined by $\alpha^*(q_1, \dots, q_n) := \sum_i \alpha_i q_i$. We denote by Rel_α the kernel of this morphism. For any $\varepsilon > 0$ and $q \in \mathbb{N}$, we define the following set:

$$\text{Rel}(\alpha, q, \varepsilon) := \left\{ \alpha' \in \mathbb{N}^n / \text{Rel}_\alpha \subset \text{Rel}_{\alpha'} \text{ and } \max_i \left| q - \frac{\alpha'_i}{\alpha_i} \right| < q\varepsilon \right\}.$$

Example 6.2. If $n = 4$, and $\alpha_1 = \sqrt{2}$, $\alpha_2 = \sqrt{3}$, $\alpha_3 = 13\sqrt{2} + \sqrt{3}$, $\alpha_4 = \sqrt{2} + 757\sqrt{3}$, then any α' of the form $(n_1, n_2, 13n_1 + n_2, n_1 + 757n_2)$, where $n_1, n_2 \in \mathbb{N}_{>0}$, will satisfy $\text{Rel}_\alpha \subset \text{Rel}_{\alpha'}$.

Remark 6.3. For α and $\beta \in \mathbb{R}_{>0}^n$ we have

$$\text{Rel}_\alpha \subset \text{Rel}_\beta \iff \beta \in V \otimes_{\mathbb{Q}} \mathbb{R}$$

where $V := (\text{Ker } \alpha^*)^\perp \subset \mathbb{Q}^n$. By definition we have that $\alpha \in V \otimes_{\mathbb{Q}} \mathbb{R}$. Since V is dense in $V \otimes_{\mathbb{Q}} \mathbb{R}$ there exists $\beta \in V$ such that

$$\max_{1 \leq i \leq n} \left| 1 - \frac{\beta_i}{\alpha_i} \right| < \varepsilon.$$

Let us write $\beta_i = \frac{\alpha'_i}{q}$ where the α'_i and q are positive integers. This implies that

$$\max_{1 \leq i \leq n} \left| q - \frac{\alpha'_i}{\alpha_i} \right| < q\varepsilon.$$

Since $\beta \in V$ we have that $\alpha' \in V$ thus $\text{Rel}_\alpha \subset \text{Rel}_{\alpha'}$. This shows that for any given $\alpha \in \mathbb{R}_{>0}^n$ and $\varepsilon > 0$ there always exists $q \in \mathbb{N}$ such that $\text{Rel}(\alpha, q, \varepsilon) \neq \emptyset$.

Moreover if $\alpha \in \mathbb{N}^n$ then $\text{Rel}(\alpha, q, \varepsilon) = \{q\alpha\}$ if $0 < \varepsilon < \frac{1}{q \max\{\alpha_i\}}$. Indeed in this case the only $\alpha' \in \mathbb{N}^n$ satisfying $\max_i |q\alpha_i - \alpha'_i| < q\alpha_i\varepsilon$ is $\alpha' = q\alpha$.

Lemma 6.4. *Let $\alpha, \alpha' \in \mathbb{R}_{>0}^n$. Then $\text{Rel}_\alpha \subset \text{Rel}_{\alpha'}$ if and only if every (α) -homogeneous polynomial is a (α') -homogeneous polynomial.*

Moreover if $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ and if $a(\mathbf{x})$ is a (α) -homogeneous polynomial then

$$q(1 - \varepsilon)\nu_\alpha(a(\mathbf{x})) \leq \nu_{\alpha'}(a(\mathbf{x})) \leq q(1 + \varepsilon)\nu_\alpha(a(\mathbf{x})).$$

Proof. First let us assume that $\text{Rel}_\alpha \subset \text{Rel}_{\alpha'}$ and let $a(\mathbf{x})$ be a (α) -homogeneous polynomial. This means that for any $p, q \in \mathbb{N}^n$, if \mathbf{x}^p and \mathbf{x}^q are two nonzero monomials of $a(\mathbf{x})$, then $\sum_i \alpha_i p_i = \sum_i \alpha_i q_i$. In particular $p - q \in \text{Ker}(\alpha^*)$, thus $\sum_i \alpha'_i p_i = \sum_i \alpha'_i q_i$. Thus $a(\mathbf{x})$ is a (α') -homogeneous.

On the other hand let us assume that every (α) -homogeneous polynomial is a (α') -homogeneous polynomial. Let $r \in \text{Rel}_\alpha$. We can write $r = p - q$ where $p, q \in \mathbb{Q}_{>0}^n$. By multiplying r by a positive integer m , we may assume that $mp, mq \in \mathbb{N}^n$. By assumption on r , the polynomial $\mathbf{x}^{mp} + \mathbf{x}^{mq}$ is (α) -homogeneous. Thus it is (α') -homogeneous. This means that $\sum_i \alpha'_i mp_i = \sum_i \alpha'_i mq_i$. Hence $\sum_i \alpha'_i (p_i - q_i) = 0$ and $r = p - q \in \text{Rel}_{\alpha'}$.

Now let \mathbf{x}^p be a monomial. Then

$$\nu_{\alpha'}(\mathbf{x}^p) = \sum_i \alpha'_i p_i.$$

But $q(1 - \varepsilon)\alpha_i \leq \alpha'_i \leq q(1 + \varepsilon)\alpha_i$ for any $1 \leq i \leq n$. This proves both inequalities. \square

Example 6.5. Let $\alpha \in \mathbb{N}^n$ and $\alpha' \in \mathbb{R}_{>0}^n$. Then $\text{Rel}_\alpha \subset \text{Rel}_{\alpha'}$ if and only if there exists $\lambda \in \mathbb{R}$ such that $\alpha' = \lambda\alpha$. Indeed we have $\dim_{\mathbb{Q}}(\text{Rel}_\alpha) = n - 1$ hence either $\dim_{\mathbb{Q}}(\text{Rel}_{\alpha'}) = n$ and $\alpha' = 0$, either $\dim_{\mathbb{Q}}(\text{Rel}_{\alpha'}) = n - 1$ and there exists $\lambda \in \mathbb{R}^*$ such that $\alpha' = \lambda\alpha$.

Lemma 6.6. *Let $\alpha \in \mathbb{R}_{>0}^n$ and let $A \in \mathcal{V}_\alpha$. Let us write*

$$A = \sum_{i \in \Lambda} \frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}}$$

where Λ is a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ and $i \mapsto m(i)$ is bounded by an affine function. Then there exists $\varepsilon_A > 0$ such that for all $0 < \varepsilon \leq \varepsilon_A$, for all $q \in \mathbb{N}$, for all $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$, the element $\sum_{i \in \Lambda} \frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}}$ is in the fraction field of $\mathcal{V}_{\alpha'}$.

Moreover if $A \in \mathcal{V}_\alpha$ is not invertible, i.e. $\nu_\alpha(A) > 0$, then we may even choose $\varepsilon_A > 0$ such that for all $0 \leq \varepsilon \leq \varepsilon_A$, for all $q \in \mathbb{N}$, for all $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$, $\sum_{i \in \Lambda} \frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}} \in \mathcal{V}_{\alpha'}$ and this element is not invertible in $\mathcal{V}_{\alpha'}$.

Proof. Let $a, b \geq 0$ such that $m(i) \leq ai + b$ for any $i \in \Lambda$. By Lemma 6.4 we have

$$\begin{aligned} \nu_{\alpha'} \left(\frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}} \right) &= \nu_{\alpha'}(a_i(\mathbf{x})) - m(i)\nu_{\alpha'}(\delta(\mathbf{x})) \geq q(1 - \varepsilon)\nu_\alpha(a_i(\mathbf{x})) - q(1 + \varepsilon)m(i)\nu_\alpha(\delta(\mathbf{x})) \\ &= q(1 - \varepsilon)i - 2q\varepsilon m(i)\nu_\alpha(\delta(\mathbf{x})). \end{aligned}$$

Let ε_A be a positive real number such that $\varepsilon_A < \frac{1}{1 + 2a\nu_\alpha(\delta(\mathbf{x}))}$ and set

$$\eta := 1 - \varepsilon_A(1 + 2a\nu_\alpha(\delta(\mathbf{x}))) > 0.$$

Then for any $0 \leq \varepsilon \leq \varepsilon_A$, any $q \in \mathbb{N}$ and any $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ we have

$$\nu_{\alpha'} \left(\frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}} \right) \geq \eta qi - 2qb\varepsilon\nu_\alpha(\delta(\mathbf{x})) \quad \forall i \in \Lambda.$$

This proves that $\sum_{i \in \Lambda} \frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}}$ is in the fraction field of $\mathcal{V}_{\alpha'}$.

If $\nu_{\alpha}(A) > 0$, then $a_0(\mathbf{x}) = 0$. Let $i_0 := \nu_{\alpha}(A)$. Let $\varepsilon \geq 0$ be such that $\varepsilon \leq \varepsilon_A$ and

$$i_0 > \varepsilon((1 + 2a\nu_{\alpha}(\delta(\mathbf{x})))i_0 + 2b\nu_{\alpha}(\delta(\mathbf{x}))).$$

In this case $\nu_{\alpha'}\left(\frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}}\right) > 0$ for any $i \in \Lambda$, $i \geq i_0$. This proves the second assertion. \square

Definition 6.7. Let $\alpha \in \mathbb{R}_{>0}^n$ and $\alpha' \in \text{Rel}_{\alpha} \cap \mathbb{N}^n$. Then every (α) -homogeneous polynomial $p(\mathbf{x})$ is (α') -homogenous by Lemma 6.4. In particular if $\delta(\mathbf{x})$ is an other (α) -homogeneous polynomial and $s \in \mathbb{N}$ then

$$p(x_1\delta(\mathbf{x})^{\alpha'_1 s}, \dots, x_n\delta(\mathbf{x})^{\alpha'_n s}) = p(\mathbf{x})\delta(\mathbf{x})^{s\nu_{\alpha'}(p(\mathbf{x}))}$$

is also a (α) -homogeneous polynomial.

If $A = \sum_{i \in \Lambda} \frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}} \in \mathcal{V}_{\alpha, \delta}$ and $\alpha' \in \text{Rel}_{\alpha} \cap \mathbb{N}^n$, we will set

$$\varphi_{\alpha', s}(A) := \sum_{i \in \Lambda} \frac{a_i}{\delta_i^{m(i)}} (x_1\delta(\mathbf{x})^{\alpha'_1 s}, \dots, x_n\delta(\mathbf{x})^{\alpha'_n s}).$$

Then $\varphi_{\alpha, s} : \mathcal{V}_{\alpha, \delta} \rightarrow \mathcal{V}_{\alpha, \delta}$ is a ring morphism. We also define

$$\psi_{\alpha', s}(A) := \delta^s \varphi_{\alpha', s}(A) \quad \forall A \in \mathcal{V}_{\alpha, \delta}.$$

Lemma 6.8. Let $\alpha \in \mathbb{R}_{>0}^n$ and $A \in \mathcal{V}_{\alpha, \delta}$. For any $\varepsilon > 0$ small enough there exists $s(\varepsilon) \in \mathbb{N}$ such that for every $q \in \mathbb{N}$, $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ and $s \geq s(\varepsilon)$:

$$\psi_{\alpha', s}(A) \in \mathbb{k}[\mathbf{x}].$$

If $\nu_{\alpha}(A) > 0$ we may even assume that $\varphi_{\alpha', s}(A) \in \mathbb{k}[\mathbf{x}]$ for every $q \in \mathbb{N}$, $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ and $s \geq s(\varepsilon)$.

Proof. Let $a(\mathbf{x}), \delta(\mathbf{x}) \in \mathbb{k}[\mathbf{x}]$ be (α) -homogeneous polynomials and let $m \in \mathbb{N}$ such that $\nu_{\alpha}\left(\frac{a(\mathbf{x})}{\delta(\mathbf{x})^m}\right) = i$. Let $s \in \mathbb{N}$ and $a' \in \mathbb{N}^n$ such that $\text{Rel}_{\alpha} \subset \text{Rel}_{\alpha'}$. By Lemma 6.4 we have

$$(9) \quad \frac{a(x_1\delta(\mathbf{x})^{\alpha'_1 s}, \dots, x_n\delta(\mathbf{x})^{\alpha'_n s})}{\delta(x_1\delta(\mathbf{x})^{\alpha'_1 s}, \dots, x_n\delta(\mathbf{x})^{\alpha'_n s})^m} = a(\mathbf{x})\delta(\mathbf{x})^{s[\nu_{\alpha'}(a(\mathbf{x})) - \nu_{\alpha'}(\delta(\mathbf{x}))m] - m}.$$

Now let $A = \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}} \in \mathcal{V}_{\alpha, \delta}$ with $m(i) \leq ai + b$ for any $i \in \Lambda$, Λ being a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$. Set $d_{\alpha} := \nu_{\alpha}(\delta)$. Thus $\nu_{\alpha}(a_i) = d_{\alpha}m(i) + i$ for any $i \in \Lambda$. Hence by Lemma 6.4 we have that

$$\nu_{\alpha'}(a_i) - m(i)\nu_{\alpha'}(\delta) \geq q(1 - \varepsilon)[d_{\alpha}m(i) + i] - q(1 + \varepsilon)m(i)d_{\alpha}$$

$$(10) \quad \nu_{\alpha'}(a_i) - m(i)\nu_{\alpha'}(\delta) \geq q(1 - \varepsilon)i - 2q\varepsilon d_{\alpha}m(i).$$

Since $(1 - \varepsilon)i - 2\varepsilon d_{\alpha}m(i) \geq (1 - \varepsilon)i - 2\varepsilon d_{\alpha}(ai + b)$, for every ε small enough there exists $a_{\varepsilon} > 0$ such that

$$\nu_{\alpha'}(a_i) - m(i)\nu_{\alpha'}(\delta) \geq qa_{\varepsilon}i$$

for all $q \in \mathbb{N}$, all $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ and all $i \in \Lambda$, $i > 0$. Thus for $s \in \mathbb{N}$ and $i \in \Lambda \setminus \{0\}$ we have that

$$s(\nu_{\alpha'}(a_i) - m(i)\nu_{\alpha'}(\delta)) - m(i) \geq sq a_{\varepsilon}i - m(i) \geq (sq a_{\varepsilon} - a)i - b \geq \left(sq a_{\varepsilon} - a - \frac{b}{\min \Lambda \setminus \{0\}} \right) i.$$

In particular if $s \geq \left(a + \frac{b}{\min \Lambda \setminus \{0\}} \right) / a_{\varepsilon}$ then

$$s(\nu_{\alpha'}(a_i) - m(i)\nu_{\alpha'}(\delta)) - m(i) \geq 0$$

and $\frac{a_i}{\delta_i^{m(i)}}(x_1\delta(\mathbf{x})^{\alpha'_1 s}, \dots, x_n\delta(\mathbf{x})^{\alpha'_n s}) \in \mathbb{k}[\mathbf{x}]$ for all $i > 0$. Thus if $\nu_\alpha(A) > 0$, $a_0 = 0$ and

$\varphi_{\alpha', s}(A) \in \mathbb{k}[\mathbf{x}]$ for $s \geq \left(a + \frac{b}{\min \Lambda \setminus \{0\}}\right) / a_\varepsilon$.

In the general case where $a_0 \neq 0$, if we assume moreover that $s \geq b$, we have that

$$\delta(x)^s \frac{a_0}{\delta_0^{m(0)}}(x_1\delta(\mathbf{x})^{\alpha'_1 s}, \dots, x_n\delta(\mathbf{x})^{\alpha'_n s}) \in \mathbb{k}[\mathbf{x}].$$

This proves the lemma. □

When the components of α are \mathbb{Q} -linearly independent, by using Lemma 6.8, Theorem 5.12 gives the following generalization of the main result of [McD]:

Theorem 6.9. [McD] *Let \mathbb{k} be a field of characteristic zero and $\alpha \in \mathbb{R}_{>0}^n$ such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \dots + \mathbb{Q}\alpha_n) = n$. Then*

$$\mathbb{K}_{\nu_\alpha}^{\text{alg}} \subset \bigcup_{\sigma} \mathbb{k}((x^\beta, \beta \in \sigma \cap \mathbb{Z}^n))$$

where the first union runs over all rational strongly convex cones σ such that $\langle \alpha, \tau \rangle > 0$ for any $\tau \in \sigma$, $\tau \neq 0$. Moreover we have:

$$\overline{\mathbb{K}}_{\nu_\alpha}^{\text{alg}} \subset \bigcup_{\sigma} \bigcup_{\mathbb{k}'} \bigcup_{q \in \mathbb{N}} \mathbb{k}' \left(\left(x^\beta, \beta \in \sigma \cap \frac{1}{q} \mathbb{Z}^n \right) \right)$$

where the first union runs over all rational strongly convex cones σ such that $\langle \alpha, \tau \rangle > 0$ for any $\tau \in \sigma$, $\tau \neq 0$, and the second union runs over all the fields \mathbb{k}' finite over \mathbb{k} .

Proof. In order to prove the first inclusion, by Corollary 5.13 it is enough to prove that $\mathcal{K}_\alpha \subset \bigcup_{\sigma} \mathbb{k}((x^\beta, \beta \in \sigma \cap \mathbb{Z}^n))$ or $\mathcal{V}_\alpha \subset \bigcup_{\sigma} \mathbb{k}[[x^\beta, \beta \in \sigma \cap \mathbb{Z}^n]]$.

Since the α_i are \mathbb{Q} -linearly independent the only (α) -homogeneous polynomials are the monomials. Let $\omega \in \mathbb{N}^n$ and A be an element of $\mathcal{V}_{\alpha, \mathbf{x}^\omega} : A = \sum_{i \in \Lambda} \frac{\mathbf{x}^{p(i)}}{\mathbf{x}^{m(i)\omega}}$ where Λ is a finitely generated sub-semigroup of $\mathbb{R}_{>0}$. We have to prove that $A \in \bigcup_{\sigma} \mathbb{k}[[x^\beta, \beta \in \sigma \cap \mathbb{Z}^n]]$. Since $x_1 A \in \bigcup_{\sigma} \mathbb{k}[[x^\beta, \beta \in \sigma \cap \mathbb{Z}^n]]$ implies that $A \in \bigcup_{\sigma} \mathbb{k}[[x^\beta, \beta \in \sigma \cap \mathbb{Z}^n]]$ we may assume that $\nu_\alpha(A) > 0$.

By Lemma 6.8, we see that the monomial map $\varphi_{\alpha', s}$ defined by $x_j \mapsto x_j \mathbf{x}^{s\alpha'_j \omega}$ maps A onto an element of $\mathbb{k}[\mathbf{x}]$ for $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$, $\varepsilon > 0$ small enough and s large enough. Such a monomial map is induced by a linear map on the set of monomials and its matrix is

$$M_1 := \begin{pmatrix} 1 + s\omega_1\alpha'_1 & s\omega_1\alpha'_2 & s\omega_1\alpha'_3 & \cdots & s\omega_1\alpha'_n \\ s\omega_2\alpha'_1 & 1 + s\omega_2\alpha'_2 & s\omega_2\alpha'_3 & \cdots & s\omega_2\alpha'_n \\ s\omega_3\alpha'_1 & s\omega_3\alpha'_2 & 1 + s\omega_3\alpha'_3 & \cdots & s\omega_3\alpha'_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ s\omega_n\alpha'_1 & s\omega_n\alpha'_2 & s\omega_n\alpha'_3 & \cdots & 1 + s\omega_n\alpha'_n \end{pmatrix}$$

Set

$$M_2 := \begin{pmatrix} -s\omega_1\alpha'_1 & -s\omega_1\alpha'_2 & -s\omega_1\alpha'_3 & \cdots & -s\omega_1\alpha'_n \\ -s\omega_2\alpha'_1 & -s\omega_2\alpha'_2 & -s\omega_2\alpha'_3 & \cdots & -s\omega_2\alpha'_n \\ -s\omega_3\alpha'_1 & -s\omega_3\alpha'_2 & -s\omega_3\alpha'_3 & \cdots & -s\omega_3\alpha'_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -s\omega_n\alpha'_1 & -s\omega_n\alpha'_2 & -s\omega_n\alpha'_3 & \cdots & -s\omega_n\alpha'_n \end{pmatrix}$$

and let $\chi(t)$ be the characteristic polynomial of M_2 . Then $\chi(1) = \det(M_1)$. If $\chi(1) = 0$, then the vector $\omega := (\omega_1, \dots, \omega_n)$ is an eigenvector of M_2 with eigenvalue 1 since the image of M_2 is

generated by ω . Thus $-s(\omega_1\alpha'_1 + \cdots + \omega_n\alpha'_n) = 1$ which is not possible since $\omega_i \geq 0$ and $\alpha'_i > 0$ for any i . Thus $\det(M_1) \neq 0$ and M_1 is invertible. In particular $\sigma := M_1^{-1}(\mathbb{R}_{\geq 0}^n)$ is a rational strongly convex cone. Moreover, since $A \in \mathcal{V}_{\alpha, \delta}$, we have $\langle \alpha, \tau \rangle > 0$ for any $\tau \in \sigma$, $\tau \neq 0$. Hence $A \in \mathbb{k} \llbracket x^\beta, \beta \in \sigma \cap \mathbb{Z}^n \rrbracket$.

By Example 3.18 integral homogeneous elements with respect to ν_α are either finite over \mathbb{k} , either of the form $cx_1^{\frac{n_1}{q}} \cdots x_n^{\frac{n_n}{q}}$ for some integers $n_1, \dots, n_n \in \mathbb{Z}_{\geq 0}$, $q \in \mathbb{N}$ such that $\sum_{j=1}^n \alpha_j n_j > 0$.

Using Theorem 5.12 and since $\overline{\mathbb{K}}_{\nu_\alpha} = \lim_{\gamma_1, \dots, \gamma_s} \mathbb{K}_{\nu_\alpha}^{\text{alg}}[\gamma_1, \dots, \gamma_s]$ where the γ_i are homogeneous with respect to ν_α , we have the second inclusion by replacing σ by the rational strongly convex cone generated by σ and the n -uples (n_1, \dots, n_n) corresponding to the homogeneous elements $\gamma_1, \dots, \gamma_s$. □

Remark 6.10. In fact the proof shows that the field $\overline{\mathbb{K}}_\alpha$, as soon as

$$\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n,$$

is the field of Puiseux power series with support in rational strongly convex cones σ such that $\langle \alpha, \gamma \rangle \geq 0$ for all $\gamma \in \sigma$. Thus $\overline{\mathbb{K}}_\alpha$ is the field of α -positive Puiseux series according to [AI].

Lemma 6.11. Let $\alpha \in \mathbb{R}_{> 0}^n$ and $\alpha' \in \text{Rel}_\alpha \cap \mathbb{N}^n$. Then

$$\psi_{\alpha', t} \circ \psi_{\alpha', s} = \psi_{\alpha', \nu_{\alpha'}(\delta)st+s+t} \quad \forall s, t \in \mathbb{Z}_{\geq 0}.$$

Proof. Let $A = \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}} \in \mathcal{V}_{\alpha, \delta}$. Then we have (see Equation (9) in the proof of Lemma 6.8):

$$\delta^s \varphi_{\alpha', s}(A) = \sum_{i \in \Lambda} a_i(\mathbf{x}) \delta(\mathbf{x})^{s(1+\nu_{\alpha'}(a_i(\mathbf{x}))-\nu_{\alpha'}(\delta(\mathbf{x}))m(i))-m(i)}.$$

If $t \in \mathbb{Z}_{\geq 0}$ and $l \in \mathbb{Z}_{\geq 0}$, and $a(\mathbf{x})$ and $\delta(\mathbf{x})$ are (α') -homogeneous, we have that

$$\varphi_{\alpha', t}(a(\mathbf{x})\delta(\mathbf{x})^l) = a(\mathbf{x})\delta(\mathbf{x})^{t\nu_{\alpha'}(a)+l(t\nu_{\alpha'}(\delta)+1)}.$$

Thus by denoting

$$p_{\alpha'}(i) := \nu_{\alpha'}(a_i(\mathbf{x})) - \nu_{\alpha'}(\delta(\mathbf{x}))m(i) \text{ and } d_{\alpha'} := \nu_{\alpha'}(\delta(\mathbf{x}))$$

we obtain

$$\begin{aligned} \delta^t \varphi_{\alpha', t}(\delta^s \varphi_{\alpha', s}(A)) &= \\ &= \delta^t \sum_{i \in \Lambda} a_i(x_1 \delta^{\alpha'_1 t}, \dots, x_n \delta^{\alpha'_n t}) \times \\ &\quad \delta(x_1 \delta^{\alpha'_1 t}, \dots, x_n \delta^{\alpha'_n t})^{s(1+\nu_{\alpha'}(a_i(\mathbf{x}))-\nu_{\alpha'}(\delta(\mathbf{x}))m(i))-m(i)} = \\ &= \sum_{i \in \Lambda} a_i \delta^{t+\nu_{\alpha'}(a_i)+(s+sp_{\alpha'}(i)-m(i))(td_{\alpha'}+1)} \\ &= \sum_{i \in \Lambda} a_i \delta^{d_{\alpha'} t + sp_{\alpha'}(i) + tp_{\alpha'}(i) + sp_{\alpha'}(i) - m(i) + d_{\alpha'} ts + t + s}. \end{aligned}$$

In particular we have

$$(11) \quad \delta^t \varphi_{\alpha', t}(\delta^s \varphi_{\alpha', s}(A)) = \delta^{d_{\alpha'} st+s+t} \varphi_{\alpha', d_{\alpha'} st+s+t}(A) \quad \forall t \in \mathbb{Z}_{\geq 0}.$$

□

Lemma 6.12. *Let $\alpha \in \mathbb{R}_{>0}^n$ and $\alpha' \in \text{Rel}_\alpha \cap \mathbb{N}^n$. For all $s_1, s_2 \in \mathbb{N}$ there exist $t_1, t_2 \in \mathbb{N}$ such that*

$$\psi_{\alpha', t_1} \circ \psi_{\alpha', s_1} = \psi_{\alpha', t_2} \circ \psi_{\alpha', s_2}.$$

Proof. Let d denote $\nu_{\alpha'}(\delta)$. Let p be a prime number and $k \in \mathbb{N}$ such that p^k divides $ds_1 + 1$ and $ds_2 + 1$. Then $\gcd(p, d) = 1$ and p^k divides $ds_1 - ds_2$. Thus p^k divides $s_1 - s_2$. This proves that $\gcd(ds_1 + 1, ds_2 + 1)$ divides $s_1 - s_2$. Thus there exist $t_1 \in \mathbb{Z}$ and $t_2 \in \mathbb{Z}$ such that $(ds_1 + 1)t_1 - (ds_2 + 1)t_2 = s_2 - s_1$. If $t_1 t_2 < 0$, let say $t_1 > 0$ and $t_2 < 0$, then $(ds_1 + 1)t_1 - (ds_2 + 1)t_2 > s_1 + s_2 > |s_1 - s_2|$ which is not possible. Thus we have that $t_1 t_2 \geq 0$. If $t_1 \leq 0$ and $t_2 \leq 0$, we can replace t_1 (resp. t_2) by $t_1 + k(ds_2 + 1)$ (resp. by $t_2 + k(ds_1 + 1)$) for some positive integer k large enough. This will allow to assume that t_1 and t_2 are positive integers. Hence

$$\exists t_1, t_2 \in \mathbb{N}, ds_1 t_1 + s_1 + t_1 = ds_2 t_2 + s_2 + t_2.$$

This proves the lemma by Lemma 6.11. □

Definition 6.13. Now we consider a subring R of $\mathbb{k}[[\mathbf{x}]]$ that is an excellent Henselian local ring with maximal ideal \mathfrak{m}_R and satisfying the following properties:

- (A) $\mathbb{k}[x_1, \dots, x_n]_{(\mathbf{x})} \subset R$,
- (B) $\mathfrak{m}_R = (\mathbf{x})R$ and $\widehat{R} = \mathbb{k}[[\mathbf{x}]]$,
- (C) if $p(\mathbf{x}) \in \mathbb{k}[[\mathbf{x}]]$ is (α) -homogeneous for some $\alpha \in \mathbb{R}_{>0}^n$ then

$$f(\mathbf{x}) \in R \implies f(p(\mathbf{x})x_1, \dots, p(\mathbf{x})x_n) \in R.$$

Remark 6.14. If \mathbb{k} is a field, the ring of algebraic power series $\mathbb{k}\langle \mathbf{x} \rangle$ is an excellent Henselian local ring satisfying Properties (A), (B) and (C). If \mathbb{k} is a valued field, then the field of convergent power series $\mathbb{k}\{\mathbf{x}\}$ does also.

For a field \mathbb{k} , the ring $\mathbb{k}[[x_1, \dots, x_r]]\langle x_{r+1}, \dots, x_n \rangle$ for formal power series algebraic over $\mathbb{k}[[x_1, \dots, x_r]][x_{r+1}, \dots, x_n]$ is also an excellent Henselian local ring satisfying Properties (A), (B) and (C).

Definition 6.15. Let $\alpha \in \mathbb{R}_{>0}^n$ and let δ be a (α) -homogeneous polynomial. Let R be a ring satisfying Definition 6.13. We set

$$\mathcal{V}_{\alpha, \delta}^R := \left\{ A \in \widehat{V}_{\nu_\alpha} / \exists \Lambda \text{ a finitely generated sub-semigroup of } \mathbb{R}_{\geq 0}, \right.$$

$$\left. \forall i \in \Lambda \exists a_i \in \mathbb{k}[[\mathbf{x}]] \text{ } (\alpha)\text{-homogeneous, } \exists a, b \geq 0 \forall i \in \Lambda \exists m(i) \in \mathbb{N} \text{ s.t.} \right.$$

$$m(i) \leq ai + b, \nu_\alpha \left(\frac{a_i}{\delta^{m(i)}} \right) = i, A = \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}}$$

$$\text{and } \exists \varepsilon > 0 \forall q \in \mathbb{N} \forall \alpha' \in \text{Rel}(\alpha, q, \varepsilon) \exists s \in \mathbb{N} \text{ such that } \psi_{\alpha', s}(A) \in R \}.$$

Then \mathcal{V}_α^R is the union of the sets $\mathcal{V}_{\alpha, \delta}^R$ when δ runs over all the (α) -homogeneous polynomials.

Lemma 6.16. *The sets $\mathcal{V}_{\alpha, \delta}^R$ and \mathcal{V}_α^R are subrings of $\mathcal{V}_{\alpha, \delta}$ and \mathcal{V}_α .*

Proof. Let $A = \sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}}$ and $B = \sum_{i \in \Lambda} \frac{b_i}{\delta^{n(i)}}$ $\in \mathcal{V}_{\alpha, \delta}^R$. Then there exists $\varepsilon > 0$ such that $\forall q \in \mathbb{N}$, $\forall \alpha' \in \text{Rel}(\alpha, q, \varepsilon)$, there exist $s_1, s_2 \in \mathbb{N}$ such that

$$\psi_{\alpha', s_1}(A), \psi_{\alpha', s_2}(B) \in R.$$

Then by Lemma 6.11, Lemma 6.12 and condition (C) of Definition 6.13 there exists $s \in \mathbb{N}$ such that

$$\psi_{\alpha',s}(A), \psi_{\alpha',s}(B) \in R.$$

This shows that $\psi_{\alpha',s}(A+B) = \psi_{\alpha',s}(A) + \psi_{\alpha',s}(B) \in R$ and $A+B \in \mathcal{V}_{\alpha,\delta}^R$.

Now by Lemma 6.8 we can assume that there exists $s(\varepsilon) \in \mathbb{N}$ such that $\psi_{\alpha',s}(AB) \in \mathbb{k}[\mathbf{x}]$ for all $s > s(\varepsilon)$, for all $q \in \mathbb{N}$ and all $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$. On the other hand since $\psi_{\alpha',s}(A), \psi_{\alpha',s}(B) \in R$ then $\psi_{\alpha',\nu_{\alpha'}(\delta)st+s+t}(A), \psi_{\alpha',\nu_{\alpha'}(\delta)st+s+t}(B) \in R$ for all $t \in \mathbb{N}$ by Lemma 6.11 and Condition (C) of Definition 6.13. Thus there exists $s \in \mathbb{N}$ such that

$$\psi_{\alpha',s}(A), \psi_{\alpha',s}(B) \in R \text{ and } \psi_{\alpha',s}(AB) \in \mathbb{k}[\mathbf{x}].$$

But we have that

$$\psi_{\alpha',s}(A)\psi_{\alpha',s}(B) = \delta^s \psi_{\alpha',s}(AB) \in R.$$

Hence by Artin Approximation Theorem (cf. [Po], [Sp2]) $\psi_{\alpha',s}(AB) \in R$.

Thus $AB \in \mathcal{V}_{\alpha,\delta}^R$. This proves that $\mathcal{V}_{\alpha,\delta}^R$ is a ring.

Since \mathcal{V}_{α}^R is the direct limit of the $\mathcal{V}_{\alpha,\delta}^R$ it is also a ring. \square

Example 6.17. If $\alpha \in \mathbb{N}^n$ and $R = \mathbb{C}\{\mathbf{x}\}$ is the ring of convergent power series over \mathbb{C} , we claim that

$$\mathcal{V}_{\alpha,\delta}^{\mathbb{C}\{\mathbf{x}\}} = \left\{ \sum_{i \in \mathbb{Z}_{\geq 0}} \frac{a_i}{\delta^{a(i+1)}} \mid \forall i \ a_i \in \mathbb{C}[\mathbf{x}] \text{ is } (\alpha)\text{-homogeneous,} \right.$$

$$\left. \nu_{\alpha} \left(\frac{a_i}{\delta^{a(i+1)}} \right) = i, \ a \in \mathbb{Z}_{\geq 0} \right.$$

$$\left. \text{and } \exists C, r > 0 \text{ such that } |a_i(z)| \leq Cr^i \|z\|_{\alpha}^{\nu_{\alpha}(a_i)} \ \forall z \in \mathbb{C}^n \right\}$$

where $\|z\|_{\alpha} := \max_{j=1,\dots,n} \left| z_j^{\frac{1}{\alpha_j}} \right|$ for any $z \in \mathbb{C}^n$.

First of all every element A of $\mathcal{V}_{\alpha,\delta}$ is of the form

$$A = \sum_{i \in \mathbb{Z}_{\geq 0}} \frac{a_i}{\delta^{m(i)}}$$

where $\nu_{\alpha} \left(\frac{a_i}{\delta^{m(i)}} \right) = i$ and $m(i) \leq ai + b$ for some $a, b \in \mathbb{Z}_{\geq 0}$. By multiplying the numerator and the denominator of $\frac{a_i}{\delta^{m(i)}}$ by $\delta^{ai+b-m(i)}$ and replacing a_i by $a_i \delta^{ai+b-m(i)}$, we may assume that $m(i) = ai + b$. If $a > b$, we may replace $\frac{a_i}{\delta^{ai+b}}$ by $\frac{a_i \delta^{a-b}}{\delta^{ai+a}}$, if $a < b$ we may replace $\frac{a_i}{\delta^{ai+b}}$ by $\frac{a_i \delta^{(b-a)i}}{\delta^{bi+b}}$.

Thus any element of $\mathcal{V}_{\alpha,\delta}$ is of the form $\sum_{i \in \mathbb{Z}_{\geq 0}} \frac{a_i}{\delta^{a(i+1)}}$ where $\nu_{\alpha} \left(\frac{a_i}{\delta^{a(i+1)}} \right) = i$ for all $i \in \mathbb{Z}_{\geq 0}$. In

this case $\nu_{\alpha}(a_i) = (a\nu_{\alpha}(\delta) + 1)i + a\nu_{\alpha}(\delta)$ for any $i \in \mathbb{N}$.

By Remark 6.3 $\text{Rel}(\alpha, q, \varepsilon) = \{q\alpha\}$ for $\varepsilon > 0$ small enough since $\alpha \in \mathbb{N}^n$. Then we have (with $s = a$ in Lemma 6.8):

$$f(\mathbf{x}) := \psi_{\alpha,a}(A) = \delta(\mathbf{x})^a \sum_{i \in \mathbb{Z}_{\geq 0}} \frac{a_i}{\delta^{a(i+1)}} (x_1 \delta(\mathbf{x})^{\alpha_1 a}, \dots, x_n \delta(\mathbf{x})^{\alpha_n a}) = \sum_{i \in \mathbb{Z}_{\geq 0}} a_i(\mathbf{x})$$

and $f(\mathbf{x}) \in \mathbb{C}[\mathbf{x}]$. Moreover we have for every $q \in \mathbb{N}$

$$f_q(\mathbf{x}) := \psi_{q\alpha,a}(A) = \delta(\mathbf{x})^a \sum_{i \in \mathbb{Z}_{\geq 0}} \frac{a_i}{\delta^{a(i+1)}} (x_1 \delta(\mathbf{x})^{\alpha_1 q a}, \dots, x_n \delta(\mathbf{x})^{\alpha_n q a}) = \sum_{i \in \mathbb{Z}_{\geq 0}} a_i(\mathbf{x}) \delta(\mathbf{x})^{a(q-1)i}$$

Thus $f \in \mathbb{C}\{\mathbf{x}\}$ if and only if this power series is convergent on a neighborhood of the origin. This neighborhood may be chosen of the form:

$$B_\alpha(0, r) := \{z \in \mathbb{C}^n \mid |z_j| \leq r^{\alpha_j}, j = 1, \dots, n\}.$$

For any $z \in B_\alpha(0, r)$ set $t_j^{\alpha_j} = z_j$ for $j = 1, \dots, n$ and $b_i(t) = a_i(z)$ for any $i \in \mathbb{N}$. Then f is convergent on $B_\alpha(0, r)$ if and only $\sum_{i \in \mathbb{Z}_{\geq 0}} b_i(t)$ is convergent on $B(0, r) := \{t \in \mathbb{C}^n \mid |t_j| \leq r, j = 1, \dots, n\}$. But this series is convergent if and only if there exist $c \geq 0$ and $\rho < 1$ such that $|b_i(t)| \leq c\rho^i$ for all $i \in \mathbb{Z}_{\geq 0}$ and all $t \in B(0, r)$. Since $b_i(t)$ is a homogeneous polynomial of degree $\nu_\alpha(a_i) = (ad + 1)i + ad$ where $d := \nu_\alpha(\delta)$, we have

$$\sup_{|t_j| \leq r, j=1, \dots, n} |b_i(t)| = r^{(ad+1)i+ad} \sup_{|t_j| \leq 1, j=1, \dots, n} |b_i(t)|.$$

We see that f is convergent if and only if there exist $C \geq 0$ and $R > 0$ such that

$$\sup_{|z_j| \leq 1, j=1, \dots, n} |a_i(z)| = \sup_{|t_j| \leq 1, j=1, \dots, n} |b_i(t)| \leq CR^i.$$

This is equivalent to the following inequality for any $z \in \mathbb{C}^n$:

$$(12) \quad |a_i(z)| = |b_i(t)| \leq \max_{j=1, \dots, n} |t_j| \sup_{|t_j| \leq 1, j=1, \dots, n} |b_i(t)| \leq CR^i \|z\|_\alpha^{\nu_\alpha(a_i)}.$$

On the other hand if $f \in \mathbb{C}\{\mathbf{x}\}$ we have seen that there exist $C \geq 0$ and $R > 0$ such that

$$\sup_{|z_j| \leq 1, j=1, \dots, n} |a_i(z)| \leq CR^i.$$

Thus

$$\sup_{|z_j| \leq 1, j=1, \dots, n} |a_i(z) \delta(z)^{a(q-1)i}| \leq C(RS)^i$$

where $S := \max_{|z_j| \leq 1, j=1, \dots, n} |\delta(z)|^{a(q-1)}$. Hence $f_q \in \mathbb{C}\{\mathbf{x}\}$ for every $q \in \mathbb{N}$. This proves the claim.

We have the following analogue of Theorem 5.12 in the Henselian case:

Theorem 6.18. *Let \mathbb{k} be a field of characteristic zero and let R be a subring of $\mathbb{k}[\mathbf{x}]$ satisfying Definition 6.13. Let $\alpha \in \mathbb{R}_{>0}^n$ and let us set $N = \dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \dots + \mathbb{Q}\alpha_n)$.*

Let $P(Z) \in \mathcal{V}_\alpha^R[\langle \gamma_1, \dots, \gamma_s \rangle][Z]$ be a distinguished polynomial of degree d where the γ_i are homogeneous elements with respect to ν_α . Then the roots of $P(Z)$ are in $\mathcal{V}_\alpha^R[\langle \gamma'_1, \dots, \gamma'_N \rangle]$ for some integral homogeneous elements $\gamma'_1, \dots, \gamma'_N$ with respect to ν_α .

Proof. Let $P(Z) = Z^d + a_1 Z^{d-1} + \dots + a_d$ with $a_j \in \mathcal{V}_\alpha^R[\langle \gamma_1, \dots, \gamma_s \rangle]$ for $1 \leq j \leq d$. By Theorem 5.12 we may assume that $P(Z)$ has a root $z \in \mathcal{V}_{\alpha, \delta}[\langle \gamma_1, \dots, \gamma_N \rangle]$. We denote

$$a_i = \sum_{i_1, \dots, i_N} A_{i, i_1, \dots, i_N} \gamma_1^{i_1} \dots \gamma_N^{i_N} \text{ with } A_{i, i_1, \dots, i_N} \in \mathcal{V}_{\alpha, \delta},$$

$$z = \sum_{i_1, \dots, i_N} z_{i_1, \dots, i_N} \gamma_1 t^{i_1} \dots \gamma_N^{i_N} \text{ with } z_{i_1, \dots, i_N} \in \mathcal{V}_{\alpha, \delta}.$$

Let us fix $\varepsilon > 0$, $q \in \mathbb{N}$, $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ and s satisfying Lemma 6.8 for the A_{i, i_1, \dots, i_N} and for the z_{i_1, \dots, i_N} . For convenience we denote by φ the morphism $\varphi_{\alpha', s}$ defined in Definition 6.7. Then if A denotes one of the A_{i, i_1, \dots, i_N} or the Z_{i_1, \dots, i_N} we have $\varphi(A) \in \mathcal{V}_{\alpha', \delta}$ by Lemma 6.8. We set

$$R := \mathcal{V}_{\alpha, \delta} \cap \varphi^{-1}(\mathcal{V}_{\alpha', \delta})$$

and R' denotes the subring of $\mathcal{V}_{\alpha, \gamma}[\langle \gamma_1, \dots, \gamma_s \rangle]$ of elements $\sum_{i_1, \dots, i_N} A_{i_1, \dots, i_N} \gamma_1^{i_1} \cdots \gamma_N^{i_N}$ whose coefficients A_{i_1, \dots, i_N} are in R .

Of course φ induces a morphism $R \rightarrow \mathcal{V}_{\alpha', \delta}$ but we have the following lemma:

Lemma 6.19. *Let γ_i be homogeneous elements with respect to ν_α for $1 \leq i \leq N$. Then there exist homogeneous element γ'_i with respect to $\nu_{\alpha'}$, $1 \leq i \leq N$, such that, for any finite number of elements $A_{i_1, \dots, i_N} \in \mathcal{V}_{\alpha, \delta}$,*

$$\varphi \left(\sum_{i_1, \dots, i_N} A_{i_1, \dots, i_N} \gamma_1^{i_1} \cdots \gamma_N^{i_N} \right) := \sum_{i_1, \dots, i_N} \varphi(A_{i_1, \dots, i_N}) \gamma_1'^{i_1} \cdots \gamma_N'^{i_N}$$

defines an extension of φ from R' to $\mathcal{V}_{\alpha', \delta}[\langle \gamma'_1, \dots, \gamma'_N \rangle]$.

Proof of Lemma 6.19. Let us assume that γ_i is a homogeneous element of degree e_i with respect to ν_α . Let

$$Q_i(Z) := g_{i,0}(\mathbf{x})Z^{q_i} + g_{i,1}(\mathbf{x})Z^{q_i-1} + \cdots + g_{i,q_i}(\mathbf{x})$$

be a polynomial such that $Q_i(\gamma'_i) = 0$ and such that $g_{i,j}(\mathbf{x})$ is a (α) -homogeneous polynomial of degree $d_i + j e_i$ for some d_i .

Then $g_{i,j}(\mathbf{x})$ is a (α') -homogeneous polynomial of degree $d'_i + j e'_i$ for some constants d'_i and e'_i . Indeed, if a, b and c are (α) -homogeneous polynomials and $\nu_\alpha(a) - \nu_\alpha(b) = \nu_\alpha(b) - \nu_\alpha(c)$, then ac and b^2 are two (α) -homogeneous polynomials of same degree, i.e. $ac - b^2$ is (α) -homogeneous. Then, by Lemma 6.4, $ac - b^2$ is (α') -homogeneous, thus $\nu_{\alpha'}(a) - \nu_{\alpha'}(b) = \nu_{\alpha'}(b) - \nu_{\alpha'}(c)$.

Set $\overline{Q}_i(Z) = \delta^{s e'_i q_i} Q_i\left(\frac{Z}{\delta^{s e'_i}}\right)$. We have

$$\overline{Q}_i(Z) := g_{i,0}(\mathbf{x})Z^{q_i} + g_{i,1}(\mathbf{x})\delta(\mathbf{x})^{s e'_i} Z^{q_i-1} + \cdots + g_{i,q_i}(\mathbf{x})\delta(\mathbf{x})^{s e'_i q_i}.$$

For any i let γ'_i denote a root of $\overline{Q}_i(Z)$. So γ'_i is a homogeneous element of degree $e'_i(1 + \nu_{\alpha'}(\delta(\mathbf{x}))s)$ with respect to $\nu_{\alpha'}$. Then it is straightforward to check that

$$\varphi \left(\sum A_{i_1, \dots, i_N} \gamma_1^{i_1} \cdots \gamma_N^{i_N} \right) = \sum \varphi(A_{i_1, \dots, i_N}) \gamma_1'^{i_1} \cdots \gamma_N'^{i_N}$$

defines an extension of φ from R' to $\mathcal{V}_{\alpha', \delta}[\langle \gamma'_1, \dots, \gamma'_N \rangle]$. \square

By Lemmas 6.19, 6.11 and 6.12, and Property (C) we can assume that s is large enough for having that

$$\delta^{js} \varphi(A_j) \in R[\gamma'_1, \dots, \gamma'_N]$$

for $1 \leq j \leq d$. Again by applying Lemmas 6.19, 6.11 and 6.12 we may even assume that

$$\delta^s \varphi(z) \in \mathbb{k}[\mathbf{x}][\gamma'_1, \dots, \gamma'_N]$$

by taking s large enough. Thus $z' := \delta^s \varphi(z) \in \mathbb{k}[\mathbf{x}][\gamma'_1, \dots, \gamma'_N]$ is a root of the polynomial

$$\overline{P}(Z) := Z^d + \delta^s \varphi(A_1) Z^{d-1} + \cdots + \delta^{ds} \varphi(A_d) \in R[Z].$$

Let us write

$$z' := \sum_{i_1, \dots, i_N} z'_{i_1, \dots, i_N} \gamma_1'^{i_1} \cdots \gamma_N'^{i_N}$$

with $z'_{i_1, \dots, i_N} \in \mathbb{k}[\mathbf{x}]$ for any i_1, \dots, i_N . Let us set

$$Z := \sum_{i_1, \dots, i_N} Z_{i_1, \dots, i_N} \gamma_1'^{i_1} \cdots \gamma_N'^{i_N}$$

where Z_{i_1, \dots, i_N} are new variables. Solving $P(Z) = 0$ is equivalent to solve a finite system (\mathcal{S}) of polynomial equations in the variables Z_{i_1, \dots, i_N} with coefficients in R , just by replacing Z by $\sum_{i_1, \dots, i_N} Z_{i_1, \dots, i_N} \gamma_1'^{i_1} \cdots \gamma_N'^{i_N}$ and replacing the high powers of the γ_i by smaller ones using the

division by the $Q_i(Z_i)$. By Artin Approximation Theorem (cf. [Po], [Sp2]), the set of solutions of (\mathcal{S}) in R is dense in the set of solutions in $\mathbb{k}[[\mathbf{x}]]$, but since $P(Z) = 0$ has a finite number of solutions, then (\mathcal{S}) has a finite number of solutions and they are in R . Thus $z'_{i_1, \dots, i_N} \in R$ for all i_1, \dots, i_N , hence $z' \in R[\gamma'_1, \dots, \gamma'_N]$. This proves that $z \in \mathcal{V}_{\alpha, \delta}^R[\langle \gamma_1, \dots, \gamma_N \rangle]$. \square

7. A GENERALIZATION OF ABHYANKAR-JUNG THEOREM

Definition 7.1. Let $\alpha \in \mathbb{N}^n$ and let $\theta \in \mathbb{C}[\mathbf{x}]$ be a (α) -homogeneous polynomial. Let $a > 0$, $C > 0$ and $\eta > 0$. Set :

$$\mathcal{D}_{\theta, C, a, \eta} := \left(\bigcup_{\substack{K > 0, \varepsilon > 0 \\ \varepsilon < K^a C}} C_{K, \varepsilon} \right) \cap B(0, \eta)$$


where $B(0, \eta)$ is the open ball centered in 0 and of radius η and

$$C_{K, \varepsilon} := \{x \in \mathbb{C}^n / d_\alpha(x, \theta^{-1}(0)) > K \|x\|_\alpha \text{ and } \|x\|_\alpha < \varepsilon\}$$

where $\|\cdot\|_\alpha$ is defined in Example 6.17 and d_α is defined as follows: for any $x, y \in \mathbb{C}^n$ let us denote by $x_i^{\frac{1}{\alpha_i}}$ (resp. $y_i^{\frac{1}{\alpha_i}}$) a complex α_i -th root of x_i (resp. y_i) and let \mathbb{U}_i be the set of α_i -roots of unity.

Then we define $d_\alpha(x, y) := \max_i \inf_{\xi \in \mathbb{U}_i} |x_i^{\frac{1}{\alpha_i}} - \xi y_i^{\frac{1}{\alpha_i}}|$ and $d_\alpha(x, \theta^{-1}(0)) := \inf_{x' \in \theta^{-1}(0)} d_\alpha(x, x')$.

Then $\mathcal{D}_{\theta, C, a, \eta}$ is the complement of a hornshaped neighborhood of $\{\theta = 0\}$ as we can see on the following picture (here $n = 2$ and $\alpha = (1, 1)$):



Lemma 7.2. Let $a \in \mathbb{N}^n$ and $A \in \mathcal{V}_{\alpha, \theta}^{\mathbb{C}\{\mathbf{x}\}}$. Then there exist constants $a > 0$ and $C > 0$ such that A is analytic on $\mathcal{D}_{\theta, C, a, \eta}$ for every $\eta > 0$.

Proof. We write $A = \sum_i \frac{a_i}{\theta^{m(i)}}$ where a_i is (α) -homogeneous for every $i \in \mathbb{N}$. By multiplying a_i by a convenient power of θ we may even assume that there exist positive constants a and b such that $m(i) = ai + b$ for every i .

If $\nu_\alpha(a_i) = d_i$ there exist $C > 0$ and $r > 0$ such that

$$(13) \quad |a_i(x)| \leq Cr^i \|x\|_\alpha^{d_i} \quad \forall x \in \mathbb{C}^n$$

by Theorem 6.18, Example 6.17 and Inequality (12) of Example 6.17. On the other hand we claim that there exists a constant $C' > 0$ such that

$$(14) \quad |\theta(x)| \geq C' d_\alpha(x, \theta^{-1}(0))^{\nu_\alpha(\theta)} \quad \forall x \in \mathbb{C}^n.$$

Indeed if we embed $\mathbb{C}\{\mathbf{x}\}$ in $\mathbb{C}\{\mathbf{y}\}$ by sending x_i onto $y_i^{\alpha_i}$, we have

$$\theta(\mathbf{x}) = \theta(y_1^{\alpha_1}, \dots, y_n^{\alpha_n}) = \tau(y_1, \dots, y_n)$$

and τ is a homogeneous polynomial of degree $\nu_\alpha(\theta)$. After a linear change of coordinates, we may assume that τ is a monic polynomial in y_n of degree $\nu_\alpha(\theta)$ multiplied by a constant. Then, for all $y_1, \dots, y_n \in \mathbb{C}^n$, we have

$$|\tau(y_1, \dots, y_n)| = C' \left| \prod_{i=1}^{\nu_\alpha(\theta)} (y_n - \varphi_i(y_1, \dots, y_{n-1})) \right|$$

where φ_i is a homogeneous function which is locally analytic outside the discriminant locus of τ , for some constant $C' > 0$. Thus

$$\begin{aligned} |\tau(y_1, \dots, y_n)| &\geq C' \min_i |y_n - \varphi_i(y_1, \dots, y_{n-1})|^{\nu_\alpha(\theta)} \geq \\ &\geq C' \inf_{y' \in \tau^{-1}(0)} \max_k |y_k - y'_k|^{\nu_\alpha(\theta)} = C' d(y, \tau^{-1}(0))^{\nu_\alpha(\theta)} \end{aligned}$$

since $(y_1, \dots, y_{n-1}, \varphi_i(y_1, \dots, y_{n-1})) \in \tau^{-1}(0)$ for any i . This proves (14).

Hence we have (for positive constants ε , K and $x \in C_{K,\varepsilon}$):

$$\begin{aligned} \left| \frac{a_i(x)}{\theta^{m(i)}(x)} \right| &\leq \frac{C}{C'^{m(i)}} \frac{r^i \|x\|_\alpha^{d_i}}{d_\alpha(x, \theta^{-1}(0))^{\nu_\alpha(\theta)m(i)}} = \frac{C}{C'^{m(i)}} \frac{r^i \|x\|_\alpha^{i+\nu_\alpha(\theta)m(i)}}{d_\alpha(x, \theta^{-1}(0))^{\nu_\alpha(\theta)m(i)}} \leq \\ &\leq \frac{Cr^i \|x\|_\alpha^i}{C'^{m(i)} K^{\nu_\alpha(\theta)m(i)}} \leq \frac{C(r\varepsilon)^i}{C'^{m(i)} K^{\nu_\alpha(\theta)m(i)}} = \frac{C}{C'^b K^{\nu_\alpha(\theta)b}} \left(\frac{r\varepsilon}{C'^a K^{\nu_\alpha(\theta)a}} \right)^i. \end{aligned}$$

Then if $\varepsilon < K^{a\nu_\alpha(\theta)} \left(\frac{C'^a}{r} \right)$, A defines an analytic function on the domain $C_{K,\varepsilon}$. Thus A defines an analytic function on the domain $\mathcal{D}_{\theta, C'^a/r, a\nu_\alpha(\theta), \eta}$ for every $\eta > 0$. \square

This following proposition has been proven by Tougeron in the case $\alpha = (1, \dots, 1)$ (see Proposition 2.8 [To]):

Proposition 7.3. *Let $\alpha \in \mathbb{N}^n$ and let $P(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$ be a monic polynomial whose discriminant is equal to δu where $\delta \in \mathbb{C}\{\mathbf{x}\}$ is (α) -homogeneous and $u \in \mathbb{C}\{\mathbf{x}\}$ is invertible. If $P(Z)$ factors as $P(Z) = P_1(Z) \cdots P_r(Z)$ where $P_i(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$ is an irreducible monic polynomial of $\mathbb{C}\{\mathbf{x}\}[Z]$ for all i , then $P_i(Z)$ is irreducible in $\mathcal{V}_\alpha^{\mathbb{C}\{\mathbf{x}\}}[Z]$.*

Proof. Let $Q(Z)$ be an irreducible monic factor of $P(Z)$ in $\mathcal{V}_\alpha[Z]$. By Theorem 5.12 there exists a (α) -homogeneous polynomial $\theta \in \mathbb{C}[x]$ such that the coefficients of $Q(Z)$ are in $\mathcal{V}_{\alpha,\theta}$. Let us denote by A one of these coefficients.

Since $\mathcal{V}_{\alpha,\theta} \subset \mathcal{V}_{\alpha,\theta\delta}$ we may assume that δ divides θ , thus

$$\delta^{-1}(0) \cap B(0, \varepsilon) \subset \theta^{-1}(0) \cap B(0, \varepsilon)$$

for every $\varepsilon > 0$.

Let $\eta > 0$ small enough such that the roots of $P(Z)$ are locally analytic on the domain

$$D_{\theta,\eta} := B(0, \eta) \setminus \theta^{-1}(0) \subset B(0, \eta) \setminus \delta^{-1}(0).$$

Since A is a polynomial depending on the roots of $P(Z)$ it is locally analytic on $D_{\theta,\eta}$.

On the other hand by Lemma 7.2 A defines an analytic function on a domain $\mathcal{D}_{\theta, C, a, \eta}$.

Thus by Lemma 7.4 given below A is global analytic on $D_{\theta,\eta}$. Since the roots of $P(Z)$ are bounded near the origin, A is bounded near the origin, thus A extends to an analytic function near the origin. This proves that A is analytic on a neighborhood of the origin and $Q(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$. \square

Lemma 7.4. *Set $C > 0$, $a > 0$ and $\eta > 0$ and let $\theta \in \mathbb{C}[\mathbf{x}]$ be a (α) -homogeneous polynomial. Let $A : D_{\theta,\eta} \rightarrow \mathbb{C}$ be a multivalued function. Let us assume that A is analytic on $\mathcal{D}_{\theta,C,a,\eta}$ and locally analytic on $D_{\theta,\eta}$. Then A is analytic on $D_{\theta,\eta}$.*

Proof. Since A is locally analytic on $D_{\theta,\eta}$, then A extends to an analytic function on a small neighborhood of every path in $D_{\theta,\eta}$. If A is not analytic on $D_{\theta,\eta}$, then there exists a loop based at a point p of $D_{\theta,\eta}$, denoted by $\varphi : [0, 1] \rightarrow D_{\theta,\eta}$ with $\varphi(0) = \varphi(1) = p$, such that A extends to an analytic function on a neighborhood of φ but $A \circ \varphi(0) \neq A \circ \varphi(1)$. Let us write $\varphi(t) = (\varphi_1(t), \dots, \varphi_n(t))$ and let us define $\Phi : [0, 1] \times S \rightarrow \mathbb{C}^n$ by

$$\Phi(t, s) := (s^{\alpha_1} \varphi_1(t), \dots, s^{\alpha_n} \varphi_n(t))$$

where $S := \{z \in \mathbb{C} / |z| \leq 1, \Re(z) > 0\}$.

Then we have that $\delta(\Phi(t, s)) = s^{\nu_\alpha(\delta)} \delta(\varphi(t)) \neq 0$ for any $(t, s) \in [0, 1] \times S$ since $\text{Im}(\varphi) \subset D_{\theta,\eta}$ and $s \neq 0$. Thus the image of Φ is included in $D_{\theta,\eta}$. Moreover, for any $t \in [0, 1]$, let $\Phi_t : S \rightarrow D_{\theta,\eta}$ be the function defined by $\Phi_t(s) := \Phi(t, s)$. Its image is simply connected since S is simply connected and Φ_t is analytic. Thus $A \circ \Phi_t$, which is locally analytic, extends to an analytic function on S by the Monodromy Theorem.

Let us denote by h the holomorphic function on S defined by

$$h(s) := A \circ \Phi(0, s) - A \circ \Phi(1, s)$$

for any $s \in S$.

For any $s \in S$ and any $t \in [0, 1]$ we have

$$\|\Phi(t, s)\|_\alpha = |s| \|\varphi(t)\|_\alpha$$

and

$$d_\alpha(\Phi(t, s), \theta^{-1}(0)) = |s| d_\alpha(\varphi(t), \theta^{-1}(0)).$$

Let us set

$$K := \frac{1}{2} \min_{t \in [0, 1]} \frac{d_\alpha(\Phi(t, s), \theta^{-1}(0))}{\|\Phi(t, s)\|_\alpha} = \frac{1}{2} \min_{t \in [0, 1]} \frac{d_\alpha(\varphi(t), \theta^{-1}(0))}{\|\varphi(t)\|_\alpha} > 0.$$

Thus for any s belonging to the domain $S \cap \{|s| < K^a C\}$, we have $\Phi(t, s) \in \mathcal{D}_{\theta,C,a,\eta}$. Since $\Phi(t, s) \in \mathcal{D}_{\theta,C,a,\eta}$ and A is analytic on $\mathcal{D}_{\theta,C,a,\eta}$, then $A \circ \Phi(0, s) = A \circ \Phi(1, s)$, thus $h(s) = 0$ on $S \cap \{|s| < K^a C\}$. Since h is holomorphic on the connected domain S , then $h \equiv 0$ on S . This contradicts the assumption. Hence A is analytic on $D_{\theta,\eta}$. \square

Then we can extend Proposition 7.3 to the formal setting over any field of characteristic zero:

Theorem 7.5. *Let \mathbb{k} be a field of characteristic zero and $\alpha \in \mathbb{R}_{>0}^n$. Let $P(Z) \in \mathbb{k}[\mathbf{x}][Z]$ be a monic polynomial whose discriminant is equal to δu where $\delta \in \mathbb{k}[\mathbf{x}]$ is (α) -homogeneous and $u \in \mathbb{k}[\mathbf{x}]$ is a unit. If $P(Z)$ factors as $P(Z) = P_1(Z) \cdots P_s(Z)$ where the $P_i(Z)$ are irreducible monic polynomials of $\mathbb{k}[\mathbf{x}][Z]$, then the $P_i(Z)$ remain irreducible in $\mathcal{V}_\alpha[Z]$.*

Proof. Let us prove this theorem when $P(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$. If $\alpha \in \mathbb{N}^n$, this is exactly Proposition 7.3. If $\alpha \notin \mathbb{N}^n$, then by Lemma 6.6, any decomposition $P(Z) = Q_1(Z) \cdots Q_r(Z)$ in $\mathcal{V}_\alpha[Z]$ is also a decomposition in $\mathcal{V}_{\alpha'}[Z]$ for $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ where ε is small enough. Then every irreducible monic factor of $Q_i(Z)$ in $\mathcal{V}_{\alpha'}[Z]$ is in $\mathbb{C}\{\mathbf{x}\}[Z]$ by Proposition 7.3, thus $Q_i(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$ for every i . In particular since the $Q_i(Z)$ are irreducible in $\mathcal{V}_\alpha[Z]$ then they are irreducible polynomials of $\mathbb{C}\{\mathbf{x}\}[Z]$.

Now let us consider the general case. Let

$$P(Z) = Z^d + a_{d-1}(\mathbf{x})Z^{d-1} + \cdots + a_0(\mathbf{x})$$

be a polynomial satisfying the hypothesis of the theorem with $a_k(\mathbf{x}) \in \mathbb{k}[\mathbf{x}]$ for $0 \leq k \leq d-1$. Since $P(Z)$ is defined over a field extension of \mathbb{Q} generated by countably many elements and since such a field extension embeds in \mathbb{C} , we may assume that \mathbb{C} is a field extension of \mathbb{k} and $P(Z) \in \mathbb{C}[\mathbf{x}]$.

The discriminant of $P(Z)$ is a polynomial depending on the coefficients $a_0(\mathbf{x}), \dots, a_{d-1}(\mathbf{x})$ that we denote by $D(a_0(\mathbf{x}), \dots, a_{d-1}(\mathbf{x}))$. Let

$$R(A_0, \dots, A_{d-1}, U) := D(A_0, \dots, A_{d-1}) - \delta(\mathbf{x})U \in \mathbb{C}[\mathbf{x}][A_0, \dots, A_{d-1}, U].$$

Then $R(a_0(\mathbf{x}), \dots, a_{d-1}(\mathbf{x}), u(\mathbf{x})) = 0$.

On the other hand, saying that $P(Z)$ factors as $P = P_1 \cdots P_s$ is equivalent to

$$\exists b_1(\mathbf{x}), \dots, b_r(\mathbf{x}) \text{ such that } a_i(\mathbf{x}) = R_i(b_1(\mathbf{x}), \dots, b_r(\mathbf{x})) \quad \forall i$$

for some polynomials $R_i(B_1, \dots, B_r) \in \mathbb{Q}[B_1, \dots, B_r]$, $0 \leq i \leq d-1$ (these R_i are the coefficients of Z^i in the product $P_1(Z) \cdots P_s(Z)$ and the b_j are the coefficients of the $P_k(Z)$).

By Artin Approximation Theorem [Art], for any integer $c > 0$ there exist convergent power series

$$\bar{a}_{0,c}(\mathbf{x}), \dots, \bar{a}_{d-1,c}(\mathbf{x}), \bar{u}_c(\mathbf{x}), \bar{b}_{1,c}(\mathbf{x}), \dots, \bar{b}_{r,c}(\mathbf{x}) \in \mathbb{C}\{\mathbf{x}\}$$

such that

$$(15) \quad R(\bar{a}_{0,c}(\mathbf{x}), \dots, \bar{a}_{d-1,c}(\mathbf{x}), \bar{u}_c(\mathbf{x})) = 0,$$

$$(16) \quad \bar{a}_{i,c}(\mathbf{x}) - R_i(\bar{b}_{1,c}(\mathbf{x}), \dots, \bar{b}_{r,c}(\mathbf{x})) = 0 \text{ for } 0 \leq i \leq d-1$$

and

$$\bar{a}_{k,c}(\mathbf{x}) - a_k(\mathbf{x}), \bar{u}_c(\mathbf{x}) - u(\mathbf{x}), \bar{b}_{l,c}(\mathbf{x}) - b_l(\mathbf{x}) \in (\mathbf{x})^c$$

for $0 \leq k \leq d, 1 \leq l \leq r$. Set

$$P_{(c)}(Z) := Z^d + \bar{a}_{d-1,c}(\mathbf{x})Z^{d-1} + \cdots + \bar{a}_{0,c}(\mathbf{x}).$$

Then $P_{(c)}(Z)$ factors as

$$P_{(c)}(Z) = P_{1,(c)}(Z) \cdots P_{s,(c)}(Z)$$

in $\mathbb{C}\{\mathbf{x}\}[Z]$ because of Equation (16) (the coefficients of the $P_{i,(c)}(Z)$ are the $b_{k,c}$), and $P_{i,(c)}(Z) - P_i(Z) \in (\mathbf{x})^c \mathbb{k}[\mathbf{x}][Z]$ for $1 \leq i \leq s$. Moreover the discriminant of $P_{(c)}(Z)$ is of the form $\delta(\mathbf{x})u_{(c)}$ where $u_{(c)}$ is a unit in $\mathbb{C}\{\mathbf{x}\}$ if $c \geq 1$ by Equation (15). Since $P_i(Z)$ is irreducible in $\mathbb{k}[\mathbf{x}][Z]$, then $P_{i,(c)}(Z)$ is irreducible in $\mathbb{k}[\mathbf{x}][Z]$ for all i for c large enough (let us say for $c \geq c_0$). Moreover we can remark that $\nu_\alpha(a) \geq \min_i \{\alpha_i\} \text{ord}(a)$ for any $a \in \mathbb{k}[\mathbf{x}]$, thus $\nu_\alpha(\bar{b}_{k,c}(\mathbf{x}) - b_k(\mathbf{x})) \geq \min_i \{\alpha_i\} c$.

Let $c \geq c_0$ and let us assume that $P_{i,(c)}(Z)$ is not irreducible in $\mathcal{V}_\alpha[Z]$. Thus it is the product of two monic polynomials: let us say

$$P_{i,(c)}(Z) = P_{i,(c),1}(Z)P_{i,(c),2}(Z)$$

with $P_{i,(c),1}(Z), P_{i,(c),2}(Z) \in \mathcal{V}_\alpha[Z]$ and $\deg_Z(P_{i,(c),k}(Z)) > 0$ for $k = 1, 2$. In fact by Theorem 6.18 we may assume that $P_{i,(c),1}(Z), P_{i,(c),2}(Z) \in \mathcal{V}_\alpha^{\mathbb{C}\{\mathbf{x}\}}[Z]$. By Proposition 7.3 we see that $P_{i,(c),1}(Z), P_{i,(c),2}(Z) \in \mathbb{C}\{x\}[Z]$, and by Proposition 7.6 $P_{i,(c),1}(Z), P_{i,(c),2}(Z) \in \mathbb{L}\{x\}[Z]$ where \mathbb{L} is a subfield of \mathbb{C} which is finite over \mathbb{k} . Thus $\mathbb{L} = \mathbb{k}[\gamma]$ by the Primitive Element Theorem where γ is a homogeneous element of degree 0 with respect to ν_α by Example 3.19. But we have $\mathcal{V}_\alpha \cap \mathbb{k}[\gamma] = \mathbb{k}$. Thus $P_{i,(c),1}(Z), P_{i,(c),2}(Z) \in \mathbb{k}\{\mathbf{x}\}[Z] \subset \mathbb{k}[\mathbf{x}][Z]$ which contradicts the

assumption that $P_{i,(c)}$ is irreducible in $\mathbb{k}[[\mathbf{x}]]\langle Z \rangle$. Thus $P_{i,(c)}(Z)$ is irreducible in $\mathcal{V}_\alpha[Z]$. Hence, by Corollary 4.14, $P_i(Z)$ is irreducible in $\mathcal{V}_\alpha[Z]$ since $\nu_\alpha(\bar{b}_{k,c}(\mathbf{x}) - b_k(\mathbf{x}))$ increases at least linearly with c . \square

The next proposition is a generalization of a result of S. Cutkosky and O. Kashcheyeva [CK] (see also Proposition 1 [AM]) and we will use it to prove Theorem 7.7. It is again an application of Theorem 5.12.

Proposition 7.6. *Let $\mathbb{k} \rightarrow \mathbb{k}'$ be a characteristic zero field extension. Let $f \in \mathbb{k}'[[\mathbf{x}]]$ be algebraic over $\mathbb{k}[[\mathbf{x}]]$ and let \mathbb{L} be the field extension of \mathbb{k} generated by all the coefficients of f . Then $\mathbb{k} \rightarrow \mathbb{L}$ is a finite field extension.*

Proof. Let $\alpha \in \mathbb{R}_{>0}^n$ such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n$. By Theorem 5.12 the roots of the minimal polynomial of f are in $\mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_n \rangle]$ for some homogeneous elements $\gamma_1, \dots, \gamma_n$ with respect to ν_α . Let us denote by \mathcal{V}'_α the ring defined in Definition 5.1 and Lemma 5.5 where \mathbb{k} is replaced by \mathbb{k}' . Then $\mathbb{k}'[[\mathbf{x}]]$ and $\mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_n \rangle]$ are subrings of $\mathcal{V}'_\alpha[\langle \gamma_1, \dots, \gamma_n \rangle]$. Thus by unicity of the roots of the minimal polynomial of f we have that $f \in \mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_n \rangle]$. By Example 3.16 the homogeneous elements γ_i may be written as $\gamma_i = c_i \mathbf{x}^{\beta_i}$ where c_i is algebraic over \mathbb{k}' (and so over \mathbb{k}) and $\beta_i \in \mathbb{Q}^n$ for $1 \leq i \leq n$.

By expanding f either as a formal power series of $\mathbb{k}'[[\mathbf{x}]]$, $f = \sum_i b_i(\mathbf{x})$ where $b_i(\mathbf{x}) \in \mathbb{k}'[[\mathbf{x}]]$ is a (α) -homogeneous polynomial for any i , either as an element of $\mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_n \rangle]$, $f = \sum_i \frac{a_i(\mathbf{x})}{\delta(\mathbf{x})^{m(i)}} \gamma_1^{k_1(i)} \cdots \gamma_n^{k_n(i)}$, and by identifying the homogeneous terms of same valuation (which are monomials by Example 3.16), we obtain a countable number of relations of the following form:

$$(17) \quad b(\mathbf{x})\delta^m(\mathbf{x}) = \sum a_{n_1, \dots, n_s}(\mathbf{x})\gamma_1^{n_1} \cdots \gamma_n^{n_n}$$

where $b(\mathbf{x})$ (corresponding to the $b_i(\mathbf{x})$), $a_{n_1, \dots, n_s}(\mathbf{x})$ (corresponding to the $a_i(\mathbf{x})$) and δ are monomials, $b(\mathbf{x}) \in \mathbb{k}'[[\mathbf{x}]]$, $a_{n_1, \dots, n_s}(\mathbf{x}) \in \mathbb{k}[[\mathbf{x}]]$, $m \in \mathbb{N}$, and the sum is finite. By dividing Equality (17) by \mathbf{x}^β for β well chosen, we see that the coefficient of $b(\mathbf{x})$ is in $\mathbb{k}[c_1, \dots, c_n]$ and \mathbb{L} is a subfield of $\mathbb{k}[c_1, \dots, c_n]$. \square

We can strengthen Theorem 7.5 as follows:

Theorem 7.7. *Let $\alpha \in \mathbb{R}_{>0}^n$ and let $P(Z) \in \mathbb{k}[[\mathbf{x}]]\langle Z \rangle$ be a monic polynomial such that its discriminant $\Delta = \delta u$ where $\delta \in \mathbb{k}[[\mathbf{x}]]$ is (α) -homogeneous and $u \in \mathbb{k}[[\mathbf{x}]]$ is a unit. Let us set $N := \dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n)$. Then there exist $\gamma_1, \dots, \gamma_N$ integral homogeneous elements with respect to ν_α and a (α) -homogeneous polynomial $c(\mathbf{x}) \in \mathbb{k}[[\mathbf{x}]]$ such that the roots of $P(Z)$ are in $\frac{1}{c(\mathbf{x})}\mathbb{k}'[[\mathbf{x}]]\langle \gamma_1, \dots, \gamma_N \rangle$ where $\mathbb{k} \rightarrow \mathbb{k}'$ is finite.*

Remark 7.8. This result shows that for a given root z of the polynomial $P(Z)$ the other roots of $P(Z)$ are obtained from z by the action of the elements of the Galois groups of the elements $\gamma_1, \dots, \gamma_N$ on z . For instance if $\alpha \in \mathbb{N}^n$ (so $N = 1$ – we can always assume this by Lemma 6.4), then the Galois group of $P(Z)$ is a quotient of the Galois group of the minimal polynomial of γ_1 , i.e. the Galois group of one weighted homogeneous polynomial.

Proof of Theorem 7.7. If $Q(Z)$ is a monic polynomial dividing $P(Z)$ in $\mathbb{k}[[\mathbf{x}]]\langle Z \rangle$, then the discriminant of $Q(Z)$ divides the discriminant of $P(Z)$. Thus we may assume that $P(Z)$ is irreducible. We will consider three cases: first the case where the coefficients of $P(Z)$ are complex analytic with $\alpha \in \mathbb{N}^n$, then with $\alpha \in \mathbb{R}_{>0}^n$, and finally the general case.

• Let us assume that $\alpha \in \mathbb{N}^n$ and that $P(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$. By Theorem 5.12 the roots of $P(Z)$ are of the form

$$\sum_{i_1, \dots, i_s} A_{i_1, \dots, i_s} \gamma_1^{i_1} \cdots \gamma_s^{i_s}$$

where $\gamma_1, \dots, \gamma_s$ are integral homogeneous elements with respect to ν_α and $A_{i_1, \dots, i_s} \in \mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}$ for any i_1, \dots, i_s . We may even choose $s = 1$ by Proposition 3.28, but we treat here the general case $s \geq 1$ that will be used in the sequel.

We replace $\gamma_1, \dots, \gamma_s$ by other integral homogeneous elements with respect to ν_α as follows: let us denote by $\gamma_{1,1} := \gamma_1, \dots, \gamma_{1,q_1}$ the conjugates of γ_1 over \mathbb{K}_n . If $\gamma_2 \notin \mathbb{K}_n[\gamma_{1,1}, \dots, \gamma_{1,q_1}]$ we denote by $\gamma_{2,1} := \gamma_2, \dots, \gamma_{2,q_2}$ its conjugates over $\mathbb{K}_n[\gamma_{1,1}, \dots, \gamma_{1,q_1}]$ and so on. So for $1 \leq l \leq s$, q_l denotes the degree of the minimal polynomial of γ_l over $\mathbb{K}_n[\gamma_{i,j}]_{1 \leq i < l, 1 \leq j \leq q_i}$, and for $1 \leq l \leq s$, $\gamma_{l,1}, \dots, \gamma_{l,q_l}$ denote the conjugates of $\gamma_l = \gamma_{l,1}$ over $\mathbb{K}_n[\gamma_{i,j}]_{1 \leq i < l, 1 \leq j \leq q_i}$. Then we may assume that the roots of $P(Z)$ are of the form

$$\sum_{\substack{0 \leq i_1 < q_1 \\ \dots \\ 0 \leq i_s < q_s}} A_{i_1, \dots, i_s} \gamma_{1,j_1}^{i_1} \cdots \gamma_{s,j_s}^{i_s}$$

where $A_{i_1, \dots, i_s} \in \mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}$, $\nu_\alpha(A_{i_1, \dots, i_s} \gamma_{1,j_1}^{i_1} \cdots \gamma_{s,j_s}^{i_s}) \geq 0$ for any i_1, \dots, i_s , and $1 \leq j_i \leq q_i$ for any i .

Let us assume that $P(Z)$ factors into a product of monic irreducible polynomials as $P(Z) = P_1(Z) \cdots P_r(Z)$ in $\mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}[\gamma_{i,j}]_{1 \leq i < s, 1 \leq j \leq q_i}[Z]$. We write the roots of $P_1(Z)$ as $z_j = \sum_{i=0}^{q_s-1} B_i \gamma_{s,j}^i$

where

$$B_i \in \mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}[\gamma_{i,j}]_{1 \leq i < s, 1 \leq j \leq q_i}$$

for all i . Then the roots of the other $P_l(Z)$ are $\sum_{i=0}^{q_s-1} B'_i \gamma_{s,j}^i$ where $(B'_0, \dots, B'_{q_s-1})$ is the image

(B_0, \dots, B_{q_s-1}) by a $\mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}$ -automorphism of $\mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}[\gamma_{i,j}]_{1 \leq i < s, 0 \leq j < q_i}$. If the roots of $P_1(Z)$ satisfy the theorem, then we see that the roots of the other $P_l(Z)$ will also satisfy the theorem since they are conjugates of the roots of $P_1(Z)$ by $\mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}$ -automorphisms of $\mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}[\gamma_{i,j}]_{1 \leq i < s, 0 \leq j < q_i}$. Thus it is enough to prove the result for the roots of $P_1(Z)$. We have

$$\begin{pmatrix} z_1 \\ z_2 \\ \vdots \\ z_{q_s} \end{pmatrix} = \begin{pmatrix} 1 & \gamma_{s,1} & \cdots & \gamma_{s,1}^{q_s-1} \\ 1 & \gamma_{s,2} & \cdots & \gamma_{s,2}^{q_s-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \gamma_{s,q_s} & \cdots & \gamma_{s,q_s}^{q_s-1} \end{pmatrix} \begin{pmatrix} B_0 \\ B_1 \\ \vdots \\ B_{q_s-1} \end{pmatrix}.$$

Let us set $M := \begin{pmatrix} 1 & \gamma_{s,1} & \cdots & \gamma_{s,1}^{q_s-1} \\ 1 & \gamma_{s,2} & \cdots & \gamma_{s,2}^{q_s-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \gamma_{s,q_s} & \cdots & \gamma_{s,q_s}^{q_s-1} \end{pmatrix}$. The determinant of M is a homogeneous element c

with respect to ν_α where $\nu_\alpha(c) = \frac{1}{2}q_s(q_s - 1)\nu_\alpha(\gamma_s)$. Thus we have

$$B_i = \frac{1}{c} (R_{i,1}(\gamma_{s,1}, \dots, \gamma_{s,q_s})z_1 + \cdots + R_{i,s}(\gamma_{s,1}, \dots, \gamma_{s,q_s})z_{q_s})$$

where the $R_{i,j}$ are polynomials with coefficients in \mathbb{Q} and the element $R_{i,j}(\gamma_{s,1}, \dots, \gamma_{s,q_s})$ is homogeneous with respect to ν_α . By multiplying c and $R_{i,1}(\gamma_{s,1}, \dots, \gamma_{s,q_s})z_1 + \dots + R_{i,s}(\gamma_{s,1}, \dots, \gamma_{s,q_s})z_{q_s}$ by the conjugates of c over $\mathbb{k}[\mathbf{x}]$ we may assume that $c = c(\mathbf{x}) \in \mathbb{k}[\mathbf{x}]$ is a (α) -homogeneous polynomial. The z_i and the $\gamma_{s,j}$ are locally analytic on $D_{\theta,\eta} := B(0,\eta) \setminus \theta^{-1}(0)$ and bounded near the origin, where $\{\theta = 0\}$ contains the discriminant locus of $P(Z)$ and of the minimal polynomials of the γ_i and η is small enough. Thus $c(\mathbf{x})B_i$ is locally analytic on $D_{\theta,\eta}$ for $1 \leq i \leq q_s$ and is bounded near the origin. Moreover $c(\mathbf{x})B_i$ is algebraic over $\mathbb{k}[\mathbf{x}]$ since the $g_{s,j}$ and the z_k are algebraic over $\mathbb{k}[\mathbf{x}]$. By induction on s (we replace z_1, \dots, z_{q_s} by $c(\mathbf{x})B_0, \dots, c(\mathbf{x})B_{q_s-1}$ - here we just used the fact that the roots of $P(Z)$ are algebraic over $\mathbb{k}[\mathbf{x}]$ and locally analytic over a domain of the form $D_{\theta,\eta}$) we see that there exists a (α) -homogeneous polynomial $c(\mathbf{x})$ such that $c(\mathbf{x})A_{\underline{i}}$ is locally analytic on $D_{\theta,\eta}$ and bounded near the origin for any $\underline{i} := (i_1, \dots, i_s)$. Since $c(\mathbf{x})A_{\underline{i}} \in \mathcal{K}_\alpha^{\mathbb{C}\{\mathbf{x}\}}$ and it is bounded near the origin, we see that $c(\mathbf{x})A_{\underline{i}} \in \mathcal{V}_\alpha^{\mathbb{C}\{\mathbf{x}\}}$. Thus it is analytic on $\mathcal{D}_{\theta,C,a,\eta}$ for C, a and η well chosen (see Lemma 7.2). Hence by Lemma 7.4 it is analytic on $D_{\theta,\eta}$ and since it is bounded near the origin, $c(\mathbf{x})A_{\underline{i}} \in \mathbb{C}\{\mathbf{x}\}$ for any \underline{i} .

• Now let us consider any $\alpha \in \mathbb{R}_{>0}^n$ and $P(Z) \in \mathbb{C}\{\mathbf{x}\}[Z]$. Then the roots of $P(Z)$ are in $\mathcal{V}_\alpha[\langle \gamma_1, \dots, \gamma_s \rangle]$ for some integral homogeneous elements with respect to ν_α denoted by $\gamma_1, \dots, \gamma_s$. Let us denote these roots by z_1, \dots, z_d . For any $\alpha' \in \mathbb{N}^n$ such that $\text{Rel}_\alpha \subset \text{Rel}_{\alpha'}$, $\gamma_1, \dots, \gamma_s$ are integral homogeneous elements with respect to $\nu_{\alpha'}$. Thus, for any $\varepsilon > 0$ small enough (say $\varepsilon < \varepsilon_0$), for any $q \in \mathbb{N}$ and any $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$, $z_1, \dots, z_d \in \mathcal{V}_{\alpha'}[\langle \gamma_1, \dots, \gamma_s \rangle]$ by Proposition 6.6. Moreover, by the previous case, we see that

$$\begin{aligned} & \forall \varepsilon \in]0, \varepsilon_0[, \forall q \in \mathbb{N}, \forall \alpha' \in \text{Rel}(\alpha, q, \varepsilon), \\ & \exists c_{\alpha'}(\mathbf{x}) \text{ an } (\alpha')\text{-homogeneous polynomial such that} \\ & c_{\alpha'}(\mathbf{x})z_1, \dots, c_{\alpha'}(\mathbf{x})z_d \in \mathbb{C}\{\mathbf{x}\}[\gamma_1, \dots, \gamma_s]. \end{aligned}$$

Moreover we see that that $c_\alpha(\mathbf{x})$ may be chosen as being the product of the determinants of Vandermonde matrices as M depending only on $\gamma_1, \dots, \gamma_s$, thus $c_{\alpha'}(\mathbf{x})$ does not depend on α' . Let us denote $c(\mathbf{x}) := c_{\alpha'}(\mathbf{x})$. Since $c(\mathbf{x})$ is a (α') -homogeneous polynomial for all $\alpha' \in \text{Rel}(\alpha, q, \varepsilon)$ then $c(\mathbf{x})$ is a (α) -homogeneous polynomial. This proves the result.

• Now let us consider the general case, $\alpha \in \mathbb{R}_{>0}^n$ and $P(Z) \in \mathbb{k}[\mathbf{x}][Z]$ where \mathbb{k} is a field of characteristic zero.

Let us write $P(Z) = Z^d + a_{d-1}(\mathbf{x})Z^{d-1} + \dots + a_0(\mathbf{x})$. Exactly as in the proof of Theorem 7.5 we may assume that \mathbb{C} is a field extension of \mathbb{k} . Let us use the notations of the proof of Theorem 7.5. Let

$$R(A_0, \dots, A_{d-1}, U) := D(A_0, \dots, A_{d-1}) - \delta(\mathbf{x})U \in \mathbb{C}[\mathbf{x}][A_0, \dots, A_{d-1}, U]$$

where D is the universal discriminant of a monic polynomial of degree d . Then

$$R(a_0(\mathbf{x}), \dots, a_{d-1}(\mathbf{x}), u(\mathbf{x})) = 0.$$

By Artin Approximation Theorem [Art], for any integer $c > 0$, there exist convergent power series $\bar{a}_{0,c}(\mathbf{x}), \dots, \bar{a}_{d-1,c}(\mathbf{x}), \bar{u}_c(\mathbf{x}) \in \mathbb{C}\{\mathbf{x}\}$ such that

$$(18) \quad R(\bar{a}_{0,c}(\mathbf{x}), \dots, \bar{a}_{d-1,c}(\mathbf{x}), \bar{u}_c(\mathbf{x})) = 0,$$

and

$$\bar{a}_{k,c}(\mathbf{x}) - a_k(\mathbf{x}), \bar{u}_c(\mathbf{x}) - u(\mathbf{x}) \in (\mathbf{x})^c \quad \text{for } 0 \leq k \leq d.$$

Let $P_{(c)}(Z) := Z^d + \bar{a}_{d-1,c}(\mathbf{x})Z^{d-1} + \dots + \bar{a}_{0,c}(\mathbf{x})$. Then $P_{(c)}(Z)$ is irreducible for c large enough (say $c \geq c_0$). Moreover the discriminant of $P_{(c)}(Z)$ is of the form $\delta(\mathbf{x})u_{(c)}$ where $u_{(c)}$ is a unit in $\mathbb{C}\{\mathbf{x}\}$ if $c \geq 1$ by Equation (18). By the previous case, the roots of $P_{(c)}(Z)$ are in

$\frac{1}{c_c(\mathbf{x})}\mathbb{C}\{\mathbf{x}\}[\gamma_{1,c}, \dots, \gamma_{N,c}]$ where $\gamma_{1,c}, \dots, \gamma_{N,c}$ are integral homogeneous elements with respect to ν_α and $c_c(\mathbf{x})$ is a (α) -homogeneous polynomial. By Proposition 4.17 and the previous cases, we may assume that the $\gamma_{i,c}$ does not depend on c , thus let us denote $\gamma_{i,c}$ by γ_i . Moreover $c_c(\mathbf{x})$ may be chosen as being the product of the determinants of Vandermonde matrices as M depending only on $\gamma_1, \dots, \gamma_s$, thus $c_c(\mathbf{x})$ does not depend on c . Let us denote by $c(\mathbf{x})$ this common (α) -homogeneous polynomial.

Thus, when c goes to infinity, we see that the roots of $P(Z)$ are in $\frac{1}{c(\mathbf{x})}\mathbb{C}[[\mathbf{x}]][\gamma_1, \dots, \gamma_N]$. Such a root has the form $\sum_{i_1, \dots, i_N} A_{i_1, \dots, i_N} \gamma_1^{i_1} \cdots \gamma_N^{i_N}$ where i_k runs from 0 to $q_k - 1$. In this case $c(\mathbf{x})A_{i_1, \dots, i_N} \in \mathbb{C}[[\mathbf{x}]]$ is algebraic over $\mathbb{k}[[\mathbf{x}]]$, thus $c(\mathbf{x})A_{i_1, \dots, i_N} \in \mathbb{k}'[[\mathbf{x}]]$ where $\mathbb{k} \rightarrow \mathbb{k}'$ is finite by Proposition 7.6. Thus the roots of $P(Z)$ are in $\frac{1}{c(\mathbf{x})}\mathbb{k}'[[\mathbf{x}]][\gamma_1, \dots, \gamma_N]$. \square

In the case where the α_i are linearly independent over \mathbb{Q} , we can choose $c(\mathbf{x}) = 1$. This is exactly the Abhyankar-Jung Theorem:

Corollary 7.9 (Abhyankar-Jung Theorem). *Let $P(Z) \in \mathbb{k}[[\mathbf{x}]][[Z]]$ be a monic polynomial whose discriminant has the form $\mathbf{x}^\beta u(\mathbf{x})$ where $\beta \in \mathbb{N}^n$ and $u(0) \neq 0$. Then there exist an integer $q \in \mathbb{N}$ and a finite field extension $\mathbb{k} \rightarrow \mathbb{k}'$ such that the roots of $P(Z)$ are in $\mathbb{k}'[[x_1^{\frac{1}{q}}, \dots, x_n^{\frac{1}{q}}]]$.*

Proof. By the previous theorem applied to any $\alpha \in \mathbb{R}_{>0}^n$ satisfying $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n$, the roots of $P(Z)$ are in $\frac{1}{\mathbf{x}^\gamma} \mathbb{k}'[[x_1^{\frac{1}{q}}, \dots, x_n^{\frac{1}{q}}]]$ for some $\beta \in \mathbb{N}^n$, $q \in \mathbb{N}$ and $\mathbb{k} \rightarrow \mathbb{k}'$ a finite field extension. Since the discriminant of any monic factor of $P(Z)$ in $\mathbb{k}'[[x_1, \dots, x_n]][Z]$ divides the discriminant of $P(Z)$, we may assume that $P(Z)$ is irreducible in $\mathbb{k}'[[x_1, \dots, x_n]][Z]$, thus we assume that $\mathbb{k}' = \mathbb{k}$.

Let z be a root of $P(Z)$ and let us denote by $\text{NP}(z)$ its Newton polyhedron. Then $\text{NP}(z) \subset -\gamma + \mathbb{R}_{\geq 0}^n$. Let us assume that $\text{NP}(z) \not\subset \mathbb{R}_{\geq 0}^n$. This means that there exists $\gamma' \in \text{NP}(z)$ such that one its coordinates, let us say γ'_n , is negative. But since z is a root of $P(Z)$ that is a monic polynomial with coefficients in $\mathbb{k}[[\mathbf{x}]]$ then $\nu_\alpha(z) \geq 0$ for any $\alpha \in \mathbb{R}_{>0}^n$. But in this case there exists $\alpha \in \mathbb{R}_{>0}^n$ such that $\langle \alpha, \gamma' \rangle < 0$ which is a contradiction. Thus $\text{NP}(z) \subset \mathbb{R}_{\geq 0}^n$ which proves the corollary. \square

Let us finish this part by giving a few results which are analogous to the fact that if $z \in \mathbb{C}\{t^{\frac{1}{k}}\}$ for some $k \in \mathbb{N}$, t being a single variable, then its minimal polynomial over $\mathbb{C}[[t]]$ is a polynomial with convergent power series. The next result can also be seen as the converse of Theorem 6.18:

Corollary 7.10. *Let $P(Z) \in \mathbb{k}[[\mathbf{x}]][[Z]]$ be an irreducible monic polynomial whose discriminant has the form $\delta(\mathbf{x})u(\mathbf{x})$, where $\delta(\mathbf{x})$ is a (α) -homogeneous polynomial, $\alpha \in \mathbb{R}_{>0}^n$, and $u(\mathbf{x}) \in \mathbb{k}[[\mathbf{x}]]$ is invertible. Let us assume that $P(Z)$ has a root in $\mathcal{V}_\alpha^R[\langle \gamma_1, \dots, \gamma_s \rangle]$ where R is an excellent Henselian local ring satisfying Properties (A), (B) and (C) and $\gamma_1, \dots, \gamma_s$ are homogeneous elements with respect to ν_α . Then the coefficients of $P(Z)$ are in R .*

Proof. By Theorem 7.5, $P(Z)$ is irreducible in $\mathcal{V}_\alpha[Z]$. Let

$$z \in \mathcal{V}_\alpha^R[\langle \gamma_1, \dots, \gamma_s \rangle]$$

be a root of $P(Z)$ as given in the statement. We can write $z = \sum A_{i_1, \dots, i_s} \gamma_1^{i_1} \cdots \gamma_s^{i_s}$ where the sum is finite and $A_{i_1, \dots, i_s} \in \mathcal{V}_\alpha^R$. Then the others roots of $P(Z)$ are of the form $\sum A_{i_1, \dots, i_s} \sigma(\gamma_1)^{i_1} \cdots \sigma(\gamma_s)^{i_s}$ where σ is a $\mathbb{K}_{\nu_\alpha}^{\text{alg}}$ -automorphism of $\overline{\mathbb{K}}_{\nu_\alpha}^{\text{alg}}$. Thus all the roots of $P(Z)$ are in $\overline{\mathcal{V}}_\alpha^R$. Hence the coefficients of $P(Z)$ are in $\overline{\mathcal{V}}_\alpha^R \cap \mathbb{k}[[\mathbf{x}]] = R$. \square

Definition 7.11. Let \mathbb{k} be a valued field and let σ be a strongly convex rational cone of \mathbb{R}^n containing $\mathbb{R}_{\geq 0}^n$. There exists an invertible $n \times n$ matrix $M = (m_{i,j})_{1 \leq i,j \leq n}$ such that $M\gamma \in \mathbb{R}_{\geq 0}^n$ for any $\gamma \in \sigma$. We denote by $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \mathbb{Z}^n\}$ the subring of $\mathbb{k}[[\mathbf{x}^\beta, \beta \in \sigma \cap \mathbb{Z}^n]]$ of power series $f(\mathbf{x})$ such that $f(\tau(\mathbf{x})) \in \mathbb{k}\{\mathbf{x}\}$ where τ is the map defined by

$$(\tau(x_1), \dots, \tau(x_n)) = (x_1^{m_{1,1}} \cdots x_n^{m_{1,n}}, \dots, x_1^{m_{n,1}} \cdots x_n^{m_{n,n}}).$$

By Example 6.17 $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \mathbb{Z}^n\}$ is a subring of $\mathcal{V}_{\alpha, \delta}^{\mathbb{k}\{x\}}$ for any α such that $\langle \alpha, \gamma \rangle > 0$ for all $\gamma \in \sigma \setminus \{0\}$.

Let us mention the following theorem proven by A. Gabrielov and J.-Cl. Tougeron by using transcendental methods (they use in a crucial way the maximum principle for analytic functions):

Theorem 7.12. [Ga][To] *Let $P(Z) \in \mathbb{C}[[\mathbf{x}]]\langle Z \rangle$ be an irreducible monic polynomial. If one root of $P(Z)$ is in $\mathbb{C}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$ where σ is a strongly convex rational cone and $q \in \mathbb{N}$, then $P(Z) \in \mathbb{C}\{\mathbf{x}\}\langle Z \rangle$.*

Using what we have done we extend this theorem to any algebraically closed valued field of characteristic zero under the assumption that the discriminant of $P(Z)$ is close to be weighted homogeneous. First we need the following lemma:

Lemma 7.13. *Let $P(Z) \in \mathbb{k}[[\mathbf{x}]]\langle Z \rangle$ be an irreducible monic polynomial where \mathbb{k} is a characteristic zero algebraically closed valued field. Let $\alpha \in \mathbb{R}_{> 0}^n$ such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n$ and $P(Z)$ is irreducible in $\mathcal{V}_\alpha\langle Z \rangle$. By Theorem 6.9, the roots of $P(Z)$ are in $\mathbb{k}[[\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n]]$ where σ is a strongly convex rational cone such that $\langle \alpha, \gamma \rangle > 0$ for any $\gamma \in \sigma$, $\gamma \neq 0$, and $q \in \mathbb{N}$. If one root of $P(Z)$ is in $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$, then the others roots of $P(Z)$ are in $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$ and $P(Z) \in \mathbb{k}\{\mathbf{x}\}\langle Z \rangle$.*

Proof. Let $z \in \mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$ be a root of $P(Z)$. For any $\xi = (\xi_1, \dots, \xi_n)$ vector of q -th roots of unity let us denote by z_ξ the element of $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$ obtain from z by replacing $(x_1^{\frac{1}{q}}, \dots, x_n^{\frac{1}{q}})$ by $(\xi_1 x_1^{\frac{1}{q}}, \dots, \xi_n x_n^{\frac{1}{q}})$. In particular $z_\xi \in \mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$. Then for any ξ , z_ξ is a root of $P(Z)$. Let I be a subset of \mathbb{U}_q^n , where \mathbb{U}_q is the group of q -th root of unity, such that

$$\begin{aligned} z_\xi &\neq z_{\xi'} \text{ for any } \xi, \xi' \in I, \xi \neq \xi', \\ \text{and } \forall \xi \in \mathbb{U}_q^n, \exists \xi' \in I, z_{\xi'} &= z_\xi. \end{aligned}$$

Let us set $Q(Z) = \prod_{\xi \in I} (Z - z_\xi)$. Then $Q(Z)$ is a monic polynomial of $\mathcal{V}_\alpha\langle Z \rangle$ whose roots are roots of $P(Z)$. Thus it divides $P(Z)$ in $\mathcal{V}_\alpha\langle Z \rangle$ hence, since $P(Z)$ is irreducible, $Q(Z) = P(Z)$. Thus the other roots of $P(Z)$ are in $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$ and $P(Z) \in \mathbb{k}\{\mathbf{x}\}\langle Z \rangle$. \square

Corollary 7.14. *Let $P(Z) \in \mathbb{k}[[\mathbf{x}]]\langle Z \rangle$ be an irreducible monic polynomial of degree d where \mathbb{k} is a characteristic zero algebraically closed valued field. Let $\alpha \in \mathbb{R}_{> 0}^n$ such that $\dim_{\mathbb{Q}}(\mathbb{Q}\alpha_1 + \cdots + \mathbb{Q}\alpha_n) = n$. Let us assume that there exists an irreducible monic polynomial $Q(Z) \in \mathbb{k}[[\mathbf{x}]]\langle Z \rangle$ of degree d whose discriminant Δ_Q is a monomial times a unit and such that*

$$\nu_\alpha(P(Z) - Q(Z)) \geq \frac{d}{2} \nu_\alpha(\Delta_Q).$$

Let us assume moreover that one of the roots of $P(Z)$ is in $\mathbb{k}\{\mathbf{x}^\beta, \beta \in \sigma \cap \frac{1}{q}\mathbb{Z}^n\}$ for some strongly convex rational cone σ , where $\langle \alpha, \gamma \rangle > 0$ for any $\gamma \in \sigma \setminus \{0\}$, and $q \in \mathbb{N}$. Then the coefficients of $P(Z)$ are in $\mathbb{k}\{\mathbf{x}\}$.

Proof. By Remark 4.15 and Proposition 4.14, the polynomial $P(Z)$ is irreducible in $\mathcal{V}_\alpha\langle Z \rangle$. Thus we can apply the previous Lemma. \square

8. DIOPHANTINE APPROXIMATION

Here we give a necessary condition for an element of $\widehat{\mathbb{K}}_\nu$ to be algebraic over \mathbb{K}_n :

Theorem 8.1. [Ro1][II] *Let ν be an Abhyankar valuation and let $z \in \mathbb{K}_\nu^{\text{alg}}$. Then there exist two constants $C > 0$ and $a \geq 1$ such that*

$$\left| z - \frac{f}{g} \right|_\nu \geq C |g|_\nu^a \quad \forall f, g \in \mathbb{k}[\mathbf{x}].$$

Proof. Let $P(Z) := a_0 Z^d + a_1 Z^{d-1} + \cdots + a_d \in \mathbb{K}_n[Z]$ be an irreducible polynomial such that $P(z) = 0$. Let $h \in \mathbb{k}[\mathbf{x}]$ and set

$$P_h(Z) := h^d a_0^{d-1} P\left(\frac{Z}{ha_0}\right).$$

Then $P_h(Z) = Z^d + a_1 h Z^{d-1} + a_2 a_0 h^2 Z^{d-2} + \cdots + a_d a_0^{d-1} h^d$ and $z h a_d$ is a root of $P_h(Z)$. It is straightforward to check that z satisfies the theorem if and only if $z h a_d$ does. Thus we may assume that $P(Z)$ is a monic polynomial and $\nu(z) > 0$ by choosing h such that $\nu(h)$ is large enough. Let us set $Q(Z_1, Z_2) := Z_1^d P(Z_2/Z_1)$. By Theorem 3.1 [Ro1] there exist two constants $a \geq d$ and $b \geq 0$ such that

$$\text{ord}(Q(f, g)) \leq a \min\{\text{ord}(f), \text{ord}(g)\} + b \quad \forall f, g \in \mathbb{k}[\mathbf{x}].$$

Moreover, by Izumi's Theorem ([Iz], [Re], [ELS]), there exists a constant $c \geq 1$ such that for all $f \in \mathbb{k}[\mathbf{x}]$, $\text{ord}(f) \leq \nu(f) \leq c \text{ord}(f)$. Thus

$$\nu(Q(f, g)) \leq ac \min\{\nu(f), \nu(g)\} + bc \quad \forall f, g \in \mathbb{k}[\mathbf{x}].$$

Since $P(Z)$ is irreducible in $\mathbb{K}_n[Z]$ and \mathbb{K}_n is a characteristic zero field, $P(Z)$ has no multiple roots in \widehat{V}_ν and we may write

$$P(Z) = R(Z)(Z - z)$$

where $R(Z) \in \widehat{V}_\nu[Z]$ and $R(z) \neq 0$. Set $r := \nu(z)$. Let $f, g \in \mathbb{k}[\mathbf{x}]$ with $g \neq 0$. Two cases may occur: either

$$(19) \quad \left| z - \frac{f}{g} \right|_\nu \geq e^{-r}$$

either $\nu\left(z - \frac{f}{g}\right) > r$. In the last case we have $\nu\left(\frac{f}{g}\right) = \nu(z) > 0$. In particular $\nu\left(R\left(\frac{f}{g}\right)\right) \geq 0$ and $\nu(f) > \nu(g)$. Thus

$$(ac - d)\nu(g) + bc \geq \nu\left(P\left(\frac{f}{g}\right)\right) \geq \nu\left(\frac{f}{g} - z\right).$$

Thus we have

$$(20) \quad A\nu(g) + B \geq \nu\left(\frac{f}{g} - z\right) \quad \text{or} \quad \left| z - \frac{f}{g} \right|_\nu \geq e^{-B} |g|_\nu$$

with $A = ac - d$ and $B = bc$. Then (19) and (20) prove the theorem. \square

Example 8.2. Let $\sigma := (-1, 1)\mathbb{R}_{>0} + (1, 0)\mathbb{R}_{>0} \subset \mathbb{R}^2$. This is a rational strongly convex cone of \mathbb{R}^2 . Let $f(x_1, x_2)$ be a power series, $f(x_1, x_2) \in \mathbb{k}[[x_1, x_2]]$. Let us set

$$g(x_1, x_2) := \sum_{i=0}^{\infty} \left(\frac{x_2}{x_1}\right)^{i!} + f(x_1, x_2) \in \mathbb{k}[[x^\beta, \beta \in \sigma \cap \mathbb{Z}]].$$

Then $g \in \mathcal{V}_\alpha$ for any $\alpha \in \mathbb{R}_{>0}^2$ such that $\alpha_2 > \alpha_1$. Moreover

$$\nu_\alpha \left(g - f - \sum_{i=0}^n \left(\frac{x_2}{x_1} \right)^{i!} \right) = (n+1)! (\alpha_2 - \alpha_1) = \frac{\alpha_2 - \alpha_1}{\alpha_1} (n+1) \nu_\alpha(x_1^{n!}).$$

Thus there do not exist constants A and B such that

$$A \nu_\alpha(x_1^{n!}) + B \geq \nu_\alpha \left(g - f - \sum_{i=0}^n \left(\frac{x_2}{x_1} \right)^{i!} \right) \quad \forall n \in \mathbb{N}.$$

Hence $g(x_1, x_2)$ is not algebraic over \mathcal{F}_2 by Theorem 8.1.

NOTATIONS

- ν_α is the monomial valuation defined by $\nu_\alpha(x_i) := \alpha_i$ for any i (cf. Example 2.4).
- V_ν is the valuation ring associated to ν .
- \widehat{V}_ν is the completion of V_ν .
- \mathbb{K}_n is the fraction field of $\mathbb{k}[\mathbf{x}]$ and V_ν .
- $\widehat{\mathbb{K}}_\nu$ is the fraction field of \widehat{V}_ν .
- $\text{Gr}_\nu V_\nu$ is the graded ring associated to V_ν (cf. Part 3).
- V_ν^{alg} is the algebraic closure (or the Henselization) of V_ν in \widehat{V}_ν (see Lemma 2.10).
- $\mathbb{K}_\nu^{\text{alg}}$ is the fraction field of V_ν^{alg} .
- V_ν^{fg} is the subring of \widehat{V}_ν whose elements have ν -support included in a finitely generated sub-semigroup of $\mathbb{R}_{\geq 0}$ (cf. Definition 3.14).
- $\mathbb{K}_\nu^{\text{fg}}$ is the fraction field of V_ν^{fg} .
- For any $\alpha \in \mathbb{R}_{>0}^n$, a (α) -homogeneous polynomial is a weighted homogeneous polynomial for the weights $\alpha_1, \dots, \alpha_n$ (see Definition 2.8).
- $A[\langle \gamma_1, \dots, \gamma_s \rangle]$ is the valuation ring associated to $A[\gamma_1, \dots, \gamma_s]$ when $A = \widehat{V}_\nu, V_\nu^{\text{fg}}$ or V_ν^{alg} (cf. Definition 3.26).
- \overline{V}_ν is the direct limit of the rings $\widehat{V}_\nu[\langle \gamma_1, \dots, \gamma_s \rangle]$ where the γ_i are homogeneous elements with respect to ν (cf. Definition 3.27).
- $\overline{\mathbb{K}}_\nu$ is the fraction field of \overline{V}_ν .
- $\overline{V}_\nu^{\text{alg}}$ is the direct limit of the rings $V_\nu^{\text{alg}}[\langle \gamma_1, \dots, \gamma_s \rangle]$ where the γ_i are homogeneous elements with respect to ν .
- $\overline{\mathbb{K}}_\nu^{\text{alg}}$ is the fraction field of $\overline{V}_\nu^{\text{alg}}$.
- $\overline{V}_\nu^{\text{fg}}$ is the direct limit of the rings $V_\nu^{\text{fg}}[\langle \gamma_1, \dots, \gamma_s \rangle]$ where the γ_i are homogeneous elements with respect to ν .
- $\overline{\mathbb{K}}_\nu^{\text{fg}}$ is the fraction field of $\overline{V}_\nu^{\text{fg}}$.
- $\mathcal{V}_{\alpha, \delta}$ is the subring of $V_{\nu_\alpha}^{\text{fg}}$ of elements of the form $\sum_{i \in \Lambda} \frac{a_i}{\delta^{m(i)}}$ where $\Lambda \subset \mathbb{R}$ is a finitely generated semigroup, $\nu_\alpha \left(\frac{a_i}{\delta^{m(i)}} \right) = i$ and $i \mapsto m(i)$ is bounded by an affine function (see Definition 5.1).
- \mathcal{V}_α is the direct limit of the $\mathcal{V}_{\alpha, \delta}$ over all the (α) -homogeneous polynomials δ . It is a valuation ring (cf. Proposition 5.5).
- \mathcal{K}_α is the fraction field of \mathcal{V}_α (cf. Definition 5.6).
- $\overline{\mathcal{K}}_\alpha$ is the direct limit of the fields $\mathcal{K}[\langle \gamma_1, \dots, \gamma_s \rangle]$ where the γ_i are homogeneous elements with respect to ν (cf. Definition 5.6).
- $\mathcal{V}_{\alpha, \delta}^R$ is the subring $\mathcal{V}_{\alpha, \delta}$ whose elements are in the Henselian ring R after a suitable transform (cf. Definition 6.15).

- \mathcal{V}_α^R is the direct limit of the $\mathcal{V}_{\alpha,\delta}^R$ over all the (α) -homogeneous polynomials δ .

REFERENCES

- [Ab] S. Abhyankar, On the ramification of algebraic functions, *Amer. J. Math.*, **77**, (1955), 575-592.
- [AM] S. Abhyankar, T. Moh, On analytic independence, *Trans. A.M.S.*, **219**, (1976), 77-87.
- [Aro] F. Aroca, Puiseux parametric equations of analytic sets, *Proc. Amer. Math. Soc.*, **132**, (2004), no. 10, 3035-3045.
- [AI] F. Aroca, G. Ilardi, A family of algebraically closed fields containing polynomials in several variables, *Comm. Algebra*, **37**, (2009), no. 4, 1284-1296.
- [Art] M. Artin, On the solutions of analytic equations, *Invent. Math.*, **5**, (1968), 277-291.
- [BM] E. Bierstone, P. D. Milman, Arc-analytic functions, *Invent. Math.*, **101** (1990), no. 2, 411-424.
- [BK] E. Brieskorn, H. Knörrer, Plane algebraic curves, Birkhäuser, (1986).
- [Cu] S. D. Cutkosky, Resolution of Singularities, *Graduate Studies in Mathematics*, **63**, American Math. Soc., (2004).
- [CK] S. D. Cutkosky, O. Kashcheyeva, Algebraic series and valuation rings over nonclosed fields, *J. of Pure and Applied Algebra*, **212**, (2008), 1996-2012.
- [ELS] L. Ein, R. Lazarsfeld, K. Smith, Uniform approximation of Abhyankar valuation ideals in smooth function fields, *Amer. J. Math.*, **125**, (2003), no. 2, 409-440.
- [Ei] G. Eisenstein, Über eine allgemeine Eigenschaft der Reihen-Entwicklungen aller Algebraischen Funktionen, *Bericht Königl. Preuss. Akad. d. Wiss. Zu Berlin*, 441-443, (1852).
- [FJ] C. Favre, M. Jonsson, The Valutive Tree, Lecture Notes in Mathematics, 1853. Springer-Verlag, Berlin, 2004. xiv+234 pp.
- [Ga] A. Gabriellov, Formal relations between analytic functions, *Izv. Akad. Nauk. SSSR*, **37**, 1056-1088, (1973).
- [Go] P. D. González Pérez, Singularités quasi-ordinaires toriques et polyèdre de Newton du discriminant, *Canad. J. Math.*, **52**, (2000), 348-368.
- [Gr] A. Grothendieck, Étude locale des schémas et des morphismes de schémas IV, *Inst. Hautes Études Sci. Publ. Math.*, No. 32, 1967.
- [HOV] F. Herrera Govantes, M. Olalla Acosta, J. L. Vicente Córdoba, Valuations in fields of power series, *Proceedings of the International Conference on Algebraic Geometry and Singularities (Sevilla, 2001)*, *Rev. Mat. Iberoamericana*, **19**, (2003), no. 2, 467-482.
- [II] H. Ito, S. Izumi, Diophantine inequality for equicharacteristic excellent Henselian local domains, *C. R. Math. Acad. Sci. Soc. R. Can.*, **30**, (2008), no. 2, 48-55.
- [Iz] S. Izumi, A mesure of integrity for local analytic algebras, *Publ. RIMS, Kyoto Univ.*, **21**, (1985), 719-736.
- [Ju] H. E. W. Jung, Darstellung der Funktionen eines algebraischen Körpers zweier unabhängiger Veränderlichen x, y in der Umgebung einer Stelle $x = a, y = b$. *J. Reine Angew. Math.*, **133**, (1908), 289-314.
- [KV] K. Kiyek, J. L. Vicente, On the Jung-Abhyankar theorem, *Arch. Math. (Basel)*, **83**, (2004), no. 2, 123-134.
- [McC] P. J. McCarthy, Algebraic extensions of fields, Chelsea Publishing Co., New York, (1976), ix+166 pp.
- [McD] J. McDonald, Fiber polytopes and fractional power series, *Journal of Pure and Applied Algebra*, **104**, (1995), 213-233.
- [M-B] L. Moret-Bailly, An extension of Greenberg's theorem to general valuation rings, *Manuscripta Math.*, **139**, (2012), no. 1-2, 153-166.
- [MS] H. Mourtada, B. Schober, A polyhedral characterization of quasi-ordinary singularities, arXiv:1512.07507.
- [PR] A. Parusiński, G. Rond, The Abhyankar-Jung Theorem, *Journal of Algebra*, **365**, (2012), 29-41.
- [Po] D. Popescu, General Neron desingularisation and approximation, *Nagoya Math. J.*, **104**, (1986), 85-115.
- [Pu1] V. Puiseux, Recherches sur les fonctions algébriques, *J. Math. Pures Appl.*, **15**, (1850), 365-480.
- [Pu2] V. Puiseux, Nouvelles recherches sur les fonctions algébriques, *J. Math. Pures Appl.*, **16**, (1851), 228-240.
- [Re] D. Rees, Izumi's theorem, Commutative algebra (Berkeley, CA, 1987), 407-416, *Math. Sci. Res. Inst. Publ.*, **15**, (1989).
- [Ri] P. Ribenboim, Fields: algebraically closed and others, *Manuscripta Math.*, **75**, (1992), 115-150.
- [Ro1] G. Rond, Approximation diophantienne dans le corps des séries en plusieurs variables, *Ann. Institut Fourier*, vol. 56, no. 2, (2006), 299-308.
- [Ro2] G. Rond, Homomorphisms of local algebras in positive characteristic, *J. Algebra*, **322**, (2009), no. 12, 4382-4407.
- [RS] G. Rond, M. Spivakovsky, The analogue of Izumi's Theorem for Abhyankar valuations, *J. Lond. Math. Soc.*, **90** (3), (2014), 725-740.
- [Sa] A. Sathaye, Generalized Newton-Puiseux expansions and Abhyankar-Moh semigroup theorem, *Invent. Math.*, **74**, (1983), 149-157.

- [Sp1] M. Spivakovsky, Valuations in function fields of surfaces, *Amer. J. Math.*, **112**, (1990), no. 1, 107-156.
- [Sp2] M. Spivakovsky, A new proof of D. Popescu's theorem on smoothing of ring homomorphisms, *J. Amer. Math. Soc.*, **12** (1999), no. 2, 381-444.
- [SV] M. J. Soto, J. L. Vicente, The Newton procedure for several variables, *Linear Algebra Appl.*, **435**, (2011), no. 2, 255-269.
- [Te] B. Teissier, Overweight deformations of affine toric varieties and local uniformization, "Valuation theory in interaction", Proceedings of the second international conference on valuation theory, Segovia-El Escorial, 2011. European Math. Soc. Publishing House, Congress Reports Series, (2014), 474-565.
- [To] J.-Cl. Tougeron, Sur les racines d'un polynôme à coefficients séries formelles, *Real analytic and algebraic geometry (Trento 1988)*, 325-363, *Lectures Notes in Math.*, **1420**, (1990).

E-mail address: guillaume.rond@univ-amu.fr

AIX-MARSEILLE UNIVERSITÉ, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453 MARSEILLE, FRANCE