

REPRESENTATIONS OF QUANTUM GROUPS AT ROOTS OF UNITY, WHITTAKER VECTORS AND Q-W ALGEBRAS

A. SEVOSTYANOV

ABSTRACT. Let $U_\varepsilon(\mathfrak{g})$ be the standard simply connected version of the Drinfeld–Jumbo quantum group at an odd m -th root of unity ε . The center of $U_\varepsilon(\mathfrak{g})$ contains a huge commutative subalgebra isomorphic to the algebra Z_G of regular functions on (a finite covering of a big cell in) a complex connected, simply connected algebraic group G with Lie algebra \mathfrak{g} . Let V be a finite-dimensional representation of $U_\varepsilon(\mathfrak{g})$ on which Z_G acts according to a non-trivial character η_g given by evaluation of regular functions at $g \in G$. Then V is a representation of the finite-dimensional algebra $U_{\eta_g} = U_\varepsilon(\mathfrak{g})/U_\varepsilon(\mathfrak{g})\text{Ker } \eta_g$. We show that in this case, under certain restrictions on m , U_{η_g} contains a subalgebra $U_{\eta_g}(\mathfrak{m}_-)$ of dimension $m^{\frac{1}{2}\dim \mathcal{O}}$, where \mathcal{O} is the conjugacy class of g , and $U_{\eta_g}(\mathfrak{m}_-)$ has a one-dimensional representation \mathbb{C}_{χ_g} . We also prove that if V is not trivial then the space of Whittaker vectors $\text{Hom}_{U_{\eta_g}(\mathfrak{m}_-)}(\mathbb{C}_{\chi_g}, V)$ is not trivial and the algebra $W_{\eta_g} = \text{End}_{U_{\eta_g}}(U_{\eta_g} \otimes_{U_{\eta_g}(\mathfrak{m}_-)} \mathbb{C}_{\chi_g})$ naturally acts on it which gives rise to a Schur-type duality between representations of the algebra U_{η_g} and of the algebra W_{η_g} called a q-W algebra.

1. INTRODUCTION

Let \mathfrak{g}' be the Lie algebra of a semisimple algebraic group G' over an algebraically closed field \mathbf{k} of characteristic $p > 0$. Let $x \mapsto x^{[p]}$ be the p -th power map of \mathfrak{g}' into itself. The structure of the enveloping algebra of \mathfrak{g}' is quite different from the zero characteristic case. Namely, the elements $x^p - x^{[p]}$, $x \in \mathfrak{g}'$ are central. For any linear form θ on \mathfrak{g}' , let U_θ be the quotient of the enveloping algebra of \mathfrak{g}' by the ideal generated by the central elements $x^p - x^{[p]} - \theta(x)^p$ with $x \in \mathfrak{g}'$. Then U_θ is a finite-dimensional algebra. Kac and Weisfeiler proved that any simple \mathfrak{g}' -module can be regarded as a module over U_θ for a unique θ as above (this explains why all simple \mathfrak{g}' -modules are finite-dimensional).

One can identify θ with an element of \mathfrak{g}' via the Killing form and reduce the general situation to the case of nilpotent θ . In that case, among other things, Premet defines in [13, 14] a subalgebra $U_\theta(\mathfrak{m}_\theta) \subset U_\theta$ generated by a Lie subalgebra $\mathfrak{m}_\theta \subset \mathfrak{g}'$ such that $U_\theta(\mathfrak{m}_\theta)$ has dimension $p^{\frac{d}{2}}$, where d is the dimension of the coadjoint orbit of θ , and is equipped with a one-dimensional representation \mathbf{k}_{χ_θ} , $\chi_\theta : U_\theta(\mathfrak{m}_\theta) \rightarrow \mathbf{k}$. Thus for every U_θ -module V the algebra $\text{End}_{U_\theta}(U_\theta \otimes_{U_\theta(\mathfrak{m}_\theta)} \mathbf{k}_{\chi_\theta})$ naturally acts on the space $\text{Hom}_{U_\theta}(U_\theta \otimes_{U_\theta(\mathfrak{m}_\theta)} \mathbf{k}_{\chi_\theta}, V)$ called the space of Whittaker vectors in V . This leads to a Schur-type duality between representations of the algebras U_θ and $\text{End}_{U_\theta}(U_\theta \otimes_{U_\theta(\mathfrak{m}_\theta)} \mathbf{k}_{\chi_\theta})$, the latter being called a W-algebra. Tensoring V with the one-dimensional representation $\mathbf{k}_{-\chi_\theta}$ of $U_\theta(\mathfrak{m}_\theta)$ one can define the structure of a $U_0(\mathfrak{m}_\theta)$ -module (here $U_0(\mathfrak{m}_\theta)$ is the subalgebra generated by \mathfrak{m}_θ in U_0 which corresponds to $\theta = 0$) on V , which makes it a $U_\theta - U_0(\mathfrak{m}_\theta)$ -bimodule, and the two actions satisfy certain compatibility conditions. This brings the representation theory of the algebra U_θ into the context of equivariant representation theory showing also some similarity with the theory of generalized Gelfand–Graev representations (see [9]).

Another important example of finite-dimensional algebras is related to the theory of quantum groups at roots of unity. Let \mathfrak{g} be a complex finite-dimensional semisimple Lie algebra. A remarkable

property of the standard Drinfeld-Jimbo quantum group $U_\varepsilon(\mathfrak{g})$ associated to \mathfrak{g} , where ε is a primitive m -th root of unity, is that its center contains a huge commutative subalgebra isomorphic to the algebra Z_G of regular functions on (a finite covering of a big cell in) a complex algebraic group G with Lie algebra \mathfrak{g} (see [3, 4]). In this paper we consider the simply connected version of $U_\varepsilon(\mathfrak{g})$ and the case when m is odd. In that case G is the connected, simply connected algebraic group corresponding to \mathfrak{g} .

Consider finite-dimensional representations of $U_\varepsilon(\mathfrak{g})$, on which Z_G acts according to non-trivial characters η_g given by evaluation of regular functions at various points $g \in G$. Note that all irreducible representations of $U_\varepsilon(\mathfrak{g})$ are of that kind, and every such representation is a representation of the algebra $U_{\eta_g} = U_\varepsilon(\mathfrak{g})/U_\varepsilon(\mathfrak{g})\text{Ker } \eta_g$ for some η_g .

In this paper we construct certain subalgebras $U_{\eta_g}(\mathfrak{m}_-)$ in U_{η_g} which have properties similar to those of the subalgebras $U_\theta(\mathfrak{m}_\theta) \subset U_\theta$. In particular, there is a Schur-type duality between representations of the algebras U_{η_g} and of certain quantum group versions of W-algebras called q-W-algebras. U_{η_g} -modules also naturally become $U_{\eta_g} - U_{\eta_1}(\mathfrak{m}_-)$ -bimodules. We show that for such modules the corresponding spaces of Whittaker vectors are always non-trivial which can be regarded as an analogue of the Engel theorem for quantum groups at roots of unity.

It turns out that the definition of the subalgebras $U_{\eta_g}(\mathfrak{m}_-)$ is related to the existence of some special transversal slices Σ_s to the set of conjugacy classes in G . These slices Σ_s associated to (conjugacy classes of) elements s in the Weyl group of \mathfrak{g} were introduced by the author in [15]. The slices Σ_s play the role of the Slodowy slices in algebraic group theory. In the particular case of elliptic Weyl group elements these slices were also introduced later by He and Lusztig in paper [7] within a different framework.

A remarkable property of a slice Σ_s is that if g is conjugate to an element in Σ_s then U_{η_g} has a subalgebra of dimension $m^{\frac{1}{2}\text{codim } \Sigma_s}$ with a non-trivial character. If $g \in \Sigma_s$ (in fact g may belong to a larger variety) then the corresponding subalgebra $U_{\eta_g}(\mathfrak{m}_-)$ can be explicitly described in terms of quantum group analogues of root vectors. There are also analogues of subalgebras $U_{\eta_g}(\mathfrak{m}_-)$ in $U_q(\mathfrak{g})$ in case of generic q (see [16]).

Q-W-algebras can be regarded as noncommutative deformations of truncated versions of the algebras of regular functions on the slices Σ_s . In case of generic ε q-W algebras were introduced and studied in [16].

In [17], Theorem 5.2 it is shown that for every conjugacy class \mathcal{O} in G one can find a transversal slice Σ_s such that \mathcal{O} intersects Σ_s and $\dim \mathcal{O} = \text{codim } \Sigma_s$. Thus in this case for the corresponding algebra $U_{\eta_g}(\mathfrak{m}_-)$ we have $\dim U_{\eta_g}(\mathfrak{m}_-) = m^{\frac{1}{2}\dim \mathcal{O}}$, where \mathcal{O} is the conjugacy class of g .

There is, however, a major difference between Lie algebras and quantum groups: in case of Lie algebras \mathfrak{g}' over fields of prime characteristic the algebras $U_\theta(\mathfrak{m}_\theta)$ are local while in the quantum group case the algebras $U_{\eta_g}(\mathfrak{m}_-)$, which play the role of $U_\theta(\mathfrak{m}_\theta)$, are not local. In particular, each $U_{\eta_g}(\mathfrak{m}_-)$ may have several one-dimensional representations.

The author is grateful to the referee for careful reading of the text.

2. NOTATION

Fix the notation used throughout the text. Let G be a simply connected finite-dimensional complex simple Lie group, \mathfrak{g} its Lie algebra. Fix a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ and let Δ be the set of roots of $(\mathfrak{g}, \mathfrak{h})$. Let α_i , $i = 1, \dots, l$, $l = \text{rank}(\mathfrak{g})$ be a system of simple roots, $\Delta_+ = \{\beta_1, \dots, \beta_N\}$ the set of positive roots. Let H_1, \dots, H_l be the set of simple root generators of \mathfrak{h} .

Let a_{ij} be the corresponding Cartan matrix, and let d_1, \dots, d_l , $d_i \in \{1, 2, 3\}$, $i = 1, \dots, l$ be coprime positive integers such that the matrix $b_{ij} = d_i a_{ij}$ is symmetric. There exists a unique non-degenerate invariant symmetric bilinear form $(,)$ on \mathfrak{g} such that $(H_i, H_j) = d_j^{-1} a_{ij}$. It induces an isomorphism of vector spaces $\mathfrak{h} \simeq \mathfrak{h}^*$ under which $\alpha_i \in \mathfrak{h}^*$ corresponds to $d_i H_i \in \mathfrak{h}$. We denote

by α^\vee the element of \mathfrak{h} that corresponds to $\alpha \in \mathfrak{h}^*$ under this isomorphism. The induced bilinear form on \mathfrak{h}^* is given by $(\alpha_i, \alpha_j) = b_{ij}$.

Let W be the Weyl group of the root system Δ . W is the subgroup of $GL(\mathfrak{h})$ generated by the fundamental reflections s_1, \dots, s_l ,

$$s_i(h) = h - \alpha_i(h)H_i, \quad h \in \mathfrak{h}.$$

The action of W preserves the bilinear form $(,)$ on \mathfrak{h} .

For any root $\alpha \in \Delta$ we also denote by s_α the corresponding reflection.

For every element $w \in W$ one can introduce the set $\Delta_w = \{\alpha \in \Delta_+ : w(\alpha) \in -\Delta_+\}$, and the number of the elements in the set Δ_w is equal to the length $l(w)$ of the element w with respect to the system Γ of simple roots in Δ_+ .

Let \mathfrak{b}_+ be the Borel subalgebra corresponding to Δ_+ , \mathfrak{b}_- the opposite Borel subalgebra, \mathfrak{n}_\pm their nilradicals. Denote by H, N_+, N_-, B_+, B_- the maximal torus, the maximal unipotent subgroups and the Borel subgroups of G which correspond to the Lie subalgebras $\mathfrak{h}, \mathfrak{n}_+, \mathfrak{n}_-, \mathfrak{b}_+$ and \mathfrak{b}_- , respectively.

We identify \mathfrak{g} and its dual by means of the canonical invariant bilinear form. Then the coadjoint action of G on \mathfrak{g}^* is naturally identified with the adjoint one.

Let \mathfrak{g}_β be the root subspace corresponding to a root $\beta \in \Delta$, $\mathfrak{g}_\beta = \{x \in \mathfrak{g} | [h, x] = \beta(h)x \text{ for every } h \in \mathfrak{h}\}$. $\mathfrak{g}_\beta \subset \mathfrak{g}$ is a one-dimensional subspace. It is well known that for $\alpha \neq -\beta$ the root subspaces \mathfrak{g}_α and \mathfrak{g}_β are orthogonal with respect to the canonical invariant bilinear form. Moreover \mathfrak{g}_α and $\mathfrak{g}_{-\alpha}$ are non-degenerately paired by this form.

Root vectors $X_\alpha \in \mathfrak{g}_\alpha$ satisfy the following relations:

$$[X_\alpha, X_{-\alpha}] = (X_\alpha, X_{-\alpha})\alpha^\vee.$$

Note also that in this paper we denote by \mathbb{N} the set of nonnegative integer numbers, $\mathbb{N} = \{0, 1, \dots\}$.

3. QUANTUM GROUPS

In this paper we shall consider some specializations of the standard Drinfeld-Jimbo quantum group $U_h(\mathfrak{g})$ defined over the ring of formal power series $\mathbb{C}[[h]]$, where h is an indeterminate. We follow the notation of [2].

Let V be a $\mathbb{C}[[h]]$ -module equipped with the h -adic topology. This topology is characterized by requiring that $\{h^n V \mid n \geq 0\}$ is a base of the neighborhoods of 0 in V , and that translations in V are continuous.

A topological Hopf algebra over $\mathbb{C}[[h]]$ is a complete $\mathbb{C}[[h]]$ -module equipped with a structure of $\mathbb{C}[[h]]$ -Hopf algebra (see [2], Definition 4.3.1), the algebraic tensor products entering the axioms of the Hopf algebra are replaced by their completions in the h -adic topology.

The standard quantum group $U_h(\mathfrak{g})$ associated to a complex finite-dimensional simple Lie algebra \mathfrak{g} is a topological Hopf algebra over $\mathbb{C}[[h]]$ topologically generated by elements $H_i, X_i^+, X_i^-, i = 1, \dots, l$, subject to the following defining relations:

$$\begin{aligned} [H_i, H_j] &= 0, \quad [H_i, X_j^\pm] = \pm a_{ij} X_j^\pm, \quad X_i^+ X_j^- - X_j^- X_i^+ = \delta_{i,j} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}, \\ \sum_{r=0}^{1-a_{ij}} (-1)^r \begin{bmatrix} 1 - a_{ij} \\ r \end{bmatrix}_{q_i} (X_i^\pm)^{1-a_{ij}-r} X_j^\pm (X_i^\pm)^r &= 0, \quad i \neq j, \end{aligned}$$

where

$$\begin{aligned} K_i &= e^{d_i h H_i}, \quad e^h = q, \quad q_i = q^{d_i} = e^{d_i h}, \\ \begin{bmatrix} m \\ n \end{bmatrix}_q &= \frac{[m]_q!}{[n]_q! [n-m]_q!}, \quad [n]_q! = [n]_q \cdots [1]_q, \quad [n]_q = \frac{q^n - q^{-n}}{q - q^{-1}}, \end{aligned}$$

with comultiplication defined by

$$\Delta_h(H_i) = H_i \otimes 1 + 1 \otimes H_i, \quad \Delta_h(X_i^+) = X_i^+ \otimes K_i^{-1} + 1 \otimes X_i^+, \quad \Delta_h(X_i^-) = X_i^- \otimes 1 + K_i \otimes X_i^-,$$

antipode defined by

$$S_h(H_i) = -H_i, \quad S_h(X_i^+) = -X_i^+ K_i, \quad S_h(X_i^-) = -K_i^{-1} X_i^-,$$

and counit defined by

$$\varepsilon_h(H_i) = \varepsilon_h(X_i^\pm) = 0.$$

We shall also use the weight-type generators

$$Y_i = \sum_{j=1}^l d_i(a^{-1})_{ij} H_j.$$

Let $L_i^{\pm 1} = e^{\pm h Y_i}$.

Now we shall explicitly describe a linear basis for $U_h(\mathfrak{g})$. First following [2] we recall the construction of root vectors of $U_h(\mathfrak{g})$ in terms of a braid group action on $U_h(\mathfrak{g})$. Let m_{ij} , $i \neq j$ be equal to 2, 3, 4, 6 if $a_{ij} a_{ji}$ is equal to 0, 1, 2, 3, respectively. The braid group $\mathcal{B}_{\mathfrak{g}}$ associated to \mathfrak{g} has generators T_i , $i = 1, \dots, l$, and defining relations

$$T_i T_j T_i T_j \dots = T_j T_i T_j T_i \dots$$

for all $i \neq j$, where there are m_{ij} T 's on each side of the equation.

$\mathcal{B}_{\mathfrak{g}}$ acts by algebra automorphisms of $U_h(\mathfrak{g})$ as follows:

$$T_i(X_j^+) = -X_j^- e^{hd_i H_i}, \quad T_i(X_j^-) = -e^{-hd_i H_i} X_j^+, \quad T_i(H_j) = H_j - a_{ji} H_i,$$

$$T_i(X_j^+) = \sum_{r=0}^{-a_{ij}} (-1)^{r-a_{ij}} q_i^{-r} (X_i^+)^{(-a_{ij}-r)} X_j^+ (X_i^+)^{(r)}, \quad i \neq j,$$

$$T_i(X_j^-) = \sum_{r=0}^{-a_{ij}} (-1)^{r-a_{ij}} q_i^r (X_i^-)^{(r)} X_j^- (X_i^-)^{(-a_{ij}-r)}, \quad i \neq j,$$

where

$$(X_i^+)^{(r)} = \frac{(X_i^+)^r}{[r]_{q_i}!}, \quad (X_i^-)^{(r)} = \frac{(X_i^-)^r}{[r]_{q_i}!}, \quad r \geq 0, \quad i = 1, \dots, l.$$

Recall that an ordering of a set of positive roots Δ_+ is called normal if for any three roots α , β , γ such that $\gamma = \alpha + \beta$ we have either $\alpha < \gamma < \beta$ or $\beta < \gamma < \alpha$.

For any reduced decomposition $w_0 = s_{i_1} \dots s_{i_D}$ of the longest element w_0 of the Weyl group W of \mathfrak{g} the ordering

$$\beta_1 = \alpha_{i_1}, \beta_2 = s_{i_1} \alpha_{i_2}, \dots, \beta_D = s_{i_1} \dots s_{i_{D-1}} \alpha_{i_D}$$

is a normal ordering in Δ_+ , and there is a one-to-one correspondence between normal orderings of Δ_+ and reduced decompositions of w_0 (see [19]).

Fix a reduced decomposition $w_0 = s_{i_1} \dots s_{i_D}$ of w_0 and define the corresponding root vectors in $U_h(\mathfrak{g})$ by

$$(3.1) \quad X_{\beta_k}^\pm = T_{i_1} \dots T_{i_{k-1}} X_{i_k}^\pm.$$

Note that one can construct root vectors in the Lie algebra \mathfrak{g} in a similar way. Namely, if $X_{\pm \alpha_i}$ are simple root vectors of \mathfrak{g} then one can introduce an action of the braid group $\mathcal{B}_{\mathfrak{g}}$ by algebra

automorphisms of \mathfrak{g} defined on the standard generators as follows:

$$T_i(X_{\pm\alpha_i}) = -X_{\mp\alpha_i}, \quad T_i(H_j) = H_j - a_{ji}H_i,$$

$$T_i(X_{\alpha_j}) = \frac{1}{(-a_{ij})!} \text{ad}_{X_{\alpha_i}}^{-a_{ij}} X_{\alpha_j}, \quad i \neq j,$$

$$T_i(X_{-\alpha_j}) = \frac{(-1)^{a_{ij}}}{(-a_{ij})!} \text{ad}_{X_{-\alpha_i}}^{-a_{ij}} X_{-\alpha_j}, \quad i \neq j.$$

Now the root vectors $X_{\pm\beta_k} \in \mathfrak{g}_{\pm\beta_k}$ of \mathfrak{g} can be defined by

$$(3.2) \quad X_{\pm\beta_k} = T_{i_1} \dots T_{i_{k-1}} X_{\pm\alpha_{i_k}}.$$

The root vectors X_{β}^- satisfy the following relations:

$$(3.3) \quad X_{\alpha}^- X_{\beta}^- - q^{(\alpha, \beta)} X_{\beta}^- X_{\alpha}^- = \sum_{\alpha < \delta_1 < \dots < \delta_n < \beta} C(k_1, \dots, k_n) (X_{\delta_1}^-)^{k_1} (X_{\delta_2}^-)^{k_2} \dots (X_{\delta_n}^-)^{k_n}, \quad \alpha < \beta,$$

where $C(k_1, \dots, k_n) \in \mathcal{P}$, and $\mathcal{P} = \mathbb{C}[q, q^{-1}]$ if \mathfrak{g} is simply-laced, $\mathcal{P} = \mathbb{C}[q, q^{-1}, \frac{1}{[2]_q}]$ if \mathfrak{g} is of type B_l, C_l or F_4 , and $\mathcal{P} = \mathbb{C}[q, q^{-1}, \frac{1}{[2]_q}, \frac{1}{[3]_q}]$ if \mathfrak{g} is of type G_2 .

Note that by construction

$$X_{\beta}^+ \pmod{h} = X_{\beta} \in \mathfrak{g}_{\beta},$$

$$X_{\beta}^- \pmod{h} = X_{-\beta} \in \mathfrak{g}_{-\beta}$$

are root vectors of \mathfrak{g} .

$U_h(\mathfrak{g})$ is a quasitriangular Hopf algebra, i.e. there exists an invertible element $\mathcal{R} \in U_h(\mathfrak{g}) \otimes U_h(\mathfrak{g})$, called a universal R-matrix, such that

$$(3.4) \quad \Delta_h^{opp}(a) = \mathcal{R} \Delta_h(a) \mathcal{R}^{-1} \text{ for all } a \in U_h(\mathfrak{g}).$$

An explicit expression for \mathcal{R} may be written by making use of the q -exponential

$$\text{exp}_q(x) = \sum_{k=0}^{\infty} q^{\frac{1}{2}k(k+1)} \frac{x^k}{[k]_q!}$$

in terms of which the element \mathcal{R} takes the form (see e.g. [2], Theorem 8.3.9):

$$(3.5) \quad \mathcal{R} = \prod_{\beta} \text{exp}_{q_{\beta}}[(1 - q_{\beta}^{-2})X_{\beta}^- \otimes X_{\beta}^+] \text{exp} \left[h \sum_{i=1}^l (Y_i \otimes H_i) \right],$$

where the product is over all the positive roots of \mathfrak{g} , and the order of the terms is such that the α -term appears to the left of the β -term if $\alpha < \beta$ with respect to the normal ordering of Δ_+ .

One can calculate the action of the comultiplication on the root vectors $X_{\beta_k}^{\pm}$ in terms of the universal R-matrix. For instance for $\Delta_h(X_{\beta_k}^-)$ one has (see e.g. [2], Theorem 8.3.7)

$$(3.6) \quad \Delta_h(X_{\beta_k}^-) = \tilde{\mathcal{R}}_{<\beta_k} (X_{\beta_k}^- \otimes 1 + e^{h\beta^{\vee}} \otimes X_{\beta_k}^-) \tilde{\mathcal{R}}_{<\beta_k}^{-1},$$

where

$$\tilde{\mathcal{R}}_{<\beta_k} = \tilde{\mathcal{R}}_{\beta_1} \dots \tilde{\mathcal{R}}_{\beta_{k-1}}, \quad \tilde{\mathcal{R}}_{\beta_r} = \text{exp}_{q_{\beta_r}}[(1 - q_{\beta_r}^{-2})X_{\beta_r}^+ \otimes X_{\beta_r}^-].$$

4. REALIZATIONS OF QUANTUM GROUPS ASSOCIATED TO WEYL GROUP ELEMENTS

Some important ingredients that will be used in the proof of the main statement in Section 7 are certain subalgebras of the quantum group. These subalgebras are defined in terms of realizations of the algebra $U_{\hbar}(\mathfrak{g})$ associated to Weyl group elements. Following [16] we introduce these realizations in this section.

Let s be an element of the Weyl group W of the pair $(\mathfrak{g}, \mathfrak{h})$, and \mathfrak{h}' the orthogonal complement in \mathfrak{h} , with respect to the Killing form, to the subspace of \mathfrak{h} fixed by the natural action of s on \mathfrak{h} . Let \mathfrak{h}'^* be the image of \mathfrak{h}' in \mathfrak{h}^* under the identification $\mathfrak{h}^* \simeq \mathfrak{h}$ induced by the canonical bilinear form on \mathfrak{g} . The restriction of the natural action of s on \mathfrak{h}^* to the subspace \mathfrak{h}'^* has no fixed points. Therefore one can define the Cayley transform $\frac{1+s}{1-s}P_{\mathfrak{h}'^*} : \mathfrak{h}^* \rightarrow \mathfrak{h}'^* \subset \mathfrak{h}^*$ of the restriction of s to \mathfrak{h}'^* , where $P_{\mathfrak{h}'^*}$ is the orthogonal projection operator onto \mathfrak{h}'^* in \mathfrak{h}^* , with respect to the Killing form.

Recall that in the classification theory of conjugacy classes in the Weyl group W of the complex simple Lie algebra \mathfrak{g} the so-called primitive (or semi-Coxeter in another terminology) elements play a primary role. According to the results of [1] the element s of the Weyl group of the pair $(\mathfrak{g}, \mathfrak{h})$ is primitive in the Weyl group W' of a regular semisimple Lie subalgebra $\mathfrak{g}' \subset \mathfrak{g}$, $\text{rank } \mathfrak{g}' = \dim \mathfrak{h}'$, of the form

$$\mathfrak{g}' = \mathfrak{h}' + \sum_{\alpha \in \Delta'} \mathfrak{g}_{\alpha},$$

where Δ' is a root subsystem of the root system Δ of \mathfrak{g} , \mathfrak{g}_{α} is the root subspace of \mathfrak{g} corresponding to root α .

Moreover, by Theorem C in [1] s can be represented as a product of two involutions,

$$(4.1) \quad s = s^1 s^2,$$

where $s^1 = s_{\gamma_1} \dots s_{\gamma_n}$, $s^2 = s_{\gamma_{n+1}} \dots s_{\gamma_{l'}}$, the roots in each of the sets $\gamma_1, \dots, \gamma_n$ and $\gamma_{n+1}, \dots, \gamma_{l'}$ are positive and mutually orthogonal, and the roots $\gamma_1, \dots, \gamma_{l'}$ form a linear basis of \mathfrak{h}'^* , in particular l' is the rank of \mathfrak{g}' . The scalar products $\left(\frac{1+s}{1-s} P_{\mathfrak{h}'^*} \gamma_i, \gamma_j \right)$ can be computed as follows.

Lemma 4.1. ([16], **Lemma 6.2**) *Let $P_{\mathfrak{h}'^*}$ be the orthogonal projection operator onto \mathfrak{h}'^* in \mathfrak{h}^* , with respect to the Killing form. Then the scalar products $\left(\frac{1+s}{1-s} P_{\mathfrak{h}'^*} \gamma_i, \gamma_j \right)$ are of the form:*

$$(4.2) \quad \left(\frac{1+s}{1-s} P_{\mathfrak{h}'^*} \gamma_i, \gamma_j \right) = \varepsilon_{ij}(\gamma_i, \gamma_j),$$

where

$$\varepsilon_{ij} = \begin{cases} -1 & i < j \\ 0 & i = j \\ 1 & i > j \end{cases}.$$

Let γ_i^* , $i = 1, \dots, l'$ be the basis of \mathfrak{h}'^* dual to γ_i , $i = 1, \dots, l'$ with respect to the restriction of the bilinear form (\cdot, \cdot) to \mathfrak{h}'^* . Since the numbers (γ_i, γ_j) are integer each element γ_i^* has the form $\gamma_i^* = \sum_{j=1}^{l'} m_{ij} \gamma_j$, where $m_{ij} \in \mathbb{Q}$. Therefore by Lemma 4.1 and using for simple roots α_i the decomposition of the form $P_{\mathfrak{h}'^*} \alpha_i = \sum_{p=1}^{l'} (\alpha_i, \gamma_p) \gamma_p^* = \sum_{p,q=1}^{l'} (\alpha_i, \gamma_p) m_{pq} \gamma_q$ we deduce that the numbers

$$(4.3) \quad q_{ij} = \frac{1}{d_j} \left(\frac{1+s}{1-s} P_{\mathfrak{h}'^*} \alpha_i, \alpha_j \right) = \frac{1}{d_j} \sum_{k,l,p,q=1}^{l'} (\gamma_k, \alpha_i) (\gamma_l, \alpha_j) \left(\frac{1+s}{1-s} P_{\mathfrak{h}'^*} \gamma_p, \gamma_q \right) m_{kp} m_{lq}, \quad i, j = 1, \dots, l$$

are rational as all factors in the products in the sum in the right hand side are rational. Fix a positive integer d such that $q_{ij} \in \frac{1}{d}\mathbb{Z}$ for any $i < j$ or for any $i > j$, $i, j = 1, \dots, l$.

Let n be a non-zero integer number, and let $U_h^s(\mathfrak{g})$ be the topological algebra over $\mathbb{C}[[h]]$ topologically generated by elements e_i, f_i, H_i , $i = 1, \dots, l$ subject to the relations:

$$[H_i, H_j] = 0, [H_i, e_j] = a_{ij}e_j, [H_i, f_j] = -a_{ij}f_j, e_i f_j - q^{c_{ij}} f_j e_i = \delta_{i,j} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}},$$

$$c_{ij} = nd \left(\frac{1+s}{1-s} P_{\mathfrak{b}'^*} \alpha_i, \alpha_j \right), K_i = e^{d_i h H_i},$$

$$\sum_{r=0}^{1-a_{ij}} (-1)^r q^{rc_{ij}} \begin{bmatrix} 1-a_{ij} \\ r \end{bmatrix}_{q_i} (e_i)^{1-a_{ij}-r} e_j (e_i)^r = 0, i \neq j,$$

$$\sum_{r=0}^{1-a_{ij}} (-1)^r q^{rc_{ij}} \begin{bmatrix} 1-a_{ij} \\ r \end{bmatrix}_{q_i} (f_i)^{1-a_{ij}-r} f_j (f_i)^r = 0, i \neq j.$$

Note that the matrix c_{ij} is skew-symmetric.

Proposition 4.2. ([16], **Theorem 4.1**) *For every solution $n_{ij} \in \mathbb{C}$, $i, j = 1, \dots, l$ of equations*

$$(4.4) \quad d_j n_{ij} - d_i n_{ji} = c_{ij}$$

there exists an algebra isomorphism $\psi_{\{n_{ij}\}} : U_h^s(\mathfrak{g}) \rightarrow U_h(\mathfrak{g})$ defined by the formulas:

$$\psi_{\{n_{ij}\}}(e_i) = X_i^+ \prod_{p=1}^l e^{hn_{ip} Y_p}, \quad \psi_{\{n_{ij}\}}(f_i) = \prod_{p=1}^l e^{-hn_{ip} Y_p} X_i^-, \quad \psi_{\{n_{ij}\}}(H_i) = H_i, \quad i = 1, \dots, l.$$

The algebra $U_h^s(\mathfrak{g})$ is called the realization of the quantum group $U_h(\mathfrak{g})$ corresponding to the element $s \in W$.

Remark 4.3. *Let $n'_{ij} \in \mathbb{C}$ be a solution of the homogeneous system that corresponds to (4.4),*

$$(4.5) \quad d_i n'_{ji} - d_j n'_{ij} = 0.$$

Then the map defined by

$$(4.6) \quad \begin{aligned} X_i^+ &\mapsto X_i^+ \prod_{p=1}^l e^{hn'_{ip} Y_p}, \\ X_i^- &\mapsto \prod_{p=1}^l e^{-hn'_{ip} Y_p} X_i^-, \\ H_i &\mapsto H_i \end{aligned}$$

is an automorphism of $U_h(\mathfrak{g})$. Therefore for given element $s \in W$ the isomorphism $\psi_{\{n_{ij}\}}$ is defined uniquely up to automorphisms (4.6) of $U_h(\mathfrak{g})$.

The realizations $U_h^s(\mathfrak{g})$ of the quantum group $U_h(\mathfrak{g})$ are connected with quantizations of some non-standard bialgebra structures on \mathfrak{g} . At the quantum level changing bialgebra structure corresponds to the so-called Drinfeld twist (see [16], Section 4).

Equip $U_h^s(\mathfrak{g})$ with the comultiplication Δ_s given by

$$\Delta_s(H_i) = H_i \otimes 1 + 1 \otimes H_i,$$

$$\Delta_s(e_i) = e_i \otimes e^{-hd_i H_i} + e^{hd_i nd \frac{1+s}{1-s} P_{\mathfrak{b}'^*} H_i} \otimes e_i, \quad \Delta_s(f_i) = f_i \otimes 1 + e^{-hd_i nd \frac{1+s}{1-s} P_{\mathfrak{b}'^*} H_i + hd_i H_i} \otimes f_i,$$

the antipode $S_s(x)$ given by

$$S_s(e_i) = -e^{-hd_i nd \frac{1+s}{1-s} P_{\mathfrak{b}'^*} H_i} e_i e^{hd_i H_i}, \quad S_s(f_i) = -e^{hd_i nd \frac{1+s}{1-s} P_{\mathfrak{b}'^*} H_i - hd_i H_i} f_i, \quad S_s(H_i) = -H_i,$$

and counit defined by

$$\varepsilon_s(H_i) = \varepsilon_s(X_i^\pm) = 0.$$

The comultiplication Δ_s is obtained from the standard comultiplication by a Drinfeld twist (see [16], Section 4). Namely, let

$$(4.7) \quad \mathcal{F} = \exp\left(-h \sum_{i,j=1}^l \frac{n_{ij}}{d_i} Y_i \otimes Y_j\right) \in U_h(\mathfrak{h}) \otimes U_h(\mathfrak{h}),$$

where n_{ij} is the solution of the corresponding equation (4.4) used in the definition of the isomorphism $\psi_{\{n_{ij}\}}$. Then

$$(4.8) \quad \Delta_s(a) = (\psi_{\{n_{ij}\}}^{-1} \otimes \psi_{\{n_{ij}\}}^{-1}) \mathcal{F} \Delta(\psi_{\{n_{ij}\}}(a)) \mathcal{F}^{-1}.$$

We shall construct analogues of root vectors for $U_h^s(\mathfrak{g})$. It is convenient to introduce an operator $K \in \text{End } \mathfrak{h}$ defined by

$$(4.9) \quad KH_i = \sum_{j=1}^l \frac{n_{ij}}{d_i} Y_j.$$

For any solution of equation (4.4) and any normal ordering of the root system Δ_+ we define the following elements of $U_h^s(\mathfrak{g})$, $e_\beta = \psi_{\{n\}}^{-1}(X_\beta^+ e^{hK\beta^\vee})$, $f_\beta = \psi_{\{n\}}^{-1}(e^{-hK\beta^\vee} X_\beta^-)$, $\beta \in \Delta_+$.

$U_h^s(\mathfrak{g})$ is a quasitriangular topological Hopf algebra with the universal R-matrix \mathcal{R}^s ,

$$(4.10) \quad \mathcal{R}^s = \prod_{\beta} \exp_{q_\beta} \left[(1 - q_\beta^{-2}) f_\beta \otimes e_\beta e^{-hnd \frac{1+s}{1-s} P_{\mathfrak{h}'\beta^\vee}} \right] \times \\ \exp \left[h \left(\sum_{i=1}^l (Y_i \otimes H_i) - \sum_{i=1}^l nd \frac{1+s}{1-s} P_{\mathfrak{h}'\beta^\vee} H_i \otimes Y_i \right) \right],$$

where $P_{\mathfrak{h}'}$ is the orthogonal projection operator onto \mathfrak{h}' in \mathfrak{h} with respect to the Killing form, and the order of the terms is such that the α -term appears to the left of the β -term if $\alpha < \beta$ with respect to the normal ordering of Δ_+ .

Using formula (3.6) and Proposition 4.3 in [16] one can also find that

$$(4.11) \quad \Delta_s(f_{\beta_k}) = \tilde{\mathcal{R}}_{<\beta_k}^s (e^{-hnd \frac{1+s}{1-s} P_{\mathfrak{h}'\beta^\vee} + h\beta^\vee} \otimes f_{\beta_k} + f_{\beta_k} \otimes 1) (\tilde{\mathcal{R}}_{<\beta_k}^s)^{-1},$$

where

$$\tilde{\mathcal{R}}_{<\beta_k}^s = \tilde{\mathcal{R}}_{\beta_1}^s \dots \tilde{\mathcal{R}}_{\beta_{k-1}}^s, \quad \tilde{\mathcal{R}}_{\beta_r}^s = \exp_{q_{\beta_r}} \left[(1 - q_{\beta_r}^{-2}) e_{\beta_r} e^{-hnd \frac{1+s}{1-s} P_{\mathfrak{h}'\beta^\vee}} \otimes f_{\beta_r} \right],$$

and

$$(\tilde{\mathcal{R}}_{<\beta_k}^s)^{-1} = (\tilde{\mathcal{R}}_{\beta_{k-1}}^s)^{-1} \dots (\tilde{\mathcal{R}}_{\beta_1}^s)^{-1}, \quad (\tilde{\mathcal{R}}_{\beta_r}^s)^{-1} = \exp_{q_{\beta_r}^{-1}} \left[(1 - q_{\beta_r}^2) e_{\beta_r} e^{-hnd \frac{1+s}{1-s} P_{\mathfrak{h}'\beta^\vee}} \otimes f_{\beta_r} \right].$$

We shall actually need not the algebras $U_h(\mathfrak{g})$ and $U_h^s(\mathfrak{g})$ themselves but some their specializations defined over certain rings and over the field of complex numbers. They are similar to the non-restricted integral form and to its specialization for the standard quantum group $U_h(\mathfrak{g})$. The results below are slight modifications of similar statements for $U_h(\mathfrak{g})$, and we refer to [2], Ch. 9 for the proofs.

Note that by the choice of d we have $q^{c_{ij}} \in \mathbb{C}[q, q^{-1}]$. Let $\mathcal{A} = \mathbb{C}[q, q^{-1}]$ if \mathfrak{g} is simply-laced, $\mathcal{A} = \mathbb{C}[q, q^{-1}, \frac{1}{|2|_q}]$ if \mathfrak{g} is of type B_l, C_l or F_4 , and $\mathcal{A} = \mathbb{C}[q, q^{-1}, \frac{1}{|2|_q}, \frac{1}{|3|_q}]$ if \mathfrak{g} is of type G_2 .

Let $U_{\mathcal{A}}^s(\mathfrak{g})$ be the \mathcal{A} -subalgebra of $U_h^s(\mathfrak{g})$ generated by the elements $e_i, f_i, L_i^{\pm 1}$, $\frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}$, $i = 1, \dots, l$ and $U_{\mathcal{A}}(\mathfrak{g})$ the \mathcal{A} -subalgebra of $U_h(\mathfrak{g})$ generated by the elements $X_i^\pm, L_i^{\pm 1}$, $\frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}$, $i = 1, \dots, l$.

Remark 4.4. Note that the general solution of equation (4.4) is given by

$$(4.12) \quad n_{ij} = \frac{1}{2d_j}(c_{ij} + s_{ij}),$$

where $s_{ij} = s_{ji}$. If $q_{ij} \in \frac{1}{d}\mathbb{Z}$ for any $i < j$, we put

$$s_{ij} = \begin{cases} c_{ij} & i < j \\ 0 & i = j \\ -c_{ij} & i > j \end{cases}.$$

Then

$$n_{ij} = \begin{cases} \frac{1}{d_j}c_{ij} & i < j \\ 0 & i = j \\ 0 & i > j \end{cases}.$$

By the choice of d we have $c_{ij} \in d_j n\mathbb{Z}$, and hence $n_{ij} \in n\mathbb{Z}$ for $i, j = 1, \dots, l$. A similar consideration shows that if $q_{ij} \in \frac{1}{d}\mathbb{Z}$ for any $i > j$ then there exists a solution $n_{ij} \in n\mathbb{Z}$ for $i, j = 1, \dots, l$.

Since there is a solution $n_{ij} \in n\mathbb{Z}$ for $i, j = 1, \dots, l$ to equation (4.4) the map $\psi_{\{n_{ij}\}}$ associated to this solution in Proposition 4.2 induces an isomorphism of algebras $\psi_{\{n_{ij}\}} : U_{\mathcal{A}}^s(\mathfrak{g}) \rightarrow U_{\mathcal{A}}(\mathfrak{g})$.

All algebras introduced above are Hopf algebras with comultiplications induced from $U_h^s(\mathfrak{g})$ or $U_h(\mathfrak{g})$. The algebra $U_{\mathcal{A}}^s(\mathfrak{g})$ acts on itself by the left and the right adjoint representations,

$$(4.13) \quad \text{Ad}'x(y) = x_1 y S_s(x_2), \quad \text{Ad}x(y) = S_s(x_1) y x_2,$$

where we use the Sweedler notation for the comultiplication, $\Delta_s(x) = x_1 \otimes x_2$.

Denote by $U_{\mathcal{A}}(\mathfrak{n}_+)$, $U_{\mathcal{A}}(\mathfrak{n}_-)$ the subalgebras of $U_{\mathcal{A}}(\mathfrak{g})$ generated by the X_i^+ and by the X_i^- , respectively. For any $\alpha \in \Delta_+$ and the quantum root vectors X_{α}^{\pm} constructed with the help of any normal ordering in Δ_+ one has $X_{\alpha}^{\pm} \in U_{\mathcal{A}}(\mathfrak{n}_{\pm})$.

Let $U_{\mathcal{A}}^s(\mathfrak{n}_+)$, $U_{\mathcal{A}}^s(\mathfrak{n}_-)$ be the subalgebras of $U_{\mathcal{A}}^s(\mathfrak{g})$ generated by the e_i and by the f_i , $i = 1, \dots, l$, respectively.

From the definition of the quantum root vectors X_{β}^{\pm} and from the commutation relations between the generators X_i^{\pm} and $L_j^{\pm 1}$, $i, j = 1, \dots, l$ it follows that for the solution of equation (4.4) introduced in Remark 4.4 and any normal ordering of the root system Δ_+ the elements $e_{\beta} = \psi_{\{n_{ij}\}}^{-1}(X_{\beta}^+ e^{hK\beta^{\vee}})$ and $f_{\beta} = \psi_{\{n_{ij}\}}^{-1}(e^{-hK\beta^{\vee}} X_{\beta}^-)$, $\beta \in \Delta_+$ lie in the subalgebras $U_{\mathcal{A}}^s(\mathfrak{n}_+)$ and $U_{\mathcal{A}}^s(\mathfrak{n}_-)$, respectively.

The most important for us are the specializations $U_{\varepsilon}(\mathfrak{g})$ and $U_{\varepsilon}^s(\mathfrak{g})$ of $U_{\mathcal{A}}(\mathfrak{g})$ and $U_{\mathcal{A}}^s(\mathfrak{g})$, $U_{\varepsilon}(\mathfrak{g}) = U_{\mathcal{A}}(\mathfrak{g})/(q - \varepsilon)U_{\mathcal{A}}(\mathfrak{g})$, $U_{\varepsilon}^s(\mathfrak{g}) = U_{\mathcal{A}}^s(\mathfrak{g})/(q - \varepsilon)U_{\mathcal{A}}^s(\mathfrak{g})$, where $\varepsilon \in \mathbb{C}^*$, $\varepsilon^{2d_i} \neq 1$, $i = 1, \dots, l$ and $\varepsilon^4 \neq 1$ if \mathfrak{g} is of type G_2 .

Note that all specializations introduced above are invariant under the action of the braid group $\mathcal{B}_{\mathfrak{g}}$.

Denote by $U_{\varepsilon}(\mathfrak{n}_+)$, $U_{\varepsilon}(\mathfrak{n}_-)$ and $U_{\varepsilon}(\mathfrak{h})$ the subalgebras of $U_{\varepsilon}(\mathfrak{g})$ generated by the X_i^+ , by the X_i^- , and by the $L_i^{\pm 1}$, respectively.

Fix a normal ordering in Δ_+ and let X_{α}^{\pm} be the corresponding quantum root vectors. Define for $\mathbf{r} = (r_1, \dots, r_D) \in \mathbb{N}^D$,

$$(X^+)_{\mathbf{r}} = (X_{\beta_1}^+)^{r_1} \dots (X_{\beta_D}^+)^{r_D},$$

$$(X^-)_{\mathbf{r}} = (X_{\beta_D}^-)^{r_D} \dots (X_{\beta_1}^-)^{r_1},$$

and for $\mathbf{s} = (s_1, \dots, s_l) \in \mathbb{Z}^l$,

$$L^{\mathbf{s}} = L_1^{s_1} \dots L_l^{s_l}.$$

Proposition 4.5. ([2], **Proposition 9.2.2**) *The algebra $U_\varepsilon(\mathfrak{g})$ is generated over \mathbb{C} by $L_i^{\pm 1}$, X_i^\pm , $i = 1, \dots, l$.*

The elements $(X^+)^{\mathbf{r}}$, $(X^-)^{\mathbf{t}}$ and $L^{\mathbf{s}}$, for $\mathbf{r}, \mathbf{t} \in \mathbb{N}^D$, $\mathbf{s} \in \mathbb{Z}^l$, form linear bases of $U_\varepsilon(\mathfrak{n}_+)$, $U_\varepsilon(\mathfrak{n}_-)$ and $U_\varepsilon(\mathfrak{h})$, respectively, and the products $(X^+)^{\mathbf{r}}L^{\mathbf{s}}(X^-)^{\mathbf{t}}$ form a basis of $U_\varepsilon(\mathfrak{g})$. In particular, multiplication defines an isomorphism of vector spaces:

$$U_\varepsilon(\mathfrak{n}_-) \otimes U_\varepsilon(\mathfrak{h}) \otimes U_\varepsilon(\mathfrak{n}_+) \rightarrow U_\varepsilon(\mathfrak{g}).$$

The root vectors X_β^- satisfy the following relations in $U_\varepsilon(\mathfrak{g})$:

$$(4.14) \quad X_\alpha^- X_\beta^- - \varepsilon^{(\alpha, \beta)} X_\beta^- X_\alpha^- = \sum_{\alpha < \delta_1 < \dots < \delta_n < \beta} C(k_1, \dots, k_n) (X_{\delta_n}^-)^{k_n} (X_{\delta_{n-1}}^-)^{k_{n-1}} \dots (X_{\delta_1}^-)^{k_1},$$

where $\alpha < \beta$, the sum is taken over tuples of roots $\delta_1, \dots, \delta_n$ such that $\alpha < \delta_1 < \dots < \delta_n < \beta$, and over $k_i \in \mathbb{N}$, $C(k_1, \dots, k_n) \in \mathbb{C}$, and for each term in the right hand side $\sum_{i=1}^n k_i \delta_i = \alpha + \beta$.

Now we shall study the algebraic structure of $U_\varepsilon^s(\mathfrak{g})$. Denote by $U_\varepsilon^s(\mathfrak{n}_\pm)$ the subalgebra in $U_\varepsilon^s(\mathfrak{g})$ generated by e_i (f_i), $i = 1, \dots, l$. Let $U_\varepsilon^s(\mathfrak{h})$ be the subalgebra in $U_\varepsilon^s(\mathfrak{g})$ generated by $L_i^{\pm 1}$, $i = 1, \dots, l$.

We shall construct a Poincaré–Birkhoff–Witt basis for $U_\varepsilon^s(\mathfrak{g})$.

Proposition 4.6. (i) *For any normal ordering of the root system Δ_+ and for any integer valued solution of equation (4.4) the elements e_β and f_β lie in the subalgebras $U_\varepsilon^s(\mathfrak{n}_+)$ and $U_\varepsilon^s(\mathfrak{n}_-)$, respectively. The elements f_β , $\beta \in \Delta_+$ satisfy the following commutation relations*

$$(4.15) \quad f_\alpha f_\beta - \varepsilon^{(\alpha, \beta) + nd(\frac{1+s}{1-s} P_{\mathfrak{h}^*} \alpha, \beta)} f_\beta f_\alpha = \sum_{\alpha < \delta_1 < \dots < \delta_n < \beta} C'(k_1, \dots, k_n) f_{\delta_n}^{k_n} f_{\delta_{n-1}}^{k_{n-1}} \dots f_{\delta_1}^{k_1}, \quad \alpha < \beta,$$

where $C'(k_1, \dots, k_n) \in \mathbb{C}$.

(ii) *Moreover, the elements $(e)^\mathbf{r} = (e_{\beta_1})^{r_1} \dots (e_{\beta_D})^{r_D}$, $(f)^\mathbf{t} = (f_{\beta_D})^{t_D} \dots (f_{\beta_1})^{t_1}$ and $L^{\mathbf{s}} = L_1^{s_1} \dots L_l^{s_l}$ for $\mathbf{r}, \mathbf{t} \in \mathbb{N}^D$, $\mathbf{s} \in \mathbb{Z}^l$ form bases of $U_\varepsilon^s(\mathfrak{n}_+)$, $U_\varepsilon^s(\mathfrak{n}_-)$ and $U_\varepsilon^s(\mathfrak{h})$, and the products $(f)^\mathbf{t} L^{\mathbf{s}} (e)^\mathbf{r}$ form a basis of $U_\varepsilon^s(\mathfrak{g})$. In particular, multiplication defines an isomorphism of vector spaces,*

$$U_\varepsilon^s(\mathfrak{n}_-) \otimes U_\varepsilon^s(\mathfrak{h}) \otimes U_\varepsilon^s(\mathfrak{n}_+) \rightarrow U_\varepsilon^s(\mathfrak{g}).$$

The proof of this proposition is similar to the proof of Proposition 4.2 in [16].

5. NILPOTENT SUBALGEBRAS AND QUANTUM GROUPS

In this section we define the subalgebras of $U_\varepsilon(\mathfrak{g})$ which resemble nilpotent subalgebras in \mathfrak{g} and possess non-trivial characters. We start by recalling the definition of certain normal orderings of root systems associated to Weyl group elements (see [16], Section 5 for more details). The definition of subalgebras of $U_\varepsilon(\mathfrak{g})$ having non-trivial characters will be given in terms of root vectors associated to such normal orderings.

Proposition 5.1. ([16], **Proposition 5.1**) *Let $s \in W$ be an element of the Weyl group W of the pair $(\mathfrak{g}, \mathfrak{h})$, Δ the root system of the pair $(\mathfrak{g}, \mathfrak{h})$. Then there is a system of positive roots Δ_+^s such that decomposition (4.1) is reduced in the sense that $l(s) = l(s^2) + l(s^1)$, where $l(\cdot)$ is the length function in W with respect to the system of simple roots in Δ_+^s , and $\Delta_s^s = \Delta_{s^2}^s \cup s^2(\Delta_{s^1}^s)$, $\Delta_{s^{-1}}^s = \Delta_{s^1}^s \cup s^1(\Delta_{s^2}^s)$ (disjoint unions), $\Delta_{s^{1,2}}^s = \{\alpha \in \Delta_+^s : s^{1,2}\alpha \in -\Delta_+^s\}$, $\Delta_s^s = \{\alpha \in \Delta_+^s : s\alpha \in -\Delta_+^s\}$. Here s^1, s^2 are the involutions entering decomposition (4.1), $s^1 = s_{\gamma_1} \dots s_{\gamma_n}$, $s^2 = s_{\gamma_{n+1}} \dots s_{\gamma_{l'}}$, the roots in each of the sets $\gamma_1, \dots, \gamma_n$ and $\gamma_{n+1}, \dots, \gamma_{l'}$ are positive and mutually orthogonal.*

Moreover, there is a normal ordering of the root system Δ_+^s of the following form

$$(5.1) \quad \begin{aligned} & \beta_1^1, \dots, \beta_t^1, \beta_{t+1}^1, \dots, \beta_{t+\frac{p-n}{2}}^1, \gamma_1, \beta_{t+\frac{p-n}{2}+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_1}^1, \gamma_2, \\ & \beta_{t+\frac{p-n}{2}+n_1+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_2}^1, \gamma_3, \dots, \gamma_n, \beta_{t+p+1}^1, \dots, \beta_{l(s^1)}^1, \dots, \\ & \beta_1^2, \dots, \beta_q^2, \gamma_{n+1}, \beta_{q+2}^2, \dots, \beta_{q+m_1}^2, \gamma_{n+2}, \beta_{q+m_1+2}^2, \dots, \beta_{q+m_2}^2, \gamma_{n+3}, \dots, \\ & \gamma_{l'}, \beta_{q+m_{l(s^2)}+1}^2, \dots, \beta_{2q+2m_{l(s^2)}-(l'-n)}^2, \beta_{2q+2m_{l(s^2)}-(l'-n)+1}^2, \dots, \beta_{l(s^2)}^2, \\ & \beta_1^0, \dots, \beta_{D_0}^0, \end{aligned}$$

where

$$\begin{aligned} & \{\beta_1^1, \dots, \beta_t^1, \beta_{t+1}^1, \dots, \beta_{t+\frac{p-n}{2}}^1, \gamma_1, \beta_{t+\frac{p-n}{2}+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_1}^1, \gamma_2, \\ & \beta_{t+\frac{p-n}{2}+n_1+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_2}^1, \gamma_3, \dots, \gamma_n, \beta_{t+p+1}^1, \dots, \beta_{l(s^1)}^1\} = \Delta_{s^1}^s, \\ & \{\beta_{t+1}^1, \dots, \beta_{t+\frac{p-n}{2}}^1, \gamma_1, \beta_{t+\frac{p-n}{2}+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_1}^1, \gamma_2, \\ & \beta_{t+\frac{p-n}{2}+n_1+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_2}^1, \gamma_3, \dots, \gamma_n\} = \{\alpha \in \Delta_+^s \mid s^1(\alpha) = -\alpha\} = \Delta_{s^1=-1}^s, \\ & \{\beta_1^2, \dots, \beta_q^2, \gamma_{n+1}, \beta_{q+2}^2, \dots, \beta_{q+m_1}^2, \gamma_{n+2}, \beta_{q+m_1+2}^2, \dots, \beta_{q+m_2}^2, \gamma_{n+3}, \dots, \\ & \gamma_{l'}, \beta_{q+m_{l(s^2)}+1}^2, \dots, \beta_{2q+2m_{l(s^2)}-(l'-n)}^2, \beta_{2q+2m_{l(s^2)}-(l'-n)+1}^2, \dots, \beta_{l(s^2)}^2\} = \Delta_{s^2}^s, \\ & \{\gamma_{n+1}, \beta_{q+2}^2, \dots, \beta_{q+m_1}^2, \gamma_{n+2}, \beta_{q+m_1+2}^2, \dots, \beta_{q+m_2}^2, \gamma_{n+3}, \dots, \\ & \gamma_{l'}, \beta_{q+m_{l(s^2)}+1}^2, \dots, \beta_{2q+2m_{l(s^2)}-(l'-n)}^2\} = \{\alpha \in \Delta_+^s \mid s^2(\alpha) = -\alpha\} = \Delta_{s^2=-1}^s, \\ & \{\beta_1^0, \dots, \beta_{D_0}^0\} = \{\alpha \in \Delta_+^s \mid s(\alpha) = \alpha\}. \end{aligned}$$

The length of the ordered segment $\Delta_{m_+} \subset \Delta$ in normal ordering (5.1),

$$(5.2) \quad \begin{aligned} \Delta_{m_+} = & \gamma_1, \beta_{t+\frac{p-n}{2}+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_1}^1, \gamma_2, \beta_{t+\frac{p-n}{2}+n_1+2}^1, \dots, \beta_{t+\frac{p-n}{2}+n_2}^1, \\ & \gamma_3, \dots, \gamma_n, \beta_{t+p+1}^1, \dots, \beta_{l(s^1)}^1, \dots, \beta_1^2, \dots, \beta_q^2, \\ & \gamma_{n+1}, \beta_{q+2}^2, \dots, \beta_{q+m_1}^2, \gamma_{n+2}, \beta_{q+m_1+2}^2, \dots, \beta_{q+m_2}^2, \gamma_{n+3}, \dots, \gamma_{l'}, \end{aligned}$$

is equal to

$$(5.3) \quad D - \left(\frac{l(s) - l'}{2} + D_0 \right),$$


where D is the number of roots in Δ_+^s , $l(s)$ is the length of s and D_0 is the number of positive roots fixed by the action of s .

For any two roots $\alpha, \beta \in \Delta_{m_+}$ such that $\alpha < \beta$ the sum $\alpha + \beta$ cannot be represented as a linear combination $\sum_{k=1}^q c_k \gamma_{i_k}$, where $c_k \in \mathbb{N}$ and $\alpha < \gamma_{i_1} < \dots < \gamma_{i_k} < \beta$.

Remark 5.2. In case when $s = s^1$ is an involution the last root in the segment Δ_{m_+} is the root preceding β_1^0 in normal ordering (5.1).

We shall also need another system of positive roots associated to (the conjugacy class of) the Weyl group element s . In order to define it we need to recall the definition of a circular normal ordering of the root system Δ .

Let $\beta_1, \beta_2, \dots, \beta_D$ be a normal ordering of a positive root system. Then following [11] one can introduce the corresponding circular normal ordering of the root system Δ where the roots in Δ are located on a circle in the following way



Circular normal ordering of a root system.

Let $\alpha, \beta \in \Delta$. One says that the segment $[\alpha, \beta]$ of the circle is minimal if it does not contain the opposite roots $-\alpha$ and $-\beta$ and the root β follows after α on the circle above, the circle being oriented clockwise. In that case one also says that $\alpha < \beta$ in the sense of the circular normal ordering,

$$(5.4) \quad \alpha < \beta \Leftrightarrow \text{the segment } [\alpha, \beta] \text{ of the circle is minimal.}$$

Later we shall need the following property of minimal segments which is a direct consequence of Proposition 3.3 in [10].

Lemma 5.3. *Let $[\alpha, \beta]$ be a minimal segment in a circular normal ordering of a root system Δ . Then if $\alpha + \beta$ is a root we have*

$$\alpha < \alpha + \beta < \beta.$$

Note that any segment in a circular normal ordering of Δ of length equal to the number of positive roots is a system of positive roots.

Now consider the circular normal ordering of Δ corresponding to the system of positive roots Δ_+^s and to its normal ordering introduced in Proposition 5.1. The segment which consists of the roots α satisfying $\gamma_1 \leq \alpha < -\gamma_1$ is a system of positive roots in Δ as its length is equal to the number of positive roots and it is closed under addition of roots by Lemma 5.3.

The system of positive roots Δ_+ introduced this way and equipped with the normal ordering induced by the circular normal ordering is called the normally ordered system of positive roots associated to the (conjugacy class of) the Weyl group element $s \in W$.

The linear subspace of \mathfrak{g} generated by the root vectors X_α ($X_{-\alpha}$), $\alpha \in \Delta_{\mathfrak{m}_+}$ is in fact a Lie subalgebra $\mathfrak{m}_+ \subset \mathfrak{g}$ ($\mathfrak{m}_- \subset \mathfrak{g}$). Let \mathfrak{b}_+ be the Borel subalgebra associated to Δ_+ and \mathfrak{b}_- is the opposite Borel subalgebra. Denote by \mathfrak{n}_\pm the nilradicals of \mathfrak{b}_\pm . Let H, N_+, N_-, B_+, B_- be the Cartan subgroup, the maximal unipotent subgroups and the Borel subgroups of G which correspond to the Lie subalgebras $\mathfrak{h}, \mathfrak{n}_+, \mathfrak{n}_-, \mathfrak{b}_+$ and \mathfrak{b}_- , respectively. Note that by definition $\Delta_{\mathfrak{m}_+} \subset \Delta_+$, and hence $\mathfrak{m}_\pm \subset \mathfrak{n}_\pm$.

Now we can define the subalgebras of $U_\varepsilon(\mathfrak{g})$ which resemble nilpotent subalgebras in \mathfrak{g} and possess non-trivial characters.

Theorem 5.4. *Let $s \in W$ be an element of the Weyl group W of the pair $(\mathfrak{g}, \mathfrak{h})$, Δ the root system of the pair $(\mathfrak{g}, \mathfrak{h})$. Fix a decomposition (4.1) of s and let Δ_+ be a system of positive roots associated*

to s . Assume that $\varepsilon^{2d_i} \neq 1$, $\varepsilon^4 \neq 1$ if \mathfrak{g} is of type G_2 and that $\varepsilon^{nd-1} = 1$, where d and n are introduced in Section 4. Let $U_\varepsilon^s(\mathfrak{g})$ be the realization of the quantum group $U_\varepsilon(\mathfrak{g})$ associated to s . Let $f_\beta \in U_\varepsilon^s(\mathfrak{n}_-)$, $\beta \in \Delta_+$ be the root vectors associated to the corresponding normal ordering of Δ_+ .

Then elements $f_\beta \in U_\varepsilon^s(\mathfrak{n}_-)$, $\beta \in \Delta_{\mathfrak{m}_+}$, where $\Delta_{\mathfrak{m}_+} \subset \Delta$ is ordered segment (5.2), generate a subalgebra $U_\varepsilon^s(\mathfrak{m}_-) \subset U_\varepsilon^s(\mathfrak{g})$. The elements $f^{\mathbf{r}} = f_{\beta_D}^{r_D} \dots f_{\beta_1}^{r_1}$, $r_i \in \mathbb{N}$, $i = 1, \dots, D$ and r_i can be strictly positive only if $\beta_i \in \Delta_{\mathfrak{m}_+}$, form a linear basis of $U_\varepsilon^s(\mathfrak{m}_-)$.

Moreover the map $\chi^s : U_\varepsilon^s(\mathfrak{m}_-) \rightarrow \mathbb{C}$ defined on generators by

$$(5.5) \quad \chi^s(f_\beta) = \begin{cases} 0 & \beta \notin \{\gamma_1, \dots, \gamma_{l'}\} \\ c_i & \beta = \gamma_i, c_i \in \mathbb{C} \end{cases}$$

is a character of $U_\varepsilon^s(\mathfrak{m}_-)$.

Proof. The first statement of the theorem follows straightforwardly from commutation relations (4.15) and Proposition 4.6.

In order to prove that the map $\chi^s : U_\varepsilon^s(\mathfrak{m}_-) \rightarrow \mathbb{C}$ defined by (5.5) is a character of $U_\varepsilon^s(\mathfrak{m}_-)$ we show that all relations (4.15) for f_α, f_β with $\alpha, \beta \in \Delta_{\mathfrak{m}_+}$, which are defining relations in the subalgebra $U_\varepsilon^s(\mathfrak{m}_-)$ by part (ii) of Proposition 4.6, belong to the kernel of χ^s . By definition the only generators of $U_\varepsilon^s(\mathfrak{m}_-)$ on which χ^s may not vanish are f_{γ_i} , $i = 1, \dots, l'$. By the last statement in Proposition 5.1 for any two roots $\alpha, \beta \in \Delta_{\mathfrak{m}_+}$ such that $\alpha < \beta$ the sum $\alpha + \beta$ can not be represented as a linear combination $\sum_{k=1}^q c_k \gamma_{i_k}$, where $c_k \in \mathbb{N}$ and $\alpha < \gamma_{i_1} < \dots < \gamma_{i_q} < \beta$. Hence for any two roots $\alpha, \beta \in \Delta_{\mathfrak{m}_+}$ such that $\alpha < \beta$ the value of the map χ^s on the r.h.s. of the corresponding commutation relation (4.15) is equal to zero.

Therefore it suffices to prove that

$$\chi^s(f_{\gamma_i} f_{\gamma_j} - \varepsilon^{(\gamma_i, \gamma_j) + nd(\frac{1+s}{1-s} P_{\mathfrak{h}'^* \gamma_i, \gamma_j})} f_{\gamma_j} f_{\gamma_i}) = c_i c_j (1 - \varepsilon^{(\gamma_i, \gamma_j) + nd(\frac{1+s}{1-s} P_{\mathfrak{h}'^* \gamma_i, \gamma_j})}) = 0, \quad i < j.$$

Since $\varepsilon^{nd-1} = 1$ and $(\frac{1+s}{1-s} P_{\mathfrak{h}'^* \gamma_i, \gamma_j})$ are integer numbers for any $i, j = 1, \dots, l'$, the last identity always holds provided $(\gamma_i, \gamma_j) + (\frac{1+s}{1-s} P_{\mathfrak{h}'^* \gamma_i, \gamma_j}) = 0$ for $i < j$. As we saw in Lemma 4.1 this is indeed the case. This completes the proof. \square

6. QUANTUM GROUPS AT ROOTS OF UNITY

Let m be an odd positive integer number, and $m > d_i$ is coprime to all d_i for all i , ε a primitive m -th root of unity. In this section, following [2], Section 9.2, we recall some results on the structure of the algebra $U_\varepsilon(\mathfrak{g})$. We keep the notation introduced in Section 2.

Let Z_ε be the center of $U_\varepsilon(\mathfrak{g})$.

Proposition 6.1. ([3], Corollary 3.3, [4], Theorems 3.5, 7.6 and Proposition 4.5) *Fix the normal ordering in the positive root system Δ_+ corresponding a reduced decomposition $w_0 = s_{i_1} \dots s_{i_D}$ of the longest element w_0 of the Weyl group W of \mathfrak{g} and let X_α^\pm be the corresponding root vectors in $U_\varepsilon(\mathfrak{g})$, and X_α the corresponding root vectors in \mathfrak{g} . Let $x_\alpha^- = (\varepsilon_\alpha - \varepsilon_\alpha^{-1})^m (X_\alpha^-)^m$, $x_\alpha^+ = (\varepsilon_\alpha - \varepsilon_\alpha^{-1})^m T_0 (X_\alpha^-)^m$, where $T_0 = T_{i_1} \dots T_{i_D}$, $\alpha \in \Delta_+$ and $l_i = L_i^m$, $i = 1, \dots, l$ be elements of $U_\varepsilon(\mathfrak{g})$.*

Then the following statements are true.

(i) *The elements x_α^\pm , $\alpha \in \Delta_+$, l_i , $i = 1, \dots, l$ lie in Z_ε .*

(ii) *Let Z_0 (Z_0^\pm and Z_0^0) be the subalgebras of Z_ε generated by the x_α^\pm and the $l_i^{\pm 1}$ (respectively by the x_α^\pm and by the $l_i^{\pm 1}$). Then $Z_0^\pm \subset U_\varepsilon(\mathfrak{n}_\pm)$, $Z_0^0 \subset U_\varepsilon(\mathfrak{h})$, Z_0^\pm is the polynomial algebra with generators x_α^\pm , Z_0^0 is the algebra of Laurent polynomials in the l_i , $Z_0^\pm = U_\varepsilon(\mathfrak{n}_\pm) \cap Z_0$, and multiplication defines an isomorphism of algebras*

$$Z_0^- \otimes Z_0^0 \otimes Z_0^+ \rightarrow Z_0.$$

The subalgebra Z_0 is independent of the choice of the reduced decomposition $w_0 = s_{i_1} \dots s_{i_D}$.

(iii) $U_\varepsilon(\mathfrak{g})$ is a free Z_0 -module with basis the set of monomials $(X^+)^{\mathbf{r}}L^{\mathbf{s}}(X^-)^{\mathbf{t}}$ in the statement of Proposition 4.5 for which $0 \leq r_k, t_k, s_i < m$ for $i = 1, \dots, l, k = 1, \dots, D$.

(iv) $\text{Spec}(Z_0) = \mathbb{C}^{2D} \times (\mathbb{C}^*)^l$ is a complex affine space of dimension equal to $\dim \mathfrak{g}$.

(v) The subalgebra Z_0 is preserved by the action of the braid group automorphisms T_i .

(vi) Let G be the connected simply connected Lie group corresponding to the Lie algebra \mathfrak{g} and G_0^* the solvable algebraic subgroup in $G \times G$ which consists of elements of the form $(L_+, L_-) \in G \times G$,

$$(L_+, L_-) = (t, t^{-1})(n_+, n_-), \quad n_\pm \in N_\pm, \quad t \in H.$$

Then $\text{Spec}(Z_0^0)$ can be naturally identified with the maximal torus H in G , and the map

$$\tilde{\pi} : \text{Spec}(Z_0) = \text{Spec}(Z_0^+) \times \text{Spec}(Z_0^0) \times \text{Spec}(Z_0^-) \rightarrow G_0^*,$$

$$\tilde{\pi}(u_+, t, u_-) = (t\mathbf{X}^+(u_+), t^{-1}\mathbf{X}^-(u_-)^{-1}), \quad u_\pm \in \text{Spec}(Z_0^\pm), \quad t \in \text{Spec}(Z_0^0),$$

$$\mathbf{X}^\pm : \text{Spec}(Z_0^\pm) \rightarrow N_\pm,$$

$$\mathbf{X}^- = \exp(x_{\beta_D}^- X_{-\beta_D}) \exp(x_{\beta_{D-1}}^- X_{-\beta_{D-1}}) \dots \exp(x_{\beta_1}^- X_{-\beta_1}),$$

$$\mathbf{X}^+ = \exp(x_{\beta_D}^+ T_0(X_{-\beta_D})) \exp(x_{\beta_{D-1}}^+ T_0(X_{-\beta_{D-1}})) \dots \exp(x_{\beta_1}^+ T_0(X_{-\beta_1})),$$

where $x_{\beta_i}^\pm$ should be regarded as complex-valued functions on $\text{Spec}(Z_0)$, is an isomorphism of varieties independent of the choice of reduced decomposition of w_0 .

From parts (ii) and (iii) of Proposition 6.1 we immediately deduce similar properties for the algebra $U_\varepsilon^s(\mathfrak{g})$.

Proposition 6.2. (i) The subalgebra $\psi_{\{n_{ij}\}}^{-1}(Z_0) \subset U_\varepsilon^s(\mathfrak{g})$ is the tensor product of the polynomial algebra with generators $e_\alpha^m, f_\alpha^m, \alpha \in \Delta_+$ and of the algebra of Laurent polynomials in $l_i, i = 1, \dots, l$.

(ii) $U_\varepsilon^s(\mathfrak{g})$ is a free $\psi_{\{n_{ij}\}}^{-1}(Z_0)$ -module with basis the set of monomials $(f)^{\mathbf{r}}L^{\mathbf{s}}(e)^{\mathbf{t}}$ for which $0 \leq r_k, t_k, s_i < m$ for $i = 1, \dots, l, k = 1, \dots, D$.

Remark 6.3. In fact $\text{Spec}(Z_0)$ carries a natural structure of a Poisson-Lie group, and the map $\tilde{\pi}$ is an isomorphism of algebraic Poisson-Lie groups if G_0^* is regarded as the dual Poisson-Lie group to the Poisson-Lie group G equipped with the standard Sklyanin bracket (see [4], Theorem 7.6). We shall not need this fact in this paper.

Let $\mathbf{K} : \text{Spec}(Z_0^0) \rightarrow H$ be the map defined by $\mathbf{K}(h) = h^2, h \in H$.

Proposition 6.4. ([4], Corollary 4.7) Let $G^0 = N_- H N_+$ be the big cell in G . Then the map

$$\pi = \mathbf{X}^- \mathbf{K} \mathbf{X}^+ : \text{Spec}(Z_0) \rightarrow G^0$$

is independent of the choice of reduced decomposition of w_0 , and is an unramified covering of degree 2^l .

Denote by $\lambda_0 : G_0^* \rightarrow G^0$ the map defined by $\lambda_0(L_+, L_-) = L_-^{-1} L_+$. Then obviously $\pi = \lambda_0 \circ \tilde{\pi}$.

Define derivations \underline{x}_i^\pm of $U_{\mathcal{A}}(\mathfrak{g})$ by

$$(6.1) \quad \underline{x}_i^+(u) = \left[\frac{(X_i^+)^m}{[m]_{q_i}!}, u \right], \quad \underline{x}_i^-(u) = T_0 \underline{x}_i^+ T_0^{-1}(u), \quad i = 1, \dots, l, \quad u \in U_{\mathcal{A}}(\mathfrak{g}).$$

Let \widehat{Z}_0 be the algebra of formal power series in the $x_\alpha^\pm, \alpha \in \Delta_+$, and the $l_i^{\pm 1}, i = 1, \dots, l$, which define holomorphic functions on $\text{Spec}(Z_0) = \mathbb{C}^{2D} \times (\mathbb{C}^*)^l$. Let

$$\widehat{U}_\varepsilon(\mathfrak{g}) = U_\varepsilon(\mathfrak{g}) \otimes_{Z_0} \widehat{Z}_0, \quad \widehat{Z}_\varepsilon = Z_\varepsilon \otimes_{Z_0} \widehat{Z}_0.$$

Proposition 6.5. ([3], Propositions 3.4, 3.5, [4], Proposition 6.1, Theorem 6.6)

(i) On specializing to $q = \varepsilon$, (6.1) induces a well-defined derivation \underline{x}_i^\pm of $U_\varepsilon(\mathfrak{g})$.

(ii) The series

$$\exp(t\underline{x}_i^\pm) = \sum_{k=0}^{\infty} \frac{t^k}{k!} (\underline{x}_i^\pm)^k$$

converge for all $t \in \mathbb{C}$ to a well-defined automorphism of the algebra $\widehat{U}_\varepsilon(\mathfrak{g})$.

(iii) Let \mathcal{G} be the group of automorphisms generated by the one-parameter groups $\exp(t\underline{x}_i^\pm)$, $i = 1, \dots, l$. The action of \mathcal{G} on $\widehat{U}_\varepsilon(\mathfrak{g})$ preserves the subalgebras \widehat{Z}_ε and \widehat{Z}_0 , and hence \mathcal{G} acts by holomorphic automorphisms on the complex algebraic varieties $\text{Spec}(Z_\varepsilon)$ and $\text{Spec}(Z_0)$.

(iv) Let \mathcal{O} be a conjugacy class in G . The intersection $\mathcal{O}^0 = \mathcal{O} \cap G^0$ is a smooth connected variety, and the variety $\pi^{-1}(\mathcal{O}^0)$ is a \mathcal{G} -orbit in $\text{Spec}(Z_0)$.

(v) If \mathcal{P} is a \mathcal{G} -orbit in $\text{Spec}(Z_0)$ then the connected components of $\tau^{-1}(\mathcal{P})$ are \mathcal{G} -orbits in $\text{Spec}(Z_\varepsilon)$.

Given a homomorphism $\eta : Z_0 \rightarrow \mathbb{C}$, let

$$U_\eta(\mathfrak{g}) = U_\varepsilon(\mathfrak{g})/I_\eta,$$

where I_η is the ideal in $U_\varepsilon(\mathfrak{g})$ generated by elements $z - \eta(z)$, $z \in Z_0$. By part (iii) of Proposition 6.1 $U_\eta(\mathfrak{g})$ is an algebra of dimension $m^{\dim \mathfrak{g}}$ with linear basis the set of monomials $(X^+)^r L^s (X^-)^t$ for which $0 \leq r_k, t_k, s_i < m$ for $i = 1, \dots, l$, $k = 1, \dots, D$.

If $\tilde{g} \in \mathcal{G}$ then for any $\eta \in \text{Spec}(Z_0)$ we have $\tilde{g}\eta \in \text{Spec}(Z_0)$ by part (iii) of Proposition 6.5, and by part (ii) of the same proposition \tilde{g} induces an isomorphism of algebras,

$$\tilde{g} : U_\eta(\mathfrak{g}) \rightarrow U_{\tilde{g}\eta}(\mathfrak{g}).$$

7. WHITTAKER VECTORS

In this section we introduce the notion of Whittaker vectors for modules over quantum groups at roots of unity and prove an analogue of the Engel theorem for them. We start by studying some properties of quantum groups at roots of unity.

From now on we fix an element $s \in W$. Let Δ_+ be a system of positive roots associated to s^{-1} . We also fix positive integer d such that $p_{ij} \in \frac{1}{d}\mathbb{Z}$ for any $i < j$ (or $i > j$), $i, j = 1, \dots, l$, where the numbers p_{ij} are defined by formula (4.3). We shall always assume that $m > d_i$ is odd and coprime to all d_i , $i = 1, \dots, l$ and that d and m are coprime. The last condition is equivalent to the existence of an integer n such that $\varepsilon^{nd-1} = 1$. We fix an integer valued solution n_{ij} to equations (4.4) and identify the algebra $U_\varepsilon^{s^{-1}}(\mathfrak{g})$ associated to the Weyl group element s^{-1} with $U_\varepsilon(\mathfrak{g})$ using Theorem 4.2 and the solution $-n_{ij}$ to equations (4.4). Using this identification $U_\varepsilon^{s^{-1}}(\mathfrak{m}_-)$ can be regarded as a subalgebra in $U_\varepsilon(\mathfrak{g})$. Therefore for every character $\eta : Z_0 \rightarrow \mathbb{C}$ one can define the corresponding subalgebra in $U_\eta(\mathfrak{g})$. We denote this subalgebra by $U_\eta(\mathfrak{m}_-)$. By part (ii) of Proposition 6.2 we have $\dim U_\eta(\mathfrak{m}_-) = m^{\dim \mathfrak{m}_-}$.

First we study some properties of the finite dimensional algebras $U_\eta(\mathfrak{g})$ and $U_\eta(\mathfrak{m}_-)$. In order to define Whittaker vectors for quantum groups at roots of unity we shall need some auxiliary notions that we are going to discuss now.

Below we use the normal ordering of Δ_+ associated to s^{-1} . Observe that by Proposition 5.1 for any two roots $\alpha, \beta \in \Delta_{\mathfrak{m}_+}$ such that $\alpha < \beta$ the sum $\alpha + \beta$ can not be represented as a linear combination $\sum_{k=1}^q c_k \gamma_{i_k}$, where $c_k \in \mathbb{N}$ and $\alpha < \gamma_{i_1} < \dots < \gamma_{i_k} < \beta$, and hence from commutation relations (4.15) one can deduce that

$$(7.1) \quad f_\alpha f_\beta - \varepsilon^{(\alpha, \beta) - nd(\frac{1+s}{1-s} P_{\mathfrak{v}'^* \alpha, \beta})} f_\beta f_\alpha = \sum_{\alpha < \delta_1 < \dots < \delta_n < \beta} C'(k_1, \dots, k_n) f_{\delta_n}^{k_n} f_{\delta_{n-1}}^{k_{n-1}} \dots f_{\delta_1}^{k_1} \in \mathcal{J},$$

where at least one of the roots δ_i in the right hand side of the last formula belongs to $\Theta = \{\alpha \in \Delta_{\mathfrak{m}_+} : \alpha \notin \{\gamma_1, \dots, \gamma_{l'}\}\}$, \mathcal{J} is the ideal in $U_\eta(\mathfrak{m}_-)$ generated by the elements $f_\beta \in U_\eta(\mathfrak{m}_-)$, $\beta \in \Theta$. Thus from part (ii) of Proposition 6.2 and commutation relations (7.1) it follows that if $\delta_1 < \delta_2 < \dots < \delta_b$ are the roots in the segment $\Delta_{\mathfrak{m}_+}$, the elements

$$(7.2) \quad x_{k_1, \dots, k_b} = f_{\delta_b}^{k_b} f_{\delta_{b-1}}^{k_{b-1}} \dots f_{\delta_1}^{k_1}$$

for $k_i \in \mathbb{N}$, $k_i < m$ form a linear basis of $U_\eta(\mathfrak{m}_-)$, and elements (7.2) for $k_i \in \mathbb{N}$, $k_i < m$ and $k_i > 0$ for at least one $\delta_i \in \Theta$ form a linear basis of \mathcal{J} .

Lemma 7.1. *Let η be an element of $\text{Spec}(Z_0)$. Assume that $\eta(f_{\gamma_i}^m) = a_i \neq 0$ for $i = 1, \dots, l'$ and that $\eta(f_\beta^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$, and hence $f_{\gamma_i}^m = \eta(f_{\gamma_i}^m) = a_i \neq 0$ in $U_\eta(\mathfrak{m}_-)$ for $i = 1, \dots, l'$ and $f_\beta^m = 0$ in $U_\eta(\mathfrak{m}_-)$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$. Then the ideal \mathcal{J} is the Jacobson radical of $U_\eta(\mathfrak{m}_-)$ and $U_\eta(\mathfrak{m}_-)/\mathcal{J}$ is isomorphic to the truncated polynomial algebra*

$$\mathbb{C}[f_{\gamma_1}, \dots, f_{\gamma_{l'}}]/\{f_{\gamma_i}^m = a_i\}_{i=1, \dots, l'}$$

.

Proof. First we show that \mathcal{J} is nilpotent.

Let i be the largest number such that $k_j = 0$ for $j > i$ in (7.2) and $k_i \neq 0$. Then we define the degree of x_{k_1, \dots, k_b} by

$$\deg(x_{k_1, \dots, k_b}) = (k_i, i) \in \{1, \dots, m-1\} \times \{1, \dots, b\}.$$

Equip $\{1, \dots, m-1\} \times \{1, \dots, b\}$ with the order such that $(k, i) < (k', j)$ if $j > i$ or $j = i$ and $k' > k$.

For any given $(k, i) \in \{1, \dots, m-1\} \times \{1, \dots, b\}$ denote by $(U_\eta(\mathfrak{m}_-))_{(k, i)}$ the linear span of the elements x_{k_1, \dots, k_b} with $\deg(x_{k_1, \dots, k_b}) \leq (k, i)$ and define $\mathcal{J}_{(k, i)} = \mathcal{J} \cap (U_\eta(\mathfrak{m}_-))_{(k, i)}$. We also have $(U_\eta(\mathfrak{m}_-))_{(k, i)} \subset (U_\eta(\mathfrak{m}_-))_{(k', j)}$ and $\mathcal{J}_{(k, i)} \subset \mathcal{J}_{(k', j)}$ if $(k, i) < (k', j)$, and $\mathcal{J}_{(m-1, b)} = \mathcal{J}$. Note that for the first few i linear spaces $\mathcal{J}_{(k, i)}$ may be trivial, and these are all possibilities when those spaces can be trivial.

We shall prove that \mathcal{J} is nilpotent by induction over the order in $\{1, \dots, m-1\} \times \{1, \dots, b\}$. Let (k, i) be minimal possible such that $\mathcal{J}_{(k, i)}$ is not trivial. Then we must have $k = 1$. If $y \in \mathcal{J}_{(1, i)}$ then y must be of the form

$$(7.3) \quad y = f_\beta v,$$

where v is a linear combination of elements of the form $f_{\delta_{i_1}}^{k_1} \dots f_{\delta_{i_r}}^{k_r}$ for $\delta_{i_1}, \dots, \delta_{i_r} \in \{\gamma_{n+1}, \dots, \gamma_{l'}\}$, $\beta > \delta_{i_1}$, and β is the first root from the set Θ greater than $\gamma_{l'}$ in the normal ordering of Δ_+ associated to s^{-1} . Here it is assumed that $f_{\delta_1}^{k_1} \dots f_{\delta_r}^{k_r} = 1$ if the set $\{\gamma_{n+1}, \dots, \gamma_{l'}\}$ is empty.

Now equation (7.1) implies that for any $f_{\delta_{i_j}}$ which appears in the expression for v one has

$$(7.4) \quad f_\beta f_{\delta_{i_j}} - \varepsilon^{(\beta, \delta_{i_j}) - nd(\frac{1+s}{1-s} P_{\mathfrak{h}'^* \beta, \delta_{i_j}})} f_{\delta_{i_j}} f_\beta \in \mathcal{J}_{(m-1, i-1)} = 0$$

as by our choice of i $\mathcal{J}_{(m-1, i-1)} = 0$.

Formula (7.4) implies that the product of m elements of type (7.3) can be represented in the form

$$f_\beta^m v',$$

where v' is of the same form as v . Since $f_\beta^m = 0$ we deduce that $\mathcal{J}_{(1, i)}^m = 0$.

Now assume that $\mathcal{J}_{(k, i)}^K = 0$ for some $K > 0$. Let (k', i') be the smallest element of $\{1, \dots, m-1\} \times \{1, \dots, b\}$ which satisfies $(k, i) < (k', i')$. Then by Propositions 4.6 and 6.2, and by (7.1), any element of $\mathcal{J}_{(k', i')}$ is of the form $f_{\delta_{i'}} u + u'$, where $u' \in \mathcal{J}_{(k, i)}$ and if $\delta_{i'} \in \Theta$ then $u \in (U_\eta(\mathfrak{m}_-))_{(k, i)}$; if $\delta_{i'} \notin \Theta$ then $u \in \mathcal{J}_{(k, i)}$.

Now equation (7.1) together with (4.15) imply that for any $u \in (U_\eta(\mathfrak{m}_-))_{(k,i)}$ one has

$$(7.5) \quad u f_{\delta_{i'}} = c f_{\delta_{i'}} u + w,$$

where c is a non-zero constant depending on u , and $w \in \mathcal{J}_{(k,i)}$. By formula (7.5) the product of m elements $f_{\delta_{i'}} u_p + u'_p$, $p = 1, \dots, m$ of the type described above can be represented in the form

$$(7.6) \quad \sum_{j=0}^m f_{\delta_{i'}}^j c_j,$$

where $c_j \in \mathcal{J}_{(k,i)}$ for $j = 0, \dots, m-1$ and if $\delta_{i'} \in \Theta$ then $c_m \in (U_\eta(\mathfrak{m}_-))_{(k,i)}$; if $\delta_{i'} \notin \Theta$ then $c_m \in \mathcal{J}_{(k,i)}$. In the former case $f_{\delta_{i'}}^m = 0$, and the last term in sum (7.6) is zero; in the latter case $f_{\delta_{i'}}^m = \eta(f_{\delta_{i'}}^m) \neq 0$, and the last term in sum (7.6) is from $\mathcal{J}_{(k,i)}$. So we can combine it with the term corresponding to $j = 0$. In both cases sum (7.6) takes the form

$$(7.7) \quad \sum_{j=0}^{m-1} f_{\delta_{i'}}^j c'_j,$$

where $c'_j \in \mathcal{J}_{(k,i)}$. By (7.5) the product of K sums of type (7.7) is of the form

$$\sum_{j=0}^{(m-1)K} f_{\delta_{i'}}^j c''_j,$$

where each c''_j is a linear combination of elements from $\mathcal{J}_{(k,i)}^K$. By our assumption $\mathcal{J}_{(k,i)}^K = 0$, and hence the product of any mK elements of $\mathcal{J}_{(k',i')}$ is zero. This justifies the induction step and proves that $\mathcal{J}_{(m-1,b)} = \mathcal{J}$ is nilpotent. Hence \mathcal{J} is contained in the Jacobson radical of $U_\eta(\mathfrak{m}_-)$.

Using commutation relations (4.15) we also have (see the proof of Theorem 5.4)

$$f_{\gamma_i} f_{\gamma_j} - f_{\gamma_j} f_{\gamma_i} \in \mathcal{J}.$$

Therefore the quotient algebra $U_\eta(\mathfrak{m}_-)/\mathcal{J}$ is isomorphic to the truncated polynomial algebra

$$\mathbb{C}[f_{\gamma_1}, \dots, f_{\gamma_{l'}}] / \{f_{\gamma_i}^m = a_i\}_{i=1, \dots, l'}$$

which is semisimple. Therefore \mathcal{J} coincides with the Jacobson radical of $U_\eta(\mathfrak{m}_-)$. \square

In Theorem 5.4 we constructed some characters of the algebra $U_\varepsilon^s(\mathfrak{m}_-)$. Similarly one can define characters of the algebra $U_\varepsilon^{s^{-1}}(\mathfrak{m}_-)$. Now we show that the algebra $U_\eta(\mathfrak{m}_-)$ has a finite number of irreducible representations which are one-dimensional, and all those representations can be obtained from each other by twisting with the help of automorphisms of $U_\eta(\mathfrak{m}_-)$.

Proposition 7.2. *Let η be an element of $\text{Spec}(Z_0)$. Assume that $\eta(f_{\gamma_i}^m) = a_i \neq 0$ for $i = 1, \dots, l'$ and that $\eta(f_\beta^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$, and hence $f_{\gamma_i}^m = \eta(f_{\gamma_i}^m) = a_i \neq 0$ in $U_\eta(\mathfrak{m}_-)$ for $i = 1, \dots, l'$ and $f_\beta^m = 0$ in $U_\eta(\mathfrak{m}_-)$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$. Then all non-zero irreducible representations of the algebra $U_\eta(\mathfrak{m}_-)$ are one-dimensional and have the form*

$$(7.8) \quad \chi(f_\beta) = \begin{cases} 0 & \beta \notin \{\gamma_1, \dots, \gamma_{l'}\} \\ c_i & \beta = \gamma_i, i = 1, \dots, l' \end{cases},$$

where complex numbers c_i satisfy the conditions $c_i^m = a_i$, $i = 1, \dots, l'$. Moreover, all non-zero irreducible representations of $U_\eta(\mathfrak{m}_-)$ can be obtained from each other by twisting with the help of automorphisms of $U_\eta(\mathfrak{m}_-)$.

Proof. Let V be a non-zero finite-dimensional irreducible $U_\eta(\mathfrak{m}_-)$ -module. By Corollary 54.13 in [6] elements of the ideal $\mathcal{J} \subset U_\eta(\mathfrak{m}_-)$ act by zero transformations on V . Hence V is in fact an irreducible representation of the algebra $U_\eta(\mathfrak{m}_-)/\mathcal{J}$ which is isomorphic to the truncated polynomial algebra

$$\mathbb{C}[f_{\gamma_1}, \dots, f_{\gamma_{l'}}]/\{f_{\gamma_i}^m = a_i\}_{i=1, \dots, l'}.$$

The last algebra is commutative and all its complex irreducible representations are one-dimensional. Therefore V is one-dimensional, and if v is a nonzero element of V then $f_{\gamma_i}v = c_i v$, for some $c_i \in \mathbb{C}$, $i = 1, \dots, l'$. Note that $\eta(f_{\gamma_i}^m) = a_i \neq 0$, $i = 1, \dots, l'$ and hence $c_i^m = a_i \neq 0$, $i = 1, \dots, l'$. In particular, the elements f_{γ_i} act on V by semisimple automorphisms.

If we denote by $\chi : U_\eta(\mathfrak{m}_-) \rightarrow \mathbb{C}$ the character of $U_\eta(\mathfrak{m}_-)$ such that

$$\chi(f_\beta) = \begin{cases} 0 & \beta \notin \{\gamma_1, \dots, \gamma_{l'}\} \\ c_i & \beta = \gamma_i, i = 1, \dots, l' \end{cases}$$

and by \mathbb{C}_χ the corresponding one-dimensional representation of $U_\eta(\mathfrak{m}_-)$ then we have $V = \mathbb{C}_\chi$.

Now we have to prove that the representations \mathbb{C}_χ for different characters χ are obtained from each other by twisting with the help of automorphisms of $U_\eta(\mathfrak{m}_-)$.

Since $c_i^m = a_i$, $i = 1, \dots, l'$ there are only finitely many possible characters χ corresponding to the given η in the statement of this proposition. If χ and χ' are two such characters, $\chi(f_{\gamma_i}) = c_i$, $i = 1, \dots, l'$ and $\chi'(f_{\gamma_i}) = c'_i$, $i = 1, \dots, l'$ then the relations $c_i^m = c_i'^m = a_i$, $i = 1, \dots, l'$ imply that $c'_i = \varepsilon^{m_i} c_i$, $0 \leq m_i \leq m - 1$, $m_i \in \mathbb{Z}$, $i = 1, \dots, l'$.

Now observe that for any $h \in \mathfrak{h}$ the map defined by $f_\alpha \mapsto \varepsilon^{\alpha(h)} f_\alpha$, $\alpha \in \Delta_{\mathfrak{m}_+}$ is an automorphism of the algebra $U_\varepsilon^{s-1}(\mathfrak{m}_-)$ generated by elements f_α , $\alpha \in \Delta_{\mathfrak{m}_+}$ with defining relations (4.15). Here the principal branch of the analytic function ε^z is used to define $\varepsilon^{\alpha(h)}$, so that $\varepsilon^{\alpha(h)} \varepsilon^{\beta(h)} = \varepsilon^{(\alpha+\beta)(h)}$ for any $\alpha, \beta \in \Delta_{\mathfrak{m}_+}$. If in addition $\varepsilon^{m\gamma_i(h)} = 1$, $i = 1, \dots, l'$ the above defined map gives rise to an automorphism ς of $U_\eta(\mathfrak{m}_-)$. Indeed in that case $(\varepsilon^{\gamma_i(h)} f_{\gamma_i})^m = f_{\gamma_i}^m$, $i = 1, \dots, l'$ and all the remaining defining relations $f_{\gamma_i}^m = \eta(f_{\gamma_i}^m) = a_i \neq 0$, $i = 1, \dots, l'$, $f_\beta^m = \eta(f_\beta^m) = 0$, $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$ of the algebra $U_\eta(\mathfrak{m}_-)$ are preserved by the action of the above defined map ς .

Now fix $h \in \mathfrak{h}$ such that $\gamma_i(h) = m_i$, $i = 1, \dots, l'$. Obviously we have $\varepsilon^{m m_i} = 1$, $i = 1, \dots, l'$. We claim that the representation \mathbb{C}_χ twisted by the corresponding automorphism ς coincides with $\mathbb{C}_{\chi'}$. Indeed, we obtain

$$\chi(\varsigma f_{\gamma_i}) = \chi(\varepsilon^{m_i} f_{\gamma_i}) = \varepsilon^{m_i} c_i = c'_i, \quad i = 1, \dots, l'.$$

This completes the proof of the proposition. \square

Let V be a $U_\eta(\mathfrak{g})$ -module, where η is an element of $\text{Spec}(Z_0)$ such that $\eta(f_{\gamma_i}^m) = a_i \neq 0$ for $i = 1, \dots, l'$ and that $\eta(f_\beta^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$. Let $\chi : U_\eta(\mathfrak{m}_-) \rightarrow \mathbb{C}$ be a character defined in the Proposition 7.2, \mathbb{C}_χ the corresponding one-dimensional $U_\eta(\mathfrak{m}_-)$ -module. Then the space $V_\chi = \text{Hom}_{U_\eta(\mathfrak{m}_-)}(\mathbb{C}_\chi, V)$ is called the space of Whittaker vectors of V . Elements of V_χ are called Whittaker vectors.

Now we describe the space of Whittaker vectors in terms of a nilpotent action of the unital subalgebra $U_{\eta_0}(\mathfrak{m}_-)$ generated by f_α , $\alpha \in \Delta_{\mathfrak{m}_+}$ in the small quantum group $U_{\eta_0}(\mathfrak{g}) = U_\varepsilon^{s-1}(\mathfrak{g})/I_{\eta_0}$ corresponding to the trivial central character η_0 such that $\tilde{\pi}(\eta_0) = 1 \in G_0^*$ and $\eta_0(x_\alpha^\pm) = 0$, $\alpha \in \Delta_+$, $\eta_0(l_i) = 1$, $i = 1, \dots, l$. Recall that the algebra $U_\varepsilon^{s-1}(\mathfrak{g})$ is a Hopf algebra. We shall need the following formula for the action of the comultiplication on the quantum root elements f_β , $\beta = \sum_{i=1}^l c_i \alpha_i \in$

$\Delta_{\mathfrak{m}_+}, c_i \in \mathbb{N}$,

$$(7.9) \quad \begin{aligned} \Delta_{s-1}(f_\beta) = & \prod_{i=1}^l K_i^{c_i} \prod_{i,j=1}^l L_j^{\frac{nd}{d_j}(\frac{1+s}{1-s}P_{\mathfrak{h}'^*} \alpha_i, \alpha_j)c_i} \otimes f_\beta + f_\beta \otimes 1 + \\ & + \sum_i y_i \otimes x_i, \quad x_i \in U_{<\beta}, y_i \in U_{>\beta} U_\varepsilon^{s-1}(\mathfrak{h}), \end{aligned}$$

where $U_{<\beta}$ is the subalgebra (without unit) in $U_\varepsilon^{s-1}(\mathfrak{m}_-)$ generated by $f_\alpha, \alpha < \beta$ and $U_{>\beta}$ is the subalgebra (without unit) in $U_\varepsilon^{s-1}(\mathfrak{n}_-)$ generated by $f_\alpha, \alpha > \beta$.

To derive formula (7.9) we first observe that by Corollary 4.3.2 in [5] one has in $U_h^{s-1}(\mathfrak{g})$

$$(7.10) \quad \begin{aligned} \Delta(X_\beta^-) = & \prod_{i=1}^l K_i^{c_i} \otimes X_\beta^- + X_\beta^- \otimes 1 + \\ & + \sum_i y'_i \otimes x'_i, \end{aligned}$$

where x'_i are elements of the subalgebra (without unit) in $U_h(\mathfrak{g})$ generated by $X_\alpha^-, \alpha < \beta$ and y'_i are elements of the subalgebra (without unit) in $U_h(\mathfrak{g})$ generated by $X_\alpha^-, \alpha > \beta$ and by the $L_j^{\pm 1}, j = 1, \dots, l$.

Using formula (4.8), the definition of the elements f_β and the action of the comultiplication Δ on the elements H_i we obtain from formula (7.10) that

$$\begin{aligned} \Delta_{s-1}(f_\beta) = & \prod_{i=1}^l K_i^{c_i} \prod_{i,j=1}^l L_j^{\frac{nd}{d_j}(\frac{1+s}{1-s}P_{\mathfrak{h}'^*} \alpha_i, \alpha_j)c_i} \otimes f_\beta + f_\beta \otimes 1 + \\ & + \sum_i y_i \otimes x_i, \end{aligned}$$

where x_i are elements of the subalgebra (without unit) in $U_h^{s-1}(\mathfrak{g})$ generated by $f_\alpha, \alpha < \beta$ and by the $L_j^{\pm 1}, j = 1, \dots, l$ and y_i are elements of the subalgebra (without unit) in $U_h^{s-1}(\mathfrak{g})$ generated by $f_\alpha, \alpha > \beta$ and by the $L_j^{\pm 1}, j = 1, \dots, l$.

On the other hand formula (4.11) and commutation relations (4.15) imply that in fact x_i are elements of the subalgebra (without unit) in $U_h^{s-1}(\mathfrak{g})$ generated by $f_\alpha, \alpha < \beta$. Since $U_{\mathcal{A}}^{s-1}(\mathfrak{g})$ is in fact a Hopf subalgebra in $U_h^{s-1}(\mathfrak{g})$ we deduce that x_i are elements of the subalgebra (without unit) in $U_{\mathcal{A}}^{s-1}(\mathfrak{g})$ generated by $f_\alpha, \alpha < \beta$ and y_i are elements of the subalgebra (without unit) in $U_{\mathcal{A}}^{s-1}(\mathfrak{g})$ generated by $f_\alpha, \alpha > \beta$ and by the $L_j^{\pm 1}, j = 1, \dots, l$. This implies (7.9).

Formula (7.9) shows that $U_\varepsilon^{s-1}(\mathfrak{m}_-)$ is a right coideal in $U_\varepsilon^{s-1}(\mathfrak{g})$. One can also equip the algebra $U_\varepsilon^{s-1}(\mathfrak{m}_-)$ with a character given by formula (5.5), where the numbers c_i are the same as in the definition of the character χ . We denote this character by the same letter, $\chi : U_\varepsilon^{s-1}(\mathfrak{m}_-) \rightarrow \mathbb{C}$.

Note that V can be regarded as a $U_\varepsilon(\mathfrak{g})$ -module and a $U_\varepsilon^{s-1}(\mathfrak{g})$ -module assuming that the ideal I_η acts on V in the trivial way. Now observe that $\Delta_{s-1} : U_\varepsilon^{s-1}(\mathfrak{m}_-) \rightarrow U_\varepsilon^{s-1}(\mathfrak{g}) \otimes U_\varepsilon^{s-1}(\mathfrak{m}_-)$ is a homomorphism of algebras. Composing it with the tensor product $S_{s-1} \otimes \chi$ of the anti-homomorphism S_{s-1} and of the character χ , which can be regarded as an anti-homomorphism as well, one can define an anti-homomorphism, $U_\varepsilon^{s-1}(\mathfrak{m}_-) \rightarrow U_\varepsilon^{s-1}(\mathfrak{g}), x \mapsto S_{s-1}(x_1)\chi(x_2), \Delta_{s-1}(x) = x_1 \otimes x_2, x \in U_\varepsilon^{s-1}(\mathfrak{m}_-)$.

Using this anti-homomorphism one can introduce a right $U_\varepsilon^{s-1}(\mathfrak{m}_-)$ -action on V which we call the adjoint action and denote it by Ad . It is given by the formula (compare with the definition of

the adjoint action in (4.13))

$$(7.11) \quad \text{Ad } xv = S_{s-1}(x_1)\chi(x_2)v, x \in U_\varepsilon^{s-1}(\mathfrak{m}_-), v \in V,$$

where $\Delta_{s-1}(x) = x_1 \otimes x_2$.

Note that using the Sweidler notation for the comultiplication, $(\Delta_{s-1} \otimes id \otimes id)(\Delta_{s-1} \otimes id)\Delta_{s-1}(x) = x_1 \otimes x_2 \otimes x_3 \otimes x_4$, the coassociativity of the comultiplication and the definition of the antipode we have for any $x \in U_\varepsilon^{s-1}(\mathfrak{m}_-)$, $y \in U_\varepsilon^{s-1}(\mathfrak{g})$, $v \in V$ (compare with the proof of Lemma 2.2 in [8])

$$(7.12) \quad \text{Ad}x(yv) = S_{s-1}(x_1)\chi(x_2)yv = S_{s-1}(x_1)yx_2S_{s-1}(x_3)\chi(x_4)v = \text{Ad}x_1(y)\text{Ad}x_2(v).$$

Similarly to the Proposition in Section 5.6 in [4] we infer that Z_0 is a Hopf subalgebra in $U_\varepsilon^{s-1}(\mathfrak{g})$. Namely,

$$\begin{aligned} \Delta_{s-1}(f_i^m) &= K_i^m \prod_{j=1}^l L_j^{m \frac{nd}{d_j} (\frac{1+s}{1-s} P_{\mathfrak{h}'^* \alpha_i, \alpha_j})} \otimes f_i^m + f_i^m \otimes 1, \\ \Delta_{s-1}(e_i^m) &= e_i^m \otimes K_i^{-m} + \prod_{j=1}^l L_j^{-m \frac{nd}{d_j} (\frac{1+s}{1-s} P_{\mathfrak{h}'^* \alpha_i, \alpha_j})} \otimes e_i^m, \\ \Delta_{s-1}(L_i^m) &= L_i^m \otimes L_i^m. \end{aligned}$$

Therefore recalling that by the definition of χ for $x \in U_\varepsilon^{s-1}(\mathfrak{m}_-) \cap Z_0$ one has $\chi(x) = \eta(x)$ we deduce

$$\text{Ad } xv = S_{s-1}(x_1)\chi(x_2)v = \eta(S_{s-1}(x_1)x_2)v = \varepsilon_{s-1}(x)v, v \in V,$$

where ε_{s-1} is the counit of $U_\varepsilon^{s-1}(\mathfrak{g})$. Note that by the definition of the ideal I_{η_0} the ideal $U_\varepsilon^{s-1}(\mathfrak{m}_-) \cap I_{\eta_0} \subset U_\varepsilon^{s-1}(\mathfrak{m}_-)$ is generated by the elements f_α^m , $\alpha \in \Delta_{\mathfrak{m}_+}$ and $\varepsilon_{s-1}(f_\alpha^m) = 0$ for $\alpha \in \Delta_{\mathfrak{m}_+}$ by the definition of ε_{s-1} . Hence the adjoint action of $U_\varepsilon^{s-1}(\mathfrak{m}_-)$ on V induces an action of the subalgebra $U_{\eta_0}(\mathfrak{m}_-)$ of the small quantum group $U_{\eta_0}(\mathfrak{g})$. We call this action the adjoint action as well.

Note that the small quantum group $U_{\eta_0}(\mathfrak{g})$ is a Hopf algebra with the comultiplication inherited from $U_\varepsilon^{s-1}(\mathfrak{g})$. Now arguments used in the proof of Proposition 5.6 in [18] can be applied verbatim to establish the following lemma.

Lemma 7.3. *The space of Whittaker vectors V_χ coincides with the space of $U_{\eta_0}(\mathfrak{m}_-)$ -invariants for the adjoint action on V ,*

$$(7.13) \quad V_\chi = \{v \in V : \text{Ad } x(v) = \varepsilon_{s-1}(x)v \ \forall x \in U_{\eta_0}(\mathfrak{m}_-)\}.$$

Proof. Indeed, denote by T_β the factor $\prod_{i=1}^l K_i^{c_i} \prod_{i,j=1}^l L_j^{\frac{nd}{d_j} (\frac{1+s}{1-s} P_{\mathfrak{h}'^* \alpha_i, \alpha_j}) c_i}$ which appears in (7.9), $T_\beta = \prod_{i=1}^l K_i^{c_i} \prod_{i,j=1}^l L_j^{\frac{nd}{d_j} (\frac{1+s}{1-s} P_{\mathfrak{h}'^* \alpha_i, \alpha_j}) c_i}$. Then by the definition of the antipode we have from (7.9)

$$S_{s-1}(T_\beta)f_\beta + S_{s-1}(f_\beta) + \sum_i S_{s-1}(y_i)x_i = \varepsilon_{s-1}(f_\beta) = 0.$$

Since $S_{s-1}(T_\beta) = T_\beta^{-1} = \prod_{i=1}^l K_i^{-c_i} \prod_{i,j=1}^l L_j^{-\frac{nd}{d_j} (\frac{1+s}{1-s} P_{\mathfrak{h}'^* \alpha_i, \alpha_j}) c_i}$ this yields

$$(7.14) \quad S_{s-1}(f_\beta) = -S_{s-1}(T_\beta)f_\beta - \sum_i S_{s-1}(y_i)x_i.$$

Now for $\beta \in \Delta_{\mathfrak{m}_+}$, (7.9), (7.14) and definition (7.11) of the adjoint action imply

$$(7.15) \quad \begin{aligned} \text{Ad } f_\beta v &= T_\beta^{-1} \chi(f_\beta) v - T_\beta^{-1} f_\beta v - \sum_i S_{s^{-1}(y_i)} x_i v + \sum_i S_{s^{-1}(y_i)} \chi(x_i) v = \\ &= T_\beta^{-1} (\chi(f_\beta) - f_\beta) v + \sum_i S_{s^{-1}(y_i)} (\chi(x_i) - x_i) v, \quad x_i \in U_{<\beta}, y_i \in U_{>\beta} U_\varepsilon^{s^{-1}}(\mathfrak{h}). \end{aligned}$$

If $v \in V_\chi$ we immediately obtain from (7.15) that $\text{Ad } f_\beta v = 0$ for any $\beta \in \Delta_{\mathfrak{m}_+}$, i.e. v belongs to the right hand side of (7.13).

Conversely, suppose that v belongs to the right hand side of (7.13). We shall show that $xv = \chi(x)v$ for any $x \in U_\varepsilon^{s^{-1}}(\mathfrak{m}_-)$. Let $\overline{U}_{<\beta}$ be the subalgebra with unit generated by $U_{<\beta}$. We proceed by induction over the subalgebras $\overline{U}_{<\delta_k}$, $k = 1, \dots, b+1$, where as before $\delta_1 < \dots < \delta_b$ is the normally ordered segment $\Delta_{\mathfrak{m}_+}$ and we define $\overline{U}_{<\delta_{b+1}}$ to be the subalgebra $U_\varepsilon^{s^{-1}}(\mathfrak{m}_-)$.

Observe that δ_1 is a simple root and hence $U_{<\delta_1} = 0$. Therefore we deduce from (7.15) for $\beta = \delta_1$

$$\text{Ad } f_{\delta_1} v = T_{\delta_1}^{-1} (\chi(f_{\delta_1}) - f_{\delta_1}) v = 0.$$

Since $T_{\delta_1}^{-1}$ acts on V by an invertible transformation this implies $(\chi(f_{\delta_1}) - f_{\delta_1})v = 0$, and hence $xv = \chi(x)v$ for any $x \in \overline{U}_{<\delta_2}$ as $\overline{U}_{<\delta_2}$ is generated by f_{δ_1} .

Now assume that for some $k \leq b$ $xv = \chi(x)v$ for any $x \in \overline{U}_{<\delta_k}$. Then by (7.15)

$$\text{Ad } f_{\delta_k} v = T_{\delta_k}^{-1} (\chi(f_{\delta_k}) - f_{\delta_k}) v = 0.$$

As above this implies

$$(7.16) \quad (\chi(f_{\delta_k}) - f_{\delta_k})v = 0.$$

By Proposition 4.6 any element $y \in \overline{U}_{<\delta_{k+1}}$ can be uniquely represented in the form $y = f_{\delta_k} y' + y''$, where $y', y'' \in \overline{U}_{<\delta_k}$. Now by (7.16) and by the induction assumption

$$yv = (f_{\delta_k} y' + y'')v = \chi(f_{\delta_k}) \chi(y')v + \chi(y'')v = \chi(y)v,$$

i.e. $yv = \chi(y)v$ for any $x \in \overline{U}_{<\delta_{k+1}}$. This establishes the induction step and completes the proof. \square

The following proposition is an analogue of Engel theorem for quantum groups at roots of unity.

Proposition 7.4. *Let η be an element of $\text{Spec}(Z_0)$. Assume that $\eta(f_{\gamma_i}^m) = a_i \neq 0$ for $i = 1, \dots, l'$ and that $\eta(f_\beta^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$, and hence $f_{\gamma_i}^m = \eta(f_{\gamma_i}^m) = a_i \neq 0$ in $U_\eta(\mathfrak{m}_-)$ for $i = 1, \dots, l'$ and $f_\beta^m = 0$ in $U_\eta(\mathfrak{m}_-)$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$. Let $\chi : U_\eta(\mathfrak{m}_-) \rightarrow \mathbb{C}$ be any character defined in Proposition 7.2. Then any non-zero finite-dimensional $U_\eta(\mathfrak{g})$ -module contains a non-zero Whittaker vector.*

Proof. First we show that the augmentation ideal \mathcal{J}^0 of $U_{\eta_0}(\mathfrak{m}_-)$ coincides with its Jacobson radical which is nilpotent. The proof of this fact is similar to that of Lemma 7.1, and we shall keep the notation used in that proof.

We define $\mathcal{J}_{(k,i)}^0 = \mathcal{J}^0 \cap (U_{\eta_0}(\mathfrak{m}_-))_{(k,i)}$, so that $\mathcal{J}_{(k,i)}^0 \subset \mathcal{J}_{(k',j)}^0$ if $(k,i) < (k',j)$, and $\mathcal{J}_{(m-1,b)}^0 = \mathcal{J}^0$.

We shall prove that \mathcal{J}^0 is nilpotent by induction over the order in $\{1, \dots, m-1\} \times \{1, \dots, b\}$. Note that $(k,i) = (1,1)$ is minimal possible such that $\mathcal{J}_{(k,i)}$ is not trivial. If $y \in \mathcal{J}_{(1,1)}$ then y must be of the form

$$(7.17) \quad y = af_{l'}, a \in \mathbb{C}.$$

The product of m elements of type (7.17) is equal to zero,

$$f_\beta^m a^m = 0,$$

as $f_\beta^m = 0$. We deduce that $(\mathcal{J}_{(1,1)}^0)^m = 0$.

Now assume that $(\mathcal{J}_{(k,i)}^0)^K = 0$ for some $K > 0$. Let (k', i') be the smallest element of $\{1, \dots, m-1\} \times \{1, \dots, b\}$ which satisfies $(k, i) < (k', i')$. Then by Propositions 4.6 and 6.2 any element of $\mathcal{J}_{(k',i')}^0$ is of the form $f_{\delta_{i'}} u + u'$, where $u' \in \mathcal{J}_{(k,i)}^0$ and $u \in (U_{\eta_0}(\mathfrak{m}_-))_{(k,i)}$.

Now equation (4.15) implies that for any $u \in (U_{\eta_0}(\mathfrak{m}_-))_{(k,i)}$ one has

$$(7.18) \quad u f_{\delta_{i'}} = c f_{\delta_{i'}} u + w,$$

where c is a non-zero constant depending on u , and $w \in \mathcal{J}_{(k,i)}^0$. By the formula (7.18) the product of m elements $f_{\delta_{i'}} u_p + u'_p$, $p = 1, \dots, m$ of the type described above can be represented in the form

$$(7.19) \quad \sum_{j=0}^m f_{\delta_{i'}}^j c_j,$$

where $c_j \in \mathcal{J}_{(k,i)}^0$ for $j = 0, \dots, m-1$ and $c_m \in (U_{\eta_0}(\mathfrak{m}_-))_{(k,i)}$. Since $f_{\delta_{i'}}^m = 0$ the last term in sum (7.19) is zero. So sum (7.19) takes the form

$$(7.20) \quad \sum_{j=0}^{m-1} f_{\delta_{i'}}^j c'_j,$$

where $c'_j \in \mathcal{J}_{(k,i)}^0$. By (7.18) the product of K sums of type (7.20) is of the form

$$\sum_{j=0}^{(m-1)K} f_{\delta_{i'}}^j c''_j,$$

where each c''_j is a linear combination of elements from $(\mathcal{J}_{(k,i)}^0)^K$. By our assumption $(\mathcal{J}_{(k,i)}^0)^K = 0$, and hence the product of any mK elements of $\mathcal{J}_{(k',i')}^0$ is zero. This justifies the induction step and proves that $\mathcal{J}_{(m-1,b)}^0 = \mathcal{J}^0$ is nilpotent. Hence \mathcal{J}^0 is contained in the Jacobson radical of $U_{\eta_0}(\mathfrak{m}_-)$.

The quotient algebra $U_{\eta_0}(\mathfrak{m}_-)/\mathcal{J}^0$ is isomorphic to \mathbb{C} . Therefore \mathcal{J}^0 coincides with the Jacobson radical of $U_{\eta_0}(\mathfrak{m}_-)$.

Now let V be a finite-dimensional $U_\eta(\mathfrak{g})$ -module. Then V is also a finite-dimensional $U_{\eta_0}(\mathfrak{m}_-)$ -module with respect to the adjoint action. Thus V must contain a non-trivial irreducible $U_{\eta_0}(\mathfrak{m}_-)$ -submodule with respect to the adjoint action on which the Jacobson radical \mathcal{J}^0 must act trivially. From (7.13) it follows that this non-trivial irreducible submodule consists of Whittaker vectors. This completes the proof. \square

Let

$$G = \bigcup_{\mathcal{C} \in C(W)} G_{\mathcal{C}}.$$

be the Lusztig partition of G (see [12]; we use the notation of [17], Section 4). Here $C(W) \subset \underline{W}$ is a certain subset of the set of conjugacy classes \underline{W} in W .

From the above discussion and Proposition 6.2 in [17] we deduce the following theorem.

Theorem 7.5. *Let $\eta \in \text{Spec}(Z_0)$ be an element such that $\pi\eta \in G_{\mathcal{C}}$, $\mathcal{C} \in C(W)$ and $s \in \mathcal{C}$. Denote by d the number corresponding to s^{-1} and defined in Proposition 6.2 in [17]. Assume that m and d are coprime.*

Then there is a system of positive roots $\Delta_+^{s^{-1}}$ and a quantum coadjoint transformation \tilde{g} such that $\xi = \tilde{g}\eta$ satisfies $\xi(f_{\gamma_i}^m) = a_i \neq 0$ for $i = 1, \dots, l'$ and $\xi(f_\beta^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$, where $f_\alpha \in U_\xi(\mathfrak{m}_-)$ are generators of the corresponding algebra $U_\xi(\mathfrak{m}_-) \subset U_\xi(\mathfrak{g})$ defined in Section 7. Let

$\chi : U_\xi(\mathfrak{m}_-) \rightarrow \mathbb{C}$ be any character defined in Proposition 7.2. Then any finite-dimensional $U_\eta(\mathfrak{g})$ -module contains a non-zero Whittaker vector with respect to the subalgebra $U_\eta^\sim(\mathfrak{m}_-) = \tilde{g}^{-1}U_\xi(\mathfrak{m}_-)$ and the character χ^\sim given by the composition of χ and \tilde{g} , $\chi^\sim = \chi \circ \tilde{g} : U_\eta^\sim(\mathfrak{m}_-) \rightarrow \mathbb{C}$.

Moreover, $\dim U_\eta^\sim(\mathfrak{m}_-) = m^{\frac{1}{2}\dim \mathcal{O}_{\pi\eta}}$, where $\mathcal{O}_{\pi\eta}$ is the conjugacy class of $\pi\eta \in G_\mathcal{C}$.

Proof. Let

$$\Delta_0 = \{\alpha \in \Delta \mid s(\alpha) = \alpha\},$$

and Γ the set of simple roots in a system of positive roots $\Delta_+^{s^{-1}}$ defined in Proposition 5.1. We shall need the parabolic subalgebra \mathfrak{p} of \mathfrak{g} and the parabolic subgroup P associated to the subset $\Gamma_0 = \Gamma \cap \Delta_0$ of simple roots and containing the Borel subalgebra corresponding to $\Delta_+^{s^{-1}}$. Let \mathfrak{n} and \mathfrak{l} be the nilradical and the Levi factor of \mathfrak{p} , N and L the unipotent radical and the Levi factor of P , respectively.

Note that we have natural inclusions of Lie algebras $\mathfrak{p} \supset \mathfrak{n}$. We also denote by $\bar{\mathfrak{n}}$ the nilpotent subalgebra opposite to \mathfrak{n} . Denote by N the subgroup of G corresponding to the Lie subalgebra \mathfrak{n} and by \bar{N} the opposite unipotent subgroup in G with the Lie algebra $\bar{\mathfrak{n}}$. Let Z be the subgroup of G generated by the semi-simple part of the Levi factor L corresponding to the Lie subalgebra \mathfrak{l} and by the centralizer of s in H . Denote by \dot{s} a representative of s in G . Let $N_s = \{v \in N \mid \dot{s}v\dot{s}^{-1} \in \bar{N}\}$ and $H_0 \subset H$ the subgroup corresponding to the orthogonal complement \mathfrak{h}_0 of \mathfrak{h}' in \mathfrak{h} with respect to the Killing form.

By Theorem 5.2 in [17] for every $\mathcal{C} \in C(W)$ and $s \in \mathcal{C}$ there is a system of positive roots $\Delta_+^{s^{-1}}$ as in Proposition 5.1 and such that all conjugacy classes in the stratum $G_\mathcal{C}$ intersect the variety $\dot{s}H_0N_s$ which is a subvariety of the transversal slice $\Sigma_s = \dot{s}ZN_s$ to the set of conjugacy classes in G . Note that N_s is not a subgroup in N_+ . But every element of $n_s \in N_s$ can be uniquely factorized as follows $n_s = n_s^+n_s^-$, $n_s^\pm \in N_s \cap N_\pm$, $N_s \cap N_- = N \cap N_-$, $N_s \cap N_+ \subset M_+$, where $M_+ \subset N_+$ is the subgroup corresponding to the Lie subalgebra \mathfrak{m}_+ . Therefore every element $\dot{s}h_0n_s \in \dot{s}H_0N_s$ can be represented as follows $\dot{s}h_0n_s = \dot{s}h_0n_s^+n_s^-$, and conjugating by n_s^- we obtain that $\dot{s}h_0n_s$ is conjugate to $n_s^- \dot{s}h_0n_s^+$. Since the decomposition $s = s^1s^2$ is reduced

$$(7.21) \quad \dot{s}^{-1}(N_s \cap N_-)\dot{s} = \dot{s}^{-1}(N \cap N_-)\dot{s} \subset M_+.$$

Taking into account that H_0 normalizes M_+ we have $n_s^- \dot{s}h_0n_s^+ = \dot{s}h_0h_0^{-1}\dot{s}^{-1}n_s^- \dot{s}h_0n_s^+ = \dot{s}h_0m_s$, $m_s = h_0^{-1}\dot{s}^{-1}n_s^- \dot{s}h_0n_s^+ \in M_+$. We deduce that all conjugacy classes in the stratum $G_\mathcal{C}$ intersect the variety $\dot{s}H_0M_+$.

Recall that by Proposition 6.2 in [18] we have $u = z_+n'\dot{s}m$ for some $z_+ \in Z \cap N_+$, $n' \in N$, $m \in M_+$, where $u \in G$ is an element of the form

$$(7.22) \quad u = \prod_{i=1}^{l'} \exp[t_i X_{-\gamma_i}],$$

$t_i \in \mathbb{C}$ are non-zero constants depending on the choice of the representative \dot{s} and the product over roots is taken in the order opposed to the normal order associated to s^{-1} . Factorizing $n' = n_+n_-$, $n_\pm \in N \cap N_\pm$ and recalling (7.21) we obtain $u = z_+n'\dot{s}m = z_+n_+\dot{s}s^{-1}n_-\dot{s}m = z_+n_+\dot{s}m'$, $m' = \dot{s}^{-1}n_-\dot{s}m \in M_+$, and hence $\dot{s} = n_+^{-1}z_+^{-1}um'^{-1}$.

Now let $\dot{s}h_0m_s, h_0 \in H_0, m_s \in M_+$ be an element of $\dot{s}H_0M_+$. Using the previous expression for \dot{s} we can write $\dot{s}h_0m_s = n_+^{-1}z_+^{-1}um'^{-1}h_0m_s$. Conjugating this element by z_+n_+ and recalling that H_0 normalizes M_+ we infer that $\dot{s}h_0m_s$ is conjugate to

$$um'^{-1}h_0m_s n_+^{-1}z_+^{-1} = uh_0n = \lambda_0(h_0^{\frac{1}{2}}n, h_0^{-\frac{1}{2}}u^{-1}),$$

where $n = h_0^{-1} m'^{-1} h_0 m_s n_+^{-1} z_+^{-1} \in N_+$, $h_0^{\frac{1}{2}} \in H_0$ is any element such that $h_0^{\frac{1}{2}} h_0^{\frac{1}{2}} = h_0$, λ_0 is defined immediately after Proposition 6.4. We deduce that all conjugacy classes in the stratum G_C intersect the variety uH_0N_+ .

By part (iv) of Proposition 6.5 we conclude that if $\eta \in \text{Spec}(Z_0)$ satisfies $\pi\eta \in G_C$ then there is a quantum coadjoint transformation \tilde{g} such that $\tilde{\pi}(\tilde{g}\eta) = (h_0^{\frac{1}{2}} n, h_0^{-\frac{1}{2}} u^{-1})$ for some $n \in N_+$, $h_0^{\frac{1}{2}} \in H_0$.

Denote $\xi = \tilde{g}\eta$. From the definition of the map $\tilde{\pi}$ and of the element u it follows that

$$\exp(\xi(x_{\beta_D}^-)X_{-\beta_D}) \exp(\xi(x_{\beta_{D-1}}^-)X_{-\beta_{D-1}}) \dots \exp(\xi(x_{\beta_1}^-)X_{-\beta_1}) = u$$

which implies $\xi((X_{\gamma_i}^-)^m) = \frac{t_i}{(\varepsilon_{\gamma_i} - \varepsilon_{\gamma_i^{-1}})^m} \neq 0$ for $i = 1, \dots, l'$ and that $\xi((X_{\beta}^-)^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$.

By the definition of the elements f_{β} with $\beta = \sum_{i=1}^{l'} m_i \alpha_i$ we have $f_{\beta} = \prod_{i,j=1}^{l'} L_j^{m_i n_{ij}} X_{\beta}^-$. Therefore the commutation relations between elements L_j and X_{β}^- imply that $f_{\beta}^m = c_{\beta} \prod_{i,j=1}^{l'} L_j^{m m_i n_{ij}} (X_{\beta}^-)^m$, where c_{β} are non-zero constants, and hence $\xi(f_{\beta}^m) = c_{\beta} \prod_{i,j=1}^{l'} \xi(L_j^m)^{m_i n_{ij}} \xi((X_{\beta}^-)^m)$. Since $\xi(L_j) \neq 0$ for $j = 1, \dots, l$, $\xi((X_{\gamma_i}^-)^m) = \frac{t_i}{(\varepsilon_{\gamma_i} - \varepsilon_{\gamma_i^{-1}})^m} \neq 0$ for $i = 1, \dots, l'$ and $\xi((X_{\beta}^-)^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$ we deduce $\xi(f_{\gamma_i}^m) = a_i \neq 0$ for $i = 1, \dots, l'$ and $\xi(f_{\beta}^m) = 0$ for $\beta \in \Delta_{\mathfrak{m}_+}$, $\beta \notin \{\gamma_1, \dots, \gamma_{l'}\}$. Thus ξ satisfies the condition of Propositions 7.2 and 7.4. Let $U_{\tilde{g}\eta}(\mathfrak{m}_-) = U_{\xi}(\mathfrak{m}_-)$ be the corresponding subalgebra in $U_{\xi}(\mathfrak{g})$.

Note that by Theorem 5.2 in [17] for any $g \in G_C$ we have

$$\dim Z_G(g) = \dim \Sigma_s,$$

where $Z_G(g)$ is the centralizer of g in G .

By the definition of Σ_s we also have $\dim \Sigma_s = l(s) + 2D_0 + \dim \mathfrak{h}'^{\perp}$. Observe also that $\dim G = 2D + \dim \mathfrak{h}$ and $\dim \mathfrak{h} - \dim \mathfrak{h}'^{\perp} = \dim \mathfrak{h}' = l'$, and hence from (5.3) we deduce that $\dim \mathfrak{m}_- = D - D_0 - \frac{1}{2}(l(s) - l') = \frac{1}{2}(\dim G - \dim \Sigma_s) = \frac{1}{2} \dim \mathcal{O}_g$ and $\dim U_{\tilde{g}\eta}(\mathfrak{m}_-) = \dim U_{\tilde{g}\eta}(\mathfrak{m}_-) = m^{\dim \mathfrak{m}_-} = m^{\frac{1}{2} \dim \mathcal{O}_g}$, where \mathcal{O}_g is the conjugacy class of any $g \in G_C$.

In particular, $\dim U_{\tilde{g}\eta}(\mathfrak{m}_-) = m^{\frac{1}{2} \dim \mathcal{O}_{\pi\eta}}$, where $\mathcal{O}_{\pi\eta}$ is the conjugacy class of $\pi\eta \in G_C$.

The remaining statements of this theorem are consequences of Proposition 7.4. \square

For given $\eta \in \text{Spec}(Z_0)$ and $\tilde{g} \in \mathcal{G}$ as in Theorem 7.5 we denote $\mathbb{C}_{\chi_{\tilde{g}}}$ the corresponding one-dimensional representation of $U_{\tilde{g}\eta}(\mathfrak{m}_-)$. Let $Q_{\chi_{\tilde{g}}}$ be the induced left $U_{\eta}(\mathfrak{g})$ -module, $Q_{\chi_{\tilde{g}}} = U_{\eta}(\mathfrak{g}) \otimes_{U_{\tilde{g}\eta}(\mathfrak{m}_-)} \mathbb{C}_{\chi_{\tilde{g}}}$. Let $W_{\varepsilon, \eta}^s(G) = \text{End}_{U_{\eta}(\mathfrak{g})}(Q_{\chi_{\tilde{g}}})^{opp}$ be the algebra of $U_{\eta}(\mathfrak{g})$ -endomorphisms of $Q_{\chi_{\tilde{g}}}$ with the opposite multiplication. The algebra $W_{\varepsilon, \eta}^s(G)$ is called a q-W algebra associated to $s \in W$. Denote by $U_{\eta}(\mathfrak{g})\text{-mod}$ the category of finite-dimensional left $U_{\eta}(\mathfrak{g})$ -modules and by $W_{\varepsilon, \eta}^s(G)\text{-mod}$ the category of finite-dimensional left $W_{\varepsilon, \eta}^s(G)$ -modules. Observe that if $V \in U_{\eta}(\mathfrak{g})\text{-mod}$ then the algebra $W_{\varepsilon, \eta}^s(G)$ naturally acts on the finite-dimensional space $V_{\chi_{\tilde{g}}} = \text{Hom}_{U_{\tilde{g}\eta}(\mathfrak{m}_-)}(\mathbb{C}_{\chi_{\tilde{g}}}, V) = \text{Hom}_{U_{\eta}(\mathfrak{g})}(Q_{\chi_{\tilde{g}}}, V)$ by compositions of homomorphisms.

Proposition 7.6. *Let $\Phi : E \mapsto Q_{\chi_{\tilde{g}}} \otimes_{W_{\varepsilon, \eta}^s(G)} E$ be the functor from the category of finite-dimensional left $W_{\varepsilon, \eta}^s(G)$ -modules to the category $U_{\eta}(\mathfrak{g})\text{-mod}$. Its right adjoint functor is given by $\Psi : V \mapsto V_{\chi_{\tilde{g}}}$ and satisfies $\Psi \circ \Phi = \text{Id}$.*

Proof. The first statement follows from the definitions.

For the second one, let E be a finite-dimensional $W_{\varepsilon, \eta}^s(G)$ -module. First we observe that by the definition of the algebra $W_{\varepsilon, \eta}^s(G)$ we have $W_{\varepsilon, \eta}^s(G) = \text{End}_{U_{\eta}(\mathfrak{g})}(Q_{\chi_{\tilde{g}}})^{opp} = \text{Hom}_{U_{\tilde{g}\eta}(\mathfrak{m}_-)}(\mathbb{C}_{\chi_{\tilde{g}}}, Q_{\chi_{\tilde{g}}}) = (Q_{\chi_{\tilde{g}}})_{\chi_{\tilde{g}}}$ as a linear space, and hence $(Q_{\chi_{\tilde{g}}} \otimes_{W_{\varepsilon, \eta}^s(G)} E)_{\chi_{\tilde{g}}} = E$. This proves the second statement. \square

REFERENCES

- [1] Carter, R. W., Conjugacy classes in the Weyl group, *Compositio Math.* **25** (1972), 1–59.
- [2] Chari, V., Pressley, A., A guide to quantum groups, Cambridge Univ. Press (1994).
- [3] De Concini, C., Kac, V. G., Representations of quantum groups at roots of 1. Operator algebras, unitary representations, enveloping algebras, and invariant theory (Paris, 1989), *Progr. Math.* **92**, Birkhäuser Boston, Boston, MA (1990), 471–506.
- [4] De Concini, C., Kac, V. G., Procesi, C., Quantum coadjoint action, *J. Amer. Math. Soc.* **5** (1992), 151–189.
- [5] Drupieski, C. M., On injective modules and support varieties for the small quantum group, *Int. Math. Res. Not.* **2011** (2011), 2263–2294.
- [6] Curtis, C. W., Reiner, I., Representation theory of finite groups and associative algebras, Interscience Publishers (1962).
- [7] He, X., Lusztig, G., A generalization of Steinberg’s cross-section, *J. Amer. Math. Soc.* **25** (2012), 739–757.
- [8] Joseph, A., Letzter, G., Local finiteness of the adjoint action for quantized enveloping algebras, *J. Algebra* **153** (1992), 289–318.
- [9] Kawanaka, N. “Generalized Gelfand–Graev representations and Ennola duality.” In: Algebraic groups and related topics (Kyoto/Nagoya, 1983), 175–206, *Adv. Stud. Pure Math.* **6**, Amsterdam: North-Holland, 1985.
- [10] Khoroshkin, S.M., Tolstoy, V.N., Universal R–matrix for quantized (super)algebras, *Comm. Math. Phys.* **141** (1991), 599–617.
- [11] Khoroshkin, S. M., Tolstoy, V. N., The Cartan-Weyl basis and the universal R-matrix for quantum Kac-Moody algebras and superalgebras. Quantum symmetries (Clausthal, 1991), 336–351, World Sci. Publ., River Edge, NJ (1993).
- [12] Lusztig, G., On conjugacy classes in a reductive group, arXiv:1305.7168.
- [13] Premet, A., Irreducible representations of Lie algebras of reductive groups and the Kac-Weisfeiler conjecture, *Invent. Math.* **121** (1995), 79–117.
- [14] Premet, A., Special transverse slices and their enveloping algebras. With an appendix by Serge Skryabin, *Adv. Math.* **170** (2002), 1–55.
- [15] Sevostyanov, A., Algebraic group analogues of Slodowy slices and deformations of Poisson W–algebras, *Int. Math. Res. Not.* **2011** (2011), 1880–1925.
- [16] Sevostyanov, A., Conjugacy classes in Weyl groups and q-W algebras, *Adv. Math.* **228** (2011), 1315–1376.
- [17] Sevostyanov, A., Strictly transversal slices to conjugacy classes in algebraic groups, arXiv:1403.4108.
- [18] Sevostyanov, A., Localization of quantum biequivariant \mathcal{D} –modules and q-W algebras, *Proc. London. Math. Soc.* **113** (5) (2016), 583–626.
- [19] Zelobenko, D.P., Extremal cocycles on Weyl groups, *Funk. Anal. i.ego Pril.* **21**, no. 3 (1987), 11–21.

INSTITUTE OF PURE AND APPLIED MATHEMATICS, UNIVERSITY OF ABERDEEN, ABERDEEN AB24 3UE, UNITED KINGDOM, E-MAIL: A.SEVASTYANOV@ABDN.AC.UK