# POLETSKY-STESSIN HARDY SPACES ON DOMAINS BOUNDED BY AN ANALYTIC JORDAN CURVE IN $\mathbb C$

SIBEL ŞAHİN

ABSTRACT. We study Poletsky-Stessin Hardy spaces that are generated by continuous, subharmonic exhaustion functions on a domain  $\Omega \subset \mathbb{C}$ , that is bounded by an analytic Jordan curve. Different from Poletsky & Stessin's work these exhaustion functions are not necessarily harmonic outside of a compact set but have finite Monge-Ampére mass. We have showed that functions belonging to Poletsky-Stessin Hardy spaces have a factorization analogous to classical Hardy spaces and the algebra  $A(\Omega)$  is dense in these spaces as in the classical case ; however, contrary to the classical Hardy spaces, composition operators with analytic symbols on these Poletsky-Stessin Hardy spaces need not always be bounded.

## INTRODUCTION

The aim of this paper is to study the behavior of the Hardy spaces  $H^p_u(\Omega)$  defined by Poletsky-Stessin in 2008, in a setting where  $\Omega$  is a domain in  $\mathbb C$  containing 0. bounded by an analytic Jordan curve and u is a continuous subharmonic exhaustion function for  $\Omega$  which has finite Monge-Ampére mass but not necessarily harmonic out of a compact set. The main interest in Poletsky- Stessin's work ([4]) is to study the behavior of these new spaces when the exhaustion function, say  $\varphi$  is harmonic out of a compact set (i.e the measure  $\Delta \varphi$  has compact support) whereas our main focus will be on these Poletsky-Stessin Hardy spaces where the associated exhaustion function u has finite mass but the measure  $\Delta u$  does not have compact support. One of the main consequences of this choice of exhaustion function is that the new Hardy spaces and the classical ones do not coincide and so that we have new Banach spaces to be explored inside the classical Hardy spaces. Throughout this study we will discuss some properties of classical Hardy spaces that can be generalized to this extent like factorization and approximation also as far as the composition operators on these Hardy spaces are concerned we will see that these classes behave different than the classical Hardy spaces.

Now we will briefly discuss the content of this paper. In Section 1 we will first give the definitions of classical spaces on  $\Omega$ , namely Hardy and Hardy-Smirnov classes and then we will give the necessary background information about the construction of the Poletsky-Stessin Hardy spaces  $H_u^p(\Omega)$ . We will also introduce Demailly's Lelong-Jensen formula which is one of the most powerful tools that we use in our results. We will then compare all those classes of holomorphic functions and will see that Poletsky-Stessin Hardy spaces are not always equal to the classical ones. Main results of this study are given in the following sections, in the first one we will characterize Poletsky-Stessin Hardy classes  $H_u^p(\Omega)$  through their boundary values

Date: March 10, 2013.

Key words and phrases. Monge-Ampére measure, Hardy Space, Jordan domain, exhaustion.

and the corresponding Monge-Ampére boundary measure. This boundary value characterization will enable us to prove factorization properties analogous to classical case and next we will show the algebra  $A(\Omega)$  is dense in these spaces. Finally we will examine the composition operators induced by holomorphic self maps and we will see that even on the simplest of such domains , namely the unit disc, not all composition operators are bounded contrary to the classical Hardy space case.

## Acknowledgments

This paper is based on the first part of my PhD dissertation and I would like to express my sincere gratitude to my advisor Prof.Aydm Aytuna for his valuable guidance, support and his endless patience. This study would not have been possible without his help. I would also like to thank Prof. Evgeny A.Poletsky for the valuable discussions that we had during my visit at Syracuse University and I would like to thank Dr. Nihat Gökhan Göğüş and Dr. Muhammed Ali Alan for their suggestions and comments about this study.

# 1. Poletsky-Stessin Hardy Spaces on Domains Bounded By An Analytic Jordan Curve

In this section we will give the preliminary definitions and some important results that we will use throughout this paper. Let  $\Omega$  be a domain in  $\mathbb{C}$  containing 0 and bounded by an analytic Jordan curve. There are different classes of holomorphic functions studied by various authors on domains in  $\mathbb{C}$ , here we will mention first the classical Hardy space  $H^p(\Omega)$  which is defined as follows:

# (1) $H^{p}(\Omega) = \{ f \in \mathcal{O}(\Omega) \mid |f|^{p} has \ a \ harmonic \ majorant \ in \ \Omega \}$

The second class we will mention is the Hardy-Smirnov class  $E^p(\Omega)$ : A holomorphic function f on  $\Omega$  is said to be of class  $E^p(\Omega)$  if there exists a sequence of rectifiable Jordan curves  $C_1, C_2, \ldots$  in  $\Omega$  tending to boundary in the sense that  $C_n$  eventually surrounds each compact subdomain of  $\Omega$  such that

$$\int_{C_n} |f(z)|^p ds \le M < \infty$$

In ([4]), Poletsky & Stessin introduced new Hardy type classes of holomorphic functions on a large class of domains in  $\mathbb{C}^n$  (see also ([1])) but our focus will be the domains in  $\mathbb{C}$  containing 0 and bounded by an analytic Jordan curve. Before defining these new classes let us first give some preliminary definitions. Let  $\varphi$ :  $\Omega \to [-\infty, 0)$  be a negative, continuous, subharmonic exhaustion function for  $\Omega$ . Following Demailly define pseudoball:

$$B(r) = \{ z \in \Omega : \varphi(z) < r \} \quad , \quad r \in [-\infty, 0) \}$$

and pseudosphere:

$$S(r) = \{ z \in \Omega : \varphi(z) = r \} \quad , \quad r \in [-\infty, 0),$$

and set

$$\varphi_r = \max\{\varphi, r\} \quad , \quad r \in (-\infty, 0)$$

In ([2]) Demailly introduced the Monge-Ampére measures in the sense of currents as :

$$\mu_{\varphi,r} = (dd^c \varphi_r) - \chi_{\Omega \setminus B(r)} (dd^c \varphi) \quad r \in (-\infty, 0)$$

These measures are supported on S(r) and  $dd^c \varphi = \frac{1}{4} \triangle \varphi dz d\overline{z}$  in  $\mathbb{C}$ . Demailly had proved a general formula in ([3]) called Lelong-Jensen Formula which is a fundamental tool used in most of the results and in our setting it is given as follows:

**Theorem 1.1.** Let r < 0 and  $\phi$  be a subharmonic function on  $\Omega$  then for any negative, continuous, subharmonic exhaustion function u

(2) 
$$\int_{S_u(r)} \phi d\mu_{u,r} = \int_{B_u(r)} \phi (dd^c u) + \int_{B_u(r)} (r-u) dd^c \phi$$

One of the main results of this study is to characterize the Poletsky-Stessin Hardy spaces through their boundary values and for this we also need boundary measures introduced by Demailly in ([3]). Now let  $\varphi : \Omega \to [-\infty, 0)$  be a continuous subharmonic exhaustion for  $\Omega$ . And suppose that the total Monge-Ampére mass is finite, i.e.

(3) 
$$MA(\varphi) = \int_{\Omega} (dd^c \varphi) < \infty$$

Then when r approaches to 0,  $\mu_r$  converge to a positive measure  $\mu$  weak\*-ly on  $\Omega$  with total mass  $\int_{\Omega} (dd^c \varphi)$  and supported on  $\partial \Omega$ . One calls the measure  $\mu$  as the **Monge-Ampére measure on the boundary associated with the exhaustion**  $\varphi$ .

Now we can introduce the Poletsky-Stessin Hardy classes which will be our main focus throughout this study, in ([4]), Poletsky & Stessin gave the definition of new Hardy spaces using Monge-Ampére measures as :

**Definition 1.**  $H^p_{\varphi}(\Omega)$ , p > 0, is the space of all holomorphic functions f on  $\Omega$  such for

$$\limsup_{r \to 0^-} \int_{S_{\varphi,}(r)} |f|^p d\mu_{\varphi,r} < \infty$$

The norm on these spaces is given by:

$$\|f\|_{H^p_{\varphi}} = \left(\lim_{r \to 0^-} \int_{S_{\varphi}(r)} |f|^p d\mu_{\varphi,r}\right)^{\frac{1}{p}}$$

and with respect to these norm the spaces  $H^p_{\varphi}(\Omega)$  are Banach spaces ([4],pg:16). Moreover on these Banach spaces point evaluations are continuous.

Now we will compare new Hardy spaces with the classical Hardy spaces given as in (1):

**Theorem 1.2.** Let  $\Omega$  be a domain in  $\mathbb{C}$  containing 0 and bounded by an analytic Jordan curve. Suppose  $\varphi$  is a continuous, negative, subharmonic exhaustion function for  $\Omega$  such that  $\varphi$  is harmonic out of a compact set  $K \subset \Omega$ . Then for a holomorphic function  $f \in \mathcal{O}(\Omega)$ ,  $f \in H^p_{\varphi}(\Omega)$  if and only if  $|f|^p$  has a harmonic majorant.

*Proof.* Let  $|f|^p$  has a harmonic majorant u on  $\Omega$ . Then

(4) 
$$\int_{S(r)} |f|^p d\mu_{\varphi,r} \le \int_{S(r)} u d\mu_{\varphi,r} = \int_{B(r)} u (dd^c \varphi)$$

by Lelong-Jensen formula and we know that  $\varphi$  is harmonic out of the compact set K so

(5) 
$$\int_{B(r)} u(dd^c \varphi) \le \int_K u(dd^c \varphi) \le C_K ||u||_{L^{\infty}(K)}$$

for some constant  $C_K$  and this bound is independent of r. Hence

(6) 
$$\sup_{r<0} \int_{S(r)} |f|^p d\mu_{\varphi,r} \le M < \infty$$

for some M

 $\Rightarrow f\in H^p_\varphi(\Omega).$  For the converse, suppose  $f\in H^p_\varphi(\Omega)$  and  $|f|^p$  has no harmonic majorant. Then

(7) 
$$\frac{1}{2\pi} \int_{\Omega} (-g_{\Omega}(z,w)) \Delta |f|^p = \infty$$

where  $g_{\Omega}(z, w)$  is the Green function of the domain  $\Omega$  ([6]). Then from Lelong-Jensen formula

$$\frac{1}{2\pi} \int_{S(r)} |f|^p d\mu_{\varphi,r} \ge \frac{1}{2\pi} \int_{B(r)} (r-\varphi) \Delta |f|^p$$

Note that left hand side is bounded independent from r since  $f \in H^p_{\varphi}(\Omega)$ . Let us take a compact set  $F \subset \Omega$  containing the support of  $\Delta \varphi$  and  $\{w\}$  such that both  $\varphi$  and  $g_{\Omega}(z, w)$  are bounded on  $\partial F$  and  $bg_{\Omega}(z, w) \leq \varphi \leq cg_{\Omega}(z, w)$  holds on  $\partial F$  for some numbers b, c > 0. By the maximality of both  $\varphi$  and  $g_{\Omega}(z, w)$  on  $\Omega \setminus F$ this inequality holds on  $\Omega \setminus F$  also hence near boundary we have

$$\varphi \le cg_{\Omega}(z,w)$$

Then by Theorem 3.1 of ([4], pg:11) for a positive constant a > 0 we have the following

$$\frac{1}{2\pi} \int_{B(ar)} (ar - g_{\Omega}(z, w)) \Delta |f|^p \le \frac{1}{2\pi} \int_{S(ar)} |f|^p d\mu_{g,r} \le \frac{1}{2\pi} \int_{S(r)} |f|^p d\mu_{\varphi,r}$$

and as  $r \to 0$  by Fatou lemma we have

$$\int_{\mathbb{D}} (-g_{\Omega}(z,w))\Delta |f|^p \le \lim_{r \to 0} \frac{1}{2\pi} \int_{B(ar)} (ar - g_{\Omega}(z,w))\Delta |f|^p \le \lim_{r \to 0} \frac{1}{2\pi} \int_{S(r)} |f|^p d\mu_{\varphi,r}$$

the last limit is bounded since  $f \in H^p_{\varphi}(\Omega)$  but the first integral goes to infinity by (7). This contradiction proves the assertion therefore  $|f|^p$  has a harmonic majorant.

**Corollary 1.1.** Let  $\Omega$  be a domain in  $\mathbb{C}$  containing 0 and bounded by an analytic Jordan curve. Suppose  $g_{\Omega}(z, w)$  is the Green function of  $\Omega$  with the logarithmic pole at  $w \in \Omega$ . Then  $H^p(\Omega) = H^p_{g_{\Omega}}(\Omega)$  for  $p \ge 1$ .

*Proof.* The Green function  $g_{\Omega}(z, w)$  is harmonic out of the compact set  $\{w\}$  by definition so by previous theorem we have  $f \in H^p_{g_{\Omega}}(\Omega)$  is equivalent to the condition that  $|f|^p$  has a harmonic majorant which is by definition means  $f \in H^p(\Omega)$ .  $\Box$ 

Now we would like to compare Poletsky-Stessin Hardy classes with the classical Hardy and Hardy-Smirnov spaces when  $\Omega$  is a domain in  $\mathbb{C}$  that is bounded by an analytic Jordan curve. First of all on  $\Omega$  we have  $E^p(\Omega) = H^p(\Omega)$  by (Theorem 10.2,[7]) and using Theorem 1.1 we see that for any exhaustion function  $\varphi$  which is harmonic out of a compact set we have  $E^p(\Omega) = H^p(\Omega) = H^p_{\varphi}(\Omega)$ . However this need **not** be the case when our exhaustion function u has finite Monge-Ampére mass yet not harmonic out of a compact set. When an exhaustion function u has finite Monge-Ampére mass then by (Theorem 3.1, [4]) we have  $H^p_u(\Omega) \subset H^p_{g_\Omega}(\Omega) = H^p(\Omega)$ where the last equality is given in Corollary 1.1 however by explicitly constructing an exhaustion function u on the unit disc  $\mathbb{D}$ , we will show that  $H^p_u(\Omega)$  need not be equal to  $H^p(\Omega)$ :

**Theorem 1.3.** There exists an exhaustion function u with finite Monge-Ampére mass such that the Hardy space  $H^p_u(\mathbb{D}) \subsetneq H^p(\mathbb{D})$ .

*Proof.* (Without loss of generality assume p=1) In order to prove this result we will first construct an exhaustion function u with finite mass:

Let  $\mathbb{D}$  be the unit disc in  $\mathbb{C}$ , and let  $\rho$  be the solution of the Dirichlet problem in the unit disc such that  $\Delta \rho = 0$  in  $\mathbb{D}$  and  $\rho = f$  on  $\partial \mathbb{D}$  where  $f(z) = -(1-x)^{\frac{3}{4}}$ . Then define  $u = f - \rho$ . We will first show that u is a continuous, subharmonic, exhaustion function for  $\mathbb{D}$ :

*u* is continuous:  $u(z) = f(z) + \int_{\partial \mathbb{D}} P(z, e^{i\theta})(1 - \cos \theta)^{\frac{3}{4}} d\theta$  and both parts on the right hand side are in  $C^2(\mathbb{D}) \cap C(\mathbb{D})$  hence *u* is a continuous function.

*u* is subharmonic: *u* is a  $C^2(\mathbb{D})$  function and  $\Delta u = \Delta(-(1-x)^{\frac{3}{4}}) = (1-x)^{\frac{-5}{4}} \ge 0$  hence *u* is subharmonic.

u is an exhaustion: For this we should show that  $A_c = \{x \mid u(x) < c\}$  is relatively compact in  $\mathbb{D}$  for all c < 0. Suppose not then there exists a sequence  $x_n$  in  $A_c$  such that it has a subsequence  $x_{n_k}$  converging to boundary of  $\mathbb{D}$ . Now  $x_{n_k} \to x$ , |x| = 1and u is continuous so  $u(x_{n_k}) \to u(x)$  but  $u(x_{n_k}) < c$  so u(x) < c. This contradicts the fact that u = 0 on the boundary. Hence  $A_c$  is relatively compact in  $\mathbb{D}$  for all cand u is an exhaustion.

Since we have a negative, continuous, subharmonic exhaustion function in  $\mathbb{D}$  we can define Monge-Ampére measure  $\mu_u$  associated with u and we will show that total mass  $\|\mu_u\|$ , of  $\mu_u$  is finite :

$$\|\mu_u\| = \int_{\mathbb{D}} dd^c u = \int_{-1}^1 \int_{-\sqrt{(1-x)(1+x)}}^{\sqrt{(1-x)(1+x)}} (1-x)^{-\frac{5}{4}} dy dx = \int_{-1}^1 2(1+x)^{\frac{1}{2}} (1-x)^{-\frac{3}{4}} dx$$
$$\leq 2\sqrt{2} \int_{-1}^1 (1-x)^{-\frac{3}{4}} dx$$

say t = 1 - x then

$$= 2\sqrt{2} \int_0^2 \frac{1}{t^{\frac{3}{4}}} dt < \infty$$

Hence  $\mu_u$  has finite mass.

We know that for any continuous, negative, subharmonic exhaustion function u the Hardy space  $H^p_u(\mathbb{D}) \subset H^1(\mathbb{D})$  ([4],pg:13) but now we will show that the converse is not true by using the u(z) that we constructed above as exhaustion. Take  $F(z) = \frac{1}{(1-z)^{2q}}$  for  $q < \frac{1}{2}$ 

First of all we want to show that  $F(z) \in H^1(\mathbb{D})$  so we will show the following growth condition is satisfied:

$$\sup_{0 < r < 1} \int_{0}^{2\pi} |F(re^{i\theta}|d\theta < \infty \quad \text{for} \quad 2q < 1$$
$$|F(z)| = \frac{1}{|1 - re^{i\theta}|^{2q}} = \frac{1}{(1 + r^2 - 2Re(re^{i\theta}))^q} = \frac{1}{(1 + r^2 - 2r\cos(\theta))^q}$$

Now,

$$\int_0^{2\pi} \frac{1}{(1+r^2-2r\cos(\theta))^q} d\theta = \int_0^{2\pi} \frac{1}{(\sin^2\theta + (\cos\theta - r)^2)^q} d\theta \le \int_0^{2\pi} \frac{1}{(\cos\theta - r)^{2q}} d\theta$$
  
and as  $r \to 1$  this integral converges for  $2q < 1$  and

and as  $r \to 1$  this integral converges for 2q < 1 and

$$\sup_{0 < r < 1} \int_0^{2\pi} |F(re^{i\theta}|d\theta < \infty \quad \text{for} \quad q < \frac{1}{2}$$

Now we will show that  $F(z) \notin H^1_u(\mathbb{D})$ :

$$F(z)| = \frac{1}{|1-z|^{2q}} = \frac{1}{((1-x)^2 + y^2)^q} = ((1-x)^2 + y^2)^{-q}$$

from the Lelong-Jensen formula,

$$\int_{S(r)} |F(z)| d\mu_{u,r} = \int_{B(r)} |F(z)| \Delta u + \int_{B(r)} (r-u) \Delta |F(z)|$$

second integral on the right hand side is non-negative so

$$\int_{S(r)} |F(z)| d\mu_{u,r} \ge \int_{B(r)} |F(z)| \Delta u$$

now as  $r \to 0$ 

$$\|F(z)\|_{H^1_u} \ge \lim_{r \to 0} \int_{B(r)} |F(z)| \triangle u \ge \int_{\mathbb{D}} |F(z)| \triangle u$$

where the last inequality is due to Fatou lemma and on  $\mathbb{D}$ ,  $y^2 \leq 1 - x^2 \leq 2(1 - x)$  so

$$\begin{split} \|F(z)\|_{H^{1}_{u}} &\geq \int_{\mathbb{D}} |F(z)| \triangle u = \int_{-1}^{1} \int_{-\sqrt{(1-x)(1+x)}}^{\sqrt{(1-x)(1+x)}} ((1-x)^{2} + y^{2})^{-q} (1-x)^{-\frac{5}{4}} dy dx \\ &\geq \int_{-1}^{1} \int_{-\sqrt{(1-x)(1+x)}}^{\sqrt{(1-x)(1+x)}} ((1-x)^{2} + 2(1-x))^{-q} (1-x)^{-\frac{5}{4}} dy dx \\ &\geq \int_{-1}^{1} (1-x)^{\frac{1}{2}} (1-x)^{-q} (3-x)^{-q} (1-x)^{-\frac{5}{4}} dx \\ &\geq \int_{-1}^{1} \frac{1}{(1-x)^{q+\frac{3}{4}}} dx \end{split}$$

so for  $q > \frac{1}{4}$  this integral diverges. Hence for  $\frac{1}{4} < q < \frac{1}{2}$   $F(z) \in H^1(\mathbb{D})$  but  $F(z) \notin H^1_u(\mathbb{D})$   $\Box$ 

Remark 1. Moreover from this result we also deduce that if u is an arbitrary exhaustion function with finite Monge-Ampére mass  $H^p_u(\Omega)$ 's are not always closed subspaces of  $H^p(\Omega)$  because as Banach spaces the inclusion  $H^p_u(\Omega) \hookrightarrow H^p(\Omega)$  is continuous which can be deduced from Closed Graph Theorem and the fact that point evaluations are continuous ([4]). However the range is not closed because range includes all bounded functions (hence polynomials) and polynomials are dense in  $H^p(\Omega)$  but  $H^p(\Omega) \neq H^p_u(\Omega)$  in general.

# 2. BOUNDARY MONGE-AMPÉRE MEASURE & BOUNDARY VALUE CHARACTERIZATION OF POLETSKY-STESSIN HARDY SPACES

Let  $\Omega$  be a domain in  $\mathbb{C}$  containing 0 and bounded by an analytic Jordan curve and u be a continuous, negative, subharmonic exhaustion function for  $\Omega$  with finite Monge-Ampére mass. In the classical Hardy space theory on the unit disc  $\mathbb{D}$  we can characterize the  $H^p$  spaces through their boundary values inside the  $L^p$  spaces of the unit circle and since we have  $H^p_u(\Omega) \subset H^p(\Omega)$ , any holomorphic function  $f \in H^p_u(\Omega)$ has the boundary value function  $f^*$  from the classical theory (Theorem 10, [13]). In this section we will give an analogous characterization of the Poletsky-Stessin Hardy spaces through these boundary value functions and boundary Monge-Ampére measure. Moreover we will show the relation between boundary Monge-Ampére measure and Euclidean measure on the boundary  $\partial\Omega$ .

**Proposition 2.1.** Let u be a continuous, negative, subharmonic exhaustion function for  $\Omega$  with finite Monge-Ampére mass then the boundary Monge-Ampére measure  $\mu_u$  and the Euclidean measure on  $\partial\Omega$  are mutually absolutely continuous.

*Proof.* Let  $\varphi$  be a continuous function on  $\partial \Omega$  and let the Poisson integral of  $\varphi$  be

$$H(z) = \int_{\partial\Omega} P(z,\xi)\varphi(\xi)d\sigma(\xi)$$

then by Lelong-Jensen formula we have

$$\int_{\partial\Omega} \varphi d\mu_u = \int_{\Omega} H(z) dd^c u = \int_{\Omega} \int_{\partial\Omega} P(z,\xi) \varphi(\xi) d\sigma(\xi) dd^c u$$

and since  $\varphi$  is a continuous function on the boundary and  $\mu_u$  has finite mass we can use Fubini theorem to get

$$\int_{\partial\Omega} \varphi d\mu_u = \int_{\Omega} \int_{\partial\Omega} P(z,\xi) \varphi(\xi) d\sigma(\xi) dd^c u = \int_{\partial\Omega} \varphi(\xi) \left( \int_{\Omega} P(z,\xi) dd^c u(z) \right) d\sigma(\xi)$$
 Now define

Now define

$$\beta(\xi) = \int_{\Omega} P(z,\xi) dd^c u(z)$$

We will show that  $\beta(\xi)$  is  $d\sigma$ -integrable: First we see that  $\beta(\xi) \ge 0$  and

$$\int_{\partial\Omega} |\beta(\xi)| d\sigma(\xi) = \int_{\partial\Omega} \beta(\xi) d\sigma(\xi) = \int_{\partial\Omega} \int_{\Omega} P(z,\xi) dd^c u(z) d\sigma(\xi) = \int_{\Omega} \int_{\partial\Omega} P(z,\xi) d\sigma(\xi) dd^c u(z)$$

and since

$$\int_{\partial\Omega} P(z,\xi) d\sigma(\xi) = 1$$

we have

$$\int_{\partial\Omega} |\beta(\xi)| d\sigma(\xi) = \int_{\Omega} dd^c u = MA(u) < \infty$$

Hence  $\beta \in L^1(d\sigma)$  and we have

$$\int_{\partial\Omega}\varphi d\mu_u = \int_{\partial\Omega}\varphi\beta d\sigma$$

 $\Rightarrow d\mu_u = \beta d\sigma$ 

*Remark* 2. From the previous proof we see that for a fixed  $\xi \in \partial\Omega$ ,  $\beta_{\tilde{r}}(\xi)$  which is defined as

$$\beta_{\tilde{r}}(\xi) = \int_{B_u(\tilde{r})} P(z,\xi) dd^c u = \int_{S_u(\tilde{r})} P(z,\xi) d\mu_{u,\tilde{r}}$$

converges to  $\beta(\xi) = \int_{\Omega} P(z,\xi) dd^c u(z)$  by Monotone Convergence Theorem since  $\beta_{\tilde{r}} \geq 0$  for all  $\tilde{r}$  and  $\beta_{\tilde{r}}$  is increasing with respect to  $\tilde{r}$ .

Now we will give the characterization of Poletsky-Stessin Hardy spaces  $H^p_u(\Omega)$  through boundary value functions:

**Theorem 2.1.** Let  $f \in H^p(\Omega)$  be a holomorphic function and u be a continuous, negative, subharmonic exhaustion function for  $\Omega$ . Then  $f \in H^p_u(\Omega)$  if and only if  $f^* \in L^p(d\mu_u)$  for  $1 \le p < \infty$ . Moreover  $\|f^*\|_{L^p(d\mu_u)} = \|f\|_{H^p_u(\Omega)}$ .

*Proof.* (Without loss of generality assume p=1) Now let  $f \in H^1_u(\Omega) \subset H^1(\Omega)$  we want to show that  $f^* \in L^1(d\mu_u)$ . First of all

$$\int_{\partial\Omega} |f^*(\xi)| d\mu_u = \int_{\partial\Omega} |f^*(\xi)| \left( \int_{\Omega} P(z,\xi) dd^c u(z) \right) d\sigma(\xi)$$

and using Fubini-Tonelli theorem we change the order of integration and get

$$\int_{\Omega} \left( \underbrace{\int_{\partial \Omega} |f^*(\xi)| P(z,\xi) d\sigma(\xi)}_{H(z)} \right) dd^c u(z)$$

Then the harmonic function being the Poisson integral of  $|f^*|$  is the least harmonic majorant of |f| so by ([6], Theorem 4.5.4) we have

$$H(z) = |f(z)| - \int_{\Omega} g_{\Omega}(w, z) dd^{c} |f(w)|$$

where  $g_{\Omega}(w, z)$  is the Green function of  $\Omega$  with the logarithmic pole at the point z and by ([3], Theorem 4.14),  $g_{\Omega}(w, z)$  is continuous on  $\overline{\Omega}$  and subharmonic in  $\Omega$  hence we have

$$\int_{\Omega} H(z) dd^{c} u = \int_{\Omega} |f(z)| dd^{c} u - \int_{\Omega} \left( \int_{\Omega} g_{\Omega}(w, z) dd^{c} u \right) dd^{c} |f(w)|$$

Now using the boundary version of Lelong- Jensen formula ([3], Theorem 3.3) we get

$$\int_{\Omega} g_{\Omega}(w, z) dd^{c} u(z) = \int_{\Omega} u(z) dd^{c} g_{\Omega}(w, z) = u(w)$$

therefore

$$\begin{split} \int_{\partial\Omega} |f^*(\xi)| d\mu_u &= \int_{\Omega} H(z) dd^c u = \int_{\Omega} |f(z)| dd^c u - \int_{\Omega} u(z) dd^c |f| = \|f\|_{H^1_u(\Omega)} < \infty \end{split}$$
 so we have  $f^* \in L^1(d\mu_u).$ 

For the converse since  $f \in H^1(\Omega)$  we have

$$f(z) = \int_{\Omega} P(z,\xi) f^*(\xi) d\sigma(\xi)$$

now

$$\int_{S_u(r)} |f(z)| d\mu_{u,r} = \int_{S_u(r)} \left| \int_{\Omega} P(z,\xi) f^*(\xi) d\sigma(\xi) \right| d\mu_{u,r} \le \int_{\Omega} \left( \int_{S_u(r)} P(z,\xi) d\mu_{u,r} \right) |f^*(\xi)| d\sigma(\xi)$$

by Remark 2 we know  $\int_{S_u(r)} P(z,\xi) d\mu_{u,r}$  is increasing and using Monotone Convergence Theorem as  $r \to 0$  we get

$$\|f\|_{H^1_u(\Omega)} \le \int_{\partial\Omega} |f^*(\xi)| d\mu_u(\xi) < \infty$$

$$\Pi^1_u(\Omega) \qquad \Box$$

so  $f \in H^1_u(\Omega)$ 

Remark 3. The existence of boundary values for holomorphic functions depends on conditions related to domain itself and the growth of the functions. However throughout this study we have been exploiting the fact that Poletsky-Stessin Hardy classes are inside the classical ones so we already have the non tangential boundary values. In his 2011 paper ([5]) Poletsky gave definitions for two types of boundary values namely, strong and weak limit values for sequences of functions in an abstract setting that involves sequences of weakly convergent measures on increasing chains of compact sets (See [5] for details). If we specify his abstract setup to the functions  $f \in H^p_u(\Omega)$  together with Monge-Ampére measures  $d\mu_{u,r}$  and the boundary Monge-Ampére measure  $d\mu_u$ , it is possible to prove that the boundary value function  $f^*$  is both weak and strong limit value for  $f \in H^p_u(\Omega)$  (Details will be given in the thesis [18])

## 3. FACTORIZATION

Let  $\Omega$  be a domain in  $\mathbb{C}$  containing 0 and bounded by an analytic Jordan curve. Suppose  $\psi : \mathbb{D} \to \Omega$  is the conformal map such that  $\psi(0) = 0$  and  $\varphi = \psi^{-1}$ . By Carathéodory theorem for a domain like  $\Omega$  we have  $0 < m \leq |\psi'| \leq M < \infty$  for some m, M > 0. Following ([16]) we have the following :

**Definition 2.** A holomorphic function h on  $\Omega$  is  $\Omega$ -inner if  $h \circ \psi$  is inner in the classical sense i.e.  $|h \circ \psi| = 1$  for almost all  $\xi \in \partial \mathbb{D}$  and  $\Omega$ -outer if  $h \circ \psi$  is outer in the classical sense i.e.

$$\log |h \circ \psi(0)| = \int_{\partial \mathbb{D}} \log |h \circ \psi| d\sigma$$

For the classical Hardy space on the unit disc  $\mathbb{D}$  we have the following canonical factorization theorem ([7],pg:24):

**Theorem 3.1.** Every function  $f \neq 0$  of class  $H^p(\mathbb{D})$  has a unique factorization of the form f = BSF where B is a Blaschke product, S is a singular inner function and F is an outer function for the class  $H^p(\mathbb{D})$ .

Using this result and the conformal mapping the following is shown in ([16]):

**Proposition 3.1.** Every  $f \in H^p(\Omega)$  can be factored uniquely up to a unimodular constant as f = IF where I is  $\Omega$ -inner and F is  $\Omega$ -outer.

Inspired by this result we have the following corollary

**Corollary 3.1.** Let  $f \in H^p_u(\Omega)$  where u is a continuous exhaustion function with finite Monge-Ampére mass. Then f can be factored as f = IF where I is  $\Omega$ -inner and F is  $\Omega$ -outer. Moreover  $I \in H^p_u(\Omega)$  and  $F \in H^p_u(\Omega)$ .

*Proof.* First of all since  $H^p_u(\Omega) \subset H^p(\Omega)$  by the above proposition it is obvious that f can be factored as f = IF where I is  $\Omega$ -inner and F is  $\Omega$ -outer. Now define the

measure  $\hat{\mu}$  on  $\partial \mathbb{D}$  as  $\hat{\mu}(E) = \mu_u(\psi(E))$  for any measurable set  $E \subset \partial \mathbb{D}$  then it is clear from the change of variables formula that  $|I \circ \psi| = 1 \ d\hat{\mu}$ -a.e. so

$$\int_{\partial\Omega} |I| d\mu_u = \int_{\partial\mathbb{D}} |I \circ \psi| d\hat{\mu} = \int_{\partial\mathbb{D}} d\hat{\mu} = \int_{\partial\Omega} d\mu_u = MA(u) < \infty$$

since u has finite Monge-Ampére mass and for the outer part

$$\int_{\partial\Omega} |F| d\mu_u = \int_{\partial\mathbb{D}} |F \circ \psi| d\hat{\mu} = \int_{\partial\mathbb{D}} |f^* \circ \psi| d\hat{\mu} = \int_{\partial\Omega} |f^*| d\mu_u < \infty$$
  
since  $f \in H^p_u(\Omega)$ . Hence we have  $I \in H^p_u(\Omega)$  and  $F \in H^p_u(\Omega)$ .

In fact in the particular case of unit disc  $\mathbb D$  we can say more about this factorization :

**Theorem 3.2.** Let f be analytic function in  $\mathbb{D}$  such that  $f \in H^p_u(\mathbb{D})$ . Then f can be factored into a Blaschke product B, a singular inner function S and an outer function F such that  $B, S, F \in H^p_u(\mathbb{D})$  for  $1 \le p < \infty$ .

*Proof.* (Without loss of generality assume p=1) Let  $f \in H^1_u(\mathbb{D})$  then  $f \in H^1(\mathbb{D})$ so f has canonical decomposition f = BSF where B is a Blaschke product, S is a singular inner function and F is an outer function. The exhaustion u has finite mass so bounded functions belong to class  $H^1_u(\mathbb{D})$ . Since B and S are bounded ([7],pg:24),  $B, S \in H^1_u(\mathbb{D})$ . As the outer function F(z) is concerned, we know  $F \in H^1(\mathbb{D})$  so we have

$$F(z) = \frac{1}{2\pi} \int_0^{2\pi} P(r, \theta - t) F^*(e^{it}) dt$$

Now

 $\Rightarrow F(z)$ 

$$\int_{S(\widehat{r})} |F(z)| d\mu_{u,\widehat{r}} = \frac{1}{2\pi} \int_{S(\widehat{r})} \left| \int_0^{2\pi} P(r,\theta-t) F^*(e^{it}) dt \right| d\mu_{u,\widehat{r}}$$

$$\leq \frac{1}{2\pi} \int_0^{2\pi} \left( \int_{S(\widehat{r})} P(r,\theta-t) d\mu_{u,\widehat{r}} \right) |F^*(e^{it})| dt = \frac{1}{2\pi} \int_0^{2\pi} \left( \int_{S(\widehat{r})} P(r,\theta-t) d\mu_{u,\widehat{r}} \right) |f^*(e^{it})| dt$$
and using the previous result, monotone convergence theorem and the fact that

and using the previous result, monotone convergence theorem and the fact that  $|f^*| = |F^*|$  (dt)- a.e. we get as  $\hat{r} \to 0$ :

$$\|F(z)\|_{H^1_u(\mathbb{D})} \le \int_{\partial \mathbb{D}} |f^*(e^{it})| d\mu_u(t) < \infty$$
  
  $\in H^1_u(\mathbb{D})$ 

In the classical  $H^p$  space theory a useful tool for the proofs is that any analytic function  $f \in H^1(\mathbb{D})$  can be expressed as a product of two functions, f = gh where both factors g and  $h \in H^2(\mathbb{D})$ . Now we will show that there is a similar factorization in the spaces  $H^p_u(\Omega)$ .

**Corollary 3.2.** Suppose  $1 \le p < \infty$ ,  $f \in H^p_u(\Omega)$ ,  $f \ne 0$ . Then there is a zero-free function  $h \in H^2_u(\Omega)$  such that  $f = Ih^{\frac{2}{p}}$ . In particular every  $f \in H^1_u(\Omega)$  is a product of f = gh in which both factors are in  $H^2_u(\Omega)$ .

*Proof.* By the previous theorem  $f/I \in H^p_u(\Omega)$ . Since f/I has no zero in  $\Omega$  and  $\Omega$  is simply connected, there exists  $\varphi \in \mathcal{O}(\Omega)$  such that  $\exp(\varphi) = f/I$ . Define  $h = \exp(p\varphi/2)$  then  $h \in \mathcal{O}(\Omega)$  and  $|h|^2 = |f/I|^p$  and  $h \in H^2_u(\Omega)$  and  $f = Ih^{\frac{2}{p}}$ . To obtain f = gh for  $f \in H^1_u(\Omega)$  write  $f = Ih^2$  in the form f = (Ih)h.

#### 4. Approximation

Let  $A(\Omega)$  denote the algebra of holomorphic functions on  $\Omega$  which are continuous on  $\partial\Omega$ . We know that the algebra of holomorphic functions  $A(\Omega)$  is dense in the classical Hardy spaces when  $\Omega$  is a domain bounded by an analytic Jordan curve and we will show an analogous approximation result on  $H^p_u(\Omega)$  where u is a negative, continuous, subharmonic exhaustion function on  $\Omega$  with finite Monge-Ampére mass but before the result we should first mention some classes of analytic functions from the classical theory on the unit disc which will help us in the proof of approximation result:

**Definition 3.** An analytic function  $f \in \mathcal{O}(\mathbb{D})$  is said to be of class N if the integrals

$$\int_0^{2\pi} \log^+ |f(re^{i\theta})| d\theta$$

are bounded for r < 1.

**Definition 4.** An analytic function  $f \in \mathcal{O}(\mathbb{D})$  is in class  $N^+$  if it has the form f = BSF where B is a Blaschke product, S is a singular inner function and F is an outer function for the class N

It is clear that  $N \supset N^+ \supset H^p(\mathbb{D})$  for all p > 0.(For details see [7]). We will also use the following result (Theorem 2.11, [7]), which plays a crucial role in our approximation result:

**Theorem 4.1.** If a holomorphic function  $f \in N^+$  and  $f^* \in L^p$  for some p > 0, then  $f \in H^p(\mathbb{D})$ .

Now we can give the approximation result for the Poletsky-Stessin Hardy classes  $H^p_u(\Omega)$ :

**Theorem 4.2.** The algebra  $A(\Omega)$  is dense in  $H^p_u(\Omega)$ ,  $1 \le p < \infty$ .

*Proof.* (Case 1: p > 1)Let L be a linear functional on  $H^p_u(\Omega)$  such that L vanishes on  $A(\Omega)$ . Then  $L(f) = \int_{\partial\Omega} f^* \overline{g^*} d\mu_u$  for some non-zero  $g \in L^q(d\mu_u)$  hence  $\int_{\partial\Omega} \gamma \overline{g^*} d\mu_u = 0$  for all  $\gamma \in A(\Omega)$  (\*). Now we need the following lemma:

**Lemma 4.1.** Let  $\mu$  be a measure on the boundary  $\partial\Omega$  of  $\Omega$  which is orthogonal to  $A(\Omega)$ . Then  $\mu$  is absolutely continuous with respect to  $d\mu_{g_{\Omega,0}}$  which is the boundary Monge-Ampére measure with respect to the Green function with pole at 0.

Proof. The homomorphism "evaluation at 0" of  $A(\Omega)$  has the representing measure  $d\mu_{g_{\Omega,0}}$  on  $\partial\Omega$ . Then by generalized F. and M. Riesz Theorem (Theorem 7.6, [14]) the singular part  $\mu_s$  of  $\mu$  with respect to  $d\mu_{g_{\Omega,0}}$  is orthogonal to  $A(\Omega)$  that is  $\int_{\partial\Omega} f d\mu_s = 0$  for all  $f \in A(\Omega)$  but on a domain like  $\Omega$ ,  $A(\Omega)$  is dense in  $C(\partial\Omega)$  (Theorem 2.7, [14]) so  $\int_{\partial\Omega} g d\mu_s = 0$  for all  $g \in C(\partial\Omega)$ . Hence  $d\mu_s = 0$  and  $d\mu < < d\mu_{g_{\Omega,0}}$ .

Now by the above lemma and (\*) we have  $\overline{g^*}d\mu_u \ll d\mu_{g_{\Omega,0}}$  so by Radon-Nikodym Theorem we have  $\overline{g^*}d\mu_u = h^*d\mu_{g_{\Omega,0}}$  for some  $h^* \in L^1(d\mu_{g_{\Omega,0}})$  and on a domain like  $\Omega$  we have  $c_1d\mu_{g_{\Omega,0}} \leq d\sigma \leq c_2d\mu_{g_{\Omega,0}}$  so we have  $h^* \in L^1(d\sigma)$ . Now consider the function  $H^* = h^* \circ \psi$  on  $\partial \mathbb{D}$  then since  $h^* \in L^1(d\sigma)$  we have

 $h^* \in L^1(\partial \mathbb{D})$ , and since  $\psi(0) = 0$  and  $\mu_{g_{\Omega,0}}$  is in fact the harmonic measure we have  $d\mu_{g_{\Omega,0}}(e^{i\theta}) = d\mu_{g_{\Omega,0}}(\psi(e^{i\theta}))$  and using  $\varphi = \psi^{-1}$  we get

$$\int_{\partial \mathbb{D}} e^{in\theta} (h^* \circ \psi(e^{i\theta})) d\theta = \int_{\partial \Omega} (\varphi(z))^n h^*(z) d\mu_{g_{\Omega,0}}(z) = \int_{\partial \Omega} (\varphi(z))^n \overline{g^*} d\mu_u = 0$$

for all n as a consequence of (\*). Hence  $H^*$  is the boundary value of an  $H^1(\mathbb{D})$ function H then h which is defined as  $H = h \circ \psi$  is in the class  $E^1(\Omega)$  by the corollary of (Theorem 10.1, [7]). Moreover since  $E^1(\Omega) = H^1(\Omega)$  we have  $h \in H^1(\Omega)$  and since  $\psi(0) = 0$  we have h(0) = 0.

Now take  $\alpha \in H^p_{\mu}(\Omega)$  and consider the analytic function  $\alpha h$ 

$$\int_{\partial\Omega} |\alpha^*h^*|^{\frac{1}{2}} d\sigma \le \left(\int_{\partial\Omega} |\alpha^*| d\sigma\right)^{\frac{1}{2}} \left(\int_{\partial\Omega} |h^*| d\sigma\right)^{\frac{1}{2}} \le \|\alpha\|_{H^1}^{\frac{1}{2}} \|h\|_{H^1}^{\frac{1}{2}}$$

since  $h \in H^1(\Omega)$  and  $\alpha \in H^p_u(\Omega) \subset H^p(\Omega) \subset H^1(\Omega)$  so  $\alpha h \in H^{\frac{1}{2}}(\Omega)$ . On the other hand

$$\int_{\partial\Omega} |\alpha^* h^*| d\sigma \le c_2 \int_{\partial\Omega} |\alpha^*| |h^*| d\mu_{g_{\Omega,0}} = c_2 \int_{\partial\Omega} |\alpha^*| |\overline{g^*}| d\mu_u \le \left( \int_{\partial\Omega} |\alpha^*|^p d\mu_u \right)^{\frac{1}{p}} \left( \int_{\partial\Omega} |g^*|^q d\mu_u \right)^{\frac{1}{q}} < \infty$$

since  $\alpha \in H^p_u(\Omega)$  and  $g^* \in L^q(d\mu_u)$ . Hence we have  $\alpha^*h^* \in L^1(\partial\Omega)$ . Now since  $\alpha h \in H^{\frac{1}{2}}(\Omega)$  by the corollary of (Theorem 10.1, [7]) the function  $AH = \alpha h(\psi(w))[\psi'(w)]^2 \in H^{\frac{1}{2}}(\mathbb{D}) \subset N^+$  but since  $\alpha^*h^* \in L^1(\partial\Omega)$ ,  $AH^* \in L^1(\partial\mathbb{D})$ hence again by the corollary of (Theorem 10.1, [7]) we have  $\alpha h \in H^1(\Omega)$ . Finally

$$0 = \alpha h(0) = \int_{\partial \Omega} \alpha h d\mu_{g_{\Omega,0}} = \int_{\partial \Omega} \alpha \overline{g^*} d\mu_u = L(\alpha)$$

for all  $\alpha \in H^p_u(\Omega)$  hence  $A(\Omega)$  is dense in  $H^p_u(\Omega)$ .

(Case 2: p = 1): By the previous corollary we know that if  $f \in H^1_u(\Omega)$  then we can factor it out like f = gh where  $g, h \in H^2_u(\Omega)$  and from the first part of the proof we know there exists sequences  $\{g_n\}, \{h_n\} \in A(\Omega)$  such that  $\{g_n\} \to g$  and  $\{h_n\} \to h$  in  $H^2_u(\Omega)$ . Now for f consider the sequence  $\{g_nh_n\}$  then,

$$\int_{\partial\Omega} |f - g_n h_n| d\mu_u = \int_{\partial\Omega} |gh - g_n h_n| d\mu_u = \int_{\partial\Omega} |gh - gh_n + gh_n - g_n h_n| d\mu_u$$

$$\leq \left(\int_{\partial\Omega} |g|^2 d\mu_u\right)^{\frac{1}{2}} \left(\int_{\partial\Omega} |h - h_n|^2 d\mu_u\right)^{\frac{1}{2}} + \left(\int_{\partial\Omega} |h_n|^2 d\mu_u\right)^{\frac{1}{2}} \left(\int_{\partial\Omega} |g - g_n|^2 d\mu_u\right)^{\frac{1}{2}}$$
But right hand side goes to 0 singe  $\{g_n\} \to g$  and  $\{h_n\} \to h$  in  $H^2(\Omega)$  have

But right hand side goes to 0 since  $\{g_n\} \to g$  and  $\{h_n\} \to h$  in  $H^2_u(\Omega)$  hence  $\{g_nh_n\} \in A(\Omega)$  converges to f in  $H^1_u(\Omega)$ .

Combining these two cases we see that  $A(\Omega)$  is dense in  $H^p_u(\Omega)$  for  $1 \le p < \infty$ .

## 5. Composition Operators With Analytic Symbols

Let  $\phi : \Omega \to \Omega$  be a holomorphic self map of  $\Omega$ . The linear composition operator induced by the symbol  $\phi$  is defined by  $C_{\phi}(f) = f \circ \phi$ ,  $f \in \mathcal{O}(\Omega)$ . In 2003 Shapiro and Smith ([17]) showed that every holomorphic self map  $\phi$  of  $\Omega$  induces a bounded composition operator on the classical Hardy space  $H^p(\Omega)$ . Moreover we know that being in the class  $H^p_v(\Omega)$  where v is harmonic out of a compact set is equivalent to having a harmonic majorant hence any composition operator on a Hardy class generated by this sort of exhaustion function is also bounded. As a consequence of

Closed Graph Theorem continuity of a composition operator on  $H^p_u(\Omega)$  is in fact determined by whether it takes functions from  $H^p_u(\Omega)$  to  $H^p_u(\Omega)$  or not. However this does **not** always hold when exhaustion function has finite Monge-Ampére mass but not harmonic out of a compact set as we see from the following example :

**Example 1.** Suppose u is the exhaustion function that we constructed in Theorem 1.3 then we know again from the proof of Theorem 1.3 that the function  $\frac{1}{(z-1)^{\frac{3}{4}}} \notin$ 

 $H^1_u(\mathbb{D})$ . Now consider the operator with symbol  $\phi(z) = ze^{i\frac{\pi}{2}}$ , and take the function  $f(z) = \frac{1}{(z-i)^{\frac{2}{4}}}$ , then

$$\|f\|_{H^1_u(\mathbb{D})} = \int_{\partial \mathbb{D}} \frac{1}{(\xi - i)^{\frac{3}{4}}} \beta(\xi) d\sigma(\xi) < \infty$$

since the singularities of  $\beta(\xi)$  and  $f^*(\xi)$  do not overlap and they are both integrable functions on the boundary so  $f(z) \in H^1_u(\mathbb{D})$  but

$$C_{\phi}(f) = f \circ \phi = \frac{1}{e^{i\frac{3\pi}{8}}(z-1)^{\frac{3}{4}}}$$

and  $C_{\phi}(f) \notin H^1_u(\mathbb{D})$ . Therefore not every composition operator is bounded on Poletsky-Stessin Hardy classes even though the symbol function is a nice and simple one like in our example, namely a rotation.

In the next result we will examine the necessary and sufficient conditions for the composition operator  $C_{\varphi}$  to be bounded on this rather interesting space  $H^p_u(\mathbb{D})$ where u is the exhaustion function constructed in the proof of Theorem 1.3 and  $\varphi$ is an automorphism of the unit disc:

**Theorem 5.1.** Let  $\varphi$  be a Mobius transformation such that  $\varphi(z) = e^{i\theta} \frac{z-a}{1-\overline{a}z}$  where  $a \in \mathbb{D}$  and u is the exhaustion function constructed in the proof of Theorem 1.3. Then the following are equivalent:

(i)  $C_{\varphi}$  is a bounded operator on the space  $H^p_u(\mathbb{D})$ 

(ii) There exists a constant K > 0 such that  $\int_E \beta(\varphi^{*-1}(\eta)) d\sigma(\eta) \leq K \int_E \beta(\eta) d\sigma(\eta)$ for all measurable  $E \subset \partial \mathbb{D}$  where  $d\mu_u = \beta d\sigma$ (iii)  $\varphi(1) = 1$ 

*Proof.* (It is sufficient to prove the result for p = 1)  $(i \Leftrightarrow ii)$  Let  $f \in H^p_u(\mathbb{D})$  and  $\varphi$  be a Mobius transformation then

$$\|f \circ \varphi\|_{H^p_u(\mathbb{D})} = \int_{\partial \mathbb{D}} |f^* \circ \varphi^*| \beta(\xi) d\sigma(\xi) = \int_{\partial \mathbb{D}} |f^*(\eta)| \beta(\varphi^{*-1}(\eta))| (\varphi^{*-1})' |d\sigma(\eta)|$$

Suppose  $C_{\varphi}$  is bounded on  $H^p_u(\mathbb{D})$  then  $||f \circ \varphi||_{H^p_u(\mathbb{D})} \leq M||f||_{H^p_u(\mathbb{D})}$  for all  $f \in H^p_u(\mathbb{D})$ . Now since bounded functions are in  $H^p_u(\mathbb{D})$  we have  $f(z) \equiv 1 \in H^p_u(\mathbb{D})$  and we will write the above inequality for  $f(z) \equiv 1$ . Since  $|(\varphi^{*-1})'| < N < \infty$  on  $\partial \mathbb{D}$  we get

$$\int_{\partial \mathbb{D}} \beta(\varphi^{*-1}(\eta)) | (\varphi^{*-1})' | d\sigma(\eta) \le N \int_{\partial \mathbb{D}} \beta(\varphi^{*-1}(\eta)) d\sigma(\eta) \le NM \int_{\partial \mathbb{D}} \beta(\eta) d\sigma(\eta)$$

For the converse direction suppose that there exists a constant K > 0 such that  $\int_{\partial \mathbb{D}} \beta(\varphi^{*-1}(\eta)) d\sigma(\eta) \leq K \int_{\partial \mathbb{D}} \beta(\eta) d\sigma(\eta)$  for all measurable  $E \subset \partial \mathbb{D}$ . Then for any characteristic function  $\chi_E, E \subset \partial \mathbb{D}$  we have

$$\int_{\partial \mathbb{D}} \chi_E \beta(\varphi^{*-1}(\eta)) | (\varphi^{*-1})' | d\sigma(\eta) = \int_E \beta(\varphi^{*-1}(\eta)) d\sigma(\eta) \le K \int_E \beta(\eta) d\sigma(\eta)$$

$$= K \int_{\partial \mathbb{D}} \chi_E \beta(\eta) d\sigma(\eta)$$

Hence by monotone convergence theorem for any positive integrable function g we have

$$\begin{split} &\int_{\partial \mathbb{D}} g(\eta)\beta(\varphi^{*-1}(\eta))|(\varphi^{*-1})'|d\sigma(\eta) \leq K \int_{\partial \mathbb{D}} g(\eta)\beta(\eta)d\sigma(\eta) \\ \text{so} \\ \|f \circ \varphi\|_{H^p_u(\mathbb{D})} &= \int_{\partial \mathbb{D}} |f^*|^p \beta(\varphi^{*-1}(\eta))|(\varphi^{*-1})'|d\sigma(\eta) \leq N \int_{\partial \mathbb{D}} |f^*|^p \beta(\varphi^{*-1}(\eta))d\sigma(\eta) \\ &\leq C \int_{\partial \mathbb{D}} |f^*|^p \beta(\eta)d\sigma(\eta) = C \|f\|_{H^p_u(\mathbb{D})} \end{split}$$

Hence  $C_{\varphi}$  is bounded.

 $(i \Leftrightarrow iii)$  Suppose  $C_{\varphi}$  is bounded and  $\varphi(1) \neq 1$  then  $\exists \xi \in \partial \mathbb{D}$ ,  $\xi \neq 1$  such that  $\varphi(\xi) = 1$  and take the function  $f(z) = \frac{1}{(1-z)^{\frac{3}{4}}}$  then we know that  $f(z) \notin H^1_u(\mathbb{D})$ . Now consider the function  $F(z) = f \circ \varphi^{-1}(z)$  then

$$\|F(z)\|_{H^{1}_{u}(\mathbb{D})} = \int_{\partial \mathbb{D}} \frac{1}{|1 - \varphi^{-1}(\eta)|^{\frac{3}{4}}} d\mu_{u}(\eta)$$
$$= \int_{\partial \mathbb{D} \setminus B_{\gamma}(1)} \frac{1}{|1 - \varphi^{-1}(\eta)|^{\frac{3}{4}}} d\mu_{u}(\eta) + \int_{B_{\gamma}(1)} \frac{1}{|1 - \varphi^{-1}(\eta)|^{\frac{3}{4}}} d\mu_{u}(\eta) < \infty$$

for some  $\gamma > 0$ . The first integral in the last line is bounded because on  $\partial \mathbb{D} \setminus B_{\gamma}(1)$ ,  $d\mu_u$  and  $d\sigma$  are mutually absolutely continuous and  $\frac{1}{|1-\varphi^{-1}(\eta)|^{\frac{3}{4}}}$  is  $d\sigma$  integrable and the second integral is bounded because on  $B_{\gamma}(1)$ ,  $\frac{1}{|1-\varphi^{-1}(\eta)|^{\frac{3}{4}}}$  is a bounded function and hence it is  $d\mu_u$  integrable.

Hence  $F(z) \in H^1_u(\mathbb{D})$  but  $F \circ \varphi = f \notin H^1_u(\mathbb{D})$  which contradicts with the boundedness of  $C_{\varphi}$ .

Suppose now  $\varphi(1) = 1$  from the  $(i \Leftrightarrow ii)$  part of the proof we know that if  $\frac{\beta(\varphi^{-1}(\eta))}{\beta(\eta)} < M < \infty$  then  $C_{\varphi}$  is bounded and for the case  $\varphi(1) = 1$  we have  $\frac{\beta(\varphi^{-1}(\eta))}{\beta(\eta)}$  bounded hence the result follows.

We can generalize this result to a slightly wider class of symbols as follows:

**Theorem 5.2.** Let  $\varphi : \overline{\mathbb{D}} \to \overline{\mathbb{D}}$  be a locally univalent self map of  $\mathbb{D}$  such that  $\varphi$ is differentiable in a neighborhood of  $\overline{\mathbb{D}}$ . Then  $C_{\varphi}$  is bounded on  $H^1_u(\mathbb{D})$  if and only if  $\varphi(1) = 1$  and  $N^{\varphi}_{\beta}(\eta) \leq K\beta(\eta)$  for some K > 0 and all  $\eta \in \partial \mathbb{D}$  where  $N^{\varphi}_{\beta}(\eta) = \sum_{j>1} \beta(\xi_j(\eta))$  and  $\{\xi_j(\eta)\}$  are the zeros of  $\varphi(z) - \eta$ .

*Proof.* ( $\Rightarrow$ ) Suppose  $C_{\varphi}$  is bounded and  $\varphi(1) \neq 1$  then there exists a  $\xi \in \partial \mathbb{D}$  such that  $\varphi(1) = \xi$  now consider the function  $f(z) = \frac{1}{(\xi-z)^{\frac{3}{4}}}, f(z) \in H^1_u(\mathbb{D})$  and

$$\|f \circ \varphi\|_{H^1_u(\mathbb{D})} = \int_{\partial \mathbb{D}} |f^* \circ \varphi| d\mu_u = \int_{\partial \mathbb{D} \setminus \overline{B_{\gamma}(1)}} |f^* \circ \varphi| d\mu_u + \int_{\overline{B_{\gamma}(1)}} |f^* \circ \varphi| d\mu_u$$

where  $\overline{B_{\gamma}(1)} = \partial \mathbb{D} \cap B_{\gamma}(1)$  for some small  $\gamma > 0$ . The first integral in the sum is bounded since over  $\partial \mathbb{D} \setminus \overline{B_{\gamma}(1)}$ ,  $d\mu_u = Cd\sigma$  for some C > 0 and  $f \in H^1_u(\mathbb{D}) \subset H^1(\mathbb{D})$ so boundedness over this region is guaranteed by classical  $H^p$  theory but

$$\int_{\overline{B_{\gamma}(1)}} |f^* \circ \varphi| d\mu_u = \int_{\overline{B_{\gamma}(1)}} \frac{1}{|\xi - \varphi|^{\frac{3}{4}}} d\mu_u$$

and  $\varphi$  has finite derivative near  $\{1\}$  so

$$\int_{\overline{B_{\gamma}(1)}} \frac{1}{|\xi - \varphi|^{\frac{3}{4}}} d\mu_u \ge M \int_{\overline{B_{\gamma}(1)}} \frac{1}{|1 - \eta|^{\frac{3}{4}}} d\mu_u \to \infty$$

contradicting  $C_{\varphi}$  being bounded. Hence  $\varphi(1) = 1$ .

The inequality  $N_{\beta}^{\varphi}(\eta) \leq K\beta(\eta)$  is trivially true for  $\eta = 1$  so we will consider the case where  $\eta \neq 1$  and assume for a contradiction that  $N_{\beta}^{\varphi}(\eta_0) > K\beta(\eta_0)$ for all K, for some  $\eta_0 \neq 1$ . Then from the definition of  $N_{\beta}^{\varphi}(\eta_0)$  we see that  $\beta(\varphi(\eta_0)) \to \infty$  which gives  $\varphi(\eta_0) = 1$ . Then consider the function  $f(z) = \frac{1}{(\eta_0 - z)^{\frac{3}{4}}}$ then by the same argument above  $f \circ \varphi \notin H_u^1(\mathbb{D})$  contradicting  $C_{\varphi}$  being bounded hence  $N_{\beta}^{\varphi}(\eta) \leq K\beta(\eta)$  for all  $\eta \in \partial \mathbb{D}$  for some K > 0.

( $\Leftarrow$ ) Since  $\varphi$  is locally univalent we can find a countable collection of disjoint open arcs  $\Omega_j$  with  $\sigma(\partial \mathbb{D} \setminus \bigcup \Omega_j) = 0$  and the restriction of  $\varphi$  to each  $\Omega_j$  is univalent. Write  $\psi_j(w)$  for the inverse of  $\varphi$  taking  $\varphi(\Omega_j)$  onto  $\Omega_j$ . Then change of variables formula gives

$$\int_{\Omega_j} |f^* \circ \varphi| |\varphi'(\xi)| \beta(\xi) d\sigma(\xi) = \int_{\varphi(\Omega_j)} |f^*(w)| \beta(\psi_j(w)) d\sigma(w)$$

where  $\xi = \psi_j(w)$ . Now denoting the characteristic function of  $\varphi(\Omega_j)$  by  $\chi_j$  we get

$$\begin{split} \int_{\partial \mathbb{D}} |f^* \circ \varphi| |\varphi'(\xi)| \beta(\xi) d\sigma(\xi) &= \int_{\varphi(\partial \mathbb{D})} |f^*(w)| \left( \sum_{j \ge 1} \chi_j(w) \beta(\psi_j(w)) \right) d\sigma(w) \\ &= \int_{\varphi(\partial \mathbb{D})} |f^*(w)| \left( \sum_{j \ge 1} \beta(\xi_j(w)) \right) d\sigma(w) \\ \int_{\mathcal{O}} |f^* \circ \varphi| d\mu_u \le M \int_{\mathcal{O}} |f^* \circ \varphi| |\varphi'(\xi)| \beta(\xi) d\sigma(\xi) \end{split}$$

 $\mathbf{SO}$ 

$$\int_{\partial \mathbb{D}} |f^* \circ \varphi| d\mu_u \leq M \int_{\partial \mathbb{D}} |f^* \circ \varphi| |\varphi'(\xi)| \beta(\xi) d\sigma(\xi)$$
$$= M \int_{\varphi(\partial \mathbb{D})} |f^*(w)| N_{\beta}^{\varphi}(w) d\sigma(w) \leq KM \int_{\partial \mathbb{D}} |f^*(w)| d\sigma(w)$$

hence  $C_{\varphi}$  is a bounded operator.

### References

- Muhammed Ali Alan, Hardy Spaces on Hyperconvex Domains, Ankara: Middle East Technical University, 2003.
- [2] Jean-Pierre Demailly, Mesures de Monge-Ampére et Caractérisation Géométrique des Variétés Algébraiques Affines, Mémoire de la Société Mathématique de France 19(1985),1-124.
- [3] Jean-Pierre Demailly, Mesures de Monge-Ampére et Mesures Pluriharmoniques, Matematische Zeitschrift (1987), no:194,519-564.
- [4] Evgeny A. Poletsky, Michael I. Stessin, Hardy and Bergman Spaces on Hyperconvex Domains and Their Composition Operators Indiana Univ. Math. J. 57 (2008), 2153-2201.
- [5] Evgeny A. Poletsky, Weak and Strong Limit Values arXiv:1105.1365v1, (2011).
- [6] Thomas Ransford, Potential Theory in the Complex Plane, Cambridge University Press, London Mathematical Society Student Texts 28,1995.
- [7] Peter L. Duren, Theory of H<sup>p</sup> Spaces, Academic Press, Inc, 1970.

- [8] N. Kerzman, J.P. Rosay; Fonctions plurisousharmoniques d'exhaustion bornées et domaines taut, Math. Ann., 257, 1981, 171-184.
- [9] Walter Rudin, Function Theory in Polydiscs, W.A. Benjamin Inc, 1969.
- [10] Walter Rudin, Function Theory in the Unit Ball of  $\mathbb{C}^n$ , Springer-Verlag, 1980.
- [11] A. Sadullaev, Plurisubharmonic measures and capacities on complex manifolds, Russian Mathematical Surveys, 36:4, 1981, 61-119.
- [12] Maciej Klimek, Pluripotential Theory, Clarendon Press, 1991.
- [13] Elias M. Stein, The Boundary Behaviour of Holomorphic Functions of Several Complex Variables, Princeton University Press, Princeton, 1972.
- [14] Theodore W. Gamelin, Uniform Algebras, Prentice Hall, 1969.
- [15] Carl C. Cowen, Barbara D. Maccluer, Composition Operators on Spaces of Analytic Functions, CRC Press, 1995.
- [16] Alexandru Aleman, Nathan S. Feldman, William T. Ross, The Hardy Space of a Slit Domain, Frontiers in Mathematics, Birkhäuser Basel, 2009.
- [17] Joel H. Shapiro, Wayne Smith, Hardy Spaces That Support No Compact Composition Operators, Journal of Functional Analysis, 2003.
- [18] Sibel Şahin, Hardy Spaces on Bounded Hyperconvex Domains and Monge-Ampére measures, PhD Dissertation, Sabanci University.

Faculty of Engineering and Natural Sciences, Sabanci University  $E\text{-}mail\ address:\ \texttt{sahinsibel@sabanciuniv.edu}$