

On the gravitational self-energy of a spherical shell

G. Dillon¹

*Dipartimento di Fisica, Università di Genova
INFN, Sezione di Genova*

Abstract: According to Einstein’s mass-energy equivalence, a body with a given mass extending in a large region of space, will get a smaller mass when confined into a smaller region, because of its own gravitational energy. The classical self-energy problem has been studied in the past in connection with the renormalization of a charged point particle. Still exact consistent solutions have not been thoroughly discussed in the simpler framework of Newtonian gravity. Here we exploit a spherical symmetrical shell model and find two possible solutions, depending on some additional assumption. The first solution goes back to Arnowitt, Deser and Misner (1960). The second is new and yields a new vanishingly small value ($10^{-55}cm$) for the classical electron radius.

1 Introduction

In classical Newtonian physics, the gravitational energy of a simple spherically symmetric shell of mass M_0 and radius R is

$$U_0(R) = -G \frac{M_0^2}{2R} \quad (1)$$

where G is the gravitational constant.

Taking into account mass-energy equivalence from special relativity, this binding energy (negative) is equivalent to a mass defect. Furthermore the equality of inertial and gravitational masses has been tested experimentally even in presence of sizeable mass defects due to large binding energies [1, 2]. Hence the mass of the shell will be different from M_0 and, in turn, (1) should be reexpressed in terms of the new “renormalized” mass. In the following we shall refer to M_0 as the “bare” mass while the renormalized $M(R)$ is the resulting mass when M_0 is uniformly distributed on a spherical shell of radius R . $M(R)$ takes into account the gravitational self-energy, while M_0 corresponds to the sum of all the masses that one would obtain tearing the sphere in many small pieces and moving them away apart².

¹e-mail: dillon@ge.infn.it

²Here we are disregarding any form of kinetic energy or internal pressure or stress that can also contribute to the mass.

The renormalized mass at the first order in c^{-2} is immediately obtained from (1)

$$M(R) \approx M_0 + U_0(R)/c^{-2} = M_0(1 - G \frac{M_0}{2Rc^2}) \quad (2)$$

However the exact calculation of $M(R)$ from a given bare mass M_0 is not trivial and rests upon some additional assumption.

The classical self-energy problem has been studied in the past by R. Arnowitt, S. Deser and C.W. Misner (ADM) [3, 4, 5] in the framework of a canonical formulation of General Relativity. Here we wish to consider the problem in the simpler context of Newtonian physics with minimal assumptions. In this sense the present paper has some pedagogical character. In Sec.2 we revisit the solution given by ADM. We find however that this solution is inconsistent with the following expression

$$\delta U(R) = -G \frac{M(R)\delta m_0}{R} \quad (3)$$

for the gravitational interaction energy between the shell and a test-particle δm_0 settled on its surface. In Sec.3 we relate the reason for the inconsistency to the lack of an appropriate mass renormalization of the test-particle due to $\delta U(R)$ itself. In Sec.4 we propose an alternative solution, consistent with (3), which allows also for a new definition of the classical electron radius, which adds to the well known one.

2 A consistent solution for the spherical shell and an inconsistency

Given the expression (1) for the gravitational energy of a spherical shell, it seems quite natural to write down the following consistent equation for the renormalized mass $M(R)$

$$M(R) = M_0 - \frac{G}{2} \frac{M(R)^2}{Rc^2} \quad (4)$$

or equivalently for the gravitational self-energy of the shell

$$U(R) = -G \frac{(M_0 + U(R)/c^2)^2}{2R} \quad (5)$$

Defining

$$R_0 = \frac{GM_0}{c^2} \quad (6)$$

one gets the (positive) solution for the mass

$$M(R) = M_0(-1 + \sqrt{1 + 2R_0/R})R/R_0 \quad (7)$$

with the corresponding gravitational self-energy

$$U(R) = -G \frac{M(R)^2}{2R} = -M_0c^2(1 + R_0/R - \sqrt{1 + 2R_0/R})R/R_0 \quad (8)$$

The equation (4) has been considered [5, 6] in the framework of the classical theory of the electron. In fact, adding to (4) the contribution to the mass of the electromagnetic energy $e^2/2R$ (this time positive), the ensuing solution tends to a finite value when $R \rightarrow 0$:

$$M(R \rightarrow 0) = |e|/\sqrt{G} \equiv m_G \quad (9)$$

independent of M_0 . This elegant result exhibits a nice feature of the gravitational self-energy as a natural cutoff for the Coulomb self-energy of a point charge. The result is however numerically too big ($m_G \approx 10^{21}m_e$) compared to the electron mass. We shall further comment on this.

Now one could naively think that the (now renormalized) shell yields a gravitational field identical to that of a spherical shell with the trivial substitution $M_0 \rightarrow M(R)$, i.e. a gravitational potential function (for $r \geq R$)

$$\Phi(r) = -G \frac{M(R)}{r} \quad (10)$$

Hence the interaction energy with a test particle settled on its surface should be given by (3). If so, the total mass of the system (spherical shell of renormalized mass $M(R)$ plus test-particle δm_0 on its surface) is

$$M_t(R) = M(R) + \delta m_0 + \delta U(R)/c^2 = M(R) + \delta m_0 \left(1 - G \frac{M(R)}{c^2 R}\right) \quad (11)$$

Substituting $M(R)$ from (7) one has

$$M_t(R) = M(R) + \delta m_0 \left(2 - \sqrt{1 + 2R_0/R}\right) \quad (12)$$

Here we come to a contradiction. Indeed, suppose we want to deposit a test particle δm_0 on the surface of $M(R)$ and let us think about this test mass as being uniformly distributed on a thin spherical shell of radius r centered on the origin, just as $M(R)$. (Note that, neglecting higher orders in δm_0 , we do not worry about self-energy of δm_0 on its own. In other words: $\delta m(r) \approx \delta m_0$.) Now imagine to bring r to R and to stick δm_0 as a thin film on $M(R)$. Then, viewing the system as a new shell of bare mass $M_0 + \delta m_0$, one has again from (7)

$$M_t(R) = M(R) + \frac{\partial M(R)}{\partial M_0} \delta m_0 = M(R) + \frac{\delta m_0}{\sqrt{1 + 2R_0/R}} \quad (13)$$

that does not agree with (12), but at the first order in R_0/R (i.e. in c^{-2}). So there is a mistake somewhere.

3 Renormalizing the test-mass

In Sec.2 we thought about the test-particle as being spherically distributed on a thin shell concentric with the shell $M(R)$. We argued that we should not worry

about its own self-energy (since we work at first order in δm_0); however we left aside the possibility of further renormalization consequences on the masses due to the interaction energy $\delta U(R)$ itself. Let us assume, by now, that the whole $\delta U(R)$ be attributed to the test-mass, i.e.

$$\delta m_0 \rightarrow \delta m = \delta m_0 + \delta U(R)/c^2 \quad (14)$$

This assumption amounts rewriting (3) as a self-consistent equation for $\delta U(R)$

$$\delta U(R) = -G \frac{M(R)(\delta m_0 + \delta U(R)/c^2)}{R} \quad (15)$$

From (15)

$$\delta U(R) = -G \frac{M(R)\delta m_0}{R(1 + G \frac{M(R)}{Rc^2})} \quad (16)$$

and

$$M_t(R) = M(R) + \frac{\delta m_0}{1 + G \frac{M(R)}{Rc^2}} \quad (17)$$

at variance with (11). Substituting (7) we get

$$1 + G \frac{M(R)}{Rc^2} = \sqrt{1 + 2R_0/R} \quad (18)$$

so (17) now agrees with (13) and there is no inconsistency.

In fact the starting equation (4) may be obtained from (17) as follows: Viewing the total system as a new shell with increased bare mass ($M_0 + \delta m_0$), one can write

$$M_t(R) - M(R) \equiv dM(R) \quad ; \quad \delta m_0 \equiv dM_0$$

Hence (17) is a differential equation that yields the mass $M(R)$ of a spherical shell of radius R as a function of its bare mass M_0 :

$$\frac{dM(R)}{dM_0} = \frac{1}{1 + G \frac{M(R)}{Rc^2}} \quad (19)$$

whose solution is just (4).

Note that the renormalization of δm_0 may be equivalently described in terms of a suitable modification of the gravitational potential, that, instead of (10), should be written as (for $r > R$)

$$\Phi(r) \rightarrow \tilde{\Phi}(r) = -\frac{G}{(1 + G \frac{M(R)}{rc^2})} \frac{M(R)}{r} \quad (20)$$

with $M(R)$ given by (7).

4 An alternative solution and a new classical electron radius

In Sec.3 it was proved that renormalizing the mass of the test-particle by the interaction energy with the massive shell leads to (7,8). However this assumption may appear somewhat arbitrary. Moreover it seems not conciliable with additivity. In fact let us think about the shell $M(R)$ as made up by the sum of a large number N of light overlapping shells each of mass $\delta m'$

$$\sum_i \delta m'_i = N\delta m' \equiv M(R)$$

and rewrite (3) as the sum of the interaction energies with each sub-shells

$$\delta U(R) = -G \sum_i \frac{\delta m'_i \delta m_0}{R} \equiv \sum_i \delta u_i(R) = N\delta u(R) \quad (21)$$

Now for each individual term in the sum in (21) let us perform the renormalization of the test mass as before ($\delta m_0 \rightarrow \delta m_0 + \delta u(R)/c^2$). Then

$$\delta U(R) = -G \sum_i \frac{\delta m'_i (\delta m_0 + \delta u(R)/c^2)}{R} \approx -G \frac{M(R) \delta m_0}{R} \quad (22)$$

for large N . Moreover the final result (22) will not change even if the little mass equivalent to the interaction energy were attributed (fully or in part) to the shell $M(R)$ rather than to the test-particle. This is because the ensuing correction to $\delta U(R)$ comes to be at a higher order in δm_0 in this case.

Then, assuming additivity, one should not change (3) and look for a solution other than (7,8). In fact we may exploit the method outlined in Sec.3. Starting from (11) we get the differential equation

$$\frac{dM(R)}{dM_0} = 1 - G \frac{M(R)}{Rc^2} \quad (23)$$

whose solution is

$$\ln \left(1 - G \frac{M(R)}{Rc^2} \right) = -G \frac{M_0}{Rc^2} \quad (24)$$

$$M(R) = \frac{Rc^2}{G} \left(1 - \exp \left[-G \frac{M_0}{Rc^2} \right] \right) \equiv M_0 \left(1 - \exp \left[-\frac{R_0}{R} \right] \right) \frac{R}{R_0} \quad (25)$$

Instead of (12) we have now

$$M_t(R) = M(R) + \delta m_0 \exp \left[-\frac{R_0}{R} \right] \quad (26)$$

and there is, of course, no contradiction.

The alternative solution (25) deserves a couple of comments in connection with the classical theory of the electron [7]. To obtain the mass of a spherical shell of

radius R , charge e and bare mass $M_0 = 0$, it is enough to make the substitution $M_0 \rightarrow e^2/2Rc^2$ in (25). One gets

$$M_e(R) = \frac{c^2 R}{G} \left(1 - \exp\left[-G \frac{e^2}{2R^2 c^4}\right]\right) \quad (27)$$

Again we may appreciate the regularizing role of the gravitational self-energy that heals the divergency of the Coulomb self-energy when $R \rightarrow 0$. Instead of the previously quoted result $M_e(R \rightarrow 0) = m_G = |e|/\sqrt{G}$ [5, 6], this time we obtain $M_e(R \rightarrow 0) = 0$.

Let us now define

$$R_e \equiv Gm_e/c^2 \quad (28)$$

and rewrite (27) in terms of R_e and $r_e = e^2/m_e c^2$ (the usual expression for the classical electron radius [7]). We get

$$M_e(R) = m_e \frac{R}{R_e} \left(1 - \exp\left[-\frac{r_e R_e}{2R^2}\right]\right) \quad (29)$$

In order that $M_e(R) \equiv m_e$, one has to find solutions of

$$\frac{R}{R_e} \left(1 - \exp\left[-\frac{r_e R_e}{2R^2}\right]\right) = 1 \quad (30)$$

Numerically for the electron it is: $r_e = 2.82 \cdot 10^{-13} cm$ and $R_e = 0.7 \cdot 10^{-55} cm$. Then it is immediately seen that (30) has two distinct solutions (R_1, R_2), almost exactly coincident with $r_e/2$ and R_e . Note that, while for the first solution ($R_1 \approx r_e/2$) gravitational self-energy effects are totally negligible, the opposite happens for $R_2 \approx R_e$ (see (28)).

5 Concluding remarks

In this paper, taking due account of mass-energy equivalence (and of the equality of the inertial and gravitational masses), we faced the seemingly trivial problem of calculating the gravitational self-energy of a given distribution of mass. We exploited the simple model of a spherically symmetrical distribution over a shell of radius R . The problem may equivalently be stated as how to calculate the renormalized mass of the given distribution $M(R)$ from its bare mass M_0 . We displayed two possible solutions. The first one was known since 1960 [5, 6], the second is new. The difference between the two is related to specific assumptions about mass renormalization. The first one requires a modification (see (20)) of the gravitational potential generated by the shell with respect to the well known expression of Newtonian theory (10). The second solution does not need any modification of (10) and follows from a simple assumption of additivity (see Sec.4).

When the bare mass M_0 is substituted with $e^2/2Rc^2$ one gets the gravitationally renormalized e.m. mass of a pure static electric charge uniformly distributed on a spherical shell of radius R . Defining

$$R_G = |e|\sqrt{G}/c^2 \equiv Gm_G/c^2 \quad (31)$$

it is for the solution (7)

$$M_e(R) = m_G \frac{R}{R_G} \left(-1 + \sqrt{1 + (R_G/R)^2} \right) \quad (32)$$

which reaches its maximum value $m_G = |e|/\sqrt{G}$ at $R = 0$. On the other hand from the alternative solution (27) one gets

$$M_e(R) = m_G \frac{R}{R_G} \left(1 - \exp\left[-\frac{R_G^2}{2R^2}\right] \right) \quad (33)$$

that has a maximum of the order of m_G at $R \approx R_G$. Since (33) goes to zero when $R \rightarrow 0$, one finds a further solution for the electron mass at a vanishingly small radius ($10^{-55} cm$).

Indeed in [6] it was stated that a system cannot have pointlike dimensions at the mass m_G and the minimum possible dimensions for that value are just given by (31). It is remarkable that m_G may also be written as $m_G = \sqrt{\alpha} M_{Planck}$; likewise $R_G = \sqrt{\alpha} l_{Planck}$. As emphasized in [8], these quantities are purely classical (the square root of the fine-structure constant α cancels \hbar) and define the scale at which gravitational and electromagnetic self-energies become comparable.

References

- [1] L. Sauthern: Proc. Roy. Soc. S.A **84**, (1910) 325
- [2] A. L. Zeeman: Proc. Sect. Sci. K. ned. Akad. Wet. **20**, (1917) 542
- [3] R. Arnowitt, S. Deser and C.W. Misner: Phys. Rev. Lett. **4**, (1960) 375
- [4] R. Arnowitt, S. Deser and C.W. Misner: Phys. Rev. **118**, (1960) 1100
- [5] R. Arnowitt, S. Deser and C.W. Misner: Phys. Rev. **120**, (1960) 313
- [6] M.A. Markov: Sov. Phys. JETP **37**, (1972) 561
- [7] See for instance: J.D. Jackson: *Classical electrodynamics* (Wiley, New York 1975) ch. 17
- [8] M. Visser, Phys. Lett. A **139** (1989) 99