AN ASYMPTOTIC ESTIMATE OF THE VARIANCE OF THE SELF-INTERSECTIONS OF A PLANAR PERIODIC LORENTZ PROCESS

FRANÇOISE PÈNE

ABSTRACT. We consider a \mathbb{Z}^2 -periodic planar Lorentz process with strictly convex obstacles and finite horizon. This process describes the displacement of a particle moving in the plane with unit speed and with elastic reflection on the obstacles. We call number of self-intersections of this Lorentz process the number V_n of couples of integers (k, ℓ) smaller than n such that the particle hits a same obstacle both at the kth and at the ℓ th collision times. The aim of this article is to prove that the variance of V_n is equivalent to n^2 (such a result has recently been proved for simple planar random walks in [6]).

1. INTRODUCTION

We consider a finite number of convex open sets $O_1, ..., O_I \subset \mathbb{R}^2$ with boundary C^3 -smooth and with non null curvature. We repeat these sets \mathbb{Z}^2 -periodically by defining $U_{i,\ell} = O_i + \ell$ for every $(i, \ell) \in \{1, ..., I\} \times \mathbb{Z}^2$. We suppose that the closures of the $U_{i,\ell}$ are pairwise disjoint. We assume that the horizon is finite, which means that every line meets the boundary of at least one obstacle (i.e. there is no infinite free flight). We consider a particle moving in the domain $Q := \mathbb{R}^2 \setminus \bigcup_{i=1}^{I} \bigcup_{\ell \in \mathbb{Z}^2} U_{i,\ell}$ with unit speed and with respect to the Descartes reflection law at its reflection times (reflected angle=incident angle). We assume that the particle starts from $[0,1]^2 \cap Q$ with uniform distribution in position and in speed. The Lorentz process describes the evolution of the particle in Q. Because of the \mathbb{Z}^2 -periodicity, it is strongly related to the Sinai billiard, the ergodic properties of which have been studied namely by Sinai in [9] (for its ergodicity), Bunimovich and Sinai [2, 3], Bunimovich, Chernov and Sinai [4, 5] (for central limit theorems), Young [11] (for exponential rate of decorrelation). The similarity of behaviour of the Lorentz process with a simple planar random walk has been investigated by many authors ([10, 7],...). The number of auto-intersections up to time n of a random walk $(S_n)_n$ is $\tilde{V}_n := \sum_{k,\ell=1}^n \mathbf{1}_{\tilde{S}_k = \tilde{S}_\ell}$. This quantity is linked with random walks in random sceneries [1, 6]. Recently, in [6], Deligiannidis and Utev proved that $Var(\tilde{V}_n) \sim \tilde{c}n^2$ with an explicit \tilde{c} . This improved the estimation in $O(n^2 \log n)$ by Bolthausen [1]. For the Lorentz process, we define (\mathcal{I}_k, S_k) in $\{1, ..., I\} \times \mathbb{Z}^2$ for the index of the obstacle hit at the k-th reflection time $((I_0, S_0)$ being the index of the obstacle at the reflection time just before time 0). Recall that $(k^{-1/2}S_k)_{k\geq 1}$ admits an asymptotic positive variance matrix Σ^2 . We call number of self-intersections of the Lorentz process up to the *n*-th reflection time the quantity $V_n := \sum_{k,\ell=1}^n \mathbf{1}_{S_k = S_\ell, \mathcal{I}_k = \mathcal{I}_\ell}$. In [8], we proved that $\mathbb{E}[V_n] \sim c_0 n \log n$ with $c_0 := \frac{\sum_{i=1}^{I} (|\partial O_i|^2)}{(\sum_{i=1}^{I} |\partial O_i|)^2 \pi \sqrt{\det \Sigma^2}}$, where $|\partial O_i|$ stands for the length of ∂O_i . In [8], $Var(V_n) = O(n^2 \log n)$ was enough for our study of the planar Lorentz

Date: November 13, 2017.

²⁰¹⁰ Mathematics Subject Classification. 60F99,37D50.

Key words and phrases. Sinai billiard, Lorentz process, self intersection

Françoise Pène is supported by the french ANR projects GEODE (ANR-10-JCJC-0108) and PERTURBATIONS (ANR-10-BLAN-0106).

process in random scenery. Our proof of the following result uses decorrelation and precised local limit theorems established in [8]. It provides an alternative strategy to the one of [6].

Theorem 1. $Var(V_n) \sim cn^2$ with

$$c := c_0^2 \left(1 + 2J - \frac{\pi^2}{6} \right) \text{ and } J := \int_{[0,1]^3} \frac{(1 - (u + v + w))\mathbf{1}_{\{u + v + w \le 1\}} \, du \, dv \, dw}{uv + uw + vw}$$

2. Proof of Theorem 1

Observe that the distribution of $(S_k - S_0, \mathcal{I}_k)_k$ under \mathbb{P} and under $\bar{\nu}$ considered in [8] are the same (by \mathbb{Z}^2 -periodicity and by construction of $\bar{\nu}$). We write $E_{k,\ell} := \{S_k = S_\ell, \mathcal{I}_k = \mathcal{I}_\ell\}$. According to [8], we have

$$\mathbb{P}(E_{k,\ell}) = \mathbb{P}(E_{0,|\ell-k|}) = c_1|\ell-k|^{-1} + O(|\ell-k|^{-2}), \text{ with } c_1 := \frac{\sum_i^I (\mathbb{P}(\mathcal{I}_0 = i))^2}{2\pi\sqrt{\det \Sigma^2}} = \frac{c_0}{2}.$$
 (1)

Observe that we have $V_n = n + 2 \sum_{1 \le k \le \ell \le n} \mathbf{1}_{S_k = S_\ell, \mathcal{I}_k = \mathcal{I}_\ell}$ and so

$$Var(V_n) = 4 \sum_{1 \le k_1 < \ell_1 \le n} \sum_{1 \le k_2 < \ell_2 \le n} D_{k_1, \ell_1, k_2, \ell_2} = 8A_1 + 8A_2 + 8A_3 + 4A_4,$$

with $D_{k_1,\ell_1,k_2,\ell_2} := \mathbb{P}(E_{k_1,\ell_1} \cap E_{k_2,\ell_2}) - \mathbb{P}(E_{k_1,\ell_1})\mathbb{P}(E_{k_2,\ell_2})$ and

$$A_1 := \sum_{1 \le k_1 < \ell_1 \le k_2 < \ell_2 \le n} D_{k_1, \ell_1, k_2, \ell_2}, \ A_2 := \sum_{1 \le k_1 \le k_2 < \ell_1 \le \ell_2 \le n} D_{k_1, \ell_1, k_2, \ell_2}$$

$$A_3 := \sum_{1 \le k_1 < k_2 < \ell_2 < \ell_1 \le n} D_{k_1, \ell_1, k_2, \ell_2}, \ A_4 := \sum_{1 \le k < \ell \le n} [\mathbb{P}(E_{k, \ell}) - (\mathbb{P}(E_{k, \ell}))^2].$$

• Control of A_1 . Due to [8], if $k_1 < \ell_1 \le k_2 < \ell_2$, then $|D_{k_1,\ell_1,k_2,\ell_2}| \le C_1 \tau_1^{k_2-\ell_1}/((\ell_1-k_1)(\ell_2-k_2))$ for some $C_1 > 0$ and some $\tau_1 \in (0,1)$. Hence $A_1 = O(n \log^2 n) = o(n^2)$.

- Control of A_4 . Due to (1) or [10], $A_4 \le C_2 \sum_{1 \le k \le \ell \le n} (\ell - k)^{-1} = O(n \log n) = o(n^2).$ • Control of A_2 .
 - According to [8], we have

$$A_{2} = \sum_{1 \le k_{1} < k_{2} \le \ell_{1} < \ell_{2} \le n} c_{1}^{2} \left[\left(\sum_{x} \frac{e^{-\frac{\langle (\Sigma^{2})^{-1}x, x \rangle}{2} \left(\frac{1}{k_{2} - k_{1}} + \frac{1}{\ell_{1} - k_{2}} + \frac{1}{\ell_{2} - \ell_{1}}\right)}}{2\pi \sqrt{\det \Sigma^{2}} (k_{2} - k_{1})(\ell_{1} - k_{2})(\ell_{2} - \ell_{1})} \right) - \frac{1}{(\ell_{1} - k_{1})(\ell_{2} - k_{2})} \right] + o(n^{2}),$$

$$(2)$$

where $\sum_{x \in \mathbb{Z}^2 : |x| \leq ||S_1||_{\infty} \min(k_2 - k_1, \ell_1 - k_2, \ell_2 - \ell_1)}$ and $\langle \cdot, \cdot \rangle$ is the usual scalar product in \mathbb{R}^2 . $(\Sigma^2)^{-1}x.x$ (1) (1) (1)

$$- \text{ First } A_{2,0} := \sum_{1 \le k_1 < k_2 < \ell_1 < \ell_2 \le n} \sum_x \frac{e^{-\frac{(\Sigma^2)^{-1} \cdot x, x)}{2} \left(\frac{1}{k_2 - k_1} + \frac{1}{\ell_1 - k_2} + \frac{1}{\ell_2 - \ell_1}\right)}}{2\pi \sqrt{\det \Sigma^2} (k_2 - k_1) (\ell_1 - k_2) (\ell_2 - \ell_1)}$$
$$= \sum_{(k_1, m_0, m_1, m_2) \in E_n} \sum_{|x| \le ||S_1||_{\infty} \min(m_0, m_1, m_2)} \frac{e^{-\frac{((\Sigma^2)^{-1} \cdot x, x)}{2} \left(\frac{1}{m_0} + \frac{1}{m_1} + \frac{1}{m_2}\right)}}{2\pi \sqrt{\det \Sigma^2} m_0 m_1 m_2},$$

with $E_n := \{(k_1, m_0, m_1, m_2) \in \mathbb{Z}_+ : k_1 + m_0 + m_1 + m_2 \le n\}$. Observe that, using a comparison series-integral, we obtain

$$\sup_{||S_1||_{\infty} \le a \le 3} \left| |S_1||_{\infty} \left| \sum_{x \in \mathbb{Z}^2 : |x| \le am} e^{-\frac{\langle (\Sigma^2)^{-1} x, x \rangle}{2m}} - 2\pi m \sqrt{\det \Sigma^2} \right| = O(\sqrt{m}).$$
(3)

So
$$A_{2,0} = \sum_{(k_1,m_0,m_1,m_2)\in E_n} \frac{1+O(\min(m_0,m_1,m_2)^{-1/2})}{m_0m_1+m_0m_2+m_1m_2} \sim n^2 J.$$

- Second $\sum_{1 \le k_1 < k_2 < \ell_1 < \ell_2 \le n} \frac{1}{(\ell_1 - k_1)(\ell_2 - k_2)} = A_{2,1} + 2A_{2,2},$

with
$$A_{2,1} := \sum_{k=1}^{n} \sum_{\max(1,2k-n) \le m \le k} \frac{n - (2k - m) + 1}{k^2} \le \sum_{k=1}^{n} \sum_{m=0}^{k} \frac{n}{k^2} = O(n \log n) = o(n^2),$$

and
$$A_{2,2} := \sum_{1 \le k < \ell \le n} \sum_{\max(0,k+\ell-n) \le m \le k} \frac{n - (k+\ell-m) + 1}{k\ell}$$

$$= \sum_{\ell=1}^{\lfloor n/2 \rfloor} \sum_{k=1}^{\ell-1} \sum_{m=0}^{k} \dots + \sum_{\ell=\lfloor n/2 \rfloor+1}^{n} \sum_{k=1}^{n-\ell} \sum_{m=0}^{k} \dots + \sum_{\ell=\lfloor n/2 \rfloor+1}^{n} \sum_{k=n-\ell+1}^{\ell-1} \sum_{m=k+\ell-n}^{k} \dots$$

$$= o(n^2) + \sum_{\ell=1}^{\lfloor n/2 \rfloor} \sum_{k=1}^{\ell-1} \frac{2(n-\ell)-k}{2\ell} + \sum_{\ell=\lfloor n/2 \rfloor+1}^{n} \sum_{k=1}^{n-\ell} \frac{2(n-\ell)-k}{2\ell} + \sum_{\ell=\lfloor n/2 \rfloor+1}^{n} \sum_{k=n-\ell+1}^{\ell-1} \frac{(n-\ell)^2}{2k\ell}$$

$$= o(n^2) + \sum_{\ell=1}^{\lfloor n/2 \rfloor} \frac{4n-5\ell}{4} + \sum_{\ell=\lfloor n/2 \rfloor+1}^{n} \frac{3(n-\ell)^2}{4\ell} + \sum_{\ell=\lfloor n/2 \rfloor+1}^{n} \frac{(n-\ell)^2}{2\ell} \log\left(\frac{\ell}{n-\ell}\right)$$

$$\sim n^2 \left(-\frac{1}{8} + \frac{3}{4} \log 2 + \frac{I}{2}\right),$$

with

$$I := \int_{1/2}^{1} \frac{(1-u)^2}{u} \log\left(\frac{u}{1-u}\right) du$$

= $\left[Li_2(u) + \frac{1}{2}\left(u + \log u(u^2 + \log u - 4u) + \log(1-u)(-u^2 + 4u - 3)\right)\right]_{1/2}^{1},$

with $Li_2(z) := \sum_{k \ge 1} \frac{z^k}{k^2}$. So $I = Li_2(1) - Li_2(1/2) + \frac{1}{4} - \frac{\log^2 2}{2} - \frac{3}{2}\log 2 = \frac{\pi^2}{6} - (\frac{\pi^2}{12} - \frac{\log^2 2}{2}) + \frac{1}{4} - \frac{\log^2 2}{2} - \frac{3}{2}\log 2 = \frac{\pi^2}{12} + \frac{1}{4} - \frac{3}{2}\log 2$. Hence we have $A_{2,1} + 2A_{2,2} \sim \frac{\pi^2}{12}n^2$. • Control of A_3 . Notice that $\sum_{1 \le k_1 < k_2 < \ell_2 < \ell_1 \le n} \mathbb{P}(E_{k_1,\ell_1} \cap E_{k_2,\ell_2})$ and $\sum_{1 \le k_1 < k_2 < \ell_2 < \ell_1 \le n} \mathbb{P}(E_{k_1,\ell_1})\mathbb{P}(E_{k_2,\ell_2})$ are in $n^2 \log n$. But we will see that their difference is in n^2 . According to [8] and to (1),

we have:

$$A_{3} = c_{1} \sum_{1 \le k_{1} \le k_{2} < \ell_{2} < \ell_{1} \le n} \left[\left(\sum_{x} \frac{e^{-\frac{\langle (\Sigma^{2})^{-1}x, x \rangle}{2} \left(\frac{1}{k_{2}-k_{1}} + \frac{1}{\ell_{1}-\ell_{2}}\right)}}{2\pi \sqrt{\det \Sigma^{2}} (k_{2}-k_{1})(\ell_{1}-\ell_{2})} \right) - \frac{1}{(\ell_{1}-k_{1})} \right] \mathbb{P}(E_{k_{2},\ell_{2}}) + o(n^{2}) \quad (4)$$

with the same notations as for (2). Using again (3) and (1), we obtain

$$\begin{aligned} A_3 &= o(n^2) + c_1^2 \sum_{1 \le k_1 < k_2 < \ell_2 \le \ell_1 \le n} \frac{1}{\ell_2 - k_2} \left[\frac{1}{(\ell_1 - k_1) - (\ell_2 - k_2)} - \frac{1}{(\ell_1 - k_1)} \right] \\ &= o(n^2) + c_1^2 \sum_{1 \le k_1 < k_2 < \ell_2 \le \ell_1 \le n} \frac{1}{(\ell_1 - k_1)[(\ell_1 - k_1) - (\ell_2 - k_2)]} \\ &\sim c_1^2 n^2 \int_{[0,1]^4} \frac{1_{\{t+u+v+w < 1\}} dt \, du \, dv \, dw}{(u+w)(u+v+w)} = c_1^2 n^2 \int_{0 \le u \le r \le s \le 1} \frac{(1-s) \, du \, dr \, ds}{rs} = \frac{c_1^2}{2} n^2. \end{aligned}$$

References

- E. Bolthausen. A central limit theorem for two-dimensional random walks in random sceneries. Ann. Probab. 17 (1989) 108115. MR 0972774
- [2] L. A. Bunimovich and Ya. G. Sinai. Markov partitions for dispersed billiards. Commun. Math. Phys. 78 (1980) 247280. MR 0597749
- [3] L. A. Bunimovich and Ya. G. Sinai. Statistical properties of Lorentz gas with periodic configuration of scatterers. Commun. Math. Phys. 78 (1981) 479497. MR 0606459
- [4] L. A. Bunimovich, Ya. G. Sinai and N. I. Chernov. Markov partitions for two-dimensional hyperbolic billiards. Russian Math. Surveys 45 (1990) 105152. (Translation from Uspekhi Mat. Nauk 45 (1990) 97134.) MR 1071936
- [5] L. A. Bunimovich, Ya. G. Sinai and N. I. Chernov. Statistical properties of two-dimensional hyperbolic billiards. Russian Math. Surveys 46 (1991) 47106. (Translation from Usp. Mat. Nauk 46 (1991) 4392.) MR 1138952
- [6] G. Deligiannidis and S. Utev. An asymptotic variance of the self-intersections of random walks. To appear in Annals of Probab.
- [7] D. Dolgopyat, D. Szász and T. Varjú. Recurrence properties of Lorentz gas. Duke Math. J. 142 (2008) 241281. MR 240162
- [8] F. Pène. Planar Lorentz process in a random scenery. Ann. l'Inst. Henri Poincaré, Probabilités et Statistiques 45 (3), (2009) 818–839.
- [9] Ya. G. Sinai. Dynamical systems with elastic reflections. Russian Math. Surveys 25 (1970) 137189.
- [10] D. Szász, T. Varjú. Local limit theorem for the Lorentz process and its recurrence in the plane. Ergodic Theory Dynam. Systems 24 (2004) 257–278. MR 2041271
- [11] L. -S. Young. Statistical properties of dynamical systems with some hyperbolicity. Ann. Math. 147 (1998) 585650. MR 1637655

UNIVERSITÉ DE BREST, UMR CNRS 6205, LABORATOIRE DE MATHÉMATIQUE DE BRETAGNE ATLANTIQUE, 6 AVENUE LE GORGEU, 29238 BREST CEDEX, FRANCE

E-mail address: francoise.pene@univ-brest.fr