Exact sequences in the cohomology of a group extension

J. Huebschmann

USTL, UFR de Mathématiques CNRS-UMR 8524 Labex CEMPI (ANR-11-LABX-0007-01) 59655 Villeneuve d'Ascq Cedex, France Johannes.Huebschmann@math.univ-lille1.fr

October 13, 2018

Dedicated to the memory of my friend and colleague Chih-Han Sah

Abstract

In [DHW12] the authors constructed a seven term exact sequence in the cohomology of a group extension $1 \to N \to G \to Q \to 1$ with coefficients in a *G*-module *M*. However, they were unable to establish the precise link between the maps in that sequence and the corresponding maps arising from the spectral sequence associated to the group extension and the *G*-module *M*. In this paper, we show that there is a close connection between [DHW12] and the two earlier papers [Hue81a] and [Hue81b]. In particular, we show that the results in [Hue81a] and [Hue81b] entail that the maps of [DHW12] other than the obvious inflation and restriction maps do correspond to the corresponding ones arising from the spectral sequence.

2010 Mathematics Subject Classification: 18G40 20J05 20J06 55R20 Keywords and Phrases: Group extension, group cohomology, low degree exact sequences, spectral sequence of a group extension

Contents

1	Introduction	2
2	$d_2 \colon \mathrm{H}^1(N, M)^Q \longrightarrow \mathrm{H}^2(Q, M^N)$	3
	2.1 Description in [Hue81a]	3
	2.2 Description in [Hue81b]	4
	2.3 Relationship between the two previous descriptions	5
	2.4 Construction in [DHW12]	6
	2.5 Comparison of the constructions in [Hue81b] and [DHW12] and proof of The-	
	orem 1	6
3	$\mathrm{H}^{2}(G,M)_{1} \longrightarrow \mathrm{H}^{1}(Q,\mathrm{H}^{1}(N,M))$	7
	3.1 Construction of the map $\rho: \mathrm{H}^2(G, M)_1 \to \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ in [DHW12]	7
	3.2 Crossed pairs	7
	3.3 Proof of Theorem 2	10

References

1 Introduction

Consider an extension of (discrete) groups

$$e_G: 1 \longrightarrow N \xrightarrow{i^N} G \xrightarrow{\pi} Q \longrightarrow 1$$
 (1.1)

and a *G*-module *M*. Let $(E_r^{p,q}(M), d_r)$ denote the spectral sequence associated to the group extension (1.1) and the *G*-module *M*. As usual, for K = N as well as for K = Q, we will use the notation $-^K$ for the invariants relative to the group *K*, and the notation "res" and "inf" will refer to the maps in cohomology induced by the injection of *N* into *G* and by the projection from *G* to *Q*, respectively. Let $H^2(G, M)_1$ denote the kernel of res: $H^2(G, M) \to H^2(N, M)^Q$. It is a straightforward exercise to show that the canonical homomorphism

$$\operatorname{can} \colon \mathrm{H}^{2}(G, M)_{1} \longrightarrow \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M)) \cong \mathrm{E}_{2}^{1,1}(M)$$

induced by the canonical surjection $\mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \to \mathrm{E}^{1,1}_{\infty}(M)$, cf. (3.23) below, extends the classical five term exact sequence in the cohomology of the group extension (1.1) with coefficients in M by two more terms to an exact sequence of the kind

$$0 \longrightarrow \mathrm{H}^{1}(Q, M^{N}) \xrightarrow{\mathrm{inf}} \mathrm{H}^{1}(G, M) \xrightarrow{\mathrm{res}} \mathrm{H}^{1}(N, M)^{Q}$$
$$\xrightarrow{\mathrm{d}_{2}} \mathrm{H}^{2}(Q, M^{N}) \xrightarrow{\mathrm{inf}} \mathrm{H}^{2}(G, M)_{1} \xrightarrow{\mathrm{can}} \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M))$$
$$\xrightarrow{\mathrm{d}_{2}} \mathrm{H}^{3}(Q, M^{N}).$$
$$(1.2)$$

In [Sah74, (3) p. 257], C. H. Sah spelled out this sequence, with the notation "tg" rather than "can".

Let

$$\lambda \colon \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M)) \longrightarrow \mathrm{H}^{3}(Q, M^{N})$$
(1.3)

be the map constructed in [Hue81a] that yields an explicit description of the differential

$$d_2 \colon \mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \longrightarrow \mathrm{H}^3(Q, M^N).$$

In [DHW12], the authors constructed explicit maps

$$\operatorname{tr}: \operatorname{H}^{1}(N, M)^{Q} \longrightarrow \operatorname{H}^{2}(Q, M^{N})$$
(1.4)

$$\rho \colon \mathrm{H}^{2}(G, M)_{1} \longrightarrow \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M))$$
(1.5)

such that, with tr, ρ , and λ substituted for, respectively,

$$d_2 \colon \mathrm{H}^1(N, M)^Q \longrightarrow \mathrm{H}^2(Q, M^N), \tag{1.6}$$

$$\operatorname{can} \colon \mathrm{H}^{2}(G, M)_{1} \longrightarrow \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M)), \tag{1.7}$$

$$d_2 \colon \mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \longrightarrow \mathrm{H}^3(Q, M^N), \tag{1.8}$$

an exact sequence of the kind (1.2) results. They noted [DHW12, p. 71, 4th paragraph] that they did not know whether the maps tr and ρ in their sequence coincided with the maps $d_2: H^1(N, M)^Q \to H^2(Q, M^N)$ and can: $H^2(G, M)_1 \to H^1(Q, H^1(N, M))$, respectively,

that are obtained from the spectral sequence. In the addendum [DHW13] to [DHW12], they compared the conceptual description of (1.4) given in their paper with a conceptual description of a map of the same kind given in [Hue81b] (Section 1.5) and concluded that the descriptions obtained were different.

In this note we shall show that the maps (1.4) and (1.5) are special cases of maps given in [Hue81a] and [Hue81b], see Subsection 2.4 and Section 3 below, that the requisite explicit descriptions may be found in [Hue81a] and [Hue81b], and that the results in [Hue81a] and [Hue81b] entail the following.

Theorem 1. The homomorphism (1.4) constructed in [DHW12] coincides with the differential $d_2: H^1(N, M)^Q \to H^2(Q, M^N)$ of the spectral sequence $(E_r^{p,q}(M), d_r)$.

Theorem 2. The homomorphism (1.5) constructed in [DHW12] coincides with the homomorphism can: $\mathrm{H}^2(G, M)_1 \to \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ arising from the spectral sequence $(\mathrm{E}_r^{p,q}(M), d_r)$.

This settles the questions left open in [DHW12] and [DHW13]. It also shows that the claim in [DHW12, p. 71] to the effect that, in the exact sequence constructed in that paper, there were no explicit descriptions available for the maps other than the inflation map, restriction map and the map λ is unfounded. In particular, [Hue81a, Proposition 3.1] entails an explicit conceptual description of the differential (1.6), and another conceptual description of this differential is given in [Hue81b, Theorem 6, Section 5]. Moreover, in Subsection 2.5 below we shall show that the description of (1.4) given in [DHW12] is essentially the same as that in our paper [Hue81b]. For intelligibility we note that the main result in [Hue81b] is an extension of the classical five term exact sequence by three more terms, viz.

The description of (1.5) given in [DHW12] can be deduced from the construction of the map j, see Section 3 below. The conceptual descriptions of the differentials in [Hue81a] and the construction of the exact sequence (1.9) rely on the interpretation of group cohomology in terms of crossed modules developed in [Hue80].

We dedicate this paper to the memory of Chih-Han Sah, see, e.g., [DPR⁺98]. Sah contributed substantially to the understanding of the spectral sequence of a group extension, cf. [Sah74], [Sah77].

$\mathbf{2} \quad \mathbf{d}_2 \colon \mathrm{H}^1(N, M)^Q \longrightarrow \mathrm{H}^2(Q, M^N)$

We adjust the notation to the present discussion. We denote the group ring of G by $\mathbb{Z}G$, the standard augmentation map by $\varepsilon \colon \mathbb{Z}G \to \mathbb{Z}$, and the augmentation ideal ker(ε) by IG, and we use the same kind of notation for the group ring of Q and the augmentation ideal of Q.

2.1 Description in [Hue81a]

Let $T = \ker(\operatorname{Hom}_N(IG, M) \to \operatorname{Ext}^1_N(\mathbb{Z}, M) \cong \operatorname{H}^1(N, M))$; with respect to the obvious Q-module structures on $\operatorname{Hom}_N(IG, M)$ and $\operatorname{Ext}^1_N(\mathbb{Z}, M)$, the map from $\operatorname{Hom}_N(IG, M)$ to $\operatorname{Ext}^1_N(\mathbb{Z}, M)$ is a morphism of Q-modules, and hence T inherits a Q-module structure.

Consider an extension

$$e_M: 0 \longrightarrow M \xrightarrow{i_M} E_M \xrightarrow{p_M} \mathbb{Z} \longrightarrow 0$$
 (2.1)

of N-modules that represents a member of $\operatorname{Ext}_N(\mathbb{Z}, M)^Q \cong \operatorname{H}^1(N, M)^Q$. Lift the identity of \mathbb{Z} to a commutative diagram

in the category of N-modules. For $x \in G$, let α_x denote the associated operator on IG as well as that on M where the notation is slightly abused; given $q \in Q$, pick $x \in G$ such that $\pi(x) = q$, and define the derivation

$$d: Q \longrightarrow \operatorname{Hom}_{N}(\operatorname{IG}, M), \ d(q) = \alpha_{x} \mu \alpha_{x}^{-1} - \mu, \ q \in Q.$$

$$(2.3)$$

Each value d(q), as q ranges over Q, is well defined, that is, does not depend on the choice of $x \in G$ such that $\pi(x) = q$, and the values of d lie in T since $[e_M] \in \text{Ext}_N(\mathbb{Z}, M)$ is Q-invariant. Consider the semi-direct fiber product group

$$\operatorname{Hom}_{N}(\mathbb{Z}G, M) \rtimes_{T} Q = \{(\varphi, q); \varphi|_{\mathrm{I}G} = d(q)\} \subseteq \operatorname{Hom}_{N}(\mathbb{Z}G, M) \rtimes Q.$$
(2.4)

The projection to Q yields the group extension

$$\widehat{\mathbf{e}}_M : 0 \longrightarrow M^N \longrightarrow \operatorname{Hom}_N(\mathbb{Z}G, M) \rtimes_T Q \longrightarrow Q \longrightarrow 1.$$
 (2.5)

In view of [Hue81a, Proposition 3.1], the assignment to e_M of the class $[\widehat{e}_M] \in H^2(Q, M^N)$ represented by \widehat{e}_M yields a conceptual description of the differential (1.6).

Remark 3. A cocycle description of the differential (1.6) is given in [Hue81a, Remark 3.2]. For the special case where N acts trivially on M, a similar cocycle description can be found in [Sah77, p. 21]. In the latter paper, C. H. Sah explored some of the differentials in the spectral sequence of a split group extension via certain characteristic classes. These characteristic classes were extended in [Hue89]; in that paper, the interested reader will also find more references related with these characteristic classes.

2.2 Description in [Hue81b]

We now recall the construction in [Hue81b, Section 5]. Let $\operatorname{Aut}(\mathbf{e}_M)$ denote the group of pairs (α, x) such that α is an automorphism of E_M whose restriction α_M to M is an automorphism of M and coincides with the automorphism x_M of M induced by $x \in G$ via the G-action on M. Since the class $[\mathbf{e}_M] \in \operatorname{Ext}_N(\mathbb{Z}, M)$ is Q-invariant, every $x \in G$ lifts to a member (α, x) of $\operatorname{Aut}(\mathbf{e}_M)$. Moreover, given $m \in M$, let $\alpha_m \colon E_M \to E_M$ be the automorphism of E_M given by

$$\alpha_m(y) = y + p_M(y)m, \ y \in E_M.$$

The assignment to $m \in M$ of $(\alpha_m, e) \in \overline{\operatorname{Aut}(e_M)}$ and to $(\alpha, x) \in \overline{\operatorname{Aut}(e_M)}$ of $x \in G$ yields a group extension

$$0 \longrightarrow M \xrightarrow{i_M} \overline{\operatorname{Aut}(\mathbf{e}_M)} \xrightarrow{p_M} G \longrightarrow 1.$$
(2.6)

The extension e_M being one of N-modules, the N-action $n \mapsto \alpha_n \colon E_M \to E_M$ $(n \in N)$ on E_M yields the injection

$$s: N \longrightarrow \overline{\operatorname{Aut}(\mathbf{e}_M)}, \ n \longmapsto (\alpha_n, n),$$
 (2.7)

and the composite of this injection with the projection to G coincides with the injection $N \to G$ in (1.1). By construction, the normalizer $\operatorname{Aut}(\mathbf{e}_M)$ of s(N) in $\overline{\operatorname{Aut}(\mathbf{e}_M)}$ consists of the pairs $(\alpha, x) \in \overline{\operatorname{Aut}(\mathbf{e}_M)}$ such that

$$\alpha \alpha_n \alpha^{-1} = \alpha_{xnx^{-1}}, \ n \in N.$$

By [Hue81b, Proposition 1.7], $Aut(e_M)$ still maps onto G, and the surjection onto G yields a group extension

$$0 \longrightarrow M^N \longrightarrow \operatorname{Aut}(\mathbf{e}_M) \xrightarrow{\pi_{\mathbf{e}_M}} G \longrightarrow 1.$$
(2.8)

Passing to quotients, with the notation $Out(e_M) = Aut(e_M)/s(N)$, we obtain the group extension

$$\widetilde{\mathbf{e}}_M \colon 0 \longrightarrow M^N \longrightarrow \operatorname{Out}(\mathbf{e}_M) \longrightarrow Q \longrightarrow 1.$$
 (2.9)

By [Hue81b, Theorem 6], the assignment of \tilde{e}_M to e_M yields a homomorphism

$$\Delta \colon \mathrm{H}^{1}(N, M)^{Q} \longrightarrow \mathrm{Opext}(Q, M^{N})$$
(2.10)

and, via the canonical isomorphism $\text{Opext}(Q, M^N) \cong \text{H}^2(Q, M^N)$, the homomorphism Δ yields a conceptual description of the differential (1.6), viz. $d_2 \colon \text{H}^1(N, M)^Q \longrightarrow \text{H}^2(Q, M^N)$.

2.3 Relationship between the two previous descriptions

Pulling back the group (2.4) through the projection $G \to Q$ in (1.1), we obtain the group

$$\operatorname{Hom}_{N}(\mathbb{Z}G, M) \rtimes_{T} G = \{(\varphi, x); \varphi|_{IG} = \alpha_{x}\mu\alpha_{x}^{-1} - \mu\} \subseteq \operatorname{Hom}_{N}(\mathbb{Z}G, M) \rtimes G.$$
(2.11)
Given $(\varphi, y) \in \operatorname{Hom}_{N}(\mathbb{Z}G, M) \rtimes G$, define

$$\alpha_1 \colon \mathbb{Z}G \to E_M, \ \alpha_1(x) = \varphi(yx) + \nu(yx), \ x \in \mathbb{Z}G,$$
$$\alpha_2 \colon E_M \to E_M, \ \alpha_2(m) = ym.$$

Since the left-hand square of (2.2) is a pushout diagram of abelian groups, α_1 and α_2 induce a unique automorphism

$$\alpha_{\varphi,y}\colon E_M\longrightarrow E_M.$$

By [Hue81b, Lemma 5.1], the assignment of $\alpha_{\varphi,y}$ to $(\varphi, y) \in \text{Hom}_N(\mathbb{Z}G, M) \rtimes_T G$ yields a morphism

of group extensions. The resulting morphism

of group extensions identifies $Out(e_M)$ with $Hom_N(\mathbb{Z}G, M) \rtimes_T Q$. In view of [Hue81a, Proposition 3.1], diagram (2.13) entails a proof of Theorem 6 in [Hue81b].

2.4 Construction in [DHW12]

The construction of (1.4) in [DHW12, Section 5] comes down to this: Consider the split extension

$$0 \longrightarrow M \longrightarrow M \rtimes G \longrightarrow G \longrightarrow 1.$$
 (2.14)

A section $s: N \to M \rtimes G$ whose composite with the projection to G coincides with the injection $N \to G$ in $e_G: 1 \to N \xrightarrow{i^N} G \xrightarrow{\pi} Q \to 1$ is determined by its component $\varphi: N \to M$, and φ is a 1-cocycle or, equivalently, derivation, on N with values in M that represents a Q-invariant cohomology class $[\varphi] \in H^1(N, M)^Q$. Given such a section s or, equivalently, 1-cocycle φ , the normalizer $N_{M \rtimes G}(sN)$ of sN in $M \rtimes G$ maps onto G, the sequence

$$0 \longrightarrow M^N \longrightarrow N_{M \rtimes G}(sN) \longrightarrow G \longrightarrow 1$$
 (2.15)

is exact, and so is

$$\widetilde{\mathbf{e}}: 0 \longrightarrow M^N \longrightarrow N_{M \rtimes G}(sN)/sN \longrightarrow Q \longrightarrow 1.$$
 (2.16)

In [DHW12] the map (1.4) is essentially defined by the assignment to an M-valued 1-cocycle φ on N representing a Q-invariant cohomology class in $\mathrm{H}^1(N, M)$ of the class of the extension $\widetilde{\mathrm{e}}$. The construction in [DHW12] is actually couched in what is referred to as "partial semi-direct complements". The 1-cocycle φ is lurking behind the choice of the subgroup of $M \rtimes G$ denoted in [DHW12] by H, indeed, it is the component into M of the resulting section $s: N \to M \rtimes G$ such that H = sN.

2.5 Comparison of the constructions in [Hue81b] and [DHW12] and proof of Theorem 1

We maintain the notation established before. Thus $e_M: 0 \to M \xrightarrow{i_M} E_M \xrightarrow{p_M} \mathbb{Z} \to 0$ is an extension of N-modules of the kind (2.1) that represents a class

$$[\mathbf{e}_M] \in \operatorname{Ext}^1_N(\mathbb{Z}, M)^Q \cong \mathrm{H}^1(N, M)^Q,$$

and $s: N \to M \rtimes G$ is a section as in Subsection 2.4 above.

As an abelian group, E_M decomposes as a direct sum $M \oplus \mathbb{Z}$. Such a decomposition induces an obvious *G*-module structure on E_M that restricts to the *G*-module structure on M. This *G*-module structure induces a section $\sigma: G \to \overline{\operatorname{Aut}}(\mathbf{e}_M)$ for the extension (2.6) and hence an isomorphism $\overline{\operatorname{Aut}}(\mathbf{e}_M) \to M \rtimes G$. This isomorphism induces a morphism

of group extensions. Hence, cf. the introductory phrase of Subsection 2.2 above, the construction in [DHW12, Section 5] that underlies the map (1.4) is exactly the same as that in [Hue81b] which underlies the map (2.10). The only difference is that the interpretation of the naive semi-direct product $M \rtimes G$ as the group $\overline{\operatorname{Aut}(e_M)}$, not noticed in [DHW12], enabled us to show in [Hue81b] that (2.10) actually yields the differential (1.6). In particular, the map (1.4) coincides with the differential (1.6). This establishes Theorem 1. Furthermore, the claim in [DHW13] to the effect that the construction of the map (1.4) in [DHW12] was much more elementary than the construction of (2.10) in [Hue81b] is unfounded.

$\mathbf{3} \quad \mathrm{H}^2(G, M)_1 \longrightarrow \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$

We note first that, the actions of the various groups on a module being written as juxtaposition, the standard G-module structure on Der(N, M) is given by the familiar formula

$$(yd)(n) = y(d(y^{-1}ny)), \ d \in \text{Der}(N, M), \ y \in G, \ n \in N;$$

this G-module structure induces, through the projection from E to G, an E-module structure on Der(N, M), and this structure, in turn, induces the standard Q-module structure on $\text{H}^1(N, M)$.

3.1 Construction of the map $\rho: H^2(G, M)_1 \to H^1(Q, H^1(N, M))$ in [DHW12]

Consider a group extension

$$e_G^M : 0 \longrightarrow M \xrightarrow{i} E \xrightarrow{p} G \longrightarrow 1.$$
 (3.1)

Suppose that the restriction of e_G^M to N splits, and let $s: N \to E$ be a homomorphism such that $p \circ s$ coincides with the injection of N into G. Given $x \in E$, we denote its image $p(x) \in G$ by [x]. The assignment to $x \in E$ of

$$\widehat{\delta}_x \colon N \longrightarrow M, \ i(\widehat{\delta}_x(n)) = xs([x^{-1}]n[x])x^{-1}s(n^{-1}) \in i(M) \subseteq E, \ n \in N,$$
(3.2)

yields a derivation $\widehat{\delta}_{e_G^M} : E \to \text{Der}(N, M)$. Given $x \in E_N = p^{-1}(N)$, the member $y = (s[x])x^{-1}$ of E_N lies in the image i(M) of M in E_N , and

$$i(\widehat{\delta}_x(n)) = y^{-1}s(n)ys(n^{-1}) \in i(M) \subseteq E_N, \ n \in N,$$

that is, $\hat{\delta}_x$ is an inner derivation. Consequently the derivation $\hat{\delta}_{e^M_G}$, in turn, passes to a derivation

$$\delta_{\mathbf{e}_{G}^{M}} \colon Q \longrightarrow \mathrm{H}^{1}(N, M) \tag{3.3}$$

and, in [DHW12, Section 6], the map $\rho: \mathrm{H}^2(G, M)_1 \to \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ is defined by the assignment of $\delta_{\mathrm{e}_G^M}$ to e_G^M . While the derivations $\widehat{\delta}_{\mathrm{e}_G^M}$ and $\delta_{\mathrm{e}_G^M}$ depend on the choice of the homomorphism $s: N \to E$, the class $[\delta_{\mathrm{e}_G^M}] \in \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ is independent of that choice.

3.2 Crossed pairs

At the referee's request, for intelligibility, we will now briefly reproduce some material from [Hue81b]. To this end, we write the G-action on M as

$$G \times M \longrightarrow M, \ (x, y) \longmapsto {}^{x}y, \ x \in G, \ y \in M.$$

Let

$$e: 0 \longrightarrow M \longrightarrow E_N \xrightarrow{\pi_N} N \longrightarrow 1$$

be a group extension whose class $[e] \in H^2(N, M)$ is fixed under the standard Q-action on $H^2(N, M)$. Given $x \in G$, we write

$$\ell_x(y) = {}^x y, \ y \in M, \quad i_x(n) = xnx^{-1}, \ n \in N.$$

Write $\operatorname{Aut}_G(e)$ for the subgroup of $\operatorname{Aut}(E_N) \times G$ that consists of pairs (κ, x) which make the diagram

commutative.

The homomorphism

$$\beta \colon E_N \longrightarrow \operatorname{Aut}_G(\mathbf{e}), \ \beta(y) = (i_y, i^N(\pi_N(y))), \ y \in E_N,$$

together with the obvious action of $\operatorname{Aut}_G(e)$ on E_N , yields a crossed module $(E_N, \operatorname{Aut}_G(e), \beta)$ whence, in particular, $\beta(E_N)$ is a normal subgroup of $\operatorname{Aut}_G(e)$; with the notation $\operatorname{Out}_G(e)$ for the cokernel of β , the resulting crossed 2-fold extension has the form

 $0 \longrightarrow M^N \longrightarrow E_N \xrightarrow{\beta} \operatorname{Aut}_G(e) \longrightarrow \operatorname{Out}_G(e) \longrightarrow 1.$ (3.4)

The association

$$\operatorname{Der}(N,M) \ni d \longmapsto (\kappa_d, 1), \ \kappa_d(y) = (d\pi_N(y))y, \ y \in E_N,$$
(3.5)

yields an injective homomorphism $\vartheta \colon \text{Der}(N, M) \longrightarrow \text{Aut}_G(e)$; this homomorphism and the obvious map $\text{Aut}_G(e) \to G$ yield a group extension

$$0 \longrightarrow \operatorname{Der}(N, M) \xrightarrow{\vartheta} \operatorname{Aut}_{G}(e) \longrightarrow G \longrightarrow 1,$$
(3.6)

the map $\operatorname{Aut}_G(\mathbf{e}) \to G$ being surjective, since the class $[\mathbf{e}] \in \operatorname{H}^2(N, M)$ is supposed to be fixed under Q. Further, the formula $(\zeta(m))(n) = m(nm)^{-1}$, as m ranges over M and n over N, defines a homomorphism $\zeta \colon M \to \operatorname{Der}(N, M)$. With these preparations out of the way, the various groups under discussion fit into the commutative diagram

with exact rows and columns. We shall use the notation

$$\overline{\mathbf{e}} \colon 0 \longrightarrow \mathrm{H}^1(N, M) \longrightarrow \mathrm{Out}_G(\mathbf{e}) \longrightarrow Q \longrightarrow 1$$
 (3.8)

for the bottom row extension of (3.7). This extension is the cokernel, in the category of group extensions with abelian kernel, of the morphism (ζ, β, i) of group extensions.

Suppose now that the group extension $\overline{\mathbf{e}}$ splits; we then say that the group extension e: $0 \to M \to E_N \xrightarrow{\pi_N} N \to 1$ admits a crossed pair structure with respect to the group extension (1.1), and we refer to a section $\psi: Q \to \operatorname{Out}_G(\mathbf{e})$ of $\overline{\mathbf{e}}$ as a crossed pair structure on the group extension \mathbf{e} with respect to the group extension (1.1). By definition, a crossed pair (\mathbf{e}, ψ) with respect to the group extension (1.1) and the *G*-module *M* consists of a group extension $\mathbf{e}: 0 \to M \to E_N \to N \to 1$ whose class $[\mathbf{e}] \in \operatorname{H}^2(N, M)$ is fixed under *Q* such that the associated extension $\overline{\mathbf{e}}$ splits, together with a section $\psi: Q \to \operatorname{Out}_G(\mathbf{e})$ of $\overline{\mathbf{e}}$ [Hue81b, p. 152].

Remark 4. By [Hue81a, Theorem 1], the association $e \mapsto \overline{e}$ yields a conceptual description of the differential $d_2: E_2^{0,2}(M) \to E_2^{2,1}(M)$ of the spectral sequence $(E_r^{p,q}(M), d_r)$ associated to the group extension (1.1) and the *G*-module *M*.

Under a kind of generalized Baer sum, suitable classes of crossed pairs with respect to (1.1) and the *G*-module *M* constitute an abelian group Xpext(G, N; M) [Hue81b, Theorem 1]. Moreover, cf. [Hue81b, Theorem 2], suitably defined homomorphisms

$$j: \mathrm{H}^2(G, M) \longrightarrow \mathrm{Xpext}(G, N; M), \ \Delta: \mathrm{Xpext}(G, N; M) \longrightarrow \mathrm{H}^3(Q, M^N)$$

yield an extension of the classical five term exact sequence to an eight term exact sequence of the kind (1.9).

To recall the construction of j, consider a group extension

$$e_G^M : 0 \longrightarrow M \xrightarrow{i} E \xrightarrow{p} G \longrightarrow 1;$$

the restriction of e_G^M to N has the form $e: 0 \to M \to E_N \to N \to 1$, and conjugation in E induces a crossed pair structure $\psi_E: Q \to \text{Out}_G(e)$ on e (section for (3.8)) with respect to (1.1). The map j is given by the assignment to a group extension of the kind e_G^M of the crossed pair

$$(e: 0 \to M \to E_N \to N \to 1, \psi_E: Q \to \operatorname{Out}_G(e))$$

with respect to (1.1).

To recall the construction of Δ , given a crossed pair

(e:
$$0 \to M \to E_N \to N \to 1, \ \psi : Q \to \operatorname{Out}_G(e)$$
)

with respect to the group extension (1.1) and the *G*-module M, let B^{ψ} denote the fiber product group $\operatorname{Aut}_{G}(\mathbf{e}) \times_{\operatorname{Out}_{G}(\mathbf{e})} Q$ with respect to the crossed pair structure map $\psi \colon Q \to \operatorname{Out}_{G}(\mathbf{e})$ and, furthermore, let $\partial^{\psi} \colon E_{N} \to B^{\psi}$ denote the obvious homomorphism; together with the obvious action of B^{ψ} on E_{N} induced by the canonical homomorphism $B^{\psi} \to \operatorname{Aut}_{G}(\mathbf{e})$, the exact sequence

$$e_{\psi} \colon 0 \longrightarrow M^N \longrightarrow E_N \xrightarrow{\partial^{\psi}} B^{\psi} \longrightarrow Q \longrightarrow 1$$
 (3.9)

is a crossed 2-fold extension and hence represents a class in $\mathrm{H}^3(Q, M^N)$. We shall refer to e_{ψ} as the crossed 2-fold extension associated to the crossed pair (e, ψ) . The homomorphism Δ : Xpext $(G, N; M) \to \mathrm{H}^3(Q, M^N)$ is given by the assignment to a crossed pair (e, ψ) with respect to the group extension (1.1) and the G-module M of its associated crossed 2-fold extension e_{ψ} . Next we recall the injection $\alpha: \mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \longrightarrow \mathrm{Xpext}(G, N; M)$ constructed in [Hue81b, Subsection 1.3]. To this end, let e_s denote the split extension

 $\mathbf{e}_s \colon 0 \longrightarrow M \longrightarrow M \rtimes N \longrightarrow N \longrightarrow 1.$

The obvious action of G on $M \rtimes N$ induces, on \mathbf{e}_s , a canonical crossed pair structure $\psi_s \colon Q \to \operatorname{Out}_G(\mathbf{e}_s)$ with respect to (1.1). Given a derivation $\delta \colon Q \to \operatorname{H}^1(N, M)$, define the crossed pair structure

$$\psi_{\delta} \colon Q \longrightarrow \operatorname{Out}_{G}(\mathbf{e}_{s})$$

on e_s with respect to (1.1) by

$$\psi_{\delta}(x) = \delta(x)\psi_s(x), \ x \in Q; \tag{3.10}$$

here we identify in notation the member $\delta(x)$ of $\mathrm{H}^1(N, M)$ with its image in $\mathrm{Out}_G(\mathbf{e})$ under the injection $\mathrm{H}^1(N, M) \to \mathrm{Out}_G(\mathbf{e})$ in (3.8). The assignment to a derivation $\delta \colon Q \to \mathrm{H}^1(N, M)$ of the crossed pair $(\mathbf{e}_s, \psi_\delta)$ with respect to (1.1) and the *G*-module *M* induces a homomorphism

$$\alpha \colon \mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \longrightarrow \mathrm{Xpext}(G, N; M).$$

Let

$$\xi \colon \operatorname{Xpext}(G, N; M) \longrightarrow \operatorname{H}^0(Q, \operatorname{H}^2(N, M))$$

denote the canonical forgetful map. By [Hue81b, Theorem 3], the sequence

$$0 \to \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M)) \xrightarrow{\alpha} \mathrm{Xpext}(G, N; M) \xrightarrow{\xi} \mathrm{H}^{0}(Q, \mathrm{H}^{2}(N, M)) \xrightarrow{d_{2}} \mathrm{H}^{2}(Q, \mathrm{H}^{1}(N, M))$$
(3.11)

is exact and, furthermore, natural in the data. The exactness of (3.11) at Xpext(G, N; M) entails that, via α , the group $\mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ acts faithfully and transitively on the classes of crossed pair structures $\psi: Q \to \mathrm{Out}_G(\mathbf{e}_s)$ on the split extension

$$\mathbf{e}_s \colon 0 \longrightarrow M \longrightarrow M \rtimes N \longrightarrow N \longrightarrow 1.$$

Somewhat more explicitly, this action is characterized by the bijective correspondence between classes of crossed pair structures on e_s and cohomology classes of derivations $Q \to H^1(N, M)$ induced by (3.10) above.

3.3 Proof of Theorem 2

Below we will use the notation "can" for any map that is canonically defined.

Proposition 3.1. The diagram

is commutative with exact rows.

Proof. This is a consequence of the commutativity of diagram [Hue81b, (1.10)].

Proposition 3.2. The diagram

$$\begin{array}{cccc} \mathrm{H}^{2}(G,M)_{1} & \stackrel{\rho}{\longrightarrow} & \mathrm{H}^{1}(Q,\mathrm{H}^{1}(N,M)) & \stackrel{d_{2}}{\longrightarrow} & \mathrm{H}^{3}(Q,M^{N}) \\ & & & & & \\ \mathrm{can} & & & & & \\ \mathrm{H}^{2}(G,M) & \stackrel{j}{\longrightarrow} & \mathrm{Xpext}(G,N;M) & \stackrel{\Delta}{\longrightarrow} & \mathrm{H}^{3}(Q,M^{N}) \end{array}$$

$$(3.13)$$

is commutative with exact rows.

Proof. The commutativity of the right-hand square is implied by the commutativity of diagram (3.12).

We will now show that the left-hand square in (3.13) is commutative. To this end, consider a group extension

$$e_G^M \colon 0 \longrightarrow M \xrightarrow{i} E \xrightarrow{p} G \longrightarrow 1$$

that represents a member of $\mathrm{H}^2(G, M)_1$. Then the restriction $e: 0 \to M \to E_N \to N \to 1$ of e_G^M to N splits; choose a section $s: N \to E_N$, and write E_N as $M \rtimes (sN)$. By the construction of j, the group $M \rtimes (sN)$ being normal in E, conjugation in E induces a crossed pair structure $\psi_E: Q \to \mathrm{Out}_G(e)$ on e (section for (3.8)) with respect to (1.1), and the crossed pair (e, ψ_E) with respect to (1.1) represents the image $j[\mathrm{e}_G^M] \in \mathrm{Xpext}(G,N;M)$ of the class $[\mathrm{e}_G^M] \in \mathrm{H}^2(G,M)_1$. By construction, $\xi(j[\mathrm{e}_G^M]) = 0$ whence

$$j[\mathbf{e}_G^M] \in \ker(\xi) = \alpha(\mathrm{H}^1(Q, \mathrm{H}^1(N, M))) \subseteq \mathrm{Xpext}(G, N; M).$$

The canonical action of G on $M \rtimes (sN)$ induces a section $\Psi_s \colon G \to \operatorname{Aut}_G(e)$ for (3.6). On the other hand, conjugation in E induces likewise a section $\Psi_{e_G^M} \colon E \to \operatorname{Aut}_G(e)$ for (3.6), and this section, in turn, induces the crossed pair structure $\psi_E \colon Q \to \operatorname{Out}_G(e)$ on e with respect to (1.1). Recall the injection $\vartheta \colon \operatorname{Der}(N, M) \to \operatorname{Aut}_G(e)$ characterized by (3.5).

Proposition 3.3. The derivation $\widehat{\delta}_{\mathbf{e}_{C}^{M}}: E \to \mathrm{Der}(N, M)$ satisfies the identity

$$\Psi_{\mathbf{e}_{G}^{M}}(x) = \vartheta(\widehat{\delta}_{\mathbf{e}_{G}^{M}}(x))\Psi_{s}([x]), \ x \in E.$$
(3.14)

Proof. Define the derivation $\widetilde{\delta}_{\mathbf{e}_{C}^{M}} : E \to \operatorname{Der}(N, M)$ by the identity

$$\Psi_{\mathbf{e}_{G}^{M}}(x) = \vartheta(\widetilde{\delta}_{\mathbf{e}_{G}^{M}}(x))\Psi_{s}([x]), \ x \in E.$$
(3.15)

We shall show that $\widehat{\delta}_{\mathbf{e}_{G}^{M}}=\widetilde{\delta}_{\mathbf{e}_{G}^{M}}.$

By construction, an explicit expression for $\Psi_s \colon G \to \operatorname{Aut}_G(e)$ is given by

$$(\Psi_s(x))(m, s(n)) = ({}^xm, s(xnx^{-1})), \ x \in G, \ m \in M, \ n \in N.$$

Let $x \in E$, $m \in M$, and $n \in N$. Since $i([x]m) = xi(m)x^{-1}$, in view of the identity (3.5) characterizing the injection $\vartheta \colon \text{Der}(N, M) \to \text{Aut}_G(e)$,

$$\begin{split} \vartheta(\widetilde{\delta}_{\mathbf{e}_{G}^{M}}(x))\Psi_{s}([x])(i(m)s(n)) &= \vartheta(\widetilde{\delta}_{\mathbf{e}_{G}^{M}}(x))(xi(m)x^{-1}s([x]n[x]^{-1})) \\ &= i(\widetilde{\delta}_{\mathbf{e}_{G}^{M}}(x)([x]n[x]^{-1}))xi(m)x^{-1}s([x]n[x]^{-1}). \end{split}$$

Since, by construction,

$$(\Psi_{\mathbf{e}_{G}^{M}}(x))(i(m)s(n)) = i({}^{[x]}m)xs(n)x^{-1},$$
(3.16)

from (3.15), we deduce the identity

$$i({}^{[x]}m)xs(n)x^{-1} = i(\widetilde{\delta}_{\mathbf{e}_{G}^{M}}(x)([x]n[x]^{-1}))xi(m)x^{-1}s([x]n[x]^{-1}).$$
(3.17)

To simplify the notation, we will now write $\tilde{\delta} = \tilde{\delta}_{e_G^M}$ and $\tilde{\delta}_x = \tilde{\delta}(x)$. Then the identity (3.17) takes the form

$$i({}^{[x]}m)xs(n)x^{-1} = i(\widetilde{\delta}_x([x]n[x]^{-1}))xi(m)x^{-1}s([x]n[x]^{-1})$$
(3.18)

or, since M is abelian,

$$i({}^{[x]}m)xs(n)x^{-1} = i({}^{[x]}m)i(\widetilde{\delta}_x([x]n[x]^{-1}))s([x]n[x]^{-1}),$$
(3.19)

and thence

$$xs(n)x^{-1} = i(\widetilde{\delta}_x([x]n[x]^{-1}))s([x]n[x]^{-1}), \qquad (3.20)$$

whence

$$i(\widetilde{\delta}_x([x]n[x]^{-1})) = xs(n)x^{-1}s([x]n^{-1}[x]^{-1}), \qquad (3.21)$$

or, equivalently,

$$i(\tilde{\delta}_x(n)) = x(s([x]^{-1}n[x]))x^{-1}s(n^{-1}).$$
(3.22)

In view of the identity (3.2) characterizing $\hat{\delta}_{e_G^M}$, we conclude that $\hat{\delta}_{e_G^M} = \tilde{\delta}_{e_G^M}$.

We now complete the proof of Proposition 3.2. The commutativity of the left-hand square in (3.13) is a consequence of Proposition 3.3. Indeed, once a choice of section s for the restriction $e: 0 \to M \to E_N \to N \to 1$ of $e_G^M: 0 \to M \xrightarrow{i} E \xrightarrow{p} G \to 1$ to N has been made, the derivation $\delta_{e_G^M}: Q \to H^1(N, M)$ represents the image $\rho([e_G^M]) \in H^1(Q, H^1(N, M))$ of $[e_G^M] \in H^2(G, M)_1$. Now the section $\Psi_{e_G^M}: E \to \operatorname{Aut}_G(e)$ for (3.6) induces the crossed pair structure $\psi_E: Q \to \operatorname{Out}_G(e)$ on e that yields the image $j([e_G^M]) \in \operatorname{Xpext}(G, N; M)$. On the other hand, the derivation $\widehat{\delta}_{e_G^M}: E \to \operatorname{Der}(N, M)$ induces the derivation $\delta_{e_G^M}: Q \to H^1(N, M)$. Hence the identity (3.14) entails that, by its very construction, the injective homomorphism $\alpha: \operatorname{H}^1(Q, \operatorname{H}^1(N, M)) \to \operatorname{Xpext}(G, N; M)$ recovers the class of the crossed pair structure $\psi_E: Q \to \operatorname{Out}_G(e)$ on e from $\rho([e_G^M]) = [\delta_{e_G^M}] \in \operatorname{H}^1(Q, \operatorname{H}^1(N, M))$.

The homomorphism $\alpha \colon \mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \longrightarrow \mathrm{Xpext}(G, N; M)$ is injective whence, by the commutativity of the left-hand square in (3.13), $\rho \colon \mathrm{H}^2(G, M)_1 \to \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ arises as the restriction of $j \colon \mathrm{H}^2(G, M) \to \mathrm{Xpext}(G, N; M)$ to $\mathrm{H}^2(G, M)_1$, and the exact sequence in [DHW12] can be deduced from the exact sequence (1.9) constructed in [Hue81b]. The commutativity of (3.13) and the exactness of the bottom row of that diagram entail that the homomorphism $\rho \colon \mathrm{H}^2(G, M)_1 \to \mathrm{H}^1(Q, \mathrm{H}^1(N, M))$ coincides with the canonical homomorphism

$$\operatorname{can} \colon \mathrm{H}^{2}(G, M)_{1} \longrightarrow \mathrm{E}^{1,1}_{\infty}(M) \longrightarrow \mathrm{E}^{1,1}_{2}(M) \cong \mathrm{H}^{1}(Q, \mathrm{H}^{1}(N, M))$$
(3.23)

that comes from the spectral sequence $(\mathcal{E}_r^{p,q}(M), d_r)$. This establishes Theorem 2. In particular, $\rho \colon \mathcal{H}^2(G, M)_1 \to \mathcal{H}^1(Q, \mathcal{H}^1(N, M))$ maps $\mathcal{H}^2(G, M)_1$ onto

$$\mathbf{E}_{\infty}^{1,1} = \ker(d_2 \colon \mathrm{H}^1(Q, \mathrm{H}^1(N, M)) \to \mathrm{H}^3(Q, M^N)).$$

Remark 5. A derivation or 1-cocycle of the kind (3.3) or of the kind δ in (3.10) can be seen as a kind of *Eilenberg difference cocycle* [Eil40, p. 236]. The identity (3.2) spells out $\hat{\delta}_x$ as the "difference" of the two sections xs and s from N to E for the restriction of the group extension $e_G^M: 0 \to M \to E \to G \to 1$ to N. The key step in the proof of Theorem 2 consists in making the requisite difference cocycle explicit via the identity (3.14); this identity displays the 1-cocycle $\hat{\delta}_{e_G^M}: E \to \text{Der}(N, M)$ as the "difference" of the two sections $\Psi_{e_G^M}$ and Ψ_s for the corresponding extension (3.6).

References

- [DHW12] Karel Dekimpe, Manfred Hartl, and Sarah Wauters. A seven-term exact sequence for the cohomology of a group extension. J. Algebra, 369:70–95, 2012. arXiv:1103.4052.
- [DHW13] Karel Dekimpe, Manfred Hartl, and Sarah Wauters. Addendum to "A seven-term exact sequence for the cohomology of a group extension" [J. Algebra 369 (1) (2012) 70–95]. J. Algebra, 373:439–440, 2013.
- [DPR+98] Johan Dupont, Anthony Phillips, Vladimir Retakh, Judith Roitman, and Mark Saul. Chih-Han Sah (1934–1997). Notices Amer. Math. Soc., 45(1):79–83, 1998.
- [Eil40] Samuel Eilenberg. Cohomology and continuous mappings. Ann. of Math. (2), 41:231–251, 1940.
- [Hue80] Johannes Huebschmann. Crossed *n*-fold extensions of groups and cohomology. Comment. Math. Helv., 55(2):302–313, 1980.
- [Hue81a] Johannes Huebschmann. Automorphisms of group extensions and differentials in the Lyndon-Hochschild-Serre spectral sequence. J. Algebra, 72(2):296–334, 1981.
- [Hue81b] Johannes Huebschmann. Group extensions, crossed pairs and an eight term exact sequence. J. Reine Angew. Math., 321:150–172, 1981.
- [Hue89] Johannes Huebschmann. Change of rings and characteristic classes. Math. Proc. Cambridge Philos. Soc., 106(1):29–56, 1989.
- [Sah74] Chih Han Sah. Cohomology of split group extensions. J. Algebra, 29:255–302, 1974.
- [Sah77] Chih Han Sah. Cohomology of split group extensions. II. J. Algebra, 45(1):17–68, 1977.