

RELATIVE HOMOLOGICAL ALGEBRA, WALDHAUSEN K -THEORY, AND QUASI-FROBENIUS CONDITIONS.

ANDREW SALCH

ABSTRACT. We study the question of the existence of a Waldhausen category on any (relative) abelian category in which the contractible objects are the (relatively) projective objects. The associated K -theory groups are “stable algebraic G -theory,” which in degree zero form a certain stable representation group. We prove both some existence and nonexistence results about such Waldhausen category structures, including the fact that, while it was known that the category of R -modules admits a model category structure if R is quasi-Frobenius, that assumption is required even to get a Waldhausen category structure with cylinder functor—i.e., Waldhausen categories do not offer a more general framework than model categories for studying stable representation theory of rings. We study multiplicative structures on these Waldhausen categories, and we relate stable algebraic G -theory to algebraic K -theory and we compute stable algebraic G -theory for finite-dimensional quasi-Frobenius nilpotent extensions of finite fields. Finally, we show that the connective stable G -theory spectrum of $\mathbb{F}_p[x]/x^{p^n}$ is a complex oriented ring spectrum, partially answering a question of J. Morava about complex orientations on algebraic K -theory spectra.

CONTENTS

1. Introduction.	1
2. Waldhausen category structures from allowable classes on abelian categories.	4
2.1. Definitions.	4
2.2. The Waldhausen category structure on an abelian category associated to a pair of allowable classes.	9
3. Absolute and relative quasi-Frobenius conditions.	13
3.1. Definitions.	13
3.2. Existence of cylinder functors.	14
3.3. Multiplicative structure.	17
4. Applications.	19
4.1. The relationship between stable G -theory and other G -theories and K -theories.	19
4.2. Applications to algebras.	21
References	25

1. INTRODUCTION.

Waldhausen’s paper [12] defines several kinds of categorical structure which are meaningful for algebraic K -theory. A *category with cofibrations and weak equivalences*, also called a *Waldhausen category*, has just enough structure for Waldhausen’s machinery to

produce an associated K -theory infinite loop space. A Waldhausen category which satisfies additional axioms and/or has additional structure will have better properties which e.g. make the problem of actually computing the associated K -theory more tractable. For example, a Waldhausen category satisfying the “extension axiom” and the “saturation axiom” and equipped with an additional structure called a “cylinder functor” admits Waldhausen’s *Localization Theorem* (see [12]), a computationally powerful result that describes the sense in which localizations of the Waldhausen category induce long exact sequences in the K -theory groups.

Meanwhile, in stable representation theory, one regards the projective modules over a ring as “contractible,” and maps of modules that factor through projective modules are regarded as “nulhomotopic.” This suggests that the category of modules over a ring perhaps has a Waldhausen category structure in which the objects weakly equivalent to zero—the contractible objects—are precisely the projective modules. More generally, one has the tools of *relative homological algebra*: if one chooses a sufficiently well-behaved class of objects in an abelian category, one can do a form of homological algebra in which the chosen class of objects plays the role of projective objects. Given an abelian category and a class of relative projective objects, one wants to know if there is a natural Waldhausen category structure on that abelian category, such that the contractible objects are precisely the relative projectives. One also wants to know how many extra axioms are satisfied by, and how much additional structure is admitted by, such a Waldhausen category.

In this paper we prove the following theorems that answer the above questions, and explain fundamental properties of the relationships between Waldhausen K -theory, relative homological algebra, and stable representation theory:

- (1) Definition-Proposition 2.2.5: Given an abelian category C and a sufficiently nice pair of allowable classes E, F in C , there exists a Waldhausen category structure on C whose weak equivalences are the E -stable equivalences and whose cofibrations are the F -monomorphisms. In particular, the E -projective objects are precisely the contractible objects in this Waldhausen category. This Waldhausen category satisfies the saturation axiom and the extension axiom.
- (2) Theorem 3.2.3: If C has enough injectives, then C has a cylinder functor satisfying the cylinder axiom if and only if C obeys a certain generalized quasi-Frobenius condition: every object must functorially embed in an E -projective object by an F -monomorphism.
- (3) As a consequence, we have Corollary 3.2.4: any quasi-Frobenius abelian category with enough projectives and functorially enough injectives admits the structure of a Waldhausen category whose weak equivalences are the stable equivalences and whose cofibrations are the monomorphisms. This Waldhausen category has a cylinder functor, and it satisfies the saturation, extension, and cylinder axioms.
- (4) As a consequence, we have Corollary 3.2.5: if R is a finite-dimensional quasi-Frobenius algebra over a finite field, then the category of finitely generated (left) R -modules admits the structure of a Waldhausen category in which the cofibrations are the monomorphisms and the weak equivalences are the stable equivalences. This Waldhausen category satisfies the saturation and extension axioms, and it admits a cylinder functor satisfying the cylinder axiom.
- (5) In Proposition 3.3.4 we show that, for a finite-dimensional co-commutative Hopf algebra over a finite field, this Waldhausen category is even better: it has a multiplicative structure coming from the tensor product of modules over the base

- field, and this multiplicative structure gives rise to the structure of a homotopy-commutative ring spectrum on the Waldhausen K -theory spectrum of the category.
- (6) We use the theorems described above to prove our Proposition 4.2.1, in which we show that, for finite-dimensional quasi-Frobenius nilpotent extensions of finite fields, stable G -theory is a delooping of relative algebraic K -theory.
 - (7) As a consequence, we get our Theorem 4.2.3, in which we prove that, under the same assumptions on the ring, stable G -theory in positive degrees vanishes ℓ -adically and is isomorphic to topological cyclic homology with a degree shift p -adically, where p is the characteristic of the base field.
 - (8) Finally, in Theorem 4.2.6, we compute the homotopy groups of the connective cover of the stable G -theory spectrum $g_{st}(\mathbb{F}_p[x]/x^{p^n})$ of truncated polynomial algebras over \mathbb{F}_p , in terms of the Hesselholt-Madsen description of the topological cyclic homology of truncated polynomial algebras. We also provide one possible answer to a question (unpublished) of J. Morava: under what circumstances does algebraic K -theory, or a localization or other modification thereof, admit the structure of a complex oriented ring spectrum? Our answer is that the connective stable G -theory spectrum $g_{st}(\mathbb{F}_p[x]/x^{p^n})$ is, for all p and n , a complex oriented ring spectrum.

“Quasi-Frobenius conditions” appear prominently throughout this paper. Recall that a ring R is said to be quasi-Frobenius if every projective R -module is injective and vice versa. The appearance of these conditions in connection with Waldhausen K -theory stems from the theorem of Faith and Walker (see [2] for a good account of this and related theorems):

Theorem 1.0.1. (Faith-Walker.) *A ring R is quasi-Frobenius if and only if every R -module embeds in a projective R -module.*

Here is one point of view on the significance of Theorem 3.2.3. It has been known for a long time, e.g. as described in [4], that when R is a quasi-Frobenius ring, there exists a model category structure on the category of R -modules in which the cofibrations are the injections and the weak equivalences are the stable equivalences of modules. Constructing this model category structure uses the quasi-Frobenius condition in an essential way. But a Waldhausen category structure on R -modules is weaker, less highly-structured, than a model category structure; so one might hope that, even in the absence of the quasi-Frobenius condition on R , one could put the structure of a Waldhausen category on R -modules, such that the cofibrations are the injections and the weak equivalences are the stable equivalences of modules. As a consequence of Theorem 3.2.3, one only gets a Waldhausen category structure with cylinder functor on the category of R -modules if R is quasi-Frobenius. So *the category of R -modules admits a model category structure as desired if and only if it admits a Waldhausen category structure with cylinder functor as desired.* (But our results, such as Corollary 3.2.4 on existence of the cylindrical Waldhausen category structure, also have the virtue of applying to quasi-Frobenius abelian categories that are not categories of modules over a ring.)

So one knows that, when R is a quasi-Frobenius ring, then one has the model category of R -modules with cofibrations inclusions and weak equivalences the stable equivalences, and from the theorems in this paper which we have described above, one knows that relaxing the condition that R be quasi-Frobenius does not enable one to get cylindrical Waldhausen category structures in any greater generality. Then one asks the natural question: restricting to the finitely-generated R -modules, *what are the K -groups of this Waldhausen category?* That leads us to the computations of Theorem 4.2.3 and Theorem 4.2.6.

We remark that our stable G -theory Waldhausen category poses an alternative to a construction by G. Garkusha in [3], who constructs a Waldhausen category which models the cofiber (on the spectrum level) of the Cartan map from K -theory to G -theory. Garkusha's construction is a Waldhausen category structure on chain complexes of R -modules, and when R is quasi-Frobenius, our Proposition 4.1.3 also describes the cofiber of the Cartan map but with a much smaller model than Garkusha's (the stable G -theory Waldhausen category structure on R -modules, rather than on chain complexes of R -modules). Our stable G -theory, as a model for the cofiber of the Cartan map, also has the advantage of multiplicative structure, as in Proposition 3.3.4.

We would not have written this paper or thought about any of these issues if not for conversations we had with Crichton Ogle, who taught us a great deal about Waldhausen K -theory during the summer of 2012. We are grateful to C. Ogle for his generosity in teaching us about this subject.

2. WALDHAUSEN CATEGORY STRUCTURES FROM ALLOWABLE CLASSES ON ABELIAN CATEGORIES.

2.1. Definitions. This subsection, mostly consisting of definitions, is entirely review and there are no new results or definitions in it, with the exception of Definition 2.1.8 and Definition-Proposition 2.1.10.

Throughout this subsection, let \mathcal{C} be an abelian category.

We begin with the definition of an allowable class. An allowable class is the structure one needs to specify on \mathcal{C} in order to have a notion of relative homological algebra in \mathcal{C} .

Definition 2.1.1. *An allowable class in \mathcal{C} consists of a collection E of short exact sequences in \mathcal{C} which is closed under isomorphism of short exact sequences and which contains every short exact sequence in which at least one object is the zero object of \mathcal{C} . (See section IX.4 of [5] for this definition and basic properties.)*

The usual "absolute" homological algebra in an abelian category \mathcal{C} is recovered by letting the allowable class E consist of *all* short exact sequences in \mathcal{C} .

Once one chooses an allowable class E , one has the notion of monomorphisms relative to E , or " E -monomorphisms," and epimorphisms relative to E , or " E -epimorphisms."

Definition 2.1.2. *Let E be an allowable class in \mathcal{C} . A monomorphism $f : M \rightarrow N$ in \mathcal{C} is called an E -monomorphism or an E -monic if the short exact sequence*

$$0 \rightarrow M \xrightarrow{f} N \rightarrow \operatorname{coker} f \rightarrow 0$$

is in E .

Dually, an epimorphism $g : M \rightarrow N$ is called an E -epimorphism or an E -epic if the short exact sequence

$$0 \rightarrow \ker f \rightarrow M \xrightarrow{f} N \rightarrow 0$$

is in E .

In the absolute case, the case that E is all short exact sequences in \mathcal{C} , the E -monomorphisms are simply the monomorphisms, and the E -epimorphisms are simply the epimorphisms.

Projective and injective objects are at the heart of homological algebra. In relative homological algebra, one has the notion of relative projectives, or E -projectives: these are simply the objects which lift over every E -epimorphism. The E -injectives are defined dually.

Definition 2.1.3. Let E be an allowable class in \mathcal{C} . An object X of \mathcal{C} is said to be an E -projective if, for every diagram

$$\begin{array}{ccc} & & X \\ & & \downarrow \\ M & \xrightarrow{f} & N \end{array}$$

in which f is an E -epic, there exists a morphism $X \rightarrow M$ making the above diagram commute.

Dually, an object X of \mathcal{C} is said to be an E -injective if, for every diagram

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \downarrow & & \\ X & & \end{array}$$

in which f is an E -monic, there exists a morphism $N \rightarrow X$ making the above diagram commute.

When the allowable class E is clear from context we sometimes refer to E -projectives and E -injectives as relative projectives and relative injectives, respectively.

In the absolute case, the case that E is all short exact sequences in \mathcal{C} , the E -projectives are simply the projectives, and the E -injectives are simply the injectives.

Once one has a notion of relative projectives, one has a reasonable notion of a stable equivalence or, loosely, a “homotopy” between maps, as studied (usually in the absolute case, where E -projectives are simply projectives) in stable representation theory:

Definition 2.1.4. Let E be an allowable class in \mathcal{C} . Let $f, g : M \rightarrow N$ be morphisms in \mathcal{C} . We say that f and g are E -stably equivalent and we write $f \simeq g$ if $f - g$ factors through an E -projective object of \mathcal{C} .

One then has the notion of stable equivalence of objects, or loosely, “homotopy equivalence”:

Definition 2.1.5. We say that a map $f : M \rightarrow N$ is a E -stable equivalence if there exists a map $h : N \rightarrow M$ such that $f \circ h \simeq \text{id}_N$ and $h \circ f \simeq \text{id}_M$.

In the absolute case where E consists of all short exact sequences in \mathcal{C} , this is the usual notion of stable equivalence of modules over a ring. Over a Hopf algebra over a field, stably equivalent modules have isomorphic cohomology in positive degrees, so if one is serious about computing the cohomology of all finitely-generated modules over a particular Hopf algebra—such as the Steenrod algebra or the group ring of a Morava stabilizer group—it is natural to first compute the representation ring modulo stable equivalence. See [7] for this useful perspective (which motivates much of the work in this paper).

We now define the relative-homological-algebraic generalizations of an abelian categories having enough projectives or enough injectives. We provide an extra twist on this definitions as well, which we will need for certain theorems: sometimes we will need to know that, for example, not only does every object embed in an injective, but that we can choose such embeddings in a functorial way.

Definition 2.1.6. Let E be an allowable class in \mathcal{C} . We say that \mathcal{C} has enough E -projectives if, for any object M of \mathcal{C} , there exists an E -epic $N \rightarrow M$ with N an E -projective. We say that \mathcal{C} has functorially enough E -projectives if \mathcal{C} has enough E -projectives and the choice

of E -epimorphisms from E -projectives to each object of \mathcal{C} can be made functorially, i.e., there exists a functor $P : \mathcal{C} \rightarrow \mathcal{C}$ together with a natural transformation $\epsilon : P \rightarrow \text{id}_{\mathcal{C}}$ such that $P(X)$ is E -projective and $\epsilon(X) : P(X) \rightarrow X$ is an E -epimorphism for all objects X of \mathcal{C} , and such that, if $f : X \rightarrow Y$ is an E -epimorphism, then so is $P(f) : P(X) \rightarrow P(Y)$.

Dually, we say that \mathcal{C} has enough E -injectives if, for any object M of \mathcal{C} , there exists an E -monic $M \rightarrow N$ with N an E -injective. We say that \mathcal{C} has functorially enough E -injectives if \mathcal{C} has enough E -injectives and the choice of E -monomorphisms from E -injectives to each object of \mathcal{C} can be made functorially, i.e., there exists a functor $I : \mathcal{C} \rightarrow \mathcal{C}$ together with a natural transformation $\eta : \text{id}_{\mathcal{C}} \rightarrow I$ such that $I(X)$ is E -injective and $\eta(X) : X \rightarrow I(X)$ is an E -monomorphism for all objects X of \mathcal{C} , and such that, if $f : X \rightarrow Y$ is an E -monomorphism, then so is $I(f) : I(X) \rightarrow I(Y)$.

Our need to have abelian categories with *functorially* enough injectives or projectives is only due to Waldhausen’s definitions of cylinder functors and resulting theorems, in [12], demanding that cylinder functors actually be *functors*. It seems likely that one can do away with this assumption and still prove analogues of Waldhausen’s results that use cylinders (e.g. the Fibration Theorem) by mimicking the situation in model category theory: there, one knows that any morphism has a factorization into a cofibration followed by an acyclic fibration, but such factorizations are not provided in a functorial way. We don’t pursue that angle in this paper, however.

Finally, we have our first definition of a quasi-Frobenius condition:

Definition 2.1.7. *Let E be an allowable class in \mathcal{C} . We will call E a quasi-Frobenius allowable class if the E -projectives are exactly the E -injectives. If the allowable class consisting of all short exact sequences in \mathcal{C} is a quasi-Frobenius class, then we will simply say that \mathcal{C} is quasi-Frobenius.*

Here are some important examples of allowable classes in abelian categories:

- As described above, the usual “absolute” homological algebra in an abelian category \mathcal{C} is recovered by letting the allowable class E consist of *all* short exact sequences in \mathcal{C} ; then the E -projectives are the usual projectives, etc. Note that, if E is an arbitrary allowable class in \mathcal{C} , then any projective object is an E -projective object, but the converse is not necessarily true.
- There is another “trivial” case of an allowable class: if we let E consist of only the short exact sequences in \mathcal{C} in which at least one of the objects is the zero object, then the E -epics consists of all identity maps as well as all projections to the zero object, and the E -monics consist of all identity maps as well as all inclusions of the zero object. As a consequence all objects are both E -injectives and E -projectives, and E is a quasi-Frobenius allowable class.
- Suppose \mathcal{C}, \mathcal{D} are abelian categories and $F : \mathcal{C} \rightarrow \mathcal{D}$ is an additive functor. Then we can let E be the allowable class in \mathcal{C} consisting of the short exact sequences which are sent by F to split short exact sequences in \mathcal{D} . If F has a left (resp. right) adjoint G then objects of \mathcal{C} of the form GFX (resp. FGX) are E -projectives (resp. E -injectives) and the counit map $GFX \rightarrow X$ of the comonad GF (resp. the unit map $X \rightarrow GFX$ of the monad GF) is an E -epic (resp. E -monic), hence \mathcal{C} has enough E -projectives (resp. enough E -injectives). These ideas are in [5].

For example, if R is a ring and \mathcal{C} the category of R -modules and \mathcal{D} the category of abelian groups, and F the forgetful functor, then E is the class of short exact sequences of R -modules whose underlying short exact sequences of abelian

groups are split. The R -modules of the form $R \otimes_{\mathbb{Z}} M$, for M an R -module, are E -projectives.

Here is a new definition which makes many arguments involving allowable classes substantially smoother:

Definition 2.1.8. *An allowable class E is said to have retractile monics if, whenever $g \circ f$ is an E -monic, f is also an E -monic.*

Dually, an allowable class E is said to have sectile epics if, whenever $g \circ f$ is an E -epic, g is also an E -epic.

The utility of the notion of “having sectile epics” comes from the following fundamental theorem of relative homological algebra, due to Heller (see [5]), whose statement is slightly cleaner is one is willing to use the phrase “having sectile epics.” The consequence of Heller’s theorem is that, in order to specify a “reasonable” allowable class in an abelian category, it suffices to specify the relative projective objects associated to it.

Theorem 2.1.9. *If C is an abelian category and E is an allowable class in C with sectile epics and enough E -projectives, then an epimorphism $M \rightarrow N$ in C is an E -epic if and only if the induced map $\text{hom}_C(P, M) \rightarrow \text{hom}_C(P, N)$ of abelian groups is an epimorphism for all E -projectives P .*

Heller’s theorem suggests the following construction, which as far as we know, is new (but unsurprising): if E is an allowable class, we can construct a “sectile closure of E ” which has the same relative projectives and the same stable equivalences but which has sectile epics. Here are the specific properties of this construction (we have neglected to write out proofs of these properties because the proofs are so elementary):

Definition-Proposition 2.1.10. *Let C be an abelian category, E an allowable class in C . Let E_{sc} be the allowable class in C consisting of the exact sequences*

$$X \rightarrow Y \rightarrow Y/X$$

such that the induced map

$$\text{hom}_C(P, Y) \rightarrow \text{hom}_C(P, Y/X)$$

is a surjection of abelian groups for every E -projective P . We call E_{sc} the sectile closure of E . The allowable class E_{sc} has the following properties:

- E_{sc} has sectile epics.
- An object of C is an E -projective if and only if it is an E_{sc} -projective.
- If f, g are two morphisms in C then f and g are E -stably equivalent if and only if they are E_{sc} -stably equivalent.
- If X, Y are two objects in C then X and Y are E -stably equivalent if and only if they are E_{sc} -stably equivalent.
- $(E_{sc})_{sc} = E_{sc}$.
- If E, F are two allowable classes in C and $F \subseteq E$ then $F_{sc} \subseteq E_{sc}$.

Of course there is a construction dual to the sectile closure, a *retractile closure*, with dual properties, but with a less straightforward relationship to stable equivalence, since stable equivalence is defined in terms of projectives, not injectives.

We now recall Waldhausen’s definitions:

Definition 2.1.11. *A pointed category C with finite pushouts equipped with a specified class of cofibrations and a specified class of weak equivalences, both closed under composition, is called a Waldhausen category if the following axioms are satisfied:*

- **(Cof 1.)** The isomorphisms in C are cofibrations.
- **(Cof 2.)** For every object X of C , the map $\text{pt.} \rightarrow C$ is a cofibration. (We write pt. for the zero object of C .)
- **(Cof 3.)** If $X \rightarrow Y$ is a morphism in C and $X \rightarrow Z$ is a cofibration, then the canonical map $Y \rightarrow Y \amalg_X Z$ is a cofibration.
- **(Weq 1.)** The isomorphisms in C are weak equivalences.
- **(Weq 2.)** If

(2.1.1)

$$\begin{array}{ccccc} Y & \longleftarrow & X & \longrightarrow & Z \\ \downarrow & & \downarrow & & \downarrow \\ Y' & \longleftarrow & X' & \longrightarrow & Z' \end{array}$$

is a commutative diagram in C in which the maps $X \rightarrow Y$ and $X' \rightarrow Y'$ are cofibrations and all three vertical maps are weak equivalences, then the induced map $Y \amalg_X Z \rightarrow Y' \amalg_{X'} Z'$ is a weak equivalence.

Ultimately, if C is a Waldhausen category, then what one typically wants to understand is $|\mathcal{wS}.C|$, the geometric realization of the simplicial category $\mathcal{wS}.C$ constructed by Waldhausen in [12]. The K -groups of C are defined as the homotopy groups of the loop space $\Omega|\mathcal{wS}.C|$:

$$\begin{aligned} \pi_{n+1}(|\mathcal{wS}.C|) &\cong \pi_n(\Omega|\mathcal{wS}.C|) \\ &\cong K_n(C). \end{aligned}$$

If C is a Waldhausen category then the following axioms may or may not be satisfied:

- Definition 2.1.12.**
- **(Saturation axiom.)** If f, g are composable maps in C and two of $f, g, g \circ f$ are weak equivalences then so is the third.
 - **(Extension axiom.)** If

$$\begin{array}{ccccc} X & \longrightarrow & Y & \longrightarrow & Y/X \\ \downarrow & & \downarrow & & \downarrow \\ X' & \longrightarrow & Y' & \longrightarrow & Y'/X' \end{array}$$

is a map of cofiber sequences and the maps $X \rightarrow X'$ and $Y/X \rightarrow Y'/X'$ are weak equivalences then so is the map $Y \rightarrow Y'$.

Definition 2.1.13. If C is a Waldhausen category, a cylinder functor on C is a functor from the category of arrows $f : X \rightarrow Y$ in C to the category of diagrams of the form

$$\begin{array}{ccccc} X & \xrightarrow{j_1} & T(f) & \xleftarrow{j_2} & Y \\ & \searrow f & \downarrow p & \nearrow \text{id} & \\ & & Y & & \end{array}$$

in C satisfying the three conditions:

- **(Cyl 1.)** If

$$\begin{array}{ccc} X' & \xrightarrow{f'} & Y' \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$$

is a commutative diagram in \mathcal{C} in which the vertical maps are weak equivalences (resp. cofibrations), then the map $T(f') \rightarrow T(f)$ is a weak equivalence (resp. cofibration and

$$X \coprod_{X'} T(f') \coprod_{Y'} Y \rightarrow T(f)$$

is a cofibration).

- **(Cyl 2.)** $T(\text{pt.} \rightarrow Y) = Y$ and the maps p and j_2 are the identity map on Y .
- **(Cyl 3.)** The map $j_1 \coprod j_2 : A \coprod B \rightarrow T(f)$ is a cofibration.

A Waldhausen category with cylinder functor may or may not satisfy the additional condition:

- **(Cylinder axiom.)** For any map f in \mathcal{C} , the map p is a weak equivalence.

The idea here is that a cylinder functor satisfying the cylinder axiom acts very much like the mapping cylinder construction from classical homotopy theory—or, more generally, like fibrant replacement in a model category. Some of Waldhausen’s most powerful results in [12] have proofs of a sufficiently homotopy-theoretic flavor that they require that every Waldhausen category in sight has a cylinder functor obeying the cylinder axioms. A good example of this is Waldhausen’s Fibration Theorem, which we now recall:

Theorem 2.1.14. Fibration Theorem (Waldhausen). *Suppose $\mathcal{C}, \mathcal{C}_0$ are Waldhausen categories with the same underlying category and the same underlying class of cofibrations. Suppose all of the following conditions are satisfied:*

- Every weak equivalence in \mathcal{C} is also a weak equivalence in \mathcal{C}_0 .
- \mathcal{C}_0 admits a cylinder functor satisfying the cylinder axiom.
- The weak equivalences in \mathcal{C}_0 satisfy the saturation and extension axioms.

Then

$$|wS.X| \rightarrow |wS.C| \rightarrow |wS.C_0|$$

is a homotopy fibre sequence, where X is the full sub-Waldhausen-category of \mathcal{C} generated by the objects that are weakly equivalent to pt. in \mathcal{C}_0 . As a consequence, after looping and taking homotopy groups, we get the long exact sequence of K -groups:

$$\cdots \rightarrow K_{n+1}(\mathcal{C}_0) \rightarrow K_n(X) \rightarrow K_n(\mathcal{C}) \rightarrow K_n(\mathcal{C}_0) \rightarrow K_{n-1}(X) \rightarrow \cdots$$

The question of when our Waldhausen categories given by allowable classes on abelian categories satisfy the required conditions for the Fibration Theorem to hold is the subject of most of this paper.

2.2. The Waldhausen category structure on an abelian category associated to a pair of allowable classes. In this subsection, we will prove that an abelian category equipped with a pair of allowable classes E, F admits a Waldhausen category structure in which the cofibrations are the F -monomorphisms and the weak equivalences are the E -stable equivalences. In this subsection we will also find conditions under which this Waldhausen category satisfies the extension and saturation axioms (see Definition 2.1.12 for definitions of these axioms). To get anywhere, we will need some lemmas:

Lemma 2.2.1. *A pullback of a surjective map of abelian groups is surjective.*

Proof. The forgetful functor from abelian groups to sets is a right adjoint, hence preserves limits. It also clearly preserves surjections. So the lemma is true if a pullback of a surjective maps of sets is surjective, which is an elementary exercise. \square

Lemma 2.2.2. *Let \mathcal{C} be an abelian category and let E be an allowable class with retractile monics. Then E -monics are closed under pushout in \mathcal{C} . That is, if $X \rightarrow Z$ is an E -monic and $X \rightarrow Y$ is any morphism in \mathcal{C} , then the canonical map $Y \rightarrow Y \amalg_X Z$ is an E -monic.*

Proof. Suppose $f : X \rightarrow Z$ is an E -monic and $X \rightarrow Y$ any morphism. We have the commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & X & \xrightarrow{f} & Z & \longrightarrow & \text{coker } f \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & Y & \longrightarrow & Y \amalg_X Z & \longrightarrow & \text{coker } f \longrightarrow 0 \end{array}$$

and hence, for every E -injective I , the induced commutative diagram of abelian groups

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{hom}_{\mathcal{C}}(\text{coker } f, I) & \longrightarrow & \text{hom}_{\mathcal{C}}(Z, I) & \longrightarrow & \text{hom}_{\mathcal{C}}(X, I) \longrightarrow 0 \\ & & \uparrow \cong & & \uparrow & & \uparrow \\ 0 & \longrightarrow & \text{hom}_{\mathcal{C}}(\text{coker } f, I) & \longrightarrow & \text{hom}_{\mathcal{C}}(Y \amalg_X Z, I) & \longrightarrow & \text{hom}_{\mathcal{C}}(Y, I). \end{array}$$

Exactness of the top row follows from f being an E -monic together with E having retractile monics, hence E is its own retractile closure, hence E -monics are precisely the maps which induce a surjection after applying $\text{hom}_{\mathcal{C}}(-, I)$ for every E -injective I . Now in particular we have a commutative square in the above commutative diagram:

$$\begin{array}{ccc} \text{hom}_{\mathcal{C}}(Z, I) & \longrightarrow & \text{hom}_{\mathcal{C}}(X, I) \\ \uparrow & & \uparrow \\ \text{hom}_{\mathcal{C}}(Y \amalg_X Z, I) & \longrightarrow & \text{hom}_{\mathcal{C}}(Y, I), \end{array}$$

which is a pullback square of abelian groups, by the universal property of the pushout. The top map in the square is a surjection, hence so is the bottom map, Lemma 2.2.1. So $\text{hom}_{\mathcal{C}}(Y \amalg_X Z, I) \rightarrow \text{hom}_{\mathcal{C}}(Y, I)$ is a surjection for every E -injective I . Again since E is its own retractile closure, this implies that $Y \rightarrow Y \amalg_X Z$ is an E -monic. \square

Lemma 2.2.3. *Suppose \mathcal{C} is an abelian category, E an allowable class in \mathcal{C} with retractile monics. Suppose \mathcal{C} has enough E -injectives. A composite of E -monomorphisms is an E -monomorphism.*

Proof. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be E -monomorphisms. Let I be an E -injective object equipped with a map $X \rightarrow I$. Then, since f is an E -monomorphism, $X \rightarrow I$ extends through f to a map $Y \rightarrow I$, which in turn extends through g since g is an E -monomorphism. So every map to an E -injective from Z extends through $g \circ f$. Now, by the dual of Heller's theorem 2.1.9, $g \circ f$ is an E -monomorphism. \square

Lemma 2.2.4. *Let \mathcal{C} be an abelian category and let E be an allowable class in \mathcal{C} . Then a composite of two E -stable equivalences in \mathcal{C} is an E -stable equivalence.*

Proof. Let $X \xrightarrow{f} Y \xrightarrow{g} Z$ be a pair of E -stable equivalences. Then there exist E -projective objects P_X, P_Y of \mathcal{C} and morphisms

$$\begin{array}{ccc} Y & \xrightarrow{f'} & X, \\ Z & \xrightarrow{g'} & Y, \\ X & \xrightarrow{i_X} & P_X, \\ Y & \xrightarrow{i_Y} & P_Y, \\ P_X & \xrightarrow{s_X} & X, \text{ and} \\ P_Y & \xrightarrow{s_Y} & Y \end{array}$$

such that

$$\begin{aligned} f' \circ f - \text{id}_X &= s_X \circ i_X \text{ and} \\ g' \circ g - \text{id}_Y &= s_Y \circ i_Y. \end{aligned}$$

Then we have

$$f' \circ g' \circ g \circ f - \text{id}_X = s_X \circ i_X + f' \circ s_Y \circ i_Y \circ f$$

so $f' \circ g' \circ g \circ f - \text{id}_X$ factors through the E -projective $P_X \oplus P_Y$. A similar argument applies to showing that $g \circ f \circ f' \circ g' - \text{id}_Z$ factors through an E -projective. So $g \circ f$ is an E -stable equivalence. \square

Definition-Proposition 2.2.5. *Let \mathcal{C} be an abelian category, let E, F be allowable classes in \mathcal{C} with $F \subseteq E$. Suppose each of the following conditions are satisfied:*

- F has retractile monics.
- E has retractile monics and sectile epics.
- \mathcal{C} has enough F -injectives.
- \mathcal{C} has enough E -projectives and enough E -injectives.
- Every E -projective object is E -injective.

Then there exists a Waldhausen category structure on \mathcal{C} in which the cofibrations are the F -monomorphisms and the weak equivalences are the E -stable equivalences. We write $\mathcal{C}_{E\text{-we}}^{F\text{-cof}}$ for this Waldhausen category. This Waldhausen category satisfies the saturation axiom and the extension axiom.

Proof. We check Waldhausen's axioms from Definition 2.1.11. In the case of an abelian category \mathcal{C} and allowable classes E, F with the stated classes of cofibrations and weak equivalences, axioms (Cof 1) and (Cof 2) and (Weq 1) are immediate. That the class of cofibrations is closed under composition is Lemma 2.2.3. That the class of weak equivalences is closed under composition is Lemma 2.2.4. We show that the remaining axioms are satisfied:

- Axiom (Cof 3) is a consequence of Lemma 2.2.2.
- Axiom (Weq 2) is actually fairly substantial and takes some work to prove—enough so that we moved this work into a paper of its own, [11]. In Corollary 4.4 of that paper, we prove that, if $E = F$, \mathcal{C} has enough E -projectives and enough E -injectives, E has sectile epics and retractile monics, and every E -projective object is E -injective, then \mathcal{C} satisfies axiom (Weq 2). We refer the reader to that paper for the proof, which requires some work and a number of preliminary results, and would make the present paper much longer if we included it here.

Once we have the result for $F = E$, it follows for $F \subseteq E$, since if diagram 2.1.1 has its horizontal maps F -monomorphisms, the horizontal maps are also then E -monomorphisms.

- The saturation axiom follows easily from Lemma 2.2.4 together with the observation that, if $X \xrightarrow{f} Y$ is a E -stable equivalence and $Y \xrightarrow{f'} X$ is a morphism such that $f \circ f' - \text{id}_Y$ and $f' \circ f - \text{id}_X$ both factor through E -projective objects, then f' is also a E -stable equivalence. That is, E -stable equivalences have “up-to-equivalence inverses.”
- Finally, we handle the extension axiom. We begin by assuming that $E = F$, and that

$$(2.2.1) \quad \begin{array}{ccccc} X & \longrightarrow & Y & \longrightarrow & Y/X \\ \downarrow f & & \downarrow g & & \downarrow h \\ X' & \longrightarrow & Y' & \longrightarrow & Y'/X' \end{array}$$

is a map of cofiber sequences (i.e., short exact sequences in E) and the maps $X \rightarrow X'$ and $Y/X \rightarrow Y'/X'$ are E -stable equivalences. Then, for any object M of C , we have the commutative diagram with exact columns

$$\begin{array}{ccc} \text{Ext}_{C/E}^1(X', M) & \xrightarrow{\cong} & \text{Ext}_{C/E}^1(X, M) \\ \downarrow & & \downarrow \\ \text{Ext}_{C/E}^2(Y'/X', M) & \xrightarrow{\cong} & \text{Ext}_{C/E}^2(Y/X, M) \\ \downarrow & & \downarrow \\ \text{Ext}_{C/E}^2(Y', M) & \longrightarrow & \text{Ext}_{C/E}^2(Y, M) \\ \downarrow & & \downarrow \\ \text{Ext}_{C/E}^2(X', M) & \xrightarrow{\cong} & \text{Ext}_{C/E}^2(X, M) \\ \downarrow & & \downarrow \\ \text{Ext}_{C/E}^3(Y'/X', M) & \xrightarrow{\cong} & \text{Ext}_{C/E}^3(Y/X, M). \end{array}$$

The horizontal maps marked as isomorphisms are isomorphisms because an E -stable equivalence $A \rightarrow B$ induces a natural equivalence of functors $\text{Ext}_{C/E}^i(B, -) \xrightarrow{\cong} \text{Ext}_{C/E}^i(A, -)$ for all $i \geq 1$; this is Lemma 3.6 of [11]. By the Five Lemma, we then have a natural isomorphism of functors $\text{Ext}_{C/E}^2(Y', -) \xrightarrow{\cong} \text{Ext}_{C/E}^2(Y, -)$. But since every E -projective is E -injective, this natural transformation being a natural isomorphism implies that $Y \rightarrow Y'$ is an E -stable equivalence; this is Lemma 4.2 of [11]. Hence the extension axiom is satisfied if $E = F$. Now if we do not have $E = F$ but instead $F \subseteq E$, then the extension axiom remains satisfied, as we have fewer diagrams to check the extension axiom for.

□

3. ABSOLUTE AND RELATIVE QUASI-FROBENIUS CONDITIONS.

3.1. Definitions. To our knowledge these definitions are all new. They are variants of the condition that every object embeds in a projective object, which Faith and Walker showed (see Theorem 1.0.1) to be equivalent, for categories of modules over a ring, to the ring being quasi-Frobenius.

Definition 3.1.1. • Let C be an abelian category, E, F a pair of allowable classes in C . We say that C is cone-Frobenius relative to E, F if, for any object X of C , there exists an F -monomorphism from X to an E -projective object of C .

We say that C is functorially cone-Frobenius relative to E, F if there exists a functor $J : C \rightarrow C$ and a natural transformation $\eta : \text{id}_C \rightarrow J$ such that:

- (1) $J(X)$ is E -projective for every object X of C ,
- (2) $\eta(X) : X \rightarrow J(X)$ is an F -monomorphism for every object X of C , and
- (3) if $f : X \rightarrow Y$ is an F -monomorphism then so is the map $J(f) : J(X) \rightarrow J(Y)$, and so is the universal map

$$(3.1.1) \quad Y \coprod_X J(X) \rightarrow J(Y).$$

We sometimes call the pair J, η a cone functor on C relative to E, F . When E, F are understood from context we simply call J, η a relative cone functor.

- **(The absolute case.)** If $E = F$ is the class of all short exact sequences in C and C is cone-Frobenius relative to E, F , then we simply say that C is cone-Frobenius.

If $E = F$ is the class of all short exact sequences in C and C is functorially cone-Frobenius relative to E, F , then we simply say that C is functorially cone-Frobenius.

The idea behind this definition is that the F -monomorphism and E -projective object together are a kind of “mapping cone,” in the sense of homotopy theory, on X : an embedding into a contractible object.

These definitions are very reasonable and in fact are very generally satisfied in the presence of much milder-looking quasi-Frobenius conditions on the abelian category. We present some theorems to that effect from our paper [10], using the following definition:

Definition 3.1.2. Let C be an abelian category, and let E be an allowable class in C . We will say that a set of objects $\{X_s\}_{s \in S}$ of C is an E -monic generating set for C if the following condition is satisfied:

- If $f : M \rightarrow N$ is a morphism in C such that $f \circ g \neq 0$ for every E -monomorphism $g : X_s \rightarrow M$, then f is E -monic.

If an E -monic generating set for C exists, then we say that C is E -monically small. If κ is a cardinal number and an E -monic generating set for C exists with $\leq \kappa$ objects in it, then we say that C is E -monically κ -small.

For example, we prove the following in [10]:

Proposition 3.1.3. Let R be a ring and let $\text{Mod}(R)$ be the category of left R -modules. Let E be the absolute allowable class on C , i.e., E is the class of all short exact sequences in C . Then $\text{Mod}(R)$ is E -monically κ -small, where κ is the number of left ideals of R .

Then we have the theorem, also from [10]:

Theorem 3.1.4. Suppose C is an abelian category, E, F allowable classes in C . Suppose F has retractile monics, and suppose that C is F -monically κ -small and C has all coproducts

of cardinality $\leq \kappa$. Suppose that every object of C embeds in an F -injective object by an F -monomorphism, i.e., for every object X there exists an F -monomorphism from X to an F -injective object. Suppose every F -injective object of C is E -projective. Then C is functorially cone-Frobenius relative to E, F .

As a special case:

Corollary 3.1.5. *Suppose C is an abelian category with enough injectives, with all small coproducts, and which is monically small. Suppose that every injective object in C is projective. Then C is functorially cone-Frobenius.*

As an even more special case:

Corollary 3.1.6. *Let R be a quasi-Frobenius ring. Then the category of R -modules is functorially cone-Frobenius.*

Finally, there is the question of when we have the functorial cone-Frobenius condition on the *finitely generated* modules over a ring. This is another result from [10]:

Corollary 3.1.7. *Let k be a finite field and let R be an quasi-Frobenius associative k -algebra which is finite-dimensional as a k -vector space. Then the category of finitely generated (left) R -modules is functorially cone-Frobenius.*

So the functorial cone-Frobenius condition seems to be a reasonable and often-satisfied one.

We also want to consider a relative form of the cone-Frobenius condition:

Definition 3.1.8. • *If C, C' are abelian categories and $i : C' \rightarrow C$ is an additive functor with right adjoint r , let E denote the allowable class of all short exact sequences in C and let E' the allowable class of all short exact sequences in C' . Then we say that C' is relatively quasi-Frobenius over C (resp. functorially relatively quasi-Frobenius over C) if C' is cone-Frobenius relative to i^*E, r_*E (resp. functorially cone-Frobenius relative to i^*E, r_*E).*

The idea here is that, in C' , any object embeds into some other object in such a way that, once one applies i , one gets a mapping cone (an E -monomorphic embedding into a contractible, i.e., E -projective, object) in C . If the reader is wondering whether such a thing is really more than a cone functor on C' or a cone functor on C , the reader might try letting C and C' be module categories over rings, and let i be the base-change/extension-of-scalars functor induced by a *surjection* of rings. The fact that ring surjections are rarely flat and hence that i is not generally left exact, i.e., i does not generally preserve monomorphisms, means that most attempts one might make to produce a mapping cone on C' do not give mapping cones on C after applying i . So the relative Frobenius conditions really are expressing something nontrivial.

Now the Faith-Walker Theorem, stated above as Theorem 1.0.1, can be restated using our definitions: *a ring R is quasi-Frobenius if and only if the category of R -modules is cone-Frobenius.*

3.2. Existence of cylinder functors. Now we show the equivalence of the functorial cone-Frobenius condition with the existence of cylinder functors satisfying the cylinder axioms.

First, we need a couple of lemmas:

Lemma 3.2.1. *Suppose C is an abelian category, E an allowable class in C with sectile epics. Suppose C has enough E -projectives. Then any finite direct sum of members of E is in E .*

Proof. Let I be a finite set and

$$0 \rightarrow X_i \rightarrow Y_i \rightarrow Z_i \rightarrow 0$$

be a member of E for every $i \in I$. Then, for any E -projective object P of C , we have the commutative diagram

$$\begin{array}{ccc} \mathrm{hom}_C(P, \oplus_i Y_i) & \longrightarrow & \mathrm{hom}_C(P, \oplus_i Z_i) \\ \downarrow \cong & & \downarrow \cong \\ \oplus_i \mathrm{hom}_C(P, Y_i) & \longrightarrow & \oplus_i \mathrm{hom}_C(P, Z_i). \end{array}$$

The bottom horizontal map is a surjection of abelian groups, so the top horizontal map is as well. Now by Heller's theorem 2.1.9, the map $\oplus_i Y_i \rightarrow \oplus_i Z_i$ is an E -epimorphism. So the short exact sequence

$$0 \rightarrow \oplus_i X_i \rightarrow \oplus_i Y_i \rightarrow \oplus_i Z_i \rightarrow 0$$

is in E . □

Lemma 3.2.2. (Shearing E -monics.) *Let C be an abelian category and let E be an allowable class in C . Suppose X, Y, Z are objects in C and suppose we have E -monomorphisms $e : X \rightarrow Y$ and $f : Z \rightarrow Y$. Let s be the morphism*

$$s : X \oplus Z \rightarrow Y \oplus Z$$

given by the matrix of maps

$$s = \begin{bmatrix} e & f \\ 0 & \mathrm{id}_Z \end{bmatrix}.$$

Then $\mathrm{coker} s$ is naturally isomorphic to $\mathrm{coker} e$. Furthermore, if C has enough E -injectives and E has retractile monics, then s is an E -monomorphism.

Proof. We first show that $\mathrm{coker} e \cong \mathrm{coker} s$. But this follows immediately from the commutative diagram with exact rows and exact columns:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & X & \xrightarrow{e} & Y & \longrightarrow & \mathrm{coker} e & \longrightarrow & 0 \\ \downarrow & & \downarrow i & & \downarrow i & & \downarrow & & \downarrow \\ 0 & \longrightarrow & X \oplus Z & \xrightarrow{s} & Y \oplus Z & \longrightarrow & \mathrm{coker} s & \longrightarrow & 0 \\ \downarrow & & \downarrow \pi & & \downarrow \pi & & \downarrow & & \downarrow \\ 0 & \longrightarrow & Z & \xrightarrow{\mathrm{id}} & Z & \longrightarrow & 0 & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 & \longrightarrow & 0 \end{array}$$

in which the maps marked π are projections to the second summand, and the maps marked i are inclusions as the first summand.

Now assume that E has retractile monics, and let $t : Y \oplus Z \rightarrow Y \oplus Y$ be the map given by the matrix of maps

$$t = \begin{bmatrix} \text{id}_Y & -f \\ 0 & f \end{bmatrix}.$$

Then a matrix multiplication reveals that the composite map $t \circ s : X \oplus Z \rightarrow Y \oplus Y$ is the direct sum map $e \oplus f$, a direct sum of E -monomorphisms, hence by Lemma 3.2.1, itself an E -monomorphism. (Note that, by taking the opposite category and noticing that the definition of an allowable class in an abelian category is self-dual, we get the conclusion of Lemma 3.2.1 if E has retractile monics and C has enough E -injectives.) Now since $t \circ s$ is an E -monomorphism and E is assumed to have retractile monics, s is also an E -monomorphism. \square

Theorem 3.2.3. *Let C, E, F be as in Proposition-Definition 2.2.5. Then the Waldhausen category C_{E-we}^{F-cof} admits a cylinder functor satisfying the cylinder axiom if and only if C is functorially cone-Frobenius relative to E, F .*

Proof. If C_{E-we}^{F-cof} has a cylinder functor satisfying the cylinder axiom, then the cylinder functor I sends, for any object X of C , the map $X \rightarrow \text{pt.}$ to the diagram

$$\begin{array}{ccc} X & \xrightarrow{F-cof} & I(X) \longleftarrow \text{pt.} \\ & \searrow & \downarrow E-we \\ & & \text{pt.} \end{array}$$

where the map marked $F - cof$ is an F -monic and the map marked $E - we$ is an E -stable equivalence. But for an object's map to the zero object to be an E -stable equivalence, this is equivalent to that object being an E -projective. Furthermore, if $f : X \rightarrow Y$ is any F -monomorphism in C , then we have the commutative diagram

$$\begin{array}{ccc} X \amalg \text{pt.} & \longrightarrow & I(X \amalg \text{pt.}) \\ \downarrow f \amalg \text{id}_{\text{pt.}} & & \downarrow I(f \amalg \text{id}_{\text{pt.}}) \\ Y \amalg \text{pt.} & \longrightarrow & I(Y \amalg \text{pt.}) \end{array}$$

and condition (Cyl 1) in the definition of a cylinder functor requires that $I(f \amalg \text{id}_{\text{pt.}})$ be an F -monomorphism as well. This completes one direction of the proof: $X \mapsto I(X \rightarrow \text{pt.})$ is a relative cone functor.

Now suppose we have a functor J and natural transformation e as in the definition of a relative cone functor in Definition 3.1.1. We claim that the functor sending any map $X \xrightarrow{f} Y$ to the diagram

$$\begin{array}{ccc} X & \xrightarrow{(f, e(X))} & Y \oplus J(X) \longleftarrow Y \\ & \searrow f & \downarrow \pi_Y \\ & & Y \end{array},$$

where π_Y is projection to Y , is a cylinder functor on C satisfying the cylinder axiom. We check Waldhausen's conditions from Definition 2.1.11. Condition (Cyl 2) is immediate, and the cylinder axiom follows from the projection $J(X) \oplus Y \rightarrow \text{pt.} \oplus Y \cong Y$ being a direct sum of E -stable equivalences, hence itself an E -stable equivalence.

We handle condition (Cyl 1) as follows: if $X' \rightarrow X$ and $Y' \rightarrow Y$ are F -monomorphisms, then the direct sum $J(X') \oplus Y' \rightarrow J(X) \oplus Y$ is an F -monomorphism by the assumption that J sends F -monomorphisms to F -monomorphisms, and

$$X \coprod_{X'} J(f') \coprod_{Y'} Y \rightarrow J(f)$$

being an F -monomorphism is exactly equivalent to the map 3.1.1 being an F -monomorphism, which is part of the definition of the functorial cone-Frobenius condition.

Now suppose that $X' \rightarrow X$ and $Y' \rightarrow Y$ are E -stable equivalences. We also note that, since $J(X')$, $J(X)$ are E -projective, the projections $J(X') \rightarrow \text{pt.}$ and $J(X) \rightarrow \text{pt.}$ are E -stable equivalences, and Lemma 2.2.4 gives us that the composite of either one with an E -stable inverse of the other is an E -stable equivalence between $J(X)$ and $J(X')$. So $J(X') \oplus Y' \rightarrow J(X) \oplus Y$ is an E -stable equivalence.

So condition (Cyl 1) holds.

Lemma 3.2.2 implies that the direct sum map

$$\begin{bmatrix} \text{id}_Y & f \\ 0 & e(X) \end{bmatrix} : Y \oplus X \rightarrow Y \oplus J(X)$$

is an E -monomorphism, i.e., a cofibration, so condition (Cyl 3) holds.

On the other hand, one easily verifies that, if C has a cylinder functor, then that cylinder functor is precisely the functor necessary for C to be functorially quasi-Frobenius category relative to E, F , from Definition 3.1.1. \square

Corollary 3.2.4. *Suppose C is a quasi-Frobenius abelian category with enough projectives and functorially enough injectives. Then C admits the structure of a Waldhausen category in which the cofibrations are the monomorphisms and the weak equivalences are the stable equivalences. Furthermore, this Waldhausen category satisfies the saturation and extension axioms, and it admits a cylinder functor satisfying the cylinder axiom. We call this Waldhausen category the stable G -theory of C , and we call its associated K -groups the stable G -theory groups of C .*

Corollary 3.2.5. *Suppose k is a finite field and R a quasi-Frobenius k -algebra which is finite-dimensional as a k -vector space. Then the category of finitely generated (left) R -modules admits the structure of a Waldhausen category in which the cofibrations are the monomorphisms and the weak equivalences are the stable equivalences. Furthermore, this Waldhausen category satisfies the saturation and extension axioms, and it admits a cylinder functor satisfying the cylinder axiom.*

3.3. Multiplicative structure. We recall Waldhausen's construction of multiplicative structures on the K -theory of Waldhausen categories:

Theorem 3.3.1. (Waldhausen.) *Suppose $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are Waldhausen categories and $F : \mathcal{A} \times \mathcal{B} \rightarrow \mathcal{C}$ is a functor satisfying each of the following conditions:*

- *If $M \rightarrow N$ is a cofibration in \mathcal{A} and L is an object of \mathcal{B} , then $F(M, L) \rightarrow F(N, L)$ is a cofibration in \mathcal{C} .*
- *If $M \rightarrow N$ is a cofibration in \mathcal{B} and L is an object of \mathcal{C} , then $F(L, M) \rightarrow F(L, N)$ is a cofibration in \mathcal{C} .*

- If $M' \rightarrow M$ and $N' \rightarrow N$ are cofibrations in \mathcal{A} and \mathcal{B} , respectively, then the universal map

$$F(M', N) \coprod_{F(M', N')} F(M, N') \rightarrow F(M, N)$$

is a cofibration in \mathcal{C} .

Then F induces a natural pairing

$$\Omega |wS.\mathcal{A}| \times \Omega |wS.\mathcal{B}| \rightarrow \Omega \Omega |wS.S.\mathcal{C}| \xrightarrow{\cong} \Omega |wS.\mathcal{C}|.$$

Waldhausen's pairing is sufficiently natural to imply the following (see [8] for a good expository account of infinite loop spaces with ring structure, but note that the statement we can make, below, is about *homotopy-commutative* ring spaces and spectra, but *not* E_∞ -ring spaces or spectra):

Corollary 3.3.2. *Suppose \mathcal{C} is a Waldhausen category which is equipped with a symmetric monoidal product $X, Y \mapsto X \otimes Y$ satisfying the conditions for the functor F in Theorem 3.3.1. Then $\Omega |wS.\mathcal{C}|$ is a homotopy-commutative ring space, that is, the infinite loop space $\Omega |wS.\mathcal{C}|$ is equipped with a homotopy-commutative product that is compatible with its loop space structure.*

Equivalently, when regarded as a spectrum using its infinite loop space structure, $\Omega |wS.\mathcal{C}|$ is a homotopy-commutative ring spectrum.

Lemma 3.3.3. *Let k be a field and let M', M, N', N be finite-dimensional k -vector spaces. Suppose we have monomorphisms $M' \rightarrow M$ and $N' \rightarrow N$ of k -vector spaces. Then the canonical map*

$$M' \otimes_k N \coprod_{M' \otimes_k N'} M \otimes_k N' \rightarrow M \otimes_k N$$

is a monomorphism.

Proof. We will regard M' and N' as subspaces of M and N , respectively. Choose a k -linear basis $\{m_1, \dots, m_i\}$ for M and a k -linear basis $\{n_1, \dots, n_j\}$ for N such that their restrictions to initial subsequences $\{m_1, \dots, m_{i'}\}$ and $\{n_1, \dots, n_{j'}\}$ form k -linear bases for M' and N' , respectively. Then we have the short exact sequence of k -vector spaces

$$0 \rightarrow M' \otimes_k N' \xrightarrow{f} M' \otimes_k N \oplus M \otimes_k N' \rightarrow M' \otimes_k N \coprod_{M' \otimes_k N'} M \otimes_k N' \rightarrow 0.$$

We write p for the composite map

$$(3.3.1) \quad M' \otimes_k N \oplus M \otimes_k N' \rightarrow M' \otimes_k N \coprod_{M' \otimes_k N'} M \otimes_k N' \rightarrow M \otimes_k N.$$

Suppose we have an element

$$x = \left(\sum_{a=1}^{i'} \sum_{b=1}^j \alpha_{a,b} m_a \otimes n_b, \sum_{a=1}^i \sum_{b=1}^{j'} \beta_{a,b} m_a \otimes n_b \right) \in M' \otimes_k N \oplus M \otimes_k N'$$

such that $p(x) = 0$. Then we have:

$$\begin{aligned} 0 &= \sum_{a=1}^{i'} \sum_{b=1}^j \alpha_{a,b} m_a \otimes n_b - \sum_{a=1}^i \sum_{b=1}^{j'} \beta_{a,b} m_a \otimes n_b \\ &= \sum_{a=1}^{i'} \sum_{b=1}^{j'} (\alpha_{a,b} - \beta_{a,b}) m_a \otimes n_b + \sum_{a=1}^{i'} \sum_{b=j'+1}^j \alpha_{a,b} m_a \otimes n_b + \sum_{a=i'+1}^i \sum_{b=1}^{j'} -\beta_{a,b} m_a \otimes n_b, \end{aligned}$$

hence $\beta_{a,b} = 0$ for all $a > i'$, and $\alpha_{a,b} = 0$ for all $b > j'$, and $\alpha_{a,b} = \beta_{a,b}$ if $a \leq i'$ and $b \leq j'$. These conditions together imply that x is in the subspace $M' \otimes_k N'$ of $M' \otimes_k N \oplus M \otimes_k N'$, hence the kernel of the composite map 3.3.1 is contained in $M' \otimes_k N'$, hence that the map from the quotient

$$(M' \otimes_k N \oplus M \otimes_k N') / (M' \otimes_k N') \cong M' \otimes_k N \coprod_{M' \otimes_k N'} M \otimes_k N'$$

to $M \otimes_k N$ is injective. \square

Proposition 3.3.4. *Let k be a field and let A be a co-commutative Hopf algebra over k which is finite-dimensional as a k -vector space. Let $\mathcal{G}_{st}(A)$ be the stable G -theory Waldhausen category of A from Corollary 3.2.4, that is, $\mathcal{G}_{st}(A)$ is the category of finitely generated (left) A -modules, with cofibrations the monomorphisms and weak equivalences the stable equivalences. Then the tensor product \otimes_k over k gives $\mathcal{G}_{st}(A)$ a symmetric monoidal product satisfying the assumptions of Corollary 3.3.2. Hence the stable G -theory spectrum $\Omega|_{\text{wS}} \mathcal{C}$ has the structure of a homotopy-commutative ring spectrum.*

Proof. Finite-dimensional Hopf algebras over fields are known to be quasi-Frobenius; see e.g. [7] for this fact. That the tensor product over k is symmetric monoidal follows from the Hopf algebra being co-commutative. That the tensor product over k satisfies the first two conditions of Definition 3.3.1 is because every A -module is flat as a k -module, since k is a field, so the tensor product over k preserves monomorphisms. That the tensor product over k satisfies the third condition is because of Lemma 3.3.3. \square

4. APPLICATIONS.

4.1. The relationship between stable G -theory and other G -theories and K -theories.

We recall that stable G -theory was defined in Corollary 3.2.4. It sits naturally in a diagram relating it to algebraic K -theory, algebraic G -theory, and the “derived representation groups,” but the relationship between these theories does not seem to be simple enough to permit easy computation of one from the others. We describe this relationship and provide a few comments in Remark 4.1.4 about the computational task of computing one of these theories once one has computed the others.

Now we define some notations we use to describe certain Waldhausen categories associated to an abelian category \mathcal{C} :

Definition 4.1.1. *Suppose \mathcal{C} is an abelian category. We will write:*

- $\mathcal{K}^\oplus(\mathcal{C})$ for the split K -theory category of \mathcal{C} , i.e., the Waldhausen category structure on the full subcategory generated by the projective objects of \mathcal{C} , where cofibrations are split inclusions and weak equivalences are isomorphisms.
- $\mathcal{K}(\mathcal{C})$ for the nonsplit K -theory category of \mathcal{C} , i.e., the Waldhausen category structure on the full subcategory generated by the projective objects of \mathcal{C} , where cofibrations are inclusions and weak equivalences are isomorphisms.
- $\mathcal{G}^\oplus(\mathcal{C})$ for the split G -theory category of \mathcal{C} , i.e., the Waldhausen category structure on \mathcal{C} where cofibrations are split inclusions and weak equivalences are isomorphisms.
- $\mathcal{G}(\mathcal{C})$ for the G -theory category of \mathcal{C} , i.e., the Waldhausen category structure on \mathcal{C} where cofibrations are inclusions and weak equivalences are isomorphisms.
- When it exists: $\mathcal{G}_{st}^\oplus(\mathcal{C})$ for the stable split G -theory category of \mathcal{C} , i.e., the Waldhausen category structure on \mathcal{C} where cofibrations are split inclusions and weak

equivalences are stable equivalences. (In the absolute sense, i.e., E -stable equivalences where E is the allowable class of all short exact sequences in \mathcal{C} .)

- When it exists: $\mathcal{G}_{st}(\mathcal{C})$ for the stable G -theory category of \mathcal{C} , i.e., the Waldhausen category structure on \mathcal{C} where cofibrations are inclusions and weak equivalences are stable equivalences. (In the absolute sense, i.e., E -stable equivalences where E is the allowable class of all short exact sequences in \mathcal{C} .)

Remark 4.1.2. We note that, if R is a ring and \mathcal{C} is the category of finitely generated R -modules, then

$$\pi_n \Omega |wS \cdot \mathcal{K}^\oplus(\mathcal{C})| \cong K_n(R),$$

i.e., the Waldhausen K -theory of the split K -theory Waldhausen category recovers the classical algebraic K -theory of R . Meanwhile,

$$\pi_n \Omega |wS \cdot \mathcal{G}(\mathcal{C})| \cong G_n(R),$$

i.e., the Waldhausen G -theory of the (nonsplit) G -theory Waldhausen category recovers the classical algebraic G -theory of R .

The other theories are more obscure but still meaningful. In degree zero,

$$\pi_0 \Omega |wS \cdot \mathcal{G}^\oplus(\mathcal{C})| \cong \text{Rep}(R),$$

the representation group (actually ring, under tensor product; but we have not discussed any multiplicative structures on our Waldhausen categories, which is another subject entirely) of R —that is, the Grothendieck group completion of the monoid of isomorphism classes of finitely-generated R -modules. So we sometimes regard the split G -theory as the “derived representation theory” and the groups $\pi_* \Omega |wS \cdot \mathcal{G}^\oplus(\mathcal{C})|$ as the “derived representation groups” of R .

In degree zero,

$$\pi_0 \Omega |wS \cdot \mathcal{G}_{st}^\oplus(\mathcal{C})| \cong \text{StableRep}(R),$$

the stable representation group of R —that is, the Grothendieck group completion of the monoid of stable equivalence classes of finitely-generated R -modules. So we sometimes regard the split stable G -theory as the “derived stable representation theory” and the groups $\pi_* \Omega |wS \cdot \mathcal{G}_{st}^\oplus(\mathcal{C})|$ as the “derived stable representation groups” of R .

Finally, the results of this paper are really about the stable G -theory groups

$$\pi_* \Omega |wS \cdot \mathcal{G}_{st}(\mathcal{C})| \cong (G_{st})_*(\mathcal{C}),$$

as defined in Corollary 3.2.4. In degree zero, (nonsplit) stable G -theory is

$$\pi_0 \Omega |wS \cdot \mathcal{G}_{st}(\mathcal{C})| \cong \text{StableRep}(R)/A,$$

the stable representation group modulo the subgroup A generated by all elements of the form $L - M + N$ where

$$0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$$

is a short exact sequence in \mathcal{C} .

Proposition 4.1.3. For any abelian category \mathcal{C} , we have a commutative diagram of topological spaces

$$(4.1.1) \quad \begin{array}{ccccc} |wS \cdot \mathcal{K}^\oplus(\mathcal{C})| & \longrightarrow & |wS \cdot \mathcal{G}^\oplus(\mathcal{C})| & \longrightarrow & |wS \cdot \mathcal{G}_{st}^\oplus(\mathcal{C})| \\ \downarrow & & \downarrow & & \downarrow \\ |wS \cdot \mathcal{K}(\mathcal{C})| & \longrightarrow & |wS \cdot \mathcal{G}(\mathcal{C})| & \longrightarrow & |wS \cdot \mathcal{G}_{st}(\mathcal{C})| \end{array}$$

in which the horizontal composites are nulhomotopic.

Suppose further that every projective object in \mathcal{C} is also injective. Then the map

$$|wS \cdot \mathcal{K}^\oplus(\mathcal{C})| \rightarrow |wS \cdot \mathcal{K}(\mathcal{C})|$$

in the above diagram is a homotopy equivalence.

Finally, suppose \mathcal{C} is functorially quasi-Frobenius, has enough projectives, has enough injectives, and every projective object is injective. Then the bottom row in diagram 4.1.1 is a homotopy fiber sequence.

Proof. First, when every projective object in \mathcal{C} is also injective, then any injective map between projective objects splits, by the universal property of an injective object; so injections and split injections coincide in the categories of projective objects in \mathcal{C} , so the functor $\mathcal{K}^\oplus(\mathcal{C}) \rightarrow \mathcal{K}(\mathcal{C})$ is an isomorphism of Waldhausen categories.

That the bottom row, under the stated assumptions, is a homotopy fiber sequence is an immediate consequence of Waldhausen's Fibration Theorem 2.1.14 and our Theorem 3.2.3.

Everything else here is a consequence of elementary facts from [12]. \square

Remark 4.1.4. There is a computational tool we would really like to have, but don't: we would very much like to be able to compute the homotopy groups of the fiber F^1 of the map

$$|wS \cdot \mathcal{G}_{st}^\oplus(\mathcal{C})| \rightarrow |wS \cdot \mathcal{G}_{st}(\mathcal{C})|.$$

From the associated long exact sequence induced by π_* as well as the identification we have of $(G_{st}^\oplus)_0$ and $(G_{st})_0$, we know (for example) that π_0 surjects on to the subgroup of the stable representation group $\text{StabRep}(\mathcal{C})$ generated by elements of the form $M_0 - M_1 + M_2$ for all short exact sequences $0 \rightarrow M_0 \rightarrow M_1 \rightarrow M_2 \rightarrow 0$ in \mathcal{C} , but it is hard to say much about $\pi_i(F^1)$ for $i > 0$. Since $(G_{st}^\oplus)_0(\mathcal{C})$ is the stable representation group of \mathcal{C} , one should think of $(G_{st}^\oplus)_i(\mathcal{C})$ as the *derived* stable representation groups of \mathcal{C} . An identification of $\pi_*(F^1)$ would allow one to pass from a knowledge of $(G_{st})_i(\mathcal{C})$, computable using our localization results in this paper, to a knowledge of the derived stable representation groups $(G_{st}^\oplus)_i(\mathcal{C})$. This would be valuable.

The problem of identifying F^1 is a special case of the following general problem: when one changes (expands or contracts) the class of cofibrations in a Waldhausen category \mathcal{C} , how does it change the K -theory groups $\pi_*(\Omega |wS \cdot \mathcal{C}|)$? We have an approach to this problem, especially in the situation here of identifying F^1 , in which we use a relative cell decomposition to describe the homology $H_n(|wS \cdot \mathcal{C}|, |wS \cdot \mathcal{C}^\oplus|)$ and then, making use of this homology computation and the relative Hurewicz map, we use an analogue of Serre's method of computing homotopy groups of spheres to work our way up the relative Postnikov tower of the map $|wS \cdot \mathcal{C}^\oplus| \rightarrow |wS \cdot \mathcal{C}|$. This technique is difficult and in any case is beyond the scope of the present paper. We hope to return to it in a later paper.

4.2. Applications to algebras. In this section we will finally apply our results to actual rings and algebras! We will frequently assume that the rings in question are quasi-Frobenius. For an explanation of why this implies the functorial cone-Frobenius condition, see Proposition 3.1.3, Theorem 3.1.4, and Cors. 3.1.5, 3.1.6, and 3.1.7. Throughout this section, whenever we assume that an algebra is over a *finite* field, the only reason we assume finiteness of the field is so that the above construction gives a cone functor on the finitely-generated module category; if one can extend this construction to finitely-generated modules over algebras over more general fields, then one can do away with the finiteness assumption on the field.

Proposition 4.2.1. (Stable G -theory is a delooping of relative algebraic K -theory.) *Suppose k is a finite field and A is a quasi-Frobenius k -algebra which is finite-dimensional as a k -vector space. Suppose $f : A \rightarrow k$ is a surjective morphism of k -algebras with nilpotent kernel. Write $K(A \rightarrow k)$ for the relative K -theory space of f , that is, the fiber of the map*

$$\Omega |wS \cdot \mathcal{K}(A)| \rightarrow \Omega |wS \cdot \mathcal{K}(k)|.$$

Then we have a homotopy equivalence $K(A \rightarrow k) \simeq \Omega^2 |G_{st}(A)|$, hence

$$(G_{st})_n(A) \cong K_{n-1}(A \rightarrow k)$$

for all $n > 0$.

Proof. We have the commutative diagram of (pointed) spaces

$$(4.2.1) \quad \begin{array}{ccccc} K(A \rightarrow k) & \longrightarrow & \Omega |wS \cdot \mathcal{K}(A)| & \longrightarrow & \Omega |wS \cdot \mathcal{K}(k)| \\ \downarrow & & \downarrow & & \downarrow \\ G(A \rightarrow k) & \longrightarrow & \Omega |wS \cdot \mathcal{G}(A)| & \longrightarrow & \Omega |wS \cdot \mathcal{G}(k)| \\ \downarrow & & \downarrow & & \downarrow \\ \Omega |wS \cdot \mathcal{G}_{st}(A)| & \longrightarrow & \Omega |wS \cdot \mathcal{G}_{st}(A)| & \longrightarrow & \text{pt.} \end{array}$$

in which the spaces $K(A \rightarrow k)$ and $G(A \rightarrow k)$ are defined to be the homotopy fibers of the evident maps, so that the rows in diagram 4.2.1 are homotopy fiber sequences.

We note that the middle and right-hand columns in diagram 4.2.1 are also homotopy fiber sequences, by Proposition 4.1.3. Since all rows and the rightmost two columns in diagram 4.2.1 are homotopy fiber sequences, so is the left-hand column. By Quillen's filtration argument in [9], one also knows that any nilpotent extension of algebras over a field induces a homotopy equivalence in G -theory spaces, so the map $\Omega |wS \cdot \mathcal{G}(A)| \rightarrow \Omega |wS \cdot \mathcal{G}(k)|$ is a homotopy equivalence. So its fiber $G(A \rightarrow k)$ is contractible. Hence the left-hand column of diagram 4.2.1 reads, up to homotopy equivalence,

$$K(A \rightarrow k) \rightarrow \text{pt.} \rightarrow \Omega |wS \cdot \mathcal{G}_{st}(A)|.$$

Hence $K(A \rightarrow k) \simeq \Omega^2 |wS \cdot \mathcal{G}_{st}(A)|$. \square

Now we recall Gabber's rigidity theorem, from e.g. Chapter 4 of [13]:

Theorem 4.2.2. (Gabber.) *Suppose A is a commutative ring and (A, I) is a Hensel pair. Suppose $1/i \in A$. Then, for all $n > 0$, the map*

$$K_n(A; \mathbb{Z}/i\mathbb{Z}) \rightarrow K_n(A/I; \mathbb{Z}/i\mathbb{Z})$$

is an isomorphism.

Theorem 4.2.3. *Suppose k is a finite field of characteristic p and A is a quasi-Frobenius commutative k -algebra which is finite-dimensional as a k -vector space. Suppose $f : A \rightarrow k$ is a surjective morphism of k -algebras with nilpotent kernel. Then we have the following computation of the stable G -theory groups of A :*

- $(G_{st})_0(A)$ is the abelian group whose objects are the stable equivalence classes of finitely-generated A -modules, modulo the relation $[L] + [N] = [M]$ if there exists a short exact sequence $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$.
- For $n > 0$, $(G_{st})_n(A)$ is uniquely ℓ -divisible for all primes $\ell \neq p$. (So $(G_{st})_n(A)$ has no summands of the form $\mathbb{Z}/\ell^i\mathbb{Z}$ or \mathbb{Z}_ℓ or $(\mathbb{Q}/\mathbb{Z})_\ell$, but it may have summands of the form \mathbb{Q} , for example.)

- For $n > 0$, the modulo p^i stable G -theory $(G_{st})_n(A; \mathbb{Z}/p^i\mathbb{Z})$ is isomorphic to the topological cyclic homology $TC_{n-1}(A; \mathbb{Z}/p^i\mathbb{Z})$.

Proof. • The statement about $(G_{st})_0(A)$ is the usual identification of K_0 of a Waldhausen category; see Remark 4.1.2.

- Suppose ℓ is a prime distinct from p . From Gabber rigidity, Theorem 4.2.2, we know that $K_n(A \rightarrow k; \mathbb{Z}/\ell^i\mathbb{Z}) \cong 0$ for $n \geq 1$. Since A is local, we also know that the map $K_0(A) \rightarrow K_0(k)$ is an isomorphism (the identity map from \mathbb{Z} to \mathbb{Z}), so $K_0(A \rightarrow k; \mathbb{Z}/\ell^i\mathbb{Z})$ vanishes as well. From the short exact sequence

$$0 \rightarrow K_n(A \rightarrow k)/\ell^i \rightarrow K_n(A \rightarrow k; \mathbb{Z}/\ell^i\mathbb{Z}) \rightarrow \ell^i \text{ torsion in } K_{n-1}(A \rightarrow k) \rightarrow 0$$

and vanishing of the middle term for $n > 0$, we then know that $K_n(A \rightarrow k)/\ell^i \cong 0$ for $n > 0$, and $K_n(A \rightarrow k)$ is ℓ^i -torsion-free for $n \geq 0$. So, in particular, $K_n(A \rightarrow k)$ is uniquely ℓ -divisible for all $n \geq 0$. By Proposition 4.2.1, we know that $(G_{st})_n(A) \cong K_{n-1}(A \rightarrow k)$ for $n \geq 1$, so for $n \geq 1$ we conclude that $(G_{st})_n(A)$ is uniquely ℓ -divisible.

- McCarthy's theorem (see e.g. Madsen's article [6] for a good expository account) implies that $K_n(A \rightarrow k; \mathbb{Z}/p^i\mathbb{Z}) \cong TC_n(A \rightarrow k; \mathbb{Z}/p^i\mathbb{Z})$. Now use Proposition 4.2.1. \square

One of our results (in Theorem 4.2.6, below) involves proving that certain stable G -theory spectra are complex oriented. To prepare for that result, we provide the definitions of a complex orientation.

Definition 4.2.4. *Let E be a homotopy-commutative ring spectrum. By a complex orientation on E we mean a choice of element $\chi \in \tilde{E}^2(BU(1))$ with the property that χ maps to the multiplicative unit element $1 \in E^0(\text{pt.})$ under the composite*

$$\tilde{E}^2(BU(1)) \rightarrow \tilde{E}^2(S^2) \xrightarrow{\cong} \tilde{E}^0(S^0) \xrightarrow{\cong} E^0(\text{pt.}),$$

where the left-hand map is the map induced in E -cohomology by inclusion of the 2-skeleton S^2 of $BU(1)$, the classifying space of complex line bundles.

If E admits a complex orientation, we sometimes say that E is complex oriented.

Complex orientations are important for geometric purposes, since the element χ behaves essentially like a first Chern class, allowing one to carry out geometric arguments using E -cohomology that require characteristic classes for line bundles. Complex orientations also connect powerfully to number theory, via formal group laws: a complex orientation on a homotopy-commutative ring spectrum E gives rise to a one-dimensional commutative formal group law on $\pi_*(E)$, and much of the homotopy theory of complex oriented ring spectra can be described completely in terms of the moduli theory of one-dimensional formal groups. Adams's book [1] is an excellent reference for this material.

Definition 4.2.5. *Suppose k is a finite field and A is a quasi-Frobenius k -algebra which is finite-dimensional as a k -vector space. We will write $g_{st}(A)$ for the connective stable G -theory spectrum of A , that is, the connective cover of the spectrum $\Omega|_{wS} \mathcal{G}_{st}(A)$. We will write $(g_{st})_i(A)$ for the i th homotopy group $\pi_i(g_{st}(A))$.*

If A is also a co-commutative Hopf algebra over k , then by Proposition 3.3.4 and the fact that connective covers of homotopy-commutative ring spectra are also homotopy-commutative ring spectra, $g_{st}(A)$ is a homotopy-commutative ring spectrum.

A note for the reader who is not used to connective covers: the essential property of the connective cover x of a spectrum X is that $\pi_i x \cong \pi_i X$ for $i \geq 0$, and $\pi_i x \cong 0$ for $i < 0$.

Now one wants to use Proposition 3.3.4 to get a multiplicative version of Theorem 4.2.3. So we have the following, which also uses Hesselholt-Madsen's computation of the topological cyclic homology of truncated polynomial rings (see e.g. the survey article [6]):

Theorem 4.2.6. *Suppose k is a finite field of characteristic p . Let $A \cong k[x]/x^{p^n}$ for some positive integer n . Then we have the following results:*

- $(g_{st})_0(A) \cong \mathbb{Z}/p^n\mathbb{Z}$.
- For all positive integers m , $(g_{st})_{2m}(A) \cong TC_{2m-1}(A; \hat{\mathbb{Z}}_p) \cong \mathbb{W}_{mp^{n-1}}(k)/V_{p^n}\mathbb{W}_{m-1}(k)$, where $\mathbb{W}_i(k)$ is the standard filtration quotient i of the ring of big Witt vectors, that is,

$$\mathbb{W}_i(k) = (1 + Xk[[X]])^\times / (1 + X^{m+1}k[[X]])^\times,$$

and V_j is the Verschiebung morphism sending a power series $f(X)$ to the power series $f(X^j)$.

- For all positive integers m , $(g_{st})_{2m-1}(A) \cong 0$.
- $g_{st}(A)$ is a complex oriented ring spectrum.

Proof. Throughout, we implicitly use the fact that $(G_{st})_i(A) \cong (g_{st})_i(A)$ for $i \geq 0$.

- To get the isomorphism $(G_{st})_0(A) \cong \mathbb{Z}/p^n\mathbb{Z}$ we use the identification of degree zero stable G -theory, in Remark 4.1.2, as the group of stable equivalence classes of finitely generated A -modules, modulo the relation splitting all short exact sequences. Now A is a principal ideal domain, hence its finitely-generated module category is Krull-Schmidt and every indecomposable finitely-generated module is cyclic, and for every $i \leq p^n$, we have the short exact sequence

$$0 \rightarrow k \rightarrow k[x]/x^i \rightarrow k[x]/x^{i-1} \rightarrow 0$$

of A -modules, so we have the relation $[k] + [k[x]/x^{i-1}] = [k[x]/x^i]$ in $(G_{st})_0(A)$, so $[k]$ generates $(G_{st})_0(A)$. Finally, we observe that $[k[x]/x^{p^n}] = p^n[k]$ is zero in stable G -theory. So $(G_{st})_0(A) \cong \mathbb{Z}/p^n\mathbb{Z}$, generated by the stable equivalence class of the A -module k .

- Note that we have a description of the stable G -theory of A in Theorem 4.2.3 up to uniquely ℓ -divisible summands. But, by Proposition 3.3.4, the spectrum of the infinite loop space $\Omega|_{wS} \mathcal{G}_{st}(A)|$ is a homotopy-commutative ring spectrum. So each homotopy group $\pi_i \Omega|_{wS} \mathcal{G}_{st}(A)| \cong (G_{st})_i(A)$ is a module over $\pi_0 \Omega|_{wS} \mathcal{G}_{st}(A)| \cong (G_{st})_0(A) \cong \mathbb{Z}/p^n\mathbb{Z}$. The only uniquely ℓ -divisible group admitting the structure of a $\mathbb{Z}/p^n\mathbb{Z}$ -module is the trivial group. So the uniquely divisible factors whose existence is not ruled out by Theorem 4.2.3 are indeed ruled out in this case, by the multiplicative structure.

The computation of $TC_*(A; \hat{\mathbb{Z}}_p)$ is due to Hesselholt and Madsen, as in [6].

- That $(G_{st})_*(A)$ vanishes in odd positive degrees is similar to the computation in even degrees. Theorem 4.2.3 guarantees that, up to uniquely ℓ -divisible summands, $(G_{st})_*(A)$ agrees with topological cyclic homology with a degree shift, and Hesselholt and Madsen compute that the topological cyclic homology of A vanishes in even degrees. So $(G_{st})_*(A)$ is uniquely ℓ -divisible in odd positive degrees. But we know that $(G_{st})_0(A) \cong \mathbb{Z}/p^n\mathbb{Z}$, and as a consequence of Proposition 3.3.4, we know that $(G_{st})_i(A)$ is a $(G_{st})_0(A)$ -module for all i . So the uniquely ℓ -divisible summands must be trivial. So $(G_{st})_i(A)$ is trivial for odd positive i .

- We have the inclusion of the 2-skeleton $S^2 \hookrightarrow BU(1)$ into the classifying space of complex line bundles, and the map it induces of $g_{st}(A)$ -cohomological Atiyah-Hirzebruch spectral sequences:

$$\begin{array}{ccc}
 H^p(BU(1); (g_{st}(A))^q) & \Longrightarrow & g_{st}(A)^{p+q}(BU(1)) \\
 \downarrow & & \downarrow \\
 H^p(S^2; (g_{st}(A))^q) & \Longrightarrow & g_{st}(A)^{p+q}(S^2).
 \end{array}$$

Since the homotopy groups of $g_{st}(A)$ and the cohomology groups of $BU(1)$ are both concentrated in even degrees, there is no room for differentials in the spectral sequence for $BU(1)$. We conclude that the map of spectral sequences is simply projection on to the $p = 0$ and $p = 2$ columns, and that in particular, every element in $H^2(S^2; (g_{st}(A))^0) \cong g_{st}(A)^0(\text{pt.})$ —in particular, the multiplicative unit element—is in the image of the projection from $H^2(BU(1); (g_{st}(A))^0) \cong g_{st}(A)^2(BU(1))$. So $g_{st}(A)$ is complex oriented. \square

REFERENCES

- [1] J. F. Adams. *Stable homotopy and generalised homology*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1995. Reprint of the 1974 original.
- [2] Carl Faith. Embedding modules in projectives: A report on a problem. In *Advances in Non-Commutative Ring Theory*, volume 951, pages 21–40. Lecture Notes in Mathematics, 1982.
- [3] Grigory Garkusha. On the homotopy cofibre spectrum of $K(R) \rightarrow G(R)$. *J. Algebra Appl.*, 1(3):327–334, 2002.
- [4] Mark Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [5] Saunders Mac Lane. *Homology*. Classics in Mathematics. Springer-Verlag, Berlin, 1995. Reprint of the 1975 edition.
- [6] Ib Madsen. Algebraic K -theory and traces. In *Current developments in mathematics, 1995 (Cambridge, MA)*, pages 191–321. Int. Press, Cambridge, MA, 1994.
- [7] H. R. Margolis. *Spectra and the Steenrod algebra*, volume 29 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1983. Modules over the Steenrod algebra and the stable homotopy category.
- [8] J. P. May. What precisely are E_∞ ring spaces and E_∞ ring spectra? In *New topological contexts for Galois theory and algebraic geometry (BIRS 2008)*, volume 16 of *Geom. Topol. Monogr.*, pages 215–282. Geom. Topol. Publ., Coventry, 2009.
- [9] Daniel Quillen. Higher algebraic K -theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973.
- [10] Andrew Salch. A functorial construction of injective resolutions with applications in algebra and homotopy theory. *in preparation*.
- [11] Andrew Salch. Homotopy colimits in stable representation theory. *available on arXiv*.
- [12] Friedhelm Waldhausen. Algebraic K -theory of spaces. In *Algebraic and geometric topology (New Brunswick, N.J., 1983)*, volume 1126 of *Lecture Notes in Math.*, pages 318–419. Springer, Berlin, 1985.
- [13] Charles A. Weibel. *The K-book*, volume 145 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K -theory.