

Purely helical absolute equilibria and chirality of (magneto)fluid turbulence

Jian-Zhou Zhu, Weihong Yang² and Guang-Yu Zhu^{1,3}

¹WCI Center for Fusion Theory, National Fusion Research Institute, 169-148 Gwahak-ro, Daejeon, Korea

²Department of Modern Physics, University of Science and Technology of China, 230026 Hefei, China

³Life and Chinese Medicine Study Center, 47 Bayi Cun, 366025 Yong'an, Fujian, China

(Received ?; revised ?; accepted ?. - To be entered by editorial office)

Purely helical absolute equilibria of incompressible neutral fluids and plasmas (electron, single-fluid and two-fluid magnetohydrodynamics) are systematically studied with the help of helical (wave) representation and truncation, for genericities and specificities about helicity. A unique chirality selection and amplification mechanism and relevant insights, such as the one-chiral-sector-dominated states, among others, about (magneto)fluid turbulence follow.

Key words:

1. Background, technique, and basic ideas

Helical modes are basic in electromagneto- and hydrodynamics (see, *e.g.*, Moses 1971). They are left- or right-handed, signaturing *chirality*[†] which may be quantified by helicity and its relevant derivatives, such as the relative helicity, important for the statistical dynamics.

1.1. Helical turbulence and absolute equilibrium

Recognizing the importance of helicity in hydrodynamic turbulence is relatively new, though Helmholtz-Kelvin theorem is old (Moffatt 2008). Indeed, in a communication with C.-C. Lin in 1945, L. Onsager noticed that the coefficients of the Fourier modes of hydrodynamic velocity field are “‘momentoids’ in the sense of Boltzmann, and the theorem of equipartition would apply if their number were finite. Since this is not the case, we get a ‘violet catastrophe’ instead.”[‡] Statistical absolute equilibrium (AE) energy equipartition among each Fourier modes was later explicitly formulated by T.-D. Lee (1952) for both pure hydrodynamics (HD) and magneto-hydrodynamics (MHD). Neither Onsager nor Lee (who, interestingly, as well-known, however soon suggested with Yang in 1956 the chiral “world” — parity is violated in the weak interactions!) considered the invariance of helicity[¶] which makes the flow field lose mirror symmetry and which can also be involved in the generalized equipartitions (Kraichnan 1973; Frisch et al. 1975). Now, tremendous progresses with helical representation/decomposition have been made (see, *e.g.*, Yang, Su & Wu 2010, and references therein): Nature of the triadic interactions can be exposed more clearly (Waleffe 1992) and be exploited to understand better the fluctuations, for instance, of electron MHD (EMHD) and Hall MHD (Galtier & Bhattacharjee 2003; Galtier

[†] This notion is widely used in chemistry, physics and (origin of) life sciences and was called *dissymmetry*, which is still occasionally used, before Kelvin (1904) and various attempts have been made to mathematically quantify it (see, *e.g.*, the review by Petitjean 2003).

[‡] This remarkable comment adds more to Onsager’s ignored legacy on hydrodynamics than that exposed by Eyink & Sreenivasan (2006, private communication in 2008) who reproduced the letter.

[¶] Later, Betchov (1961) first tried to explore invariant helicity’s role in turbulence, contrasting a box of nails to screws with reflexion asymmetry.

2006). And, spectral dynamics can be diagnosed in a finer way (Chen, Chen & Eyink 2003, see more detailed discussions in §2.2). In particular, for the pure-helical-mode subsystem with only modes of one chiral sector of Navier-Stokes equations, one can expect a dual cascade picture. Indeed, we will see that Kraichnan (1973)'s helical spectra can be refined to allow a negative temperature state to support the dual cascades. On the other hand, approaching nearly maximally helical, *i.e.*, one-chiral-sector-dominated states (OCSDSs) with severe *chiral symmetry* breaking, *i.e.*, imbalance of positive and negative helicity, along scales have been explicitly demonstrated (*e.g.*, Meneguzzi, Frisch & Pouquet 1981; Brandenburg, Dobler & Subramanian 2002, see more detailed discussions in §2.3.1) but want a corresponding theoretical understanding, as we will offer. Recently, Meyrand & Galtier (2012) studied Hall MHD new chirality symmetry breaking in the sense of domination by whistler or ion-cyclotron waves defined by the linear wave dispersion relation, which is different to the chirality signature coming from the helical representation used in this paper; see §2.3.2 for more remarks.

The equilibrium-statistical-mechanics approach to investigate turbulence had been somewhat esoteric, but Kraichnan (1967, 1973, hereafter K67 and K73) established in a more explicit and complete way the AE for both 2D and 3D incompressible HD. Fourier modes beyond $[k_{min}, k_{max}]$ being discarded (Galerkin truncation), certain rugged quadratic invariants — for solutions regular enough to bear no dissipative anomaly (see, *e.g.*, Eyink 2008) — such as the kinetic energy (\mathcal{E}_K) and enstrophy (for 2D) or helicity (\mathcal{H}_K for 3D), are still conserved. With the constraints of these rugged invariants, Kraichnan obtained the respective energy spectral densities for 2D and 3D:

$$\bar{U}_K(\mathbf{k}) = 1/(\alpha + \beta k^2) \text{ and } U_K(\mathbf{k}) = 2\alpha/(\alpha^2 - \beta^2 k^2)$$

respectively. For symbolic convenience, from now on the vector argument \mathbf{k} will be replaced with its module k by isotropicity consideration. $U_K(k)$ for 3D, for instance, can be derived immediately from the Gibbs distribution

$$\sim \exp\{-(\alpha\mathcal{E}_K + \beta\mathcal{H}_K)\},$$

where α is the temperature parameter associated with energy and β with helicity (enstrophy in 2D).[†] K67 showed that low enstrophy state in 2D corresponds to a negative α , indicating condensation of energy at smallest k with a roughly (smoothed) \perp shape spectral density: *c.f.*, similar MHD figures in Frisch et al. (1975). In 3D, there is no such negative temperature state ($\alpha > 0$ due to the *realizability condition* from the positive definiteness of the quadratic form $\alpha\mathcal{E}_K + \beta\mathcal{H}_K$) but only \lrcorner shape spectral density, and low helicity state corresponds to vanishing β and equipartition of energy. By statistical consideration of the tendency of the interacting modes to relax towards the equilibrium state, inverse energy cascade was then argued for 2D but disputed for 3D. Note also that, as argued by L'vov et al. (2002) for 2D turbulence, in some

[†] We adopt K73 notations and definitions: $\mathcal{E}_K = \sum_{\mathbf{k}} U_K(k) \rightarrow \int dk 4\pi E_K(k)$ and $\mathcal{H}_K = \sum_{\mathbf{k}} Q_K(k) \rightarrow \int dk 4\pi H_K(k)$ in the continuous- k limit, where \bullet implies restricting to the subset of surviving modes and $E(k) = k^2 U_K(k)$ and $H(k) = k^2 Q_K(k)$ are the 1D spectra. We will always use α for energy related temperature parameter and β for helicity. Self-evident indexes, such as M for “magnetic”, when necessary for discrimination, will be added to β, U, Q, \mathcal{E} and \mathcal{H} etc. And, for simplicity we will always use Gibbs ensemble calculation and will not repeatedly formulate and explain it. The general results of this paper are not affected by the differences between an infinite domain and a finite cyclic box, so we may switch between these two descriptions, depending on which one is more convenient. Difference of a factor of 2 may arise, depending on how one treats the realizability condition (see below) and the invariant(s) (summation over the whole or half of the wavenumber domain etc.), and yet another freedom about the sign of helicity is of one's free choice. Also, spectra in this paper may be obtained in other approaches, such as finding the stationary solution of the master equation with the properties of vertices relevant to the conservation laws, which may avoid explicitly resorting to the Gibbs distribution (Private communication with E. A. Kuznetsov).

situations turbulence is actually not far from absolute equilibrium. One may even imagine a two-step scenario that the system first reached equilibrium which was then broken with the quantity, say, energy, being removed from the scales of concentration. Such a thought experiment makes good sense when the thermalization eddy turn over time scale is reasonably smaller than that of the equilibrium-breaking mechanism, such as that due to (hypo)viscosity. In modern terminology, we may say that the system is driven by the *entropic force* towards the maximum entropy state.

1.2. Helical (wave/mode) representation

For a 3D transverse vector field, such as the velocity \mathbf{u} , vorticity $\boldsymbol{\omega} = \nabla \times \mathbf{u}$ and the transverse component of vector potential \mathbf{A} of magnetic field \mathbf{B}^\dagger etc., in a cyclic box of volume, say, $\mathcal{V} = (2\pi)^3$, the helical mode/wave representation in Fourier space reads (Moses 1971)

$$\mathbf{v} = \sum_{\mathbf{k}} \hat{\mathbf{v}}(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}} = \sum_c \mathbf{v}^c = \sum_{\mathbf{k},c} \hat{\mathbf{v}}^c(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}} = \sum_{\mathbf{k},c} \hat{\mathbf{v}}^c(\mathbf{k}) \hat{\mathbf{h}}_c(\mathbf{k}) e^{i\mathbf{k}\cdot\mathbf{r}}. \quad (1.1)$$

Here $\hat{i}^2 = -1$ and $c^2 = 1$ for the chirality indexes $c = \text{“+”}$ or “-” . The helical mode bases (complex eigenvectors of the curl operator) have the following properties $i\hat{\mathbf{k}} \times \hat{\mathbf{h}}_c(\mathbf{k}) = ck\hat{\mathbf{h}}_c(\mathbf{k})$, $\hat{\mathbf{h}}_c(-\mathbf{k}) = \hat{\mathbf{h}}_c^*(\mathbf{k}) = \hat{\mathbf{h}}_{-c}(\mathbf{k})$ and $\hat{\mathbf{h}}_{c_1}(\mathbf{k}) \cdot \hat{\mathbf{h}}_{c_2}^*(\mathbf{k}) = \delta_{c_1,c_2}$, the Euclidean norm. A relation used for numerical computation, such as the numerical experiments with the (pseudo-)spectral method using the various truncation schemes to be discussed in §2.2, is $\hat{\mathbf{v}}^c(\mathbf{k}) = \hat{\mathbf{v}}(\mathbf{k}) + c\hat{\mathbf{k}} \times \hat{\mathbf{v}}(\mathbf{k})/k$ (see, *e.g.*, Lesieur 1990; Melander & Hussain 1993).

Here the new element in the theoretical formulation of the absolute equilibrium problem lies in viewing the system a gas of pure helical modes $\hat{\mathbf{v}}^c(\mathbf{k})$, representing $\hat{\mathbf{v}}^c(\mathbf{k})\hat{\mathbf{h}}_c(\mathbf{k})e^{i\mathbf{k}\cdot\mathbf{r}} + c.c.$ for simplicity, *i.e.*, the chiroids *a la* Kelvin (1904), as the working ‘momentoids’. Corresponding densities can be defined accordingly: For instance, *mean* magnetic energy $\mathcal{E}_M = \sum_{c,\mathbf{k}} U_M^c(k)$ and helicity $\mathcal{H}_M = \sum_{c,\mathbf{k}} Q_M^c(k)$ with

$$U_M^c(k) = ckQ_M^c(k) = \langle |\hat{B}^c(\mathbf{k})|^2 \rangle / 2, \quad (1.2)$$

with a reversed factor of k for the kinetic case due to the difference between magnetic and kinetic helicities by definition, where $\langle \bullet \rangle$ denotes the mean, per unit volume or in the statistical sense. Inserting Eq. (1.1) back into the rugged invariants, \mathcal{E} and \mathcal{H} , constraining the statistical ensemble, the Gibbsian one used for our calculations, we can obtain the chirally split densities $U^c(k)$ and $Q^c(k)$ which present finer physical structures than the mixed ones

$$U(k) = U^+(k) + U^-(k) \text{ and } Q(k) = Q^+(k) + Q^-(k) : \quad (1.3)$$

We remark that this result should be perceived in two perspectives. One is that the AE spectra of pure helical modes of each chiral sector present separately, independent of the existence of the other one, *i.e.*, whether the other sector is truncated or not for some \mathbf{k} s, since the truncation can be performed arbitrarily on the chiroids, except that Hermitian symmetry should be kept like the classical Fourier truncation; the other perspective is that the spectra are chirally decomposed into two sectors, if both exist, *i.e.*, the truncations for both sectors are symmetric. Note that $|Q_M(\mathbf{k})| \leq U_M(\mathbf{k})/k$, so the purely helical mode is called *maximally helical* (see, *e.g.*, Kraichnan 1973). Other derivatives such as the relative helicity $|kQ_M(k)|/[U_M(k)] = k \sum_c Q_M^c(k) / \sum_c U_M^c(k)$ (Kraichnan 1973, with a reversed factor of k as in Eq. 1.2) can be used for quantifying the degree of chirality. Only when the modes have the same wavelength and

† Note that we have used Coulomb gauge $\hat{\mathbf{A}} \cdot \mathbf{k} = 0$, so $i\hat{\mathbf{k}} \times [i\hat{\mathbf{k}} \times \hat{\mathbf{A}}(\mathbf{k})] = k^2 \hat{\mathbf{A}}(\mathbf{k}) = \hat{\mathbf{k}} \times \hat{\mathbf{B}}(\mathbf{k})$: The longitudinal component of \mathbf{A} with whatever gauge is not involved in the relevant calculations, so Coulomb gauge is not really necessary but brings symbolic convenience. Note also that helical *wave*, with the time argument included, is *truly* chiral in the sense of Barron (see, *e.g.*, Cintas & Viedma 2012).

are *homochiral*, i.e., all with same handedness, can we see the physical-space field resulting from their superposition is Beltrami, i.e., $\nabla \times \mathbf{v} = \gamma \mathbf{v}$ with constant γ , *force free* for magnetic field, in which case, nonlinearity is typically depleted completely.

1.3. Statistical ensembles of truncated chiroids

When the system is reduced to the dynamics of the helical modes, one can then consider various truncations directly on such chiroids (Waleffe 1992; Biferale, Musacchio & Toschi 2012). Detailed triadic interactions of the helical Fourier modes in various hydrodynamic-type models, such as HD, MHD, EMHD and Hall MHD etc., have been closely looked into by different authors (e.g., Waleffe 1992; Lessinnes, Plunian & Carati 2009; Galtier & Bhattacharjee 2003; Galtier 2006). And, relevant details of the AE calculation have been well described in the literature and do not require any further elaboration here. For instance, it is routinary to check Liouville theorem and rugged conservation properties after Galerkin truncation of the helical modes, which is true for all the models studied here (for HD, *c.f.*, equations 7 and/or 9 of Waleffe 1992). To be a bit more definite but without loss of generality, using indexed y s for the variables related to the real and imaginary parts of the active chiroids $\hat{v}^c(\mathbf{k})$, we can write down the dynamical equations as

$$\partial_t y_n = \sum_{l,m} Y_{lmn} y_l y_m, \quad (1.4)$$

with Y_{lmn} satisfying some specific symmetries to assure the detailed conservation of energy and relevant helicity(ies) and Liouville theorem. Note that for cases with linear terms of the original variables on the right-hand side, such as the 3D gyrokinetics (Zhu & Hammett 2010), a simple linear transformation of variables reduces them to this form which is formally the same as the well-known classical Fourier Galerkin truncated Euler case. Readers can go to the Appendixes for more discussions on the *detailed conservation laws, dynamical and topological aspects, and, tacit assumptions about ergodicity or mixing etc. in the statistical considerations.*

1.4. Plan

We progressively perform a minimal but systematic investigation, with different emphases, of EMHD with formally pure magnetic field dynamics in §2.1, HD in §2.2, and, single-fluid (§2.3.1) and two-fluid (§2.3.2) MHDs. §2.1 discusses mainly the natural chiral selection for inverse magnetic helicity transfer; §2.2 concentrates on the chirally asymmetric truncation effects, §2.3.1 on new insights to the classical dynamo issue, §2.3.2 on the two-fluid effects. Although many of the discussions in the (sub)subsections, such as the asymmetric truncations of the two chiral sectors for some \mathbf{k} s in §2.2, can be carried over to other (sub)subsections, *mutatis mutandis*, to get some relevant new insights, we won't detail such obvious points. The general purpose is to lay out the basic AE as a first step to explore some fundamentals of turbulent transfers, especially for a comprehensive basic understanding of the relevant helicity effects. It should be pointed out that there have been many other interesting AE-relevant investigations for different dynamical models and on various specific physical issues, the important one of which is relevant to a space uniform magnetic field and anisotropic fluctuations and has been continuously attacked in the last several decades. Extra particular studies should be done, though $k = 0$ mode can formally be included in the calculations and brief relevant remarks will be offered at the appropriate circumstances. The focus is the most basic new insights attached to the chirally decomposed AE, with which we will revisit the most relevant studies, rather than any other specific turbulence (closure) theories, such as the wave turbulence theories studied by Galtier and collaborators, though our results may be used as benchmarks of relevant analytical or numerical treatments.

In summary, an incompressible hydrodynamic-type system can be reduced to the dynamics of pure helical modes, the “chiroids” *a la* Kelvin, with helical wave/mode representation. Left-

and right-handed sectors of the absolute statistical equilibrium spectra are split. Either sector may present without the necessity of the existence of the other, *i.e.*, unichirally with asymmetric Galerkin truncation. One sector can dominate around its positive pole(s) with corresponding net helicity, providing a unique chirality selection and amplification mechanism. Chirally truncated systems preserve Moffatt's topological interpretation of helicity due to the detailed incompressibility for each chiroid. We obtain new insights about chirality selection and amplification, and, spectral transfers of turbulence of various neutral-fluid and plasma systems. For instance, one-chiral-sector-dominated states are naturally supported by magnetohydrodynamics (MHD) absolute equilibria with magnetic helicity, and homochiral Euler system allows negative temperature states which were excluded by Kraichnan for the non-homochiral case. A major purpose is to make a systematic comparison of the effects of various helicities, for finding genericities and specificities, and we clarify the special role of magnetic helicity for turbulent inverse magnetic helicity transfer/cascade by analyzing the electron MHD, with only magnetic field, and the two-fluid MHD, with the combination of various helicities in a symmetric way.

As said, it is not necessary to elaborate the calculations repeatedly, thus the following presentation will mainly consist in a set of brief backgrounds and theoretical frameworks, discussions of our results with careful comments on relevant studies and new explanations of documented data. Readers are suggested to go directly to the (sub)section(s) for the interested model(s)/topics first and then, before trying to read the analyses of others, to §3 for further discussions, where not only the major results are summarized, but also the genericities and differences are extracted by comparisons across different models.

2. One-chiral-sector-dominated absolute equilibrium and turbulence states

Basic (rules of) notations, definitions and calculation techniques follow §1. Readers are assumed to be familiar with the relevant ideas and techniques of K67 and K73 summarized there. We do not repeat the further detailed simple calculations of the spectra following K73 and Frisch et al. (1975) which will actually be found to be greatly simplified, since much, for single- and two-fluid MHDs, or all, for EMHD and HD, of the diagonalization work and the solenoidal constraints, have already been performed from the beginning with the helical representation while constructing our ensemble.

2.1. Pure magnetic fluid

EMHD equation

$$\partial_t \mathbf{B} + \nabla \times [(\nabla \times \mathbf{B}) \times \mathbf{B}] = 0$$

formally involves only the magnetic field and may be called pure magneto-dynamics. This fluid model corresponds to the small electron skin depth $d_e \ll 1$ limit of the more general case, which we will discuss later, and is relevant to helicons or whistler waves in solid conductor, including neutron star's solid crust, atmosphere etc (see, e.g., Biskamp et al. 1999; Galtier & Bhattacharjee 2003, and references therein). Note that \mathbf{B} is "frozen in", by definition, to the electron fluid velocity $\mathbf{u}_e = -\nabla \times \mathbf{B}$. Rugged invariants are magnetic energy and helicity:

$$\mathcal{E}_M = \frac{1}{2\mathcal{V}} \int \mathbf{B}^2 d^3 \mathbf{r} = \frac{1}{2} \sum_{\mathbf{k}, c} |\hat{B}^c(\mathbf{k})|^2 \quad \text{and} \quad \mathcal{H}_M = \frac{1}{2\mathcal{V}} \int \mathbf{A} \cdot \mathbf{B} d^3 \mathbf{r} = \frac{1}{2} \sum_{\mathbf{k}, c} c |\hat{B}^c(\mathbf{k})|^2 / k.$$

The two chiral sectors of the AE spectral densities (*c.f.*, §1) of energy and helicity are then

$$U_M^c(k) = k/(c\beta + \alpha k) \quad \text{and} \quad Q_M^c(k) = 1/(\beta + c\alpha k) = cU_M^c(k)/k. \quad (2.1)$$

From the above spectral relations, just as K67, but with energy playing the role of enstrophy there, a low energy state corresponds to condensation of Q at smallest wavenumbers, close to the

positive pole $k_p = -c\beta/\alpha$ of one of the chiral sectors, say, $c = +$, with $\beta < 0 < \alpha$. The implication for turbulence is inverse helicity and forward energy transfers. In principle, as long as there is net helicity with $\beta \neq 0$, one chiral sector can dominate at large scales, *i.e.*, OCSDS around the positive pole, though commonly done in experiment to provoke EMHD turbulence is to impose a background magnetic field (see, e.g., Stenzel 1999) which breaks the skin effect and guides the waves. If helicity is injected at some intermediate k , we should see dominant inverse helicity and forward energy transfers. If the transfers in these two regimes are approximable by self-similar local cascades and that suitable for simple dimensional analysis, energy spectra follow $k^{-5/3}$ and $k^{-7/3}$ scalings: Biskamp et al. (1999) first proposed and presented in slightly different situations such scalings, and their Fig. 8b with electron skin depth $d_e \ll 1$ does correspond to the forward energy cascade of our case. The scale separation between the dominant dynamics of the two cascade quantities of EMHD is weaker than 2D fluid turbulence, in the sense that the spectral ratio of each chiral sector is at most k , instead of k^2 of the latter. So, at finite Reynolds numbers, cascade of either definitely is accompanied by stronger (than 2D turbulence) leaking of the other. The subdominant energy transfer, accompanying the inverse helicity transfer and vanishing at high Reynolds number limit, should not be recognized to be the genuine inertial cascade. One should be particularly careful for the decaying case which is exactly the nice simulation by Cho (2011), who, by “inverse energy cascade”, meant merely the backward shift of the peak of his energy spectrum, which is not genuine in connect to the conventional notion of inertial cascade and which is not in conflict with our statement of forward energy cascade (Private communication).

Our result indicates a largest-scale nearly force-free magnetic fields. For the discrete- k case, the smallest- k modes contain most of the energy, so the whole global structure may appear to be roughly Beltrami, with smaller-scale “turbulent” fluctuations. Note that completely force-free fields, instead of ours with the scale-dependent degree of chirality measurable by the relative helicity (Kraichnan 1973), were obtained with several variational formulations in 1950s: We will come back to this in §2.3.1 for single-fluid MHD.

The basic feature, concerning the issue of magnetic helicity inverse transfers/cascades, of the EMHD results in the above is also central to other MHD models with magnetic field. Some brief remarks for the finite- d_e (which is used for scale normalization here) general EMHD model,

$$\partial_t(\nabla^2 \mathbf{B} - \mathbf{B}) + \nabla \times [(\nabla \times \mathbf{B}) \times (\nabla^2 \mathbf{B} - \mathbf{B})] = 0,$$

are in order. The “frozen-in” generalized vorticity is

$$\nabla \times \mathbf{P}_e = \nabla \times \mathbf{u}_e - \mathbf{B} \text{ with } \mathbf{P}_e = \mathbf{u}_e - \mathbf{A}.$$

The rugged invariants are now total energy and *generalized* helicity (Biskamp et al. 1999)

$$\mathcal{E} = \frac{1}{2\mathcal{V}} \int (\mathbf{B}^2 + \mathbf{u}_e^2) d^3 \mathbf{r}, \quad \mathcal{H}_G = \frac{1}{2\mathcal{V}} \int \nabla \times \mathbf{P}_e \cdot \mathbf{P}_e d^3 \mathbf{r},$$

resulting in

$$U_M^c(k) = k / \{(k^2 + 1)[c\beta(k^2 + 1) + \alpha k]\},$$

which complicates the quantitative transitional spectral behaviors (*c.f.*, §2.3.2 for more general discussions for similar situations in two-fluid MHD). In the other limit regime of scales much smaller than d_e , or in another word, when $k \gg 1$ and that $k^2 + 1$ can be replaced by k^2 , $k^2 U_M^c(k) \rightarrow 1/(c\beta k + \alpha)$, the same as the pure HD case, supporting both energy and helicity cascading forwardly (*c.f.*, Fig. 8a of Biskamp et al. 1999); † Magnetic helicity concentrating

† Note that, for convenience, in this regime one may want to study magnetic enstrophy W defined through $W(k) = k^2 U_M^c(k)$ and the other quantity S , which one might want to call magnetic *helistrophy*, defined through $S(k) = k^2 Q_M^c(k)$. Such an attempt however is conceptually not very appropriate, since neither of them are conserved quantities.

at or transferring to “large” (of course in the sense of subsidiary limit) scale can only be obtained with imposed asymmetric truncation between the two chiral sectors, such as that leaving only the $c = +$ sector (Waleffe 1992, and see §2.2 for the discussions). This should not be surprising, since electron kinetic fluid flow dominates in this limit.

2.2. Pure neutral fluids

For the classical incompressible HD, *i.e.*,

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} = -\nabla p,$$

where the pressure p can be eliminated by $\nabla \cdot \mathbf{u} = 0$, the rugged invariants are kinetic energy and helicity

$$\mathcal{E}_K = \frac{1}{2\mathcal{V}} \int \mathbf{u}^2 d^3 \mathbf{r}, \quad \mathcal{H}_K = \frac{1}{2\mathcal{V}} \int \nabla \times \mathbf{u} \cdot \mathbf{u} d^3 \mathbf{r},$$

which lead to the densities of separate chiral sectors:

$$U_K^c(k) = 1/(\alpha + c\beta k), \quad Q_K^c(k) = ckU_K^c(k). \quad (2.2)$$

Note that the above spectra can not be considered to be simply the decomposition (into two chiral sectors) of K73 densities (*c.f.*, §1) which are not valid when there is asymmetric truncation between the two chiral sectors of some \mathbf{k} , that is, when only one of the chiral sectors of some \mathbf{k} is truncated to be unichiral. For example, if there is no cancelation at some k , one can not derive from Eq. (2.2)

$$\alpha = [U_K^-(k) + U_K^+(k)]/[2U_K^+(k)U_K^-(k)], \quad \beta = [U_K^-(k) - U_K^+(k)]/[2kU_K^+(k)U_K^-(k)] \quad (2.3)$$

which are in particular not true for any k in the homochiral system with only one, say, the positive chiral sector. In such homochiral case with $c = +$, $\alpha > 0$ is not required by realizability condition (§2.2) and Eq. (2.2) shows that the low helicity state corresponds to a negative α with a sharp peak at the lowest modes $k_{min} > k_p = |\alpha/\beta|$ close to the positive simple pole k_p . Such \perp -shape energy spectral density, just as K67, indicates inverse energy and forward helicity dual transfers (Waleffe 1992, who also argued for this with his “instability assumption”) with Kraichnan’s argument of the tendency of relaxation towards equilibrium, as numerically realized by Biferale, Musacchio & Toschi (2012) with remarkable quality. Note however that such HD scenario does not work in vanishing- d_e EMHD in §2.1, neither for other more complicated MHD models as will be studied in the next section, which, for instance, when truncated to be homochiral, presents no drastic change of transfer/cascade direction; this is because the large-scale magnetic helicity concentration of OCSDS with symmetric truncation and that of the homochiral state coincide.

2.2.1. OCSDSs in the sense of fluxes: “Second order” OCSDSs

If both sectors present for every \mathbf{k} , *only* \perp shape spectral density dominated by one of the sector around the peak is allowed by the realizability conditions $\alpha > 0$ and $k_{max} < \alpha/|\beta|$ from the positive definiteness of the quadratic form: *c.f.*, §1. Large-scale condensation mechanism is absent, thus inverse cascade in HD generally needs other special treatments as reviewed by Yang, Su & Wu (2010) and Biferale, Musacchio & Toschi (2012). Although, unlike at large scales, normal dissipation would devastatingly ruin such small-scale explicit OCSDS AE structure, some residuals of such intrinsic nonlinearity effects may persist. Indeed, the fluxes of the two chiral sectors reported by Chen, Chen & Eyink (2003) do show systematic differences: According to the working conditions, their Figs. 2–5 correspond to the case with positive helicity, that is, the positive pole belongs to the positive sector with negative β . Dominance of the positive sector of AE spectrum indicates that nonlinearity should support the transfer of this sector to

be more persistent, consistent with the results and analyses of Chen, Chen & Eyink (2003). We may also call them “implicit” OCSDSs, or “second order” OCSDSs, in the sense of dominance of energy and helicity fluxes of one sector, as signatored by the lower panels of their Figs. 4 and 5. Such second-order OCSDSs may be viewed as yet another special evidence, besides the isotropization (Lee 1952) and bottleneck (Frisch et al. 2008), of the *persistence of thermalization* around the end of inertial range. The large- k viscous effect efficiently restores the reflexion symmetry, and the degree of local-in- k chirality measured by relative helicity vanishes as k^{-1} throughout the inertial range with accurate $k^{-5/3}$ scaling exponents for both energy and helicity (Kraichnan 1973). The large- k pole effect of one chiral sector nevertheless provides a prototype for other similar possible physics (*c.f.*, §2.3.2) in more complex situations, furthermore it might be possible to find its stronger activity in a non-Newtonian fluid such as the (dilute) solution of chiral polymers, say, deoxyribonucleic acid (DNA), in a normal fluid where some kind of resonance between one chiral sector of the fluid motion and the polymer’s chiral structure/activity could happen; see more relevant discussions in §3.1.

2.2.2. OCSDSs with special truncation schemes and “smooth” transitions of energy transfer directions

Negative temperature state emerges with dramatic physical indications for the homochiral case. Now, if we add just one pair of conjugate modes, with opposite helicity, negative temperature is then excluded, by the nonnegativity of U_K^c . Naturally, a kind of “phase transition” is happening in the temperature parameters, since the negative temperature must jump to some positive value. Then with superficial impression from the above analysis and from K73 one would tend to expect similarly a sharp transition of inverse energy cascade to forward energy cascade. But, such a superpowerful potential of a single alien chiroid would be shocking. Thus we need to look into the corresponding absolute equilibrium states by starting with a clean pool of homochiral $c = +$ modes and put aliens into it. *It turns out that there should be no phase-transition-like behavior concerning cascades and energy can “smoothly” switch from completely-inverse to partly-inverse-and-partly-forward and to completely-forward cascades, depending on how (many) aliens are put into the pool. For the general truncations of chiroids with asymmetry but not homochirally, K73’s argument for excluding inverse energy cascade still does not simply work, and large-scale concentration of energy, indicating inverse energy transfer in turbulence, can exist without a negative temperature state.* To see this, suppose we have only one alien chiroid, *i.e.*, one negative-helicity mode, at the objective *condensation wavenumber*, $k_c = k_{min}$ much smaller than the injection wavenumber k_{in} , and that negative temperature state is excluded for implying a conventional inverse energy transfer/cascade argument *a la* K67. Would energy abruptly turn to cascade forwardly, or there should be a transitional behavior? To get illumination, one may apply to the arguments and thought experiment presented at the end of §1.1. Consider Eqs. (2.2) and (2.3) with $\alpha > 0$ and $\beta > 0$, and suppose energy is injected at some intermediate k_{in} . This alien mode may help transfer extra positive helicity nonlocally to small scales, by which, though, its own amplitudes of (negative) helicity and energy would have to increase; this is because for the excitation of any mode in $k > k_{in}$ the injected helicity at k_{in} is not enough to support it, just by the relation $Q_K(k) = kU_K(k)$ for pure helical mode, thus some extra positive helicity should be provided from $k < k_{in}$, which is facilitated by the excitation of negative helicity in the alien mode. The growth of this alien mode is allowed, even in the sense of complete AE, with α/β approaching k_c^+ , *i.e.*, from above:

$$\alpha/\beta \rightarrow k_c^+ \text{ and that } U_K^-(k_c) \rightarrow \infty, \text{ turning } k_c \text{ into the pole } k_p. \quad (2.4)$$

Note that k can be larger than α/β even though $U_K^-(k) = 1/(\alpha - \beta k)$, since the alien mode in this sector is restricted below k_c , actually, as said, the $c = -$ sector of Eq. (2.2) being only for $k_c = k_{min}$ now, unlike the traditional symmetrically truncated system with $\alpha/\beta > k_{max}$ as discussed

by Kraichnan (1973); or, in another word, Kraichnan (1973) could not let α/β approach k_c from above, because he had other larger- k alien modes which otherwise would have negative energy for $k > \alpha/\beta$. Thus, adding such a single alien excludes the negative-temperature state, but the single alien can carry the energy condensating there as a particular form of OCSDS. The k -distribution of energy of such a state does not differ from that of the homochiral negative temperature state too much. And, one probably can carefully design a simulation by adding an alien to the smallest k of Biferale, Musacchio & Toschi (2012)'s simulation and still get inverse transfer of energy. When we gradually increase the number of aliens starting from k_{min} to the condensation wavenumber $k_c \geq k_{min}$, there may or may not be a nonzero forward nonlocal transfer of energy to small scales in the infinite Reynolds number limit; or, in another word, depending on the strength of the nonlocal kicking of the alien modes, the energy accompanying the forward helicity transfer may or may not vanish as the wavenumber goes to infinity: It may not be impossible that, on average, a single alien could remotely “kick” the small scales by “stirring” up vortex stretching over the field, making the solution rougher and that causing some finite energy dissipation. With more aliens, forward helicity transfer will be more. The injected energy must gradually be partitioned to be transferred to two opposite directions simultaneously in the infinite Reynolds number limit. And, to transit to a state with all energy completely cascading forwardly, sufficient aliens must be added to the regime of k larger than k_{in} , since if all aliens are added only to the regime below k_{in} , the nonlocal interaction across k_{in} for transferring helicity to larger k must be accompanied with some transfer of energy to $k < k_{in}$, as is also indicated by analyzing the corresponding absolute equilibria [*c.f.*, statement (2.4) for $k_c < k_{in}$]. Regarding the cascades and/or nonlocal transfers in the infinite Reynolds number limit, the approach we have taken is *tuning the degree of the regularity of the solutions to be appropriately “dissipative” by manipulating the population of the helical modes*. To our best knowledge, for the full 3D Navier-Stokes, actually even for 2D, there is not yet satisfying mathematical theory for the cascade statements in the infinite Reynolds number limit. But, it appears that we may use the absolute equilibrium states to obtain at least some fine and clear physical-picture intuitions for such an approach.

2.2.3. HD summary

We have come from *magneto-dynamics* in §2.1 to this *hydro-dynamics* which, with the great substantiation for transition to *magneto-hydro-dynamics*, needs a summary:

- (a) Kraichnan (1973)'s argument for forward energy and helicity cascades can be refined to find the relevance of his absolute equilibrium with the energy/helicity fluxes reported by Chen, Chen & Eyink (2003), as a kind of “second order” OCSDS;
- (b) as shown by our chirally split spectra, his argument however is not for the homochiral truncation where the negative temperature state, indicating inverse energy transfer/cascade, is allowed;
- (c) such a “sharp” change of cascade scenario is not shared by the pure *magneto-dynamics*;
- (d) The cascade transition is not really “sharp” but has a “smooth” dependence on how alien modes are added to the homochiral pool.

2.3. Magnetised fluids

In principle, there can be many fluid models for describing different subsets of the kinetic phase space of plasma dynamics (see, *e.g.*, a very limited list in a review by Schekochihin et al. 2009). Here we study the classical single-fluid and the most general two-fluid MHDs. From the plasma physics point of view, two-fluid MHD is for a more complete description of the kinetic effects for the dynamics/scales between those of EMHD and single-fluid MHD. Two-fluid MHD presents various helicities in a unified way.

2.3.1. Single-fluid MHD

As introduced in §1, Meneguzzi, Frisch & Pouquet (1981) found with direct numerical simulations that “the large-scale \mathbf{B} is mostly force free and produces only very little large-scale motion,” with the relative magnetic helicity density $|kQ_M(k)/U_M(k)|$ being close to 1, nearly maximally helical; and, recently, Brandenburg, Dobler & Subramanian (2002) explicitly pointed out, by their Fig. 21 with postprocessing using helical decomposition, that the simulation with similar setup also present such OCSDSs. We now turn to explain such findings with the classical single fluid MHD equations

$$\begin{aligned}\partial_t \mathbf{u} &= -(\mathbf{u} \cdot \nabla) \mathbf{u} + (\mathbf{B} \cdot \nabla) \mathbf{B} - \nabla(p + B^2/2), \\ \partial_t \mathbf{B} &= -(\mathbf{u} \cdot \nabla) \mathbf{B} + (\mathbf{B} \cdot \nabla) \mathbf{u}, \text{ where } \nabla \cdot \mathbf{u} = 0 \text{ and } \nabla \cdot \mathbf{B} = 0.\end{aligned}$$

Rugged invariants are three (see, *e.g.*, Woltjer 1959; Frisch et al. 1975), the energy, magnetic helicity and cross helicity

$$\left\{ \mathcal{E} = \frac{1}{2\mathcal{V}} \int (\mathbf{u}^2 + \mathbf{B}^2) d^3\mathbf{r}, \mathcal{H}_M = \frac{1}{2\mathcal{V}} \int \mathbf{A} \cdot \mathbf{B} d^3\mathbf{r} \text{ and } \mathcal{H}_C = \frac{1}{2\mathcal{V}} \int \mathbf{u} \cdot \mathbf{B} d^3\mathbf{r} \right\},$$

which, together with Eq. (1.1), leads to (*c.f.*, §1, also for notations)

$$U_K^c(k) = \frac{4(\alpha k + c\beta_M)}{(4\alpha^2 - \beta_C^2)k + c4\alpha\beta_M}, \quad U_M^c(k) = \frac{4(\alpha k)}{(4\alpha^2 - \beta_C^2)k + c4\alpha\beta_M}, \quad (2.5)$$

$$Q_M^c(k) = \frac{c}{k} U_M^c(k), \quad Q_K^c(k) = ck U_K^c(k), \quad Q_C^c(k) = \frac{-2\beta_C k}{(4\alpha^2 - \beta_C^2)k + c4\alpha\beta_M}. \quad (2.6)$$

Similar to the statement in §2.2 in comparing our spectra to those of Kraichnan (1973), our spectra can not be considered to be simply the decomposition of those of Frisch et al. (1975) which are not valid when there is asymmetric truncation of the two chiral sectors of some \mathbf{k} . When there is no asymmetric truncation, our spectra chirally split those of Frisch et al. (1975) following whom we start with the case of null cross helicity with $\beta_C = 0$ for discussions: $Q_M^c(k)$ with $\text{sgn}(c\beta_M) = -1$ is responsible for the condensation of Q_M at small k , around the positive simple pole $k_p = -c\beta_M/\alpha$. When the dynamics is dominated by the $c = +$ ($c = -$) sector, it is simply to say that large positive (negative) magnetic helicity state corresponds to a negative (positive) β_M with a \perp shape spectral density is favored: Frisch et al. (1975) plotted such spectra, the spectral densities multiplied by k^2 , in their Figs. 1 and 2 for illustration, by choosing several typical temperature parameters. The other sector’s pole has the opposite sign and is not reachable, thus, without such a mechanism of large-scale attraction, the energy would be transferred to small scales or simply less excited. When β_C (or \mathcal{H}_C) is nonzero, the prefactor before k in the denominators quantitatively changes, but the qualitative picture is not altered. As pointed out in §2.1, the large-scale nearly maximally helical state predicted by AE is close (see also next paragraph) but different to the purely force-free one intensively studied in 1950s (see references in Woltjer 1959). The common feature with that of Woltjer (1959) is that the invariant cross helicity does not essentially change the large-scale nearly maximally helical physics. Recently, some authors argue that cross-helicity, signature of the imbalance along and opposite to the background magnetic field, may be important to determine the (reduced) MHD forward cascade inertial range scaling exponent (see, *e.g.*, Perez et al. 2012, and references therein), which is beyond the scope of this study, although a spacially uniform \mathbf{B}_0 can be formally included in our calculation and analysis.

Thus, the OCSDSs in Meneguzzi, Frisch & Pouquet (1981) and Brandenburg, Dobler & Subramanian (2002) are related to our AE spectra. Actually, Pouquet, Frisch & Léorat (1976) had carried out systematic study of eddy-damped quasi-normal Markovian (EDQNM) MHD turbulence and non-linear dynamo. The data of their Figs. 4 and 5 around $k = 0.16$, actually starting from the be-

gining of the inertial range to ever larger scales, and, of Figs. 8 and 9 around $k = 0.1$, already clearly presented OCSDSs as can be seen from the values of the relative helicities computed from their figures. Obviously, as EDQNM shares the conservation properties of the original system, it satisfies the AE spectra and that the OCSDS arguments also work. We won't go too far into much more details, but just remark that the pertinent discussion of Pouquet, Frisch & Léorat (1976, p. 345, second paragraph) can also be elaborated: For instance, the large-scale ensemble can be understood by AE with positive magnetic helicity, with finer chiral-sector dynamics for the different chiralities in separate scale regimes. We only want to remark that relaxation of magnetic helicity to the largest scales does not necessarily indicate local cascade, nonlocal transfer also is possible (see, *e.g.* Brandenburg & Subramanian 2005); and, there is nothing in conflict with the more mechanical reasoning, such as the alpha effect.

Conceptually, a reader may quickly question whether we really learn anything more from chiral decomposition than that if helicity is large, one chirality must dominate. Isn't that obvious? The simple pole mentioned under Eqs. (2.5) and (2.6) is already present in the spectra of Frisch et al. (1975) and indeed accounts for the accumulation of magnetic helicity at large scales. Chiral decomposition is not needed to reach this conclusion? The answers are “no”s. To understand these problems, one must first understand that large net helicity does not necessarily mean OCSDS or big relative helicity. Now, suppose the spectra were not chirally decomposed and that the small- k pole might contribute to both left- and right-handed helicities. In such an ambiguous situation, one would not be able to conclude OCSDS as the scale approaching the pole, which was exactly what happened in the past studies mentioned in the above, sounding like just a hair's breadth though. Interestingly, even in the extremely strong sense of “non-helical” state with $Q_M(k) = Q_K(k) = 0$, *i.e.*, the two chiral sectors of both magnetic and velocity fields balancing at each wavenumber, AE seems to still support the so-called non-helical turbulent dynamo. The reason is that, in this situation, while energy, either kinetic or magnetic, is equipartitioned into each helical mode, magnetic helicities of both sectors with opposite signs are “attracted” by the same pole $k_p = 0$. Note that unlike EMHD in §2.1, magnetic energy itself here is not conserved and kinetic energy can be transformed to it to ease the inverse magnetic helicity transfers for the two chiral sectors simultaneously. Note also that Pouquet, Frisch & Léorat (1976) and Meneguzzi, Frisch & Pouquet (1981), and other isotropic MHD simulations with unit Prandtl number, found a slight excess of magnetic energy at small scales, which may be due to such “attraction” from large scales. Without decomposition, as net helicity at any k is seen to be zero, researchers traditionally have not seriously thought about the simultaneous backward transfer of both sectors, to our best knowledge.

2.3.2. Two-fluid MHD

Two-fluid effects, the decoupling between electrons and ions, are important in many laboratory and astrophysical situations (see, *e.g.*, Yamada et al. 2002; Brandenburg & Subramanian 2005). The ideal incompressible two-fluid MHD states that the generalized vorticities $\nabla \times \mathbf{P}_s$, with canonical momenta $\mathbf{P}_s = m_s \mathbf{u}_s + q_s \mathbf{A}$ for each species s , are “frozen in” (*c.f.* §2.1) to the respective flows (see, *e.g.*, Ruban 1999, and references therein)

$$m_s n_s \frac{d\mathbf{u}_s}{dt} = q_s n_s (\mathbf{E} + \mathbf{u}_s \times \mathbf{B}) - \nabla p_s,$$

where \mathbf{E} is the electric field vector and q_s and m_s are charge and mass. Since this model has very rich physics, to remind ourselves the relevant context and the weights of physical quantities in quantifying chirality, instead of being purely geometrical as discussed in Petitjean (2003), we now use the normalization which keeps some physical parameters explicitly, unlike those in EMHD and single-fluid MHD. The dynamics is constrained by three rugged invariants, *i.e.*, the

total energy and self-helicities:

$$\mathcal{E} = \frac{1}{2\mathcal{V}} \int [\mathbf{E}^2 + \mathbf{B}^2 + \sum_{\bar{s}} m_{\bar{s}} n_{\bar{s}} u_{\bar{s}}^2] d^3\mathbf{r} \text{ and } \mathcal{H}_s = \frac{1}{2\mathcal{V}} \int \nabla \times \mathbf{P}_s \cdot \mathbf{P}_s d^3\mathbf{r}.$$

Here, two-fluid effects are in the extra terms in the invariants compared to single-fluid MHD.

With Eq. (1.1), we are led to the following AE spectra densities (*c.f.*, §1, also for notations)

$$Q_M^c(k) = \frac{c}{k} U_M^c(k), \quad U_M^c(k) = \frac{kT(k)}{D^c(k)}, \quad U_{sK}^c(k) = \frac{N_s^c(k)}{D^c(k)}, \quad Q_{sC}^c(k) = \frac{q_s \beta_s L_{\bar{s}}(k)}{-D^c(k)/k},$$

$$Q_{sK}^c(k) = ck U_{sK}^c(k) \text{ and } Q_s^c(k) = m_s^2 Q_{sK}^c(k) + 2m_s q_s Q_{sC}^c(k) + q_s^2 Q_M^c(k) \quad (2.7)$$

with $D^c(k) = \alpha k [T(k) + (\sum_{\bar{s}} q_{\bar{s}}^2 n_{\bar{s}} m_{\bar{s}}) \prod_{\bar{s}} \beta_{\bar{s}}] + c \alpha O \prod_{\bar{s}} n_{\bar{s}}$, $L_s(k) = \alpha n_s + c \beta_s m_s k$, $m_s N_s^c(k) = k[\alpha L_{\bar{s}}(k) + q_{\bar{s}}^2 m_{\bar{s}} \prod_{\bar{s}} \beta_{\bar{s}}] + \underline{c n_{\bar{s}} O}$, $O = \alpha \sum_{\bar{s}} q_{\bar{s}}^2 \beta_{\bar{s}}$ and $T(k) = \prod_{\bar{s}} L_{\bar{s}}(k)$ where \bar{s} means the other species than s and where the index C is for the ‘‘cross’’ helicity as defined in §2.3.1. We summarize the following points:†

First. The poles, *i.e.*, roots of the third order polynomials $D^c(k)$, of the two sectors are of opposite signs, as c appears in the second and zeroth order terms.

Second. The relevant spectrum may be of \perp shape, with a positive pole on the left. This is similar to the single-fluid MHD (§2.3.1) case with OCSDS of inverse magnetic helicity transfer. Like the discussion in the end of §2.1 for the regime of scales much smaller than d_e , \perp shape spectral density as in HD (§2.2) may be relevant to the scenario of both energy and helicities cascading forwardly. Now there can be other larger positive poles, from the same chiral sector or not, which may also be physically relevant to the possible persistence or emergence of chiralities due to the dominance of different physical processes at different scale regimes. Note that like Hall MHD in Meyrand & Galtier (2012), ion MHD (IMHD) or EMHD can be identified in our two-fluid model by putting the fluid velocity of one species to zero: Asymmetries between the respective dynamics, with the opposite chirality in their sense of whistler or cyclotron waves, may present at different scale regimes, due to, say, the different m_s of the two species (unlike electron-positron plasma).‡

Third. A \sqcup shape spectral density [$Q_M(k)$, say] may be confined inbetween two distinct positive poles, belonging to the same chiral sector or not, which presents cross-scale, *e.g.*, going across the ion or electron skin depths etc., behavior: The general EMHD in §2.1 with finite electron inertial can already have such a feature as is clearly seen from $U_M^c(k)$ given there. The general effects can be understood with the combination of the two cases in the above *second* item. And, the large- k peak may also be relevant to small-scale field generation (see, *e.g.* Brandenburg & Subramanian 2005, in the context of battery mechanisms) or indicating the ‘‘second order’’ or ‘‘implicit’’ OCSDS as discussed in §2.2.

Last. If $O = 0$ in the zeroth order terms underlined below Eq. (2.7), magnetic helicity does

† The result unavoidably appearing a bit complicated, it may be helpful for readers to focus on $U_M^c(k)$ and $Q_M^c(k)$ first. Further simplification of these formulae can be made for some situations. For instance, electron-positron plasma with mass equivalence and charge conjugation enables us to take all masses and charges be normalized unity. But, for our purposes here, taking D^c as polynomials of k with the fundamental theorem of algebra and Vieta’s formulas in mind suffices. Note that we have general formula for the roots whose nature is determined by the discriminant. In the cases discussed below, the parameter regimes (constrained by the realizability condition) can be obtained with such basic knowledge by some simple but tedious manipulations and are omitted here. Electric energy distribution is omitted for two reasons: One is that it can be neglected in usual cases; and, the second is that it is decoupled from the others.

‡ The chirality of Meyrand & Galtier (2012) designated by their magnetic polarization does not depend on our sense of chiral sectors in the IMHD or EMHD linear dispersion relations exploited by them, but only on the signs of the charges. In our work, chirality refers to the definite right- or left-handed sector in the helical representation, which is more basic and works for any models, and which may be used to similarly interpret strong turbulence (not limited to the linear wave dispersion relation) as well.

not act in constraining the AE ensemble through $\sum_{\bar{s}} \beta_{\bar{s}} \mathcal{H}_{\bar{s}}$ (*c.f.*, §1). Note that $k = 0$ does not become a pole for this situation, as every spectrum in Eq. (2.7) has a factor of k to cancel it. Since now $\prod_{\bar{s}} \beta_{\bar{s}} < 0$, we can see that this case is similar to the last situation considered by Kraichnan (1967, p. 1423), with our $[T(k)]^{-1} > 0$ from the realizability conditions being eligible to act the role of k^2 there; see also $\bar{U}_K(k)$ in §1. This corresponds to a \lrcorner shape spectral density. The pole for large-scale concentration is gone now, since only Q_M^c , by definition, has $k = 0$ as the asymptotic pole as in single-fluid MHD. *We thus can infer from this point that magnetic helicity constraint under two-fluid framework is still crucial for large-scale concentration of magnetic fields, as in single-fluid MHD where it is conserved.*

Careful analyses with appropriate choices of the physical parameters can be made for detailed illustration and more subtle implications. Similarly is for other intermediate models such as Hall MHD and general EMHD with finite electron inertial in §2.1 whose result is a bit simpler, with the denominators being polynomials of second order. From the plasma physics point of view, it is very interesting to spell out all detailed effects of each physical element (skin depths, mass ratio effects etc.), which however is not the focus of this paper. We have to refrain from treating these cases in too great a detail.

3. Further discussions

3.1. General remarks

Turbulence statistics can be sharpened with helical representation, which has been well discerned since Moffatt (1970) and K73 (Kraichnan 1973) and has become fully workable since Moses (1971) and Waleffe (1992). We have merely focused on the most basic AE aspect concerning the direction of spectral transfer as well as the selection and amplification of chirality. The key point is that although the dynamics of the two chiral sectors are in general coupled, the absolute equilibrium spectra are cleanly split, with poles of opposite signs. Not only that the finer physical structures offer new insights about the “(near-)racemic mixture” (*c.f.*, the last paragraph of §2.3.1), but also that one should keep away from the misconception that the chirally decomposed quantities derived in this paper never appear in the AE ensemble by themselves, but always in combinations giving inviscid invariants. Actually, OCSDSs may emerge in natural systems due to mechanisms relevant to what we have discussed or one can work with samples of “enantiopure compounds” in §2.2. Concerning partial fraction decomposition, our results physically assures the decomposability of the spectra of the traditional Fourier modes from the hydrodynamic-type models studied here and practically solves the mathematical problem, giving also the nice “conjugate” mathematical structures in the spectra of the opposite chiral sectors, at least to the degree of two parts with poles of exactly opposite signs, which is not trivial for some models such as the two-fluid MHD in §2.3.2. Naive attempts to perform the “post” decomposition of the traditional spectrum could be formidable and confusing, for lack of physical motivation: For instance, one could think of trying further to decompose the already chirally decomposed two-fluid MHD spectra.

For the absolute equilibria themselves, both K73 and Frisch et al. (1975)’s insights were almost here as we are now. Especially K73 explicitly discussed the interactions of pure helical modes. Curiously, K73 however did not[†] study the chiroids absolute equilibria to which his

[†] Kraichnan (1973) might not have been motivated to systematically examine the dynamics in helical-mode representation, especially the Liouville theorem and detailed conservation laws, the latter of which only appeared two decades later (Waleffe 1992). There is a conceptual issue here as we will elaborate a little bit. He wrote in the second page of that paper: “The two helical waves provide an alternative to the usual Fourier decomposition into plane-wave components.” In the usual Fourier representation, with consideration of isotropy (but lack of reflexion symmetry) as he was considering, there is no reason and no way to distinguish special components of the 3D spectra or spectra of special components of the Fourier

traditional mixed ones can not be reduced by taking any limits of any of the temperature parameters, in which sense we mean, in §2.2, his results are not valid for asymmetrically truncated systems. Such a piece of thin “window paper” was not pierced probably due to the fact that conventionally the relevance of the traditional chirally symmetric Galerkin truncation were made to the classical chirally symmetric viscosity or resistivity following Lee (1952, footnote 2) and Kraichnan (1973, footnote 8). For such physical considerations, see also some recent works (Frisch et al. 2008; Zhu & Taylor 2010) where other dissipation models lead to convergence to the classical Galerkin truncations in some sense. However, with our HD in §2.2 results, we have refined Kraichnan’s argument to reveal an interesting feature of helical turbulence beyond the non-existence of negative temperature states that Kraichnan emphasized. Just as Kraichnan used the AE to suggest directions of energy transfer, so that a pile-up of energy at large scales in the 2D case can suggest the flux of energy to large scales far from equilibrium, we suggest that the preferential transfer into one chiral sector observed by Chen, Chen & Eyink (2003) might be related to our observation about his helical AE. Furthermore, if we can somehow introduce chirally asymmetric dissipation and/or resistance in (magneto)fluids[‡], then the small-scale damping as discussed in §2.2 would not be simply only for chiral symmetry restoration and that explicit small-scale chirality selection and amplification similar to those at large scales could also present. Actually, in general plasma dynamics, such as cyclotron damping, essentially a 3D analog of the classical 1D Landau damping, and plasma heating (see, *e.g.*, Chaps. 10, 11 and 17 of Stix 1992), our result may be of stronger connection, since the ion and electron cyclotron resonances are of opposite chiralities and at different scales, addressable by two-fluid MHD model in §2.3.2.

3.2. Comparisons: for genericities, specificities and beyond

A major purpose of this work is to find the genericities and differences of various helicity effects by comparisons of the different hydrodynamic-type models. The subject in the center of the comparison is that of magnetic versus kinetic helicities. The pure magnetodynamic result for OCSDS of magnetic helicity (transfer) at large scales, as represented in the vanishing- d_e EMHD, generically lies in the core also of other MHD models. Two-fluid MHD has the most general and complete elements of helicities and show convincingly the crucial role of magnetic helicity for large scales. It appears to be nothing deep but simply due to the mathematical relations

$$Q_K^c(k) = kU_K^c(k)$$

and

$$Q_M^c(k) = U_M^c(k)/k$$

by definitions, by which, one can practically assume equipartition between kinetic and magnetic energy of same chiral sector at some intermediate scale and find that magnetic helicity of that sector belongs more to larger scales. One “artificial” way to look at it is the following: The gyrofrequency of a charged particle’s helical motion around \mathbf{B} is $\Omega = qB/m$, which means

coefficients, since they are all statistically identical. And, now the helical representation, as he noted, is only an alternative to the usual Fourier representation, concerning the degrees of freedom, thus he might omit the important distinguishability of the spectra between the opposite sectors.

[‡] It might be possible to work with some chiral (conducting) polymers; or, for classical magnetofluids, some special electromagnetic techniques would be wanted. Note that conventional study of elastic polymer effects (such as Procaccia, L’vov & Benzi 2008; Steinberg 2009) have not paid attention to the chirality, that is, the possibility of a third chiral time scale τ_θ^c , over which the (chiral) torque is to be balanced, besides the transverse and longitudinal ones (τ_\perp and τ_\parallel in Hatfield & Quake 1999) of the extended coil/helix (such as DNA), and that in a simple dilute polymer solutions dynamical model (Fouxon & Lebedev 2003, whose nonlinear dynamics is exactly the same as the classical 3D single fluid MHD studied in §2.3.1!) only a single relaxation time τ is used for all modes of \mathbf{B} , the so-called “tau approximation”. It is possible that the 3D chiral property of the polymers have non-neglectable rheological effects, especially in the turbulent states where small-scale helical modes are excited.

that we can formally take the B -line as “kinetic vorticity” Ω -line, macroscopically; magnetic field is indeed a *pseudovector*, like fluid vorticity, allowing the well-known dynamical analogy between them as initiated by Batchelor (see, *e.g.*, Moffatt 2008). This then gives various helicities a kind of unified description. Magnetic helicity is thus related to the more “intrinsic” plasma particle motion. There are also other supports of the robustness of magnetic helicity (see, *e.g.*, Brandenburg & Subramanian 2005), such as analysis with more general context (Berger & Field 1984) and measurements (Ji 1999).

The HD and the large- d_e EMHD situations are different to the others, in the sense that the realizable AE spectra can only have positive pole at large k s which regime however is subject to dissipation in real physical systems. That is, reflexion symmetry breaking and restoration mechanisms meet at the same battlefield and they reach another kind of equilibrium balance which is far from our statistical absolute equilibrium, whose implications and residuals concerning chirality in conventional fluids can still be identified with careful analyses as shown in §2.2 by refining Kraichnan’s argument. Restricting to the homochiral situations (Waleffe 1992; Biferale, Musacchio & Toschi 2012), HD kinetic energy or EMHD magnetic helicity accumulating at or transferring to large scales becomes possible as indicated by the negative- α state with small- k pole, and we further find that, as long as the asymmetry is strong enough, adding aliens to exclude the negative temperature state does not necessarily drastically change the transfer picture. It is possible to control the smooth transition from completely inverse to partly inverse and partly forward and to completely forward transfers.

Concerning turbulence cascade in the infinite Reynolds number limit or some kind of thermodynamic limit in the sense of $k_{max} \rightarrow \infty$ in the conventional Galerkin truncation, full 3D Navier-Stokes’ energy and helicity both cascading to small scales indicates that the solution is singular. Note that such an indication however has not yet found rigorous mathematical support and that there is still space for opposite conjectures, such as a solution as some kind of “directional limit” without such dissipative anomalies (Zhu & Taylor 2010). *Careful examination of chiroids absolute equilibria as partly illustrated in §2.2 turns out to be able to give fine and clear intuitive pictures about the roughness of the solutions.* Now, for homochiral 3D Navier-Stokes with, say, $c = +$, it is expected that only helicity, but not energy, is transferred to small scales, which indicates that the solution is slightly less singular, with the Hölder exponent h in

$$\delta u(\ell) \sim \ell^h$$

be some value inbetween $1/3$ and $2/3$ (Eyink 2008), of course in some statistical sense as the multifractal spectrum of h spans over a wide range in realizations (Frisch 1995). The self-similar pure kinetic helicity cascade spectrum would go as (Brissaud et al. 1973; Waleffe 1992)

$$H_K^+(k) \sim k^{-4/3}$$

which, unlike Kraichnan’s 2D enstrophy spectrum $\sim k^{-1}$, is convergent when integrated over k . Note that now $h = 2/3$. This convergence so far does not bring any troubles: Unlike 2D Euler, where finite enstrophy ensures the smoothness of the solution and that in principle ensures an equilibrium statistical mechanics without truncation (Miller 1990; Robert 2003), there is no mathematical theorem to assure conservation of helicity with its finiteness (see, *e.g.*, Eyink 2008). Of course, such cascade still may have spacial intermittency, in the sense of Onsager (Eyink 2008) that the helicity dissipation appears “spotty”, in which case an anomalous part of the dissipation may be considered to be *curdling* with infinite density on some fractal sets of zero volumes (Mandelbrot 1974), described by some Dirac delta function supported by the fractal, as an extremal limit. Coming back to full Navier-Stokes and supposing both energy and helicity cascading forwardly as $k^{-5/3}$, we immediately see from Eqs. (1.2) and (1.3) that

$$H_K^c(k) = cC_1 k^{-2/3} + C_2 k^{-5/3}$$

with C_1 and C_2 being constants (Ditlevsen & Giuliani 2001). That is, the two sectors of helicity both present ultraviolet divergences, indicating more singular solutions, which is consistent with the absolute equilibrium spectra showing poles at large k , compared to the small- k poles for the homochiral case. Thus, *as indicated by the smooth transition from backward, to bi-direction and to forward cascades by manipulating the helical modes in §2.2*, how the added “alien” helical modes increase the nonlinearity to roughen the solutions is intriguing and may be relevant to the intermittency property in the sense of Onsager and Mandelbrot as mentioned above.

3.3. Conclusion and prospects

Since an incompressible hydrodynamic-type system can be reduced to the dynamics of chiroids *a la* Kelvin, it is natural that one reduces the statistical dynamics to what is based on them and “hopes that one can get some insight into the nature of more general viscous flows and even, perhaps, a deeper understanding of turbulence.” (Moses 1971) We have studied the chirality issue, starting from and essentially based on the chiroids absolute equilibria. The equilibrium spectra can also be used to guide and benchmark numerical experiments with truncation schemes such as those discussed in §2.2 with asymmetric truncation between the two chiral sectors. Hints for further theoretical considerations may also be inferred. For instance, the clear OCSDS for large-scale magnetic helicity could imply some clues to dynamical dynamo model. Looking further into anisotropic fluctuations with a background magnetic field (for such discussions under the framework of 3D gyrokinetics, see, *e.g.*, Zhu & Hammett 2010) and to more realistic laboratory situations is also a reasonable step towards a more comprehensive theory for multi-scale plasma dynamics. And, due to the cross-disciplinary popularity of the notion of chirality as the legacy of Pasteur and Lord Kelvin (see, *e.g.*, Barron 1997), one may not be able to resist a thought excursion into other fields, such as biochirality (see, *e.g.*, Blackmond 2010) among others, which is the reason why we choose the terminology “chirality” instead of “parity”, which may be thought to be associated with the symmetry of fundamental physical laws, or pure geometrical symmetry, *i.e.*, mirror symmetry, of objects, or “polarization” which is used more for (linear) waves.

In conclusion, Appendix A is written thanks to a referee who raised the questions, believed in their popularity among readers and asked for explicit answers in the paper, and who is also acknowledged for the decompression of the early version of the manuscript. This work was partially supported by the Fundamental Research Funds for the Central Universities of China and by the WCI Program of the NRF of Korea [WCI 2009-001]. We thank for the discussions with U. Frisch and S. Kurien on helical hydrodynamic closures and with M. Taylor on direct numerical simulations of helical absolute equilibria back to half a solar cycle ago, with Z.-B. Guo, Z.-W. Xia and D.-D. Zou on plasma waves, with X.-P. Hu and Z. Lin on plasma heating, and the correspondences with L. Biferale, C.-K. Chan, P. Diamond, D. Escande, M. Faganello, S. Galtier, J. Miller, M. Petitjean and V. P. Ruban, A. Schwartz during the course of this work. is grateful for the hospitality of the International Institute for Fusion Science, Université de Provence, and for the workshop “The Solar Course, the Chemic Force, and the Speeding Change of Water” at NORDITA (2011).

Appendix A. Very basic aspects of the truncated system

A.1. On the detailed conservation laws for the pure helical modes in each interacting triad

Let us outline here, for the HD case, a direct verification (*e.g.*, Waleffe 1992) and a proof (*e.g.*, Kraichnan 1973) of the detailed conservation laws for energy and helicity of the pure helical modes among each interacting triad. Both of them are simply carried over from those for the traditional Fourier modes. The direct *verification* starts from the dynamical equation of the pure helical modes, with the interactions restricted among only one triad as given by Eq. (9) of Waleffe (1992). As he shows, simple algebras by the definitions of energy and helicity using this equation

then verify the conservations of energy and helicity of the three conjugate pairs of pure helical modes, regardless the handedness of any chiroid. The alternative *proof* also needs only to change the objects of the classical Fourier modes, in the third paragraph in p. 748 of Kraichnan (1973), to pure helical modes. The idea is simply that the overall energy and helicity are formally conserved by the original dynamics without explicit truncation and the truncated modes' energy and helicity are constantly zero, due to the facts that their amplitudes are set to be nulls by definition of truncation and that the 'energy and helicity expressions are quadratic and diagonal in the wave-vector amplitudes.' Note that the expression being diagonal in the wave-vector amplitudes is also important: Suppose it is not diagonal and that the convolution involves the multiplication of modes in the truncated and un-truncated domains, then the change rate of it is not assured to be constantly zero, since the change rate constitutes a component from the multiplication of the time derivative of a mode in the truncated domain with another mode in the un-truncated domain, both of which can be non-zero; an example is the quadratic invariants of 2D gyrokinetics in the Fourier-Hankel/Bessel representation and truncation, where the phase-space "wave-vector" is extended from the conventional wave-vector to include a component from the spectral representation of the velocity variable and where those quadratic expressions not being diagonal in the extended wave-vector are not ruggedly conserved, *i.e.*, not invariant after Fourier-Hankel/Bessel truncation (see pp. 3–4 of Zhu 2011).

As some readers may feel easier to start with a degenerate trivial case to get somewhat more concrete grasp, let's suppose first that we retain only contributions from the region of wave-number space $|\mathbf{k}| < K$ (Galerkin truncation), and that we start with a single triad of pure helical modes (chiroids), using indexed c to denote the chirality of the leg, $\{[\pm\mathbf{k}, c_k]; [\pm\mathbf{p}, c_p]; [\pm\mathbf{q}, c_q]\}$ with $\mathbf{k} + \mathbf{p} + \mathbf{q} = 0$ and $K/2 < |\mathbf{k}|, |\mathbf{p}|, |\mathbf{q}| < K$, so that harmonics generated by the Euler equations are eliminated under this truncation which we can think of as providing some kind of 'artificial dynamics' and which is denoted by the index " g ". Let $\mathbf{u}_g(\mathbf{x}, t)$ and $\boldsymbol{\omega}_g(\mathbf{x}, t) = \nabla \times \mathbf{u}_g(\mathbf{x}, t)$ be the velocity and vorticity fields evolving under this artificial dynamics. Then the claim is that the mean kinetic energy $\mathcal{E}_g = \langle \mathbf{u}_g^2/2 \rangle$ and mean helicity $\mathcal{H}_g = \langle \mathbf{u}_g \cdot \boldsymbol{\omega}_g \rangle$ are invariant in time. Yet another degenerate case is the Arnold-Beltrami-Childress (ABC) flow, composed of three conjugate pairs of pure helical modes with the same wavelength, which has been used by many authors to study kinematic dynamo and which can be generalized to contain more conjugate pairs of pure helical modes of same wavelengths and that presumably to become more chaotic, in the Lagrangian/streamline sense (see, *e.g.*, Arnold & Khesin 1998, and references therein).

A.2. Dynamical and topological aspects

By definition, the Galerkin-truncation dynamics of vorticity is

$$\partial_t \boldsymbol{\omega}_g = [\nabla \times (\mathbf{u}_g \times \boldsymbol{\omega}_g)]_g. \quad (\text{A } 1)$$

We are not sure whether the fact that \mathcal{H}_g is constant assures a fictitious meaningful $\tilde{\mathbf{v}}$ to solve $\partial_t \boldsymbol{\omega}_g = \nabla \times (\tilde{\mathbf{v}} \times \boldsymbol{\omega}_g)$, *i.e.*, making an analogue of the Kelvin-Helmholtz or "frozen-in" theorem (which is sufficient but not necessary for the conservation law.) Definitely, $\tilde{\mathbf{v}} = \mathbf{u}_g$ is not the solution, otherwise the last index " g " on the right-hand side of Eq. (A 1) would have no effect.

Note that the topological interpretation of the helicity as the degree of (average) knottedness and/or linkage (Moffatt 1969) of (closed) field line(s) formally carries over to the Galerkin truncated case. Actually, the interpretation itself has not much to do with the dynamics but simply works for fields satisfying some basic properties by the definition of Gauss linking number, which has a lot of subtleties and complications when generalized to continuous fields and general boundary conditions (see, *e.g.*, Moffatt 1969; Berger & Field 1984; Arnold & Khesin 1998, among many other references cited therein and appearing later). The Galerkin-truncated dynamics is formally changed much in physical space, as partly shown in the last paragraph, while

formally unchanged (except for truncation) in Fourier space concerning triadic interactions. One formally unchanged thing, besides those such as the quadratic invariants, that clearly bridges the physical- and Fourier-space representation, is the preservation of the incompressibility of the fields, which is also due to the fact that the orthogonality between their chiroids and the wavevectors holds in detail, *i.e.*, for each chiroid $\mathbf{k} \cdot \hat{\mathbf{h}}_c(\mathbf{k}) = 0$, and which justifies the definition of flux tube(s) as the key to the topological interpretation (Moffatt 1969). Topological, especially knot-theory, approaches to the statistical dynamics of course deserves further pursue, which is however beyond the scope of this note.

A.3. On the tacit assumptions in the statistical consideration

It is difficult and quite open to establish mathematically rigorous conditions for justifying the application of statistical mechanics. For example, according to literatures (see, *e.g.*, Eyink & Sreenivasan 2006, and references therein), ergodicity, being sufficient, may not be trivially satisfied but may neither be necessary; and, the mixing time scale could be hard to estimate for evaluating the closeness of physical relevance of the equilibrium ensemble. However there is a trivial bottom line that is assumed to be met, that is, all modes should be directly or indirectly connected by forming the interacting triads to define a system. For example, now suppose instead that we start with two triads $\{[\pm\mathbf{k}_j, c_{\mathbf{k}_j}]; [\pm\mathbf{p}_j, c_{\mathbf{p}_j}]; [\pm\mathbf{q}_j, c_{\mathbf{q}_j}]\}$ with $j = 1, 2$ and all these wave-vectors in the spherical annulus $(K/2, K)$, as in Appendix A.1, and suppose that these triads are non-interacting. Let the energies in the triads be E_{g_1} and E_{g_2} respectively, and the helicities H_{g_1} and H_{g_2} . Then it is not eligible to use the ensemble defined by the total energy $E_g = E_{g_1} + E_{g_2}$ and the total helicity $H_g = H_{g_1} + H_{g_2}$ for the union of these two isolated systems, not to mention that sufficient number of modes are necessary for a statistical consideration, and in particular the application of Gibbs ensemble. Actually, in practice, when the number modes is large it is hardly possible for any triad to be isolated from others. In performing the calculations as in the main text, such tacit assumptions are made to exclude cases not describable by the canonical ensemble. In the 1970s, people already performed many numerical simulations, mostly for 2D cases (Orszag 1977; Kraichnan & Montgomery 1981), to study the ergodicity and mixing properties, to measure the difference between microcanonical and canonical ensembles, to find how many modes would be needed to reach the Gibbs state and to finally verify the corresponding energy spectrum. Matthaeus and collaborators, among others, have many followup studies, especially for 2D and 3D MHD absolute equilibrium ensembles. We cannot exhaust the list of references here, but would like to remark that there can be new findings by revisiting numerical check of the tacit assumptions relevant to the application of Gibbs ensembles with the new special truncations on the new freedoms of chiroids.

REFERENCES

- ARNOLD, V. I. & KHESIN, B. A. 1998 *Topological Methods in Hydrodynamics*. Springer.
- BARRON, L. D. 1997 From cosmic chirality to protein structure and function: Lord Kelvin's legacy. *Q. J. Med.* **90** 793–800.
- BERGER, M. A. & FIELD, G. B. 1984 The topological properties of magnetic helicity. *J. Fluid Mech.* **147**, 133–148.
- BETCHOV, R. 1961 Semi-isotropic turbulence and helicoidal flows. *Phys. Fluids* **4**, 925–926.
- BIFERALE, L., MUSACCHIO, S. & TOSCHI, F. 2012 Inverse energy cascade in three-dimensional isotropic turbulence. *Phys. Rev. Lett.* **108**, 104501–104504.
- BISKAMP, D., SCHWARZ, E., ZEILER, A. CELANI & DRAKE J. F. 1999 Electron magnetohydrodynamic turbulence. *Phys. Plasmas* **6**, 751–758.
- BLACKMOND, D. G. 2010 The Origin of Biological Homochirality *Cold Spring Harb Perspect Biol* **2**, a002147.
- BRANDENBURG, A., DOBLER W. & SUBRAMANIAN, K. 2002 Magnetic helicity in stellar dynamos: new numerical experiments. *Astron. Nachr.* **323**, 99–123.

- BRANDENBURG, A. & SUBRAMANIAN, K. 2005 Astrophysical magnetic fields and nonlinear dynamo theory. *Physics Reports* **417**, 1–209.
- BRISSAUD, A., FRISCH, U., LEORAT, J., LESIEUR, M. & MAZURE, M. 1973 Helicity cascades in isotropic turbulence. *Phys. Fluids* **16**, 1366–1367.
- CHEN, Q. N., CHEN, S. Y. & EYINK, G. L. 2003 The joint cascade of energy and helicity in three-dimensional turbulence. *Phys. Fluids* **15** (2), 361–374.
- CHEN, Q. N., CHEN, S. Y., EYINK, G. L. & HOLM, D. 2005 Resonant interactions in rotating homogeneous three-dimensional turbulence. *J. Fluid Mech.* **542** (2), 139–163.
- CHO, J. 2011 Magnetic Helicity Conservation and Inverse Energy Cascade in Electron Magnetohydrodynamic Wave Packets. *Phys. Rev. Lett.* **106**, 191104–191107.
- CINTAS, P. & VIEDMA, C. 2012 On the Physical Basis of Asymmetry and Homochirality. *Chirality* **24**, 894–908.
- DITLEVSEN, P. D. & GIULIANI, P. 2001 Dissipation in helical turbulence. *Phys. Fluids* **13**, 3508–3509.
- EYINK, G. L. 2008 Dissipative anomalies in singular Euler flows. *Physica D* **237**, 1956–1968.
- EYINK, G. L. & SREENIVASAN, K. R. 2006 Onsager and the theory of turbulence. *Review of Modern Physics* **78**, 87–135.
- FOUXON, A. & LEBEDEV, V. 2003 Spectra of turbulence in dilute polymer solutions. *Phys. Fluids* **15**, 2060–2072.
- FRISCH, U., POUQUET, A., LEORAT, J. & MAZURE, A. 1975 Possibility of an inverse magnetic helicity cascade in magnetohydrodynamic turbulence. *J. Fluid Mech.* **68**, 769–778.
- FRISCH, U., KURIEN, S., PANDIT, R., PAULS, W., RAY, S., WIRTH, A. & ZHU, J.-Z. 2008 Hyperviscosity, Galerkin Truncation and Bottleneck of Turbulence. *Phys. Rev. Lett.* **101**, 114501–114504.
- FRISCH, U. 1995 *Turbulence: The Legacy of Kolmogorov*. Cambridge University Press.
- GALTIER, S. 2006 Wave turbulence in incompressible Hall magnetohydrodynamics. *J. Plasma Physics* **72**, 721–769.
- GALTIER, S. & BHATTACHARJEE, A. 2003 Anisotropic weak whistler wave turbulence in electron magnetohydrodynamics. *Phys. Plasmas* **10**, 3065–3076.
- HATFIELD, J. W. & QUAKE, S. R. 1999 Dynamic Properties of an Extended Polymer in Solution. *Phys. Rev. Lett.* **82**, 3548–3551.
- Ji, H. 1999 Turbulent dynamos and magnetic helicity. *Phys. Rev. Lett.* **83**, 3198–3191.
- KELVIN, W. THOMSON 1904 *Baltimore Lectures on Molecular Dynamics and the Wave Theory of Light*. C. J. Clay and Sons, Cambridge University Press Warehouse: London.
- KRAICHNAN, R. H. 1967 Inertial ranges in two-dimensional turbulence. *Phys. Fluids* **10**, 1417–1423.
- KRAICHNAN, R. H. 1973 Helical turbulence and absolute equilibrium. *J. Fluid Mech.* **59**, 745–752.
- KRAICHNAN, R. H. 1958 Irreversible Statistical Mechanics of Incompressible Hydromagnetic Turbulence. *Phys. Rev.* **109**, 1407–1422.
- KRAICHNAN, R. H. & MONTGOMERY, D. 1980 Two-dimensional turbulence. *Reports Progr. Phys.* **43**, 547–619.
- LEE, T.-D. 1952 On some statistical properties of hydrodynamic and hydromagnetic fields. *Q. Appl. Math.* **10**, 69–74.
- LESSINNES, T., PLUNIAN F. & CARATI, D. 2009 Helical shell models for MHD. *Theor. Comput. Fluid Dyn.* **23**, 439–450.
- LESIEUR, M. 1990 *Turbulence in Fluids*. 2nd ed. Kluwer Academic, Dordrecht, The Netherlands.
- L'VOV, V. S., POMYALOV, A. & AND PROCACCIA, I. 2002 Quasi-Gaussian Statistics of Hydrodynamic Turbulence in $4/3 + \epsilon$ Dimensions. *Phys. Rev. Lett.* **89**, 064501–064504.
- MANDELBROT, B. B. 1974 Intermittent turbulence in self-similar cascades: divergence of high moments and dimension of the carrier. *Journal of Fluid Mechanics* **62**, 331–358.
- MELANDER, V., HUSSAIN, F. 1993 Polarized vorticity dynamics in a vortex column. *Phys. Fluids A* **5**, 1992–2003.
- MENEGUZZI, M., FRISCH, U AND POUQUET, A 1981 Helical and nonhelical turbulent dynamos. *Phys. Rev. Lett.* **47**, 1060–1063.
- MEYRAND, M. & GALTIER, S. 2012 Spontaneous Chiral Symmetry Breaking of Hall Magnetohydrodynamic Turbulence. *Phys. Rev. Lett.* **109**, 194501–1–5.
- MILLER, J. 1990 Statistical mechanics of Euler equations in two dimensions. *Phys. Rev. Lett.* **65**, 2137–2140.
- MOFFATT, H. K. 1969 The degree of knottedness of tangled vortex lines. *J. Fluid Mech.* **35**, 117–129.

- MOFFATT, H. K. 1970 Turbulent dynamo action at low magnetic Reynolds number. *J. Fluid Mech.* **41**, 435–452.
- MOFFATT, H. K. 2008 Vortex Dynamics: The Legacy of Helmholtz and Kelvin. In *Hamiltonian Dynamics, Vortex Structures, Turbulence* (ed. A.V. Borisov, V.V. Kozlov, I.S. Mamaev & M.A. Sokolovskiy), pp. 113–138. Springer.
- MOSES, H. E. 1971 Eigenfunctions of the curl operator, rotationally invariant Helmholtz theorem and applications to electromagnetic theory and fluid mechanics. *SIAM (Soc. Ind. Appl. Math.) J. Appl. Math.* **21**, 114–130.
- PEREZ, J. C., MASON, J., BOLDYREV, S. & CATTANEO, F. 2012 On the Energy Spectrum of Strong Magnetohydrodynamic Turbulence. *Phys. Rev. X* **2**, 041005.
- PETITJEAN, M. 2003 Chirality and Symmetry Measures: A Transdisciplinary Review. *Entropy* **5**, 271–312.
- POUQUET, A., FRISCH, U. & LÉORAT, J. 1976 Strong MHD helical turbulence and the nonlinear dynamo effect. *J. Fluid Mech.* **77**, 321–354.
- PROCACCIA, I., L'VOV, V. & BENZI, R. 2008 Colloquium: Theory of Drag Reduction by Polymers in Wall Bounded Turbulence. *Rev. Mod. Phys.* **80**, 225–247.
- ORSZAG, S. A. 1977 Statistical Theory of Turbulence, in *Fluid Dynamics, Les Houches 1973*, 237–374, eds. R. Balian & J.L. Peube. Gordon and Breach, New York.
- ROBERT, R. 2003 Statistical Hydrodynamics (Onsager Revisited). in: *Handbook of mathematical fluid dynamics*, Volume 2, Eds. Susan Friedlander, Denis Serre, pp. 1–54. Gulf Professional Publishing.
- STEINBERG, V. 2009 Elastic stresses in random flow of a dilute polymer solution and the turbulent drag reduction problem. *C. R. Physique* **10**, 728–739.
- STIX, T. H. 1992 *Waves in Plasmas*. American Institute of Physics.
- RUBAN, V. P. 1999 Motion of magnetic flux lines in magnetohydrodynamics. *Sov. Phys. JETP* **89**, 299–310.
- SCHEKOCHIHIN, A. A., COWLEY, S. C., DORLAND, W., HAMMETT, G. W., HOWES, G. G., QUATAERT, E. & TATSUNO, T. 2009 Astrophysical gyrokinetics: Kinetic and fluid cascades in magnetized weakly collisional plasmas. *The Astrophysical Journal Supplement Series* **182**, 310–377.
- STENZEL, R. L. 1999 Whistler waves in space and laboratory plasmas. *J. Geophys. Res.* **14**, 14379–14395.
- WALEFFE, F. 1992 The nature of triad interactions in homogeneous turbulence. *Phys. Fluids A* **4**, 350–363.
- WOLTJER, L. 1959 Hydromagnetic equilibrium II. Stability in the variational formulation. *Proc. Nat. Acad. Sci. U.S.A.* **45** 769–771.
- YAMADA, H., KATANO, T., KANAI, K., ISHIDA, A. & STEINHAEUER, L. 2002 Equilibrium analysis of a flowing two-fluid plasma. *Phys. Plasmas* **11**, 4605–4614.
- YANG, Y.-T., SU, W.-D. & WU, J.-Z. 2010 Helical-wave decomposition and applications to channel turbulence with streamwise rotation. *J. Fluid Mech.* **662**, 91–122.
- ZHU, J.-Z. 2011 Fourier-Hankel/Bessel space absolute equilibria of 2D gyrokinetics. arXiv:1109.2511v1 [nlin.CD]: under revision for later journal publication.
- ZHU, J.-Z. & HAMMETT, G. W. 2010 Gyrokinetic absolute equilibrium and turbulence. *Phys. Plasmas* **17**, 122307-1–13.
- ZHU, J.-Z. & TAYLOR, M. 2010 Intermittency and Thermalization of Turbulence. *Chinese Phys. Lett.* **27**, 054702–054705.