

A variant of nonsmooth maximum principle for state constrained problems

Md. Haider Ali Biswas and M.d.R. de Pinho

Abstract— We derive a variant of the nonsmooth maximum principle for problems with pure state constraints. The interest of our result resides on the nonsmoothness itself since, when applied to smooth problems, it coincides with known results. Remarkably, in the normal form, our result has the special feature of being a sufficient optimality condition for linear-convex problems, a feature that the classical Pontryagin maximum principle had whereas the nonsmooth version had not. This work is distinct to previous work in the literature since, for state constrained problems, we add the Weierstrass conditions to adjoint inclusions using the joint subdifferentials with respect to the state and the control. Our proofs use old techniques developed in [16], while appealing to new results in [7].

I. INTRODUCTION

It is commonly accepted that optimal control appears with the publication of the seminal book [14] where the statement and proof of the Pontryagin Maximum Principle played a crucial role (we refer the reader to the survey [13] for an interesting historic account of the pioneering results). Since then we have witnessed continuous developments.

Generalization of the classical maximum principle to problems with nonsmooth data appeared in 1970's as mainly the result of the work of Francis Clarke (see [3] and references therein). The nonsmooth maximum principle, nowadays a well established result, was then extended and refined by a number of authors. One of the first attempts to extend it to cover problems with state constraints came up in [16].

A special feature of the classical Pontryagin maximum principle is that it is also a sufficient optimality condition for the normal form of the so called linear-convex problems. Regrettably, the nonsmooth version had no such feature. Nonsmooth necessary optimality conditions in the vein of maximum principles were proposed in [8] overcoming this setback. Regrettably those necessary conditions did not include the Weierstrass condition responsible for the very name Maximum Principle. More recently the setbacks in [8] were taken care of in [6] where a new variant of the nonsmooth maximum principle is derived by appealing to [5]. As in [8], Lipschitz continuity of dynamics with respect to both state and control is assumed, the special ingredient responsible for sufficiency of the nonsmooth maximum principle when

applied to normal linear convex problems (see problem (LC) below)¹. In what follows, and for simplicity, we opt to refer to the statement of this new nonsmooth maximum principle stated as Theorem 3.1 in [7] which plays a crucial role in our developments.

Here we extend Theorem 3.1 in [7] to cover state constrained problems. In doing so we follow closely the approach of [9] and [10] where the main result in [8] is generalized to cover state constrained problems in two steps; first the convex case is treated in [9] using techniques based on [16] and then convexity is removed in [10].

In this paper we show that the proofs in [9] and [10] adapted easily to allow extension of Theorem 3.1 in [7] to state constrained problems. In this way we obtain a new variant of the nonsmooth maximum principle, improving on [10] by adding the Weierstrass condition to the previous conditions while keeping the interesting feature of being a sufficient condition for normal linear-convex problems.

II. PRELIMINARIES

A. Notation

Here and throughout \mathbb{B} represents the closed unit ball centered at the origin regardless of the dimension of the underlying space and $|\cdot|$ represents the Euclidean norm or the induced matrix norm on $\mathbb{R}^{p \times q}$. The *Euclidean distance function* with respect to a given set $A \subset \mathbb{R}^k$ is

$$d_A: \mathbb{R}^k \rightarrow \mathbb{R}, \quad y \mapsto d_A(y) = \inf \{|y - x| : x \in A\}.$$

A function $h: [0, 1] \rightarrow \mathbb{R}^p$ lies in $W^{1,1}([0, 1]; \mathbb{R}^p)$ if and only if it is absolutely continuous; in $L^1([0, 1]; \mathbb{R}^p)$ iff it is integrable; and in $L^\infty([0, 1]; \mathbb{R}^p)$ iff it is essentially bounded. The norm of $L^1([0, 1]; \mathbb{R}^p)$ is denoted by $\|\cdot\|_1$ and the norm of $L^\infty([0, 1]; \mathbb{R}^p)$ is $\|\cdot\|_\infty$.

We make use of standard concepts from nonsmooth analysis. Let $A \subset \mathbb{R}^k$ be a closed set with $\bar{x} \in A$. The *limiting normal cone to A at \bar{x}* is denoted by $N_A(\bar{x})$.

Given a lower semicontinuous function $f: \mathbb{R}^k \rightarrow \mathbb{R} \cup \{+\infty\}$ and a point $\bar{x} \in \mathbb{R}^k$ where $f(\bar{x}) < +\infty$, $\partial f(\bar{x})$ denotes the *limiting subdifferential* of f at \bar{x} . When the function f is Lipschitz continuous near x , the convex hull of the limiting subdifferential, $\text{co } \partial f(x)$, coincides with the (*Clarke*) *subdifferential*. Properties of Clarke's subdifferentials (upper semi-continuity, sum rules, etc.), can be found in [4]. For details on such nonsmooth analysis concepts, see [4], [15], [17] and [12].

¹ With respect to generalizations of [8] we also refer the reader to a different version of a nonsmooth maximum principle in [1] making use of "compatible" feedback controls.

This work has been supported by the European Union Seventh Framework Programme [FP7-PEOPLE-2010-ITN] under grant agreement n264735-SADCO. The first author is supported by the grant SFRH/BD/63707/2009, FCT, Portugal.

Haider Ali Biswas is with ISR and DEEC, Faculdade de Engenharia da Universidade do Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal dee08022@fe.up.pt

Maria do Rosário de Pinho is with ISR and DEEC, Faculdade de Engenharia da Universidade do Porto, Rua Dr. Roberto Frias, 4200-465 Porto, Portugal mrpinho@fe.up.pt

B. The Problem

Consider the problem denoted throughout by (P) of minimizing

$$l(x(a), x(b)) + \int_a^b L(t, x(t), u(t)) dt$$

subject to the differential equation

$$\dot{x}(t) = f(t, x(t), u(t)) \quad \text{a.e. } t \in [a, b],$$

the state constraint

$$h(t, x(t)) \leq 0 \quad \text{for all } t \in [a, b],$$

the boundary conditions

$$(x(a), x(b)) \in C,$$

and the control constraints

$$u(t) \in U(t) \quad \text{a.e. } t \in [a, b].$$

Here the interval $[a, b]$ is fixed. We have the state $x(t) \in \mathbb{R}^n$ and the control $u(t) \in \mathbb{R}^k$. The function describing the dynamics is $f : [a, b] \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}^n$. Moreover h and L are scalar functions $h : [a, b] \times \mathbb{R}^n \rightarrow \mathbb{R}$, $L : [a, b] \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}$, U is a multifunction and $C \subset \mathbb{R}^n \times \mathbb{R}^n$.

We shall denote by (S) the problem one obtains from (P) in the absence of the state constraint $h(t, x(t)) \leq 0$ and we refer to it as a *standard optimal control problem*.

Throughout this paper we assume that the following basic assumptions are in force:

- B1** the functions L and f are $\mathcal{L} \times \mathcal{B}$ -measurable,
- B2** the multifunction U has $\mathcal{L} \times \mathcal{B}$ -measurable graph,
- B3** the set C is closed and l is locally Lipschitz.

For (P) (or (S)) a pair (x, u) comprising an absolutely continuous function x , the state, and a measurable function u , the control, is called an *admissible process* if it satisfies all the constraints.

An admissible process (x^*, u^*) is a *strong local minimum* of (P) (or (S)) if there exists $\varepsilon > 0$ such that (x^*, u^*) minimizes the cost over all admissible processes (x, u) such that

$$|x(t) - x^*(t)| \leq \varepsilon \quad \text{for all } t \in [a, b]. \quad (1)$$

It is a *local $W^{1,1}$ -minimum* if there exists some $\varepsilon > 0$ such that it minimizes the cost to all processes (x, u) satisfying (1) and

$$\int_a^b |\dot{x}(t) - \dot{x}^*(t)| dt \leq \varepsilon.$$

Let $R : [a, b] \rightarrow]0, +\infty]$ be a given measurable function. Then the admissible process (x^*, u^*) is a *local minimum of radius R* if it minimizes the cost over all admissible processes (x, u) such that

$$|x(t) - x^*(t)| \leq \varepsilon, \quad |u(t) - u^*(t)| \leq R(t) \quad \text{a.e.}$$

$$\text{and } \int_a^b |\dot{x}(t) - \dot{x}^*(t)| dt \leq \varepsilon$$

for some $\varepsilon > 0$.

C. Assumptions

In what follows the pair (x^*, u^*) will always denote the solution of the optimal control problem under consideration.

Let us take any function ϕ defined in $[a, b] \times \mathbb{R}^n \times \mathbb{R}^k$ and taking values in \mathbb{R}^n or \mathbb{R} .

A1 There exist constants k_x^ϕ and k_u^ϕ for almost every $t \in [a, b]$ and every (x_i, u_i) ($i = 1, 2$) such that

$$x_i \in \{x : |x - x^*(t)| \leq \varepsilon\}, \quad u_i \in U(t)$$

we have

$$|\phi(t, x_1, u_1) - \phi(t, x_2, u_2)| \leq k_x^\phi |x_1 - x_2| + k_u^\phi |u_1 - u_2|.$$

A2 The set valued function $t \rightarrow U(t)$ is closed valued and there exists a constant $c > 0$ such that for almost every $t \in [a, b]$ we have

$$|u(t)| \leq c \quad \forall u \in U(t).$$

When **A1** is imposed on f and/or L , then the Lipschitz constants are denoted by k_x^f , k_u^f , k_x^L and k_u^L . Observe that if U is independent of time, then **A2** states that the set U is compact. Assumption **A2** requires the controls to be bounded, a strong hypothesis but nevertheless quite common in applications. It also simplifies the proofs of the forthcoming results where limits of sequence of controls needed to be taken.

D. Auxiliary Results

Attention now goes to problem (S) , i.e., we assume that the state constraint is now absent. We next state an adaptation of Theorem 3.1 in [7] essential to our analysis in the forthcoming sections. It is ‘‘an adaptation’’ because it holds under stronger assumptions than those in [7].

Theorem 2.1: Let (x^*, u^*) be a strong local minimum for problem (S) . If **B1–B3** are satisfied, f and L satisfy **A1** and U is closed valued, then there exist $p \in W^{1,1}([a, b]; \mathbb{R}^n)$ and a scalar $\lambda_0 \geq 0$ satisfying the nontriviality condition [NT]:

$$\|p\|_\infty + \lambda_0 > 0,$$

the Euler adjoint inclusion [EI]:

$$\begin{aligned} (-\dot{p}(t), 0) \in \partial_{x,u}^C (\langle p, f \rangle - \lambda_0 L)(t, x^*(t), u^*(t)) \\ - \{0\} \times K |p(t)| \partial_u^C d_{U(t)}(u^*(t)) \quad \text{a.e.}, \end{aligned}$$

the global Weierstrass condition [W]:

$$\forall u \in U(t),$$

$$\begin{aligned} \langle p(t), f(t, x^*(t), u) \rangle + \lambda_0 L(t, x^*(t), u) \leq \\ \langle p(t), f(t, x^*(t), u^*(t)) \rangle + \lambda_0 L(t, x^*(t), u^*(t)) \quad \text{a.e.}, \end{aligned}$$

and the transversality condition [T]:

$$(p(a), -p(b)) \in N_C^L(x^*(a), x^*(b)) + \lambda_0 \partial^L l(x^*(a), x^*(b)).$$

In the above K is a constant depending merely on k_x^f , k_x^L , k_u^f and k_u^L .

In [7] the analysis is done for *local minimum of radius R* instead of strong minimum and it holds under a weaker assumption than **A1**.

We point out that the conditions given by the classical nonsmooth maximum principle (see [5]) are [NT], [W], [T] and [EI] is replaced by

$$-\dot{p}(t) \in \partial_x^C \left(\langle p(t), f(t, x^*(t), u^*(t)) \rangle - \lambda_0 L(t, x^*(t), u^*(t)) \right). \quad (2)$$

We refer the reader to [6] for a discussion on (2) and [EI].

III. MAIN RESULTS

We now turn to problem (P). We derive a new nonsmooth maximum principle for this state constrained problem in the vein of Theorem 3.1 in [7] in two stages. Firstly the result is established under a convexity assumption on the “velocity set” (see **C** below). Then such hypothesis is removed. This is proved following an approach in [17] and similar to what is done in [10].

On h we impose the following:

A3 For all x such that $|x(t) - x^*(t)| \leq \varepsilon$ the function $t \rightarrow h(t, x)$ is continuous. Furthermore, there exists a constant $k_h > 0$ such that the function $x \rightarrow h(t, x)$ is Lipschitz of rank k_h for all $t \in [a, b]$.

The need to impose continuity of $t \rightarrow h$ instead of merely semi upper continuity is discussed in [9].

Recall that our basic assumptions **B1–B3** are in force. Suppose that f and L satisfy **A1** and that **A2** holds. For future use, observe that these assumptions also assert that following conditions are satisfied:

$$\begin{aligned} |\phi(t, x^*(t), u) - \phi(t, x^*(t), u^*(t))| &\leq \\ k_u^\phi |u - u^*(t)| &\text{ for all } u \in U(t) \text{ a.e. } t \end{aligned} \quad (3)$$

and there exists an integrable function k such that

$$|\phi(t, x^*(t), u)| \leq k(t) \text{ for all } u \in U(t) \text{ a.e. } t. \quad (4)$$

In the above ϕ is to be replaced by f and L . Moreover, it is a simple matter to see that the sets $f(t, x, U(t))$ and $L(t, x, U(t))$ are compact for all $x \in x^*(t) + \varepsilon \mathbb{B}$.

A. Convex Case

Consider the additional assumption on the “velocity set”:

C The velocity set

$$\{(v, l) = (f(t, x, u), L(t, x, u)), u \in U(t)\}$$

is convex for all $(t, x) \in [a, b] \times \mathbb{R}^n$.

Introduce the following subdifferential

$$\begin{aligned} \bar{\partial}_x h(t, x) &:= \\ \text{co}\{\lim \xi_i : \xi_i \in \partial_x h(t_i, x_i), (t_i, x_i) \rightarrow (t, x)\}. \end{aligned} \quad (5)$$

Proposition 1: Let (x^*, u^*) be a strong local minimum for problem (P). Assume that f and L satisfy **A1**, assumptions **B1–B3**, **A2** and **C** hold and h satisfies **A3**. Then there exist $p \in W^{1,1}([a, b]; \mathbb{R}^n)$, $\gamma \in L^1([a, b]; \mathbb{R})$, a measure $\mu \in C^\oplus([a, b]; \mathbb{R})$, and a scalar $\lambda_0 \geq 0$ satisfying

$$(i) \mu\{[a, b]\} + \|p\|_\infty + \lambda_0 > 0,$$

$$(ii) (-\dot{p}(t), 0) \in \partial_{x,u}^C \left(\langle q(t), f(t, x^*(t), u^*(t)) \rangle - \lambda_0 L(t, x^*(t), u^*(t)) \right) - \{0\} \times N_{U(t)}^C(u^*(t)) \text{ a.e.,}$$

$$(iii) \forall u \in U(t), \langle q(t), f(t, x^*(t), u) \rangle - \lambda_0 L(t, x^*(t), u) \leq \langle q(t), f(t, x^*(t), u^*(t)) \rangle - \lambda_0 L(t, x^*(t), u^*(t)) \text{ a.e.,}$$

$$(iv) (p(a), -q(b)) \in N_C^L(x^*(a), x^*(b)) + \lambda_0 \partial l(x^*(a), x^*(b)),$$

$$(v) \gamma(t) \in \bar{\partial} h(t, x^*(t)) \quad \mu\text{-a.e.,}$$

$$(vi) \text{supp}\{\mu\} \subset \{t \in [a, b] : h(t, x^*(t)) = 0\},$$

where

$$q(t) = \begin{cases} p(t) + \int_{[a,t)} \gamma(s) \mu(ds) & t \in [a, b) \\ p(t) + \int_{[a,b]} \gamma(s) \mu(ds) & t = b. \end{cases} \quad (6)$$

B. Maximum Principle in the Nonconvex Case

Now we replace the subdifferential $\bar{\partial}_x h$ by a more refined subdifferential $\partial_x^> h$ defined by

$$\begin{aligned} \partial_x^> h(t, x) &:= \text{co}\{\xi : \exists (t_i, x_i) \xrightarrow{h} (t, x) : \\ h(t_i, x_i) &> 0 \forall i, \partial_x h(t_i, x_i) \rightarrow \xi\}. \end{aligned} \quad (7)$$

Theorem 3.1: Let (x^*, u^*) be a strong local minimum for problem (P). Assume that f and L satisfy **A1**, h satisfies **A3** and that **A2** as well as the basic assumptions **B1–B3** hold. Then there exist an absolutely continuous function p , an integrable function γ , a non-negative measure $\mu \in C^\oplus([a, b]; \mathbb{R})$, and a scalar $\lambda_0 \geq 0$ such that conditions (i)–(vi) of Proposition 1 hold with $\partial_x^> h$ as in (7) replacing $\bar{\partial}_x h$ and where q is as defined in (6).

For the convex case see [2] for preliminary results for problems with additional mixed state control constraints. Removal of convexity will be the focus of future work.

The above theorem adapts easily when we assume (x^*, u^*) to be a weak local minimum instead of a strong local minimum (see discussion above). It is sufficient to replace $U(t)$ by $U(t) \cap \mathbb{B}_\varepsilon(u^*(t))$.

Theorem 3.1 can now be extended to deal with a local $W^{1,1}$ -minimum for (P).

Theorem 3.2: Let (x^*, u^*) be merely a local $W^{1,1}$ -minimum for problem (P). Then the conclusions of Theorem 3.1 hold.

We omit the proof of this Theorem here since it can be easily obtained mimicking what is done in [17].

C. Linear Convex Problems

The distinction between Theorem 3.1 and classical nonsmooth maximum principle (see [17]) is well illustrated by an example provided in [9]. We recover such example here

showing that Theorem 3.1 can eliminate processes whereas the classical nonsmooth maximum principle cannot.

Example: Consider the problem on the interval $[0, 1]$:

$$(L) \begin{cases} \text{Minimize } \int_0^1 (w_1|x - u_1| + w_2|x - u_2| + x)dt \\ \text{subject to} \\ \dot{x}(t) = 4w_1(t)u_1(t) + 4w_2(t)u_2(t) \text{ for a. e. } t, \\ x(t) \geq -1 \text{ for all } t, \\ u_1(t), u_2(t) \in [-1, 1] \text{ for a. e. } t, \\ (w_1(t), w_2(t)) \in W \text{ for a. e. } t, \\ x(0) = 0 \end{cases}$$

where

$$W := \{(w_1, w_2) \in \mathbb{R}^2 : w_1, w_2 \geq 0, w_1 + w_2 = 1\}.$$

The process $(x^*, u_1^*, u_2^*, w_1^*, w_2^*) := (0, 0, 0, 1, 0)$ is an admissible process with cost 0 and along the trajectory the state constraint is inactive. It is easy to see that the classical nonsmooth maximum principle holds when we take all the multipliers 0 but $\lambda_0 = 1$. However, $(x^*, u_1^*, u_2^*, w_1^*, w_2^*)$ is not optimal. In fact, if we consider the process $(x, u_1, u_2, w_1, w_2) = (-4\alpha t, -\alpha, 0, 1, 0)$, with $\alpha \in (0, 1/4)$, we see that this process has cost $-3/4\alpha$. Now let us apply Theorem 3.1 to our problem for the process $(x^*, u_1^*, u_2^*, w_1^*, w_2^*)$. Since the state constraint is inactive, we deduce that measure μ is null. Considering the Euler Lagrange equation in (ii) of Theorem 3.1 we deduce that there should exist an absolutely continuous function p and a scalar $\lambda_0 \geq 0$ satisfying (i) of Theorem 3.1 and such that $p(1) = 0$, $-\dot{p}(t) = -\lambda_0(1 + e(t))$ and $0 = 4p(t) + \lambda_0 e(t)$ where $e(t)$ takes values in $[-1, 1]^2$. A simple analysis will convince the reader that this situation is impossible. This means that Theorem 3.1 does not hold excluding $(x^*, u_1^*, u_2^*, w_1^*, w_2^*)$ as a minimum. ■

Consider the problem

$$(LC) \begin{cases} \text{Minimize } l(x(a), x(b)) + \int_a^b L(t, x(t), u(t))dt \\ \text{subject to} \\ \dot{x}(t) = A(t)x(t) + B(t)u(t) \text{ for a. e. } t \in [a, b], \\ D(t)x(t) \leq 0 \text{ for all } t \in [a, b], \\ u(t) \in U(t) \text{ for a. e. } t \in [a, b], \\ (x(a), x(b)) \in E \end{cases}$$

where E is convex, the multifunction U is convex valued, the functions l and $(x, u) \rightarrow L(t, x, u)$ are convex, the function $A : [0, 1] \rightarrow \mathbb{R}^{n \times n}$ is integrable, the function $B : [0, 1] \rightarrow \mathbb{R}^{n \times k}$ is measurable, and the function $D : [0, 1] \rightarrow \mathbb{R}^{1 \times n}$ is continuous. Then (LC) is what we refer to as a linear convex problem with state constraints.

Theorem 3.1 (and of course Theorem 3.2) keeps the significant feature of being a sufficient condition of optimality in the normal form for problem (LC) . This follows directly from the observation that the proof of Proposition 4.1 in [9]

²The function e appears from the subdifferential of the cost which is clearly nonsmooth due to the presence of the modulus.

proves our claim. No adaptation is required in this case. For completeness we state such proposition here.

We say that a process (x^*, u^*) is a **normal** extremal if it satisfies the conclusions of Theorem 3.1 with $\lambda_0 = 1$.

Proposition 2: ([9]) If the process (x^*, u^*) is a normal extremal for problem (LC) , then it is a minimum.

Let us return to our previous example. Problem (L) is what we call a linear convex problem. It is now obvious that the process $(x^*, u_1^*, u_2^*, w_1^*, w_2^*) := (0, 0, 0, 1, 0)$ does not satisfy the conclusions of Theorem 3.1, if it did, then it would be a minimum as asserted by Proposition 2 and it is not.

IV. PROOFS OF THE MAIN RESULTS

Since our proofs are based on those in [9] and [10] we only give a brief sketch of them, refereing the reader to the appropriate literature for details.

All the results are proved assuming that $L \equiv 0$. The case of $L \neq 0$ is treated by a standard and well known technique.

A. Sketch of the Proof of Proposition 1

- First the validity of the Proposition is established for the simpler problem

$$(Q) \begin{cases} \text{Minimize } l(x(b)) \\ \text{subject to} \\ \dot{x}(t) = f(t, x(t), u(t)) \text{ a.e. } t \in [a, b] \\ u(t) \in U(t) \text{ a.e. } t \in [a, b] \\ h(t, x(t)) \leq 0 \text{ for all } t \in [a, b] \\ (x(a), x(b)) \in \{x_a\} \times E_b. \end{cases}$$

Problem (Q) is a special case of (P) in which $E = \{x_a\} \times E_b$ and $l(x_a, x_b) = l(x_b)$.

Our proof consists of the following steps

- Q1 Define a sequence of problems penalizing the state-constraint violation. The sequence of problems is

$$(Q_i) \begin{cases} \text{Minimize } l(x(b)) + i \int_a^b h^+(t, x(t)) dt \\ \text{subject to} \\ \dot{x}(t) = f(t, x(t), u(t)) \text{ a.e. } t \in [a, b] \\ (x(a), x(b)) \in \{x_a\} \times E_b, \end{cases}$$

where $h^+(t, x) := \max\{0, h(t, x)\}$.

- Q2 Assume that $\liminf_{i \rightarrow \infty} \{Q_i\} = \inf\{Q\}$.

- Q3 Set W to be the set of measurable functions $u : [a, b] \rightarrow \mathbb{R}^k$, $u(t) \in U(t)$ a.e. such that a solution of the differential equation $\dot{x}(t) = f(t, x(t), u(t))$, for almost every $t \in [a, b]$, with $x(t) \in x^*(t) + \varepsilon \mathbb{B}$ for all $t \in [a, b]$ and $x(a) = x_a$ and $x(b) \in E_b$. We provide W with the L^1 metric defined by $\Delta(u, v) := \|u - v\|_{L^1}$ and set

$$J_i(u) := l(x(b)) + i \int_a^b h^+(t, x(t)) dt.$$

Then (W, Δ) is a complete metric space in which the functional $J_i : W \rightarrow \mathbb{R}$ is continuous.

Q4 Apply Ekeland's theorem to the sequence of problems of the form

$$(O_i) \begin{cases} \text{Minimize} & J_i(u) \\ \text{subject to} & u \in W \end{cases}$$

which are closely related to (Q_i) .

The conclusion of application of Ekeland's theorem shows that (x_i, u_i) solves the following optimal control problem:

$$(E_i) \begin{cases} \text{Minimize} & l(x(b)) + i \int_a^b h^+(t, x(t)) dt + \\ & \sqrt{\varepsilon_i} \int_a^b |u(t) - u_i(t)| dt \\ \text{subject to} & \\ & \dot{x}(t) = f(t, x(t), u(t)) \text{ a.e. } t \in [a, b] \\ & u(t) \in U(t) \text{ a.e. } t \in [a, b] \\ & x(a) = x_a \\ & x(b) \in E_b. \end{cases}$$

The fact that $\varepsilon_i \rightarrow 0$ allows us to prove that u_i converges strongly to u^* and x_i converges uniformly to x^* .

Q6 Rewriting these conditions and taking limits as in [9] we get the required conclusions.

Q7 Finally we show that **C** implies **IH**.

The remaining of the proof has three stages. We first extend Proposition 1 to problems where $x(a) \in E_a$, and E_a is a closed set. This is done following the lines in the end of the proof of Theorem 3.1 in [16].

Next we consider the case when the cost is $l = l(x(a), x(b))$. This is done using the technique in Step 2 of section 6 in [11]. And finally, following again the approach in section 6 in [11], we derive necessary conditions when $(x(a), x(b)) \in E$, completing the proof.

In order to proof our result, an important piece of analysis added to the proof of Theorem 3.1 in [9] concerns the Weierstrass condition (iii) of Proposition 1. The information extracted while taking limits allow us to do that without that much ado.

B. Sketch of the Proof of Theorem 3.1

We now proceed to prove our main Theorem 3.1. We recall that under our hypotheses both (3) and (4) hold and that the set $f(t, x, U(t))$ is compact.

Our proof consists of several steps. We first consider the following 'minimax' optimal control problem where the state constraint functional $\max_{t \in [a, b]} h(t, x(t))$ appears in the cost.

$$(\tilde{R}) \begin{cases} \text{Minimize} & \tilde{l}(x(a), x(b), \max_{t \in [a, b]} h(t, x(t))) \\ \text{over} & x \in W^{1,1} \text{ and measurable } u \text{ satisfying} \\ & \dot{x}(t) = f(t, x(t), u(t)) \text{ a.e. } t \in [a, b] \\ & u(t) \in U(t) \text{ a.e. } t \in [a, b] \\ & (x(a), x(b)) \in E_a \times \mathbb{R}^n. \end{cases}$$

where $\tilde{l} : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ is a given function and $E_a \subset \mathbb{R}^n$ is a given closed set. We observe that (\tilde{R}) is the optimal control problem with free endpoint constraints.

We impose here the following additional assumption **A4**, the necessity of which for the forthcoming development of our proof will become clear soon.

A4 The integrable function \tilde{l} is Lipschitz continuous on a neighbourhood of

$$(x^*(a), x^*(b), \max_{t \in [a, b]} h(t, x^*(t)))$$

and \tilde{l} is monotone in the z variable, in the sense that $z' \geq z$ implies $\tilde{l}(y, x, z') \geq \tilde{l}(y, x, z)$, for all $(y, x) \in \mathbb{R}^n \times \mathbb{R}^n$.

The following proposition is a straightforward adaptation of Proposition 9.5.4 of [17].

Proposition 3: Let (x^, u^*) be a strong local minimum for problem (\tilde{R}) . Assume the basic hypotheses, **A1**, **A2** and **A3** and the data for the problem (\tilde{R}) satisfies the hypothesis **A4**. Then there exist an absolutely continuous function $p : [a, b] \rightarrow \mathbb{R}^n$, an integrable function $\gamma : [a, b] \rightarrow \mathbb{R}^n$, a non-negative measure $\mu \in C^\oplus([a, b]; \mathbb{R})$, and a scalar $\lambda_0 \geq 0$ such that*

$$\mu\{[a, b]\} + \|p\|_\infty + \lambda_0 > 0, \quad (8)$$

$$(-\dot{p}(t), 0) \in \partial_{x, u}^C \langle q(t), f(t, x^*(t), u^*(t)) \rangle \quad (9)$$

$$-\{0\} \times N_{U(t)}^C(u^*(t)) \text{ a.e.}$$

$$(p(a), -q(b), \int_{[a, b]} \mu(ds)) \in \quad (10)$$

$$N_{E_a}^L(x^*(a)) \times \{0, 0\} +$$

$$\lambda_0 \partial \tilde{l}(x^*(a), x^*(b), \max_{t \in [a, b]} h(t, x^*(t))),$$

$$\gamma(t) \in \bar{\partial} h(t, x^*(t)) \quad \mu\text{-a.e.}, \quad (11)$$

$$\forall u \in U(t), \quad (12)$$

$$\langle q(t), f(t, x^*(t), u) \rangle \leq \langle q(t), f(t, x^*(t), u^*(t)) \rangle \text{ a.e.},$$

$$\text{supp}\{\mu\} \subset \quad (13)$$

$$\{t \in [a, b] : h(t, x^*(t)) = \max_{s \in [a, b]} h(s, x^*(s))\},$$

where q is defined as in (6).

We now turn to the derivation of Theorem 3.1. Consider the set

$$V := \{(x, u, e) : (x, u) \text{ satisfies } \dot{x}(t) = f(t, x(t), u(t)), \\ u(t) \in U(t) \text{ a.e.}, e \in \mathbb{R}^n, (x(a), e) \in C \\ \text{and } \|x - x^*\|_{L^\infty} \leq \varepsilon\} \quad (14)$$

and let $d_V : V \times V \rightarrow \mathbb{R}$ be a function defined by

$$d_V((x, u, e), (x', u', e')) = \quad (15)$$

$$|x(a) - x'(a)| + |e - e'| + \int_a^b |u(t) - u'(t)| dt$$

For all i , we choose $\varepsilon_i \downarrow 0$ and define the function

$$\tilde{l}_i(x, y, x', y', z) := \\ \max\{l(x, y) - l(x^*(a), x^*(b)) + \varepsilon_i^2, z, |x' - y'|\}.$$

Then d_V defines a metric on the set V and (V, d_V) is a complete metric space such that

- If $(x_i, u_i, e_i) \rightarrow (x, u, e)$ in the metric space (V, d_V) , then $\|x_i - x\|_{L^\infty} \rightarrow 0$,
- The function

$$(x, u, e) \rightarrow \tilde{l}_i(x(a), e, x(b), e, \max_{t \in [a, b]} h(t, x(t)))$$

is continuous on (V, d_V) .

We now consider the following optimization problem

$$\begin{aligned} \text{Minimize } & \{\tilde{l}_i(x(a), e, x(b), e, \max_{t \in [a, b]} h(t, x(t))) : \\ & (x, u, e) \in V\}. \end{aligned}$$

We observe that

$$\tilde{l}_i(x^*(a), x^*(b), x^*(b), x^*(b), \max_{t \in [a, b]} h(t, x^*(t))) = \varepsilon_i^2.$$

Since \tilde{l}_i is non-negative valued, it follows that $(x^*, u^*, x^*(b))$ is an ε_i^2 -minimizer for the above minimization problem. According to Ekeland's Theorem there exists a sequence $\{(x_i, u_i, e_i)\}$ in V such that for each i , we have

$$\begin{aligned} \tilde{l}_i(x_i(a), e_i, x_i(b), e_i, \max_{t \in [a, b]} h(t, x_i(t))) &\leq (16) \\ \tilde{l}_i(x(a), e, x(b), e, \max_{t \in [a, b]} h(t, x(t))) &+ \\ \varepsilon_i d_V((x, u, e), (x_i, u_i, e_i)) & \end{aligned}$$

for all $(x, u, e) \in V$ and we also have

$$d_V((x_i, u_i, e_i), (x^*, u^*, x^*(b))) \leq \varepsilon_i. \quad (17)$$

Thus the condition (17) implies that $e_i \rightarrow x^*(b)$ and $u_i \rightarrow u^*$ in the L^1 norm. By using subsequence extraction, we conclude that $u_i \rightarrow u^*$ a.e. and $x_i \rightarrow x^*$ uniformly.

Now we define the arc $y_i \equiv e_i$. Accordingly we get $y_i \rightarrow x^*(b)$ uniformly. From the minimization property (16), we say that $(x_i, y_i, w_i \equiv 0, u_i)$ is a strong local minimum for the optimal control problem

$$(\widetilde{R}_i) \begin{cases} \text{Minimize} \\ \tilde{l}_i(x(a), y(a), x(b), y(b), \max_{t \in [a, b]} h(t, x(t))) \\ + \varepsilon_i [|x(a) - x_i(a)| + |y(a) - y_i(a)| + w(b)] \\ \text{over } x, y, w \in W^{1,1} \text{ and measurable } u \text{ satisfying} \\ \dot{x}(t) = f(t, x(t), u(t)), \dot{y}(t) = 0, \\ \dot{w}(t) = |u(t) - u_i(t)| \text{ a.e.,} \\ u(t) \in U(t) \text{ a.e.,} \\ (x(a), y(a), w(a)) \in C \times \{0\}. \end{cases}$$

Now we observe that the cost function of (\widetilde{R}_i) satisfies all the assumptions of the Proposition 3 and thus this is an example of optimal control problem where the special case of maximum principle of Proposition 3 applies. Rewriting the conclusions of Proposition 3 and taking limits we obtained the required conditions. The remain of the proof follows closely the approach in [10].

V. CONCLUSIONS

In this work we derive a variant nonsmooth maximum principle for state constrained problems. The novelty of this work is that our results are also sufficient conditions of optimality for the normal linear-convex problems. The result presented in the main theorem is quite distinct to previous work in the literature since for state constrained problems, we add the Weierstrass conditions to adjoint inclusions using the joint subdifferentials with respect to the state and the control. The illustrated example presented in the paper justifies our results.

REFERENCES

- [1] Artstein Z. 2011. *Pontryagin Maximum Principle revisited with feedbacks*. European Journal of Control 17, 46-54.
- [2] Biswas M. H. A., de Pinho M. d. R. 2011. *A nonsmooth maximum principle for optimal control problems with state and mixed constraints-convex case*, DCDS Supplement 2011, Proceedings of the 8th AIMS International Conference, pp. 174 -183, September 2011.
- [3] Clarke F. 1976. *The maximum principle under minimal hypotheses*. SIAM J. Control Optim. 14, 1078-1091.
- [4] Clarke F. 1983. *Optimization and Nonsmooth Analysis*. John Wiley, New York.
- [5] Clarke F. 2005. *Necessary conditions in dynamic optimization*. Mem. Amer. Math. Soc.
- [6] Clarke F, de Pinho M. d. R. 2009. *The Nonsmooth Maximum Principle*. Control Cybernetics, 38, 1151-1168.
- [7] Clarke F, de Pinho M. d. R. 2010. *Optimal control problems with mixed constraints*. SIAM J. Control Optim. 48, 4500-4524.
- [8] de Pinho M. d. R., Vinter R. B. 1995. *An Euler-Lagrange inclusion for optimal control problems*. IEEE Trans. Automat. Control, 40, 1191-1198.
- [9] de Pinho M. d. R., Ferreira M. M., Fontes F. A. C. C. 2002. *An Euler-Lagrange inclusion for optimal control problems with state constraints*. Dynam. Control Systems 8, 23-45.
- [10] de Pinho M. d. R., Ferreira M. M., Fontes F. A. C. C. 2005. *Unmaximized inclusion necessary conditions for nonconvex constrained optimal control problems*. ESAIM Control Optim. Calc. Var., 11, 614-632.
- [11] de Pinho M. d. R., Loewen P, Silva G. N. 2009. *A weak maximum principle for optimal control problems with nonsmooth mixed constraints*. Set-Valued and Variational Analysis, 17, 203-2219.
- [12] Mordukhovich B. 2006. *Variational analysis and generalized differentiation. Basic Theory*. Springer-Verlag, Berlin.
- [13] Pesch H. J, Plail M. 2009. *The Maximum Principle of optimal control: A history of ingenious ideas and missed opportunities*. Control and Cybernetics, 38, 973-995.
- [14] Pontryagin L. S, Boltyanskii V. G, Gamkrelidze R. V, Mischenko E. F. 1962. *The Mathematical Theory of Optimal Processes*. John Wiley, New York.
- [15] Rockafellar R. T and Wets B., *Variational Analysis*, Grundlehren Math. Wiss. 317, Springer-Verlag, Berlin, 1998.
- [16] Vinter R.B, Pappas G. 1982. *A maximum principle for nonsmooth optimal-control problems with state constraints*. J. Math. Anal. Appl., 89, 212-232.
- [17] Vinter R. 2000. *Optimal Control*. Birkhäuser, Boston.