

Erdélyi-Kober Fractional Integral Operators from a Statistical Perspective -II

A.M. MATHAI

Centre for Mathematical Sciences,
Arunapuram P.O., Pala, Kerala-68674, India, and
Department of Mathematics and Statistics, McGill University,
Montreal, Quebec, Canada, H3A 2K6

and

H.J. HAUBOLD

Office for Outer Space Affairs, United Nations
P.O. Box 500, Vienna International Centre
A - 1400 Vienna, Austria, and
Centre for Mathematical Sciences,
Arunapuram P.O., Pala, Kerala-68674, India

Abstract

In this article we examine the densities of a product and a ratio of two real positive definite matrix-variate random variables X_1 and X_2 , which are statistically independently distributed, and we consider the density of the product $U_1 = X_2^{\frac{1}{2}} X_1 X_2^{\frac{1}{2}}$ as well as the density of the ratio $U_2 = X_2^{\frac{1}{2}} X_1^{-1} X_2^{\frac{1}{2}}$. We define matrix-variate Kober fractional integral operators of the first and second kinds from a statistical perspective, making use of the derivation in the predecessor of this paper for the scalar variable case, by deriving the densities of products and ratios where one variable has a matrix-variate type-1 beta density and the other variable has an arbitrary density. Various types of generalizations are considered, by using pathway models, by appending matrix variate hypergeometric series etc. During this process matrix-variate Saigo operator and other operators are also defined and properties studied.

1. Introduction

All the matrices appearing in this article are $p \times p$ real positive definite. Corresponding results for hermitian cases can be obtained but not given in this article. The following standard notations will be used. $A > O$ means the $p \times p$ real matrix is symmetric, $X = X'$, and further, it is positive definite. $|A|$ means the determinant of A , $\text{tr}(A)$ = trace of A , dX will stand for the wedge product of differentials in any matrix X . If X is $p \times q$ with $X = (x_{ij})$ then

$$dX = dx_{11} \wedge dx_{12} \wedge \dots \wedge dx_{pq} \text{ for a general matrix} \quad (1.1)$$

$$= \prod_{i \geq j} \wedge dx_{ij} = \prod_{j \geq i} \wedge dx_{ij} \text{ when } X = X' \text{ or when } X \text{ is symmetric.} \quad (1.2)$$

Also $\int_X f(X) dX$ will mean the integrals over all X (need not be symmetric or even square) of a real-valued scalar function $f(X)$ of matrix argument X . In the same format

$$\int_A^B f(Y) dY = \int_{O < A < Y < B} f(Y) dY$$

will mean the integration of a real-valued scalar function of the real positive definite $p \times p$ matrix Y over the space of positive definite matrices such that $A > O, X > O, X - A > O, B - X > O$ where A and B are $p \times p$ constant matrices. The notation will then imply that if $O < X < I$ then all eigenvalues of X are in the open interval $(0, 1)$. We will need some Jacobians of matrix transformations in this paper. For further results on Jacobians, see Mathai (1997).

$$Y = AXA', X = X', |A| \neq 0 \Rightarrow dY = |A|^{p+1} dX \quad (1.3)$$

$$Y = X^{-1} \Rightarrow dY = \begin{cases} |X|^{-2p} dX & \text{for a general } X \\ |X|^{-(p+1)} dX & \text{for } X = X'. \end{cases} \quad (1.4)$$

We will denote the unique positive definite square root of a positive definite matrix A by $A^{\frac{1}{2}}$. The following standard property will be used very often in this article. For $p \times p$ nonsingular matrices A and B

$$\begin{aligned} |I \pm AB| &= |I \pm BA| = |A| |A^{-1} \pm B| = |B| |B^{-1} \pm A| \quad (\text{when nonsingular}) \\ |I \pm AB| &= |I \pm A^{\frac{1}{2}} B A^{\frac{1}{2}}| = |I \pm B^{\frac{1}{2}} A B^{\frac{1}{2}}| \quad (\text{when positive definite}). \end{aligned} \quad (1.5)$$

The real matrix-variate gamma function, denoted by $\Gamma_p(\alpha)$ is defined as follows which has an integral representation when $\Re(\alpha) > \frac{p-1}{2}$.

$$\begin{aligned} \Gamma_p(\alpha) &= \pi^{\frac{p(p-1)}{4}} \Gamma(\alpha) \Gamma(\alpha - \frac{1}{2}) \dots \Gamma(\alpha - \frac{p-1}{2}), \Re(\alpha) > \frac{p-1}{2} \\ &= \int_{X>O} |X|^{\alpha - \frac{p+1}{2}} e^{-\text{tr}(X)} dX, \Re(\alpha) > \frac{p-1}{2}. \end{aligned} \quad (1.6)$$

The real matrix-variate type-1 beta density for the $p \times p$ positive definite matrix X_1 , with parameters α and β and denoted by $f_1(X_1)$, is defined as follows:

$$\begin{aligned} f_1(X_1) &= \frac{\Gamma_p(\alpha + \beta)}{\Gamma_p(\alpha) \Gamma_p(\beta)} |X_1|^{\alpha - \frac{p+1}{2}} |I - X_1|^{\beta - \frac{p+1}{2}}, O < X_1 < I \\ f_2(Y_1) &= \frac{\Gamma_p(\alpha + \beta)}{\Gamma_p(\alpha) \Gamma_p(\beta)} |Y_1|^{\beta - \frac{p+1}{2}} |I - Y_1|^{\alpha - \frac{p+1}{2}} dY_1, O < Y_1 < I \end{aligned}$$

for $\Re(\alpha) > \frac{p-1}{2}$, $\Re(\beta) > \frac{p-1}{2}$, and $f(X_1) = 0$, $f_2(Y_1) = 0$ elsewhere. Type-1 and Type-2 beta integrals and beta functions are defined and denoted as follows for $\Re(\alpha) > \frac{p-1}{2}$, $\Re(\beta) > \frac{p-1}{2}$:

$$\begin{aligned} B_p(\alpha, \beta) &= \frac{\Gamma_p(\alpha) \Gamma_p(\beta)}{\Gamma_p(\alpha + \beta)} \\ &= \int_{O < X < I} |X|^{\alpha - \frac{p+1}{2}} |I - X|^{\beta - \frac{p+1}{2}} dX \quad (\text{type-1}) \\ &= \int_{O < Y < I} |Y|^{\beta - \frac{p+1}{2}} |I - Y|^{\alpha - \frac{p+1}{2}} dY \quad (\text{type-1}) \\ &= \int_{U > O} |U|^{\alpha - \frac{p+1}{2}} |I + U|^{-(\alpha + \beta)} dU \quad (\text{type-2}) \\ &= \int_{V > O} |V|^{\beta - \frac{p+1}{2}} |I + V|^{-(\alpha + \beta)} dV \quad (\text{type-2}) \end{aligned} \quad (1.7)$$

2. Kober Operator of the Second Kind for the Real Matrix-variate Case

Definition 2.1. We will define and denote Kober operator of the second kind for the real matrix-variate case as follows:

$$K_X^{\zeta, \alpha} f(X) = \frac{|X|^{\zeta}}{\Gamma_p(\alpha)} \int_{T>X} |T - X|^{\alpha - \frac{p+1}{2}} |T|^{-\zeta - \alpha} f(T) dT, \Re(\alpha) > \frac{p-1}{2}. \quad (2.1)$$

Consider two $p \times p$ real matrix-variate random variables X_1 and X_2 , independently distributed, where X_1 has a real matrix-variate type-1 beta density $f_1(X_1)$ with parameters $(\zeta + \frac{p+1}{2}, \alpha)$, that is,

$$f_1(X_1) = \frac{\Gamma_p(\zeta + \alpha + \frac{p+1}{2})}{\Gamma_p(\zeta + \frac{p+1}{2})\Gamma_p(\alpha)} |X_1|^\zeta |I - X_1|^{\alpha - \frac{p+1}{2}}, 0 < X_1 < I$$

for $\Re(\zeta) > -1, \Re(\alpha) > \frac{p-1}{2}$ and $f_1(X_1) = 0$ elsewhere. Let X_2 have an arbitrary density $f(X_2)$. Then the joint density of X_1 and X_2 is $f_1(X_1)f(X_2)$. Let us consider the transformation $U = X_2^{\frac{1}{2}}X_1X_2^{\frac{1}{2}}, X_2 = V$ so that $X_2 = V, X_1 = V^{-\frac{1}{2}}UV^{-\frac{1}{2}}$ and the Jacobian is given by $dX_1 \wedge dX_2 = |V|^{-\frac{p+1}{2}} dU \wedge dV$. If the joint density is denoted by $f(U, V)$ then

$$\begin{aligned} f(U, V)dU \wedge dV &= \frac{\Gamma_p(\zeta + \alpha + \frac{p+1}{2})}{\Gamma_p(\alpha)\Gamma(\zeta + \frac{p+1}{2})} |V^{-\frac{1}{2}}UV^{-\frac{1}{2}}|^\zeta \\ &\times |I - V^{-\frac{1}{2}}UV^{-\frac{1}{2}}|^{\alpha - \frac{p+1}{2}} f(V) |V|^{-\frac{p+1}{2}} dU \wedge dV. \end{aligned}$$

Therefore the marginal density of U , denoted by $g(U)$, is available by integrating out V from $f(U, V)$. That is,

$$\begin{aligned} g(U) &= \int_V f_1(V^{-\frac{1}{2}}UV^{-\frac{1}{2}})f(V)|V|^{-\frac{1}{2}}dV \\ &= \frac{\Gamma_p(\zeta + \alpha + \frac{p+1}{2})}{\Gamma_p(\zeta + \frac{p+1}{2})} \int_{V>U} \frac{1}{\Gamma_p(\alpha)} |V|^{-\frac{p+1}{2}} \\ &\times |U|^\zeta |V|^{-\zeta} |V|^{-\alpha + \frac{p+1}{2}} |V - U|^{\alpha - \frac{p+1}{2}} f(V)dV \\ &= \frac{\Gamma_p(\alpha + \zeta + \frac{p+1}{2})}{\Gamma_p(\zeta + \frac{p+1}{2})} K_U^{\zeta, \alpha} f(U). \end{aligned}$$

Hence we have the following theorem:

Theorem 2.1. *When X_1 and X_2 are independently distributed $p \times p$ positive definite real matrix random variables and when $X_2 = V$ and $U = X_2^{\frac{1}{2}}X_1X_2^{\frac{1}{2}}$ or $X_1 = V^{-\frac{1}{2}}UV^{-\frac{1}{2}}$ and when X_1 has a real matrix-variate type-1 beta distribution with the parameters $(\zeta + \frac{p+1}{2}, \alpha)$ and if $g(U)$ denotes the density of U then*

$$\frac{\Gamma_p(\zeta + \frac{p+1}{2})}{\Gamma_p(\alpha + \zeta + \frac{p+1}{2})} g(U) = K_U^{\zeta, \alpha} f(U) \quad (2.2)$$

is Kober fractional integral operator of the second kind for the real matrix-variate case.

As a special case of (2.2), or independently, we can derive a result for the right sided Weyl operator for the real matrix-variate case. Let the right sided Weyl fractional integral operator be denoted by $({}_X W_\infty^{-\alpha} f)(X)$ with infinity here signifying that $T - X$ is positive definite.

Theorem 2.2. *Let X_1, X_2, U, V be as defined in Theorem 2.1. Let X_1 have a type-1 beta density with the parameters $(\frac{p+1}{2}, \alpha)$. Let the arbitrary density of X_2 be denoted by $f_2(X_2) = |X_2|^\alpha f(X_2)$, where $f(X_2)$ is arbitrary. Let the density of U be again denoted by $g(U)$. Then*

$$\begin{aligned} ({}_X W_\infty^{-\alpha} f)(X) &= \frac{1}{\Gamma(\alpha)} \int_{V>U} |V - U|^{\alpha - \frac{p+1}{2}} f(V)dV \\ &= \frac{\Gamma_p(\frac{p+1}{2})}{\Gamma_p(\alpha + \frac{p+1}{2})} g(U), \Re(\alpha) > \frac{p-1}{2}. \end{aligned} \quad (2.3)$$

2.1. A pathway generalization of Kober operator of the second kind in the matrix case

A pathway generalization, parallel to the results in the scalar case can be considered. In the pathway case when generalization to matrices is considered we take $\delta = 1$. For a general δ , there will be problems with Jacobians of transformations for X^δ even if $X > O$ and $\delta > 0$, see for example Mathai (1997) for the case $\delta = 2$ and when $X = X'$ to see the type of complications. Hence we take the case $\delta = 1$ only. Let X_1 have a pathway density

$$f_1(X_1) = C_1 |X_1|^\gamma |I - a(1-q)X|^{-\frac{\eta}{1-q}} \quad (2.4)$$

for $I - a(1-q)X > O, q < 1, \eta > 0, a > 0$ where C_1 can be seen to be the following:

$$C_1 = \frac{[a(1-q)]^{p\gamma + \frac{p(p+1)}{2}} \Gamma_p(\gamma + \frac{\eta}{1-q} + (p+1))}{\Gamma_p(\gamma + \frac{p+1}{2}) \Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2})}. \quad (2.5)$$

Let X_2 have an arbitrary density $f(X_2)$ and let X_1 and X_2 be statistically independently distributed. Let $U = X_2^{\frac{1}{2}} X_1 X_2^{\frac{1}{2}}, X_2 = V$ or $X_1 = V^{-\frac{1}{2}} U V^{-\frac{1}{2}}$ and the Jacobian is $|V|^{-\frac{p+1}{2}}$. Let $g(U)$ be the density of U . Then, going through the earlier steps we have the following:

$$g(U) = C_1 |U|^\gamma \int_{V > a(1-q)U} |V|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} |V - a(1-q)U|^{-\frac{\eta}{1-q}} f(V) dV. \quad (2.6)$$

Then

$$\begin{aligned} \Gamma_p(\gamma + \frac{p+1}{2}) g(U) &= \frac{[a(1-q)]^{p\gamma + \frac{p(p+1)}{2}} \Gamma_p(\gamma + \frac{\eta}{1-q} + \frac{(p+1)}{2})}{\Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2})} |U|^\gamma \\ &\quad \times \int_{V > a(1-q)U} |V - a(1-q)U|^{-\frac{\eta}{1-q}} |V|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} f(V) dV \\ &= K_{U,a,q}^{\gamma, \frac{\eta}{1-q} + \frac{p+1}{2}} f(U) \end{aligned} \quad (2.7)$$

where $K_{U,a,q}^{\gamma, \frac{\eta}{1-q} + \frac{p+1}{2}} f(U)$ can be called the generalized pathway Kober operator of the second kind in the real matrix-variate case. When the pathway parameter q varies from $-\infty$ to 1 it provides a pathway or a class of operators and all these operators in this pathway class will eventually go to the exponential form. For $a = 1, q = 0, \frac{\eta}{1-q} = \alpha - \frac{p+1}{2}$ and $\gamma = \zeta$ we have

$$K_{U,a,q}^{\gamma, \frac{\eta}{1-q} + \frac{p+1}{2}} f(U) = K_U^{\zeta, \alpha} f(U) \quad (2.8)$$

the Kober operator of the second kind as a constant multiple of the density of the product of two matrix-variate independently distributed random variables. Note that when $q \rightarrow 1_-$ we can evaluate the limit of $g(U)$ by using the following lemmas:

Lemma 2.1.

$$\lim_{q \rightarrow 1_-} C_1 = \frac{(a\eta)^{p\gamma + \frac{p(p+1)}{2}}}{\Gamma_p(\gamma + \frac{p+1}{2})} \quad (i)$$

Proof: Open up each $\Gamma_p(\cdot)$ in C_1 of (2.5) in terms of ordinary gamma functions. Then use the following asymptotic approximation for gamma functions. For $|z| \rightarrow \infty$ and γ a bounded quantity

$$\Gamma(z + \gamma) \approx \sqrt{2\pi} z^{z + \gamma - \frac{1}{2}} e^{-z}. \quad (ii)$$

This is the first term in the asymptotic series. This term is also known as Stirling's approximation. When $q \rightarrow 1_-$ we have $\frac{1}{1-q} \rightarrow \infty$ and hence take $|z|$ as $\frac{\eta}{1-q}$ and expand all gammas by using Stirling's approximation to see that C_1 reduces to (i) above.

Lemma 2.2.

$$\lim_{q \rightarrow 1_-} |I - a(1-q)X|^{\frac{\eta}{1-q}} = e^{-a\eta \operatorname{tr}(X)}. \quad (iii)$$

Proof: Writing the determinant in terms of eigenvalues we have

$$|I - a(1-q)X|^{\frac{\eta}{1-q}} = \prod_{j=1}^p (1 - a(1-q)\lambda_j)^{\frac{\eta}{1-q}} \quad (iv)$$

where $\lambda_1, \dots, \lambda_p$ are the eigenvalues of X . Now

$$\lim_{q \rightarrow 1_-} (1 - a(1-q)\lambda_j)^{\frac{\eta}{1-q}} = e^{-a\eta \lambda_j}. \quad (v)$$

Hence

$$\lim_{q \rightarrow 1_-} |I - a(1-q)X|^{\frac{\eta}{1-q}} = \prod_{j=1}^p e^{-a\eta \lambda_j} = e^{-a\eta \operatorname{tr}(X)}$$

which establishes (iii). Now by using Lemmas 2.1 and 2.2 we have

$$\begin{aligned} \lim_{q \rightarrow 1_-} g(U) &= \frac{(a\eta)^{p\gamma + \frac{p(p+1)}{2}}}{\Gamma_p(\gamma + \frac{p+1}{2})} |U|^\gamma \\ &\times \int_{V>O} |V|^{-\gamma - \frac{p+1}{2}} e^{-a\eta \operatorname{tr}(V^{-\frac{1}{2}}UV^{-\frac{1}{2}})} dV. \end{aligned} \quad (2.9)$$

This is the limiting form of the pathway Kober operator of the second kind in this class of pathway operators of the second kind.

In the pathway generalization, one can also replace the parameter a with a constant positive definite matrix A . In this case the model will be written as

$$f_1(X_1) = C_1(A) |X_1|^\gamma |I - (1-q)A^{\frac{1}{2}}X_1A^{\frac{1}{2}}|^{\frac{\eta}{1-q}} \quad (2.10)$$

for $q < 1, A > O, X_1 > O, I - (1-q)A^{\frac{1}{2}}X_1A^{\frac{1}{2}} > O$. The pathway parameter is still q . In this case

$$C_1(A) = \frac{(1-q)^{p\gamma + \frac{p(p+1)}{2}} |A|^{\gamma + \frac{p+1}{2}} \Gamma_p(\gamma + \frac{\eta}{1-q} + (p+1))}{\Gamma_p(\gamma + \frac{p+1}{2}) \Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2})}. \quad (2.11)$$

Then $g(U)$ of (2.6) goes to the following form, denoted by $g_A(U)$

$$\begin{aligned} g_A(U) &= C_1(A) |A|^{\frac{\eta}{1-q}} |U|^\gamma \\ &\times \int_{V^*>O} |V^{\frac{1}{2}}A^{-1}V^{\frac{1}{2}} - (1-q)U|^{\frac{\eta}{1-q}} |V|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} f(V) dV \end{aligned} \quad (2.12)$$

where

$$V^* = V^{\frac{1}{2}}A^{-1}V^{\frac{1}{2}} - (1-q)U.$$

Then one can define a pathway generalized Kober operator of the second kind as

$$\begin{aligned} K_{U,A,q}^{\gamma, \frac{\eta}{1-q} + \frac{p+1}{2}} f(U) &= \Gamma_p(\gamma + \frac{p+1}{2}) g_A(U) \\ &= \frac{(1-q)^{p\gamma + \frac{p(p+1)}{2}} |A|^{\gamma + \frac{\eta}{1-q} + \frac{p+1}{2}} \Gamma_p(\gamma + \frac{\eta}{1-q} + (p+1))}{\Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2})} |U|^\gamma \\ &\times \int_{V^*>O} |V^{\frac{1}{2}}A^{-1}V^{\frac{1}{2}} - (1-q)U|^{\frac{\eta}{1-q}} |V|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} f(V) dV \end{aligned} \quad (2.13)$$

In this case, as $q \rightarrow 1_-$ we have

$$\begin{aligned} \lim_{q \rightarrow 1^-} g_A(U) &= \frac{|A|^{\gamma + \frac{p+1}{2}} \eta^{p\gamma + \frac{p(p+1)}{2}}}{\Gamma_p(\gamma + \frac{p+1}{2})} |U|^\gamma \\ &\times \int_{V>A} |V|^{-\gamma - \frac{p+1}{2}} e^{-\eta \operatorname{tr}(A^{\frac{1}{2}} V^{-\frac{1}{2}} U V^{-\frac{1}{2}} A^{\frac{1}{2}})} f(V) dV. \end{aligned} \quad (2.14)$$

3. M-transforms of Kober Operator of the Second Kind

The generalized matrix transform or M-transform is defined and illustrated in Mathai (1997). The M-transform of Kober operator of the second kind is the following:

Theorem 3.1. *For the Kober operator of the second kind defined in (2.1) the M-transform with parameter s is given by*

$$\begin{aligned} M\{K_X^{\zeta, \alpha} f(X); s\} &= \int_{X>O} |X|^{s - \frac{p+1}{2}} \left[\int_{T>X} \frac{|X|^\zeta}{\Gamma_p(\alpha)} |T - X|^{\alpha - \frac{p+1}{2}} |T|^{-\zeta - \alpha} f(T) dT \right] dX \\ &= \frac{\Gamma_p(\zeta + s)}{\Gamma(\alpha + \zeta + s)} f^*(s), \quad \Re(\zeta + s) > \frac{p-1}{2}, \Re(\alpha) > \frac{p-1}{2} \end{aligned} \quad (3.1)$$

where $f^*(s)$ is the M-transform of $f(X)$.

Proof: Interchanging the integral we have

$$M\{K_X^{\zeta, \alpha} f(X); s\} = \int_{T>O} |T|^{-\zeta - \alpha} f(T) \left[\frac{1}{\Gamma_p(\alpha)} \int_{O<X<T} |X|^{\zeta + s - \frac{p+1}{2}} |T - X|^{\alpha - \frac{p+1}{2}} dX \right] dT.$$

Note that

$$|T - X| = |T| |I - T^{-\frac{1}{2}} X T^{-\frac{1}{2}}|, \quad Y = T^{-\frac{1}{2}} X T^{-\frac{1}{2}} \Rightarrow dY = |T|^{-\frac{p+1}{2}} dX.$$

Hence

$$\int_{X<T} |X|^{\zeta - \frac{p+1}{2}} |T - X|^{\alpha - \frac{p+1}{2}} dX = |T|^{\alpha + \zeta + s - \frac{p+1}{2}} \int_Y |Y|^{\zeta + s - \frac{p+1}{2}} |I - Y|^{\alpha - \frac{p+1}{2}} dY.$$

We can evaluate the Y -integral by using real matrix-variate type-1 beta integral.

$$\int_{O<Y<I} |Y|^{\zeta + s - \frac{p+1}{2}} |I - Y|^{\alpha - \frac{p+1}{2}} dY = \frac{\Gamma_p(\zeta + s) \Gamma_p(\alpha)}{\Gamma_p(\alpha + \zeta + s)}$$

for $\Re(\alpha) > \frac{p-1}{2}$, $\Re(\zeta + s) > \frac{p-1}{2}$. Now the T -integral gives

$$\int_{T>O} |T|^{s - \frac{p+1}{2}} f(T) dT = f^*(s)$$

where $f^*(s)$ is the M-transform of $f(X)$. Hence (3.1) follows. Note that for $p = 1$ the result agrees with that in the scalar case, which is available in the literature, see for example Mathai and Haubold (2008).

From (3.1) for $\zeta = 0$ and $\Re(\alpha) > \frac{p-1}{2}$ we have the special case of the Kober operator of the second kind

$$K_X^{0, \alpha} f(X) = \frac{1}{\Gamma(\alpha)} \int_{T>X} |T - X|^{\alpha - \frac{p+1}{2}} |T|^{-\alpha} f(T) dT. \quad (3.2)$$

But the right side of (3.2) is Weyl fractional integral of order α in the matrix case, ${}_X W_\infty^{-\alpha} f(X)$, except for the factor $|T|^{-\alpha}$. The Weyl integral in the matrix case is

$${}_X W_\infty^{-\alpha} f(X) = \frac{1}{\Gamma_p(\alpha)} \int_{T>X} |T - X|^{\alpha - \frac{p+1}{2}} f(T) dT, \quad \Re(\alpha) > \frac{p-1}{2}. \quad (3.3)$$

Hence we have the following corollary.

Corollary 3.1.1. *The M-transform of the right sided Weyl operator in the real matrix case is given by*

$$M\{{}_XW_\infty^{-\alpha}|X|^{-\alpha}f(X); s\} = \frac{\Gamma_p(s)}{\Gamma_p(\alpha+s)}f^*(s) \quad (3.4)$$

for $\Re(s) > \frac{p-1}{2}$, $\Re(\alpha) > \frac{p-1}{2}$ where $f^*(s)$ is the M-transform of $f(X)$.

The proof is parallel to that in Theorem 3.1. Let us see whether a Mellin convolution type formula holds for Kober operator of the second kind in the matrix case. Let

$$g(U) = \int_V |V|^{-\frac{p+1}{2}} f_1(V^{-\frac{1}{2}}UV^{-\frac{1}{2}})f_2(V)dV \quad (3.5)$$

where $f_1(X_1)$ is a type-1 matrix-variate beta density with parameters $(\zeta + \frac{p+1}{2}, \alpha)$. That is,

$$f_1(X_1) = \frac{\Gamma_p(\alpha + \zeta + \frac{p+1}{2})}{\Gamma_p(\alpha)\Gamma_p(\zeta + \frac{p+1}{2})}|X_1|^\zeta|I - X_1|^{\alpha - \frac{p+1}{2}}, 0 < X_1 < I \quad (3.6)$$

for $\Re(\alpha) > \frac{p-1}{2}$, $\Re(\zeta) > -1$ and $f_1(X_1) = 0$ elsewhere. Substituting (3.6) in (3.5) we have

$$\begin{aligned} \frac{\Gamma_p(\zeta + \frac{p+1}{2})}{\Gamma_p(\alpha + \zeta + \frac{p+1}{2})}g(U) &= \frac{1}{\Gamma_p(\alpha)} \int_V |V|^{-\frac{p+1}{2}}|U|^\zeta|V|^{-\zeta} \\ &\quad \times |I - V^{-\frac{1}{2}}UV^{-\frac{1}{2}}|^{\alpha - \frac{p+1}{2}} f(V)dV \\ &= \frac{|U|^\zeta}{\Gamma_p(\alpha)} \int_V |V|^{-\zeta - \alpha}|V - U|^{\alpha - \frac{p+1}{2}} f(V)dV \\ &= \frac{|U|^\zeta}{\Gamma_p(\alpha)} \int_{V>U} |V - U|^{\alpha - \frac{p+1}{2}}|V|^{-\zeta - \alpha} f(V)dV \\ &= K_U^{\zeta, \alpha} f(U) \end{aligned} \quad (3.7)$$

which is the Kober operator of the second kind. Hence we have the following theorem:

Theorem 3.2. *Kober operator of the second kind with real matrix argument can also be represented as a Mellin convolution type formula*

$$K_X^{\zeta, \alpha} f(X) = \int_V |V|^{-\frac{p+1}{2}} f_1(V^{-\frac{1}{2}}XV^{-\frac{1}{2}})f_2(V)dV$$

where $f_1(X_1)$ is a type-1 beta density with parameters $(\zeta + \frac{p+1}{2}, \alpha)$ and $f_2(V)$ is an arbitrary function or arbitrary density if the Kober operator is to be taken as a constant multiple of a statistical density.

4. Generalization in Terms of Hypergeometric Series for Kober Operator of the Second Kind in the Real Matrix Case

For introducing hypergeometric series of matrix argument we will need the definitions, notation and lemmas. Hypergeometric functions of matrix argument are defined in terms of matrix-variate Laplace transforms, M-transforms and zonal polynomials. Explicit series form for all cases is available through the definition in terms of zonal polynomials and hence we will define in terms of zonal polynomials.

$$\begin{aligned} {}_rF_s(Z) &= {}_rF_s(a_1, \dots, a_r; b_1, \dots, b_s; Z) \\ &= \sum_{k=0}^{\infty} \sum_K \frac{(a)_K \dots (a_r)_K}{(b_1)_K \dots (b_s)_K} \frac{C_K(Z)}{k!} \end{aligned} \quad (4.1)$$

where $K = (k_1, \dots, k_p)$, $k_1 + \dots + k_p = k$ is a partition of $k = 0, 1, 2, \dots$

$$(a)_K = \prod_{j=1}^p \left(a - \frac{j-1}{2}\right)_{k_j}, (b)_{k_j} = b(b+1)\dots(b+k_j-1), (b)_0 = 1, b \neq 0 \quad (4.2)$$

and $C_K(Z)$ is a zonal polynomial of order k and Z is a $p \times p$ matrix. The series is defined for the real and complex matrices. Zonal polynomials are certain symmetric functions of the eigenvalues of Z . In our discussions, Z will be real and positive definite. For more details about zonal polynomials see Mathai, Provost and Hayakawa (1995). The following basic results are needed in our discussions. A standard notation in this area is

$$\Gamma_p(\alpha, K) = \Gamma_p(\alpha)(\alpha)_K. \quad (4.3)$$

The following basic results are needed in our discussion.

Lemma 4.1.

$$\int_O^I |X|^{\alpha - \frac{p+1}{2}} |I - X|^{\beta - \frac{p+1}{2}} C_K(TX) dX = \frac{\Gamma_p(\alpha, K) \Gamma_p(\beta)}{\Gamma_p(\alpha + \beta, K)} C_K(T) \quad (4.4)$$

for $\Re(\alpha) > \frac{p-1}{2}, \Re(\beta) > \frac{p-1}{2}$.

Lemma 4.2. For $\Re(\alpha) > \frac{p-1}{2}, A > O, S > O$

$$\int_{O < S < A} |S|^{\alpha - \frac{p+1}{2}} C_K(ZS) dS = \frac{\Gamma_p(\alpha, K) \Gamma_p(\frac{p+1}{2})}{\Gamma_p(\alpha + \frac{p+1}{2}, K)} |A|^\alpha C_K(ZA). \quad (4.5)$$

Let us assume that all the parameters $a_1, \dots, a_r, b_1, \dots, b_s$ are real and positive and let the argument matrices be $p \times p$ and positive definite. For $A > O$, let the density of X_1 be

$$\begin{aligned} f_1(X_1) &= \frac{1}{c_f} {}_rF_s(a_1, \dots, a_r; b_1, \dots, b_s; AX_1) |X_1|^\zeta |I - X_1|^{\alpha - \frac{p+1}{2}} \\ &= \frac{1}{c_f} \sum_{k=0}^{\infty} \sum_K \frac{(a_1)_K \dots (a_r)_K}{(b_1)_K \dots (b_s)_K} \frac{1}{k!} C_K(AX_1) |X_1|^\zeta |I - X_1|^{\alpha - \frac{p+1}{2}} \end{aligned} \quad (4.6)$$

where the normalizing constant c_f is available by integrating out term by term with the help of Lemma 4.1. It will be available in terms of a ${}_{r+1}F_{s+1}$. Let $f(X_2)$ be an arbitrary density. As before, let $X_1 = V^{-\frac{1}{2}}UV^{-\frac{1}{2}}$, then denoting the density of U again by $g(U)$ we have

$$\begin{aligned} g(U) &= \int_v f_1(V^{-\frac{1}{2}}UV^{-\frac{1}{2}}) f(V) |V|^{-\frac{p+1}{2}} dV \\ &= \frac{1}{c_f} \frac{\Gamma_p(\alpha + \zeta + \frac{p+1}{2})}{\Gamma_p(\zeta + \frac{p+1}{2}) \Gamma_p(\alpha)} \left(\sum_{k=0}^{\infty} \sum_K \frac{(a_1)_K \dots (a_r)_K}{(b_1)_K \dots (b_s)_K} \frac{1}{k!} \right) \\ &\quad \times \int_V |V^{-\frac{1}{2}}UV^{-\frac{1}{2}}|^\zeta |I - V^{-\frac{1}{2}}UV^{-\frac{1}{2}}|^{\alpha - \frac{p+1}{2}} |V|^{-\frac{p+1}{2}} C_K(AV^{-\frac{1}{2}}UV^{-\frac{1}{2}}) f(V) dV \end{aligned} \quad (4.7)$$

This is the generalization of a constant times the Kober operator of the second kind in the matrix case. For ${}_rF_s = {}_2F_1$ one has the matrix-variate generalization of a constant times the Saigo operator of the second kind in the real matrix-variate case.

5. Kober Fractional Integral Operators of the First Kind in the Matrix Case

Definition 5.1. We will give the following definition and notation for Kober fractional integral operator of the first kind in the real matrix-variate case:

$$I_X^{\zeta, \alpha} f(X) = \frac{|X|^{-\zeta - \alpha}}{\Gamma_p(\alpha)} \int_{V < X} |X - V|^{\alpha - \frac{p+1}{2}} |V|^\zeta f(V) dV \quad (5.1)$$

for $\Re(\zeta) > -1, \Re(\alpha) > \frac{p-1}{2}$.

Theorem 5.1. For $\Re(\alpha) > \frac{p-1}{2}, \Re(\zeta) > -1$ the M -transform, with parameter s , of Kober operator of the first kind in the real matrix-variate case, is given by

$$\begin{aligned} M\{I_X^{\zeta, \alpha} f(X); s\} &= \int_{X>O} |X|^{s-\frac{p+1}{2}} \left[\frac{|X|^{-\zeta-\alpha}}{\Gamma_p(\alpha)} \int_{V<X} |X-V|^{\alpha-\frac{p+1}{2}} |V|^\zeta f(V) dV \right] dX \\ &= \frac{\Gamma_p(\zeta + \frac{p+1}{2} - s)}{\Gamma_p(\alpha + \zeta + \frac{p+1}{2} - s)} f^*(s), \Re(s) < \Re(\zeta + 1), \Re(\alpha) > \frac{p-1}{2} \end{aligned} \quad (5.2)$$

where $f^*(s)$ is the M -transform of $f(X)$.

Proof: Integrating out X first we have the X -integral

$$\begin{aligned} \int_{X>V} |X|^{s-\zeta-\alpha-\frac{p+1}{2}} |X-V|^{\alpha-\frac{p+1}{2}} dX &= \int_{Y>O} |Y+V|^{s-\zeta-\alpha-\frac{p+1}{2}} |Y|^{\alpha-\frac{p+1}{2}} dY, Y = X - V \\ &= |V|^{s-\zeta-\alpha-\frac{p+1}{2}} \int_{Y>O} |I + V^{-\frac{1}{2}} Y V^{-\frac{1}{2}}|^{s-\zeta-\alpha-\frac{p+1}{2}} |Y|^{\alpha-\frac{p+1}{2}} dY. \end{aligned}$$

Put $Z = V^{-\frac{1}{2}} Y V^{-\frac{1}{2}} \Rightarrow dZ = |V|^{-\frac{p+1}{2}} dY$. Then the X -integral is

$$\begin{aligned} |V|^{s-\zeta-\frac{p+1}{2}} \int_{Z>O} |Z|^{\alpha-\frac{p+1}{2}} |I+Z|^{-(\frac{p+1}{2}+\alpha+\zeta-s)} dZ \\ = |V|^{s-\zeta-\frac{p+1}{2}} \frac{\Gamma_p(\alpha) \Gamma_p(\frac{p+1}{2} + \zeta - s)}{\Gamma_p(\frac{p+1}{2} + \alpha + \zeta - s)} \end{aligned}$$

for $\Re(\alpha) > \frac{p-1}{2}, \Re(\zeta - s) > -1$ by evaluating the integral by using a type-2 matrix-variate beta integral in the real case. Now, the V -integral becomes

$$\int_{V>O} |V|^{s-\frac{p+1}{2}} f(V) dV = f^*(s).$$

Hence

$$M\{I_X^{\zeta, \alpha} f(X); s\} = \frac{\Gamma_p(\frac{p+1}{2} + \zeta - s)}{\Gamma_p(\frac{p+1}{2} + \alpha + \zeta - s)} f^*(s) \quad (5.3)$$

for $\Re(\alpha) > \frac{p-1}{2}, \Re(\zeta - s) > -1$. Note that for $\zeta = 0$,

$$I_X^{0, \alpha} f(X) = |X|^{-\alpha} {}_0D_X^{-\alpha} f(X) \quad (5.4)$$

where ${}_0D_X^{-\alpha}$ is the left sided Riemann-Liouville fractional integral for the matrix-variate case. Note that for the scalar case, for $p = 1$,

$$M\{I_x^{\zeta, \alpha} f(x); s\} = \frac{\Gamma(1 + \zeta - s)}{\Gamma(1 + \alpha + \zeta - s)} \quad (5.5)$$

for $\Re(\alpha) > 0, \Re(\zeta - s) > -1$ agreeing with the corresponding Mellin transform in the scalar case.

Corollary 5.1.1. The M -transform of $|X|^{-\alpha} {}_0D_X^{-\alpha} f(X)$ is given by

$$M\{|X|^{-\alpha} {}_0D_X^{-\alpha} f(X); s\} = \frac{\Gamma_p(\frac{p+1}{2} - s)}{\Gamma_p(\frac{p+1}{2} + \alpha - s)} f^*(s) \quad (5.6)$$

for $\Re(\alpha) > \frac{p-1}{2}, \Re(s) < 1$.

The proof is parallel to that in Theorem 5.1.

Let us treat a Kober operator of the first kind as a statistical density. Let X_2 have an arbitrary real matrix-variate density $f(X_2)$ and X_1 have a real matrix-variate type-1 beta density with parameters (ζ, α) . That is,

$$f_1(X_1) = \frac{\Gamma_p(\zeta + \alpha)}{\Gamma_p(\zeta)\Gamma_p(\alpha)} |X_1|^{\zeta - \frac{p+1}{2}} |I - X_1|^{\alpha - \frac{p+1}{2}}, O < X_1 < I \quad (5.7)$$

for $\Re(\zeta) > \frac{p-1}{2}$, $\Re(\alpha) > \frac{p-1}{2}$ and $f_1(X_1) = 0$ elsewhere. Let X_1 and X_2 be statistically independently distributed. Consider the transformation $X_2 = V, X_1 = V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}$. The Jacobian is given by

$$dX_1 \wedge dX_2 = |V|^{\frac{p+1}{2}} |U|^{-(p+1)} dU \wedge dV.$$

The marginal density of U , denoted by $g_r(U)$ where r designates that it is coming from a ratio, is given by

$$\begin{aligned} g_r(U) &= \frac{\Gamma_p(\zeta + \alpha)}{\Gamma_p(\zeta)\Gamma_p(\alpha)} \int_V |V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}|^{\zeta - \frac{p+1}{2}} \\ &\quad \times |I - V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}|^{\alpha - \frac{p+1}{2}} f(V) |V|^{\frac{p+1}{2}} |U|^{-(p+1)} dV \\ &= \frac{\Gamma_p(\zeta + \alpha)}{\Gamma_p(\zeta)\Gamma_p(\alpha)} |U|^{-\zeta - \alpha} \int_{V < U} |U - V|^{\alpha - \frac{p+1}{2}} |V|^{\zeta} f(V) dV. \end{aligned}$$

Therefore

$$\begin{aligned} \frac{\Gamma_p(\zeta)}{\Gamma_p(\zeta + \alpha)} g_r(U) &= \frac{|U|^{-\zeta - \alpha}}{\Gamma_p(\alpha)} \int_{V < U} |V|^{\zeta} f(V) dV \\ &= I_X^{\zeta, \alpha} f(X). \end{aligned} \quad (5.8)$$

This is Kober operator of the first kind in the real matrix-variate case and it can be considered as a constant multiple of a real matrix-variate statistical density.

One can also consider a pathway extension for the real matrix-variate Kober operator of the first kind.

5.2. Pathway Extension of Kober Operator of the First Kind in the Matrix Case

Consider the following pathway modified form of the density for X_1 . That is,

$$f_1(X_1) = C_2 |X_1|^{\gamma - \frac{p+1}{2}} |I - a(1-q)X_1|^{\frac{\eta}{1-q}}, I - a(1-q)X_1 > O \quad (5.9)$$

for $q < 1, a > 0, \eta > 0$ where

$$C_2 = \frac{[a(1-q)]^{p\gamma} \Gamma_p(\gamma + \frac{\eta}{1-q} + \frac{p+1}{2})}{\Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2}) \Gamma_p(\gamma)}. \quad (5.10)$$

Consider the same type of transformation as before: $X_2 = V, X_1 = V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}$. The marginal density of U , denoted by $g_p(U)$, is given by

$$\begin{aligned} g_p(U) &= C_2 \int_V |V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}|^{\gamma - \frac{p+1}{2}} |I - a(1-q)V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}|^{\frac{\eta}{1-q}} \\ &\quad \times f(V) |V|^{\frac{p+1}{2}} |U|^{-(p+1)} dV \\ &= C_2 |U|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} \int_{U > a(1-q)V} |U - a(1-q)V|^{\frac{\eta}{1-q}} |V|^{\gamma} f(V) dV. \end{aligned} \quad (5.11)$$

Then

$$\begin{aligned} \Gamma_p(\gamma) g_p(U) &= \frac{[a(1-q)]^{p\gamma} \Gamma_p(\gamma + \frac{\eta}{1-q} + \frac{p+1}{2})}{\Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2})} |U|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} \\ &\quad \times \int_{U > a(1-q)V} |U - a(1-q)V|^{\frac{\eta}{1-q}} |V|^{\gamma} f(V) dV \end{aligned} \quad (5.12)$$

The right side of (5.12) is the pathway extension of Kober operator of the first kind. The right side divided by $\Gamma_p(\gamma)$ is also a statistical density of a type of ratio of independently distributed matrix-variate random variables.

Note that for $a = 1, q = 0, \frac{\eta}{1-q} + \frac{p+1}{2} = \alpha$, (5.12) reduces to the special case (5.7) for $\gamma = \zeta$. Thus, (5.12) describes a vast family of operators which can all be considered as generalizations of the Kober operator of the first kind in the real matrix-variate case. The limiting form when $q \rightarrow 1_-$ is available from the structure in (5.11). Note that

$$\lim_{q \rightarrow 1_-} |I - a(1-q)V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}|^{\frac{\eta}{1-q}} = e^{-a\eta \operatorname{tr}(V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}})} \quad (5.13)$$

Hence

$$\begin{aligned} \lim_{q \rightarrow 1_-} g_p(U) &= \left(\lim_{q \rightarrow 1_-} C_2 \right) \int_{V > O} |U|^{-\gamma - \frac{p+1}{2}} |V|^\gamma \\ &\quad \times e^{-a\eta \operatorname{tr}(V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}})} f(V) dV \end{aligned} \quad (5.14)$$

where

$$\lim_{q \rightarrow 1_-} C_2 = \frac{(a\eta)^{p\gamma}}{\Gamma_p(\gamma)}.$$

That is,

$$\begin{aligned} \lim_{q \rightarrow 1_-} g_p(U) &= \frac{(a\eta)^{p\gamma}}{\Gamma_p(\gamma)} |U|^{-\gamma - \frac{p+1}{2}} \\ &\quad \times \int_{V > O} |V|^\gamma e^{-a\eta \operatorname{tr}(V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}})} f(V) dV. \end{aligned} \quad (5.15)$$

In this case also one can replace the parameter a in $f_1(X_1)$ by a constant positive definite matrix $A > O$. Then $f_1(X_1)$ denoted by $f_{1A}(X_1)$ can be written as

$$f_{1A}(X_1) = C_2(A) |X_1|^{\gamma - \frac{p+1}{2}} |I - (1-q)A^{\frac{1}{2}}X_1A^{\frac{1}{2}}|^{\frac{\eta}{1-q}}$$

where

$$C_2(A) = \frac{(1-q)^{p\gamma} |A|^\gamma \Gamma_p(\gamma + \frac{\eta}{1-q} + \frac{p+1}{2})}{\Gamma_p(\gamma) \Gamma_p(\frac{\eta}{1-q} + \frac{p+1}{2})}.$$

Then the density of $U = V^{\frac{1}{2}}U^{-1}V^{\frac{1}{2}}$, denoted by $g_A(U)$, is given by

$$\begin{aligned} g_A(U) &= C_2(A) |U|^{-\gamma - (\frac{\eta}{1-q} + \frac{p+1}{2})} \\ &\quad \times \int_{U > (1-q)V^{\frac{1}{2}}AV^{\frac{1}{2}}} |V|^\gamma |U - (1-q)V^{\frac{1}{2}}AV^{\frac{1}{2}}|^{\frac{\eta}{1-q}} f(V) dV. \end{aligned} \quad (5.16)$$

Acknowledgment

The authors would like to thank the Department of Science and Technology, Government of India, New Delhi, for the financial assistance for this work under project number SR/S4/MS:287/05.

References

- Mathai, A.M. (1997): *Jacobians of Matrix Transformations and Functions of Matrix Argument*, World Scientific Publishing, New York.
- Mathai, A.M. and Haubold, H.J. (2008): *Special Functions for Applied Scientists*, Springer, New York.
- Mathai, A.M., Provost, S.B., and Hayakawa, T. (1995): *Bilinear Forms and Zonal Polynomials*, Springer, New York.