### Fractional Operators in the Matrix Variate Case

A.M. MATHAI

Centre for Mathematical Sciences, Arunapuram P.O., Pala, Kerala-68674, India, and Department of Mathematics and Statistics, McGill University, Montreal, Quebec, Canada, H3A 2K6

and

H.J. HAUBOLD

Office for Outer Space Affairs, United Nations P.O. Box 500, Vienna International Centre A - 1400 Vienna, Austria, and

Centre for Mathematical Sciences, Arunapuram P.O., Pala, Kerala-68674, India

### [This paper is dedicated to Professor Dr. Francesco Mainardi at his 70th birthday]

### Abstract

Fractional integral operators connected with real-valued scalar functions of matrix argument are applied in problems of mathematics, statistics and natural sciences. In this article we start considering the case of a Gauss hypergeometric function with the argument being a rectangular matrix. Subsequently some fractional integral operators are introduced which complement these results available on fractional operators in the matrix variate cases. Several properties and limiting forms are derived. Then the pathway idea is incorporated to move among several different functional forms. When these are used as models for problems in the natural sciences then these can cover the ideal situations, neighborhoods, in between stages and paths leading to optimal situations.

Mathematical Subject Classification 2010: 26A33, 33C60, 33E12, 33E20

Key Words and Phrases: fractional calculus, fractional operators, multivariate and matrix-variate functions, Gauss hypergeometric function

### 1. Introduction

The importance of anomalous reaction/relaxation and transport/diffusion is well recognized in many disciplines including physics, chemistry, biology, and engineering. Despite this fact, anomalous relaxation and transport are not well understood, and there is the need to develop mathematical and statistical models with predictive power. Recent developments of fractional calculus in terms of integro-differential operators that provide a unifying framework to model key aspects of anomalous relaxation and transport like non-locality, non-Markovian (memory) effects, and non-Gaussian (Levy) processes became available (cf. Mathai et al. [11], Nair [4]). The extension of such developments to matrix-variate statistical densities in general (Mathai and Haubold [8], Mainardi [13]) and to the specifically interesting case of Mittag-Leffler functions and matrix-variate analogues (Mathai [7]) have been achieved more recently. Such results are applicable to the solution of linear coupled fractional differential equations (Lim et al. [1]) as well as to the handling of fractional Poisson probability distributions (Laskin [12]). Further use of fractional integral operators connected with real-valued scalar functions of matrix argument can be utilized for fractional matrix calculus (Phillips [3]), numerical solution of fractional diffusion-wave equations (Garg and Manohar [6]), stability analysis of fractional-order systems (Jiao and Chen [9]), and probably for the application of the matrix-variate Mellin transform in radar image processing (Anfinsen and Eltoft [10]).

Let  $X = (x_{ij})$  be a  $p \times r$ ,  $r \ge p$  matrix of real distinct scalar variables  $x_{ij}$ 's. Let A be a  $p \times p$  real positive definite constant matrix, that is, A = A' > O, prime denoting the transpose. Let B be a constant

positive definite  $r \times r$  matrix, B > O. Let  $A^{\frac{1}{2}}$  and  $B^{\frac{1}{2}}$  denote the positive definite square roots of A and B respectively. Let X be of full rank p. Then  $Z_X = A^{\frac{1}{2}}XBX'A^{\frac{1}{2}}$  is symmetric positive definite matrix. In this paper we will consider only real matrices. The corresponding results in the complex domain can be done parallel to the real case. The following standard notations will be used. Real matrix variate gamma will be denoted by

$$\Gamma_p(\alpha) = \pi^{\frac{p(p-1)}{4}} \Gamma(\alpha) \Gamma(\alpha - \frac{1}{2}) \dots \Gamma(\alpha - \frac{j-1}{2}), \qquad (1.1)$$

for  $\Re(\alpha) > \frac{p-1}{2}$  where  $\Re(\cdot)$  denotes the real part of  $(\cdot)$ . It can be shown that  $\Gamma_p(\alpha)$  has the integral representation

$$\Gamma_p(\alpha) = \int_{S>O} |S|^{\alpha - \frac{p+1}{2}} \mathrm{e}^{-\mathrm{tr}(S)} \mathrm{d}S, \ \Re(\alpha) > \frac{p-1}{2}$$
(1.2)

where |S| means the determinant of the  $p \times p$  positive definite matrix S and tr(S) denotes the trace of S. The wedge product of the  $\frac{p(p+1)}{2}$  differentials  $dx_{ij}$ 's will be denoted by

$$\mathrm{d}S = \prod_{i \ge j} \wedge \mathrm{d}s_{ij}, \ \wedge = \ \mathrm{wedge} \tag{1.4}$$

The type-1 beta integral is given by

$$B_p(\alpha,\beta) = \frac{\Gamma_p(\alpha)\Gamma_p(\beta)}{\Gamma_p(\alpha+\beta)} = \int_{O < X < I} |S|^{\alpha-\frac{p+1}{2}} |I-S|^{\beta-\frac{p+1}{2}} \mathrm{d}S, \tag{1.5}$$

for  $\Re(\alpha) > \frac{p-1}{2}$ ,  $\Re(\beta) > \frac{p-1}{2}$ , where S is  $p \times p$  positive definite and O < S < I means S > O, I - S > O or all the eigenvalues of S are in the open interval (0, 1). In general,  $\int_X$  means the integral over X. The type-2 beta integral is given by

$$B_{p}(\alpha,\beta) = \int_{S>O} |S|^{\alpha-\frac{p+1}{2}} |I+S|^{-(\alpha+\beta)} \mathrm{d}S$$
  
= 
$$\int_{U>O} |U|^{\beta-\frac{p+1}{2}} |I+U|^{-(\alpha+\beta)} \mathrm{d}U, \Re(\alpha) > \frac{p-1}{2}, \Re(\beta) > \frac{p-1}{2}$$
(1.6)

Let

$$Z_X = A^{\frac{1}{2}} X B X' A^{\frac{1}{2}}$$
 and  $Z_Y = A^{\frac{1}{2}} Y B Y' A^{\frac{1}{2}}$ 

where  $X = (x_{ij})$  and  $Y = (y_{ij})$  are  $p \times r, r \ge p$  matrices of real elements, and of full rank p. Consider the evaluation of the integral

$$I_1 = \int_X |Z_X|^a |I - Z_X|^{c-a - \frac{p+1}{2}} |I - Z_Y Z_X|^{-b} \mathrm{d}X,$$
(1.7)

for  $O < Z_X < I, O < Z_Y < I$ . This integral corresponds to the Euler integral for Gauss hypergeometric function. Let

$$U = A^{\frac{1}{2}} X B^{\frac{1}{2}} \Rightarrow dU = |A|^{\frac{r}{2}} |B|^{\frac{p}{2}} dX \text{ (see Mathai, 1997, equation (1.2.20))}$$
(1.8)

Then, after integration over the Stiefel manifold,

$$Z_X = UU' = V \Rightarrow \mathrm{d}U = \frac{\pi^{\frac{rp}{2}}}{\Gamma_p(\frac{r}{2})} |V|^{\frac{r}{2} - \frac{p+1}{2}} \mathrm{d}V \tag{1.9}$$

(see Mathai, 1997, Theorem 2.16 and Remark 2.13). Now, the integral  $I_1$  becomes,

$$I_{1} = |A|^{-\frac{r}{2}} |B|^{-\frac{p}{2}} \frac{\pi^{\frac{rp}{2}}}{\Gamma_{p}(\frac{r}{2})} \times \int_{O < V < I} |V|^{a + \frac{r}{2} - \frac{p+1}{2}} |I - V|^{c-a - \frac{p+1}{2}} |I - Z_{Y}^{\frac{1}{2}} V Z_{Y}^{\frac{1}{2}}|^{-b} \mathrm{d}V.$$
(1.10)

Let us expand the last factor in the integrand in terms of zonal polynomials. For a discussion of zonal polynomials see Mathai et al. (1995).

$$|I - Z_Y^{\frac{1}{2}} V Z_Y^{\frac{1}{2}}|^{-b} = \sum_{k=0}^{\infty} \sum_K \frac{(b)_K}{k!} C_K(Z_Y V)$$
(1.11)

where  $C_K(\cdot)$  is the zonal polynomial of order  $k, K = (k_1, k_2, ..., k_p), k_1 + ... + k_p = k$  and

$$(b)_{K} = \prod_{j=1}^{p} (b - \frac{j-1}{2})_{k_{j}} = \frac{\Gamma_{p}(b,K)}{\Gamma_{p}(b)},$$
  
$$\Gamma_{p}(b,K) = \pi^{\frac{p(p-1)}{4}} \prod_{j=1}^{p} \Gamma(b+k_{j} - \frac{j-1}{2}) = \Gamma_{p}(b)(b)_{K}$$
(1.12)

and  $(a)_m$  is the Pochhammer symbol

$$(a)_m = a(a+1)...(a+m-1), (a)_0 = 1, a \neq 0.$$

We can evaluate the integral

$$\int_{O < V < I} |V|^{a + \frac{r}{2} - \frac{p+1}{2}} |I - V|^{c-a - \frac{p+1}{2}} C_K(Z_Y V) dV = \frac{\Gamma_p(a + \frac{r}{2}, K) \Gamma_p(c-a)}{\Gamma_p(c + \frac{r}{2}, K)} C_K(Z_Y)$$
(1,13)

(see Mathai, 1997, (5.1.26)). Note that interchange of integrals and sums is valid here. Then

$$I_{1} = |A|^{-\frac{r}{2}} |B|^{-\frac{p}{2}} \frac{\pi^{\frac{rp}{2}}}{\Gamma_{p}(\frac{r}{2})} \frac{\Gamma_{p}(a+\frac{r}{2})\Gamma_{p}(c-a)}{\Gamma_{p}(c+\frac{r}{2})} \times \sum_{k=0}^{\infty} \sum_{K} \frac{(b)_{K}(a+\frac{r}{2})_{K}}{(c+\frac{r}{2})_{K}} \frac{C_{K}(Z_{Y})}{k!}.$$
(1.14)

## 2. Hypergeometric Functions of Rectangular Matrix Argument

In the notations of hypergeometric function of matrix argument the series part of (1.14) can be written as a  $_2F_1$ . That is,

$$\sum_{k=0}^{\infty} \sum_{K} \frac{(b)_{K}(a+\frac{r}{2})_{K}}{(c+\frac{r}{2})_{K}} \frac{C_{K}(Z_{Y})}{k!} = {}_{2}F_{1}(a+\frac{r}{2},b;c+\frac{r}{2};Z_{Y}), \ \|Z_{Y}\| < 1$$
(2.1)

where  $\|(\cdot)\|$  denotes a norm of  $(\cdot)$ . Hence we have the following theorem:

**Theorem 2.1.** For X, Y, A, B as defined in Section 1

$${}_{2}F_{1}(a + \frac{r}{2}, b; c + \frac{r}{2}; Z_{Y}) = \frac{|A|^{\frac{r}{2}} |B|^{\frac{p}{2}} \Gamma_{p}(\frac{r}{2}) \Gamma_{p}(c + \frac{r}{2})}{\pi^{\frac{rp}{2}} \Gamma_{p}(a + \frac{r}{2}) \Gamma_{p}(c - a)} \times \int_{X} |Z_{X}|^{a} |I - Z_{X}|^{c - a - \frac{p+1}{2}} |I - Z_{Y} Z_{X}|^{-b} \mathrm{d}X$$

$$(2.2)$$

for  $O < Z_X < I, O < Z_Y < I, \Re(c-a) > \frac{p-1}{2}, \Re(a) > -\frac{r}{2} + \frac{p-1}{2}.$ 

For establishing some limiting forms and pathways we need the following results, which will be stated as lemmas.

Lemma 2.1. For  $K = (k_1, ..., k_p), k_1 + ... + k_p = k, (a)_K = \prod_{j=1}^p (a - \frac{j-1}{2})_{k_j},$  $\lim_{q \to 1} (q-1)^k (\frac{1}{q-1})_K = 1.$ 

Note that

$$(q-1)^{k} \left(\frac{1}{q-1}\right)_{K} = \left\{\prod_{j=1}^{p} (q-1)^{k_{j}}\right\} \left\{\prod_{i=1}^{k_{j}} \left(\frac{1}{q-1} - \left(\frac{j-1}{2}\right) + i - 1\right)\right\}$$
$$= \prod_{j=1}^{p} \left[1 - (q-1)\left(\frac{j-1}{2}\right) + (q-1)(i-1)\right] = 1.$$

Lemma 2.2.

$$\lim_{q \to 1} |I + (q-1)Z_Y|^{-\frac{1}{q-1}} = e^{-\operatorname{tr}(Z_Y)}.$$
(2.4)

**Proof:** Let  $\lambda_j$ , j = 1, ..., p be the eigenvalues of  $Z_Y$ . Then

$$|I + (q-1)Z_Y| = \prod_{j=1}^p (1 + (q-1)\lambda_j)$$

But

$$\prod_{j=1}^{p} [\lim_{q \to 1} \{1 + (q-1)\lambda_j\}^{-\frac{1}{q-1}}] = \prod_{j=1}^{p} e^{-\lambda_j} = e^{-\sum_{j=1}^{p} \lambda_j}$$
$$= e^{-\operatorname{tr}(Z_Y)}.$$
(2.5)

## 3. Fractional Integral Operators

Let  $Z_X$  and  $Z_Y$  be as defined in Section 1. Let the generalized fractional integral operator of matrix argument be defined and denoted as

$$\begin{aligned} (_{0}D_{X}^{-\alpha}f)(X) &= \frac{1}{\Gamma_{p}(\alpha)} \int_{Z_{X}>Z_{Y}>O} |Z_{X} - Z_{Y}|^{\alpha - \frac{p+1}{2}} f(Z_{Y}) \mathrm{d}Y \\ &= \frac{|Z_{X}|^{\alpha - \frac{p+1}{2}}}{\Gamma_{p}(\alpha)} \int_{Z_{X}>Z_{Y}>O} |I - Z_{X}^{-\frac{1}{2}}Z_{Y}Z_{X}^{-\frac{1}{2}}|^{\alpha - \frac{p+1}{2}} f(Z_{Y}) \mathrm{d}Y. \end{aligned}$$

Make the transformations  $U = A^{\frac{1}{2}}YB^{\frac{1}{2}}$ , V = UU',  $W = Z_X^{-\frac{1}{2}}VZ_X^{-\frac{1}{2}}$ . Then integrating out over the Stiefel manifold we have

$${}_{0}D_{X}^{-\alpha}f = \frac{|Z_{X}|^{\alpha + \frac{r}{2} - \frac{p+1}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\alpha)} \frac{\pi^{\frac{rp}{2}}}{\Gamma_{p}(\frac{r}{2})} \times \int_{O < W < I} |I - W|^{\alpha - \frac{p+1}{2}} f(Z_{X}^{\frac{1}{2}}WZ_{X}^{\frac{1}{2}})|W|^{\frac{r}{2} - \frac{p+1}{2}} dW.$$
(3.1)

Consider the special cases of  $f(\cdot)$ . Consider the operator operating on a power function.

Case 1. Let  $f(Z_X) = |Z_X|^{\eta}$ . Then

$${}_{0}D_{X}^{-\alpha}|Z_{X}|^{\eta} = \frac{|Z_{X}|^{\alpha+\frac{r}{2}+\eta-\frac{p+1}{2}}\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\alpha)\Gamma_{p}(\frac{r}{2})} \int_{O
$$= \frac{|Z_{X}|^{\alpha+\frac{r}{2}+\eta-\frac{p+1}{2}}\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\frac{r}{2})} \frac{\Gamma_{p}(\frac{r}{2}+\eta)}{\Gamma_{p}(\alpha+\frac{r}{2}+\eta)}, \ \Re(\frac{r}{2}+\eta) > \frac{p-1}{2}.$$
(3.2)$$

Case 2. Zonal polynomial of order  $k, f(Z_X) = C_K(Z_X)$ 

Then going through the same steps as above

$${}_{0}D_{X}^{-\alpha}C_{K}(Z_{X}) = \frac{|Z_{X}|^{\alpha+\frac{r}{2}-\frac{p+1}{2}}\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\alpha)\Gamma_{p}(\frac{r}{2})} \int_{O < W < I} |W|^{\frac{r}{2}-\frac{p+1}{2}}$$

$$\times |I - W|^{\alpha-\frac{p+1}{2}}C_{K}(Z_{X}^{\frac{1}{2}}WZ_{X}^{\frac{1}{2}})dW, \text{ (Mathai, 1997, equation (5.1.26))}$$

$$= \frac{|Z_{X}|^{\alpha+\frac{r}{2}-\frac{p+1}{2}}\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\alpha)\Gamma_{p}(\frac{r}{2})} \frac{\Gamma_{p}(\frac{r}{2},K)\Gamma_{p}(\alpha)}{\Gamma_{p}(\alpha+\frac{r}{2},K)}C_{K}(Z_{X})$$

$$= \frac{|Z_{X}|^{\alpha+\frac{r}{2}-\frac{p+1}{2}}\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\alpha+\frac{r}{2})} \frac{(\frac{r}{2})_{K}C_{K}(Z_{X})}{(\alpha+\frac{r}{2})_{K}}$$

# 4. Extended Saigo Operators

Let

$$f(Z_X) = {}_2F_1(a,b;c;I - Z_X^{-\frac{1}{2}}Z_YZ_X^{-\frac{1}{2}})\phi(Z_Y), Z_X > Z_Y.$$

Here  $_2F_1$  is a Gauss hypergeometric function of matrix argument  $I - Z_X^{-\frac{1}{2}} Z_Y Z_X^{-\frac{1}{2}}$  and  $\phi(Z_Y)$  is an arbitrary function so that

$$\int_{Z_X > Z_Y} |I - Z_X^{-\frac{1}{2}} Z_Y Z_X^{-\frac{1}{2}}|^{\alpha - \frac{p+1}{2}} C_K (I - Z_X^{-\frac{1}{2}} Z_Y Z_X^{-\frac{1}{2}}) \phi(Z_Y) \mathrm{d}Y < \infty.$$

Opening up the  $_2F_1$  in terms of zonal polynomials and then substituting  $U = A^{\frac{1}{2}}YBA^{\frac{1}{2}}$ , V = UU' we have

$${}_{0}D_{X}^{-\alpha}[{}_{2}F_{1}(a,b;c;I-Z_{X}^{-\frac{1}{2}}Z_{Y}Z_{X}^{-\frac{1}{2}})\phi(Z_{Y})]$$

$$=\sum_{k=0}^{\infty}\sum_{K}\frac{(a)_{K}(b)_{K}}{k!(c)_{K}}\frac{|Z_{X}|^{\alpha-\frac{p+1}{2}}\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_{p}(\alpha)\Gamma_{p}(\frac{r}{2})}$$

$$\int_{O< V
(4.1)$$

Consider the special case  $\phi(V) = |V|^{\eta}$  and then make the transformation  $Z_X^{-\frac{1}{2}}VZ_X^{-\frac{1}{2}} = W$ . Then the right side of (4.1) reduces to the following:

$$\sum_{k=0}^{\infty} \sum_{K} \frac{(a)_{K}(b)_{K}}{k!(c)_{K}} \frac{|Z_{X}|^{\alpha + \frac{r}{2} + \eta - \frac{p+1}{2}} \pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}} |B|^{\frac{p}{2}} \Gamma_{p}(\alpha) \Gamma_{p}(\frac{r}{2})} \int_{O < W < I} |I - W|^{\alpha - \frac{p+1}{2}} |W|^{\eta + \frac{r}{2} - \frac{p+1}{2}} C_{K}(I - W) \mathrm{d}W.$$

The integral part reduces to the following:

$$\int_{O < W < I} |W|^{\eta + \frac{r}{2} - \frac{p+1}{2}} |I - W|^{\alpha - \frac{p+1}{2}} C_K (I - W) dW$$
  
= 
$$\int_{O < T < I} |T|^{\alpha - \frac{p+1}{2}} |I - T|^{\eta + \frac{r}{2} - \frac{p+1}{2}} C_K (T) dT$$
  
= 
$$\frac{\Gamma_p(\alpha, K) \Gamma_p(\eta + \frac{r}{2})}{\Gamma_p(\alpha + \eta + \frac{r}{2}, K)} C_K (I)$$
  
= 
$$\frac{\Gamma_p(\alpha) \Gamma_p(\eta + \frac{r}{2})}{\Gamma_p(\alpha + \eta + \frac{r}{2})} \frac{(\alpha)_K}{(\alpha + \eta + \frac{r}{2})_K} C_K (I).$$

Substituting back and denoting the left side by  $I_X$  we have

$$I_X = \frac{\pi^{\frac{rp}{2}}}{|A|^{\frac{r}{2}}|B|^{\frac{p}{2}}\Gamma_p(\frac{r}{2})} \frac{\Gamma_p(\eta + \frac{r}{2})}{\Gamma_p(\alpha + \eta + \frac{r}{2})} |Z_X|^{\alpha + \eta + \frac{r}{2} - \frac{p+1}{2}} \times {}_{3}F_2(a, b, \alpha; c, \alpha + \eta + \frac{r}{2}; I).$$

### 5. Some Statistical Considerations

Let  $U_1 = A_1^{\frac{1}{2}} X_1 B_1 X_1' A_1^{\frac{1}{2}}$ ,  $U_2 = A_2^{\frac{1}{2}} X_2 B_2 X_2' A_2^{\frac{1}{2}}$ . Let  $X_1$  and  $X_2$  be independently and exponentially distributed random matrices where  $A_1$  and  $A_2$  are  $p \times p$  real symmetric positive definite constant matrices,  $B_1$  is  $r_1 \times r_1$ ,  $r_1 \ge p$  and  $B_2$  is  $r_2 \times r_2$ ,  $r_2 \ge p$  constant positive definite matrices,  $X_1$  is  $p \times r_1$  and  $X_2$  is  $p \times r_2$  matrices of distinct real scalar random variables and the matrices be of full rank p, where a prime denotes the transpose and the square roots are the unique positive definite square roots of  $A_1$  and  $A_2$  respectively. Let the densities of  $X_1$  and  $X_2$  be denoted by  $f_1(X_1)$  and  $f_2(X_2)$  and the joint density, denoted by  $f(X_1, X_2) = f_1(X_1)f_2(X_2)$  due to independence. Further, let

$$f_j(X_j) \mathrm{d}X_j = c_j \mathrm{e}^{-\mathrm{tr}(A_j^{\frac{1}{2}} X_j B_j X_j' A_j^{\frac{1}{2}})} \mathrm{d}X_j, j = 1, 2$$
(5.1)

where  $c_j$  is the normalizing constant. Make the transformations  $V_j = A_j^{\frac{1}{2}} X_j B_j^{\frac{1}{2}} \Rightarrow dX_j = |A_j|^{-\frac{r_j}{2}} |B_j|^{-\frac{p}{2}} dV_j$ where  $dX_j$  is the wedge product of all differentials in  $X_k = (x_{ij}^{(k)})$  or  $dX_k = \prod_{i=1}^p \prod_{j=1}^{r_k} \wedge dx_{ij}^{(k)}$ ,  $\wedge =$  wedge product. Let  $W_j = V_j V_j'$  and integrate out over the Stiefel manifold. Then we have

$$dV_j = \frac{\pi^{\frac{r_j p}{2}}}{\Gamma_p(\frac{r_j}{2})} |W_j|^{\frac{r_j}{2} - \frac{p+1}{2}} dW_j$$
(5.2)

and tr(·) denotes the trace of (·). Hence the rectangular matrix  $X_j$  having the exponential density means  $W_j$  is having a real matrix-variate gamma density and further,

$$dX_j = \frac{\pi^{\frac{r_j p}{2}} |W_j|^{\frac{r_j}{2} - \frac{p+1}{2}}}{|A_j|^{\frac{r_j}{2}} |B_j|^{\frac{p}{2}} \Gamma_p(\frac{r_j}{2})} dW_j.$$
(5.3)

If the densities of  $V_j$  and  $W_j$  are denoted by  $g_j(V_j)$  and  $h_j(W_j)$  respectively then

$$h_j(W_j) \mathrm{d}W_j = \frac{1}{\Gamma_p(\frac{r_j}{2})} |W_j|^{\frac{r_j}{2} - \frac{p+1}{2}} \mathrm{e}^{-\mathrm{tr}(W_j)} \mathrm{d}W_j, W_j > O$$
(5.4)

$$g_j(V_j) dV_j = \frac{1}{\pi^{\frac{r_j p}{2}}} e^{-\operatorname{tr}(V_j V_j')} dV_j$$
(5.5)

$$f_j(X_j) \mathrm{d}X_j = \frac{|A_j|^{\frac{r_j}{2}} |B_j|^{\frac{p}{2}}}{\pi^{\frac{r_jp}{2}}} \mathrm{e}^{-\mathrm{tr}(A_j^{\frac{1}{2}} X_j B_j X_j' A_j^{\frac{1}{2}})} \mathrm{d}X_j.$$
(5.6)

#### 5.1. Density of the sum

Let us examine the density of the sum  $U_1 + U_1$ . The joint density of  $X_1$  and  $X_2$ , denoted by  $f(X_1, X_2)$ , is given by

$$\begin{split} f(X_1, X_2) \mathrm{d}X_1 \wedge \mathrm{d}X_2 &= f_1(X_1) f_2(X_2) \mathrm{d}X_1 \wedge \mathrm{d}X_2. \\ \text{Make the transformations } V_j &= A_j^{\frac{1}{2}} X_j B_j^{\frac{1}{2}}, W_j = V_j V_j' \text{ then} \\ f(X_1, X_2) \mathrm{d}X_1 \wedge \mathrm{d}X_2 &= f_1(V_1 V_1') f_2(V_2 V_2') \{\prod_{j=1}^2 |A_j|^{-\frac{r_j}{2}} |B_j|^{-\frac{p}{2}} \} \mathrm{d}V_1 \wedge \mathrm{d}V_2 \\ &= f_1(W_1) f_2(W_2) \{\prod_{j=1}^2 |A_j|^{-\frac{r_j}{2}} |B_j|^{-\frac{p}{2}} \frac{\pi^{\frac{r_j}{2}}}{\Gamma_p(\frac{r_j}{2})} |W_j|^{\frac{r_j}{2} - \frac{p+1}{2}} \} \mathrm{d}W_1 \wedge \mathrm{d}W_2. \end{split}$$

Then  $U_1 + U_2 = W_1 + W_2$ . Make the transformation  $U = W_1 + W_2$  and  $V = W_1$ , the Jacobian is unity and  $W_1 = V, W_2 = U - V$  and the integration is over the positive definite matrices U and V with U - V > O. If  $X_1$  and  $X_2$  are independently distributed as in (5.1) then denoting the marginal density of U by  $f^*(U)$  we have

$$f^{*}(U)dU = \int_{U>V>0} f_{1}(U-V)f_{2}(V)dV$$
(5.7)

$$= \frac{1}{\Gamma_p(\frac{r_1}{2})\Gamma_p(\frac{r_2}{2})} |U - V|^{\frac{r_1}{2} - \frac{p+1}{2}} |V|^{\frac{r_2}{2} - \frac{p+1}{2}} e^{-\operatorname{tr}(U - V) - \operatorname{tr}(V)} dV \wedge dV.$$
(5.8)

Make the transformation  $Z = U^{-\frac{1}{2}}VU^{-\frac{1}{2}}$  for fixed U, then

$$f^{*}(U)dU = \frac{|U|^{\frac{r_{1}+r_{2}}{2}-\frac{p+1}{2}}e^{-\operatorname{tr}(U)}}{\Gamma_{p}(\frac{r_{1}}{2})\Gamma_{p}(\frac{r_{2}}{2})} \int_{Z>O} |Z|^{\frac{r_{2}}{2}-\frac{p+1}{2}}|I-Z|^{\frac{r_{1}}{2}-\frac{p+1}{2}}dZ \wedge dU$$
$$= \frac{|U|^{\frac{r_{1}+r_{2}}{2}-\frac{p+1}{2}}e^{-\operatorname{tr}(U)}}{\Gamma_{p}(\frac{r_{1}+r_{2}}{2})}dU.$$

Thus the sum U is real matrix-variate gamma distributed. Note that the integral in (5.7) is Riemann-Liouville left-sided fractional integral for the power function or where the arbitrary function is of the form  $f(T) = |T|^{\frac{r_2}{2} - \frac{p+1}{2}}$ .

## Acknowledgment

The authors would like to thank the Department of Science and Technology, Government of India, for the financial assistance for this work under Project No. SR/S4/MS:287/05.

## References

[1] Lim, S.C., Eab, C.H., Mak, K.H., Li, M., and Chen, S.Y. (2012): Solving linear coupled fractional differential equations by direct operational method and some applications *Mathematical Problems in Engineering*, doi:10.1155/2012/653939., 28 pages.

[2] Mathai, A.M. (1997): Jacobians of Matrix Transformations and Functions of Matrix Argument, World Scientific Publishing, New York.

[3] Phillips, P.C.B. (1987): Fractional matrix calculus and the distribution of multivariate tests *Time Series* and *Econometric Modeling*, Eds. I.B. MacNeill and G.J. Umphrey, D. Reidel Publishing Company, 219-234.

[4] Nair, S.S. (2009): Pathway fractional integration operator, *Fractional Calculus and Applied Analysis*, **12**, 237-252.

[5] Mathai, A.M., Provost, S.B., and Hayakawa, T. (1995): *Bilinear Forms and Zonal Polynomials*, Springer, New York.

[6] Garg, M. and Manohar, P. (2010): Numerical solution of fractional diffusion-wave equation with two space variables by matrix method, *Fractional Calculus and Applied Analysis*, **13**, 191-207.

[7] Mathai, A.M. (2010): Some properties of Mittag-Leffler functions and Matrix-variate analogues: A statistical perspective, *Fractional Calculus and Applied Analysis*, **13**, 113-132.

[8] Mathai, A.M. and Haubold, H.J. (2011): Matrix-variate statistical distributions and fractional calculus, *Fractional Calculus and Applied Analysis*, **14**, 138-155.

[9] Jiao, Z. and Chen Y.Q. (2011): Stability analysis of fractional-order systems with double non-commensurate orders for matrix case, *Fractional Calculus and Applied Analysis*, **14**, 436-453.

[10] Anfinsen, S.N. and Eltoft, T. (2011): Application of the matrix-variate Mellin transform to analysis of polarimetric radar images, *IEEE Transactions on Geoscience and Remote Sensing*, **49**, 2281-2295.

[11] Mathai, A.M., Saxena, R.K., and Haubold, H.J. (2010): *The H-Function: Theory and Applications*, Springer, New York.

[12] Laskin, N. (2009): Some applications of the fractional Poisson probability distribution, *Journal of Mathematical Physics*, **50**, 113513.

[13] Mainardi, F. (2010): Fractional Calculus and Waves in Linear Viscoelasticity, Imperial College Press, London.