A CONSTRUCTION OF SUBSHIFTS AND A CLASS OF SEMIGROUPS

TOSHIHIRO HAMACHI AND WOLFGANG KRIEGER

ABSTRACT. Subshifts with property (A) are constructed from a class of directed graphs. As special cases the Markov-Dyck shifts are shown to have property (A). The semigroups, that are associated to \mathcal{R} -graph shifts with Property (A), are determined. Also results on the reconstruction of \mathcal{R} -graphs from their \mathcal{R} -graph shifts are obtained.

1. INTRODUCTION

Let Σ be a finite alphabet, and let S be the shift on the shift space $\Sigma^{\mathbb{Z}}$,

$$S((x_i)_{i\in\mathbb{Z}}) = (x_{i+1})_{i\in\mathbb{Z}}, \qquad (x_i)_{i\in\mathbb{Z}} \in \Sigma^{\mathbb{Z}}.$$

An S-invariant closed subset X of $\Sigma^{\mathbb{Z}}$ is called a subshift. For an introduction to the theory of subshifts see [Ki] or [LM]. In [Kr2] a Property (A) of subshifts was introduced that is an invariant of topological conjugacy. Also in [Kr2] a semigroup was constructed that is invariantly attached to a subshift with property (A) (see also [CS, Section 9]). Prototypes of subshifts with Property (A) are the Dyck shifts [Kr1]. To recall the construction of the Dyck shifts, let N > 1, and let $\alpha^{-}(n), \alpha^{+}(n), 0 \leq n < N$, be the generators of the Dyck inverse monoid (the polycyclic monoid [NP]) \mathcal{D}_N , that satisfy the relations

$$\alpha^{-}(n)\alpha^{+}(m) = \begin{cases} 1, & \text{if } n = m, \\ 0, & \text{if } n \neq m. \end{cases}$$

The Dyck shifts are defined as the subshifts

$$D_N \subset (\{\alpha^-(n) : 0 \le n < N\} \cup \{\alpha^+(n) : 0 \le n < N\})^{\mathbb{Z}}$$

with the admissible words $(\sigma_i)_{1 \le i \le I}$, $I \in \mathbb{N}$, of $D_N, N > 1$, given by the condition

$$\prod_{1 \le i \le I} \sigma_i \ne 0.$$

The Dyck inverse monoid \mathcal{D}_N is associated to the Dyck shift D_N .

We denote a finite directed graph with vertex set \mathfrak{P} and edge set \mathcal{E} by $G(\mathfrak{P}, \mathcal{E})$. As notation for the source vertex and target vertex of an edge or path in a directed graph we use s and t. We recall from [Kr4] the notion of an \mathcal{R} -graph. Let there be given a finite directed graph $G(\mathcal{V}, \mathcal{E})$. Assume also given a partition

$$\mathcal{E} = \mathcal{E}^- \cup \mathcal{E}^+$$

We set

$$\begin{split} \mathcal{E}^{-}(\mathfrak{q},\mathfrak{r}) &= \{e^{-} \in \mathcal{E}^{-} : s(e^{-}) = \mathfrak{q}, \ t(e^{-}) = \mathfrak{r}\}, \\ \mathcal{E}^{+}(\mathfrak{q},\mathfrak{r}) &= \{e^{-} \in \mathcal{E}^{+} : s(e^{+}) = \mathfrak{r}, \ t(e^{+}) = \mathfrak{q}\}, \qquad \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}. \end{split}$$

We assume that $\mathcal{E}^{-}(\mathfrak{q},\mathfrak{r}) \neq \emptyset$ if and only if $\mathcal{E}^{+}(\mathfrak{q},\mathfrak{r}) \neq \emptyset, \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}$, and we assume that the directed graph $G(\mathfrak{P}, \mathcal{E}^{-})$ is strongly connected, or, equivalently, that the

directed graph $G(\mathfrak{P}, \mathcal{E}^+)$ is strongly connected. We call $G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ a partitioned directed graph. Let there further be given relations ¹

$$\mathcal{R}(\mathfrak{q},\mathfrak{r})\subset\mathcal{E}^{-}(\mathfrak{q},\mathfrak{r}) imes\mathcal{E}^{+}(\mathfrak{q},\mathfrak{r}),\qquad\mathfrak{q},\mathfrak{r}\in\mathfrak{P},$$

and set

$$\mathcal{R} = \bigcup_{\mathfrak{q},\mathfrak{r}\in\mathfrak{P}} \mathcal{R}(\mathfrak{q},\mathfrak{r}).$$

The resulting structure, that we call an \mathcal{R} -graph, we denote by $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$. We also recall the construction of a semigroup (with zero) $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ from an \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ as described in [Kr3]. The semigroup $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ contains idempotents $\mathbf{1}_{\mathfrak{p}}, \mathfrak{p} \in \mathfrak{P}$, and has \mathcal{E} as a generating set. Besides $\mathbf{1}_{\mathfrak{p}}^2 = \mathbf{1}_{\mathfrak{p}}, \mathfrak{p} \in \mathfrak{P}$, the defining relations are:

$$f^-g^+ = \mathbf{1}_{\mathfrak{q}}, \quad f^- \in \mathcal{E}^-(\mathfrak{q}, \mathfrak{r}), g^+ \in \mathcal{E}^+(\mathfrak{q}, \mathfrak{r}), (f^-, g^+) \in \mathcal{R}(\mathfrak{q}, \mathfrak{r}), \quad \mathfrak{q}, \mathfrak{r} \in \mathfrak{P},$$

and

$$\begin{split} \mathbf{1}_{\mathfrak{q}} e^- &= e^- \mathbf{1}_{\mathfrak{r}} = e^-, \quad e^- \in \mathcal{E}^-(\mathfrak{q}, \mathfrak{r}), \\ \mathbf{1}_{\mathfrak{r}} e^+ &= e^+ \mathbf{1}_{\mathfrak{q}} = e^+, \quad e^+ \in \mathcal{E}^+(\mathfrak{q}, \mathfrak{r}), \quad \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}, \end{split}$$

$$f^{-}g^{+} = \begin{cases} \mathbf{1}_{\mathfrak{q}}, & \text{if } (f^{-}, g^{+}) \in \mathcal{R}(\mathfrak{q}, \mathfrak{r}), \\ 0, & \text{if } (f^{-}, g^{+}) \notin \mathcal{R}(\mathfrak{q}, \mathfrak{r}), \end{cases} f^{-} \in \mathcal{E}^{-}(\mathfrak{q}, \mathfrak{r}), g^{+} \in \mathcal{E}^{+}(\mathfrak{q}, \mathfrak{r}), \ \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}, \end{cases}$$

and

$$\mathbf{1}_{\mathfrak{q}}\mathbf{1}_{\mathfrak{r}}=0, \quad \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}, \mathfrak{q} \neq \mathfrak{r}.$$

We call $\mathcal{S}_{\mathcal{R}}(G(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}))$ an \mathcal{R} -graph semigroup. We write $\mathcal{S}^{-}(\mathfrak{P}, \mathcal{E}^{-})(\mathcal{S}^{+}(\mathfrak{P}, \mathcal{E}^{+}))$ for the set of non-zero elements of the subsemigroup of $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}))$, that is generated by $\mathcal{E}^{-}(\mathcal{E}^{+})$.

Special cases are the graph inverse semigroups of finite directed graphs $G(\mathfrak{P}, \mathcal{E})$ ([AH],[L, Section 10.7.]). With the edge set $\mathcal{E}^- = \{e^- : e_\circ \in \mathcal{E}_\circ\}$ of a copy of $G(\mathfrak{P}, \mathcal{E})$, and with the edge set $\mathcal{E}^+ = \{e^- : e \in \mathcal{E}\}$ of the reversal of $G(\mathfrak{P}, \mathcal{E})$, the graph inverse semigroup $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}))$ of $G(\mathfrak{P}, \mathcal{E})$ is the \mathcal{R} -graph semigroup of the partitioned graph $G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ with the relations

$$\mathcal{R}(\mathfrak{q},\mathfrak{r}) = \{(e^-,e^+) : e \in \mathcal{E}, s(e) = \mathfrak{q}, t(e) = \mathfrak{r}\}, \quad \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}.$$

In [HI] a criterion was given for the existence of an embedding of an irreducible subshift of finite type into a Dyck shift and this result was extended in [HIK] to a larger class of target shifts with Property (A). These target shifts were constructed by a method that presents the subshifts by means of a suitably structured irreducible finite labeled directed graph with labels taken from the inverse semigroup of an irreducible finite directed graph, in which every vertex has at least two incoming edges. This method was extended in [Kr4] by the use of \mathcal{R} -graph semigroups. Following [HIK, Kr4] we describe this construction.

We denote a finite directed labelled graph with vertex set \mathcal{V} , edge set Σ and a label map λ by $G(\mathcal{V}, \Sigma, \lambda)$. Let there be given an \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ and a finite strongly connected labeled directed graph $G(\mathcal{V}, \Sigma, \lambda)$ such that

$$(G 1) \qquad \lambda(\sigma) \in S^{-}(\mathfrak{P}, \mathcal{E}^{-}) \cup \{\mathbf{1}_{\mathfrak{p}} : \mathfrak{p} \in \mathfrak{P}\} \cup S^{+}(\mathfrak{P}, \mathcal{E}^{+}), \qquad \sigma \in \Sigma.$$

The label map λ extends to finite paths $(\sigma_i)_{1 \le i \le I}$ in the graph $G(\mathcal{V}, \Sigma)$ by

$$\lambda((\sigma_i)_{1 \le i \le I}) = \prod_{1 \le i \le I} \lambda(\sigma_i).$$

¹We consider complete heterogeneous relations.

Denoting for $\mathfrak{p} \in \mathfrak{P}$ by $\mathcal{V}(\mathfrak{p})$ the set of $V \in \mathcal{V}$ such that there is a cycle $(\sigma_i)_{1 \leq i \leq I}, I \in \mathbb{N}$, in the graph $G(\mathcal{V}, \Sigma)$ from V to V such that

$$\lambda((\sigma_i)_{1\leq i\leq I})=\mathbf{1}_{\mathfrak{p}}$$

we require the following conditions (G 2 - 5) to be satisfied:

(G 2)
$$\mathcal{V}(\mathfrak{p}) \neq \emptyset, \quad \mathfrak{p} \in \mathfrak{P},$$

(G 3)
$$\{\mathcal{V}(\mathfrak{p}) : \mathfrak{p} \in \mathfrak{P}\}$$
 is a partition of \mathcal{V} .

(G 4) For $V \in \mathcal{V}(\mathfrak{p}), \mathfrak{p} \in \mathfrak{P}$, and for all edges σ that leave $V, \mathbf{1}_{\mathfrak{p}}\lambda(\sigma) \neq 0$, and for all edges σ that enter $V, \lambda(\sigma)\mathbf{1}_{\mathfrak{p}} \neq 0$,

(G 5) For $f \in \mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+})), \mathfrak{q}, \mathfrak{r} \in \mathfrak{P}$, such that $\mathbf{1}_{\mathfrak{q}}f\mathbf{1}_{\mathfrak{r}} \neq 0$, and for $U \in \mathcal{V}(\mathfrak{q}), W \in \mathcal{V}(\mathfrak{r})$, there exists a path b in the labeled directed graph $G(\mathcal{V}, \Sigma, \lambda)$ from U to W such that $\lambda(b) = f$.

A finite labeled directed graph $G(\mathcal{V}, \Sigma, \lambda)$, that satisfies conditions (G 1 - 5), gives rise to a subshift $X(G(\mathcal{V}, \Sigma, \lambda))$, that has as its language of admissible words the set of finite paths b in the graph $G(\mathcal{V}, \Sigma, \lambda)$ such that $\lambda(b) \neq 0$. We call this subshift an $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentation. Given an \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ and using the injection of the edge set $\mathcal{E}^- \cup \mathcal{E}^+$ into $\mathcal{S}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ as label map, one obtains a particular case of an $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentation, that we denote by $X(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$, and that we call the \mathcal{R} -graph shift of $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$. In the case of the graph inverse semigroups $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}))$ of strongly connected finite directed graphs $G(\mathfrak{P}, \mathcal{E})$ the subshifts $X(G(\mathfrak{P}, \mathcal{E}))$ are the Markov-Dyck shifts [M].The Dyck shifts D_N can be obtained in this way from the one-vertex directed graph with N > 1 loops. Also the Markov-Motzkin shifts [KM1] of strongly connected finite directed graphs $G(\mathfrak{P}, \mathcal{E})$ can be written as $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}))$ -presentations.

Béal, Blockelet and Dima [BBD1, BBD2] have introduced the notions of a Dyck automaton and of a sofic Dyck shift. Strengthening the Condition (G 1) to

(1.1)
$$\lambda(\sigma) \in \mathcal{E}^- \cup \{\mathbf{1}_{\mathfrak{p}} : \mathfrak{p} \in \mathfrak{P}\} \cup \mathcal{E}^+,$$

one obtains directed labelled graphs $G(\mathcal{V}, \Sigma, \lambda)$, that are Dyck-automata, with $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentations $X(\mathcal{V}, \Sigma, \lambda)$, that are sofic Dyck shifts. The alphabet of a sofic Dyck shift is partitioned into a set of call symbols, a set of internal symbols and a set of return symbols. The corresponding partition of the alphabet of an \mathcal{S} -presentation, that satisfies (1.1), is the partition of its alphabet into the sets $\mathcal{E}^-, \{\mathbf{1}_{\mathfrak{p}}, \mathfrak{p} \in \mathfrak{P}\}, \mathcal{E}^+$. The set of matched edges, that appears in the construction of a Dyck automaton, is provided by the relation \mathcal{R} . The \mathcal{R} -graph shifts are finite type Dyck shifts in the sense of [BBD3].

Given finite sets \mathcal{E}^- and \mathcal{E}^+ and a relation $\mathcal{R} \subset \mathcal{E}^- \times \mathcal{E}^+$, we set

$$\mathcal{E}^-(\mathcal{R}) = \{e^- \in \mathcal{E}^- : \{e^-\} \times \mathcal{E}^+ \subset \mathcal{R}\}, \quad \mathcal{E}^+(\mathcal{R}) = \{e^+ \in \mathcal{E}^+ : \mathcal{E}^- \times \{e^+\} \subset \mathcal{R}\}.$$

For a partitioned directed graph $G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ denote by $\mathfrak{P}^{(1)}$ the set of vertices in \mathfrak{P} that have a single predecessor vertex in \mathcal{E}^- , or, equivalently, that have a single successor vertex in \mathcal{E}^+ . For $\mathfrak{p} \in \mathfrak{P}^{(1)}$ the predecessor vertex of \mathfrak{p} in \mathcal{E}^- , which is identical to the successor vertex of \mathfrak{p} in \mathcal{E}^+ , will be denoted by $\eta(\mathfrak{p})$. For an \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ we set

$$\mathcal{E}_{\mathcal{R}}^{-} = \bigcup_{\mathfrak{p} \in \mathfrak{P}^{(1)}} \mathcal{E}^{-}(\mathcal{R}(\eta(\mathfrak{p}), \mathfrak{p})), \quad \mathcal{E}_{\mathcal{R}}^{+} = \bigcup_{\mathfrak{p} \in \mathfrak{P}^{(1)}} \mathcal{E}^{+}(\mathcal{R}(\eta(\mathfrak{p}), \mathfrak{p})),$$

and

$$\mathfrak{P}^{(1)}_{\mathcal{R}} = \{\mathfrak{p} \in \mathfrak{P}^{(1)} : \mathcal{R}(\eta(\mathfrak{p}), \mathfrak{p}) = \mathcal{E}^{-}(\eta(\mathfrak{p}), \mathfrak{p}) imes \mathcal{E}^{+}(\eta(\mathfrak{p}), \mathfrak{p})\}$$

An $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentation $X(G(\mathcal{V}, \Sigma, \lambda))$ is a Markov shift if and only if $\mathfrak{P}^{(1)} = \mathfrak{P}^{(1)}_{\mathcal{R}}$. We formulate two conditions (I) and (II) on \mathcal{R} -graphs $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$, such that

$$\mathfrak{P}^{(1)} \setminus \mathfrak{P}^{(1)}_{\mathcal{R}} \neq \emptyset.$$

Condition (II) comes in two parts (II–) and (II+) that are symmetric to one another:

- (I) For $\mathfrak{p} \in \mathfrak{P}^{(1)} \setminus \mathfrak{P}^{(1)}_{\mathcal{R}}, \mathcal{E}^{-}(\mathcal{R}(\eta(\mathfrak{p}),\mathfrak{p})) = \emptyset$, or $\mathcal{E}^{+}(\mathcal{R}(\eta(\mathfrak{p}),\mathfrak{p})) = \emptyset$.
- (II–) There is no cycle in the directed graph $G(\mathfrak{P}, \mathcal{E}_{\mathcal{P}}^{-})$.
- (II+) There is no cycle in the directed graph $G(\mathfrak{P}, \mathcal{E}_{\mathcal{P}}^+)$.

We show in Section 2 that an $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}))$ -presentation has Property (A) if and only if the \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+})$ satisfies conditions (I) and (II). In particular the \mathcal{R} -graph shifts $X(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}))$ have Property (A) if and only if the \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-}, \mathcal{E}^{+})$ satisfies Conditions (I) and (II). This implies that Markov-Dyck shifts of strongly connected finite directed graphs have Property (A). Also the Markov-Motzkin shifts of strongly connected finite directed graphs have Property (A). Concerning the invariance under flow-equivalence of Property (A), and of the associated semigroup, in particular for \mathcal{R} -graph shifts, see [Kr3].

In Section 3 we describe how one can obtain from an \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$, such that $\mathfrak{P}^{(1)} \setminus \mathfrak{P}^{(1)}_{\mathcal{R}} \neq \emptyset$, that satisfies conditions (I) and (II), an \mathcal{R} -graph $G_{\widehat{\mathcal{R}}}(\widehat{\mathfrak{P}}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+)$, such that the \mathcal{R} -graph semigroup $\mathcal{S}(G_{\widehat{\mathcal{R}}}(\widehat{\mathfrak{P}}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+))$ is associated to all $\mathcal{S}(G_{\mathcal{R}}(\mathcal{V}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentation. To obtain the \mathcal{R} -graph $G_{\widehat{\mathcal{R}}}(\mathfrak{P}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+)$ we apply a procedure, that extends a procedure for Markov-Dyck shifts, that was described in [HK2] and [KM2].

In Section 5 we consider examples. We show, that the isomorphism class of a one-vertex \mathcal{R} -graph can be recovered from the topological conjugacy class of its \mathcal{R} -graph shift. For certain \mathcal{R} -graph semigroups of a one-vertex graph see [HK1, Section 4]. We also consider a class of examples of \mathcal{R} -graph shifts, to which the \mathcal{R} -graph semigroups of one-vertex \mathcal{R} -graphs are associated, and that have the graph of a renewal system as an underlying graph $G(\mathfrak{P}, \{(\mathfrak{q}, \mathfrak{r}) \in \mathfrak{P} \times \mathfrak{P} : \mathcal{E}^-(\mathfrak{q}, \mathfrak{r}) \neq \emptyset)\})$. For \mathcal{R} -graphs in this class we show, that the isomorphism class of the \mathcal{R} -graph can be recovered from the topological conjugacy class of its \mathcal{R} -graph shift. This result covers certain Markov-Dyck shifts. For other results on the reconstruction of a directed graph from its Markov-Dyck shift see [KM1, Section 3] and [HK2].

2. $\mathcal{S}_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ -presentations

We denote the length of a directed path in a directed graph by ℓ . Given an an \mathcal{R} -graph

$$G_{\mathcal{R}} = G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^-),$$

we denote by $\mathcal{S}^{-}(G_{\mathcal{R}})(\mathcal{S}^{+}(G_{\mathcal{R}}))$ the set of non-zero elements of the subsemigroup of $\mathcal{S}^{-}(G_{\mathcal{R}})(\mathcal{S}^{+}(G_{\mathcal{R}}))$, that is generated by $\mathcal{E}^{-}(\mathcal{E}^{+})$. There is the one-to-one correspondence between the paths in the directed graph $G(\mathfrak{P}, \mathcal{E}^{-})$ ($G(\mathfrak{P}, \mathcal{E}^{+})$) and the elements of $\mathcal{S}^{-}(G_{\mathcal{R}})$ ($\mathcal{S}^{+}(G_{\mathcal{R}})$). We will use the same symbol to denote a path in $G(\mathfrak{P}, \mathcal{E}^{-})$ ($G(\mathfrak{P}, \mathcal{E}^{+})$) and the corresponding element of $\mathcal{S}^{-}(G_{\mathcal{R}})$ ($\mathcal{S}^{+}(G_{\mathcal{R}})$) (as we have already done for the edges in $\mathcal{E}^{-} \cup \mathcal{E}^{+}$). It will be clear from the context, which one is meant. For the elements of $\mathcal{S}^{-}(G_{\mathcal{R}})$ ($\mathcal{S}^{+}(G_{\mathcal{R}})$) the notations ℓ, s, t are also used. An element g of $\mathcal{S}(G_{\mathcal{R}})$ determines uniquely

$$\mathfrak{q}(g) \in \mathfrak{P}, \quad u^+(g) \in \{\mathbf{1}_{\mathfrak{q}(g)}\} \cup \mathcal{S}^+(G_{\mathcal{R}}), \ u^-(g) \in \{\mathbf{1}_{\mathfrak{q}(g)}\} \cup \mathcal{S}^-(G_{\mathcal{R}}),$$

such that its normal form is given by

$$g = u^+(g)\mathbf{1}_{\mathfrak{q}(g)}u^-(g).$$

We write the normal forms of elements g^- of $\mathcal{S}^-(G_{\mathcal{R}})$, and of elements g^+ of $\mathcal{S}^+(G_{\mathcal{R}})$ as

$$g^- = \prod_{1 \leq i(-) \leq \ell(g^-)} e^-_{i(-)}[g(-)], \quad g^+ = \prod_{1 \leq i(+) \leq \ell(g^+)} e^-_{i(+)}[g(+)].$$

We denote the set of non-zero elements of the subsemigroup of $\mathcal{S}(G_{\mathcal{R}})$, that is generated by $\mathcal{E}_{\mathcal{R}}^{-}(\mathcal{E}_{\mathcal{R}}^{-})$ by $\mathcal{S}_{\mathcal{R}}^{-}(G_{\mathcal{R}})(\mathcal{S}_{\mathcal{R}}^{+}(G_{\mathcal{R}}))$.

2.1. Context in $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}))$. In this subsection we consider an \mathcal{R} -graph $G_{\mathcal{R}} = G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}),$

such that

(2.1)
$$\mathfrak{P}^{(1)} \setminus \mathfrak{P}^{(1)}_{\mathcal{R}} \neq \emptyset$$

that satisfies conditions (I) and (II). For $f \in \mathcal{S}(G_{\mathcal{R}})$ we set

$$\Gamma^{-}(f) = \{ g \in \mathcal{S} : gf \neq 0 \}, \quad \Gamma^{+}(f) = \{ g \in \mathcal{S} : fg \neq 0 \},$$

$$\Gamma(f) = \{ (g(-), g(+)) \in \mathcal{S}^{2} : g(-)fg(+) \neq 0 \},$$

and we refer to $\Gamma(f)$ as the context of f.

We denote for $\mathfrak{q}, \mathfrak{r} \in \mathfrak{P}$ by $\ell_{-}(\mathfrak{q}, \mathfrak{r})$ ($\ell_{+}(\mathfrak{q}, \mathfrak{r})$) the length of a path in $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^{-})$ ($G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^{+})$) from \mathfrak{q} to \mathfrak{r} , provided such a path exists. By Condition (II) this notation is meaningful.

We denote for $\mathbf{q} \in \mathfrak{P}_{\mathcal{R}}^{(1)}$ by $D_{+}(\mathbf{q})$ the maximal length of a path in $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^{+})$ that leaves \mathbf{q} , and by $D_{-}(\mathbf{q})$ the maximal length of a path in $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^{-})$ that enters \mathbf{q} . We also set

$$D_{\circ}(\mathfrak{q}) = \begin{cases} \min\{D_{+}(\mathfrak{q}), D_{-}(\mathfrak{q})\}, & \text{if } \mathfrak{q} \in \mathfrak{P}_{\mathcal{R}}^{(1)}, \\ 0, & \text{if } \mathfrak{q} \in \mathfrak{P}^{(1)} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)} \end{cases}$$

We set inductively

$$\eta^k(\mathfrak{q}) = \eta(\eta^{k-1}(\mathfrak{q})), \qquad 1 < k < \max\{D_-(\mathfrak{q}), D_+(\mathfrak{q})\}, \mathfrak{q} \in \mathfrak{P}^{(1)}.$$

We remark, that a path b in $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^+)$, that starts at \mathfrak{q} , and that has length less then or equal to $D_{\circ}(\mathfrak{q})$, transverses the vertices $\eta^k, 1 \leq k < \ell(b)$, before entering its target vertex. A similar remark applies to paths in $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^+)$, that enter \mathfrak{q} .

We set $\eta^0(\mathfrak{q}) = \mathfrak{q}$, and for $\mathfrak{q} \in \mathfrak{P}^{(1)}$ we set

$$\mathfrak{R}_{-}(\mathfrak{q}) = \{\eta^{k}(\mathfrak{q}) : 0 \le k < D_{-}(\mathfrak{q})\}, \quad \mathfrak{R}_{+}(\mathfrak{q}) = \{\eta^{k}(\mathfrak{q}) : 0 \le k < D_{+}(\mathfrak{q})\},$$

and for $q(-), q(+) \in \mathfrak{P}^{(1)}$, such that

$$\mathfrak{q}(-) \neq \mathfrak{q}(+), \quad \mathfrak{R}_+(\mathfrak{q}(-)) \cap \mathfrak{R}_-(\mathfrak{q}(+)) \neq \emptyset,$$

we denote by $H_+(\mathfrak{q}(-),\mathfrak{q}(+))(H_-(\mathfrak{q}(-),\mathfrak{q}(+)))$ the minimal length of a path in $G(\mathfrak{P}, \mathcal{E}^+_{\mathcal{R}})$ $(G(\mathfrak{P}, \mathcal{E}^-_{\mathcal{R}}))$, that has $\mathfrak{q}(-)(\mathfrak{q}(+))$ as source vertex and a vertex in $\mathfrak{R}_+(\mathfrak{q}(+)))$ $(\mathfrak{R}_+(\mathfrak{q}(-)))$ as target vertex.

Lemma 2.1. For $\mathfrak{q}(-), \mathfrak{q}(+) \in \mathfrak{P}^{(1)}$, such that

$$\mathfrak{q}(-) \neq \mathfrak{q}(+), \quad \mathfrak{R}_+(\mathfrak{q}(-)) \cap \mathfrak{R}_-(\mathfrak{q}(+)) \neq \emptyset,$$

one has that

(2.2)
$$\eta^{H_{+}(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(-)) = \eta^{H_{-}(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(+)).$$

Proof. The inequality

 $\eta^{H_+(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(-)) \neq \eta^{H_-(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(+)),$

would imply the existence of a path in $G(\mathfrak{P}, \mathcal{E}^+)$, as well as in $G(\mathfrak{P}, \mathcal{E}^+)$, from $\eta^{H_+(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(-))$ to $\eta^{H_-(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(+))$, and also the existence of a path in $G(\mathfrak{P}, \mathcal{E}^+)$, as well as in $G(\mathfrak{P}, \mathcal{E}^+)$, from $\eta^{H_-(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(+))$ to $\eta^{H_+(\mathfrak{q}(-),\mathfrak{q}(+))}(\mathfrak{q}(-))$, contradicting the assumption (2.1).

We denote the vertex, that appears in (2.2) by $\mathfrak{p}(\mathfrak{q}(-), \mathfrak{q}(+))$.

Lemma A. Let $\mathfrak{q} \in \mathfrak{P}$. For

(2.3) $f^+ \in \{\mathbf{1}_{\mathfrak{q}}\} \cup \mathcal{S}^+(G_{\mathcal{R}}), \quad f^- \in \{\mathbf{1}_{\mathfrak{q}}\} \cup \mathcal{S}^-(G_{\mathcal{R}}),$ such that

~ ~ ~

(2.4)
$$s(f^+) = \mathbf{1}_{\mathfrak{q}} = t(f^-), \quad t(f^+) = s(f^-),$$

all elements of $\mathcal{S}(G_{\mathcal{R}})$ of the form f^+f^- have the same context.

Proof. One notes, that

$$0 \le \ell(f^+) = \ell(f^-) \le D_{\circ}(\mathfrak{q}).$$

Set

$$\Gamma_{\circ}^{-}(\mathfrak{q}) = \{g(-) \in \Gamma^{-}(\mathfrak{q}) : \ell(h^{-}(g(-))) \leq D_{\circ}(\mathfrak{q})\},$$

$$\Gamma_{\circ}^{+}(\mathfrak{q}) = \{g(+) \in \Gamma^{+}(\mathfrak{q}) : \ell(h^{+}(g(-))) \leq D_{\circ}(\mathfrak{q})\},$$

$$\Gamma_{\circ}(\mathfrak{q}) = \{ (g(-), g(+)) \in (\Gamma^{-}(\mathfrak{q}) \setminus \Gamma_{\circ}^{-}(\mathfrak{q})) \times (\Gamma^{+}(\mathfrak{q}) \setminus \Gamma_{\circ}^{+}(\mathfrak{q})) : \\
(\prod_{1 \leq i(+) < \ell(h^{-}(g(-))) - D_{\circ}(\mathfrak{q}))} e_{i(+)}^{+}[g(-)]) \mathbf{1}_{\eta^{D_{\circ}(\mathfrak{q})}(\mathfrak{q})} \\
(\prod_{\ell(h^{+}(g(+)) - D_{\circ}(\mathfrak{q}) < i(+) \leq \ell(h^{+}(g(+)))} e_{i(+)}^{+}[g(-)]) \neq 0.$$

By Condition (II) one has for f^+, f^- as in (2.3) and (2.4) $\Gamma(f^+f^-) = (\Gamma_{\circ}^-(\mathfrak{q}) \times \Gamma^+(\mathfrak{q})) \cup (\Gamma^-(\mathfrak{q}) \times \Gamma_{\circ}^+(\mathfrak{q})) \cup \Gamma_{\circ}(\mathfrak{q}).$

Lemma B. Let $\mathfrak{q}(-), \mathfrak{q}(+) \in \mathfrak{P}^{(1)}$, such that

$$\mathfrak{q}(-) \neq \mathfrak{q}(+).$$

(B-) Let there exist a path in $S^+(G_{\mathcal{R}})$ from $\mathfrak{q}(-)$ to $\mathfrak{q}(+)$. Then for (2.5) $h^+ \in \mathcal{S}(G_{\mathcal{R}}), \quad f^+ \in \{\mathbf{1}_{\mathfrak{q}(+)}\} \cup \mathcal{S}^+(G_{\mathcal{R}}), \quad f^- \in \{\mathbf{1}_{\mathfrak{q}(+)}\} \cup \mathcal{S}^-(G_{\mathcal{R}}),$ such that

(2.6) $s(h^+) = \mathbf{1}_{\mathfrak{g}(-)}, \quad t(h^+) = \mathbf{1}_{\mathfrak{g}(+)},$

$$s(f^+) = \mathbf{1}_{q(-)}, \quad t(f^+) = s(f^-), \quad t(f^-) = \mathbf{1}_{q(+)},$$

all elements of $S(G_{\mathcal{R}})$ of the form $h^+f^+f^-$ have the same context. (B+) Let there exist a path in from $\mathfrak{q}(+)$ to $\mathfrak{q}(-)$. Then for

$$h^+ \in \mathcal{S}(G_{\mathcal{R}}), \quad f^+ \in \{\mathbf{1}_{\mathfrak{q}(+)}\} \cup \mathcal{S}^+(G), \quad f^- \in \{\mathbf{1}_{\mathfrak{q}(+)}\} \cup \mathcal{S}^-(G_{\mathcal{R}}),$$

such that

$$\begin{split} s(h^+) &= \mathbf{1}_{\mathfrak{q}(-)}, \quad t(h^+) = \mathbf{1}_{\mathfrak{q}(+)}, \\ s(f^+) &= \mathbf{1}_{\mathfrak{q}(-)}, \quad t(f^+) = s(f^-), \quad t(f^-) = \mathbf{1}_{\mathfrak{q}(+)}, \end{split}$$

all elements of $\mathcal{S}(G_{\mathcal{R}})$ of the form $f^+f^-h^-$ have the same context.

Proof. We prove (B-). We note that

$$0 \le \ell(f^+) = \ell(f^-) \le D_{\circ}(\mathfrak{q}(+)).$$

 Set

$$\begin{split} &\Gamma_{\circ}^{-}(\mathfrak{q}(-),\mathfrak{q}(+)) = \{g(-) \in \Gamma^{-}(\mathbf{1}_{\mathfrak{q}(-)}) : \ell(h^{+}(g(-))) \leq \ell(\mathfrak{q}(-),\mathfrak{q}(+)) + D_{\circ}(\mathfrak{q}(+))\}, \\ &\Gamma_{\circ}^{+}(\mathfrak{q}(+)) = \{g(+) \in \Gamma^{+}(\mathbf{1}_{\mathfrak{q}(+)}) : \ell(h^{-}(g(+))) \leq D_{\circ}(\mathfrak{q}(+))\}, \end{split}$$

$$\begin{split} \Gamma_{\circ}(\mathfrak{q}(-),\mathfrak{q}(+)) &= \\ (g(-),g(+)) \in (\Gamma^{-}(\mathfrak{q}(-)) \setminus \Gamma_{\circ}^{-}(\mathfrak{q}(-),\mathfrak{q}(+)) \times (\Gamma^{+}(\mathfrak{q}(+)) \setminus \Gamma_{\circ}^{+}(\mathfrak{q}(+))) : \end{split}$$

$$(\prod_{1 \le i(-) < \ell(h^-(g(-))) - \ell_+(\mathfrak{q}(-),\mathfrak{q}(+)) - D_{\diamond}(\mathfrak{q}(+))} e^-_{i(-)}[g(-)]) \mathbf{1}_{\eta^{D_{\diamond}(\mathfrak{q}(+))}(\mathfrak{q}(+))} (\prod_{D_{\diamond}(\mathfrak{q}(+)) < i(+) \le \ell(h^+(g(+))} e^+_{i(+)}[g(+)]) \ne 0.$$

By Condition (II) one has for h^+, f^+, f^- , as in (2.5) and (2.6), that $\Gamma(h^+f^+f^-) =$

$$(\Gamma_{\circ}^{-}(\mathfrak{q}(-),\mathfrak{q}(+)) \times \Gamma^{+}(\mathfrak{q}(+)) \cup (\Gamma^{-}(\mathfrak{q}(-)) \times \Gamma_{\circ}^{+}(\mathfrak{q}(+))) \cup \Gamma_{\circ}(\mathfrak{q}(-),q(+)).$$

The proof of (B+) is symmetric.

Lemma C. Let $\mathfrak{g}(-), \mathfrak{g}(+) \in \mathfrak{P}^{(1)}$, such that

na C. Let
$$\mathfrak{q}(-), \mathfrak{q}(+) \in \mathfrak{P}^{(-)}$$
, such that $\mathfrak{q}(-) \neq \mathfrak{q}(+),$

and

$$\mathfrak{R}(\mathfrak{q}(-)) \cap \mathfrak{R}(\mathfrak{q}(+)) \neq \emptyset, \quad \mathfrak{p}(\mathfrak{q}(-), \mathfrak{q}(+)) \notin \{\mathfrak{q}(-), \mathfrak{q}(+)\}.$$
Then for
$$(2.7) \qquad h^+, f^+ \in \mathcal{S}^+(G_{\mathcal{R}}), \quad f^-, h^- \in \mathcal{S}^-(G_{\mathcal{R}}),$$

such that

(2.8)

$$s(h^+) = \mathbf{1}_{\mathfrak{q}(-)}, \ t(h^+) = \mathbf{1}_{\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+))},$$

$$s(f^+) = \mathbf{1}_{\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+))}, \ t(f^+) = s(f^-), \ t(f^-) = \mathbf{1}_{\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+))},$$

 $s(h^-) = \mathbf{1}_{\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+))}, \ t(h^-) = \mathbf{1}_{\mathfrak{q}(+)},$

all elements of $S(G_R)$ of the form $h^+f^+f^-h^-$ have the same context. Proof. One notes that

$$0 \le \ell(f^+) = \ell(f^-) \le D_{\circ}(\mathfrak{p}(\mathfrak{q}(-), \mathfrak{q}(+))).$$

 Set

$$\begin{split} \Gamma_{\circ}^{-}(\mathfrak{q}(-),\mathfrak{q}(+)) &= \\ \{g(-) \in \Gamma^{-}(\mathbf{1}_{\mathfrak{q}(-)}) : \ell(h^{-}(g(-))) \leq H_{-}(\mathfrak{q}(-),\mathfrak{q}(+)) + D_{\circ}(\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+)))\}, \end{split}$$

$$\begin{split} \Gamma_{\circ}^{-}(\mathfrak{q}(-),\mathfrak{q}(+)) &= \\ \{g(+) \in \Gamma^{+}(\mathbf{1}_{\mathfrak{q}(+)}) : \ell(h^{+}(g(+))) \leq H_{+}(\mathfrak{q}(-),\mathfrak{q}(+)) + D_{\circ}(\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+)))\}, \\ \Gamma_{\circ}(\mathfrak{q}(-),\mathfrak{q}(+)) &= \\ \{(g(-),g(+)) \in (\Gamma^{-}(\mathbf{1}_{\mathfrak{q}(-)}) \setminus \Gamma_{\circ}^{-}(\mathfrak{q}(-),\mathfrak{q}(+)) \times (\Gamma^{+}(\mathbf{1}_{\mathfrak{q}(+)}) \setminus \Gamma_{\circ}^{+}(\mathfrak{q}(+),\mathfrak{q}(+))) : \end{split}$$

$$(\prod_{1 \le i(-) < \ell(h^{-}(g(-))) - \ell_{+}(\mathfrak{q}(-),\mathfrak{q}(+)) - D_{\circ}(\mathfrak{q}(+))} e_{i(-)}^{-}[g(-)]) \mathbf{1}_{\eta^{D_{\circ}(\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+)))}(\mathfrak{p}(\mathfrak{q}(-),\mathfrak{q}(+)))} (\prod_{D_{\circ}(\mathfrak{q}(+)) < i(+) \le \ell(h^{+}(g(+)))} e_{i(+)}^{+}[g(+)]) \ne 0.$$

By Condition (II) one has for that for h^+ , f^- , h^- as in (2.7), (2.8), that $\Gamma(h^+f^+f^-h^-) =$ $(\Gamma_{\circ}^-(\mathfrak{q}(-),\mathfrak{q}(+)) \times \Gamma^+(\mathbf{1}_{\mathfrak{q}(+)}) \cup (\Gamma^-(\mathbf{1}_{\mathfrak{q}(-)}) \times \Gamma_{\circ}^+(\mathfrak{q}(-),\mathfrak{q}(+))) \cup \Gamma_{\circ}(\mathfrak{q}(-),q(+)).$

2.2. Property (A) and \mathcal{R} -graph shifts. We introduce notation and terminology for subshifts. The set of periodic points of a subshift X we denote by P(X). The smallest period of $p \in P(X)$, we denote by $\pi(p)$. We denote the language of admissible words of a subshift $X \subset \Sigma^{\mathbb{Z}}$ by $\mathcal{L}(X)$. The context of a word $b \in \mathcal{L}(X)$ is defined as the set

$$\Gamma(b) = \{ c(-), c(+) \} \in \mathcal{L}(X)^2 : C(-)bc(+) \in \mathcal{L}(X) \}.$$

Concerning $\mathcal{S}(G(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentations $X(\mathcal{V}, \Sigma, \lambda))$ we remark, that

Given a subshift $X \subset \Sigma^{\mathbb{Z}}$ we set

$$x_{[i,k]} = (x_j)_{i \le j \le k}, \quad x \in X, i, k \in \mathbb{Z}, i \le k,$$

and

$$X_{[i,k]} = \{x_{[i,k]} : x \in X\}, i, k \in \mathbb{Z}, i \le k.$$

We set

$$\Gamma(a) = \bigcup_{n,m \in \mathbb{N}} \{ (b,c) \in X_{[i-n,i]} \times X_{[k,k+m]} : (b,a,c) \in X_{[i-n,k+m]} \},\$$
$$a \in X_{[i,k]}, i, k \in \mathbb{Z}, i \le k.$$

and call $\Gamma(a)$ the context of the block a. We set

$$\Gamma_n^+(a) = \{ b \in X_{(k,k+n]} : (a,b) \in X_{[i,k+n]} \},\$$

$$n \in \mathbb{N}, a \in X_{[i,k]}, i, k \in \mathbb{Z}, i \le k.$$

 Γ^{-} has the symmetric meaning. We set

$$\omega^+(a) = \bigcup_{n \in \mathbb{N}} \bigcap_{n \in \mathbb{N}} \bigcap_{c \in \Gamma_n^-(a)} \{ b \in X_{(k,k+n]} : (c,a,b) \in X_{[i-n,k+m]} \},$$
$$a \in X_{[i,k]}, i, k \in \mathbb{Z}, i \le k.$$

 ω^{-} has the symmetric meaning.

Given a subshift $X \subset \Sigma^{\mathbb{Z}}$ we define a subshift of finite type (more precicely, an n-step Markov shift) $A_n(X)$ by

$$A_n(X) = \bigcap_{i \in \mathbb{Z}} (\{x \in X : x_i \in \omega^+(x_{[i-n,i)})\} \cap (\{x \in X : x_i \in \omega^-(x_{(i,i+n]})\}) \quad n \in \mathbb{N},$$

and we set

$$A(X) = \bigcup_{n \in \mathbb{N}} A_n(X).$$

We recall from [Kr2] the definition of Property (A). For $n \in \mathbb{N}$ a subshift $X \subset \Sigma^{\mathbb{Z}}$, has property $(a, n, H), H \in \mathbb{N}$, if for $h, \tilde{h} \geq 3H$ and for $I_{-}, I_{+}, \tilde{I}_{-}, \tilde{I}_{+} \in \mathbb{Z}$, such that

$$I_{+} - I_{-}, I_{+} - I_{-} \ge 3H,$$

and for

$$a \in A_n(X)_{(I_-,I_+]}, \quad \widetilde{a} \in A_n(X)_{(\widetilde{I}_-,\widetilde{I}_+]},$$

such that

$$a_{(I_{-},I_{-}+H]} = \tilde{a}_{(\tilde{I}_{-},\tilde{I}_{+}H]}, \quad a_{(I_{+}-H,I_{+}]} = \tilde{a}_{(\tilde{I}_{+}-H,\tilde{I}_{+}]},$$

one has that a and \tilde{a} have the same context. A subshift $X \subset \Sigma^{\mathbb{Z}}$ has property (A) if there are $H_n, n \in \mathbb{N}$, such that X has the properties $(a, n, H_n), n \in \mathbb{N}$.

Theorem 2.2. Let

$$G_{\mathcal{R}} = G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+),$$

be an \mathcal{R} -graph such that

(2.9)
$$\mathfrak{P}^{(1)} \setminus \mathfrak{P}^{(1)}_{\mathcal{R}} \neq \emptyset.$$

For an $\mathcal{S}_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ -presentation $X(\mathcal{V}, \Sigma, \lambda)$ to have Property (A) it is necessary and sufficient, that $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ satisfies Conditions (I) and (II).

Proof. We prove necessity. We choose vertices $V_{\mathfrak{p}} \in \mathcal{V}(\mathfrak{p}), \mathfrak{p} \in \mathfrak{P}$, and cycles $c_{\mathfrak{p}}, \mathfrak{p} \in \mathfrak{P}$ in the graph $G(\mathcal{V}, \Sigma)$, such that

$$\mathfrak{s}(c_{\mathfrak{p}}) = t(c_{\mathfrak{p}}) = V_{\mathfrak{p}}, \quad \lambda(c_{\mathfrak{p}}) = \mathbf{1}_{\mathfrak{p}}, \qquad \mathfrak{p} \in \mathfrak{P}.$$

For $k \in \mathbb{N}$ we denote by $c_{\mathfrak{p}}^k$ the cycle, that transverses k-times the cycle $c_{\mathfrak{p}}$. Also we choose for all $e^- \in \mathcal{E}^-$ a path a_{e^-} in the graph $G(\mathcal{V}, \Sigma)$, such that

$$s(a_{e^-}) = V_{s(e^-)}, t(a_{e^-}) = V_{t(e^-)}, \quad \lambda(a_{e^-}) = e^-.$$

and we make similar choices for all $e^+ \in \mathcal{E}^+$. We set

$$M = \max(\{\ell(a_{e^-}) : e^- \in \mathcal{E}^-\} \cup \{\ell(c_{\mathfrak{p}}) : \mathfrak{p} \in \mathfrak{P}\} \cup \{\ell(a_{e^+}) : e^+ \in \mathcal{E}^+\}).$$

One has that

$$c_{\mathfrak{p}}^{k} \in \mathcal{L}(A_{M}(X(\mathcal{V}, \Sigma, \lambda))), \qquad k \in \mathbb{N}, \mathfrak{p} \in \mathfrak{P}.$$

(I). Assume that the \mathcal{R} -graph $\mathcal{G}_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ does not satisfy Condition (I). Under this assumption we can choose a vertex $\mathfrak{p} \in \mathfrak{P}^1 \setminus \mathfrak{P}^1_{\mathcal{R}}$, and edges

(2.10)
$$e^- \in \mathcal{E}^-_{\mathcal{R}}(\eta(\mathfrak{p}), \mathfrak{p}), \qquad e^+ \in \mathcal{E}^+_{\mathcal{R}}(\eta(\mathfrak{p}), \mathfrak{p}).$$

such that

(2.11) $(e^-, e^+) \notin \mathcal{R}(\eta(\mathfrak{p}), \mathfrak{p}).$

By construction

$$(c_{\mathfrak{p}}^k, a_{e^+}, a_{e^-}, c_{\mathfrak{p}}^k) \in \mathcal{L}(A_M(X(\mathcal{V}, \Sigma, \lambda))), \quad k \in \mathbb{N}.$$

It follows from (2.10), that

$$(a_{e^{-}}, a_{e^{+}}) \in \Gamma(c_{\mathfrak{p}}^{k}, a_{e^{+}}, a_{e^{-}}, c_{\mathfrak{p}}^{k}), \quad k \in \mathbb{N}.$$

and it follows from (2.11), that

$$(a_{e^-}, a_{e^+}) \notin \Gamma(c_{\mathfrak{p}}^{2k}), \quad k \in \mathbb{N}$$

We have shown, that $X(G(\mathcal{V}, \Sigma, \lambda))$ does not have Property (A).

(II) Assume that the \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ does not satisfy Condition (II -). By (2.9) every cycle in $G(\mathfrak{P}, \mathcal{E}^-)$ transverses at least one vertex in $\mathfrak{P}^{(1)} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}$. We can therefore choose a path

$$f^{-} = (e_l^{-})_{1 \le l \le \ell(f^{-})},$$

in the graph $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^{-})$, and a vertex $\widetilde{\mathfrak{q}} \in \mathfrak{P}^{(1)} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}$, such that

 $s(f^-) = \widetilde{\mathfrak{q}} = t(f^-),$

together with edges

 $\widetilde{e}^- \in \mathcal{E}_{\mathcal{R}}^-, \qquad \widetilde{e}^+ \in \mathcal{E}_{\mathcal{R}}^+,$ (2.12)such that $t(\tilde{e}^-) = \tilde{\mathfrak{q}} = s(\tilde{e}^+),$ and such that $(\tilde{e}^-, \tilde{e}^+) \notin \mathcal{R}.$ (2.13)We set $a_{f^-} = (a_{e_l^-})_{1 \le l \le \ell(f^-)}.$ By construction

$$(c_{\mathfrak{p}}^k, a_{f^-}, c_{\mathfrak{p}}^k) \in \mathcal{L}(A_M(X(G))), \quad k \in \mathbb{N}$$

It follows from (2.12), that

$$(a_{e^{-}}, a_{e^{+}}) \in \Gamma(c_{\mathfrak{p}}^{k}, a_{e^{+}}, a_{e^{-}}, c_{\mathfrak{p}}^{k}), \quad k \in \mathbb{N}.$$

and it follows from (2.13), that

$$(a_{e^-}, a_{e^+}) \notin \Gamma(c_{\mathfrak{p}}^{2k}), \quad k \in \mathbb{N}$$

We have shown, that $X(G(\mathcal{V}, \Sigma, \lambda))$ does not have Property (A). The case (II +) is symmetric.

We prove sufficiency. Let $n \in \mathbb{N}$, and let $a(-), a(+) \in \mathcal{L}(X(\mathcal{V}, \Sigma, \lambda)), \ell(a(-)), \ell(a(+)) =$ n. For $m \in [1, n]$ denote by by $a(-)_{[1,m]}(a(+)_{[m,n]})$ the prefix (suffix) of length m (n - m + 1) of a(-)(a(+)). Set

$$\begin{split} M(-) &= \\ & \left\{ \begin{aligned} & \max\{m \in [1,n] : \lambda(a(-)_{[1,m]}) = u^+(\lambda(a(-))\} & \text{if } u^+(\lambda(a(-)) \in \mathcal{S}^+(G_{\mathcal{R}}), \\ & 0, & \text{if } u^+(\lambda(a(-)) = \mathbf{1}_{\mathfrak{q}(\lambda(a(-)))}, \end{aligned} \right. \end{split}$$

M(+) = $\begin{cases} \max\{m \in [1,n] : \lambda(a(+)_{[n-m,n]}) = u^{-}(\lambda(a(-)))\} & \text{if } u^{-}(\lambda(a(+)) \in \mathcal{S}^{-}(G_{\mathcal{R}}), \\ & \text{if } u^{-}(\lambda(a(+)) = 1, \alpha < \infty) \end{cases}$ if $u^-(\lambda(a(+)) = \mathbf{1}_{\mathfrak{q}(\lambda(a(+)))}$. 0,

Let

$$K(-), K(+), \bar{K}(-), \bar{K}(+) \in \mathbb{Z},$$

 $K(+) - K(-), \bar{K}(+) - \bar{K}(-) > 2n,$

and let

(2.14)
$$b \in A_n(X(G))_{[K(-),K(+)]}, \quad \bar{b} \in A_n(X(G))_{[\bar{K}(-),\bar{K}(+)]}$$

be such that

$$b_{[K(-),K(-)+n]} = \bar{b}_{[\bar{K}(-),\bar{K}(-)+n]} = a(-),$$

$$b_{[K(+)-n,K(+)]} = \bar{b}_{[\bar{K}(+)-n,\bar{K}(+)]} = a(+),$$

Set

be

$$c = b_{[K(-)+M(-),K(+)-M(+)]}, \quad \bar{c} = b_{[\bar{K}(-)+M(-),\bar{K}(+)-M(+)]}.$$

We consider the four cases (A), (B-), (B+) and (C):

(A) In the case, that $q(\lambda(a(-))) = q(a(-))$, let set

$$f^+ = h^+(\lambda(c)), f^- = h^-(\lambda(c)), \quad \bar{f}^+ = \bar{h}^+(\lambda(\bar{c})), \bar{f}^- = \bar{h}^-(\lambda(\bar{c})),$$

It follows from (2.14), that

$$f^+, \bar{f}^+ \in \mathcal{S}^+_{\mathcal{R}}(G_{\mathcal{R}}), \quad f^-, \bar{f}^- \in \mathcal{S}^-_{\mathcal{R}}(G_{\mathcal{R}}).$$

By Lemma A

(2.15)
$$\Gamma(f^+f^-) = \Gamma(\bar{f}^+\bar{f}^-).$$

(B -) In the case, that

$$\mathfrak{q}(\lambda(a(-)))\neq\mathfrak{q}(\lambda(a(+)),\quad \mathfrak{q}(\lambda(a(-)))=\mathfrak{p}(\mathfrak{q}(\lambda(a(-))),\mathfrak{q}(\lambda(a(+))),$$

one has

$$\ell(h^+(\lambda(c)), \ell(h^+(\lambda(c)) \ge \ell_+(\mathfrak{q}(\lambda(a(-)), \mathfrak{q}(\lambda(a(+)))).$$

Let

$$h^+, f^+, f^-, \bar{h}^+, \bar{f}^-,$$

be given by

$$\begin{split} \ell(h^+) &= \ell(\bar{h}^+) = \ell_+(\mathfrak{q}(\lambda(a(-)),\mathfrak{q}(\lambda(a(+)),\\ h^+ &= u^+(\lambda(c))f^+, \quad \bar{h}^+ = u^+(\lambda(\bar{c}))f^+. \end{split}$$

By (2.14)

$$h^+, \bar{h}^+, f^+, \bar{f}^+ \in \mathcal{S}^-_{\mathcal{R}}(G_{\mathcal{R}}), \quad f^-, \bar{f}^- \in \mathcal{S}^+_{\mathcal{R}}(G_{\mathcal{R}}).$$

By Lemma B,

(2.16)
$$\Gamma(h^+ f^+ f^-) = \Gamma(\bar{h}^+ \bar{f}^+ \bar{f}^-).$$

(C): In the case, that

$$\mathfrak{q}(\lambda(a(-))) \neq \mathfrak{q}(\lambda(a(+))),$$

$$\mathfrak{p}(\mathfrak{q}(\lambda(a(-))),\mathfrak{q}(\lambda(a(+)))\notin \{\mathfrak{q}(\lambda(a(-))),\mathfrak{q}(\lambda(a(+)))\},$$

Let

$$h^+\bar{h}^+, f^+, \bar{f}^+, \in \mathcal{S}^+_{\mathcal{R}}(G_{\mathcal{R}}), \quad f^-, \bar{f}^-, h^-\bar{h}^-, \in \mathcal{S}^-_{\mathcal{R}}(G_{\mathcal{R}}),$$

be given by

$$\begin{split} \ell(h^+) &= \ell(\bar{h}^+) = \ell_+(\mathfrak{q}(\lambda(a(-))), \mathfrak{p}(\mathfrak{q}(\lambda(a(-))), \mathfrak{q}(\lambda(a(+))))), \\ \\ \ell(h^-) &= \ell(\bar{h}^-) = \ell_-(\mathfrak{p}(\mathfrak{q}(\lambda(a(-)))), \mathfrak{q}(\lambda(a(+))), \mathfrak{q}(\lambda(a(+)))), \\ \\ u^+(c) &= h^+f^+, \ u^+(\bar{c}) = \bar{h}^+\bar{f}^+, \quad u^-(c) = h^-f^-, \ u^-(\bar{c}) = \bar{h}^-\bar{f}^-. \end{split}$$

By (2.14)

$$h^+, \bar{h}^+, f^+, \bar{f}^+ \in \mathcal{S}^-_{\mathcal{R}}(G_{\mathcal{R}}), \quad f^-, \bar{f}^-, h^+, \bar{h}^+ \in \mathcal{S}^+_{\mathcal{R}}(G_{\mathcal{R}}).$$

By Lemma C,

(2.17)
$$\Gamma(h^+f^+f^-h^-) = \Gamma(\bar{h}^+\bar{f}^+\bar{f}^-\bar{h}^-).$$

It follows from (2.15 - 17), that in all cases the context of b is equal to the context or \overline{b} . It is shown, that X has properties $a(n,n), n \in \mathbb{N}$.

3. The \mathcal{R} -graph semigroup associated to an $\mathcal{S}(G_{\mathcal{R}}(\mathcal{P}, \mathcal{E}^- \cup \mathcal{E}^+))$ -presentation

Following the terminology, that was introduced in [HI], we say for an \mathcal{R} -graph $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ and an $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^-, \mathcal{E}^+))$ -presentation $X(\mathcal{V}, \Sigma, \lambda)$, that a periodic point p in $X(\mathcal{V}, \Sigma, \lambda)$) is neutral, if there exist $I \in \mathbb{Z}$ and $\mathfrak{p} \in \mathfrak{P}$ such that $\lambda(p_{[I,I+\pi(p)]}) = \mathbf{1}_{\mathfrak{p}}$, and we say that a periodic point p in $X(\mathcal{V}, \Sigma, \lambda)$ has negative (positive) multiplier, if there exist $I \in \mathbb{Z}$, such that $\lambda(p_{[I,I+\pi(p)]}) \in \mathcal{S}^-(\mathfrak{P}, \mathcal{E}^+ \cup \mathcal{E}^-)(\mathcal{S}^+(\mathfrak{P}, \mathcal{E}^+ \cup \mathcal{E}^-))$.

Lemma 3.1. Let $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ be an \mathcal{R} -graph, such that $\mathfrak{P} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)} \neq \emptyset$, that satisfies Conditions (II), and let $X(\mathcal{V}, \Sigma, \lambda)$ be an $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^-, \mathcal{E}^+))$ -presentation. Then a periodic point of $X(\mathcal{V}, \Sigma, \lambda)$ is in $A(X(\mathcal{V}, \Sigma, \lambda))$ if and only if it is neutral.

Proof. Let

$$(3.1) p \in A(X(\mathcal{V}, \Sigma, \lambda)).$$

and let $I \in \mathbb{Z}$ be such that

$$\lambda(p_{[I,I+\pi(p)}) \in \mathcal{S}^{-}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}) \cup \{\mathbf{1}_{\mathfrak{p}} : \mathfrak{p} \in \mathfrak{P}\} \cup \mathcal{S}^{+}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}).$$

If

$$\mathcal{N}(p_{[I,I+\pi(p))}) \in \mathcal{S}^{-}_{\mathcal{R}}(\mathfrak{P},\mathcal{E}^{-}\cup\mathcal{E}^{+})$$

then it follows from (3.1) that $\lambda(p_{[I,I+\pi(p)})$ is given by a cycle in the directed graph $G(\mathfrak{P}, \mathcal{E}_{\mathcal{R}}^{-})$, contradicting Condition (II-). For the case that

$$\lambda(p_{[I,I+\pi(p))}) \in \mathcal{S}^+(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+),$$

one has the symmetric argument.

For the converse, note that $\lambda(p_{[I,I+\pi(p))}) = \mathbf{1}_{\mathfrak{p}}$, implies $p \in A_{\pi(p)}(X(\mathcal{V}, \Sigma, \lambda))$.

Given finite sets \mathcal{E}^- and \mathcal{E}^- and a relation $\mathcal{R} \subset \mathcal{E}^- \times \mathcal{E}^+$, we say that $e^- \in \mathcal{E}^$ and $\tilde{e}^- \in \mathcal{E}^-$ are $\sim (\mathcal{R}, -)$ -equivalent if

$$\{\tilde{e}^+ \in \mathcal{E}^+ : (e^-, \tilde{e}^+) \in \mathcal{R}\} = \{\tilde{e}^+ \in \mathcal{E}^+ : (\tilde{e}^-, \tilde{e}^+) \in \mathcal{R}\}$$

An equivalence relation $\sim (\mathcal{R}, +)$ on \mathcal{E}^+ is defined symmetrically. Given an \mathcal{R} -graph $\mathcal{G}_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ we we construct an \mathcal{R} -graph $\mathcal{G}_{\bar{\mathcal{R}}}(\mathfrak{P}, \bar{\mathcal{E}}^- \cup \bar{\mathcal{E}}^+)$, by setting

$$\begin{split} \bar{\mathcal{E}}^-(\mathfrak{q},\mathfrak{r}) &= [\mathcal{E}^-(\mathfrak{q},\mathfrak{r})]_{\sim(\mathcal{R},-)}, \qquad \bar{\mathcal{E}}^+(\mathfrak{q},\mathfrak{r}) = [\mathcal{E}^-(\mathfrak{q},\mathfrak{r})]_{\sim(\mathcal{R},+)}, \\ \bar{\mathcal{R}}(\mathfrak{q},\mathfrak{r}) &= \{(\bar{e}^-,\bar{e}^+) \in \bar{\mathcal{E}}^-(\mathfrak{q},\mathfrak{r}) \times \bar{\mathcal{E}}^+(\mathfrak{q},\mathfrak{r}) : \bar{e}^- \times \bar{e}^+ \subset \mathcal{R}(\mathfrak{q},\mathfrak{r})\}, \qquad \mathfrak{q},\mathfrak{r} \in \mathfrak{P}. \end{split}$$

We denote by $\bar{\mathcal{F}}_{\bar{\mathcal{R}}}^-$ the set of edges in $\bar{\mathcal{E}}^-$, that are the single incoming edges of their target vertices, and we denote by $\bar{\mathcal{F}}_{\bar{\mathcal{R}}}^+$ the set of edges in $\bar{\mathcal{E}}^+$, that are the eingle outgoing edges of their source vertices. Observe that the set $\mathfrak{P}_{\mathcal{R}}^{(1)}$ is the set of target vertices of the edges in $\bar{\mathcal{F}}_{\bar{\mathcal{R}}}^-$, which is equal to the set of source vertices of the edges in $\bar{\mathcal{F}}_{\bar{\mathcal{R}}}^+$, and that $\mathfrak{P} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}$ is the set of vertices, that have at least two incoming edges in $\bar{\mathcal{E}}^-$, or, equivalently, that have at least two outgoing edges in $\bar{\mathcal{E}}^+$. We set

$$\widehat{\mathfrak{R}} = \mathfrak{P} \setminus \mathfrak{P}_{\mathcal{P}}^{(1)},$$

and we denote by $\widehat{\mathfrak{R}}^{\bullet}$ the set of vertices in $\widehat{\mathfrak{R}}$, that are source vertices of an edge in $\overline{\mathcal{F}}_{\overline{\mathcal{R}}}^-$, or, equivalently, that are target vertices of an edge in $\overline{\mathcal{F}}_{\overline{\mathcal{R}}}^+$. For $\mathfrak{r} \in \mathfrak{R}_{\overline{\mathcal{R}}}^{\bullet}$ we denote by $\mathfrak{P}(\mathfrak{r})$ the set of $\mathfrak{p} \in \mathfrak{P}$, that are target vertices of a path in the graph $G(\mathfrak{P}, \overline{\mathcal{F}}_{\overline{\mathcal{R}}}^{-})$, that leaves \mathfrak{r} , or, equivalently, that are source vertices of a path in the graph $G(\mathfrak{P}, \overline{\mathcal{F}}_{\overline{\mathcal{R}}}^{+})$, that enters \mathfrak{r} . For $\mathfrak{r} \in \mathfrak{R} \setminus \mathfrak{R}^{\bullet}$ we set $\mathfrak{P}(\mathfrak{r}) = {\mathfrak{r}}$. We also set

$$\begin{split} \bar{\mathcal{F}}^{-}_{\bar{\mathcal{R}}}(\mathfrak{r}) &= \{f^{-} \in \bar{\mathcal{F}}^{-}_{\bar{\mathcal{R}}} : t(f^{-}) \in \mathfrak{P}(\mathfrak{r})\}, \\ \bar{\mathcal{F}}^{+}_{\bar{\mathcal{R}}}(\mathfrak{r}) &= \{f^{+} \in \bar{\mathcal{F}}^{+}_{\bar{\mathcal{R}}} : s(f^{+}) \in \mathfrak{P}(\mathfrak{r})\}, \qquad \mathfrak{r} \in \widehat{\mathfrak{R}}^{\bullet}, \end{split}$$

and have obtained partitions

$$\mathfrak{P} = \bigcup_{\mathfrak{r} \in \widehat{\mathfrak{R}}} \mathfrak{P}(\mathfrak{r}), \quad \bar{\mathcal{E}}^- = (\bar{\mathcal{E}}^- \setminus \bar{\mathcal{F}}_{\bar{\mathcal{R}}}^-) \cup \bigcup_{\mathfrak{r} \in \widehat{\mathfrak{R}}} \bar{\mathcal{F}}_{\bar{\mathcal{R}}}^-(\mathfrak{r}), \quad \bar{\mathcal{E}}^+ = (\bar{\mathcal{E}}^- \setminus \bar{\mathcal{F}}_{\bar{\mathcal{R}}}^+) \cup \bigcup_{\mathfrak{r} \in \widehat{\mathfrak{R}}} \bar{\mathcal{F}}_{\bar{\mathcal{R}}}^-(\mathfrak{r}).$$

For all $\mathbf{r} \in \mathfrak{R}^{\bullet}_{\overline{\mathcal{R}}}$ the directed graph $G({\mathbf{r}} \cup \mathfrak{P}(\mathbf{r}), \overline{\mathcal{F}}^{-}_{\overline{\mathcal{R}}}(\mathbf{r}))$ is an outward directed tree, and the directed graph $G({\mathbf{r}} \cup \mathfrak{P}(\mathbf{r}), \overline{\mathcal{F}}^{+}_{\overline{\mathcal{R}}}(\mathbf{r}))$ is an inward directed tree. The directed trees $G({\mathbf{r}} \cup \mathfrak{P}(\mathbf{r}), \overline{\mathcal{F}}^{-}_{\overline{\mathcal{R}}}(\mathbf{r}))$ and $G({\mathbf{r}} \cup \mathfrak{P}(\mathbf{r}), \overline{\mathcal{F}}^{+}_{\overline{\mathcal{R}}}(\mathbf{r}))$ are reversals of one another, and the set $\mathfrak{L}(\mathbf{r})$ of their leaves o is given by the set of vertices in $\mathfrak{P}(\mathbf{r})$, that are source vertices of an edge in $\overline{\mathcal{E}}^{-} \setminus \overline{\mathcal{F}}^{-}$, or, equivalently, that are target vertices of an edge in $\overline{\mathcal{E}}^{+} \setminus \overline{\mathcal{F}}^{+}$. To a vertex $\mathbf{r} \in \mathfrak{R}_{\overline{\mathcal{R}}} \setminus \mathfrak{R}^{\bullet}_{\overline{\mathcal{R}}}$ we associate the degenerate tree $G({\mathbf{r}}, \emptyset)$ with its leaf \mathbf{r} . For $\mathbf{r} \in \widehat{\mathfrak{R}}$, if the source (target) vertex of an edge $e^{-} \in \overline{\mathcal{E}}^{-} \setminus \overline{\mathcal{F}}^{-}_{\overline{\mathcal{R}}}$ ($e^{+} \in \overline{\mathcal{E}}^{+} \setminus \overline{\mathcal{F}}^{-}_{\overline{\mathcal{R}}}$) is in $\mathfrak{P}(\mathbf{r})$, then this source (target) vertex is necessarily in $\mathfrak{L}(\mathbf{r})$.

From the \mathcal{R} -graph $\mathcal{G}_{\mathcal{R}}(\mathfrak{P}, \mathcal{\bar{E}}^- \cup \mathcal{\bar{E}}^+)$ we construct an \mathcal{R} -graph $\mathcal{G}_{\mathcal{R}}(\mathfrak{P}, \mathcal{\hat{E}}^- \cup \mathcal{\hat{E}}^+)$. We postulate that there are bijections

$$e^- \to \widehat{e}^- \in \widehat{\mathcal{E}}^- \ (e^- \in \overline{\mathcal{E}}^- \setminus \overline{\mathcal{F}}^-), \quad e^+ \to \widehat{e}^+ \in \widehat{\mathcal{E}}^+ \ (e^+ \in \overline{\mathcal{E}}^- \setminus \overline{\mathcal{F}}^-).$$

We specify the source and target mappings of the graph $\mathcal{G}_{\widehat{\mathcal{R}}}(\mathfrak{P}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+)$ by setting

$$\begin{split} t(\widehat{e}^-) &= t(e^-), \qquad e^- \in \bar{\mathcal{E}}^-, \\ s(\widehat{e}^+) &= s(e^+), \qquad e^+ \in \bar{\mathcal{E}}^-, \end{split}$$

and by assigning for $\mathfrak{r} \in \widehat{\mathfrak{R}}$, to an edge $e^- \in \overline{\mathcal{E}}^-$, such that $s(\widehat{e}^-) \in \mathfrak{L}(\mathfrak{r})$, the vertex \mathfrak{r} as source vertex, and to an edge $e^+ \in \overline{\mathcal{E}}^+$, such that $t(\widehat{e}^-) \in \mathfrak{L}(\mathfrak{r})$, the vertex \mathfrak{r} as target vertex. We set

$$\widehat{\mathcal{R}} = \{ (\widehat{e}^{-}, \widehat{e}^{+}) \in \widehat{\mathcal{E}}^{-} \times \widehat{\mathcal{E}}^{+} : (e^{-}, e^{+}) \in \overline{\mathcal{R}} \cap \left((\overline{\mathcal{E}}^{-} \setminus \overline{\mathcal{F}}^{-}) \times \overline{\mathcal{E}}^{+} \setminus \overline{\mathcal{F}}^{+})) \right) \}.$$

In the \mathcal{R} -graph $\mathcal{G}_{\widehat{\mathcal{R}}}(\mathfrak{P}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+)$ every vertex has at least two incoming edges in $\widehat{\mathcal{E}}^-$, which means that every vertex in $\widehat{\mathcal{E}}^+$ has at least two outgoing edges in $\widehat{\mathcal{E}}^+$.

We introduce additional notation for subshifts. We denote the set of periodic points in A(X) by P(A(X)). The subshifts $X \subset \Sigma^{\mathbb{Z}}$: that we consider in this paper are such that P(A(X)) is dense in X. We introduce a preorder relation \gtrsim into the set P(A(X)) where for $q, r \in P(A(X))$ means that there exists a point in A(X)that is left asymptotic to the orbit of q and right asymptotic to the orbit of r. The equivalence relation on P(A(X)) that results from the preorder relation \gtrsim we denote by \approx .

The proof of the following theorem is similar to a proof for Markov-Dyck shifts in [KM2].

Theorem 3.2. Let $\mathcal{G}_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-}, \mathcal{E}^{+})$ be an \mathcal{R} -graph, that satisfies conditions (I) and (II), and let

$$\mathfrak{P}\setminus\mathfrak{P}_{\mathcal{R}}^{(1)}\neq\emptyset.$$

Let $X(\mathcal{V}, \Sigma, \lambda)$ be an $\mathcal{S}_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-}, \mathcal{E}^{+})$ -presentation, let q, r be neutral periodic points of $X(\mathcal{V}, \Sigma, \lambda)$, and let $\mathfrak{q}, \mathfrak{r} \in \mathfrak{P}$, and $I(q), I(r) \in \mathbb{Z}$ be such that

$$p_{(I(q),I(q)+\pi(q)]} = \mathbf{1}_{\mathfrak{q}}, \quad p_{(I(r),I(r)+\pi(r)]} = \mathbf{1}_{\mathfrak{r}}.$$

Then $q \approx r$ if and only if \mathfrak{q} and \mathfrak{r} are in the same element of the partition $\{\mathfrak{P}(\mathfrak{p}) :$ $\mathfrak{p} \in \widehat{\mathcal{R}}$.

Proof. Assume that $p \approx q$. Let $N \in \mathbb{N}$ and

(3.2)
$$x^{(q,r)}, x^{(r,q)} \in A_N(X(\mathcal{V}, \Sigma, \lambda)),$$

and let

$$I_{-}(q,r), I_{+}(q,r), \quad I_{-}(r,q), I_{+}(r,q) \in \mathbb{Z},$$
$$I_{-}(q,r) < I_{+}(q,r), \quad I_{-}(r,q) < I_{+}(r,q),$$

$$I_{-}(q,r) < I_{+}(q,r), \quad I_{-}(r,q) < I_{+}(r,q)$$

be such that

$$\begin{aligned} x_{(-\infty,I_{-}(q,r)]}^{(q,r)} &= q_{(-\infty,I(q)]}, \quad x_{(I_{+}(q,r),\infty)}^{(q,r)} = r_{(I(r),\infty)}, \\ x_{(-\infty,I_{-}(r,q)]}^{(r,q)} &= r_{(-\infty,I(r)]}, \quad x_{(I_{+}(r,q),\infty)}^{(r,q)} = q_{(I(q),\infty)}. \end{aligned}$$

By (3.2)

(3.3)
$$(x_{(-\infty,I_{+}(q,r)]}^{(q,r)}, r_{(I(r),I(r)+N\pi(r)]}, x_{(I_{-}(q,r),\infty)}^{(r,q)}) \in X(\mathcal{V}, \Sigma, \lambda),$$

and

(3.4)
$$(x_{(-\infty,I_{+}(r,q)]}^{(r,q)}, q_{(I(q),I(q)+N\pi(q)]}, x_{(I_{-}(r,q),\infty)}^{(q,r)}) \in X(\mathcal{V}, \Sigma, \lambda).$$

As a consequence of (3.2) there are also

$$\mathfrak{p}(q,r) \in \mathfrak{P}, \quad J_{-}(q,r), J_{+}(q,r) \in \mathbb{N},$$

and

$$\begin{split} e^+_{j_+(q,r)}(q,r) &\in \mathcal{E}^+_{\mathcal{R}}, \quad J_+(q,r) \ge j_+(q,r) > 0, \\ e^-_{j_-(q,r)}(q,r) &\in \mathcal{E}^-_{\mathcal{R}}, \quad 0 < j_-(q,r) \le J_-(q,r), \end{split}$$

such that

(3.5)
$$\lambda(x_{(I_{-}(q,r),I_{+}(q,r)]}^{(q,r)}) = \mathbf{1}_{\mathfrak{q}}(\prod_{J_{+}(q,r)\geq j_{+}(q,r)>0} e_{j_{+}}^{+}(q,r))\mathbf{1}_{\mathfrak{p}^{(q,r)}}(\prod_{0< j_{-}(q,r)\leq J_{-}(q,r)} e_{j_{-}(q,r)}^{-}(q,r))\mathbf{1}_{\mathfrak{r}},$$

and there are also

$$\mathfrak{p}(r,q) \in \mathfrak{P}, \quad J_{-}(r,q), J_{+}(r,q) \in \mathbb{N},$$

and

$$\begin{aligned} e^+_{j_+(r,q)}(q,r) &\in \mathcal{E}^+_{\mathcal{R}}, \quad J_+(r,q) \ge j_+(r,q) > 0, \\ e^-_{j_-(r,q)}(q,r) &\in \mathcal{E}^-_{\mathcal{R}}, \quad 0 < j_-(r,q) \le J_-(r,q), \end{aligned}$$

such that

$$(3.6) \quad \lambda(x_{(J_{-}(r,q),J_{+}(r,q)]}^{(r,q)}) = \\ \mathbf{1}_{\mathfrak{r}}(\prod_{J_{+}(r,q)\geq j_{+}(r,q)>0} e_{j_{+}}^{+}(r,q))\mathbf{1}_{\mathfrak{p}^{(r,q)}}(\prod_{0< j_{-}(r,q)\leq J_{-}(r,q)} e_{j_{-}(r,q)}^{-}(r,q))\mathbf{1}_{\mathfrak{q}}.$$

By (3.3), and in case that $J_{+}(q,r) \geq J_{+}(r,q)$ $\mathbf{1}_{\mathfrak{q}}(\prod_{J_{+}(q,r)\geq j_{+}(q,r)>0} e_{j_{+}}^{+}(q,r))\mathbf{1}_{\mathfrak{p}^{(q,r)}}(\prod_{0< j_{-}(q,r)\leq J_{-}(q,r)} e_{j_{-}(q,r)}^{-}(q,r))\mathbf{1}_{\mathfrak{r}}$ $\mathbf{1}_{\mathfrak{r}}(\prod_{J_{+}(r,q)\geq j_{+}(r,q)>0} e_{j_{+}}^{+}(r,q))\mathbf{1}_{\mathfrak{p}^{(r,q)}}(\prod_{0< j_{-}(r,q)\leq J_{-}(r,q)} e_{j_{-}(r,q)}^{-}(r,q))\mathbf{1}_{\mathfrak{q}} =$ $\mathbf{1}_{\mathfrak{q}}(\prod_{J_{+}(q,r)\geq j_{+}(q,r)>0} e_{j_{+}}^{+}(q,r))\mathbf{1}_{\mathfrak{p}^{(q,r)}}(\prod_{0< j_{-}(r,q)\leq J_{-}(r,q)} e_{j_{-}(r,q)}(r,q))\mathbf{1}_{\mathfrak{q}} \neq 0.$

From this it follows by Condition (I) that

$$(\widehat{e}_{j_{+}}^{+}(q,r))_{J_{+}(q,r) \ge j_{+}(q,r) > 0}$$

is a path in $\widehat{\mathcal{E}}^+(1)$ from \mathfrak{q} to $\mathfrak{p}^{(q,r)}$, and

$$((\widehat{e}^{-})_{0 < j_{-}(q,r) \le J_{-}(q,r) - J_{+}(r,q)}), (\widehat{e}^{-})_{0 < j_{-}(r,q) \le J_{-}(r,q)}))$$

is a path in $\widehat{\mathcal{E}}^{-}(1)$ from $\mathfrak{p}^{(q,r)}$ to \mathfrak{r} that passes through $\mathfrak{p}^{(r,q)}$ and one sees that $\mathfrak{p}^{(q,r)}, \mathfrak{p}^{(r,q)}$ and \mathfrak{q} are in the same element of the partition $\{\mathfrak{P}_{\mathfrak{p}} : \mathfrak{p} \in \mathfrak{P} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}\}$. By the same argument for the case that $J_{+}(q,r) \leq J_{+}(q,r)$, and by the symmetric argument that uses (3.6), one sees that in fact $\mathfrak{p}^{(q,r)}, \mathfrak{p}^{(r,q)}$ and \mathfrak{q} and \mathfrak{r} are in the same element of the partition $\{\mathfrak{P}_{\mathfrak{p}} : \mathfrak{p} \in \mathfrak{P} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}\}$.

For the proof of the converse let $p \in \mathfrak{P} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}$ and let $\mathfrak{q}, \mathfrak{r} \in \mathfrak{P}_{\mathfrak{p}}$. There is a path $(\widehat{e}_{j_{+}(q)}^{+})_{J_{+}(q) \geq j_{+}(q) > 0}$ in $\widehat{\mathcal{E}}^{+}(1)$ from \mathfrak{q} to \mathfrak{p} , a path $(\widehat{e}_{j_{-}}^{-}(r))_{0 < j_{-}(r) \leq J_{-}(r)}$ in $\widehat{\mathcal{E}}^{-}(1)$ from \mathfrak{p} to \mathfrak{r} , a path $(\widehat{e}_{j_{+}(r)}^{+})_{J_{+}(r) \geq j_{+}(r) > 0}$ in $\widehat{\mathcal{E}}^{+}(1)$ from \mathfrak{r} to \mathfrak{p} , and a path $(\widehat{e}_{j_{-}}^{-}(q))_{0 < j_{-}(r) \leq J_{-}(q)}$ in $\widehat{\mathcal{E}}^{-}(1)$ from \mathfrak{p} to \mathfrak{q} . Choose a vertex $V(p) \in \mathcal{V}(\mathfrak{p})$ and choose

$$e_{j_+(q)}^+ \in \widehat{e}_{j_+(q)}^+, \quad J_+(q) \ge j_+(q) > 0$$

and choose a path $b^+(q)$ in the directed graph $G(\mathcal{V}, \Sigma)$ from the target vertex of $q_{(-\infty,I(q)]}$ to V(p), such that

$$\lambda(b^+(q)) = \prod_{J_+(q) \ge j_+(q) > 0} e^+_{j_+(q)},$$

and choose

$$e_{j_{-}(r)}^{+} \in \hat{e}_{j_{-}(r)}^{+}, \quad 0 < j_{-}(r) \le J_{-}(r)$$

and also a path $b^{-}(r)$ from V(p) to the source vertex of $q_{(I(r),\infty)}$, such that

$$\lambda(b^{-}(r)) = \prod_{0 < j_{-}(r) \le J_{-}(r)} e^{-}_{j_{-}}(r),$$

choose

$$e_{j_+(r)}^+ \in \widehat{e}_{j_+(r)}^+, \quad J_+(q) \ge j_+(r) > 0$$

and choose a path $b^+(q)$ in the directed graph $G(\mathcal{V}, \Sigma)$ from the target vertex of $q_{(-\infty,I(q)]}$ to V(p), such that

$$\lambda(b^+(r)) = \prod_{J_+(r) \ge j_+(r) > 0} e^+_{j_+(r)},$$

and choose

$$e_{j_{-}(q)}^{-} \in \widehat{e}_{j_{-}(q)}^{-}, \quad 0 < j_{-}(q) \le J_{-}(q)$$

and also a path $b^-(q)$ from V(p) to the source vertex of $q_{(I(q),\infty)}$, such that

$$\lambda(b^{-}(q)) = \prod_{0 < j_{-}(q) \le J_{-}(q)} e^{-}_{j_{-}}(q),$$

Then

$$(q_{(-\infty,I(q)]}, b^+(q), b^-(r), r_{(I(r),\infty)}) \in A(X(\mathcal{V}, \Sigma, \lambda)),$$

$$(r_{(-\infty,I(r)]}, b^+(r), b^-(q), q_{(I(q),\infty)}) \in A(X(\mathcal{V}, \Sigma, \lambda)).$$

We recall at this point the construction of the associated semigroup. For a property (A) subshift $X \subset \Sigma^{\mathbb{Z}}$ we denote by Y(X) the set of points in X that are left asymptotic to a point in P(A(X)) and also right-asymptotic to a point in P(A(X)). Let $y, \tilde{y} \in Y(X)$, let y be left asymptotic to $q \in P(A(X))$ and right asymptotic to $r \in P(A(X))$, and let \tilde{y} be left asymptotic to $\tilde{q} \in P(A(X))$ and right asymptotic to $\tilde{r} \in P(A(X))$. Given that X has the properties $(a, n, H_n), n \in \mathbb{N}$, we say that y and \tilde{y} are equivalent, $y \approx \tilde{y}$, if $q \approx \tilde{q}$ and $r \approx \tilde{r}$, and if for $n \in \mathbb{N}$ such that $q, r, \tilde{q}, \tilde{r} \in P(A_n(X))$ and for $I, J, \tilde{I}, \tilde{J} \in \mathbb{Z}, I < J, \tilde{I} < \tilde{J}$, such that

$$\begin{split} y_{(-\infty,I]} &= q_{(-\infty,0]}, \quad y_{(J,\infty)} = r_{(0,\infty)}, \\ \tilde{y}_{(-\infty,\tilde{I}]} &= \tilde{q}_{(-\infty,0]}, \quad \tilde{y}_{(\tilde{J},\infty)} = \tilde{r}_{(0,\infty)}, \end{split}$$

one has for $h \geq 3H_n$ and for

$$a \in X_{(I-h,J+h]}, \quad \tilde{a} \in X_{(\tilde{I}-h,\tilde{J}+h]},$$

such that

$$a_{(I-H_n,J+H_n]} = y_{(I-H_n,J+H_n]}, \quad \tilde{a}_{(\tilde{I}-H_n,\tilde{J}+H_n]} = \tilde{y}_{(\tilde{I}-H_n,\tilde{J}+H_n]},$$
$$a_{(I-h,I-h+H_n)} = \tilde{a}_{(\tilde{I}-h,\tilde{I}-h+H_n)},$$
$$a_{(J+h-H_n,J+h]} = \tilde{a}_{(\tilde{J}+h-H_n,\tilde{J}+h]},$$

and such that

$$a_{(I-h,I]} \in A_n(X)_{(I-h,I]}, \quad \tilde{a}_{(\tilde{J}-h,\tilde{I}]} \in A_n(X)_{(\tilde{J}-h,\tilde{I}]}, \\ a_{(J,J+h]} \in A_n(X)_{(J,J+h]}, \quad \tilde{a}_{(\tilde{J},\tilde{J}+h]} \in A_n(X)_{(\tilde{J},\tilde{J}+h]},$$

that a and \tilde{a} have the same context. To give $[Y(X)]_{\approx}$ the structure of a semigroup, let $u, v \in Y(X)$, let u be right asymptotic to $q \in P(A(X))$ and let v be left asymptotic to $r \in P(A(X))$. If here $q \gtrsim r$, then $[u]_{\approx}[v]_{\approx}$ is set equal to $[y]_{\approx}$ where y is any point in Y such that there are $n \in \mathbb{N}, I, J, \hat{I}, \hat{J} \in \mathbb{Z}, I < J, \hat{I} < \hat{J}$, such that $q, r \in A_n(X)$, and such that

$$u_{(I,\infty)} = q_{(I,\infty)}, \quad v_{(-\infty,J]} = r_{(-\infty,J]},$$
$$y_{(-\infty,\hat{I}+H_n]} = u_{(-\infty,I+H_n]}, \quad y_{(\hat{J}-H_n,\infty)} = v_{(J-H_n,\infty)},$$

and

$$y_{(\hat{I},\hat{J}]} \in A_n(X)_{(\hat{I},\hat{J}]},$$

provided that such a point y exists. If such a point y does not exist, $[u]_{\approx}[v]_{\approx}$ is set equal to zero. Also, in the case that one does not have $q \gtrsim r, [u]_{\approx}[v]_{\approx}$ is set equal to zero.

Theorem 3.3. Let $G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^- \cup \mathcal{E}^+)$ be an \mathcal{R} -graph, such that

$$\mathfrak{P} \setminus \mathfrak{P}_{\mathcal{P}}^{(1)} \neq \emptyset,$$

that satisfies conditions (I) and (II), and let $X(\mathcal{V}, \Sigma, \lambda)$ be an $\mathcal{S}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^+ \cup \mathcal{E}^-))$ -presentation. Then the semigroup $\mathcal{S}(G_{\widehat{\mathcal{R}}}(\widehat{\mathfrak{P}}, \widehat{\mathcal{E}}^+ \cup \widehat{\mathcal{E}}^-))$ is associated to $X(\mathcal{V}, \Sigma, \lambda)$, and $X(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^+ \cup \mathcal{E}^-))$ has an $\mathcal{S}(G_{\widehat{\mathcal{R}}}(\widehat{\mathfrak{P}}, \widehat{\mathcal{E}}^+ \cup \widehat{\mathcal{E}}^-))$ -presentation.

Proof. We choose for $\mathfrak{r} \in \widehat{\mathfrak{R}}$ a periodic point $y^{\circ}(\mathfrak{r}) \in X(\mathfrak{P}, \Sigma, \lambda)$, such that

$$\lambda(_{[0,\pi(y^{\circ}(\mathfrak{r})))}=\mathbf{1}_{\mathfrak{r}}.$$

By Lemma (3.1) and by Theorem (3.2) we can choose a system of representatives Y° of the equivalence relation \approx such that $Y^{\circ} \supset \{y^{\circ}(\mathfrak{r}) : \mathfrak{r} \in \mathfrak{R}_{\widehat{\mathcal{R}}}\}$, and such that every point in Y° is left asymptotic to a point in $\{y^{\circ}(\mathfrak{r}) : \mathfrak{r} \in \mathfrak{R}_{\widehat{\mathcal{R}}}\}$ and also right asymptotic to a point in $\{y^{\circ}(\mathfrak{r}) : \mathfrak{r} \in \mathfrak{R}_{\widehat{\mathcal{R}}}\}$.

We set

$$\begin{split} \varphi(e^{-}) &= [e^{-}]_{\sim(\mathcal{R},-)}, \qquad e^{-} \in \mathcal{E}^{-}, \\ \varphi(f^{-}) &= \mathbf{1}_{\mathfrak{r}}, \qquad [f^{-}]_{\sim(\mathcal{R},-)} \in \bar{\mathcal{F}}_{\bar{\mathcal{R}}}^{-}(\mathfrak{r}), \mathfrak{r} \in \widehat{\mathfrak{R}}, \\ \varphi(\mathbf{1}_{\mathfrak{p}}) &= \mathbf{1}_{\mathfrak{r}}, \qquad \mathfrak{p} \in \mathfrak{P}_{\mathfrak{r}}, \mathfrak{r} \in \widehat{\mathfrak{R}}, \\ \varphi(f^{+}) &= \mathbf{1}_{\mathfrak{r}}, \qquad [f^{+}]_{\sim(\mathcal{R},+)} \in \bar{\mathcal{F}}_{\bar{\mathcal{R}}}^{+}(\mathfrak{r}), \mathfrak{r} \in \widehat{\mathfrak{R}}, \\ \varphi(e^{+}) &= [\widehat{e^{+}}]_{\sim(\mathcal{R},+)}, \qquad e^{+} \in \mathcal{E}^{+}. \end{split}$$

and for

$$g(-) \in \mathcal{S}^{-}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+})), \qquad g(+) \in \mathcal{S}^{+}(G_{\mathcal{R}}(\mathfrak{P}, \mathcal{E}^{-} \cup \mathcal{E}^{+}))$$

we set

$$\varphi(g^{-}) = \prod_{1 \le i(-) \le \ell(g(-))} \varphi(e_{i(-)}[g(-)]), \quad \varphi(g^{+}) = \prod_{1 \le i(+) \le \ell(g(+))} \varphi(e_{i(+)}[g(+)]).$$

We set

$$\widehat{\lambda}(\sigma) = \varphi(\lambda(\sigma)), \qquad \sigma \in \Sigma,$$

and

$$\begin{aligned} J_{-}(y^{\circ}) &= \max\{J < 0 : y^{\circ}_{(-\infty,J]} \in \{y^{\circ}(\mathfrak{r})_{(-\infty,J]} : \mathfrak{r} \in \mathfrak{R}\}\},\\ J_{-}(y^{\circ}) &= \min\{J > 0 : y^{\circ}_{[J,-\infty)} \in \{y^{\circ}(\mathfrak{r})_{[J,-\infty)} : \mathfrak{r} \in \mathfrak{R}\}\}, \qquad y^{\circ} \in Y^{\circ}. \end{aligned}$$

An isomorphism Ψ of $\mathcal{S}(X(\lambda))$ onto $\mathcal{S}(\widehat{G}(\lambda))$ is given by

$$\Psi([(y^{\circ})]_{\approx}) = \prod_{J_{-}(y^{\circ}) < j < J_{+}(y^{\circ})} \widehat{\lambda}(y^{\circ}_{j}), \qquad y^{\circ} \in Y^{\circ}. \quad \Box$$

ore precisely, with $I, J \in \mathbb{Z}, I < J$, such that

4. Examples

Let there be given an \mathcal{R} -graph

$$G_{\widehat{\mathcal{R}}} = G_{\widehat{R}}(\{\widehat{\mathfrak{p}}\}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+),$$

such that

$$\begin{aligned} & \operatorname{card}(\widehat{\mathcal{E}}^{-}) > 1, \\ & [\widehat{e}^{-}]_{\sim \widehat{\mathcal{R}}, -} = \{\widehat{e}^{-}\}, \qquad \widehat{e}^{-} \in \widehat{\mathcal{E}}^{-}, \\ & [\widehat{e}^{+}]_{\sim \widehat{\mathcal{R}}, -+} = \{\widehat{e}^{-}\}, \qquad \widehat{e}^{-} \in \widehat{\mathcal{E}}^{+}. \end{aligned}$$

Let there also be given a subshift X with Property (A) and associated semigroup $\mathcal{S}(G_{\widehat{R}})$. For $\widehat{e}^- \in \widehat{\mathcal{E}}^-(\widehat{e}^- \in \widehat{\mathcal{E}}^-)$ we denote by $\Lambda(\widehat{e}^-)(\Lambda(\widehat{e}^+))$ the minimal length of a periodic point in X with multiplier $\widehat{e}^-(\widehat{e}^+)$, and we denote by $P_l^{(\widehat{e}^-)}(P_l^{(\widehat{e}^+)})$ the set of periodic points of X of length $l \ge \Lambda(\widehat{e}^-)$ $(l \ge \Lambda(\widehat{e}^+))$, with multiplier $\widehat{e}^-(\widehat{e}^+)$. We set

$$P^{(-)} = \bigcup_{l \ge \Lambda(\widehat{e}^{-})} P_l^{(\widehat{e}^{-})}, \qquad P^{(+)} = \bigcup_{l \ge \Lambda(\widehat{e}^{+})} P_l^{(\widehat{e}^{+})}.$$

4.1. One-vertex \mathcal{R} -graphs.

Theorem 4.1. For \mathcal{R} -graphs

$$G_{\mathcal{R}} = G_{\mathcal{R}}(\{\mathfrak{p}\}, \mathcal{E}^- \cup \mathcal{E}^+),$$

such that

$$\mathfrak{p} \notin \mathfrak{P}_{\mathcal{R}}^{(1)}, \qquad \mathcal{E}_{\mathcal{R}}^{-}(\mathfrak{p},\mathfrak{p}) = \mathcal{E}_{\mathcal{R}}^{+}(\mathfrak{p},\mathfrak{p}) = \emptyset,$$

the topological conjugacy class of the \mathcal{R} -graph shift of $G_{\mathcal{R}}$ determines the isomorphism class of the \mathcal{R} -graph $G_{\mathcal{R}}$.

Proof. We define a relation $\mathcal{Q} \subset P_1^{(-)} \times P_1^{(+)}$ that contains the pairs $(p(-), p(+)) \in P_1^{(-)} \times P_1^{(+)}$, such that the subshift $X(G_{\mathcal{R}})$ has a point, that is left asymptotic to p(-) and right asymptotic to p(+).

To obtain mappings

$$\widehat{\nu}_{-}:\widehat{\mathcal{E}}^{-}\to\mathbb{N},\ \widehat{\nu}_{+}:\widehat{\mathcal{E}}^{+}\to\mathbb{N},$$

such that one has for

$$p(-) \in P^{(\widehat{e}^-)}, p(+) \in P^{(\widehat{e}^+)}, \qquad (\widehat{e}^-, \widehat{e}^+) \in \widehat{\mathcal{R}},$$

that

(4.1)
$$\operatorname{card}(\{\widetilde{p}(+) \in P_1^{(\widehat{e}^-)}(X) : (p(-), \widetilde{p}(+)) \in \mathcal{Q}\}) = \widehat{\nu}_-(\widehat{e}^-)$$
$$\operatorname{card}(\{\widetilde{p}(-) \in P_1^{(\widehat{e}^-)}(X) : (\widetilde{p}(-), p(+)) \in \mathcal{Q}\}) = \widehat{\nu}_+(\widehat{e}^+)$$

set

$$\begin{aligned} \widehat{\nu}_{-}(\widehat{e}^{-}) &= \operatorname{card}(\widehat{e}^{-}), \qquad \widehat{e}^{-} \in \widehat{\mathcal{E}}^{-}, \\ \widehat{\nu}_{+}(\widehat{e}^{+}) &= \operatorname{card}(\widehat{e}^{+}). \qquad \widehat{e}^{+} \in \widehat{\mathcal{E}}^{+}. \end{aligned}$$

The existence of mappings

$$\widehat{\nu}_{-}:\widehat{\mathcal{E}}^{-}\to\mathbb{N},\ \widehat{\nu}_{+}:\widehat{\mathcal{E}}^{+}\to\mathbb{N}$$

such that (4.1) holds is an invariant of topological conjugacy. Given the mappings $\widehat{\nu}_{-}: \widehat{\mathcal{E}}^{-} \to \mathbb{N}, \ \widehat{\nu}_{+}: \widehat{\mathcal{E}}^{+} \to \mathbb{N},$

such that (4.1) holds, together with the \mathcal{R} -graph

$$G_{\widehat{R}} = G_{\widehat{R}}(\{\mathfrak{p}\}, [\mathcal{E}^{-}]_{\sim(\mathcal{R}, -)} \cup [\mathcal{E}^{+}]_{\sim(\mathcal{R}, +)})$$

that is derived from $G_{\mathcal{R}}$, one can reconstruct the \mathcal{R} -graph $G_{\mathcal{R}}$ from the relation \widehat{R} .

4.2. A class of Examples of \mathcal{R} -graph shifts, to which the \mathcal{R} -graph semigroups of one-vertex \mathcal{R} -graphs are associated. Let $K \in \mathbb{N}$. We set

$$\begin{split} &\widehat{\mathcal{E}}_l^-(X) = \{\widehat{e}^- \in \widehat{\mathcal{E}}^- : \Lambda(\widehat{e}^-) = l+1\}, \\ &\widehat{\mathcal{E}}_l^+(X) = \{\widehat{e}^+ \in \widehat{\mathcal{E}}^+ : \Lambda(\widehat{e}^+) = l+1\}, \quad 0 < l \leq K, \end{split}$$

and

$$\widehat{\mathcal{R}}_l(X) = \widehat{\mathcal{R}} \cap (\widehat{\mathcal{E}}_l^-(X) \times \widehat{\mathcal{E}}_l^+(X)), \quad 0 < l \le K.$$

We also define relations

$$Q_l(X) \subset P_{l+1}^{(-)}(X) \times P_{l+1}^{(+)}(X), \quad 0 < l \le K,$$

 $Q_l(X)$ containing the pairs

$$(p(-), p(+)) \in P_{l+1}^{(-)}(X) \times P_{l+1}^{(+)}(X),$$

such that the subshift X has a point, that is left asymptotic to p(-) and right asymptotic to p(+).

We denote by $\mathcal{H}(G_{\widehat{R}})$ the family of subshifts X with Property (A) and associated semigroup $\mathcal{S}(G_{\widehat{R}})$, that satisfy the following conditions (c1-4), that are designed to be invariant under topological conjugacy:

(c1)
$$\widehat{\mathcal{E}}^{-} = \prod_{0 < l \le K} \widehat{\mathcal{E}}_{l}^{-}(X), \qquad \widehat{\mathcal{E}}^{+} = \prod_{0 < l \le K} \widehat{\mathcal{E}}_{l}^{+}(X).$$

(c2)
$$\widehat{\mathcal{R}} = \bigcup_{0 < l \le K} \widehat{\mathcal{R}}_l(X).$$

(c3) There are $\Xi_{K,l} \in \mathbb{N}, 0 < l \leq K$, such that

$$\Xi_l = \operatorname{card}(P_{l+3}^{(e^-)}) = \operatorname{card}(P_{l+3}^{(e^+)}), \quad \widehat{e}^- \in \widehat{\mathcal{E}}^-, \ \widehat{e}^- \in \widehat{\mathcal{E}}^-, \ 0 < l \le K.$$

(c3) There are mappings

$$\widehat{\nu}_{-,l}:\widehat{\mathcal{E}}_l^- \to \mathbb{N}, \quad \widehat{\nu}_{+,l}:\widehat{\mathcal{E}}_l^+ \to \mathbb{N}, \qquad 0 < l \leq K.$$

such that it holds for

$$p(-) \in P_l^{(\widehat{e}^-)}(X), p(+) \in P_l^{(\widehat{e}^+)}(X), \qquad (\widehat{e}^-, \widehat{e}^+) \in \mathcal{R}_l(X)(X),$$

that

$$\operatorname{card}(\{\widetilde{p}(+) \in P_l^{(\widehat{e}^-)}(X) : (p(-), \widetilde{p}(+)) \in \mathcal{Q}_l(X)\}) = \widehat{\nu}_{-,l}(\widehat{e}^-),$$
$$\operatorname{card}(\{\widetilde{p}(-) \in P_l^{(\widehat{e}^-)}(X) : (\widetilde{p}(-), p(+)) \in \mathcal{Q}_l(X)\}) = \widehat{\nu}_{+,l}(\widehat{e}^+).$$

Let $K \in \mathbb{N}$, and let there be given an \mathcal{R} -graph

$$G_{\mathcal{R}} = G_{\mathcal{R}}(\{\mathfrak{p}_l : 0 \le l \le K\}, \mathcal{E}^- \cup \mathcal{E}^-),$$

such that

(4.2)
$$\mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0}) \neq \emptyset, \quad \mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l}) \neq \emptyset, \qquad 0 \leq l \leq K,$$

(4.3)
$$\mathcal{E}^{-} = \bigcup_{0 \le l \le K} (\mathcal{E}^{-}(\mathfrak{p}_{l}, \mathfrak{p}_{0}) \cup \mathcal{E}^{-}(\mathfrak{p}_{l-1}, \mathfrak{p}_{l})).$$

(4.4)
$$\mathfrak{p}_l \in \mathfrak{P}_{\mathcal{R}}^{(1)}, \qquad 0 \le l \le K,$$

In the case that K = 1, also assume, that

$$\mathfrak{p}_0 \in \mathfrak{P}^{(1)} \setminus \mathfrak{P}^{(1)}_{\mathcal{R}}.$$

Let $G_{\widehat{R}} = G_{\widehat{R}}(\{\widehat{\mathfrak{p}}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+\})$ be the \widehat{R} -graph, that is derived from $G_{\mathcal{R}}$. We set $\widehat{R}_l = \widehat{R} \cap ([\mathcal{E}^-(\mathfrak{p}_l, \mathfrak{p}_0)]_{\sim (\mathcal{R}, +)} \times [\mathcal{E}^+(\mathfrak{p}_l, \mathfrak{p}_0)]_{\sim (\mathcal{R}, -)}), \quad 0 \le l \le K,$

and we introduce a condition (D) on the \mathcal{R} -graph $G_{\mathcal{R}}$, that consists of two parts, (Da) and (Db):

(Da)
$$gcd(card(\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})card(\mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})), \sum_{(\widehat{e}^{-},\widehat{e}^{+})\in\widehat{R}_{l}}card(\widehat{e}^{-})card(\widehat{e}^{+})) = 1,$$

(Db)
$$gcd\{\operatorname{card}(\widehat{e}^{-}): \widehat{e}^{-} \in [\mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0})]_{\sim(\mathcal{R},-)}\} =$$

 $gcd\{\operatorname{card}(\widehat{e}^{-}): \widehat{e}^{-} \in [\mathcal{E}^{+}(\mathfrak{p}_{l},\mathfrak{p}_{0})]_{\sim(\mathcal{R},+)}\} = 1, \quad 0 \le l \le K.$

Theorem 4.2. Let $K \in \mathbb{N}$, and let

$$G_{\mathcal{R}} = G_{\mathcal{R}}(\{\mathfrak{p}_l : 0 \le l \le K\}, \mathcal{E}^- \cup \mathcal{E}^-),$$

be an \mathcal{R} -graph, that satisfies (4.2), (4.3) and (4.4) and such that, in the case that K = 1, also $\mathfrak{p}_0 \in \mathfrak{P}^{(1)} \setminus \mathfrak{P}_{\mathcal{R}}^{(1)}$.

Condition (D) is an invariant of topological conjugacy of the \mathcal{R} -graph shift of $G_{\mathcal{R}}$. If the \mathcal{R} -graph shift of $G_{\mathcal{R}}$ satisfies Condition (D), then its topological conjugacy class determines the isomorphism class of the \mathcal{R} -graph $G_{\mathcal{R}}$.

Proof. Let $G_{\widehat{R}} = G_{\widehat{R}}(\{\widehat{\mathfrak{p}}\}, \widehat{\mathcal{E}}^- \cup \widehat{\mathcal{E}}^+)$ be the \widehat{R} -graph, that is derived from $G_{\mathcal{R}}$. The \mathcal{R} -graph shift of $G_{\mathcal{R}}$ belongs to the family $\mathcal{H}_K(G_{\widehat{R}})$: By construction (c1) and (c2) are satisfied. One has, that

$$\widehat{\mathcal{E}}_{l}^{-}(X(G_{\mathcal{R}})) = [\mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0})]_{\sim(\mathcal{R},+)}, \quad \widehat{\mathcal{E}}_{l}^{+}(X(G_{\mathcal{R}})) = [\mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0})]_{\sim(\mathcal{R},-)}, \\
\widehat{\mathcal{R}}_{l}(X(G_{\mathcal{R}})) = \widehat{\mathcal{R}} \cap (\widehat{\mathcal{E}}_{l}^{-}(X(G_{\mathcal{R}})) \times \widehat{\mathcal{E}}_{l}^{+}(X(G_{\mathcal{R}}))), \qquad 0 \le l \le K.$$

Condition (c3) is also satisfied: One has that

(4.5)
$$\Xi_{K,l}(X(G_{\mathcal{R}})) =$$

$$\sum_{0 < m \le l} (\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{m-1},\mathfrak{p}_{m}))\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{m-1},\mathfrak{p}_{m})) + \sum_{(\widehat{e}^{-},\widehat{e}^{+})\in\widehat{\mathcal{R}}_{m}} \operatorname{card}(\widehat{e}^{-})\operatorname{card}(\widehat{e}^{+})),$$

$$0 < l \le K.$$

The Condition (c4) is also satisfied. One has, that

$$\nu_{-,l}^{X(G_{\mathcal{R}})}(\widehat{e}^{-}) = \operatorname{card}(\widehat{e}^{-}) \prod_{0 < m \le l} \operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{m-1}, \mathfrak{p}_m)), \quad \widehat{e}^{-} \in \mathcal{E}_l^{-},$$
$$\nu_{-,l}^{X(G_{\mathcal{R}})}(\widehat{e}^{+}) = \operatorname{card}(\widehat{e}^{+}) \prod_{0 < m \le l} \operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{m-1}, \mathfrak{p}_m)), \quad \widehat{e}^{+} \in \mathcal{E}_l^{+}, \qquad 0 < l \le K.$$

We set

$$\begin{split} \rho_{-,l}^{X(G_{\mathcal{R}})} &= gcd\{\nu_{-,l}^{X(G_{\mathcal{R}})}(\widehat{e}^{-}): \widehat{e}^{-} \in \mathcal{E}_{l}^{-}(X(G_{\mathcal{R}}))\},\\ \rho_{+,l}^{X(G_{\mathcal{R}})} &= gcd\{\nu_{+,l}^{X(G_{\mathcal{R}})}(\widehat{e}^{+}): \widehat{e}^{+} \in \mathcal{E}_{l}^{+}(X(G_{\mathcal{R}}))\}. \end{split}$$

The proof of the theorem is by induction in K steps. For the *l*-th step of the induction we split Condition (D) into conditions (Dal) and (Dbl):

$$(Dal) \quad gcd(\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})), \sum_{(\widehat{e}^{-},\widehat{e}^{+})\in\widehat{R}_{l}}\operatorname{card}(\widehat{e}^{-})\operatorname{card}(\widehat{e}^{+})) = 1$$
$$(Dbl) \quad gcd\{\widehat{e}^{-}:\widehat{e}^{-}\in[\mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0})]_{\sim(\mathcal{R},-)}\} = gcd\{\widehat{e}^{+}:\widehat{e}^{+}\in[\mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0})]_{\sim(\mathcal{R},-)}\} = 1,$$
$$0 < l \leq K.$$

At the l-th step of the induction the cardinalities of the sets

$$\mathcal{E}^{-}(\mathfrak{p}_{m-1},\mathfrak{p}_m), \ \mathcal{E}^{+}(\mathfrak{p}_{m-1},\mathfrak{p}_m), \quad \mathcal{E}^{-}(\mathfrak{p}_m,\mathfrak{p}_0), \ \mathcal{E}^{+}(\mathfrak{p}_m,\mathfrak{p}_0), \qquad 0 < m < l,$$

are known. In the *l*-th step of the induction it is shown, that the validity of (Dal) is determined by the topological conjugacy class of $X(G_{\mathcal{R}})$. Under the assumption, that (Dal) holds, it is then also shown in the *l*-th step of the induction, that the validity of (Dbl) is determined by the topological conjugacy class of $X(G_{\mathcal{R}})$. Under the assumption, that (Dal) and (Dbl) hold, it is then shown, in the *l*-th step of the induction, how the cardinality of the edge sets

$$\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l}), \ \mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}), \quad \mathcal{E}^{-}(\mathfrak{p}_{l},\mathfrak{p}_{0}), \ \mathcal{E}^{+}(\mathfrak{p}_{l},\mathfrak{p}_{0}),$$

can be obtained from topological conjugacy invariants of $X(G_{\mathcal{R}})$.

We describe for $l \in [1, K]$ the *l*-th step of the induction. One has, that (4.6)

$$\operatorname{card}(\mathcal{Q}_{l}(X(G_{\mathcal{R}}))) = (\prod_{0 < m \leq l} (\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{m-1},\mathfrak{p}_{m})\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{m-1},\mathfrak{p}_{m})))(\sum_{(\widehat{e}^{-},\widehat{e}^{+})\in\widehat{\mathcal{R}}_{l}} \operatorname{card}(\widehat{e}^{-})\operatorname{card}(\widehat{e}^{+})) > \rho_{-,l}X(G_{\mathcal{R}}))\rho_{-,l}X(G_{\mathcal{R}}))\rho_{-,l}X(G_{\mathcal{R}}).$$
Set $\Xi_{0}(X(G_{\mathcal{R}}) = 0, \text{ and set}$

$$\sum_{X(G_{\mathcal{R}})} \sum_{i=1}^{X(G_{\mathcal{R}})} \sum_{i=1}^{X($$

$$\alpha_l^{X(G_{\mathcal{R}})} = \Xi_l(X(G_{\mathcal{R}}))) - \Xi_{l-1}(X(G_{\mathcal{R}}))),$$

$$\beta_l^{X(G_{\mathcal{R}})} = \frac{\operatorname{card}(\mathcal{Q}_l(X(G_{\mathcal{R}}))))}{\prod_{0 < m \le l} (\operatorname{card}(\mathcal{E}^-(\mathfrak{p}_{m-1},\mathfrak{p}_m)\operatorname{card}(\mathcal{E}^+(\mathfrak{p}_{m-1},\mathfrak{p}_m)))}, \quad 0 < l \le K.$$

From (4.5) one has, that

(4.7)

$$\alpha_l^{X(G_{\mathcal{R}})} = \operatorname{card}(\mathcal{E}^-(\mathfrak{p}_{l-1},\mathfrak{p}_l)\operatorname{card}(\mathcal{E}^+(\mathfrak{p}_{l-1},\mathfrak{p}_l)) + \sum_{(\widehat{e}^-,\widehat{e}^+)\in\widehat{\mathcal{R}}_l}\operatorname{card}(\widehat{e}^-)\operatorname{card}(\widehat{e}^+),$$

and from (a) one has, that

(4.8)
$$\beta_l^{X(G_{\mathcal{R}})} = \operatorname{card}(\mathcal{E}^-(\mathfrak{p}_{l-1},\mathfrak{p}_l)\operatorname{card}(\mathcal{E}^+(\mathfrak{p}_{l-1},\mathfrak{p}_l)) \sum_{(\widehat{e}^-,\widehat{e}^+)\in\widehat{\mathcal{R}}_l} \operatorname{card}(\widehat{e}^-)\operatorname{card}(\widehat{e}^+).$$

Set

$$\Delta_l^{X(G_{\mathcal{R}})} = (\alpha_l^{X(G_{\mathcal{R}})})^2 - 4\beta_l^{X(G_{\mathcal{R}})}.$$

It follows from (4.7) and (4.8), that

 $\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}_{l}))\in$

$$\{\frac{1}{2}(\alpha_l^{X(G_{\mathcal{R}})} - \sqrt{\Delta_l^{X(G_{\mathcal{R}})}}), \frac{1}{2}(\alpha_l^{X(G_{\mathcal{R}})} + \sqrt{\Delta_l^{X(G_{\mathcal{R}})}})\}.$$

At this stage the cardinalities of the sets

$$\mathcal{E}^{-}(\mathfrak{p}_{m-1},\mathfrak{p}_m), \mathcal{E}^{+}(\mathfrak{p}_{m-1},\mathfrak{p}_m), \qquad 0 < m < l,$$

are known, and are also known to be invariants of topological conjugacy. This means that $\beta_l^{X(G_{\mathcal{R}})}$ is also an invariant of topological conjugacy, as is the set of roots

$$\{\frac{1}{2}(\alpha_l^{X(G_{\mathcal{R}})} - \sqrt{\Delta_l^{X(G_{\mathcal{R}})}}), \frac{1}{2}(\alpha_l^{X(G_{\mathcal{R}})} + \sqrt{\Delta_l^{X(G_{\mathcal{R}})}})\}$$

It follows, that the validity of Condition (Dal) is an invariant of topological conjugacy.

Under the assumption, that (Dal) holds, one has, that either $\alpha_l^{X(G_{\mathcal{R}})} - \sqrt{\Delta_l^{X(G_{\mathcal{R}})}}$ or $\alpha_l^{X(G_{\mathcal{R}})} + \sqrt{\Delta_l^{X(G_{\mathcal{R}})}}$ is a divisor of $\rho_{-,l}^{X(G_{\mathcal{R}})} \rho_{+,l} X(G_{\mathcal{R}})$), since by (4.6) $\frac{\rho_{-,l}^{X(G_{\mathcal{R}})} \rho_{+,l}^{X(G_{\mathcal{R}})}}{\operatorname{card}(\mathcal{Q}_l^{X(G_{\mathcal{R}})})} < 1.$

Therefore one has, under the assumption, that (Dal) holds, that

(4.9)
$$\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})) = \begin{cases} \alpha_{l}^{X(G_{\mathcal{R}})} - \sqrt{\Delta_{l}^{X(G_{\mathcal{R}})}}, & \text{if } \alpha_{l}^{X(G_{\mathcal{R}})} - \sqrt{\Delta_{l}^{X(G_{\mathcal{R}})}} \mid \rho_{-,l}^{X(G_{\mathcal{R}})}\rho_{+,l}^{X(G_{\mathcal{R}})}, \\ \alpha_{l}^{X(G_{\mathcal{R}})} - \sqrt{\Delta_{l}^{X(G_{\mathcal{R}})}}, & \text{if } \alpha_{l}^{X(G_{\mathcal{R}})} - \sqrt{\Delta_{l}^{X(G_{\mathcal{R}})}} \mid \rho_{-,l}^{X(G_{\mathcal{R}})}\rho_{+,l}^{X(G_{\mathcal{R}})}. \end{cases}$$

Condition (Dbl) is equivalent to

$$\rho_{-,l}^{X(G_{\mathcal{R}})}\rho_{+,l}^{X(G_{\mathcal{R}})} = \prod_{0 < m \le l} [\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l})\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}_{l}))].$$

From this and from (4.9) it is seen, that under the assumption, that Condition (Dal) holds, the validity of Condition (Dbl) is an invariant under topological conjugacy. From Condition (Dal) together with Condition (Dbl) it follows that

$$\operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{l-1},\mathfrak{p}_{l}) = \frac{\rho_{-,l}^{X(G_{\mathcal{R}})}}{\prod_{0 < m < l} \operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{m-1},\mathfrak{p}_{m}))},$$
$$\operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{l-1},\mathfrak{p}_{l}) = \frac{\rho_{+,l}^{X(G_{\mathcal{R}})}}{\prod_{0 < m < l} \operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{m-1},\mathfrak{p}_{m}))},$$

and

$$\operatorname{card}(\widehat{e}^{-}) = \frac{\nu_{-,l}^{X(G_{\mathcal{R}})}}{\prod_{0 < m < l} \operatorname{card}(\mathcal{E}^{-}(\mathfrak{p}_{m-1}, \mathfrak{p}_{m}))}, \qquad \widehat{e}^{-} \in \widehat{\mathcal{E}}^{-},$$
$$\operatorname{card}(\widehat{e}^{+}) = \frac{\nu_{+,l}^{X(G_{\mathcal{R}})}}{\prod_{0 < m < l} \operatorname{card}(\mathcal{E}^{+}(\mathfrak{p}_{m-1}, \mathfrak{p}_{m}))}, \qquad \widehat{e}^{+} \in \widehat{\mathcal{E}}^{+},$$

and the relation \mathcal{R}_l can be reconstructed from

$$(\operatorname{card}(\widehat{e}^+))_{\widehat{e}^+\in\widehat{\mathcal{E}}_l^-}, \quad (\operatorname{card}(\widehat{e}^-))_{\widehat{e}^-\in\widehat{\mathcal{E}}_l^-}$$

and from the relation $\widehat{\mathcal{R}}_l$.

References

- [AH] C. J. ASH, T. E. HALL, Inverse semigroups on graphs, Semigroup Forum 11 (1975), 140–145.
- [BBD1] M. P. BÉAL, M. BLOCKELET, C. DIMA, Finite-type-Dyck shift spaces, arXiv:1311.4223 (2013)
- [BBD2] M. P. BÉAL, M. BLOCKELET, C. DIMA, Sofic-Dyck shifts, MFCS 2014, Part I, LNCS 8634, 63 - 74, (2014)
- [BBD3] M. P. BÉAL, M. BLOCKELET, C. DIMA, Sofic-Dyck shifts, Theoretical Computer Science 609 (2016), 226 – 244.
- [CS] A. COSTA, B. STEINBERG, A categorical invariant of flow equivalence of shifts, Ergod. Th. & Dynam. Sys. 36 (2016), 470 – 513.
- [CK] J. CUNTZ, W. KRIEGER, A class of C*-algebras and topological Markov chains, Inventiones Math. 56 (1980), 251 – 268.
- [HI] T. HAMACHI, K. INOUE, Embeddings of shifts of finite type into the Dyck shift, Monatsh. Math. 145 (2005), 107 – 129.
- [HIK] T. HAMACHI, K. INOUE, W. KRIEGER, Subsystems of finite type and semigroup invariants of subshifts, J. Reine Angew. Math. 632 (2009), 37 – 61.
- [HK1] T. HAMACHI, W. KRIEGER, On certain subshifts and their associated monoids, Ergod. Th. & Dynam. Sys. 36 (2016), 96 – 107
- [HK2] T. HAMACHI, W. KRIEGER, Families of directed graphs and topological conjugacy of the associated Markov-Dyck shifts, arXiv:1806.09051 (2018)
- [Ki] B. P. KITCHENS, Symbolic dynamics, Springer, Berlin, Heidelberg, New York (1998),
- [Kr1] W. KRIEGER, On the uniqueness of the equilibrium state, Math. Systems Theory 8 (1974), 97 104
- [Kr2] W. KRIEGER, On a syntactically defined invariant of symbolic dynamics, Ergod. Th. & Dynam. Sys. 20 (2000), 501 – 516
- [Kr3] W. KRIEGER, On flow-equivalence of *R*-graph shifts, Münster J. of Math. 8 (2015) 229 239.
- [Kr4] W. KRIEGER, On subshift presentations, Ergod. Th. & Dynam. Sys. 37 (2017) 1253 1290.
- [KM1] W. KRIEGER, K. MATSUMOTO, A notion of synchronization of symbolic dynamics and a class of C*-algebras, Acta Appl. Math. 126 (2013), 263 – 275.
- [KM2] W. KRIEGER, K. MATSUMOTO, Markov-Dyck shifts, neutral periodic points and topological conjugacy, Discrete and Continuous Dynamical systems 39 (2019), 1 – 18

- M. V. LAWSON, *Inverse semigroups*, World Scientific, Singapore, New Jersey, London and Hong Kong (1998).
- [LM] D. LIND AND B. MARCUS, An introduction to symbolic dynamics and coding, Cambridge University Press, Cambridge (1995).
- [M] K. MATSUMOTO, C*-algebras arising from Dyck systems of topological Markov chains, Math. Scand. 109 (2011), 31 – 54.
- [NP] M. NIVAT, J.-F. PERROT, Une généralisation du monoide bicyclique, C. R. Acad. Sc. Paris, Ser. A. 271 (1970), 824 – 827.

Toshihiro Hamachi Faculty of Mathematics Kyushu University 744 Motooka, Nishi-ku, Fukuoka 819-0395 Japan t.hamachi.796@m.kyushu-u.ac.jp

Wolfgang Krieger Institute for Applied Mathematics University of Heidelberg Im Neuenheimer Feld 205, 69120 Heidelberg Germany krieger@math.uni-heidelberg.de