

# HYPERELLIPTIC CURVES COVERING AN ELLIPTIC CURVE TWICE

XAVIER XARLES

*To the memory of F. Momose*

Let  $K$  be a field and let  $E$  be an elliptic curve over  $K$ . A natural question that have been considered by Mestre in [1] is the existence of an hyperelliptic curve  $H$  defined over  $K$  with two independent maps from  $H$  to  $E$ , hence such that its Jacobian  $\text{Jac}(H)$  is isogenous to  $E^2 \times A$ , for some abelian variety  $A$ . Mestre constructs in (op. cit.) such a curve  $H$  with genus 6 (for a field with characteristic 0). In this short note we show that there exists such a curve, but with genus 5, and for any field of characteristic  $\neq 2, 3$ . We don't know if the characteristic 2 and 3 cases can be solved using the same methods. We also show that, if we want such a curve  $H$  to be just geometrically hyperelliptic, so having a degree two map to a conic, then there is one with genus 3. It is known that there does not exist such a curve with genus 2 for a general elliptic curve over a general field.

These results have consequences on the distribution of rank  $\geq 2$  twists of elliptic curves over  $\mathbb{Q}$ , as showed by Steward and Top ([3], see also [2]).

We start with the second result. The following theorem was already been shown (but not in this form) by Mestre in a remark in [1] and by Steward and Top in [3] during the proof of their Theorem 4. The proof is elementary and we leave it to the reader.

**Theorem 1.** *Let  $E$  be an elliptic curve over  $K$  given by a Weierstrass equation of the form  $y^2 = x^3 - Ax + B$ , for some  $A \neq 0$  and  $B$  in  $K$ . Then the curve  $H$  obtained from the desingularization of the projectivization of the curve in the affine space  $\mathbb{A}^3$  given by the equations*

$$\left. \begin{aligned} y^2 &= x^3 - Ax + B \\ x^2 + xz + z^2 &= A \end{aligned} \right\}$$

*is geometrically hyperelliptic and it has two independent natural maps to  $E$ , given by  $f_1(x, y, z) = (x, y)$  and  $f_2(x, y, z) = (z, y)$ .*

**Remark.** It is a short computation to show that the abelian variety  $\text{Jac}(C)$  is isogenous to  $E^2 \times E'$ , where  $E'$  is the elliptic curve given by the Weierstrass equation  $y^2 = x^3 - 27Bx^2 + 27A^3x$ .

Next theorem is based on a similar idea of the previous theorem, but using a quartic equation instead of a cubic equation for an elliptic curve.

---

The author was partially supported by the grant MTM2009-10359.

**Theorem 2.** *Let  $K$  be a field of characteristic  $\neq 2$  and  $3$ , and take  $j \in K$ ,  $j \neq 0$  and  $1728$ . Consider  $A := \frac{3^3 j}{2^2(j-1728)}$ . Then the elliptic curve  $E$  over  $K$  given by the Weierstrass equation  $y^2 = x^3 - Ax + A$  has  $j$ -invariant  $j$  and the hyperelliptic curve  $H$  of genus 5 given by the equation*

$$y^2 = A(x+1)^4(x^2+1)^4 - 2^6 x^3(x^2+x+1)^3$$

*has two independent maps to  $E$ .*

**Proof.** The assertion on the  $j$ -invariant is easy by using the known formulae of the  $j$ -invariant. We are going to find the two maps from  $H$  to  $E$  by writing  $H$  in a different form.

First of all, we write the genus 1 curve  $E$  as the curve  $D$  with equation of the form  $y^2 = x^4 + x^3 + B$ , where  $B = \frac{A}{2^6}$ . It is standard to show that both curves are isomorphic. Now, consider the affine (singular) curve given by the equation

$$F(x, z) = \frac{(x^4 - z^4)}{(x - z)} + \frac{(x^3 - z^3)}{(x - z)} = 0$$

in the affine plane  $\mathbb{A}^2$ . This curve has genus 0 and can be parametrized by

$$x = -\frac{t^3 - 1}{t^4 - 1}, \quad z = -t \frac{t^3 - 1}{t^4 - 1}$$

By using this formulae we get that the curve  $C$  obtained from the desingularization of the projectivization of the curve in the affine space  $\mathbb{A}^3$  given by the equations

$$\left. \begin{aligned} y^2 &= x^4 + x^3 + B \\ \frac{(x^4 - z^4)}{(x - z)} + \frac{(x^3 - z^3)}{(x - z)} &= 0 \end{aligned} \right\}$$

is isomorphic to the hyperelliptic curve  $H$  given by the equation

$$y^2 = A(x+1)^4(x^2+1)^4 - 2^6 x^3(x^2+x+1)^3.$$

Now, the two independent maps from  $H$  to  $E$  are described easily as maps from  $C$  to  $D$  given by  $f_1(x, y, z) = (x, y)$  and  $f_2(x, y, z) = (z, y)$ . Both maps are clearly well defined. To show they are independent, it is sufficient to prove that there does not exist two distinct endomorphisms  $m_1$  and  $m_2$  of  $E$  as genus 1 curve such that  $m_1 \circ f_1 = m_2 \circ f_2$ . First, recall that  $f_1$  and  $f_2$  send the only point at infinity to the 0 point of  $E$ , hence  $m_1$  and  $m_2$  must be endomorphisms as elliptic curves. Moreover, both  $f_1$  and  $f_2$  have the same degree (equal to 3), hence the only possibility is that  $f_1 = -f_2$ , which is clearly not the case.  $\square$

**Remark.** It's not difficult to show that the curve  $H$  is an unramified degree 2 covering of the genus 3 curve  $H_1$  given by the equation

$$y^2 = (x^2 - 4)(A(x+2)^2 x^4 - 2^6(x+1)^3)$$

and it has an independent degree 2 covering to the genus 2 curve  $H_2$  given by the equation  $y^2 = A(x+2)^2 x^4 - 2^6(x+1)^3$ . The curve  $H_2$  has degree 2 maps to the curve  $E$  defined above, and the curve  $E'$  given by the Weierstrass equation  $y^2 = x^3 + Ax^2 + 2Ax + A$ . The curve  $H_1$  has a map to the curve  $E$ , and its kernel is simple in general (a fact that can be shown for some value

of  $A$  and modulo some prime  $p$  by computing its zeta function). Hence the abelian variety  $\text{Jac}(H)$  is isogenous to  $E^2 \times E' \times F$ , where  $F$  is an abelian surface.

For any (hyper)elliptic curve  $E$  defined over a field  $K$  given by an equation of the form  $y^2 = p(x)$ , where  $p(x) \in K[X]$ , and for any  $d \in K^*$ , we will denote by  $E_d$  the quadratic twist of  $E$ , which is the (hyper)elliptic curve given by the equation  $dy^2 = p(x)$ .

**Corollary 3.** *Let  $K$  be a field with characteristic  $\neq 2, 3$ , and let  $E$  be an elliptic curve over  $K$ . Then there exists a hyperelliptic curve  $H$  of genus  $\leq 5$  and defined over  $K$  with two independent maps from  $H$  to  $E$  defined over  $K$ .*

**Proof.** If the curve  $E$  has  $j$ -invariant equal to 0 or 1728, the result is well known (and in fact one can find such a curve with genus 2, see for example [3]).

Now, if the  $j$ -invariant is  $\neq 0$  and 1728, there exists a quadratic twist of  $E$  isomorphic to the elliptic curve given by the equation  $y^2 = x^3 - Ax + A$ , for  $A := \frac{3^3 j}{2^2(j-1728)}$ . Then the same quadratic twist of the curve  $H$  given by the previous theorem gives the answer.  $\square$

We can apply this result to give lower bounds for the number of quadratic twists with rank at least 2 for elliptic curves over  $\mathbb{Q}$ , improving slightly the results by Steward and Top in [3]. The result is a direct application of their theorem 2, but using our theorem 2 instead of Mestre's construction.

**Corollary 4.** *Let  $E$  be an elliptic curve over  $\mathbb{Q}$ . Consider, for any positive real number  $X \in \mathbb{R}$ , the function*

$$N_{\geq 2}(X) := \#\{\text{squarefree } d \in \mathbb{Z} : |d| \leq X \text{ and } \text{rank}_{\mathbb{Z}}(E_d(\mathbb{Q})) \leq 2\}.$$

*Then  $N_{\geq 2}(X) \gg X^{1/6} / \log^2(X)$ .*

#### REFERENCES

- [1] J.-F. Mestre, "Rang de courbes elliptiques d'invariant donné", C. R. Acad. Sci. Paris Sér. I Math. 314:12 (1992), 919-922.
- [2] K. Rubin, A. Silverberg, Ranks of elliptic curves. Bull. Amer. Math. Soc. (N.S.) 39 (2002), no. 4, 455-474
- [3] C. L. Stewart and J. Top, On ranks of twists of elliptic curves and power-free values of binary forms, J. Amer. Math. Soc. 8:4 (1995), 943-973.

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193  
BELLATERRA, BARCELONA, CATALONIA  
*E-mail address:* xarles@mat.uab.cat