

A 5-ENGEL ASSOCIATIVE ALGEBRA WHOSE GROUP OF UNITS IS NOT 5-ENGEL

GALINA DERYABINA AND ALEXEI KRASILNIKOV

ABSTRACT. Let R be an associative ring with unity and let $[R]$ and $U(R)$ denote the associated Lie ring (with $[a, b] = ab - ba$) and the group of units of R , respectively. In 1983 Gupta and Levin proved that if $[R]$ is a nilpotent Lie ring of class c then $U(R)$ is a nilpotent group of class at most c . The aim of the present note is to show that, in general, a similar statement does not hold if $[R]$ is n -Engel. We construct an algebra R over a field of characteristic $\neq 2, 3$ such that the Lie algebra $[R]$ is 5-Engel but the group $U(R)$ is not.

1. INTRODUCTION

Let R be an associative ring with unity and let $[R]$ and $U(R)$ denote the associated Lie ring (with $[a, b] = ab - ba$) and the group of units of R , respectively. It is known that if $[R]$ is a nilpotent Lie ring of class c then $U(R)$ is a nilpotent group of class at most c (Gupta and Levin [7]). Also, if $[R]$ is metabelian then $U(R)$ is metabelian as well (Krasilnikov [9] and Sharma and Srivastava [17]).

If R is an associative ring such that $[R]$ is centre-by-metabelian then $U(R)$, in general, is not centre-by-metabelian. For instance, if F is an infinite field of characteristic 2 and $R = M_2(F)$ is the algebra of all 2×2 matrices over F then it is well-known that $[R]$ is centre-by-metabelian but $U(R)$ does not satisfy any non-trivial identity; in particular, $U(R)$ is not centre-by-metabelian.

However, suppose that R is a unital associative algebra over a field of characteristic 0 generated (as a unital algebra) by its nilpotent elements. Then if $[R]$ is centre-by-metabelian then $U(R)$ is also centre-by metabelian (Krasilnikov and Riley [11]). Moreover, for such an algebra R , if $[R]$ satisfies an arbitrary multilinear Lie commutator identity then $U(R)$ satisfies the corresponding group commutator identity (see [11] for precise definitions); for example, if $[R]$ is solvable of length n then $U(R)$ is also solvable of length at most n .

It is natural to ask whether a similar result holds for Lie commutator identities that are not multilinear; in particular, whether it holds for the Engel identity. We show that this is, in general, not the case.

More precisely, let $[x, y] = [x, {}_{(1)}y] = xy - yx$ and let $[x, {}_{(k+1)}y] = [[x, {}_{(k)}y], y]$ for $k \geq 1$. Recall that a Lie ring L is n -Engel if $[u, {}_{(n)}v] = 0$

for all $u, v \in L$. Similarly, a group G is n -Engel if $(u,_{(n)} v) = 1$ for all $u, v \in G$, where $(x, y) = (x,_{(1)} y) = x^{-1}y^{-1}xy$ and $(x,_{(k+1)} y) = ((x,_{(k)} y), y)$ for $k \geq 1$. We are concerned with the following question.

Question. *Let F be a field of characteristic 0 and let R be a unital associative F -algebra. Suppose that the Lie algebra $[R]$ is n -Engel. Is the group $U(R)$ also n -Engel?*

If $n = 2$ then the answer to this question is “yes”. Indeed, it is well-known (see, for instance, [18, Theorem 3.1.1]) that if $[R]$ is 2-Engel and $\text{char} F \neq 3$ then $[R]$ is nilpotent of class 2. Hence, by [7], $U(R)$ is nilpotent of class at most 2 and, therefore, is 2-Engel. If $n = 3$ and the algebra R is generated (as a unital algebra) by its nilpotent elements then the answer is “yes” as well; this can be deduced from the results of [4] and [5] (see also [1]).

However, in general the answer to the question above is “no”. Our result is as follows.

Theorem 1.1. *Let F be a field of characteristic $\neq 2, 3$. Then there is a unital associative F -algebra R such that $[R]$ is a 5-Engel Lie algebra but $U(R)$ is not a 5-Engel group. This algebra R is generated (as a unital F -algebra) by 2 nilpotent elements.*

Note that if R is an associative unital ring and $[R]$ is n -Engel then $U(R)$ is m -Engel for some $m = m(n)$ (Riley and Wilson [16] and independently Amberg and Sysak [1]). If R is an algebra over a field of characteristic 0 then the existence of such $m = m(n)$ follows also from the results of Zelmanov [19] and Gupta and Levin [7]. One can check that in our example below the group $U(R)$ is 6-Engel (and nilpotent of class 7).

We obtain Theorem 1.1 as a corollary of the result below about the adjoint groups of associative algebras.

Let R be an associative ring with or without unity. It can be easily checked that R is a monoid with respect to *adjoint multiplication* defined by $u \circ v = u + v + uv$ ($u, v \in R$). The group of units of this monoid is called the *adjoint group* R° of R . It is well-known that if R is nilpotent, that is, if $R^n = \{0\}$ for some positive integer n , then $R^\circ = R$. On the other hand, if R is a ring with unity 1 then R° is isomorphic to the group of units $U(R)$ (the mapping $R^\circ \rightarrow R$ such that $a \rightarrow 1 + a$ is an isomorphism of R° onto $U(R)$).

Note that one can easily deduce from the results of [7, 9, 17] that, for an associative ring R , if the Lie ring $[R]$ is nilpotent of class c or metabelian then the adjoint group R° is also nilpotent of class at most c or metabelian, respectively. Furthermore, if $R^\circ = R$ then the converse also holds: if R° is nilpotent of class c then $[R]$ is nilpotent of class c (Du [6]) and if R° is metabelian then $[R]$ is metabelian (Amberg and Sysak [3]).

Let F be a field and let A be the free associative F -algebra without 1 on free generators x, y . Let $m(x, y), n(x, y) \in A$ be monic monomials

in x, y . If $n(x, y) = m_1(x, y)m(x, y)m_2(x, y)$ for some monic monomials $m_1(x, y), m_2(x, y) \in A \cup \{1\}$ we say that $m(x, y)$ divides $n(x, y)$ and $n(x, y)$ is a multiple of $m(x, y)$.

Let I be the ideal in A generated by the following elements:

- i) all monomials of degree 8;
- ii) all monomials of degree greater than 2 in x ;
- iii) all monic monomials of degree 7 except xyx^3xy and y^2xyxy^2 ;
- iv) all monic monomials of degree less than 7 which do not divide the monomials xyx^3xy and y^2xyxy^2 ;
- v) the polynomial $2xy^3xy - 5yxyxy^2 - 2yxy^3x + 5y^2xyxy$;
- vi) the polynomial $2yxy^3xy - 5y^2xyxy^2$.

Let $B = A/I$. It can be easily seen that $B^8 = 0$. Thus, the associative algebra B is nilpotent and, therefore, $B^\circ = B$.

Theorem 1.2. *Let F be a field of characteristic $\neq 2, 3$ and let $B = A/I$. Then $[B]$ is a 5-Engel Lie algebra but the adjoint group B° is not 5-Engel.*

To deduce Theorem 1.1 from Theorem 1.2, we embed a non-unital F -algebra B into its unital hull. Let $B_1 = F \oplus B$ be a direct sum of F -vector spaces F and B . Then B_1 has a natural associative algebra structure in which the elements of F act on B by scalar multiplication. Further, the element 1 of F is unity for B_1 and the set $1 + B$ forms a group under multiplication isomorphic to the adjoint group B° (the mapping $B^\circ \rightarrow B_1$ such that $u \rightarrow 1 + u$ is an isomorphism of B° onto $1 + B$).

It can be easily checked that, since $[B]$ is 5-Engel, so is the Lie ring $[B_1]$. On the other hand, since B° is not 5-Engel, so are the subgroup $(1 + B) \simeq B^\circ$ of $U(B_1)$ and the group $U(B_1)$. Thus, we have

Corollary 1.3. *Let F be a field of characteristic $\neq 2, 3$ and let B_1 be the unital hull of B . Then the Lie algebra $[B_1]$ is 5-Engel but the group $U(B_1)$ is not 5-Engel.*

Theorem 1.1 follows immediately from Corollary 1.3 (with $R = B_1$).

Recall that if L is a nilpotent Lie algebra over a field of characteristic 0 then L is a group with the multiplication $*$ defined by the Baker-Campbell-Hausdorff formula: $x * y = \log(e^x e^y) = x + y + \frac{1}{2}[x, y] + \dots$ (for details see, for example, [8, §9.2]). We denote this group by L^* . If L^* is an n -Engel group for some $n \geq 1$ then L is an n -Engel Lie algebra; this is well-known and can be deduced, for instance, from [8, Lemma 10.12 (d)]. However, the converse statement is false.

Indeed, if B is a nilpotent associative algebra over a field of characteristic 0 then it is well-known that $[B]^* \simeq B^\circ$ (see, for instance, [11]). Thus, by Theorem 1.2 we have

Corollary 1.4. *Let F be a field of characteristic 0 and let $B = A/I$ be the associative F -algebra defined above. Let $L = [B]$. Then L is a 5-Engel Lie algebra such that the group L^* is not 5-Engel.*

Remarks. 1. Theorems 1.1, 1.2, Corollary 1.3 and their proofs remain valid for algebras over a unital associative and commutative ring F such that $6 \neq 0$ in F .

2. One can check that if R is a nilpotent associative algebra over an infinite field and the group R° is n -Engel then the Lie algebra $[R]$ is also n -Engel.

3. For an associative ring R , the Lie ring $[R]$ is nilpotent of class c if and only if the adjoint semigroup (R, \circ) is nilpotent of class c in the sense of Mal'cev [12] or Neumann-Taylor [13]. The “only if” part of this statement has been proved independently by Krasilnikov [10] and Riley and Tasic [15] and the “if” part by Amberg and Sysak [2]. Note that if a group G , viewed as a semigroup, is Mal'cev or Neumann-Taylor nilpotent of class c then G is a nilpotent group of class c in the usual sense [12, 13].

In [14] Riley posed the following problem:

Given any positive integer n , does there exist a semigroup variety \mathcal{P}_n with the property that, for every associative ring R , the Lie ring $[R]$ is n -Engel if and only if the adjoint semigroup (R, \circ) lies in \mathcal{P}_n ?

This problem is not yet solved. Theorem 1.2 shows that, if such a variety of semigroups \mathcal{P}_n exists, the groups that belong to \mathcal{P}_n are not necessarily n -Engel.

2. PROOF OF THEOREM 1.2

Let I_0 be the two-sided ideal in A generated by the polynomials i)–iv) above. Let $C = A/I_0$. It is clear that $C = \bigoplus_{k=1}^7 C_{(k)}$ where $C_{(k)}$ is the linear span of the monomials of degree k in $x + I_0, y + I_0$.

It follows easily from the item iv) that

$$(1) \quad x^2, xy^2x, y^4, y^2xy^2, xyxy^3, y^3xyx \in I_0.$$

On the other hand, it can be easily checked that all monic monomials in x, y satisfying iv) are multiples of the monomials above. Note that every monomial in x, y of degree 1 in x and of degree 5 or 6 in y belongs to I_0 because such a monomial is a multiple of either y^4 or y^2xy^2 and the latter monomials belong to I_0 . It is straightforward to check that an F -basis of $C_{(6)}$ is formed by the images of

$$xy^3xy, yxyxy^2, yxy^3x, y^2xyxy$$

and an F -basis of $C_{(7)}$ is formed by the images of

$$yxy^3xy, y^2xyxy^2.$$

By the definition of I , I/I_0 is the ideal of C generated by the images of the polynomials

$$h_1 = 2xy^3xy - 5yxyxy^2 - 2yxy^3x + 5y^2xyxy$$

and

$$h_2 = 2yxy^3xy - 5y^2xyxy^2.$$

Note that $xh_1 \equiv h_1x \equiv 0 \pmod{I_0}$, $yh_1 \equiv -h_1y \equiv h_2 \pmod{I_0}$ and $xh_2 \equiv h_2x \equiv yh_2 \equiv h_2y \equiv 0 \pmod{I_0}$. Hence, $h_1 + I_0$ and $h_2 + I_0$ form an F -basis of the ideal I/I_0 . In particular, $C_{(7)} \cap I/I_0$ is a one-dimensional vector subspace in $C_{(7)}$ generated by the image of h_2 and $C_{(7)}/(C_{(7)} \cap I/I_0)$ is a one-dimensional vector space generated by the image of y^2xyxy^2 .

Let $a = x + I$, $b = y + I$. Then, by (1), we have

$$(2) \quad a^2 = ab^2a = b^4 = b^2ab^2 = abab^3 = b^3aba = 0.$$

It is clear that $B = \oplus_{k=1}^7 B_{(k)}$ where $B_{(k)}$ is the linear span of monomials of degree k in a, b . Note that $B_{(7)} \simeq C_{(7)}/(C_{(7)} \cap I/I_0)$ is a one-dimensional vector subspace in B generated by b^2abab^2 .

To prove that $[B]$ is 5-Engel it suffices to check that $[u, {}_{(5)}v] = 0$ for all $u, v \in B$. Let

$$u = \alpha_1 a + \beta_1 b + \gamma_1 ab + \delta_1 ba + \mu_1 b^2 + u', \quad v = \alpha_2 a + \beta_2 b + \gamma_2 ab + \delta_2 ba + \mu_2 b^2 + v',$$

where u' and v' are linear combinations of monomials in a, b of degree at least 3. Then it is straightforward to check that

$$\begin{aligned} [u, {}_{(5)}v] &= \alpha_1 [a, {}_{(5)}v] + \beta_1 [b, {}_{(5)}v] + \gamma_1 [ab, {}_{(5)}v] + \delta_1 [ba, {}_{(5)}v] + \mu_1 [b^2, {}_{(5)}v] \\ &= \alpha_1 \beta_2^5 f_0 + \alpha_1 \alpha_2 \beta_2^4 f_1 + \alpha_1 \beta_2^4 \gamma_2 f_2 + \alpha_1 \beta_2^4 \delta_2 f_3 + \alpha_1 \beta_2^4 \mu_2 f_4 + \alpha_1 \alpha_2 \beta_2^3 \mu_2 f_5 \\ &\quad + \beta_1 \alpha_2 \beta_2^4 f_6 + \beta_1 \alpha_2^2 \beta_2^3 f_7 + \beta_1 \beta_2^4 \gamma_2 f_8 + \beta_1 \alpha_2 \beta_2^3 \gamma_2 f_9 + \beta_1 \beta_2^4 \delta_2 f_{10} \\ &\quad + \beta_1 \alpha_2 \beta_2^3 \delta_2 f_{11} + \beta_1 \alpha_2 \beta_2^3 \mu_2 f_{12} + \beta_1 \alpha_2^2 \beta_2^2 \mu_2 f_{13} + \gamma_1 \beta_2^5 f_{14} \\ &\quad + \gamma_1 \alpha_2 \beta_2^4 f_{15} + \delta_1 \beta_2^5 f_{16} + \delta_1 \alpha_2 \beta_2^4 f_{17} + \mu_1 \alpha_2 \beta_2^4 f_{18} + \mu_1 \alpha_2^2 \beta_2^3 f_{19}, \end{aligned}$$

where f_i are multihomogeneous polynomials in a, b of (total) degree 6 or 7.

Recall that every monomial in x, y of degree 1 in x and of degree 5 or 6 in y belongs to I_0 . Hence, every monomial in a, b of degree 1 in a and of degree 5 or 6 in b is equal to 0. It follows immediately that $f_0 = f_4 = f_6 = f_8 = f_{10} = f_{12} = f_{14} = f_{16} = f_{18} = 0$.

It is straightforward to check that

$$f_1 = [a, b, a, b, b, b] + [a, b, b, a, b, b] + [a, b, b, b, a, b] + [a, b, b, b, b, a],$$

$$\begin{aligned} f_2 &= [a, ab, b, b, b, b] + [a, b, ab, b, b, b] + [a, b, b, ab, b, b] \\ &\quad + [a, b, b, b, ab, b] + [a, b, b, b, b, ab], \end{aligned}$$

$$\begin{aligned} f_3 &= [a, ba, b, b, b, b] + [a, b, ba, b, b, b] + [a, b, b, ba, b, b] \\ &\quad + [a, b, b, b, ba, b] + [a, b, b, b, b, ba], \end{aligned}$$

$$\begin{aligned} f_5 &= [a, b^2, a, b, b, b] + [a, b^2, b, a, b, b] + [a, b^2, b, b, a, b] + [a, b^2, b, b, b, a] \\ &\quad + [a, b, b^2, a, b, b] + [a, b, b^2, b, a, b] + [a, b, b^2, b, b, a] + [a, b, a, b^2, b, b] \\ &\quad + [a, b, b, b^2, a, b] + [a, b, b, b^2, b, a] + [a, b, a, b, b^2, b] + [a, b, b, a, b^2, b] \\ &\quad + [a, b, b, b, b^2, a] + [a, b, a, b, b, b^2] + [a, b, b, a, b, b^2] + [a, b, b, b, a, b^2], \end{aligned}$$

$$f_7 = [b, a, a, b, b, b] + [b, a, b, a, b, b] + [b, a, b, b, a, b] + [b, a, b, b, b, a] = -f_1,$$

$$f_9 = [b, a, ab, b, b, b] + [b, a, b, ab, b, b] + [b, a, b, b, ab, b] + [b, a, b, b, b, ab] \\ + [b, ab, a, b, b, b] + [b, ab, b, a, b, b] + [b, ab, b, b, a, b] + [b, ab, b, b, b, a]$$

$$f_{11} = [b, a, ba, b, b, b] + [b, a, b, ba, b, b] + [b, a, b, b, ba, b] + [b, a, b, b, b, ba] \\ + [b, ba, a, b, b, b] + [b, ba, b, a, b, b] + [b, ba, b, b, a, b] + [b, ba, b, b, b, a],$$

$$f_{13} = [b, a, b^2, a, b, b] + [b, a, b^2, b, a, b] + [b, a, b^2, b, b, a] + [b, a, a, b^2, b, b] \\ + [b, a, b, b^2, a, b] + [b, a, b, b^2, b, a] + [b, a, a, b, b^2, b] + [b, a, b, a, b^2, b] \\ + [b, a, b, b, b^2, a] + [b, a, a, b, b, b^2] + [b, a, b, a, b, b^2] + [b, a, b, b, a, b^2],$$

$$f_{15} = [ab, a, b, b, b, b] + [ab, b, a, b, b, b] + [ab, b, b, a, b, b] \\ + [ab, b, b, b, a, b] + [ab, b, b, b, b, a],$$

$$f_{17} = [ba, a, b, b, b, b] + [ba, b, a, b, b, b] + [ba, b, b, a, b, b] \\ + [ba, b, b, b, a, b] + [ba, b, b, b, b, a],$$

$$f_{19} = [b^2, a, a, b, b, b] + [b^2, a, b, a, b, b] + [b^2, a, b, b, a, b] + [b^2, a, b, b, b, a].$$

To proceed further we need the following lemma which is well-known and can be easily proved by induction.

Lemma 2.1. $[x, ({}_k y)] = \sum_{i=0}^k \binom{k}{i} (-1)^i y^i x y^{(k-i)}.$

Now we will check that $f_1 = 0$. We have

$$[a, b, a] = [ab - ba, a] = aba - ba^2 - a^2b + aba = 2aba$$

because, by (2), $a^2 = 0$. Therefore,

$$[a, b, a, b, b, b] = 2[aba, b, b, b].$$

By Lemma 2.1,

$$[aba, b, b, b] = abab^3 - 3babab^2 + 3b^2abab - b^3aba,$$

where, by (2), $abab^3 = b^3aba = 0$. Hence,

$$[aba, b, b, b] = -3babab^2 + 3b^2abab$$

and

$$(3) \quad [a, b, a, b, b, b] = -6babab^2 + 6b^2abab.$$

It is straightforward to check that $[a, b, b, a] = [a, b, a, b]$ so

$$(4) \quad [a, b, b, a, b, b] = [a, b, a, b, b, b].$$

Further, by Lemma 2.1,

$$[a, b, b, b] = ab^3 - 3bab^2 + 3b^2ab - b^3a$$

so

$$[a, b, b, b, a] = ab^3a - 3bab^2a + 3b^2aba - b^3a^2 - a^2b^3 + 3abab^2 - 3ab^2ab + ab^3a.$$

Since, by (2), $bab^2a = b^3a^2 = a^2b^3 = ab^2ab = 0$, we have

$$[a, b, b, b, a] = 2ab^3a + 3b^2aba + 3abab^2.$$

Therefore,

$$[a, b, b, b, a, b] = 2ab^3ab + 3b^2abab + 3abab^3 - 2bab^3a - 3b^3aba - 3babab^2.$$

By (2), $abab^3 = b^3aba = 0$ so

$$(5) \quad [a, b, b, b, a, b] = 2ab^3ab + 3b^2abab - 2bab^3a - 3babab^2.$$

Finally, by Lemma 2.1 and (2),

$$[a, b, b, b, b] = ab^4 - 4bab^3 + 6b^2ab^2 - 4b^3ab + b^4a = -4bab^3 - 4b^3ab$$

so, again by (2), we have

$$(6) \quad [a, b, b, b, b, a] = -4bab^3a - 4b^3aba + 4abab^3 + 4ab^3ab = -4bab^3a + 4ab^3ab.$$

Thus, by (3), (4), (5) and (6), we have

$$\begin{aligned} f_1 &= -12babab^2 + 12b^2abab + 2ab^3ab + 3b^2abab - 2bab^3a - 3babab^2 - 4bab^3a + 4ab^3ab \\ &= -15babab^2 + 15b^2abab - 6bab^3a + 6ab^3ab. \end{aligned}$$

By the item v) of the definition of the ideal I , $f_1 = 0$. Since $f_7 = -f_1$, we have $f_7 = 0$ as well.

One can check in a similar way using Lemma 2.1, (2) and the item vi) of the definition of the ideal I that $f_2 = f_3 = f_5 = f_9 = f_{11} = f_{13} = f_{15} = f_{17} = f_{19} = 0$.

More precisely, one can check using the relations (2) that

$$[b^2, a, a, b, b, b] = [b^2, a, b, a, b, b] = [b^2, a, b, b, a, b] = [b^2, a, b, b, b, a] = 0.$$

It follows that $f_{19} = 0$. Similarly, $f_5 = f_{13} = 0$ because f_5 and f_{13} are sums of certain commutators and one can check using (2) that all these commutators are equal to 0.

Further, it can be checked using (2) that $f_2 = f_3 = 0$ although the commutator summands of f_2 and f_3 are not, in general, equal to 0. Finally, one needs the relations (2) as well as the item vi) of the definition of the ideal I to check that $f_9 = f_{11} = f_{15} = f_{17} = 0$.

Thus, $[B]$ is a 5-Engel Lie ring, as required.

Now we prove that the group B° is not 5-Engel. We will check that in $U(B_1)$

$$((1+a)_{(5)}(1+b)) = 1 + 6b^2abab^2 \neq 1.$$

Hence, the subgroup $1+B$ of $U(B_1)$ is not 5-Engel. Since B° is isomorphic to $1+B$, the group B° is not 5-Engel as well.

Note that if $u \in B$ then $u^8 = 0$ so $(1+u)^{-1} = 1 - u + u^2 - \dots - u^7$. It is straightforward to check that, for all $u, v \in B$,

$$(7) \quad ((1+u), (1+v)) = 1 + [u, v] - u^2v + uvu + v^2u - vuv + w$$

where w is a linear combination of monomials of degree at least 4 in u, v . It follows that if $u = u(a, b)$ is a linear combination of some monomials of degree $k \geq 2$ in a, b and, possibly, some monomials of degree $> k$ then

$$(8) \quad ((1+u), (1+b)) = 1 + [u, b] + b^2u - bub + w' = 1 + [u, b] - b[u, b] + w'$$

where $w' \in B^{k+3}$.

By (7), we have

$$\begin{aligned} ((1+a), (1+b)) &= 1 + [a, b] + aba - a^2b + b^2a - bab + w_1 \\ &= 1 + [a, b] + aba + b^2a - bab + w_1 \end{aligned}$$

where $w_1 \in B^4$.

Let $u_1 = [a, b] + aba + b^2a - bab + w_1$; then $((1+a), (1+b)) = 1 + u_1$. By (8), we have

$$\begin{aligned} ((1+a),_{(2)} (1+b)) &= ((1+u_1), (1+b)) = 1 + [u_1, b] - b[u_1, b] + w'_2 \\ &= 1 + [a, b, b] + [(aba + b^2a - bab), b] - b[a, b, b] + w_2 \\ &= 1 + [a, b, b] + abab - baba - 2b[a, b, b] + w_2 \end{aligned}$$

where $w'_2, w_2 \in B^5$.

Similarly, one can check that

$$\begin{aligned} ((1+a),_{(3)} (1+b)) &= 1 + [a, b, b, b] + abab^2 - 2babab \\ &\quad + b^2aba - 3b[a, b, b, b] + w_3 \end{aligned}$$

where $w_3 \in B^6$,

$$((1+a),_{(4)} (1+b)) = 1 + [a, b, b, b, b] - 3babab^2 + 3b^2abab - 4b[a, b, b, b, b] + w_4$$

where $w_4 \in B^7$ and

$$((1+a),_{(5)} (1+b)) = 1 + [a, b, b, b, b, b] + 6b^2abab^2 - 5b[a, b, b, b, b, b] + w_5$$

where $w_5 = 0$ because $w_5 \in B^8$ and $B^8 = 0$ and $[a, b, b, b, b, b] = 0$ because the Lie algebra $[B]$ is 5-Engel.

Thus, $((1+a),_{(5)} (1+b)) = 1 + 6b^2abab^2$. Since $B_{(7)}$ is a one-dimensional vector subspace in B generated by b^2abab^2 , we have $b^2abab^2 \neq 0$ and $((1+a),_{(5)} (1+b)) \neq 1$. It follows that B° is not 5-Engel group, as required.

The proof of Theorem 1.2 is completed.

REFERENCES

- [1] B. Amberg and Ya.P. Sysak, *Radical rings with Engel conditions*, J. Algebra **231** (2000), 364–373.
- [2] B. Amberg and Ya. Sysak, *Associative rings whose adjoint semigroup is locally nilpotent*, Arch. Math. (Basel) **76** (2001), 426–435.
- [3] B. Amberg and Ya. Sysak, *Associative rings with metabelian adjoint group*, J. Algebra **277** (2004), 456–473.
- [4] Yu. Billig, D. Riley, V. Tasic, *Nonmatrix varieties and nil-generated algebras whose units satisfy a group identity*, J. Algebra **190** (1997), 241–252.

- [5] O. Dickenschied, *Endlichkeits- und Nilpotenzbedingungen für radikale Ringe*, dissertation, Universität Mainz, 1997.
- [6] X. Du, *The centers of a radical ring*, Canad. Math. Bull. **35** (2) (1992), 174–179.
- [7] N.D. Gupta and F. Levin, *On the Lie ideals of a ring*, J. Algebra **81** (1983), 225–231.
- [8] E. I. Khukhro, *p*-automorphisms of finite *p*-groups, London Math. Soc., Lecture Notes Series **246**, Cambridge University Press, 1998.
- [9] A.N. Krasilnikov, *On the group of units of a ring whose associated Lie ring is metabelian*, Russian Math. Surveys **47** (1992), no. 6, 214–215.
- [10] A.N. Krasil’nikov, *On the semigroup nilpotency and the Lie nilpotency of associative algebras*, Math. Notes **62** (1997), 426–433.
- [11] A.N. Krasilnikov and D.M. Riley, *The transfer of a commutator identity in a nil-generated algebra*, Int. J. Algebra Comput. **12** (2002), 437–443.
- [12] A.I. Malcev, *Nilpotent semigroups*, Ivanov. Gos. Ped. Inst. Uchen. Zap. Fiz.-Mat. Nauki **4** (1953), 107–111 (in Russian).
- [13] B.H. Neumann and T. Taylor, *Subsemigroups of nilpotent groups*, Proc. Roy. Soc. Ser. A **274** (1963), 1–4.
- [14] D.M. Riley, *Engel varieties of associative rings and the number of Mersenne primes*, J. Algebra **261** (2003), 19–30.
- [15] D.M. Riley and V. Tasic, *Malcev nilpotent algebras*, Arch. Math. (Basel) **72** (1999), 22–27.
- [16] D.M. Riley and M.C. Wilson, *Associative rings satisfying the Engel condition*, Proc. Amer. Math. Soc. **127** (1999), 973–976.
- [17] R.K. Sharma and J.B. Srivastava, *Lie centrally metabelian group rings*, J. Algebra **151** (1992), 476–486.
- [18] M.R. Vaughan-Lee. The Restricted Burnside Problem, second edition. Oxford University Press (1993).
- [19] E.I. Zel’manov. *Engel Lie algebras*, Dokl. Akad. Nauk SSSR **292** (1987), 265–268.

DEPARTMENT OF COMPUTATIONAL MATHEMATICS AND MATHEMATICAL PHYSICS (FS-11), BAUMAN MOSCOW STATE TECHNICAL UNIVERSITY, 2-ND BAUMANSKAYA STREET, 5, 105005 MOSCOW, RUSSIA

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE BRASÍLIA, 70910-900 BRASÍLIA, DF, BRAZIL
E-mail address: alexei@unb.br