A Deterministic Affine-Quadratic Optimal Control Problem^{*}

Yuanchang $Wang^{a,b}$ and Jiongmin Yong^b

^a School of Mathematics, Yunnan Normal University, Kunming, 650500, China. ^b Department of Mathematics, University of Central Florida, Orlando, FL 32816, USA.

October 5, 2018

Abstract

A Deterministic affine quadratic optimal control problem is considered. Due to the nature of the problem, optimal controls exist under some very mild conditions. Further, it is shown that under some assumptions, the value function is differentiable and therefore satisfies the corresponding Hamilton-Jacobi-Bellman equation in the classical sense. Moreover, the so-called quasi-Riccati equation is derived and any optimal control admits a state feedback representation.

Keywords. affine quadratic optimal control, dynamic programming, Hamilton-Jacobi-Bellman equation, quasi-Riccati equation, state feedback representation.

AMS Mathematics subject classification. 49J15, 49K15, 49L20, 49N10.

1 Introduction.

Consider the following controlled ordinary differential equation (ODE, for short):

(1.1)
$$\begin{cases} \dot{X}(s) = A(s, X(s)) + B(s, X(s))u(s), & s \in [t, T], \\ X(t) = x, \end{cases}$$

with cost functional

(1.2)
$$J(t,x;u(\cdot)) = \int_{t}^{T} \left[Q(s,X(s)) + \langle S(s,X(s)), u(s) \rangle + \frac{1}{2} \langle R(s,X(s))u(s), u(s) \rangle \right] ds + G(X(T)),$$

where $A : [0,T] \times \mathbb{R}^n \to \mathbb{R}^n$, $B : [0,T] \times \mathbb{R}^n \to \mathbb{R}^{n \times m}$, $Q : [0,T] \times \mathbb{R}^n \to \mathbb{R}$, $S : [0,T] \times \mathbb{R}^n \to \mathbb{R}^m$, $R : [0,T] \times \mathbb{R}^n \to \mathbb{S}^m_+$ (\mathbb{S}^m is the set of all symmetric matrices, and \mathbb{S}^m_+ is the set of all positive semi-definite matrices), and $G : [0,T] \times \mathbb{R}^n \to \mathbb{R}$ are some given maps. Let $\mathcal{U}[t,T]$ be the set of all admissible controls (which will be specified in the next section) on [t,T]. Under some mild conditions, for any $(t,x) \in [0,T) \times \mathbb{R}^n$ and $u(\cdot) \in \mathcal{U}[t,T]$, the state equation (1.1) admits a unique solution $X(\cdot) \equiv X(\cdot;t,x,u(\cdot))$ and the cost functional (1.2) is well-defined. Then we can pose the following optimal control problem.

Problem (AQ). For any given $(t, x) \in [0, T) \times \mathbb{R}^n$, find a $u^*(\cdot) \in \mathcal{U}[t, T]$ such that

(1.3)
$$J(t,x;u^*(\cdot)) = \inf_{u(\cdot)\in\mathcal{U}[t,T]} J(t,x;u(\cdot)) \equiv V(t,x).$$

^{*}The first author was supported in part by NSFC under Grant 71163046 and China State Scholarship Fund under Grant [2009]5004, the second author was supported in part by NSF under Grant DMS-1007514.

Any $u^*(\cdot)$ satisfying the above is called an *optimal control* for (t, x), and the corresponding $X^*(\cdot) \equiv X(\cdot; t, x, u^*(\cdot))$ is called an *optimal trajectory* for (t, x). The pair $(X^*(\cdot), u^*(\cdot))$ is called an *optimal pair* of Problem (AQ) for the initial pair (t, x). The function $V(\cdot, \cdot)$ is called the *value function* of Problem (AQ).

We note that the right hand side of the state equation is *affine* with respect to the control and the integrand in the cost functional is up to quadratic with respect to the control. Therefore, we call such a problem an *affine-quadratic* optimal control problem (AQ problem, for short). We see that if

$$(1.4) \qquad \begin{cases} A(t,x) = A(t)x, \quad B(t,x) = B(t), \quad Q(t,x) = \frac{1}{2} \langle Q(t)x, x \rangle, \\ S(t,x) = S(t)x, \quad R(t,x) = R(t), \quad G(x) = \frac{1}{2} \langle Gx, x \rangle, \end{cases} \qquad \forall (t,x) \in [0,T] \times \mathbb{R}^n$$

for some matrix-valued functions $A(\cdot)$, $B(\cdot)$, $Q(\cdot)$, $S(\cdot)$, $R(\cdot)$, and some matrix G, then our Problem (AQ) is reduced to a standard linear-quadratic optimal control problem (LQ problem, for short).

It is well-known that for LQ problem, under suitable conditions, one has the existence of a unique optimal control which admits a state feedback representation via the solution of a differential Riccati equation ([7], see also [11]). On the other hand, for optimal control problem of general nonlinear ordinary differential equation with a Bolza type cost functional, one generally does not expect the existence of an optimal control; However, under some mild conditions, one can characterize the value function of the optimal control problem as the unique viscosity solution to the so-called Hamilton-Jacobi-Bellman (HJB, for short) equation ([1], see also [2], [8], and the references cited therein). Note that our Problem (AQ) is between general (nonlinear) optimal control problems and LQ problems. Therefore, one expects some results "between" the results for the abovementioned two kinds of problems. A little more precisely, under certain conditions, we will have the existence of optimal controls. Further, it is possible to have state feedback representation of optimal control via a solution to the so-called quasi-Riccati equation. We would like to mention that Problem (AQ) with the state equation being linear and with the maps $x \mapsto Q(t, x)$ and $x \mapsto G(x)$ being convex, and $S(t, x) \equiv 0$ was studied in [12] and [13] by means of the quasi-Riccati equations. Also, without giving details, Problem (AQ) for stochastic differential equations was briefly discussed in [10].

Our approach is a combination of variational method and dynamic programming method. The key is to obtain, under certain hypotheses, the convexity of the map $u(\cdot) \mapsto J(t, x; u(\cdot))$ which will lead to the differentiability of the value function V(t, x). Then the Hamilton-Jacobi-Bellman (HJB, for short) equation will be satisfied in the classical sense. Furthermore, by differentiating the HJB equation we obtain a quasi-Riccati equation.

We refer to [6] and [1] for excellent surveys on the value function of optimal control theory. See also [3, 5, 2, 9] for some relevant results concerning the differentiability of value functions.

The rest of the paper is organized as follows. Section 2 collects some preliminary results. In Section 3, we present the existence of optimal controls for our Problem (AQ) and recall a Pontryagin type minimum principle. In Section 4, we derive the first and the second order variations of the cost functional with respective to the control. The invertibility of the Hessian $D_{uu}J(t, x; u(\cdot))$ of the cost functional with respect to the control variable is obtained in Section 5, under certain sufficient conditions. In Section 6, we derive the so-called quasi-Riccati equation in a very natural way, via which a state feedback representation of the optimal control is obtained. A couple of illustrative examples are presented as well. Finally, some concluding remarks are collected in Section 7.

2 Preliminaries.

Throughout this paper, we let $U \subseteq \mathbb{R}^m$ be a nonempty convex and closed set, not necessarily bounded. For convenience, we assume hereafter that $0 \in U$. Note that it could be $U = \mathbb{R}^m$. Now, we introduce the following standing assumptions.

(H1) The maps $A: [0,T] \times \mathbb{R}^n \to \mathbb{R}^n$ and $B: [0,T] \times \mathbb{R}^n \to \mathbb{R}^{n \times m}$ are continuous. There exist constants $L_A, L_B, \tilde{L}_B > 0$ such that

(2.1)
$$|A(t,x) - A(t,\bar{x})| \le L_A |x - \bar{x}|, \quad \forall t \in [0,T], \ x, \bar{x} \in \mathbb{R}^n,$$

$$(2.2) |B(t,x) - B(t,\bar{x})| \le \widetilde{L}_B |x - \bar{x}|, \forall t \in [0,T], \ x, \bar{x} \in \mathbb{R}^n,$$

and

(2.3)
$$\langle [B(t,x) - B(t,\bar{x})]^T(x-\bar{x}), u \rangle \leq L_B |x-\bar{x}|^2, \quad \forall (t,u) \in [0,T] \times U, \ x, \bar{x} \in \mathbb{R}^n.$$

Note that condition (2.3) is equivalent to the following:

(2.4)
$$\sup_{u \in U, \ x, \bar{x} \in \mathbb{R}^n, \ x \neq \bar{x}} \frac{\langle [B(t,x) - B(t,\bar{x})]^T(x-\bar{x}), u \rangle}{|x-\bar{x}|^2} \le L_B$$

On the other hand, under (2.2), the set

(2.5)
$$\mathcal{X} = \left\{ \frac{[B(t,x) - B(t,\bar{x})]^T (x - \bar{x})}{|x - \bar{x}|^2} \mid x, \bar{x} \in \mathbb{R}^n, \ x \neq \bar{x} \right\} \subseteq \mathcal{B}^m_{\tilde{L}_B}(0),$$

where $\mathcal{B}_r^m(0)$ is the ball in \mathbb{R}^m centered at 0 with radius r. Therefore, in the case U is bounded, (2.4) is satisfied with

$$L_B \ge \widetilde{L}_B \sup_{u \in U} |u|.$$

In the case $U = \mathbb{R}^m$, (2.4) is equivalent to the following:

(2.6)
$$[B(t,x) - B(t,\bar{x})]^T (x-\bar{x}) = 0, \qquad \forall t \in [0,T], \ x, \bar{x} \in \mathbb{R}^n.$$

If we denote

$$B(t,x) = \left(B^1(t,x), B^2(t,x), \cdots, B^m(t,x)\right), \quad B^i: [0,T] \times \mathbb{R}^n \to \mathbb{R}^n, \ 1 \le i \le m,$$

then (2.6) is equivalent to the following:

$$\langle B^i(t,x) - B^i(t,\bar{x}), x - \bar{x} \rangle = 0, \qquad 1 \le i \le m,$$

This is the case if $B_x^i(t,x)$ is skew symmetric, for each $1 \le i \le m$. In particular, this is the case, of course, if B(t,x) is independent of x. Note that even if B(t,x) is independent of x, due to the fact that $x \mapsto A(t,x)$ is not necessarily linear, we still have a nonlinear state equation.

Next, we introduce the following hypothesis for the functions appearing in the cost functional.

(H2) Maps $Q : [0,T] \times \mathbb{R}^n \to \mathbb{R}, S : [0,T] \times \mathbb{R}^n \to \mathbb{R}^m, R : [0,T] \times \mathbb{R}^n \to \mathbb{S}^m$, and $G : \mathbb{R}^n \to \mathbb{R}$ are continuous. There are constants $L, Q_0, G_0, S_0 > 0, \varepsilon_0 \in (0,1)$, and a continuous function $\rho : \mathbb{R}^n \to [\rho_0, \infty)$

with $\rho_0 > 0$ such that

(2.7)
$$\begin{cases} R(t,x) \ge \rho(x)I, \\ (1-\varepsilon_0)Q(t,x) - \frac{1}{2}S(t,x)^T R(t,x)^{-1}S(t,x), \ G(x) \ge -L, \\ Q(t,x) \le Q_0(1+|x|^2), \quad G(x) \le G_0(1+|x|^2), \quad |S(t,x)| \le S_0(1+|x|), \\ \forall (t,x) \in [0,T] \times \mathbb{R}^n. \end{cases}$$

(2.8)
$$\begin{cases} |A(t,x) + B(t,x)R(t,x)^{-1}S(t,x)| \le L(1+|x|), \\ |B(t,x)R(t,x)^{-1}B(t,x)^T| \le L, \end{cases} \quad (t,x) \in [0,T] \times \mathbb{R}^n \end{cases}$$

We also need the following assumption later.

(H3) The map $(t, x) \mapsto (A(t, x), B(t, x), Q(t, x), S(t, x), R(t, x), G(x))$ is twice continuously differentiable. For any $0 \le t < T$, let

(2.9)
$$\mathcal{U}[t,T] = \Big\{ u(\cdot) \in L^2(t,T;\mathbb{R}^m) \mid u(s) \in U, \quad \text{a.e. } s \in [t,T] \Big\}.$$

Any $u(\cdot) \in \mathcal{U}[t,T]$ is called an admissible control on [t,T]. We denote

$$\|u(\cdot)\|_{L^2(t,s)} = \left(\int_t^s |u(r)|^2 dr\right)^{\frac{1}{2}}, \qquad \forall u(\cdot) \in \mathcal{U}[t,s]$$

The following simple result is concerned with the well-posedness of the state equation (1.1), whose proof is straightforward.

Proposition 2.1. Let (H1) hold. Then for any initial pair $(t, x) \in [0, T] \times \mathbb{R}^n$ and $u(\cdot) \in \mathcal{U}[t, T]$, equation (1.1) admits a unique solution $X(\cdot) \equiv X(\cdot; t, x, u(\cdot))$, and the following estimate holds:

(2.10)
$$|X(s;t,x,u(\cdot))| \le K \Big[1 + |x| + ||u(\cdot)||_{L^2(t,s)} \Big], \quad \forall s \in [t,T],$$

and

(2.11)
$$|X(s;t,x,u(\cdot)) - x| \le K \Big[1 + |x| + ||u(\cdot)||_{L^2(t,s)} \Big] \Big[\sqrt{s-t} + ||u(\cdot)||_{L^2(t,s)} \Big] \sqrt{s-t},$$

hereafter, K > 0 denotes a generic constant which can be different from line to line. Further, for any $t \in [0,T], x, \bar{x} \in \mathbb{R}^n$, and $u(\cdot) \in \mathcal{U}[t,T]$, it holds

(2.12)
$$|X(s;t,x,u(\cdot)) - X(s;t,\bar{x},u(\cdot))| \le e^{(L_A + L_B)(T-t)} |x - \bar{x}|, \qquad s \in [t,T].$$

As a consequence of the above, using the technique found in [8], we have the following result on the value function.

Proposition 2.2. Let (H1)–(H2) hold. Then the value function $V(\cdot, \cdot)$ is continuous and there exists a constant K > 0 such that

(2.13)
$$-L(T-t+1) \le V(t,x) \le K(1+|x|^2), \quad \forall (t,x) \in [0,T] \times \mathbb{R}^n,$$

and

(2.14)
$$|V(t,x) - V(t,\bar{x})| \le K(|x| \lor |\bar{x}|)|x - \bar{x}|, \qquad \forall t \in [0,T], \ x, \bar{x} \in \mathbb{R}^n,$$

where $|x| \vee |\bar{x}| = \max\{|x|, |\bar{x}|\}$. Moreover, the value function $V(\cdot, \cdot)$ is the unique viscosity solution to the following HJB equation:

(2.15)
$$\begin{cases} V_t(t,x) + \langle V_x(t,x), A(t,x) \rangle + Q(t,x) \\ + \inf_{u \in U} \left[\langle B(t,x)^T V_x(t,x) + S(t,x), u \rangle + \frac{1}{2} \langle R(t,x)u, u \rangle \right] = 0, \quad (t,x) \in [0,T] \times \mathbb{R}^n, \\ V(T,x) = G(x). \end{cases}$$

Note that in the case $U = \mathbb{R}^m$, the above HJB equation can be written as

(2.16)
$$\begin{cases} V_t(t,x) + H(t,x,V_x(t,x)) = 0, & (t,x) \in [0,T] \times \mathbb{R}^n, \\ V(T,x) = G(x), & x \in \mathbb{R}^n, \end{cases}$$

with

(2.17)
$$H(t,x,p) = Q(t,x) + \langle p, A(t,x) \rangle - \frac{1}{2} \Big(B(t,x)^T p + S(t,x) \Big)^T R(t,x)^{-1} \Big(B(t,x)^T p + S(t,x) \Big), \\ (t,x,p) \in [0,T] \times \mathbb{R}^n \times \mathbb{R}^n.$$

Further, in the case that V(t, x) is differentiable, it is the classical solution to the above HJB equation and the optimal control admits the following representation:

$$u(s) = -R(s, X(s))^{-1} \Big[B(s, X(s))^T V_x(s, X(s)) + S(s, X(s)) \Big], \qquad s \in [t, T],$$

with $X(\cdot)$ being the solution to the closed-loop system:

$$\begin{cases} \dot{X}(s) = A(s, X(s)) - B(s, X(s))R(s, X(s))^{-1} \Big[B(s, X(s))^T V_x(s, X(s)) + S(s, X(s)) \Big], & s \in [t, T], \\ X(t) = x. \end{cases}$$

From [8], we note that to guarantee the uniqueness of viscosity solution to the HJB equation, we need

(2.18)
$$\begin{cases} |H(t,x,p) - H(t,y,p)| \leq \omega(|x| + |y|, |p|, |x - y|), & t \in [0,T], x, y, p \in \mathbb{R}^n, \\ |H(t,x,p) - H(t,x,q)| \leq K_0 \sum_{i=1}^k \langle x \rangle^{\lambda_i} (|p| \vee |q|)^{\nu_i} |p - q|, & t \in [0,T], x, p, q \in \mathbb{R}^n, \\ |G(x) - G(y)| \leq K_0 (\langle x \rangle \vee \langle y \rangle)^{\mu - 1} |x - y|, & \forall x, y \in \mathbb{R}^n, \\ \lambda_i, \nu_i \geq 0, & \lambda_i + (\mu - 1)\nu_i \leq 1, & 1 \leq i \leq k, \end{cases}$$

with $\langle x \rangle = \sqrt{1 + |x|^2}$. For the current case, we may let $\mu = 2$. Then

$$|G(x) - G(y)| \le L(\langle x \rangle \lor \langle y \rangle)|x - y|, \qquad \forall x, y \in \mathbb{R}^n.$$

When $U = \mathbb{R}^m$, the Hamiltonian has the explicit form (2.17). Clearly, the first condition in (2.18) holds. For the second condition, we observe that

$$|H_p(t,x)| \le |A(t,x) + B(t,x)R(t,x)^{-1}S(t,x)| + |B(t,x)R(t,x)^{-1}B(t,x)^T p| \le K_0(\langle x \rangle + |p|),$$

which is implied by (2.8). Thus, the second condition holds with

$$\lambda_1 = \nu_2 = 1, \qquad \lambda_2 = \nu_1 = 0.$$

3 Existence of Optimal Controls and Minimum Principle.

We first present the following result.

Proposition 3.1. Under (H1)–(H2), for any initial pair $(t, x) \in [0, T) \times \mathbb{R}^n$, Problem (AQ) admits an optimal control.

Proof. Let $(t,x) \in [0,T) \times \mathbb{R}^n$ be given. Let $X_0(\cdot) = X(\cdot;t,x,0)$. According to (2.10), we have

(3.1)
$$|X_0(s)| \le K(1+|x|), \quad \forall s \in [t,T],$$

for some K > 0. Let $u^k(\cdot) \in \mathcal{U}[t,T]$ be a minimizing sequence with the corresponding state trajectory $X^k(\cdot) \equiv X(\cdot;t,x,u^k(\cdot))$. Then we may assume that

$$\begin{split} J(t,x;0) &+ 1 \geq J(t,x;u^{k}(\cdot)) \\ &= \int_{t}^{T} \left[Q(s,X^{k}(s)) + \langle S(s,X^{k}(s)),u^{k}(s) \rangle + \frac{1}{2} \langle R(s,X^{k}(s))u^{k}(s),u^{k}(s) \rangle \right] ds + G(X^{k}(T)) \\ &= \int_{t}^{T} \left[\frac{1}{1-\varepsilon_{0}} \Big((1-\varepsilon_{0})Q(s,X^{k}(s)) - \frac{1}{2}S(s,X^{k}(s))^{T}R(s,X^{k}(s))^{-1}S(s,X^{k}(s)) \Big) \\ &\quad + \frac{1}{2} \big| (1-\varepsilon_{0})^{\frac{1}{2}}R(s,X^{k}(s))^{\frac{1}{2}}u^{k}(s) + (1-\varepsilon_{0})^{-\frac{1}{2}}R(s,X^{k}(s))^{-\frac{1}{2}}S(s,X^{k}(s)) \big|^{2} \\ &\quad + \frac{\varepsilon_{0}}{2} \langle R(s,X^{k}(s))u^{k}(s),u^{k}(s) \rangle \Big] ds + G(X^{k}(T)) \\ &\geq -\frac{L(T-t)}{1-\varepsilon_{0}} + \frac{\varepsilon_{0}\rho_{0}}{2} \int_{t}^{T} |u^{k}(s)|^{2} ds - L. \end{split}$$

Thus,

(3.2)
$$\int_{t}^{T} |u^{k}(s)|^{2} ds \leq K, \quad \forall k \geq 1$$

Consequently,

$$|X^{k}(s)| \le K \Big((1+|x|+\|u^{k}(\cdot)\|_{\mathcal{U}[t,T]} \Big) \le K, \quad \forall s \in [t,T], \ k \ge 1.$$

Then for any $t \leq s < \tau \leq T$,

$$|X^{k}(\tau) - X^{k}(s)| \leq \int_{s}^{\tau} \left(A_{0} + L_{A} |X^{k}(r)| + (B_{0} + \widetilde{L}_{B} |X^{k}(r))| \right) |u^{k}(r)| dr$$

$$\leq K(\tau - s) + K(\tau - s)^{\frac{1}{2}} ||u^{k}(\cdot)||_{\mathcal{U}[t,T]} \leq K(\tau - s)^{\frac{1}{2}}.$$

Thus, $\{X^k(\cdot)\}$ is uniformly bounded and equicontinuous. Hence, we may assume that $X^k(\cdot) \to X^*(\cdot)$ in $C([t,T];\mathbb{R}^n)$. Then a standard argument applies to get the existence of an optimal control (see [4]).

Now, we have the following necessary conditions for any optimal pair of Problem (AQ).

Proposition 3.2. Let (H1)–(H3) hold and $(t, x) \in [0, T) \times \mathbb{R}^n$ be given. Let $(X^*(\cdot), u^*(\cdot))$ be an optimal pair of Problem (AQ) for (t, x). Then the following adjoint equation admits a unique solution

(3.3)
$$\begin{cases} \dot{Y}(s) = -\left[A_x(s, X^*(s)) + \sum_{j=1}^m u_j^*(s)B_x^j(s, X^*(s))\right]^T Y(s) - Q_x(s, X^*(s))^T \\ -S_x(s, X^*(s))^T u^*(s) - \frac{1}{2}\sum_{j,k=1}^m u_j^*(s)u_k^*(s)R_x^{jk}(s, X^*(s))^T, \quad s \in [t, T], \\ Y(T) = G_x(X^*(T))^T, \end{cases}$$

and the following minimum condition holds:

(3.4)
$$\begin{bmatrix} B(s, X^*(s))^T Y(s) + S(s, X^*(s)) \end{bmatrix} u^*(s) + \frac{1}{2} u^*(s)^T R(s, X^*(s)) u^*(s) \\ = \min_{u \in U} \Big\{ \Big[B(s, X^*(s))^T Y(s) + S(s, X^*(s)) \Big] u + \frac{1}{2} u^T R(s, X^*(s)) u \Big\}, \quad s \in [t, T]$$

In the above, $B(s,x) = (B^1(s,x), B^2(s,x), \cdots, B^m(s,x))$ with $B^i : [0,T] \times \mathbb{R}^n \to \mathbb{R}^n$, and $B^i_x : [0,T] \times \mathbb{R}^n \to \mathbb{R}^{n \times n}$. In particular, if $U = \mathbb{R}^m$, we have

(3.5)
$$u^*(s) = -R(s, X^*(s))^{-1} \Big[B(s, X^*(s))^T Y(s) + S(s, X^*(s)) \Big], \qquad s \in [t, T].$$

From the above result, we see that under (H1)–(H3) with $U = \mathbb{R}^m$, for any $(t, x) \in [0, T) \times \mathbb{R}^n$, the following coupled two-point boundary value problem admits a solution $(X(\cdot), Y(\cdot))$:

$$(3.6) \begin{cases} \dot{X}(s) = A(s, X(s)) - B(s, X(s))R(s, X(s))^{-1} \Big[B(s, X(s))^{T}Y(s) + S(s, X(s)) \Big], \\ \dot{Y}(s) = - \Big[A_{x}(s, X(s)) - \sum_{j=1}^{m} e_{j}^{T}R(s, X(s))^{-1} \Big\{ B(s, X(s))^{T}Y(s) + S(s, X(s)) \Big\} B_{x}^{j}(s, X(s)) \Big]^{T}Y(s) \\ - Q_{x}(s, X(s))^{T} + S_{x}(s, X(s))^{T}R(s, X(s))^{-1} \Big[B(s, X(s))^{T}Y(s) + S(s, X(s)) \Big] \\ - \frac{1}{2} \sum_{j,k=1}^{m} \Big[B(s, X(s))^{T}Y(s) + S(s, X(s)) \Big]^{T}R(s, X(s))^{-1} e_{j}e_{k}^{T}R(s, X(s))^{-1} \\ \cdot \Big[B(s, X(s))^{T}Y(s) + S(s, X(s)) \Big] R_{x}^{jk}(s, X(s))^{T}, \quad s \in [t, T], \\ X(t) = x, \qquad Y(T) = G_{x}(X(T))^{T}, \end{cases}$$

where $e_j \in \mathbb{R}^m$ is the vector with entry 1 at the *i*-th position and all other entries are zero. If $(X(\cdot), Y(\cdot))$ is the unique solution to the above, then $X(\cdot) = X^*(\cdot)$ must be the optimal trajectory and the optimal control $u^*(\cdot)$ is given by (3.5).

4 Variations of the Cost Functional.

In the rest of this paper, we let $U = \mathbb{R}^m$. In this case, $\mathcal{U}[t,T]$ is a Hilbert space whose dual $\mathcal{U}[t,T]^*$ can be identified with $\mathcal{U}[t,T]$ by the Riesz representation theorem. Let us first make an observation. Define

(4.1)
$$F(t, x, u(\cdot)) = D_u J(t, x; u(\cdot)), \qquad \forall (t, x, u(\cdot)) \in [0, T] \times \mathbb{R}^n \times \mathcal{U}[0, T].$$

Then $F: [0,T] \times \mathbb{R}^n \times \mathcal{U}[0,T] \to \mathcal{U}[0,T]^* = \mathcal{U}[0,T]$. For any fixed initial pair $(t,x) \in [0,T] \times \mathbb{R}^n$, consider the following equation:

(4.2)
$$F(t, x, u(\cdot)) = 0.$$

Under (H1)–(H3), from Proposition 3.1, $u(\cdot) \mapsto J(t, x; u(\cdot))$ admits a minimum $u^*(\cdot) \equiv u^*(\cdot; t, x) \in \mathcal{U}[t, T]$, i.e.,

(4.3)
$$V(t,x) = J(t,x;u^*(\cdot)) = \inf_{u(\cdot) \in \mathcal{U}[t,T]} J(t,x;u(\cdot)).$$

Then it is necessary that $u^*(\cdot)$ is a solution to equation (4.2), and

(4.4)
$$F_u(t, x; u^*(\cdot)) = D_{uu}J(t, x; u^*(\cdot)) \ge 0.$$

Now, suppose $F_u(t, x; u^*(\cdot))^{-1} : \mathcal{U}[t, T] \to \mathcal{U}[t, T]$ exists and suppose it is a bounded operator, which, by combining (4.4), is equivalent to the following:

(4.5)
$$F_u(t,x;u^*(\cdot)) = D_{uu}J(t,x;u^*(\cdot)) \ge \delta I,$$

for some $\delta > 0$. Then, by implicit function theorem, we have that $u^*(\cdot) \equiv u^*(\cdot; t, x)$ is differentiable and

(4.6)
$$u_{(t,x)}^*(\cdot;t,x) = F_u(t,x;u^*(\cdot))^{-1}F_{(t,x)}(t,x;u^*(\cdot))$$
$$\equiv -D_{uu}J(t,x;u^*(\cdot;t,x))^{-1}[D_uJ]_{(t,x)}(t,x;u^*(\cdot;t,x)).$$

Therefore, under (H1)–(H3), as long as $D_{uu}J(t, x; u^*(\cdot; t, x))$ is uniformly positive definite, $(t, x) \mapsto u^*(\cdot; t, x)$ is differentiable, which implies that

$$V(t,x) \equiv J(t,x;u^*(\cdot;t,x))$$

is differentiable.

We now try to find conditions under which (4.5) holds. To this end, let us calculate $D_u J(t, x; u(\cdot))$ and $D_{uu} J(t, x; u(\cdot))$. Denote

$$x = \begin{pmatrix} x^1 \\ x^2 \\ \vdots \\ x^n \end{pmatrix}, y = \begin{pmatrix} y^1 \\ y^2 \\ \vdots \\ y^n \end{pmatrix} \in \mathbb{R}^n, \qquad u = \begin{pmatrix} u^1 \\ u^2 \\ \vdots \\ u^m \end{pmatrix} \in \mathbb{R}^m,$$

and

$$\begin{split} A(t,x) &= \begin{pmatrix} A^{1}(t,x) \\ A^{2}(t,x) \\ \vdots \\ A^{n}(t,x) \end{pmatrix}, \quad B(t,x) &= \begin{pmatrix} B^{11}(t,x) & B^{12}(t,x) & \cdots & B^{1m}(t,x) \\ B^{21}(t,x) & B^{22}(t,x) & \cdots & B^{2m}(t,x) \\ \vdots & \vdots & \ddots & \vdots \\ B^{n1}(t,x) & B^{n2}(t,x) & \cdots & B^{nm}(t,x) \end{pmatrix}, \\ S(t,x) &= \begin{pmatrix} S^{1}(t,x) \\ S^{2}(t,x) \\ \vdots \\ S^{m}(t,x) \end{pmatrix}, \quad R(t,x) &= \begin{pmatrix} R^{11}(t,x) & R^{12}(t,x) & \cdots & R^{1m}(t,x) \\ R^{21}(t,x) & R^{22}(t,x) & \cdots & R^{2m}(t,x) \\ \vdots & \vdots & \ddots & \vdots \\ R^{m1}(t,x) & R^{m2}(t,x) & \cdots & R^{mm}(t,x) \end{pmatrix}, \end{split}$$

 $A^i, B^{ij}, S^j, R^{jk}: [0,T]\times \mathbb{R}^n \to \mathbb{R}, \qquad 1\leq i\leq n, \ 1\leq j,k\leq m.$

Next, we denote

$$B^{j}(t,x) = \begin{pmatrix} B^{1j}(t,x) \\ B^{2j}(t,x) \\ \vdots \\ B^{nj}(t,x) \end{pmatrix}, \quad \widetilde{B}^{i}(t,x) = \begin{pmatrix} B^{i1}(t,x) \\ B^{i2}(t,x) \\ \vdots \\ B^{im}(t,x) \end{pmatrix}, \quad \forall (t,x) \in [0,T] \times \mathbb{R}^{n}, \ 1 \le j \le m, \ 1 \le i \le n.$$

Then,

$$\begin{cases} B(t,x) = (B^{1}(t,x), B^{2}(t,x), \cdots, B^{m}(t,x)), \\ B(t,x)^{T} = (\widetilde{B}^{1}(t,x), \widetilde{B}^{2}(t,x), \cdots, \widetilde{B}^{n}(t,x)), \end{cases} (t,x) \in [0,T] \times \mathbb{R}^{n}.$$

We have the following result.

Proposition 4.1. Let (H1)–(H3) hold. Then for any $(t, x) \in [0, T) \times \mathbb{R}^n$ and $u(\cdot) \in \mathcal{U}[t, T]$,

(4.7)
$$[D_u J(t, x; u(\cdot))](s) = R(s, X(s))u(s) + S(s, X(s)) + B(s, X(s))^T Y(s), \qquad s \in [t, T],$$

with $(X(\cdot), Y(\cdot))$ being the solution to the following decoupled two-point boundary value problem:

(4.8)
$$\begin{cases} \dot{X}(s) = A(s, X(s)) + B(s, X(s))u(s), \\ \dot{Y}(s) = -\left[A_x(s, X(s)) + \sum_{j=1}^m u^j(s)B_x^j(s, X(s))\right]^T Y(s) - Q_x(s, X(s))^T \\ -S_x(s, X(s))^T u(s) - \frac{1}{2}\sum_{j,k=1}^m u^j(s)u^k(s)R_x^{jk}(s, X(s))^T, \\ X(t) = x, \qquad Y(T) = G_x(X(T))^T. \end{cases}$$

Further, for any $v(\cdot) \in \mathcal{U}[t,T]$,

(4.9)
$$[D_{uu}J(t,x;u(\cdot))v(\cdot)](s) = R(s,X(s))v(s) + B(s,X(s))^T Y_1(s) + \mathbf{C}(s)X_1(s), \qquad s \in [t,T],$$

where

(4.10)
$$\begin{cases} \begin{pmatrix} \dot{X}_1(s) \\ \dot{Y}_1(s) \end{pmatrix} = \begin{pmatrix} \mathbf{A}(s) & 0 \\ -\mathbf{A}_1(s) & -\mathbf{A}(s)^T \end{pmatrix} \begin{pmatrix} X_1(s) \\ Y_1(s) \end{pmatrix} + \begin{pmatrix} B(s, X(s)) \\ -\mathbf{C}(s)^T \end{pmatrix} v(s), \quad s \in [t, T], \\ X_1(t) = 0, \quad Y_1(T) = G_{xx}(X(T))X_1(T), \end{cases}$$

with

$$(4.11) \qquad \begin{cases} \mathbf{A}(s) = A_x(s, X(s)) + \sum_{j=1}^m u^j(s) B_x^j(s, X(s)), \\ \mathbf{A}_1(s) = \sum_{i=1}^n Y^i(s) A_{xx}^i(s, X(s)) + Q_{xx}(s, X(s)) + \frac{1}{2} \sum_{j,k=1}^m u^j(s) u^k(s) R_{xx}^{jk}(s, X(s)) \\ + \sum_{j=1}^m u^j(s) \Big[\sum_{i=1}^n Y^i(s) B_{xx}^{ij}(s, X(s)) + S_{xx}^j(s, X(s)) \Big], \\ \mathbf{C}(s) = \sum_{j=1}^m u^j(s) R_x^j(s, X(s)) + S_x(s, X(s)) + \sum_{i=1}^n Y^i(s) \widetilde{B}_x^i(s, X(s)). \end{cases}$$

Proof. Let $(t,x) \in [0,T) \times \mathbb{R}^n$ be fixed and $u(\cdot), v(\cdot) \in \mathcal{U}[t,T]$, let

(4.12)
$$X(\cdot) = X(\cdot; t, x, u(\cdot)), \quad X^{\varepsilon}(\cdot) = X(\cdot; t, x, u(\cdot) + \varepsilon v(\cdot)),$$

with $\varepsilon > 0$. Let

(4.13)
$$X_1(\cdot) = \lim_{\varepsilon \to 0} \frac{X^{\varepsilon}(\cdot) - X(\cdot)}{\varepsilon}.$$

Then

$$\dot{X}_1(s) = \lim_{\varepsilon \to 0} \left\{ \frac{A(s, X^\varepsilon(s)) - A(s, X(s))}{\varepsilon} + \sum_{j=1}^m u^j(s) \frac{B^j(s, X^\varepsilon(s)) - B^j(s, X(s))}{\varepsilon} \right\} + B(s, X(s))v(s)$$
$$= \left[A_x(s, X(s)) + \sum_{j=1}^m u^j(s) B_x^j(s, X(s)) \right] X_1(s) + B(s, X(s))v(s).$$

Thus, $X_1(\cdot)$ solves the following:

$$\begin{cases} \dot{X}_1(s) = \mathbf{A}(s)X_1(s) + B(s, X(s))v(s), & s \in [t, T], \\ X_1(t) = 0, \end{cases}$$

with $\mathbf{A}(\cdot)$ being defined in (4.11). We have

$$\langle D_{u}J(t,x;u(\cdot)),v(\cdot)\rangle = \lim_{\varepsilon \to 0} \frac{J(t,x;u(\cdot) + \varepsilon v(\cdot)) - J(t,x;u(\cdot))}{\varepsilon}$$

$$= \int_{t}^{T} \left[Q_{x}(s,X(s))X_{1}(s) + \langle S(s,X(s)),v(s)\rangle + \langle S_{x}(s,X(s))X_{1}(s),u(s)\rangle \right.$$

$$+ \langle R(s,X(s))u(s),v(s)\rangle + \frac{1}{2} \sum_{j,k=1}^{m} u^{j}(s)u^{k}(s)R_{x}^{jk}(s,X(s))X_{1}(s) \right] ds + G_{x}(X(T))X_{1}(T)$$

$$= \int_{t}^{T} \left[\langle Q_{x}(s,X(s))^{T} + S_{x}(s,X(s))^{T}u(s) + \frac{1}{2} \sum_{j,k=1}^{m} u^{j}(s)u^{k}(s)R_{x}^{jk}(s,X(s))^{T},X_{1}(s)\rangle \right.$$

$$+ \langle S(s,X(s)) + R(s,X(s))u(s),v(s)\rangle \right] ds + G_{x}(X(T))X_{1}(T).$$

Let $(X(\cdot), Y(\cdot))$ be the solution to (4.8). Then (note (4.11))

$$\frac{d}{ds} \langle Y(s), X_1(s) \rangle = \langle \dot{Y}(s), X_1(s) \rangle + \langle Y(s), \mathbf{A}(s)X_1(s) \rangle + \langle Y(s), B(s, X(s))v(s) \rangle
= \langle \dot{Y}(s) + \mathbf{A}(s)^T Y(s), X_1(s) \rangle + \langle B(s, X(s))^T Y(s), v(s) \rangle.$$

Noting $X_1(t) = 0$, one has

$$G_x(X(T))X_1(T) = \langle Y(T), X_1(T) \rangle$$

= $\int_t^T \left\{ \langle \dot{Y}(s) + \mathbf{A}(s)^T Y(s), X_1(s) \rangle + \langle B(s, X(s))^T Y(s), v(s) \rangle \right\} ds.$

Consequently,

$$\langle D_{u}J(t,x;u(\cdot)),v(\cdot)\rangle = \int_{t}^{T} \Big\{ \langle \dot{Y}(s) + \mathbf{A}(s)^{T}Y(s) + Q_{x}(s,X(s))^{T} + S_{x}(s,X(s))^{T}u(s) + \frac{1}{2}\sum_{j,k=1}^{m} u^{j}(s)u^{k}(s)R_{x}^{jk}(s,X(s))^{T}, X_{1}(s)\rangle + \langle B(s,X(s))^{T}Y(s) + S(s,X(s)) + R(s,X(s))u(s),v(s)\rangle \Big\} ds$$

$$= \int_{t}^{T} \langle R(s,X(s))u(s) + S(s,X(s)) + B(s,X(s))^{T}Y(s),v(s)\rangle ds.$$

This proves (4.7).

Next, we calculate $D_{uu}J(t,x;u(\cdot))$. To this end, for any $\varepsilon \in (0,1)$, let $(X^{\varepsilon}(\cdot), Y^{\varepsilon}(\cdot))$ be the solution to the following:

$$(4.15) \qquad \begin{cases} \dot{X}^{\varepsilon}(s) = A(s, X^{\varepsilon}(s)) + B(s, X^{\varepsilon}(s)) \left[u(s) + \varepsilon v(s) \right], \\ \dot{Y}^{\varepsilon}(s) = - \left[A_x(s, X^{\varepsilon}(s)) + \sum_{j=1}^{m} [u^j(s) + \varepsilon v^j(s)] B_x^j(s, X^{\varepsilon}(s)) \right]^T Y^{\varepsilon}(s) \\ - Q_x(s, X^{\varepsilon}(s))^T - S_x(s, X^{\varepsilon}(s))^T [u(s) + \varepsilon v(s)] \\ - \frac{1}{2} \sum_{j,k=1}^{m} [u^j(s) + \varepsilon v^j(s)] [u^k(s) + \varepsilon v^k(s)] R_x^{jk}(s, X^{\varepsilon}(s))^T, \\ X^{\varepsilon}(t) = x, \qquad Y^{\varepsilon}(T) = G_x (X^{\varepsilon}(T))^T. \end{cases}$$

Then

$$[D_u J(t, x; u(\cdot) + \varepsilon v(\cdot))](s) = R(s, X^{\varepsilon}(s))[u(s) + \varepsilon v(s)] + S(s, X^{\varepsilon}(s)) + B(s, X^{\varepsilon}(s))^T Y^{\varepsilon}(s),$$

$$s \in [t, T].$$

Hence,

$$\begin{split} & [D_{uu}J(t,x;u(\cdot))v(\cdot)](s) = \lim_{\varepsilon \to 0} \frac{[D_uJ(t,x;u(\cdot) + \varepsilon v(\cdot))](s) - [D_uJ(t,x;u(\cdot))](s)}{\varepsilon} \\ & = \lim_{\varepsilon \to 0} \left\{ R(s,X^{\varepsilon}(s))v(s) + \frac{R(s,X^{\varepsilon}(s)) - R(s,X(s))}{\varepsilon} u(s) + \frac{S(x,X^{\varepsilon}(s)) - S(s,X(s))}{\varepsilon} \\ & + B(s,X^{\varepsilon}(s))^T \frac{Y^{\varepsilon}(s) - Y(s)}{\varepsilon} + \frac{B(s,X^{\varepsilon}(s))^T - B(s,X(s))^T}{\varepsilon} Y(s) \right\} \\ & = R(s,X(s))v(s) + \sum_{j=1}^m u^j(s)R_x^j(s,X(s))X_1(s) + S_x(s,X(s))X_1(s) \\ & + B(s,X(s))^TY_1(s) + \sum_{i=1}^n Y^i(s)\widetilde{B}_x^i(s,X(s))X_1(s) \\ & = R(s,X(s))v(s) + B(s,X(s))^TY_1(s) \\ & + \left[\sum_{j=1}^m u^j(s)R_x^j(s,X(s)) + S_x(s,X(s)) + \sum_{i=1}^n Y^i(s)\widetilde{B}_x^i(s,X(s))\right]X_1(s) \\ & = R(s,X(s))v(s) + B(s,X(s))^TY_1(s) + C(s)X_1(s), \end{split}$$

where

(4.16)
$$Y_1(s) = \lim_{\varepsilon \to 0} \frac{Y^{\varepsilon}(s) - Y(s)}{\varepsilon},$$

and $\mathbf{C}(\cdot)$ is defined in (4.11). Then to complete the proof, we need only to derive the equation for $Y_1(\cdot)$. First of all,

(4.17)
$$Y_1(T) = \lim_{\varepsilon \to 0} \frac{Y^{\varepsilon}(T) - Y(T)}{\varepsilon} = \lim_{\varepsilon \to 0} \frac{G_x(X^{\varepsilon}(T))^T - G_x(X(T))^T}{\varepsilon}$$
$$= G_{xx}(X(T)) \left[\lim_{\varepsilon \to 0} \frac{X^{\varepsilon}(T) - X(T)}{\varepsilon}\right] = G_{xx}(X(T))X_1(T).$$

Next,

$$\begin{split} \dot{Y}_{1}(s) &= \lim_{\varepsilon \to 0} \frac{\dot{Y}^{\varepsilon}(s) - \dot{Y}(s)}{\varepsilon} \\ &= -\lim_{\varepsilon \to 0} \Big\{ A_{x}(s, X^{\varepsilon}(s))^{T} \frac{Y^{\varepsilon}(s) - Y(s)}{\varepsilon} + \sum_{i=1}^{n} Y^{i} \frac{A_{x}^{i}(s, X^{\varepsilon}(s))^{T} - A_{x}^{i}(s, X(s))^{T}}{\varepsilon} \\ &+ \sum_{j=1}^{m} v^{j}(s) B_{x}^{j}(s, X^{\varepsilon}(s))^{T} Y^{\varepsilon}(s) + \sum_{j=1}^{m} u^{j}(s) B_{x}^{j}(s, X^{\varepsilon}(s))^{T} \frac{Y^{\varepsilon}(s) - Y(s)}{\varepsilon} \\ &+ \sum_{j=1}^{m} \sum_{i=1}^{n} u^{j}(s) \frac{B_{x}^{ij}(s, X^{\varepsilon}(s))^{T} - B_{x}^{ij}(s, X(s))^{T}}{\varepsilon} Y^{i}(s) + \frac{Q_{x}(X^{\varepsilon}(s))^{T} - Q_{x}(X(s))^{T}}{\varepsilon} \\ &+ S_{x}(s, X^{\varepsilon}(s))^{T} v(s) + \frac{S_{x}(s, X^{\varepsilon}(s))^{T} - S_{x}(s, X(s))^{T}}{\varepsilon} u(s) \\ &+ \frac{1}{2} \sum_{j,k=1}^{m} u^{j}(s) u^{k}(s) \frac{R_{x}^{jk}(s, X^{\varepsilon}(s))^{T} - R_{x}^{jk}(s, X(s))^{T}}{\varepsilon} + \sum_{j,k=1}^{m} u^{j}(s) v^{k}(s) R_{x}^{jk}(s, X^{\varepsilon}(s))^{T} \Big\}. \end{split}$$

Hence,

$$\begin{split} \dot{Y}_{1}(s) &= -A_{x}(s,X(s))^{T}Y_{1}(s) - \sum_{i=1}^{n} Y^{i}(s)A_{xx}^{i}(s,X(s))X_{1}(s) - \sum_{j=1}^{m} v^{j}(s)B_{x}^{j}(s,X(s))^{T}Y(s) \\ &- \sum_{j=1}^{m} u^{j}(s)B_{x}^{j}(s,X(s))^{T}Y_{1}(s) - \sum_{j=1}^{m} \sum_{i=1}^{n} u^{j}(s)Y^{i}(s)B_{xx}^{ij}(s,X(s))X_{1}(s) \\ &- Q_{xx}(s,X(s))X_{1}(s) - S_{x}(s,X(s))^{T}v(s) - \sum_{j=1}^{m} u^{j}(s)S_{xx}^{j}(s,X(s))X_{1}(s) \\ &- \frac{1}{2}\sum_{j,k=1}^{m} u^{j}(s)u^{k}(s)R_{xx}^{jk}(s,X(s))X_{1}(s) - \sum_{j,k=1}^{m} u^{j}(s)v^{k}(s)R_{x}^{jk}(s,X(s))^{T} \\ &= - \left[A_{x}(s,X(s)) + \sum_{j=1}^{m} u^{j}(s)B_{x}^{j}(s,X(s))\right]^{T}Y_{1}(s) \\ &- \left[\sum_{i=1}^{n} Y^{i}(s)A_{xx}^{i}(s,X(s)) + \sum_{j=1}^{m} \sum_{i=1}^{n} u^{j}(s)Y^{i}(s)B_{xx}^{ij}(s,X(s)) + Q_{xx}(s,X(s)) \right. \\ &+ \sum_{j=1}^{m} u^{j}(s)S_{xx}^{j}(s,X(s)) + \frac{1}{2}\sum_{j,k=1}^{m} u^{j}(s)u^{k}(s)R_{xx}^{jk}(s,X(s))\right]X_{1}(s) \\ &- \left[S_{x}(s,X(s)) + \sum_{i=1}^{n} Y^{i}(s)\tilde{B}_{x}^{i}(s,X(s)) + \sum_{j=1}^{m} u^{j}(s)R_{xx}^{j}(s,X(s))\right]^{T}v(s) \\ &= -\mathbf{A}(s)^{T}Y_{1}(s) - \mathbf{A}_{1}(s)X_{1}(s) - \mathbf{C}(s)^{T}v(s), \end{split}$$

where $\mathbf{A}(\cdot)$, $\mathbf{A}_1(\cdot)$, and $\mathbf{C}(\cdot)$ are given by (4.11). Thus, $(X_1(\cdot), Y_1(\cdot))$ solves (4.10).

Note that for given $(t, x) \in [0, T) \times \mathbb{R}^n$ and $u(\cdot) \in \mathcal{U}[t, T]$, both $X_1(\cdot)$ and $Y_1(\cdot)$ depend on $v(\cdot)$. It will be desirable to have a representation of $[D_{uu}J(t, x; u(\cdot))v(\cdot)]$ explicitly in terms of $v(\cdot)$. The following is such a result.

Proposition 4.2. Let (H1)–(H3) hold. For any $(t, x) \in [0, T) \times \mathbb{R}^n$ and $u(\cdot) \in \mathcal{U}[t, T]$, let $\mathbf{A}(\cdot)$, $\mathbf{A}_1(\cdot)$, and $\mathbf{C}(\cdot)$ be defined by (4.11). Then

(4.18)
$$[D_{uu}J(t,x;u(\cdot))v(\cdot)](s) = R(s,X(s))v(s) + \int_t^T \mathbf{F}(s,r)v(r)ds, \quad \forall v(\cdot) \in \mathcal{U}[t,T].$$

where

(4.19)

$$\mathbf{F}(s,r) = B(s,X(s))^{T} \Phi_{\mathbf{A}}(T,s)^{T} G_{xx}(X(T)) \Phi_{\mathbf{A}}(T,r) B(r,X(r)) + \int_{s\vee r}^{T} B(s,X(s))^{T} \Phi_{\mathbf{A}}(r',s)^{T} \mathbf{A}_{1}(r') \Phi_{\mathbf{A}}(r',r) B(r,X(r)) dr' + \mathbf{C}(s) \Phi_{\mathbf{A}}(s,r) B(r,X(r)) I_{[t,s]}(r) + B(s,X(s))^{T} \Phi_{\mathbf{A}}(r,s)^{T} \mathbf{C}(r)^{T} I_{[s,T]}(r),$$

and $\Phi_{\mathbf{A}}(\cdot, \cdot)$ is the fundamental matrix of $\mathbf{A}(\cdot)$, i.e., for any $\tau \in [t, T)$,

$$\begin{cases} \frac{d}{ds} \Phi_{\mathbf{A}}(s,\tau) = \mathbf{A}(s) \Phi_{\mathbf{A}}(s,\tau), & s \in [\tau,T], \\ \Phi_{\mathbf{A}}(\tau,\tau) = I. \end{cases}$$

Proof. Let $\Phi_{\mathbf{A}}(\cdot, \cdot)$ be the fundamental matrix of $\mathbf{A}(\cdot)$. Then

$$X_1(s) = \int_t^s \Phi_{\mathbf{A}}(s, r) B(r, X(r)) v(r) dr, \qquad s \in [t, T],$$

and

$$\begin{split} Y_1(s) &= \Phi_{\mathbf{A}}(T,s)^T G_{xx}(X(T)) X_1(T) + \int_s^T \Phi_{\mathbf{A}}(r,s)^T \Big[\mathbf{A}_1(r) X_1(r) + \mathbf{C}(r)^T v(r) \Big] dr \\ &= \Phi_{\mathbf{A}}(T,s)^T G_{xx}(X(T)) \int_t^T \Phi_{\mathbf{A}}(T,r) B(r,X(r)) v(r) dr \\ &+ \int_s^T \Phi_{\mathbf{A}}(r,s)^T \mathbf{A}_1(r) \int_t^r \Phi_{\mathbf{A}}(r,r') B(r',X(r')) v(r') dr' dr + \int_s^T \Phi_{\mathbf{A}}(r,s)^T \mathbf{C}(r)^T v(r) dr \\ &= \Phi_{\mathbf{A}}(T,s)^T G_{xx}(X(T)) \int_t^T \Phi_{\mathbf{A}}(T,r) B(r,X(r)) v(r) dr \\ &+ \int_t^T \Big[\int_{s\vee r}^T \Phi_{\mathbf{A}}(r',s)^T \mathbf{A}_1(r') \Phi_{\mathbf{A}}(r',r) dr' \Big] B(r,X(r)) v(r) dr + \int_s^T \Phi_{\mathbf{A}}(r,s)^T \mathbf{C}(r)^T v(r) dr \end{split}$$

Hence,

$$\begin{split} & \left[D_{uu}J(t,x;u(\cdot))v(\cdot)\right](s) = R(s,X(s))v(s) + B(s,X(s))^T Y_1(s) + \mathbf{C}(s)X_1(s) \\ &= R(s,X(s))v(s) + \mathbf{C}(s)\int_t^s \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr \\ &\quad + B(s,X(s))^T \Big[\Phi_{\mathbf{A}}(T,s)^T G_{xx}(X(T))\int_t^T \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \\ &\quad + \int_t^T \Big(\int_{s\vee r}^T \Phi_{\mathbf{A}}(r',s)^T \mathbf{A}_1(r')\Phi_{\mathbf{A}}(r',r)dr'\Big)B(r,X(r))v(r)dr + \int_s^T \Phi_{\mathbf{A}}(r,s)^T \mathbf{C}(r)^T v(r)dr\Big] \\ &= R(s,X(s))v(s) + \int_t^T \mathbf{F}(s,r)v(r)ds, \end{split}$$

proving (4.18).

We note that $\mathbf{F}(s, r)$ is depending on the given $u(\cdot)$ and is independent of $v(\cdot)$.

5 Invertibility of $D_{uu}J(t,x;u(\cdot))$.

Having calculated $D_{uu}J(t, x; u(\cdot))$, we now would like to look at conditions under which it admits a bounded inverse. The following is a general result whose proof is straightforward.

Proposition 5.1. Let (H1)–(H3) hold and let $(t, x) \in [0, T) \times \mathbb{R}^n$, $u(\cdot) \in \mathcal{U}[t, T]$ be given. Define $\mathbf{F}(\cdot, \cdot)$ by (4.19), with $(X(\cdot), Y(\cdot))$ being the solution to (4.8). Then $D_{uu}J(t, x; u(\cdot))$ admits a bounded inverse operator if and only if for any $w(\cdot) \in \mathcal{U}[t, T]$, the following second kind Fredholm integral equation is well-posed:

(5.1)
$$w(s) = R(s, X(s))v(s) + \int_t^T \mathbf{F}(s, r)v(r)dr, \qquad s \in [t, T].$$

A sufficient condition for the above is

(5.2)
$$|R(s, X(s))^{-1}\mathbf{F}(s, r)| \le \alpha < \frac{1}{T}, \qquad s, r \in [0, T].$$

Practically, to use the above result, we need to first solve a (decoupled) two-point boundary value problem (4.8) to get $(X(\cdot), Y(\cdot))$. Then calculate $\mathbf{A}(\cdot)$, $\mathbf{A}_1(\cdot)$ and $\mathbf{C}(\cdot)$, etc., followed by $\Phi_{\mathbf{A}}(\cdot, \cdot)$. Next, construct $\mathbf{F}(\cdot, \cdot)$ and then check see if the Fredholm integral equation (5.1) is well-posed or sufficiently look at if (5.2) holds. Apparently, some more direct sufficient conditions are desirable for $D_{uu}J(t, x; u(\cdot))$ to be invertible.

Recall from the previous section that the invertibility of $D_{uu}J(t, x; u(\cdot))$ is equivalent to the uniform positive definiteness (see (4.5)):

$$(5.3) D_{uu}J(t,x;u(\cdot)) \ge \delta I,$$

for some $\delta > 0$. Thus, we now would like to look for some sufficient conditions under which (5.3) is satisfied. To approach this, we first present the following proposition.

Proposition 5.2. Let (H1)–(H3) hold. Let $(t, x) \in [0, T) \times \mathbb{R}^n$, and $u(\cdot) \in \mathcal{U}[t, T]$ be given. Let $(X(\cdot), Y(\cdot))$ be the solution to (4.8) and $\mathbf{A}(\cdot)$, $\mathbf{A}_1(\cdot)$, and $\mathbf{C}(\cdot)$ be defined by (4.11). Then

(5.4)
$$\int_{t}^{T} \langle [D_{uu}J(t,x;u(\cdot))v(\cdot)](s),v(s) \rangle ds = \int_{t}^{T} \langle R(s,X(s))v(s),v(s) \rangle ds + \langle G_{xx}(X(T)) \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \rangle + \int_{t}^{T} \langle \mathbf{A}_{1}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr \rangle ds + 2 \int_{t}^{T} \langle \mathbf{C}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, v(s) \rangle ds.$$

Further, suppose $\bar{G} \in \mathbb{S}^n_+$ and $\bar{Q}: [0,T] \to \mathbb{S}^n_+$ such that for some $\alpha \in (0,1)$,

(5.5)
$$\begin{cases} G_{xx}(X(T)) + \bar{G} \ge 0, \\ \mathbf{A}_1(s) + \bar{Q}(s) - \alpha^{-1} \mathbf{C}(s)^T R(s, X(s))^{-1} \mathbf{C}(s) \ge 0, \quad s \in [t, T], \end{cases}$$

and

(5.6)
$$(1-\alpha)R(s,X(s)) - \left[\widehat{G}(t) + \widehat{Q}(s,t)\right]I \ge \delta I, \qquad s \in [t,T],$$

for some $\delta > 0$, with

(5.7)
$$\begin{cases} \widehat{G}(t) = \left[\int_{t}^{T} \int_{t}^{T} |B(s, X(s))^{T} \Phi_{\mathbf{A}}(T, s)^{T} \overline{G} \Phi_{\mathbf{A}}(T, r) B(r, X(r))|^{2} dr ds\right]^{\frac{1}{2}}, \\ \widehat{Q}(s, t) = \int_{s}^{T} \left[\int_{t}^{\tau} \int_{t}^{\tau} |B(r, X(r))^{T} \Phi_{\mathbf{A}}(\tau, r)^{T} \overline{Q}(\tau) \Phi_{\mathbf{A}}(\tau, r') B(r', X(r'))|^{2} dr' dr\right]^{\frac{1}{2}} d\tau, \end{cases}$$

then

(5.8)
$$D_{uu}J(t,x;u(\cdot)) \ge \delta I.$$

Proof. Let $(t,x) \in [0,T) \times \mathbb{R}^n$ and $u(\cdot) \in \mathcal{U}[t,T]$ be given. We have

$$\begin{split} &\int_{t}^{T}\int_{t}^{T} \left\langle \mathbf{F}(s,r)v(r),v(s)\right\rangle drds \\ &= \left\langle G_{xx}(X(T))\int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr,\int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr\right\rangle \\ &+ \int_{t}^{T} \left\langle \mathbf{A}_{1}(s)\int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr,\int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr\right\rangle ds \\ &+ \int_{t}^{T}\!\!\!\int_{t}^{s} \left\langle \mathbf{C}(s)\Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r),v(s)\right\rangle drds \\ &+ \int_{t}^{T}\!\!\!\int_{s}^{T} \left\langle B(s,X(s))^{T}\Phi_{\mathbf{A}}(r,s)^{T}\mathbf{C}(r)^{T}v(r),v(s)\right\rangle drds \end{split}$$

$$= \langle G_{xx}(X(T)) \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \rangle$$

+
$$\int_{t}^{T} \langle \mathbf{A}_{1}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr \rangle ds$$

+
$$2 \int_{t}^{T} \langle \mathbf{C}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, v(s) \rangle ds.$$

This proves (5.4). From this, one further has

$$\begin{split} &\int_{t}^{T} \left\langle [D_{uu}J(t,x;u(\cdot))v(\cdot)](s),v(s)\right\rangle ds = \int_{t}^{T} \left\langle R(s,X(s))v(s),v(s)\right\rangle ds \\ &+ \left\langle G_{xx}(X(T))\right\rangle \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr\right\rangle \\ &+ \int_{t}^{T} \left\langle \mathbf{A}_{1}(s)\right\rangle \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr\right\rangle ds \\ &+ 2\int_{t}^{T} \left\langle \mathbf{C}(s)\right\rangle \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, v(s)\right\rangle ds \\ &= \int_{t}^{T} \left\langle (1-\alpha)R(s,X(s))v(s),v(s)\right\rangle ds + \int_{t}^{T} \left(\left|\alpha^{\frac{1}{2}}R(s,X(s))^{\frac{1}{2}}v(s)\right|^{2}ds \\ &+ 2\left\langle \alpha^{-\frac{1}{2}}R(s,X(s))^{-\frac{1}{2}}\mathbf{C}(s)\int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, \alpha^{\frac{1}{2}}R(s,X(s))^{\frac{1}{2}}v(s)\right\rangle \\ &+ \left|\alpha^{-\frac{1}{2}}R(s,X(s))^{-\frac{1}{2}}\mathbf{C}(s)\int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr\right|^{2}\right)ds \\ &+ \left\langle G_{xx}(X(T))\int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr\right\rangle \\ &+ \int_{t}^{T} \left\langle \left[\mathbf{A}_{1}(s)-\alpha^{-1}\mathbf{C}(s)^{T}R(s,X(s))^{-1}\mathbf{C}(s)\right]\int_{t}^{s} \Phi_{A}(s,r)B(r,X(r))v(r)dr\right\rangle ds \end{split}$$

$$\begin{split} &= \int_{t}^{T} \left\langle (1-\alpha)R(s,X(s))v(s),v(s)\right\rangle ds \\ &+ \int_{t}^{T} \left| \alpha^{\frac{1}{2}}R(s,X(s))^{\frac{1}{2}}v(s) + \alpha^{-\frac{1}{2}}R(s,X(s))^{-\frac{1}{2}}\mathbf{C}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr \right|^{2} ds \\ &+ \left\langle \left[G_{xx}(X(T)) + \bar{G}\right] \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \right\rangle \\ &+ \int_{t}^{T} \left\langle \left[\mathbf{A}_{1}(s) + \bar{Q}(s) - \alpha^{-1}\mathbf{C}(s)^{T}R(s,X(s))^{-1}\mathbf{C}(s)\right] \int_{t}^{s} \Phi_{A}(s,r)B(r,X(r))v(r)dr, \\ &\qquad \int_{t}^{s} \Phi_{A}(s,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \right\rangle ds \\ &- \left\langle \bar{G} \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \right\rangle \\ &- \int_{t}^{T} \langle \bar{Q}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr \right\rangle ds \\ &\geq \int_{t}^{T} \langle (1-\alpha)R(s,X(s))v(s),v(s) \rangle ds - \left\langle \bar{G} \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \right\rangle \\ &- \int_{t}^{T} \langle \bar{Q}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr \right\rangle ds. \end{split}$$

Note that

$$\begin{split} \langle \bar{G} \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \rangle \\ &= \int_{t}^{T} \langle \int_{t}^{T} B(s,X(s))^{T} \Phi_{\mathbf{A}}(T,s)^{T} \bar{G} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, v(s) \rangle \, ds \\ &\leq \left(\int_{t}^{T} \left| \int_{t}^{T} B(s,X(s))^{T} \Phi_{\mathbf{A}}(T,s)^{T} \bar{G} \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \right|^{2} ds \right)^{\frac{1}{2}} \left(\int_{t}^{T} |v(s)|^{2} ds \right)^{\frac{1}{2}} \\ &\leq \left[\int_{t}^{T} \left(\int_{t}^{T} |B(s,X(s))^{T} \Phi_{\mathbf{A}}(T,s)^{T} \bar{G} \Phi_{\mathbf{A}}(T,r)B(r,X(r))|^{2} dr ds \right) \left(\int_{t}^{T} |v(r)|^{2} dr \right) \right]^{\frac{1}{2}} \left(\int_{t}^{T} |v(r)|^{2} dr \right)^{\frac{1}{2}} \\ &= \left[\int_{t}^{T} \int_{t}^{T} |B(s,X(s))^{T} \Phi_{\mathbf{A}}(T,s)^{T} \bar{G} \Phi_{\mathbf{A}}(T,r)B(r,X(r))|^{2} dr ds \right]^{\frac{1}{2}} \int_{t}^{T} |v(r)|^{2} dr \equiv \int_{t}^{T} \langle \hat{G}(t)v(s),v(s) \rangle \, ds, \end{split}$$

and similarly,

$$\begin{split} &\int_{t}^{T} \langle \bar{Q}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r)) \rangle v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr \rangle ds \\ &= \int_{t}^{T} \int_{t}^{s} \langle \int_{t}^{s} B(r',X(r'))^{T} \Phi_{\mathbf{A}}(s,r')\bar{Q}(s)\Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr,v(r') \rangle dr'ds \\ &\leq \int_{t}^{T} \left[\int_{t}^{s} \int_{t}^{s} |B(r,X(r))^{T} \Phi_{\mathbf{A}}(s,r)^{T}\bar{Q}(s)\Phi_{A}(s,r')B(r',X(r'))|^{2}dr'dr \right]^{\frac{1}{2}} \int_{t}^{s} |v(r)|^{2}drds \\ &= \int_{t}^{T} \left\{ \int_{\tau}^{T} \left[\int_{t}^{s} \int_{t}^{s} |B(r,X(r))^{T} \Phi_{\mathbf{A}}(t,x)^{T}\bar{Q}(s)\Phi_{\mathbf{A}}(s,r')B(r',X(r'))|^{2}dr'dr \right]^{\frac{1}{2}} ds \right\} |v(\tau)|^{2}d\tau \\ &\equiv \int_{t}^{T} \hat{Q}(s,t)|v(s)|^{2}ds, \end{split}$$

where $\widehat{G}(t)$ and $\widehat{Q}(s,t)$ are given by (5.7). Consequently,

$$\begin{split} &\int_{t}^{T} \left\langle \left[D_{uu} J(t,x;u(\cdot))v(\cdot) \right](s),v(s) \right\rangle ds \\ &\geq \int_{t}^{T} \left\langle (1-\alpha)R(s,X(s))v(s),v(s) \right\rangle ds - \left\langle \bar{G} \! \int_{t}^{T} \! \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr, \int_{t}^{T} \! \Phi_{\mathbf{A}}(T,r)B(r,X(r))v(r)dr \right\rangle \\ &\quad - \int_{t}^{T} \left\langle \bar{Q}(s) \! \int_{t}^{s} \! \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr, \int_{t}^{s} \! \Phi_{\mathbf{A}}(s,r)B(r,X(r))v(r)dr \right\rangle ds \\ &\geq \int_{t}^{T} \left\langle \left((1-\alpha)R(s,X(s)) - [\widehat{G}(t) + \widehat{Q}(s,t)]I \right)v(s),v(s) \right\rangle ds. \end{split}$$

Hence, (5.8) follows.

Let us point out that we actually do not need the invertibility of $D_{uu}J(t, x; u(\cdot))$ for all $u(\cdot) \in \mathcal{U}[t, T]$, instead, it will be enough for us to have the invertibility of $D_{uu}J(t, x; u(\cdot))$ for $u(\cdot)$ given by (see (3.5))

$$u(s) = -R(s, X(s))^{-1} \Big[B(s, X(s))^T Y(s) + S(s, X(s)) \Big], \qquad s \in [t, T],$$

with $(X(\cdot), Y(\cdot))$ being any solution to (3.6) whose existence is guaranteed by the existence of optimal controls and the Pontryagin's minimum principle.

We now look at some interesting cases.

5.1 Linear quadratic case.

Let

(5.9)
$$\begin{cases} A(t,x) = A(t)x, & B(t,x) = B(t), & Q(t,x) = \frac{1}{2} \langle Q(t)x, x \rangle \\ S(t,s) = S(t)x, & R(t,x) = R(t), & G(x) = \frac{1}{2} \langle Gx, x \rangle. \end{cases}$$

This is a classical LQ case. In this case,

$$A(s) = A(s), \quad A_1(s) = Q(s), \quad C(s) = S(s), \quad s \in [0, T].$$

Then (5.5) holds if

$$\begin{cases} G + \bar{G} \ge 0, \\ Q(s) + \bar{Q}(s) - \alpha^{-1} S(s)^T R(s)^{-1} S(s) \ge 0, \quad s \in [0, T]. \end{cases}$$

for some $\bar{G} \in \mathbb{S}^n_+$, $\bar{Q} : [0,T] \to \mathbb{S}^n_+$, and $\alpha \in (0,1)$. In this case, $\Phi_A(\cdot, \cdot)$, the fundamental matrix of $A(\cdot)$, is independent of $u(\cdot)$, $X(\cdot)$ and $Y(\cdot)$. Consequently,

are independent of $u(\cdot)$, $X(\cdot)$, and $Y(\cdot)$. Then, by Proposition 5.2, we obtain the following result.

Proposition 5.3. Let (H1)–(H3) and (5.9) hold. Suppose there exist $\alpha \in (0,1)$, and $\bar{G} \in \mathbb{S}^n_+$, $\bar{Q} : [t,T] \to \mathbb{S}^n_+$ such that

(5.11)
$$Q(s) + \bar{Q}(s) - \alpha^{-1} S(s)^T R(s)^{-1} S(s) \ge 0, \quad G + \bar{G} \ge 0, \quad s \in [t, T],$$

and

(5.12)
$$(1-\alpha)R(s) - [\widehat{G}(t) + \widehat{Q}(s,t)]I \ge \delta I, \qquad s \in [t,T],$$

for some $\delta > 0$, where $\widehat{G}(t)$ and $\widehat{Q}(s,t)$ are defined by (5.10). Then

$$D_{uu}J(t,x;u(\cdot)) \ge \delta I.$$

We point out that under the following classical conditions for LQ problems:

(5.13)
$$R(s) \ge \delta I, \quad Q(s) - S(s)^T R(s)^{-1} S(s) \ge 0, \quad G \ge 0,$$

we need only take

$$\alpha = 0, \quad \bar{G} = 0, \quad \bar{Q}(\cdot) = 0$$

Therefore, the above result covers the classical LQ problem. Further, Proposition 5.3 shows that for LQ problems, the failure of the last two conditions in (5.13) can be compensated by the sufficient positive definiteness of R(s). On the other hand, we see that due to the nature of LQ problem, the positive definiteness of $D_{uu}J(t, x; u(\cdot))$ obtained above is automatically uniform in $u(\cdot)$.

5.2 Linear semi-convex case.

Let us first assume the following:

(5.14)
$$\begin{cases} A(t,x) = A(t)x, \quad B(t,x) = B(t), \quad S(t,x) = 0, \quad R(t,x) = R(t) \ge \delta I, \\ x \mapsto Q(t,x), \quad x \mapsto G(x) \quad \text{are convex.} \end{cases}$$

In the above case, we have a linear state equation and a convex cost functional. This is a natural generalization of LQ case and we refer to it as *linear-convex* problem. Such kind of problems were carefully studied in [12, 13] by means of the so-called quasi-Riccati equation.

Note that under (5.14), it is straightforward that $u(\cdot) \mapsto J(t, x; u(\cdot))$ is uniformly convex. In our framework, one has

$$\mathbf{A}(s) = A(s), \quad \mathbf{A}_1(s) = Q_{xx}(s, X(s)), \quad \mathbf{C}(s) = 0, \qquad s \in [0, T]$$

Then

$$\begin{split} &\int_{t}^{T} \left\langle \left[D_{uu} J(t,x;u(\cdot))v(\cdot)\right](s),v(s) \right\rangle ds \\ &= \int_{t}^{T} \left\langle R(s)v(s),v(s) \right\rangle ds + \left\langle G_{xx}(X(T)) \int_{t}^{T} \Phi_{A}(T,r)B(r)v(r)dr, \int_{t}^{T} \Phi_{A}(T,r)B(r)v(r)dr \right\rangle \\ &+ \int_{t}^{T} \left\langle Q_{xx}(s,X(s)) \int_{t}^{s} \Phi_{A}(s,r)B(r)v(r)dr, \int_{t}^{s} \Phi_{A}(s,r)B(r)v(r)dr \right\rangle ds \\ &\geq \int_{t}^{T} \left\langle R(s)v(s),v(s) \right\rangle ds \geq \delta \int_{t}^{T} |v(s)|^{2} ds, \end{split}$$

proving the uniform convexity of the map $u(\cdot) \mapsto J(t, x; u(\cdot))$.

We can actually do a little bit more. Here is the result.

Proposition 5.4. Let (H1)–(H3) hold such that for some $\bar{Q}: [0,T] \to \mathbb{S}^n_+$ and $\bar{G} \in \mathbb{S}^n_+$,

(5.15)
$$A(t,x) = A(t)x, \quad B(t,x) = B(t), \quad S(t,x) = S(t)x, \quad R(t,x) = R(t),$$

and

(5.16)
$$\begin{cases} Q_{xx}(s,x) + \bar{Q}(s) - \alpha^{-1} S(s)^T R(s)^{-1} S(s) \ge 0, \quad (s,x) \in [t,T] \times \mathbb{R}^n, \\ G_{xx}(x) + \bar{G} \ge 0, \quad s \in [t,T], \end{cases}$$

and

(5.17)
$$(1-\alpha)R(s) - [\widehat{G}(t) + \widehat{Q}(s,t)]I \ge \delta I, \qquad s \in [t,T],$$

for some $\delta > 0$, where $\widehat{G}(t)$ and $\widehat{Q}(s,t)$ are defined by (5.10). Then $D_{uu}J(t,x;u(\cdot))$ is uniformly positive definite.

Proof. We note that under our conditions, one has

$$\mathbf{A}(s) = A(s), \quad \mathbf{A}_1(s) = Q_{xx}(s, X(s)), \quad \mathbf{C}(s) = S(s), \quad s \in [0, T].$$

Then by (5.16)–(5.17), we can apply Proposition 5.2 to obtain the uniform positive definiteness of $D_{uu}J(t,x;u(\cdot))$.

Recall that maps $x \mapsto Q(t, x)$ and $x \mapsto G(x)$ as semi-convex maps if there is a constant K > 0 such that

$$x \mapsto Q(t, x) + K|x|^2, \qquad x \mapsto Q(x) + K|x|^2$$

are convex. It is clear that under (5.16), $x \mapsto Q(t, x)$ and $x \mapsto G(x)$ are semi-convex. Hence, the associated problem is referred to as a *linear semi-convex* problem. Our result basically shows that the possible deviation from the convexity of the maps $x \mapsto Q(t, x)$ and $x \mapsto G(x)$ could be possibly compensated by the sufficient positive definiteness of $R(\cdot)$.

5.3 A more general case.

We now impose the following conditions:

(5.18)
$$B(t,x) = B(t), \quad R(t,x) = R(t), \quad (t,x) \in [0,T] \times \mathbb{R}^n.$$

Note that we still allow $x \mapsto (A(t,x), Q(t,x), S(t,x), G(x))$ to be nonlinear. In the current case, we have

(5.19)
$$\begin{cases} \mathbf{A}(s) = A_x(s, X(s)), \\ \mathbf{A}_1(s) = \sum_{i=1}^n Y^i(s) A^i_{xx}(s, X(s)) + Q_{xx}(s, X(s)) + \sum_{j=1}^m u^j(s) S^j_{xx}(s, X(s)), \\ \mathbf{C}(s) = S_x(s, X(s)). \end{cases}$$

Also,

$$\begin{split} &\int_{t}^{T} \left\langle \left[D_{uu} J(t,x;u(\cdot))v(\cdot)\right](s),v(s) \right\rangle ds \\ &\geq \int_{t}^{T} \left\langle (1-\alpha)R(s)v(s),v(s) \right\rangle ds - \left\langle \bar{G} \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r)v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r)v(r)dr \right\rangle \\ &- \int_{t}^{T} \left\langle \bar{Q}(s) \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr \right\rangle ds \\ &+ \left\langle (G_{xx}(X(T)) + \bar{G}) \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r)v(r)dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T,r)B(r)v(r)dr \right\rangle \\ &+ \int_{t}^{T} \left\langle \left[\mathbf{A}_{1}(s) + \bar{Q}(s) - \alpha^{-1}S_{x}(s,X(s))^{T}R(s)^{-1}S_{x}(s,X(s))\right] \right\rangle_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s,r)B(r)v(r)dr \rangle \\ &= \int_{t}^{T} \left\langle \left\{ (1-\alpha)R(s) - [\widehat{G}(t) + \widehat{Q}(s,t)]I \right\} v(s), v(s) \right\rangle ds + \mathbb{I}_{2} + \mathbb{I}_{3}, \end{split}$$

where $\widehat{G}(\cdot)$ and $\widehat{Q}(\cdot)$ are defined by the following:

(5.20)
$$\begin{cases} \widehat{G}(t) = \left[\int_{t}^{T} \int_{t}^{T} |B(s)^{T} \Phi_{\mathbf{A}}(T,s)^{T} \overline{G} \Phi_{\mathbf{A}}(T,r) B(r)|^{2} dr ds \right]^{\frac{1}{2}}, \\ \widehat{Q}(s,t) = \int_{s}^{T} \left[\int_{t}^{\tau} \int_{t}^{\tau} |B(r)^{T} \Phi_{\mathbf{A}}(\tau,r)^{T} \overline{Q}(\tau) \Phi_{\mathbf{A}}(\tau,r') B(r')|^{2} dr' dr \right]^{\frac{1}{2}} d\tau, \end{cases}$$

and

$$\begin{cases} \mathbb{I}_{2} = \langle (G_{xx}(X(T)) + \bar{G}) \int_{t}^{T} \Phi_{\mathbf{A}}(T, r) B(r) v(r) dr, \int_{t}^{T} \Phi_{\mathbf{A}}(T, r) B(r) v(r) dr \rangle \\ \mathbb{I}_{3} = \int_{t}^{T} \langle \left[\mathbf{A}_{1}(s) + \bar{Q}(s) - \alpha^{-1} S_{x}(s, X(s))^{T} R(s)^{-1} S_{x}(s, X(s)) \right] \int_{t}^{s} \Phi_{\mathbf{A}}(s, r) B(r) v(r) dr, \int_{t}^{s} \Phi_{\mathbf{A}}(s, r) B(r) v(r) dr \rangle \end{cases}$$

Naturally, we may still assume

$$G_{xx}(x) + \bar{G} \ge 0, \qquad \forall x \in \mathbb{R}^n$$

To ensure $\mathbb{I}_3 \geq 0$, we need to take a closer look at the involved terms. Note that in the current case, $\mathbf{A}_1(s)$ involves $(X(\cdot), Y(\cdot))$ and $u(\cdot)$, unless $A_{xx}^i(s, x) = 0$ and $S_{xx}^j(s, x) = 0$. Basically, we hope to get a uniform boundedness from below. On the other hand, it is known that we need the positive definiteness of $D_{uu}J(t, x; u(\cdot))$ for any candidate $u(\cdot)$ of optimal control only, rather than any admissible control. Hence, we restrict $u(\cdot)$ as follows:

$$u(s) = -R(s)^{-1} \Big[B(s)^T Y(s) + S(s, X(s)) \Big], \qquad s \in [t, T],$$

where $(X(\cdot), Y(\cdot))$ is a solution to the following:

(5.21)
$$\begin{cases} \dot{X}(s) = A(s, X(s)) - B(s)R(s)^{-1} \Big[B(s)^T Y(s) + S(s, X(s)) \Big], \\ \dot{Y}(s) = - \Big[A_x(s, X(s)) - B(s)R(s)^{-1}S_x(s, X(s)) \Big]^T Y(s) \\ - \Big[Q_x(s, X(s)) - S(s, X(s))^T R(s)^{-1}S_x(s, X(s)) \Big]^T, \quad s \in [t, T], \\ X(t) = x, \qquad Y(T) = G_x(X(T))^T. \end{cases}$$

Consequently, we have

$$\begin{aligned} \mathbf{A}_{1}(s) &= \sum_{i=1}^{n} Y^{i}(s) \Big[A^{i}_{xx}(s, X(s)) - \sum_{j=1}^{m} e^{T}_{j} R(s)^{-1} \widetilde{B}^{i}(s) S^{i}_{xx}(s, X(s)) \Big] + Q_{xx}(s, X(s)) \\ &- \sum_{j=1}^{m} \Big[e^{T}_{j} R(s)^{-1} S(s, X(s)) \Big] S^{j}_{xx}(s, X(s)). \end{aligned}$$

We now would like to explore the possibility of

$$0 \leq \mathbf{A}_{1}(s) + \bar{Q}(s) - \alpha^{-1}S_{x}(s, X(s))^{T}R(s)^{-1}S_{x}(s, X(s))$$

$$= Q_{xx}(s, X(s)) + \bar{Q}(s) - \alpha^{-1}S_{x}(s, X(s))^{T}R(s)^{-1}S_{x}(s, X(s))$$

$$- \sum_{j=1}^{m} \left[e_{j}^{T}R(s)^{-1}S(s, X(s)) \right] S_{xx}^{j}(s, X(s))$$

$$+ \sum_{i=1}^{n} Y^{i}(s) \left[A_{xx}^{i}(s, X(s)) - \sum_{j=1}^{m} e_{j}^{T}R(s)^{-1}\tilde{B}^{i}(s)S_{xx}^{i}(s, X(s)) \right],$$

for some $\bar{Q}(\cdot)$. If we are above to show that

(5.23)
$$|Y(s)| \le K_0 (1 + |X(s)|), \quad s \in [t, T],$$

then (5.22) is guaranteed by the following:

(5.24)

$$Q_{xx}(s,x) + \bar{Q}(s) - \alpha^{-1}S_x(s,x)^T R(s)^{-1}S_x(s,x) - \sum_{j=1}^m \left[e_j^T R(s)^{-1}S(s,x) \right] S_{xx}^j(s,x) + \sum_{i=1}^n K_0 \left(1 + |x| \right) \left[A_{xx}^i(s,x) - \sum_{j=1}^m e_j^T R(s)^{-1} \widetilde{B}^i(s) S_{xx}^i(s,x) \right] \ge 0, \quad \forall (s,x) \in [0,T],$$

which is practically checkable. Interestingly, in the case that $x \mapsto (A(s,x), S(s,x))$ is linear, the above is reduced to the first condition in (5.16), and (5.23) is not necessary.

Now, let us look at conditions under which (5.23) holds.

Lemma 5.5. Let (H1)-(H3) and (5.18) hold. Let

(5.25)
$$\begin{cases} |A_x(t,x) - B(t)R(t)^{-1}S_x(t,x)| \leq \bar{L}_A, & |B(t)R(t)^{-1}B(t)^T| \leq \bar{L}_B, \\ |Q_x(t,x) - S(t,x)^T R(t)^{-1}S_x(t,x)| \leq Q_0 + \bar{L}_Q |x|, & |G_x(x)| \leq |G_x(0)| + \bar{L}_G |x|, \\ \forall (t,x) \in [0,T] \times \mathbb{R}^n, \end{cases}$$

for some constants $\bar{L}_A, \bar{L}_B, \bar{L}_Q, \bar{L}_G \geq 0$. Suppose

(5.26)
$$T\left[\bar{L}_A + \bar{L}_B(\bar{L}_Q + \bar{L}_G)e^{\bar{L}_A T}\right] < 1.$$

Then there exists an absolute constant $K_0 > 0$, independent of (t, x), such that for any $(t, x) \in [0, T) \times \mathbb{R}^n$, any solution $(X(\cdot), Y(\cdot))$ of (5.21) satisfies

(5.27)
$$|Y(s)| \le K_0 \Big(1 + |X(s)| \Big), \quad s \in [t, T].$$

Proof. In what follows, we denote

$$\begin{cases} A_0 = \max_{t \in [0,T]} |A(t,0) - B(t)R(t)^{-1}S(t,0)|, \\ Q_0 = \max_{t \in [0,T]} |Q_x(t,0) - S(t,0)^T R(t)^{-1}S_x(t,0)| \end{cases}$$

For given $(t, x) \in [0, T) \times \mathbb{R}^n$, let $(X(\cdot), Y(\cdot))$ be a solution to (5.21). Let $\Psi(\cdot, \cdot)$ be the fundamental matrix of $A_x(\cdot, X(\cdot)) - B(\cdot)R(\cdot)^{-1}S_x(\cdot, X(\cdot))$, i.e.,

$$\begin{cases} \dot{\Psi}(s,\tau) = \left[A_x(s,X(s)) - B(s)R(s)^{-1}S_x(s,X(s))\right]\Psi(s,\tau), & \tau, s \in [t,T], \\ \Psi(\tau,\tau) = I, \end{cases}$$

Then by the first condition in (5.25), we have

$$|\Psi(s,\tau)| \le e^{\bar{L}_A(s-\tau)}, \qquad \forall t \le \tau \le s \le T.$$

Next, $Y(\cdot)$ admits the following representation:

$$Y(s) = \Psi(T,s)^{T}G_{x}(X(T))^{T} + \int_{s}^{T} \Psi(r,s)^{T} \Big[Q_{x}(r,X(r)) - S(r,X(r))^{T}R(r)^{-1}S_{x}(r,X(r)) \Big]^{T} dr.$$

Thus,

$$\begin{split} |Y(s)| &\leq e^{\bar{L}_A(T-s)} |G_x(0)| + \frac{Q_0(e^{\bar{L}_A(T-s)}-1)}{\bar{L}_A} + \bar{L}_G e^{\bar{L}_A(T-s)} |X(T)| + \bar{L}_Q \int_s^T e^{\bar{L}_A(r-s)} |X(r)| dr \\ &\leq K_1 + \bar{L}_G e^{\bar{L}_A T} |X(T)| + \bar{L}_Q e^{\bar{L}_A T} \int_s^T |X(r)| dr, \end{split}$$

where

$$K_1 = e^{\bar{L}_A T} |G_x(0)| + \frac{Q_0(e^{\bar{L}_A T} - 1)}{\bar{L}_A}$$

On the other hand, from

$$\begin{cases} \dot{X}(s) = A(s, X(s)) - B(s)R(s)^{-1} \Big[B(s)^T Y(s) + S(s, X(s)) \Big], & s \in [t, T], \\ X(t) = x, \end{cases}$$

for any $t \leq s \leq \tau \leq T$, we have

$$\begin{split} |X(\tau)| &\leq |X(s)| + \int_{s}^{\tau} \Big(|A(r,X(r)) - B(r)R(r)^{-1}S(r,X(r))| + |B(r)R(r)^{-1}B(r)^{T}| |Y(r)| \Big) dr \\ &\leq |X(s)| + \int_{s}^{\tau} \Big[A_{0} + \bar{L}_{A} |X(r)| + \bar{L}_{B} \Big(K_{1} + \bar{L}_{G} e^{\bar{L}_{A}T} |X(T)| + \bar{L}_{Q} e^{\bar{L}_{A}T} \int_{r}^{T} |X(r')| dr' \Big) \Big] dr \\ &\leq |X(s)| + (A_{0} + \bar{L}_{B}K_{1})(\tau - s) + \bar{L}_{A} \int_{s}^{\tau} |X(r)| dr + \bar{L}_{B}\bar{L}_{G} e^{\bar{L}_{A}T} (\tau - s)|X(T)| \\ &\quad + \bar{L}_{B}\bar{L}_{Q} \int_{s}^{\tau} e^{\bar{L}_{A}T} \int_{r}^{T} |X(r')| dr' dr \\ &\leq (A_{0} + \bar{L}_{B}K_{1})T + |X(s)| + (\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}T e^{\bar{L}_{A}T}) \int_{s}^{T} |X(r)| dr + \bar{L}_{B}\bar{L}_{G}T e^{\bar{L}_{A}T} |X(T)|. \end{split}$$

Then with $\tau = T$, we have

$$|X(T)| \le (A_0 + \bar{L}_B K_1)T + |X(s)| + (\bar{L}_A + \bar{L}_B \bar{L}_Q T e^{\bar{L}_A T}) \int_s^T |X(r)| dr + \bar{L}_B \bar{L}_G T e^{\bar{L}_A T} |X(T)|.$$

Hence, under condition (5.26), one has

$$|X(T)| \le \frac{(A_0 + \bar{L}_B K_1)T}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} + \frac{1}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} |X(s)| + \frac{\bar{L}_A + \bar{L}_B \bar{L}_Q T e^{\bar{L}_A T}}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} \int_s^T |X(r)| dr.$$

Consequently,

$$\begin{split} &\int_{s}^{T} |X(r)| dr \leq (A_{0} + \bar{L}_{B}K_{1})T^{2} + T|X(s)| + (\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}Te^{\bar{L}_{A}T})T \int_{s}^{T} |X(r)| dr + \bar{L}_{B}\bar{L}_{G}T^{2}e^{\bar{L}_{A}T}|X(T)| \\ &\leq (A_{0} + \bar{L}_{B}K_{1})T^{2} + T|X(s)| + (\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}Te^{\bar{L}_{A}T})T \int_{s}^{T} |X(r)| dr \\ &\quad + \bar{L}_{B}\bar{L}_{G}T^{2}e^{\bar{L}_{A}T} \Big[\frac{(A_{0} + \bar{L}_{B}K_{1})T}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} + \frac{1}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} |X(s)| + \frac{\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}Te^{\bar{L}_{A}T}}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} \int_{s}^{T} |X(r)| dr \Big] \\ &= (A_{0} + \bar{L}_{B}K_{1})T^{2} \Big[1 + \frac{\bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} \Big] + T \Big[1 + \frac{\bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} \Big] |X(s)| \\ &\quad + (\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}Te^{\bar{L}_{A}T})T \Big[1 + \frac{\bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} \Big] \int_{s}^{T} |X(r)| dr \\ &= \frac{(A_{0} + \bar{L}_{B}K_{1})T^{2}}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} + \frac{T}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} |X(s)| + \frac{(\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}e^{\bar{L}_{A}T})T}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}} \int_{s}^{T} |X(r)| dr. \end{split}$$

Therefore, under condition (5.26), one has

$$\int_{s}^{T} |X(r)| dr \le \frac{(A_0 + \bar{L}_B K_1) T^2 + T |X(s)|}{1 - T \left[\bar{L}_A + \bar{L}_B (\bar{L}_Q + \bar{L}_G) e^{\bar{L}_A T} \right]} \equiv K_2 (A_0 + \bar{L}_B K_1) T + K_2 |X(s)|,$$

where

$$K_{2} = \frac{T}{1 - T \left[\bar{L}_{A} + \bar{L}_{B} (\bar{L}_{Q} + \bar{L}_{G}) e^{\bar{L}_{A} T} \right]}.$$

Consequently,

$$\begin{split} |X(T)| &\leq \frac{(A_0 + \bar{L}_B K_1)T}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} + \frac{1}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} |X(s)| + \frac{\bar{L}_A + \bar{L}_B \bar{L}_Q T e^{\bar{L}_A T}}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} \int_s^T |X(r)| dr \\ &\leq \frac{(A_0 + \bar{L}_B K_1)T}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} + \frac{1}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} |X(s)| \\ &\quad + \frac{\bar{L}_A + \bar{L}_B \bar{L}_Q T e^{\bar{L}_A T}}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} \Big[K_2(A_0 + \bar{L}_B K_1)T + K_2 |X(s)| \Big] \\ &= \frac{(A_0 + \bar{L}_B K_1)T}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} \Big[1 + K_2(\bar{L}_A + \bar{L}_B \bar{L}_Q T e^{\bar{L}_A T}) \Big] + \frac{1 + K_2(\bar{L}_A + \bar{L}_B \bar{L}_Q T e^{\bar{L}_A T})}{1 - \bar{L}_B \bar{L}_G T e^{\bar{L}_A T}} |X(s)| \\ &= K_3(A_0 + \bar{L}_B K_1)T + K_3 |X(s)|, \end{split}$$

with

$$K_{3} = \frac{1 + K_{2}(\bar{L}_{A} + \bar{L}_{B}\bar{L}_{Q}Te^{\bar{L}_{A}T})}{1 - \bar{L}_{B}\bar{L}_{G}Te^{\bar{L}_{A}T}}$$

Hence,

$$\begin{aligned} |Y(s)| &\leq K_1 + \bar{L}_G e^{\bar{L}_A T} |X(T)| + \bar{L}_Q e^{\bar{L}_A T} \int_s^T |X(r)| dr \\ &\leq K_1 + \bar{L}_G e^{\bar{L}_A T} \Big[K_3(A_0 + \bar{L}_B K_1) T + K_3 |X(s)| \Big] + \bar{L}_Q e^{\bar{L}_A T} \Big[K_2(A_0 + \bar{L}_B K_1) T + K_2 |X(s)| \Big] \\ &= K_1 + e^{\bar{L}_A T} (A_0 + \bar{L}_B K_1) T (\bar{L}_G K_3 + \bar{L}_Q K_2) + e^{\bar{L}_A T} (\bar{L}_G K_3 + \bar{L}_Q K_2) |X(s)| \equiv K_4 + K_5 |X(s)|, \end{aligned}$$

with

$$\begin{cases} K_4 = K_1 + (\bar{L}_G K_3 + \bar{L}_Q K_2)(A_0 + \bar{L}_B K_1)Te^{\bar{L}_A T}, \\ K_5 = (\bar{L}_G K_3 + \bar{L}_Q K_2)e^{\bar{L}_A T}. \end{cases}$$

This proves our lemma with $K_0 = K_4 \vee K_5$.

Condition (5.26) tells us that (5.23) holds if T > 0 is not too large, and the found constant K_0 depends on all the constants $\bar{L}_A, \bar{L}_B, \bar{L}_Q, \bar{L}_G, A_0, Q_0$, as well as the time duration T. The following result, under some different conditions, shows that sometimes, T > 0 could be arbitrarily large.

Lemma 5.6. Let (H1)-(H3) and (5.18) hold. Let

$$(5.28) \begin{cases} \left[A_x(t,x) - B(t)R(t)^{-1}S_x(t,x)\right] + \left[A_x(t,x) - B(t)R(t)^{-1}S_x(t,x)\right]^T \leq -2L_0I, \\ \left|A_x(t,x) - B(t)R(t)^{-1}S_x(t,x)\right| \leq \bar{L}_A, \quad |B(t)R(t)^{-1}B(t)^T| \leq \bar{L}_B, \qquad \forall (t,x) \in [0,T] \times \mathbb{R}^n, \\ \left|Q_x(t,x) - S(t,x)^T R(t)^{-1}S_x(t,x)\right| \leq \bar{L}_Q |x|, \quad |G_x(x)| \leq \bar{L}_G |x|, \end{cases}$$

for some constants $\bar{L}_A, \bar{L}_B, \bar{L}_G, \bar{L}_Q \ge 0$ and $L_0 > 0$. Suppose

(5.29)
$$A(t,0) = S(t,0) = 0, \quad \forall t \in [0,T],$$

and

(5.30)
$$\begin{cases} 2L_0^2 - \bar{L}_B^2 \bar{L}_G^2 > 0, \qquad L_0^4 - \bar{L}_B^2 \bar{L}_Q^2 > 0, \\ 2(2L_0^2 - \bar{L}_B^2 \bar{L}_G^2)(L_0^4 - \bar{L}_B^2 \bar{L}_Q^2) - \bar{L}_B^4 \bar{L}_Q^2 \bar{L}_G^2 > 0. \end{cases}$$

For any $(t,x) \in [0,T) \times \mathbb{R}^n$, let $(X(\cdot), Y(\cdot))$ be a solution of (5.21). Then there exists an absolute constant $K_0 > 0$, independent of (t,x) and T, such that

(5.31)
$$|Y(s)| \le K_0 |X(s)|, \quad s \in [t, T].$$

Proof. For given $(t, x) \in [0, T) \times \mathbb{R}^n$, let $(X(\cdot), Y(\cdot))$ be a solution to (5.21). Then making use of the first condition in (5.28),

$$\begin{split} |Y(s)|^2 &= |G_x(X(T))|^2 + 2\int_s^T \left(\langle Y(r), \left[A_x(r, X(r)) - B(r)R(r)^{-1}S_x(r, X(r)) \right]^T Y(r) \rangle \right. \\ &+ \langle Y(r), \left[Q_x(r, X(r)) - S(r, X(r))^T R(r)^{-1}S_x(r, X(r)) \right]^T \rangle \right) dr \\ &\leq \bar{L}_G^2 |X(T)|^2 - 2L_0 \int_s^T |Y(r)|^2 dr + 2\bar{L}_Q \int_s^T |Y(r)| |X(r)| dr \\ &\leq \bar{L}_G^2 |X(T)|^2 - L_0 \int_s^T |Y(r)|^2 + \int_s^T \frac{\bar{L}_Q^2}{L_0} |X(r)|^2 dr. \end{split}$$

Hence, by Gronwall's inequality,

$$|Y(s)|^{2} \leq \bar{L}_{G}^{2} e^{-L_{0}(T-s)} |X(T)|^{2} + \frac{\bar{L}_{Q}^{2}}{L_{0}} \int_{s}^{T} e^{-L_{0}(r-s)} |X(r)|^{2} dr.$$

Next, by the first condition in (5.28) again, together with (5.29), we have

$$\langle x, A(t,x) - B(t)R(t)^{-1}S(t,x) \rangle = \langle \left(\int_0^1 \left[A_x(t,\beta x) - B(t)R(t)^{-1}S_x(t,\beta x) \right] d\beta \right) x, x \rangle \le -L_0 |x|^2.$$

Thus, from

$$\begin{cases} \dot{X}(s) = A(s, X(s)) - B(s)R(s)^{-1} \Big[B(s)^T Y(s) + S(s, X(s)) \Big], & s \in [t, T], \\ X(t) = x, \end{cases}$$

for any $t \leq s \leq \tau \leq T$, we have

$$\begin{split} |X(\tau)|^2 &= |X(s)|^2 + 2\int_s^\tau \Big(\langle X(r), A(r, X(r)) - B(r)R(r)^{-1}S(r, X(r)) \rangle \\ &- \langle X(r), B(r)R(r)^{-1}B(r)^TY(r) \rangle \Big) dr \\ &\leq |X(s)|^2 - 2\int_s^\tau \Big(L_0 |X(r)|^2 - \bar{L}_B |X(r)| |Y(r)| \Big) dr \\ &\leq |X(s)|^2 + \int_s^\tau \Big(-L_0 |X(r)|^2 + \frac{\bar{L}_B^2}{L_0} |Y(r)|^2 \Big) dr. \end{split}$$

Then, by Gronwall's inequality,

$$\begin{split} |X(\tau)|^{2} &\leq e^{-L_{0}(\tau-s)}|X(s)|^{2} + \frac{\bar{L}_{B}^{2}}{L_{0}}\int_{s}^{\tau}e^{-L_{0}(\tau-r)}|Y(r)|^{2}dr \\ &\leq e^{-L_{0}(\tau-s)}|X(s)|^{2} + \frac{\bar{L}_{B}^{2}}{L_{0}}\int_{s}^{\tau}e^{-L_{0}(\tau-r)}\Big(\bar{L}_{G}^{2}e^{-L_{0}(T-r)}|X(T)|^{2} + \frac{\bar{L}_{Q}^{2}}{L_{0}}\int_{r}^{T}e^{-L_{0}(r'-r)}|X(r')|^{2}dr'\Big)dr \\ &= e^{-L_{0}(\tau-s)}|X(s)|^{2} + \frac{\bar{L}_{B}^{2}\bar{L}_{G}^{2}(e^{-L_{0}(T-\tau)} - e^{-L_{0}(T+\tau-2s)})}{2L_{0}^{2}}|X(T)|^{2} \\ (5.32) \qquad &+ \frac{\bar{L}_{B}^{2}\bar{L}_{Q}^{2}}{L_{0}^{2}}\int_{s}^{\tau}e^{-L_{0}(\tau-r)}\Big(\int_{r}^{T}e^{-L_{0}(r'-r)}|X(r')|^{2}dr'\Big)dr \\ &= e^{-L_{0}(\tau-s)}|X(s)|^{2} + \frac{\bar{L}_{B}^{2}\bar{L}_{G}^{2}(e^{-L_{0}(T-\tau)} - e^{-L_{0}(T+\tau-2s)})}{2L_{0}^{2}}|X(T)|^{2} \\ &+ \frac{\bar{L}_{B}^{2}\bar{L}_{Q}^{2}}{2L_{0}^{3}}\int_{s}^{T}(e^{-L_{0}[\tau+r-2(\tau\wedge r)]} - e^{-L_{0}(\tau+r-2s)})|X(r)|^{2}dr \\ &\leq e^{-L_{0}(\tau-s)}|X(s)|^{2} + \frac{\bar{L}_{B}^{2}\bar{L}_{G}^{2}e^{-L_{0}(T-\tau)}}{2L_{0}^{2}}|X(T)|^{2} + \frac{\bar{L}_{B}^{2}\bar{L}_{Q}^{2}}{2L_{0}^{3}}\int_{s}^{T}e^{-L_{0}[\tau+r-2(\tau\wedge r)]}|X(r)|^{2}dr. \end{split}$$

Integrating the above over [s, T], we obtain

$$\begin{split} \int_{s}^{T} |X(\tau)|^{2} d\tau &\leq \int_{s}^{T} \left[e^{-L_{0}(\tau-s)} |X(s)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{G}^{2} e^{-L_{0}(T-\tau)}}{2L_{0}^{2}} |X(T)|^{2} \\ &\quad + \frac{\bar{L}_{B}^{2} \bar{L}_{Q}^{2}}{2L_{0}^{3}} \int_{s}^{T} e^{-L_{0}[\tau+r-2(\tau\wedge r)]} |X(r)|^{2} dr \right] d\tau \\ &\leq \frac{1-e^{L_{0}(T-s)}}{L_{0}} |X(s)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{G}^{2} (1-e^{-L_{0}(T-s)})}{2L_{0}^{3}} |X(T)|^{2} \\ &\quad + \frac{\bar{L}_{B}^{2} \bar{L}_{Q}^{2}}{2L_{0}^{3}} \int_{s}^{T} \Big(\int_{s}^{\tau} e^{-L_{0}(\tau-r)} |X(r)|^{2} dr + \int_{\tau}^{T} e^{-L_{0}(\tau-\tau)} |X(r)|^{2} dr \Big) d\tau \\ &\leq \frac{1}{L_{0}} |X(s)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{G}^{2}}{2L_{0}^{3}} |X(T)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{Q}^{2}}{2L_{0}^{3}} \Big[\int_{s}^{T} \Big(\int_{s}^{\tau} e^{-L_{0}(\tau-\tau)} d\tau \Big) |X(r)|^{2} dr \\ &\quad + \int_{s}^{T} \Big(\int_{s}^{\tau} e^{-L_{0}(\tau-\tau)} d\tau \Big) |X(r)|^{2} dr \Big] \\ &= \frac{1}{L_{0}} |X(s)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{G}^{2}}{2L_{0}^{3}} |X(T)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{Q}^{2}}{2L_{0}^{4}} \Big[\int_{s}^{T} \Big((1-e^{-L_{0}(\tau-\tau)} + (1-e^{-L_{0}(\tau-s)}) \Big) |X(r)|^{2} dr \Big] \\ &\leq \frac{1}{L_{0}} |X(s)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{G}^{2}}{2L_{0}^{3}} |X(T)|^{2} + \frac{\bar{L}_{B}^{2} \bar{L}_{Q}^{2}}{L_{0}^{4}} \int_{s}^{T} |X(r)|^{2} dr. \end{split}$$

Thus, under condition (5.30), one has

$$\int_{s}^{T} |X(r)|^{2} dr \leq \frac{L_{0}^{3}}{L_{0}^{4} - \bar{L}_{B}^{2} \bar{B}_{Q}^{2}} |X(s)|^{2} + \frac{L_{0} \bar{L}_{B}^{2} \bar{L}_{G}^{2}}{2(L_{0}^{4} - \bar{L}_{B}^{2} \bar{L}_{Q}^{2})} |X(T)|^{2}.$$

On the other hand, taking $\tau = T$ in (5.32), we get

$$|X(T)|^{2} \leq e^{-L_{0}(T-s)}|X(s)|^{2} + \frac{\bar{L}_{B}^{2}\bar{L}_{G}^{2}}{2L_{0}^{2}}|X(T)|^{2} + \frac{\bar{L}_{B}^{2}\bar{L}_{Q}^{2}}{2L_{0}^{3}}\int_{s}^{T}e^{-L_{0}(T-r)}|X(r)|^{2}dr.$$

Thus, by (5.30),

$$\begin{split} |X(T)|^2 &\leq \frac{2L_0^2 e^{-L_0(T-s)}}{2L_0^2 - \bar{L}_B^2 \bar{L}_G^2} |X(s)|^2 + \frac{\bar{L}_B^2 \bar{L}_Q^2}{L_0(2L_0^2 - \bar{L}_B^2 \bar{L}_G^2)} \int_s^T e^{-L_0(T-r)} |X(r)|^2 dr \\ &\leq \frac{2L_0^2}{2L_0^2 - \bar{L}_B^2 \bar{L}_G^2} |X(s)|^2 + \frac{\bar{L}_B^2 \bar{L}_Q^2}{L_0(2L_0^2 - \bar{L}_B^2 \bar{L}_G^2)} \int_s^T |X(r)|^2 dr \\ &\leq \frac{2L_0^2}{2L_0^2 - \bar{L}_B^2 \bar{L}_G^2} |X(s)|^2 + \frac{\bar{L}_B^2 \bar{L}_Q^2}{L_0(2L_0^2 - \bar{L}_B^2 \bar{L}_G^2)} \Big[\frac{L_0^3}{L_0^4 - \bar{L}_B^2 \bar{B}_Q^2} |X(s)|^2 + \frac{L_0 \bar{L}_B^2 \bar{L}_G^2}{2L_0^2 - \bar{L}_B^2 \bar{L}_G^2} |X(s)|^2 + \frac{L_0 \bar{L}_B^2 \bar{L}_G^2}{L_0^4 - \bar{L}_B^2 \bar{L}_Q^2} |X(r)|^2 \Big] \\ &= \Big[\frac{2L_0^2}{2L_0^2 - \bar{L}_B^2 \bar{L}_G^2} + \frac{L_0^2 \bar{L}_B^2 \bar{L}_Q^2}{(2L_0^2 - \bar{L}_B^2 \bar{L}_Q^2)} \Big] |X(s)|^2 + \frac{\bar{L}_B^4 \bar{L}_Q^2 \bar{L}_G^2}{2(L_0^4 - \bar{L}_B^2 \bar{L}_Q^2)} |X(T)|^2. \end{split}$$

Hence, by (5.30), we obtain

$$|X(T)|^{2} \leq \frac{4L_{0}^{2}(L_{0}^{4} - \bar{L}_{B}^{2}\bar{L}_{Q}^{2}) + 2L_{0}^{2}\bar{L}_{B}^{2}\bar{L}_{Q}^{2}}{2(L_{0}^{4} - \bar{L}_{B}^{2}\bar{L}_{Q}^{2})(2L_{0}^{2} - \bar{L}_{B}^{2}\bar{L}_{G}^{2}) - \bar{L}_{B}^{4}\bar{L}_{Q}^{2}\bar{L}_{G}^{2}}|X(s)|^{2},$$

and

$$\int_{s}^{T} |X(r)|^{2} dr \leq \Big[\frac{L_{0}^{3}}{L_{0}^{4} - \bar{L}_{B}^{2} \bar{B}_{Q}^{2}} + \frac{4L_{0}^{2} (L_{0}^{4} - \bar{L}_{B}^{2} \bar{L}_{Q}^{2}) + 2L_{0}^{2} \bar{L}_{B}^{2} \bar{L}_{Q}^{2}}{2(L_{0}^{4} - \bar{L}_{B}^{2} \bar{L}_{Q}^{2})(2L_{0}^{2} - \bar{L}_{B}^{2} \bar{L}_{G}^{2}) - \bar{L}_{B}^{4} \bar{L}_{Q}^{2} \bar{L}_{G}^{2}} \Big] |X(s)|^{2}.$$

Combining the above, we finally get

$$\begin{split} |Y(s)|^2 &\leq \bar{L}_G^2 |X(T)|^2 + \frac{\bar{L}_Q^2}{L_0} \int_s^T |X(r)|^2 dr \\ &\leq \Big[\frac{\bar{L}_G^2 [4L_0^2(L_0^4 - \bar{L}_B^2 \bar{L}_Q^2) + 2L_0^2 \bar{L}_B^2 \bar{L}_Q^2]}{2(L_0^4 - \bar{L}_B^2 \bar{L}_Q^2)(2L_0^2 - \bar{L}_B^2 \bar{L}_Q^2) - \bar{L}_B^4 \bar{L}_Q^2 \bar{L}_G^2} \\ &\quad + \frac{L_0^2 \bar{L}_Q^2}{L_0^4 - \bar{L}_B^2 \bar{L}_Q^2} + \frac{4L_0 \bar{L}_Q^2 (L_0^4 - \bar{L}_B^2 \bar{L}_Q^2) + 2L_0 \bar{L}_B^2 \bar{L}_Q^4}{2(L_0^4 - \bar{L}_B^2 \bar{L}_Q^2)(2L_0^2 - \bar{L}_B^2 \bar{L}_Q^2) - \bar{L}_B^4 \bar{L}_Q^2 \bar{L}_G^2} \Big] |X(s)|^2 \\ &= \Big[\frac{L_0^2 \bar{L}_Q^2}{L_0^4 - \bar{L}_B^2 \bar{L}_Q^2} + \frac{(\bar{L}_G^2 + L_0 \bar{L}_Q^2) [4L_0^2 (L_0^4 - \bar{L}_B^2 \bar{L}_Q^2) + 2L_0^2 \bar{L}_B^2 \bar{L}_Q^2]}{2(L_0^4 - \bar{L}_B^2 \bar{L}_Q^2)(2L_0^2 - \bar{L}_B^2 \bar{L}_Q^2) - \bar{L}_B^4 \bar{L}_Q^2 \bar{L}_Q^2} \Big] |X(s)|^2. \end{split}$$

Then our conclusion follows.

If (5.29) is not assumed and the last line in (5.28) is replaced by the second line of (5.25), then, with more complicated-looking estimates, we will have

$$|Y(s)| \le K_0(1+|X(s)|), \quad s \in [t,T],$$

instead of (5.31), with K_0 also independent of T. For the simplicity of presentation, we prefer not to give the details here. Having the above two lemmas, we may state the following result whose proof is clear.

Theorem 5.7. Let conditions of Lemma 5.5 or Lemma 5.6 hold. Let $\bar{G} \in \mathbb{S}^n$ and $\bar{Q} : [0,T] \to \mathbb{S}^n$ such that for some $\alpha \in (0,1)$ and $\delta > 0$,

(5.33)
$$G_{xx}(x) + \bar{G} \ge 0, \qquad x \in \mathbb{R}^n,$$

(5.34)
$$R(s) - [\widehat{G}(t) + \widehat{Q}(s)]I \ge \delta I, \qquad 0 \le t \le s \le T$$

with $\widehat{G}(\cdot)$ and $\widehat{Q}(\cdot)$ defined as in (5.10), and

(5.35)
$$Q_{xx}(s,x) + \bar{Q}(s) - K_0(1+|x|) \Big| \sum_{i=1}^n A^i_{xx}(s,x) - \sum_{j=1}^m e^T_j R(s)^{-1} \tilde{B}^i(s)^T S^j_{xx}(s,x) \Big| I - \sum_{j=1}^m \Big[e^T_j R(s)^{-1} S(s,x) \Big] S^j_{xx}(s,x) - (1-\alpha)^{-1} S_x(s,x)^T R(s)^{-1} S_x(s,x) \ge 0,$$

with $K_0 > 0$ being obtained in Lemma 5.5 or Lemma 5.6.

From (5.35), we see that due to the nonlinearity of $x \mapsto (A(t,x), S(t,x))$, we basically need the semiconvexity of $x \mapsto Q(t,x)$ and the sufficient positive definiteness of $R(\cdot)$ to compensate.

6 Quasi-Riccati Equation.

Let us keep condition (5.18). We have seen that under certain conditions $D_{uu}J(t, x; u^*(\cdot))$ admits a bounded inverse at any optimal control $u^*(\cdot) \equiv u^*(\cdot; t, x)$. When this is the case, we have the following: Denoting $x_0 = t$,

$$V(x_0, x) = J(x_0, x; u^*(\cdot; x_0, x)), \qquad D_u J(x_0, x; u^*(\cdot; x_0, x)) = 0.$$

Thus, for $0 \leq i \leq n$,

$$u_{x_i}^*(\cdot;x_0,x) = -D_{uu}J(x_0,x;u^*(\cdot;x_0,x))^{-1}D_uJ_{x_i}(x_0,x;u^*(\cdot;x_0,x)).$$

Hence,

$$V_{x_i}(x_0, x) = J_{x_i}(x_0, x; u^*(\cdot ; x_0, x)) + D_u J(x_0, x; u^*(\cdot ; x_0, x)) u^*_{x_i}(\cdot ; x_0, x) = J_{x_i}(x_0, x; u^*(\cdot ; x_0, x)).$$

Further, for $0 \leq i, j \leq n$,

$$\begin{aligned} V_{x_i x_j}(x_0, x) &= J_{x_i x_j}(x_0, x; u^*(\cdot; x_0, x)) + D_u J_{x_i}(x_0, x; u^*(\cdot; x_0, x)) u^*_{x_j}(\cdot; x_0, x) \\ &+ [D_{uu} J(x_0, x; u^*(\cdot; x_0, x)) u^*_{x_i}(\cdot; x_0, x)] u^*_{x_j}(\cdot; x_0, x) \\ &= J_{x_i x_j}(x_0, x; u^*(\cdot; x_0, x)). \end{aligned}$$

Therefore, $V(\cdot, \cdot)$ is actually twice continuously differentiable. Consequently, $V(\cdot, \cdot)$ satisfies the HJB equation in the classical sense, and by the smoothness of the coefficients, we can differentiate the equation once. Note that in the current case, our HJB equation reads:

$$\begin{cases} V_t(t,x) + V_x(t,x)A(t,x) + Q(t,x) \\ -\frac{1}{2}[V_x(t,x)B(t) + S(t,x)^T]R(t)^{-1}[B(t,x)^T V_x(t,x)^T + S(t,x)] = 0, \quad (t,x) \in [0,T) \times \mathbb{R}^n, \\ V(T,x) = G(x), \qquad x \in \mathbb{R}^n. \end{cases}$$

Now, we define

$$P(t,x) = V_x(t,x)^T.$$

Then

$$P_x(t,x) = V_{xx}(t,x) = P_x(t,x)^T, \qquad \forall (t,x) \in [0,T) \times \mathbb{R}^n,$$

and the following holds:

(6.1)
$$\begin{cases} P_t(t,x) + P_x(t,x)A(t,x) + A_x(t,x)^T P(t,x) + Q_x(t,x)^T \\ -[P_x(t,x)B(t) + S_x(t,x)^T]R(t)^{-1}[B(t)^T P(t,x) + S(t,x)] = 0, \quad (t,x) \in [0,T] \times \mathbb{R}^n, \\ P(T,x) = G_x(x), \qquad x \in \mathbb{R}^n. \end{cases}$$

The above is called a *Quasi-Riccati equation* of Problem (AQ). This is an extension of that presented in [13] for linear-convex problems. We now have the following result.

Proposition 6.1. Let (H1)–(H3) hold. Let $(X^*(\cdot), u^*(\cdot))$ be an optimal pair of Problem (AQ). Suppose the value function $V(\cdot, \cdot)$ of Problem (AQ) is twice differentiable. Then $P(\cdot, \cdot) \equiv V_x(\cdot, \cdot)^T$ is a solution to the quasi-Riccati equation (6.1), and the optimal control $u^*(\cdot)$ admits the following state feedback representation:

(6.2)
$$u^*(s) = -R(s)^{-1} \Big[B(s)^T P(s, X^*(s)) + S(x, X^*(s)) \Big], \qquad s \in [t, T].$$

Proof. It is known that if $(X^*(\cdot), u^*(\cdot))$ is an optimal pair of Problem (AQ) for the initial pair $(t, x) \in [0, T) \times \mathbb{R}^n$, and $Y(\cdot)$ is the solution to the corresponding adjoint equation, then

$$Y(s) = V_x(s.X^*(s))^T = P(s,X^*(s)), \qquad s \in [t,T],$$

and

(6.3)
$$u^{*}(s) = -R(s)^{-1} \Big[B(s)^{T} Y(s) + S(x, X^{*}(s)) \Big] \\ = -R(s)^{-1} \Big[B(s)^{T} P(s, X^{*}(s)) + S(x, X^{*}(s)) \Big], \qquad s \in [t, T]$$

This proves our conclusion.

In the case

$$A(t,x) = A(t)x, \quad Q(t,x) = \frac{1}{2} \langle Q(t)x, x \rangle, \quad S(t,x) = S(t)x, \quad G(x) = \frac{1}{2} \langle Gx, x \rangle$$

we see that

$$P(t,x) = P(t)x, \qquad (t,x) \in [0,T] \times \mathbb{R}^n,$$

with $P(\cdot)$ being the solution to the following:

$$\begin{cases} \dot{P}(t) + P(t)A(t) + A(t)^{T}P(t) + Q(t) - [P(t)B(t) + S(t)^{T}]R(t)^{-1}[B(t)^{T}P(t) + S(t)] = 0, \quad t \in [0, T], \\ P(T) = G, \end{cases}$$

which is the Riccati equation for a standard LQ problem.

To conclude this section, we present two illustrative examples.

Example 6.2. Consider the following one-dimensional linear controlled system:

$$\begin{cases} \dot{X}(s) = X(s) + u(s), & s \in [t, T], \\ X(t) = x, \end{cases}$$

with cost functional:

$$J(t,x;u(\cdot)) = \int_{t}^{T} \left(-\frac{1}{2}\cos^{2}X(s) + \frac{1}{2}\rho X(s)^{2} \right) ds - \frac{1}{2}\sin^{2}X(T),$$

where $\rho > 0$. This is a linear-semi-convex problem. According to Proposition 5.4, we may choose

$$\bar{Q}(t) \equiv \bar{G} = 1$$

Then

$$\widehat{G}(t) = \int_{t}^{T} e^{2(T-s)} ds = \frac{e^{2(T-t)} - 1}{2} \le \widehat{G}(0) = \frac{e^{2T} - 1}{2},$$

and

$$\widehat{Q}(s) = \int_{s}^{T} \int_{s}^{\tau} e^{2(\tau-r)} dr d\tau = \frac{1}{2} \int_{s}^{T} \left(e^{2(\tau-s)} - 1 \right) ds$$
$$= \frac{e^{2(T-s)} - 1}{4} - \frac{T-s}{2} \le \widehat{Q}(0) = \frac{e^{2T} - 1}{4} - \frac{T}{2}.$$

Hence, under condition

$$\rho > \frac{e^{2T} - 1}{4} - \frac{T}{2} + \frac{e^{2T} - 1}{2} = \frac{3(e^{2T} - 1)}{4} - \frac{T}{2}$$

we have the strict convexity of $u(\cdot) \mapsto J(t, x; u(\cdot))$. Therefore, optimal control unique exists and the value function is differentiable. In this case the optimal control $u^*(\cdot)$ admits a state feedback representation:

$$u^*(s) = -\rho^{-1}P(s, X^*(s)), \qquad s \in [t, T],$$

with $P(\cdot, \cdot)$ solves the following quasi-Riccati equation:

$$\begin{cases} P_t(t,x) + xP_x(t,x) + P(t,x) + \sin 2x - \rho^{-1}P_x(t,x)P(t,x) = 0, & t \in [0,T], \\ P(T,x) = -\sin 2x. \end{cases}$$

Example 6.3. Consider a one-dimensional controlled affine system

$$\begin{cases} \dot{X}(s) = \sqrt{1 + |X(s)|^2} + u(s), & s \in [t, T], \\ X(t) = x, \end{cases}$$

with cost functional

$$J(t, x; u(\cdot)) = \int_{t}^{T} \left(\frac{Q}{2} |X(s)|^{2} + \frac{R}{2} |u(s)|^{2}\right) ds.$$

In this case, we may take

$$A_0 = 1$$
, $Q_0 = 0$, $\bar{L}_A = 1$, $\bar{L}_B = \frac{1}{R}$, $\bar{L}_G = 0$, $\bar{L}_Q = q$.

Then the first condition in (5.17) automatically holds, and the second condition reads

$$\frac{2R + Q(e^{2T} - 1)}{2R} < \frac{1}{T}$$

This will be true if T < 1 is small. Next, by looking at the proof of Lemma 5.5, we see that

$$K_{1} = 0, \qquad K_{2} = \frac{RT}{2R(1-T) - QT(e^{2T} - 1)},$$

$$K_{3} = \frac{2R(1-T) - QT(e^{2T} - 1) + RT}{2R(1-T) - QT(e^{2T} - 1)},$$

$$K_{4} = \frac{QRT^{2}e^{T}}{2R(1-T) - QT(e^{2T} - 1)},$$

$$K_{5} = \frac{QRTe^{T}}{2R(1-T) - QT(e^{2T} - 1)}.$$

Then

$$|Y(s)| \le K_4 + K_5|X(s)| = \frac{QRTe^T(T+|X(s)|)}{2R(1-T) - QT(e^{2T}-1)}.$$

Now, if we assume

$$Q - \frac{QRTe^T}{2R(1-T) - QT(e^{2T} - 1)} \frac{T + |x|}{(1+|x|^2)^{\frac{3}{2}}} \ge 0, \qquad \forall x \in \mathbb{R}$$

which can be achieved if T > 0 is small, then Theorem 5.7 applies. In the current case, the quasi-Riccati equation reads:

(6.4)
$$\begin{cases} P_t(t,x) + \sqrt{1+|x|^2} P_x(t,x) + \frac{x}{\sqrt{1+|x|^2}} P(t,x) + Q - P(t,x) P_x(t,x) = 0, \quad (t,x) \in [0,T] \times \mathbb{R}^n, \\ P(T,x) = 0, \qquad x \in \mathbb{R}^n. \end{cases}$$

According to our result, under certain conditions (involving the constant K_0), the above quasi-Riccati equation admits a solution via which an optimal control admits a state feedback representation.

7 Concluding Remarks.

We have presented some very primitive results concerning what we call the affine-quadratic optimal control problems, which are a natural generalization of classical LQ problems, and also contains linear-convex problems and linear-semi-convex problems. Our results for linear state equation cover and substantially extend the known results for LQ problems and linear-convex problems. Further, we have some results for affine state equations. However, we see that there are a lot challenging problems left open. Here are a couple of these:

(i) Under our conditions, optimal controls exist and the optimality system which is a two-point boundary value problem is always solvable. It is a natural question if the corresponding quasi-Riccati equation is always solvable? A technical question relevant to this problem is: When the two-point boundary value problem is always solvable, can one obtain an estimate

$$|Y(s)| \le K(1 + |X(s)|), \qquad s \in [t, T]$$

without additional restrictive conditions?

(ii) What happens if the dependence of B(t, x) and R(t, x) on x is allowed? For such a situation, some new techniques might need to be developed.

We expect to report some further relevant results in our future publications.

References

- M. Bardi and I. Capuzzo-Dolcetta, Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations, Birkhäuser, Boston, 1997.
- M. Bardi and F. Da Lio, On the Bellman equation for some unbounded control problems, Nonlinear Diff. Eqs. Appl., 4 (1997), 491–510.
- [3] L. M. Benveniste and J. A. Scheinkman, On the differentiability of the value function in dynamic models of economics, Econometrica, 47 (1979), 727–732.
- [4] L. D. Berkovitz, Optimal Control Theory, Springer-Verlag, New York, 1974.
- [5] P. Cannarsa and H. Frankowska, Some characterizatins of optimal trajecotries in control theory, SIAM J. Control Optim., 29 (1991), 1322–1347.
- [6] H. Frankowska, Value Function in Optimal Control, Mathematical Control Theory, Part 1,2 (Trieste, 2001), 516–653.
- [7] R. E. Kalman, Contributions to the theory of optimal control, Bol. Soc. Mat. Mexicana, 5 (1960), 102–119.
- [8] H. Qiu and J. Yong, Hamilton-Jacobi equations and two-person zero-sum differential games with unbounded controls, COCV, to appear. See also arXiv:1011v2. Math.
- [9] J. P. Rincón-Zapatero and M. S. Santos, Differentiability of the value function in continuous-time economic models, preprint.
- [10] J. Yong, Stochastic optimal control and forward-backward stochastic differential equations, Computational Appl. Math., 21 (2002), 369–403.
- [11] J. Yong and X. Y. Zhou, Stochastic Control: Hamiltonian Systems and HJB Equations, Springer-Verlag, 1999.
- [12] Y. You, A nonquadratic Bolza problem and a quasi-Riccati equation for distributed parameter systems, SIAM J. Control Optim. 25 (1987), 905–920.
- [13] Y. You, Synthesis of time-variant optimal control with nonquadratic criteria, J. Math. Anal. Appl., 209 (1997), 662–682.