

THE STRONG GLOBAL DIMENSION OF PIECEWISE HEREDITARY ALGEBRAS

EDSON RIBEIRO ALVARES, PATRICK LE MEUR, AND EDUARDO N. MARCOS

In memory of Dieter Happel

ABSTRACT. Let A be a finite-dimensional piecewise hereditary algebra over an algebraically closed field. This text investigates the strong global dimension of A . This invariant is characterised in terms of the lengths of sequences of tilting mutations relating A to a hereditary abelian category, in terms of the generating hereditary abelian subcategories of the derived category of A , and in terms of the Auslander-Reiten structure of that derived category.

INTRODUCTION

Let A be a finite-dimensional algebra over an algebraically closed field k . Its category of finitely generated (left) modules is denoted by $\text{mod } A$. Then, A is called *piecewise hereditary* if the bounded derived category $\mathcal{D}^b(\text{mod } A)$ is equivalent as a triangulated category to $\mathcal{D}^b(\mathcal{H})$ where \mathcal{H} is a hereditary abelian (k -linear) category with split idempotents, finite-dimensional Hom-spaces, and with tilting objects. In the particular case where $A \simeq \text{End}_{\mathcal{H}}(T)^{\text{op}}$ for some tilting object $T \in \mathcal{H}$, the algebra A is called *quasi-tilted*. It is called *tilted* when, in addition, $\mathcal{H} \simeq \text{mod } H$ for some finite-dimensional hereditary algebra H . In [13], Happel proved that a hereditary abelian category as above is equivalent to the category of finitely generated modules over a hereditary algebra or to the category of coherent sheaves over a weighted projective line [10].

In the representation theory of finite-dimensional algebras, piecewise hereditary algebras play a particular and important role. On the one hand, this is due to information that is already known on the representation theory of hereditary algebras, of tilted algebras (see [16]) or of quasi-tilted algebras of canonical type (see [24]), and also to Happel's description of the bounded derived category of hereditary abelian categories (see below). On the other hand, these algebras are used in many parts of representation theory. For instance, in order to develop the representation theory of other classes of algebras such as the selfinjective algebras (see [28]) or the cluster tilted algebras (see [4]), in order to investigate singularity theory (see [20]), or in order to categorify cluster algebras (see [8]).

The homological characterisation of quasi-tilted algebras [14] and the Liu-Skowroński criterion for tilted algebras (see [5]) suggest that the quasi-tilted algebras are the closest piecewise hereditary algebras to hereditary ones, and it is the main objective of this text to give theoretical and numerical criteria to determine how far a piecewise hereditary algebra is from being hereditary.

Recall the description of $\mathcal{D}^b(\mathcal{H})$ made by Happel in [11]: Any object is the direct sum of (finitely many) stalk complexes $X[i]$ ($X \in \mathcal{H}$ and $i \in \mathbb{Z}$); And, for every $i, j \in \mathbb{Z}$ and $X, Y \in \mathcal{H}$, the morphism space $\text{Hom}(X[i], Y[j])$ is naturally isomorphic to $\text{Hom}_{\mathcal{H}}(X, Y)$ if $i = 0$, to $\text{Ext}_{\mathcal{H}}^1(X, Y)$ if $i = 1$, and is

Date: October 7, 2018.

The first named author acknowledges support from DMAT-UFPR and CNPq-Universal 477880/2012-6.

The second named author acknowledges financial support from FAPESP 2014/09310-5, CAPES, MathAmSud, and RFBM.

The third named author acknowledges financial support from CNPq, FAPESP, MathAmSud, and Prosul-CNPq n°490065/2010-4.

equal to zero otherwise. Hence, when $\mathcal{D}^b(\text{mod } A) \simeq \mathcal{D}^b(\mathcal{H})$, then there exists a tilting object $T \in \mathcal{D}^b(\mathcal{H})$ (that is, an object such that $\text{Hom}(T, T[i]) = 0$ for $i \in \mathbb{Z} \setminus \{0\}$, and such that $\mathcal{D}^b(\mathcal{H})$ is the smallest full triangulated subcategory of $\mathcal{D}^b(\mathcal{H})$ containing T and stable under taking direct summands) such that $A \simeq \text{End}(T)^{\text{op}}$ as k -algebras. In particular, there exists a minimal $\ell \in \mathbb{N}$ and there exists $s \in \mathbb{Z}$ such that T lies in the additive closure $\bigvee_{i=0}^{\ell} \mathcal{H}[s+i]$ of the union $\bigcup_{i=0}^{\ell} \mathcal{H}[s+i]$. When $\ell = 0$, then A is quasi-tilted. And one may expect that the larger ℓ , the further A is from being quasi-tilted. Note however that there exist examples where $\ell = 1$ and A is hereditary (see [12]).

Recall also the characterisation proved by Happel, Rickard and Schofield [15]: $\mathcal{D}^b(\text{mod } A)$ is equivalent to bounded derived category of the module category of a finite-dimensional hereditary algebra if and only if there exists a sequence of algebras with first term such a hereditary algebra, last term A , and where each term is isomorphic to the opposite of the endomorphism algebra of a tilting module over the preceding term. If the sequence has $\ell + 2$ terms, then $\text{gl.dim. } A \leq \ell + 2$. Again, one expects that, the larger ℓ , the further is A from being hereditary. However, in many examples, $\text{gl.dim. } A$ appears to be small whereas ℓ is large.

In the previous considerations, the parameter ℓ fails to give a precise measure of how far a piecewise hereditary algebra is from being quasi-tilted. Recently a new invariant for piecewise hereditary algebras has emerged and the present text aims at giving some evidence of its relevance to give such a measure. This invariant is the *strong global dimension*. It was first defined in [27] in terms of *width* of complexes. The present text makes use of a slightly different definition given by Happel and Zacharia in [17] and expressed in terms of *length* of complexes. The original definition may be recovered from the latter one by adding 1 to the invariant. Define the strong global dimension, $\text{s.gl.dim. } A \in \mathbb{N} \cup \{+\infty\}$, as follows. Let X be an indecomposable object in the homotopy category of bounded complexes of finitely generated projective A -modules. Let

$$P: \cdots \rightarrow 0 \rightarrow 0 \rightarrow P^r \rightarrow P^{r+1} \rightarrow \cdots \rightarrow P^{s-1} \rightarrow P^s \rightarrow 0 \rightarrow 0 \rightarrow \cdots,$$

be a minimal projective resolution of X , where $P^r \neq 0$ and $P^s \neq 0$. Then define the length of X as

$$\ell(X) = s - r.$$

The definition of the strong global dimension used in this text is

$$\text{s.gl.dim. } A = \sup_X \ell(X)$$

where X runs through all such indecomposable objects. It follows from the definition that $\text{s.gl.dim. } A = 1$ if and only if A is hereditary and not semi-simple. In [19, Problem 1], the question was asked whether A is piecewise hereditary when it has finite strong global dimension. This has been studied by several authors. The case of radical square-zero algebras was treated in [19]. Moreover, it follows from [6, 7] and [27] (see also [19]) that the class of algebras having both finite strong global dimension and a tame repetitive algebra coincides with the class of piecewise hereditary algebras having nonnegative Euler form. The equivalence conjectured in [19, Problem 1] was proved in the general case by Happel and Zacharia ([17]) getting as a byproduct that $\text{s.gl.dim. } A = 2$ if and only if A is quasi-tilted and not hereditary. It is worth noticing the following characterisation of the finiteness of the strong global dimension. By the main result of [27], the algebra A has finite strong global dimension if and only if the repetitive algebra \widehat{A} is locally support finite in the sense of [9]. Hence, if A has finite strong global dimension, then the push-down functor $\text{mod } \widehat{A} \rightarrow \text{mod } T(A)$ is dense (here $T(A)$ denotes the trivial extension). The question to know whether the converse implication holds was asked in [19, Problem 2]. So far, that question remains open (see [7, Theorem (B)] for a positive answer when \widehat{A} is tame).

Let \mathcal{T} be a triangulated category which is triangle equivalent to the bounded derived category of a hereditary abelian category (which is always assumed to be Hom-finite, to have split idempotents and

tilting objects). Let $T \in \mathcal{T}$ be a tilting object and let A be the piecewise hereditary algebra $\text{End}(T)^{\text{op}}$. The purpose of this text is therefore to answer the following questions:

- To what extent does $\text{s.gl.dim. } A$ measure how far A is from being quasi-tilted?
- Is it possible to compute the strong global dimension or to characterise it?

The first main result of this text gives an answer in terms of Happel's description of the bounded derived category of a hereditary abelian category. Note that part (1) is a direct consequence of that description.

Theorem 1. *Let \mathcal{T} be a triangulated category which is triangle equivalent to the bounded derived category of a hereditary abelian category. Let $T \in \mathcal{T}$ be a tilting object. Assume that $\text{End}(T)^{\text{op}}$ is not a hereditary algebra. There exists a full and additive subcategory $\mathcal{H} \subseteq \mathcal{T}$ which is hereditary and abelian, such that the embedding $\mathcal{H} \hookrightarrow \mathcal{T}$ extends to a triangle equivalence $\mathcal{D}^b(\mathcal{H}) \simeq \mathcal{T}$, and such that $T \in \bigvee_{i=0}^{\ell} \mathcal{H}[i]$ for some integer $\ell \geq 0$. Moreover*

- (1) $\text{s.gl.dim. End}(T)^{\text{op}} \leq \ell + 2$ for any such pair (\mathcal{H}, ℓ) , and
- (2) there exists such a pair (\mathcal{H}, ℓ) verifying $\text{s.gl.dim. End}(T)^{\text{op}} = \ell + 2$.

The second main result of this text gives an answer to the above questions in terms of Happel, Rickard and Schofield's theorem and using the concept of *tilting mutations* in triangulated categories ([1]): Let $T \in \mathcal{T}$ be a tilting object, let $T = T_1 \oplus T_2$ be a direct sum decomposition such that $\text{Hom}(T_2, T_1) = 0$ and consider a triangle $T'_2 \rightarrow M \rightarrow T_2 \rightarrow T'_2[1]$ where $M \rightarrow T_2$ is a minimal right add T_1 -approximation, then $T' = T_1 \oplus T'_2$ is a tilting object and called *obtained from T by tilting mutation*.

Theorem 2. *Let \mathcal{T} be a triangulated category which is triangle equivalent to the bounded derived category of a hereditary abelian category. Let $T \in \mathcal{T}$ be a tilting object. Assume that $\text{End}(T)^{\text{op}}$ is not hereditary. Then there exists an integer $\ell \geq 0$ and a sequence $T^{(0)}, T^{(1)}, \dots, T^{(\ell)}$ of tilting objects in \mathcal{T} such that*

- $\text{End}(T^{(0)})^{\text{op}}$ is a quasi-tilted algebra, $T^{(\ell)} = T$, and
- for every i the object $T^{(i)}$ is obtained from $T^{(i-1)}$ by a tilting mutation.

For any such sequence, $\text{s.gl.dim. End}(T)^{\text{op}} \leq \ell + 2$. Moreover, there exists such a sequence such that $\text{s.gl.dim. End}(T^{(i)})^{\text{op}} = 2 + i$ for every i (and, in particular $\text{s.gl.dim. End}(T)^{\text{op}} = 2 + \ell$).

This theorem is related to the second fundamental result mentioned above in the following way. Let A and H be algebras such that H is hereditary and $\mathcal{D}^b(\text{mod } A) \simeq \mathcal{D}^b(\text{mod } H)$. Assume that $A_0 = H, \dots, A_{\ell+1} = A$ is a sequence of algebras such that $A_i = \text{End}_{A_{i-1}}(M^{(i-1)})^{\text{op}}$ for a tilting A_{i-1} -module $M^{(i-1)}$ for every i . Then there exist tilting objects $T^{(0)}, \dots, T^{(\ell)}$ in $\mathcal{D}^b(\text{mod } H)$ such that $A_i \simeq \text{End}(T^{(i-1)})^{\text{op}}$ for every i , and which correspond to the tilting modules $M^{(0)}, \dots, M^{(\ell)}$ under suitable triangle equivalences $\mathcal{D}^b(\text{mod } H) \simeq \mathcal{D}^b(\text{mod } A_i)$. Then $\text{End}(T^{(0)})^{\text{op}}$ is tilted, and it follows from [18, Thm. 4.2] that $\text{s.gl.dim. End}(T^{(i-1)})^{\text{op}} \leq i + 2$ for every i . When H is of finite representation type the sequence $A_0, \dots, A_{\ell+1}$ may be chosen such that $M^{(i)}$ is an APR tilting module for every i . In such a situation $T^{(i)}$ is obtained from $T^{(i-1)}$ by a tilting mutation. From this point of view, Theorem 2 expresses the strong global dimension as the infimum of the number $\ell + 2$ of terms in all possible sequences $A_0, \dots, A_{\ell+1}$.

The proofs of Theorem 1 and Theorem 2 are based on the description of $\text{s.gl.dim. End}(T)^{\text{op}}$ in terms of the connected components of the Auslander-Reiten quiver of \mathcal{T} in which specific direct summands of T lie. The Auslander-Reiten structure of \mathcal{T} is described in [11].

The strategy of these proofs is described in Section 1. In particular, the structure of the text is given in 1.4. The text uses the following notation and general setup. Keep \mathbf{k} and \mathcal{T} as set previously. Given a full subcategory \mathcal{H} of \mathcal{T} which is hereditary, abelian and stable under taking direct summands, the embedding $\mathcal{H} \hookrightarrow \mathcal{T}$ extends to an equivalence of triangulated categories $\mathcal{D}^b(\mathcal{H}) \xrightarrow{\sim} \mathcal{T}$ if and only if \mathcal{H} generates \mathcal{T} as a triangulated category. Subcategories satisfying all these conditions are used frequently in this text.

They are called *hereditary abelian generating subcategories*. When \mathcal{H} arises from a weighted projective line, the full subcategory consisting of torsion objects (or torsion free objects) is denoted by \mathcal{H}_0 (or \mathcal{H}_+ , respectively). Given $X, Y \in \mathcal{T}$, the space $\text{Hom}_{\mathcal{T}}(X, Y)$ is denoted by $\text{Hom}(X, Y)$, and the more convenient notation $\text{Ext}^i(X, Y)$ will stand for $\text{Hom}(X, Y[i])$ ($i \in \mathbb{Z}$) whenever X and Y lie in a same hereditary abelian generating subcategory. Given an additive category \mathcal{A} , the class of indecomposable objects in \mathcal{A} is denoted by $\text{ind } \mathcal{A}$. The standard duality functor $\text{Hom}(-, \mathbb{k})$ is denoted by D . By an *Auslander-Reiten component* (of \mathcal{T}) is meant a connected component of the Auslander-Reiten quiver of \mathcal{T} . The Auslander-Reiten translation of \mathcal{T} is denoted by τ . By a *transjective component* is meant an Auslander-Reiten component which has only finitely many τ -orbits. Let $\mathcal{A} \subseteq \mathcal{T}$ be a full additive subcategory stable under taking direct summands. It is called a *one-parameter family of pairwise orthogonal tubes* if it is convex and its indecomposable objects form a (disjoint) union of pairwise orthogonal tubes in the Auslander-Reiten quiver of \mathcal{T} , and if it is maximal for these properties; equivalently, there exists a hereditary abelian generating subcategory $\mathcal{H} \subseteq \mathcal{T}$ arising from a weighted projective line and such that $\mathcal{A} = \mathcal{H}_0$ (see A.5). Here two tubes \mathcal{U}, \mathcal{V} are called *orthogonal* whenever $\text{Hom}(X, Y[i]) = 0$ for every $X \in \mathcal{U}, Y \in \mathcal{V}, i \in \mathbb{Z}$. On the other hand, \mathcal{A} is called a *maximal convex family of $\mathbb{Z}A_\infty$ components* if it is convex and its indecomposable objects form a (disjoint) union of Auslander-Reiten components of shape $\mathbb{Z}A_\infty$ in the Auslander-Reiten quiver of \mathcal{T} , and if it is maximal for these properties. This is the case if and only if exactly one of the following situations occurs for some hereditary abelian generating subcategory $\mathcal{H} \subseteq \mathcal{T}$

- \mathcal{H} arises from a weighted projective line with negative Euler characteristic and $\mathcal{A} = \mathcal{H}_+$,
- or else, \mathcal{H} arises from a hereditary algebra of wild representation type and \mathcal{A} consists of the objects obtained as direct sums of regular indecomposable modules over that algebra.

The reader is referred to [25, Chap. XIII] and [5, Chap. XVII] for a general account on the Auslander-Reiten structure (tubes, quasi-simples, components of shape $\mathbb{Z}A_\infty$) of hereditary algebras of tame and wild representation types, respectively. Note that in this text all tubes are stable.

1. OVERVIEW OF THE PROOFS OF THE MAIN THEOREMS AND STRUCTURE OF THE TEXT

1.1. An alternative definition for the strong global dimension. The whole proof makes use of the following characterisation of s.gl.dim. due to [3, Lem. 5.6]. Let $T, X \in \mathcal{T}$. Define $\ell_T^+(X), \ell_T^-(X) \in \mathbb{Z} \cup \{-\infty, +\infty\}$ as follows

$$\begin{cases} \ell_T^+(X) = \sup \{n \in \mathbb{Z} \mid \text{Hom}(X, T[n]) \neq 0\}, \\ \ell_T^-(X) = \inf \{n \in \mathbb{Z} \mid \text{Hom}(T[n], X) \neq 0\}. \end{cases}$$

Proposition ([3]). *Let $T \in \mathcal{T}$ be a tilting object. Let $A = \text{End}(T)^{\text{op}}$. Then $-\infty < \ell_T^-(X) \leq \ell_T^+(X) < +\infty$ for every $X \in \mathcal{T}$ indecomposable, and $\text{s.gl.dim. } A = \sup \{\ell_T^+(X) - \ell_T^-(X) \mid X \in \text{ind } \mathcal{T}\}$.*

In the sequel, if T is a tilting object in \mathcal{T} , and if $X \in \text{ind } \mathcal{T}$, then $\ell_T(X)$ denotes $\ell_T^+(X) - \ell_T^-(X)$. Note that $\ell_T(X) = \ell_T(X[i])$ for every $X \in \mathcal{T}$ and $i \in \mathbb{Z}$. Hence $\text{s.gl.dim. } A = \sup \{\ell_T(X) \mid X \in \text{ind } \mathcal{T}, \ell_T^-(X) = 0\}$.

1.2. Upper and lower bounds on the strong global dimension. The starting point of the proof of Theorem 1 is the following simple observation.

Proposition. *Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory. Let $T \in \mathcal{T}$ be a tilting object. If $T \in \bigvee_{i=0}^{\ell} \mathcal{H}[i]$ for some $\ell \in \mathbb{N}$, then $\text{s.gl.dim. } \text{End}(T)^{\text{op}} \leq \ell + 2$.*

Proof. This follows directly from Happel's description of $\mathcal{D}^b(\mathcal{H})$ and from the previous characterisation of the strong global dimension (see 1.1). \square

Hence, the nontrivial part of Theorem 1 is about lower bounds on the strong global dimension. Proving that theorem amounts to finding a hereditary abelian generating subcategory $\mathcal{H} \subseteq \mathcal{T}$ such that $T \in \bigvee_{i=0}^{\ell} \mathcal{H}[i]$ and $\ell+2 = \text{s.gl.dim. End}(T)^{\text{op}}$. Should this hold true, whenever $X \in \text{ind } \mathcal{T}$ is such that $\ell_T(X) = \text{s.gl.dim End}(T)^{\text{op}}$ and (up to shift) $\ell_T^-(X) = 0$, then $X \in \mathcal{H}[\ell+1]$, and there should exist indecomposable summands M_0, M_1 of T lying respectively in \mathcal{H} and $\mathcal{H}[\ell]$, together with non-zero morphisms $M_1 \rightarrow X$ and $X \rightarrow M_0[\ell+2]$. In order to get lower bounds on $\text{s.gl.dim. End}(T)^{\text{op}}$, one may therefore start from M_0 and M_1 and look for X and for nonzero morphisms $M_0 \rightarrow X$ and $X \rightarrow M_0[\ell+2]$. Using the Auslander-Reiten structure of \mathcal{T} , this is possible provided that M_0 and M_1 lie on specific Auslander-Reiten components.

1.3. The subcategories in which tilting objects start or end. In view of the previous considerations, it is useful to have some *a priori* knowledge on the Auslander-Reiten components of \mathcal{T} containing indecomposable direct summands of T lying in \mathcal{H} or $\mathcal{H}[\ell]$. In this text, these relevant Auslander-Reiten components are determined in terms of subcategories in which T *starts* or *ends*. Let $\mathcal{A} \subseteq \mathcal{T}$ be a full, additive and convex subcategory stable under taking direct summands. Here *convex* means that any path $X_0 \rightarrow \cdots \rightarrow X_n$ (of non-zero morphisms between indecomposable objects) is contained in \mathcal{A} as soon as $X_0, X_n \in \mathcal{A}$. In this text T is said to *start in* \mathcal{A} if the two following conditions hold true

- T has at least one indecomposable direct summand in \mathcal{A} ,
- for every indecomposable direct summand X of T there exists a path $X_0 \rightarrow \cdots \rightarrow X_n$ such that $X_0 \in \mathcal{A}$ and $X_n = X$.

Of course T is said to *end in* \mathcal{A} if the dual properties hold true. There are obviously many such subcategories. This text concentrates on three specific ones, namely

- (1) when $\text{ind } \mathcal{A}$ consists of a transjective Auslander-Reiten component (in which case T is said to start in a transjective component, for short), or
- (2) when \mathcal{A} is a one-parameter family of pairwise orthogonal tubes, or
- (3) when \mathcal{A} is a maximal convex family of $\mathbb{Z}A_{\infty}$ components.

1.4. Structure of the text. Consider a tilting object $T \in \mathcal{T}$. Section 2 establishes lower bounds on the strong global dimension making hypotheses on the indecomposable direct summands of T , as outlined in 1.2. For both Theorem 1 and Theorem 2, the proofs reduce the problem of determining $\text{s.gl.dim. End}(T)^{\text{op}}$ by replacing T by an adequate tilting object obtained by tilting mutation. In order to make a proper choice for that object, Section 3 determines the subcategories in which a given tilting mutation of T may start or end. Once those subcategories are known, it is necessary to determine the Auslander-Reiten components of \mathcal{T} that contain indecomposable direct summands of the involved tilting objects. This task is performed in Section 4. Using all this information, Section 5 determines $\text{s.gl.dim. End}(T)^{\text{op}}$ according to the subcategories in which T starts or ends. Finally, Section 6 proves Theorem 1 and Theorem 2. Appendix A collects technical material on bounded derived categories of weighted projective lines needed for the proofs of the main theorems.

2. LOWER BOUNDS ON THE STRONG GLOBAL DIMENSION

Given a tilting object $T \in \mathcal{T}$, this section gives lower bounds for $\text{s.gl.dim. End}(T)^{\text{op}}$. These form the technical heart of the proofs of Theorem 1 and Theorem 2. They are based on the existence of certain indecomposable direct summands of T . A separate subsection is devoted to each one of these results according to the following situations: the considered summands lie in transjective Auslander-Reiten components; or they lie in non-transjective Auslander-Reiten components; or they satisfy specific vanishing assumptions on morphism spaces.

2.1. Lower bounds using transjective Auslander-Reiten components. The first result on lower bounds on the strong global dimension is based on the existence of certain indecomposable direct summands of T lying in transjective Auslander-Reiten components. The corresponding setting is as follows.

Let Γ be a transjective Auslander-Reiten component of \mathcal{T} . Let Σ be a slice of Γ . Let S_1, \dots, S_n be the sources of Σ . Let $\mathcal{H} \subseteq \mathcal{T}$ be the full subcategory

$$\mathcal{H} = \{X \in \mathcal{T} \mid (\forall S \in \Sigma) (\forall i \neq 0) \operatorname{Hom}(S, X[i]) = 0\}.$$

Let $T \in \mathcal{T}$ be a tilting object and let $\ell \geq 1$ be an integer such that

- S_1, \dots, S_n are all indecomposable summands of T ,
- there exists an indecomposable summand L of T lying in $\mathcal{H}[\ell]$.

Lemma. *Under the previous setting there exists an object $M \in \tau^{-1}\Sigma[\ell + 1]$ together with nonzero morphisms $L \rightarrow M$ and $M \rightarrow \bigoplus_{i=1}^n S_i[\ell + 2]$. Hence $\ell_T(M) \geq \ell + 2$. In particular, $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$.*

Proof. It is useful to prove first that $L \in \tau^{-2}\mathcal{H}[\ell]$. For this purpose note that

$$(\operatorname{ind} \mathcal{H}[\ell]) \setminus (\operatorname{ind} \tau^{-2}\mathcal{H}[\ell]) = \Sigma[\ell] \cup \tau^{-1}\Sigma[\ell],$$

as sets of indecomposable objects. Since $L \in \mathcal{H}[\ell]$, the claim therefore deals with proving that $L \notin \Sigma[\ell]$ and $L \notin \tau^{-1}\Sigma[\ell]$. Using Serre duality and that T is tilting implies that

$$\operatorname{Hom}\left(\bigoplus_{i=1}^n S_i[\ell], L\right) = 0 \quad \text{and} \quad \operatorname{Hom}\left(\bigoplus_{i=1}^n \tau^{-1}S_i[\ell], L\right) = 0.$$

Since $S_1[\ell], \dots, S_n[\ell]$ (or $\tau^{-1}S_1[\ell], \dots, \tau^{-1}S_n[\ell]$) are the sources of the slice $\Sigma[\ell]$ (or $\tau^{-1}\Sigma[\ell]$, respectively), this entails that $L \notin \Sigma[\ell]$ and $L \notin \tau^{-1}\Sigma[\ell]$. Thus $L \in \tau^{-2}\mathcal{H}[\ell]$.

The category $\tau^{-2}\mathcal{H}[\ell]$ is abelian and its indecomposable injectives are the objects in $\tau^{-1}\Sigma[\ell + 1]$, up to isomorphism. Hence

$$(\exists M \in \tau^{-1}\Sigma[\ell + 1]) \operatorname{Hom}(L, M) \neq 0.$$

Besides $\operatorname{Hom}\left(\bigoplus_{i=1}^n \tau^{-1}S_i[\ell + 1], M\right) \neq 0$ for $\tau^{-1}S_i[\ell + 1], \dots, \tau^{-1}S_n[\ell + 1]$ are the sources of the slice $\tau^{-1}\Sigma[\ell + 1]$ of $\Gamma[\ell + 1]$. Serre duality then implies that $\operatorname{Hom}\left(M, \bigoplus_{i=1}^n S_i[\ell + 2]\right) \neq 0$. \square

2.2. Lower bounds using non-transjective Auslander-Reiten components. The second result on lower bounds on the strong global dimension is based on the existence of certain indecomposable direct summands of T not lying in a transjective Auslander-Reiten component. Here is the precise setting.

Let $\mathcal{H} \subseteq \mathcal{T}$ be hereditary abelian generating subcategory. Let $T \in \mathcal{T}$ be a tilting object and $\ell \in \mathbb{N}$ be such that there exist indecomposable direct summands M_0, M_1 of T satisfying the following

- $M_0 \in \mathcal{H}$ and
- $M_1 \in \mathcal{H}[\ell]$ and M_1 lies in a non-transjective Auslander-Reiten component of $\mathcal{H}[\ell]$.

Recall that the non-transjective Auslander-Reiten components are either tubes or of shape $\mathbb{Z}A_\infty$. Let $X = \tau^{-1}M_0[\ell + 1]$ and $Y = \tau M_1[1]$. These lie in $\mathcal{H}[\ell + 1]$.

Lemma. *Under the setting described previously assume that both M_0 and M_1 lie in non-transjective Auslander-Reiten components of \mathcal{T} .*

- (1) *If \mathcal{T} contains a transjective Auslander-Reiten component then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 1$.*
- (2) *If there exists a tube $\mathcal{U} \subseteq \mathcal{T}$ such that $M_0 \in \mathcal{U}$ and $M_1 \in \mathcal{U}[\ell]$ then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$.*
- (3) *If \mathcal{H} arises from a weighted projective line, $M_0 \in \mathcal{H}_+$ and $M_1 \in \mathcal{H}_0[\ell]$, then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$.*
- (4) *If M_0, M_1 lie in Auslander-Reiten components of shape $\mathbb{Z}A_\infty$, then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$.*

Proof. (1) Let \mathcal{C} be the Auslander-Reiten component of \mathcal{T} such that $M_0[\ell + 1] \in \mathcal{C}$. Let Γ be the unique transjective Auslander-Reiten component of \mathcal{T} such that

$$(\forall V \in \mathcal{C}) (\exists U \in \Gamma) \quad \text{Hom}(U, V) \neq 0.$$

Let \mathcal{R} be the disjoint union of the non-transjective Auslander-Reiten components such that

$$(\forall V \in \mathcal{R}) (\exists U \in \Gamma) \quad \text{Hom}(U, V) \neq 0.$$

Therefore

- $\mathcal{C} \subseteq \mathcal{R}$,
- \mathcal{R} is the family of regular Auslander-Reiten components of $\mathcal{H}[\ell + 1]$, and
- $\mathcal{R}[-1]$ is the family of regular Auslander-Reiten components of $\mathcal{H}[\ell]$ and $M_1 \in \mathcal{R}[-1]$.

In order to prove that $\text{s.gl.dim. End}(T)^{\text{op}} \geq \ell + 1$, it is sufficient to find $S_0 \in \Gamma$ such that

$$\text{Hom}(M_1, S_0) \neq 0 \quad \text{and} \quad \text{Hom}(S_0, M_0[\ell + 1]) \neq 0.$$

First there exists a slice Σ in Γ such that

$$(\forall S \in \Sigma) \quad \text{Hom}(S, M_0[\ell + 1]) \neq 0.$$

Define the full subcategory $\mathcal{H}' \subseteq \mathcal{T}$ as $\mathcal{H}' = \{V \in \mathcal{T} \mid \text{Hom}(V, S[i]) = 0 \text{ if } i \neq 0 \text{ and } S \in \Sigma\}$. Then

- \mathcal{H}' is hereditary and abelian,
- the indecomposable injectives of \mathcal{H}' are the objects in Σ up to isomorphism, and
- $\mathcal{R}[-1]$ is the family of regular Auslander-Reiten components of \mathcal{H}' ; in particular $M_1 \in \mathcal{H}'$.

Thus there exists $S_0 \in \Sigma$ such that $\text{Hom}(M_1, S_0) \neq 0$. By hypothesis, $\text{Hom}(S_0, M_0[\ell + 1]) \neq 0$.

(2) There exists a tube $\mathcal{U} \subseteq \mathcal{T}$ such that $Y = \tau M_1[1] \in \mathcal{U}$ and $X = \tau^{-1} M_0[\ell + 1] \in \mathcal{U}$. Moreover there exist infinite sectional paths in \mathcal{U}

$$X \rightarrow \bullet \rightarrow \bullet \rightarrow \cdots \quad \text{and} \quad \cdots \rightarrow \bullet \rightarrow \bullet \rightarrow Y.$$


Since \mathcal{U} is a tube, the two paths intersect. Hence there exist $S \in \mathcal{U}$ and sectional paths in \mathcal{U}

$$\tau^{-1} M_0[\ell + 1] \rightarrow \cdots \rightarrow S \quad \text{and} \quad S \rightarrow \cdots \rightarrow \tau M_1[1].$$

Since the composition of morphisms along a sectional path does not vanish, there exist nonzero morphisms $\tau^{-1} M_0[\ell + 1] \rightarrow S$ and $S \rightarrow \tau M_1[1]$. Using Serre duality, this implies that $\text{Hom}(S, M_0[\ell + 2]) \neq 0$ and $\text{Hom}(M_1, S) \neq 0$. Thus $\ell_T(S) \geq \ell + 2$.

(3) Let $\mathcal{U} \subseteq \mathcal{H}_0[\ell + 1]$ be the tube such that $M_1[1] \in \mathcal{U}$. Applying A.2 (part (1)) yields an infinite sectional path $S = X_0 \rightarrow \cdots \rightarrow X_n \rightarrow \cdots$ such that $\text{Hom}(\tau^{-1} X_n, M_0[\ell + 2]) \neq 0$, for every $n \geq 0$. Since \mathcal{U} is a tube there also exists an infinite sectional path $\cdots \rightarrow \bullet \rightarrow \bullet \rightarrow \tau^2 M_1[1]$, and the two paths must intersect. Therefore there exists $n \geq 0$ together with a sectional path $X_n \rightarrow \cdots \rightarrow \tau^2 M_1[1]$. The composite morphism $X_n \rightarrow \tau^2 M_1[1]$ is thus nonzero. Using Serre duality entails that $\text{Hom}(M_1, \tau^{-1} X_n) \neq 0$. Thus $\ell_T(\tau^{-1} X_n) \geq \ell + 2$.

(4) The proof of the statement is better understood using the following diagram the details of which are explained below and where all the arrows represent nonzero morphisms.



Note that $X = \tau^{-1}M_0[\ell + 1]$ and $Y = \tau M_1[1]$ lie in $\mathcal{H}[\ell + 1]$.

- S is any quasi-simple object in any Auslander-Reiten component of $\mathcal{H}[\ell + 1]$ with shape $\mathbb{Z}A_\infty$ and \mathcal{C} is that Auslander-Reiten component,
- n is any integer large enough such that $\text{Hom}(X, \tau^n S) \neq 0$ and $\text{Hom}(\tau^{-n} S, Y) \neq 0$ (see [26, Chap. XVIII, 2.6] or [22, Prop. 10.1] according to whether \mathcal{H} arises from a hereditary algebra of wild representation type or from a weighted projective line with negative Euler characteristic, respectively), whence the curved arrows in the diagram,
- M is the (unique) object in \mathcal{C} such that there exist sectional paths of irreducible morphisms $\tau^n S \rightarrow \cdots \rightarrow M$ and $M \rightarrow \cdots \rightarrow \tau^{-n} S$ in $\mathcal{H}[\ell + 1]$. The arrows in the former path (or, in the latter path) are all monomorphisms (or, epimorphisms, respectively) in $\mathcal{H}[\ell + 1]$.

Since the diagram lies in $\mathcal{H}[\ell + 1]$, the composite morphisms $X \rightarrow M$ and $M \rightarrow Y$ arising from the paths $X \rightarrow \tau^n S \rightarrow \cdots \rightarrow M$ and $M \rightarrow \cdots \rightarrow \tau^{-n} S \rightarrow Y$ are nonzero. Using Serre duality and the definition of X and Y yields: $\text{Hom}(M_1, M) \neq 0$ and $\text{Hom}(X, M_0[\ell + 2]) \neq 0$. Thus, $\ell_T(M) \geq \ell + 2$. \square

2.3. Lower bounds using non-vanishing morphism spaces. The setting is the same as in 2.2. In particular the same notation (M_0, M_1, X, Y) is used here. Let $Z \rightarrow M_1$ be a minimal right almost split morphism in \mathcal{T} . As usual M_1 is called *quasi-simple* whenever Z is indecomposable.

Lemma. *Under the setting described previously, the following hold true.*

- (1) If $\text{Hom}(X, Y) \neq 0$, then $\ell_T(\tau^{-1}M_0) \geq \ell + 2$. In particular $\text{s.gl.dim. End}(T)^{\text{op}} \geq \ell + 2$.
- (2) If $\text{Ext}^1(Y, X) \neq 0$, then $\ell_T(\tau Z) \geq \ell + 2$ or $\ell_T(\tau^2 M_1) \geq \ell + 2$ according to whether M_1 is quasi-simple or not. In particular, $\text{s.gl.dim. End}(T)^{\text{op}} \geq \ell + 2$.
- (3) If $\text{Hom}(X, Y) = 0$ and $\text{Hom}(Y, X) \neq 0$, then $\ell_T(\tau Z) \geq \ell + 1$ or $\ell_T(\tau^2 M_1) \geq \ell + 1$ according to whether M_1 is quasi-simple or is not. In particular $\text{s.gl.dim. End}(T)^{\text{op}} \geq \ell + 1$.
- (4) If $\text{Ext}^1(X, Y) \neq 0$, then $\ell_T(\tau M_1) \geq \ell + 1$. In particular $\text{s.gl.dim. End}(T)^{\text{op}} \geq \ell + 1$.

Proof. (1) Serre duality gives

$$\begin{cases} 0 \neq \text{Hom}(X, Y) = \text{Hom}(\tau^{-1}M_0[\ell + 1], \tau M_1[1]) \simeq \text{DHom}(M_1, \tau^{-1}M_0[\ell + 1]), \\ 0 \neq \text{Hom}(\tau^{-1}M_0[\ell + 1], \tau^{-1}M_0[\ell + 1]) \simeq \text{DHom}(\tau^{-1}M_0[\ell + 1], M_0[\ell + 2]). \end{cases}$$

Thus $\ell_T(\tau^{-1}M_0[\ell + 1]) \geq \ell + 2$.

- (2) The hypothesis implies that $0 \neq \text{Hom}(\tau^2 M_1, M_0[\ell + 1])$, and hence $\ell_T^+(\tau^2 M_1) \geq \ell + 1$.

Assume first that M_1 is not quasi-simple. Therefore $\text{Hom}(\tau M_1, M_1) \neq 0$. Serre duality then implies $\text{Hom}(M_1[-1], \tau^2 M_1) \neq 0$. Hence $\ell_T^-(\tau^2 M_1) \leq -1$, and thus $\ell_T(\tau^2 M_1) \geq \ell + 2$.

Assume now that M_1 is quasi-simple. Since M_1 lies in an Auslander-Reiten component of $\mathcal{H}[\ell]$ which is a tube or of shape $\mathbb{Z}A_\infty$, there exists an almost split triangle $\tau^2 M_1 \rightarrow \tau Z \rightarrow \tau M_1 \rightarrow \tau^2 M_1[1]$. Since

$M_0[\ell + 1] \in \mathcal{H}[\ell + 1]$ and $\tau^2 M_1 \in \mathcal{H}[\ell]$, it follows that

$$0 \neq \text{Ext}^1(Y, X) = \text{Hom}(\tau^2 M_1, M_0[\ell + 1]) \subseteq \text{rad}(\tau^2 M_1, M_0[\ell + 1]).$$

In particular there exists a nonzero morphism $\tau^2 M_1 \rightarrow M_0[\ell + 1]$ which factors through $\tau^2 M_1 \rightarrow \tau Z$. Hence $\text{Hom}(\tau Z, M_0[\ell + 1]) \neq 0$, and therefore $\ell_T^+(\tau Z) \geq \ell + 1$. Moreover, using Serre duality yields

$$0 \neq \text{Hom}(\tau Z, \tau M_1) \simeq \text{DHom}(M_1[-1], \tau Z).$$

Hence $\ell_T^-(\tau Z) \leq -1$, and thus $\ell_T(\tau Z) \geq \ell + 2$.

(3) The hypotheses imply that $0 \neq \text{Hom}(Y, X) = \text{Hom}(\tau^2 M_1, M_0[\ell])$. Hence $\ell_T^+(\tau^2 M_1) \geq \ell$.

Assume first that M_1 is not quasi-simple. The argument used in (2) to study $\ell_T^-(\tau^2 M_1)$ also applies here and shows that $\ell_T^-(\tau^2 M_1) \leq -1$. Thus $\ell_T(\tau^2 M_1) \geq \ell + 1$.

Assume now that M_1 is quasi-simple. It follows from the hypotheses that

$$0 = \text{Hom}(X, Y) = \text{Hom}(\tau^{-1} M_0[\ell + 1], \tau M_1[1]) = \text{Hom}(M_0[\ell], \tau^2 M_1).$$

In particular $M_0[\ell] \not\cong \tau^2 M_1$, and therefore (see above)

$$0 \neq \text{Hom}(Y, X) = \text{Hom}(\tau^2 M_1, M_0[\ell]) \subseteq \text{rad}(\tau^2 M_1, M_0[\ell]).$$

Hence there exists a nonzero morphism $\tau^2 M_1 \rightarrow M_0[\ell]$ which factors through $\tau^2 M_1 \rightarrow \tau Z$. Therefore $\text{Hom}(\tau Z, M_0[\ell]) \neq 0$, and thus $\ell_T^+(\tau Z) \geq \ell$. The arguments used in (2) to prove that $\ell_T^-(\tau Z) \leq -1$ also apply here. Thus $\ell_T(\tau Z) \geq \ell + 1$.

(4) Serre duality gives

$$\begin{cases} 0 \neq \text{Ext}^1(X, Y) = \text{Hom}(\tau^{-1} M_0[\ell + 1], \tau M_1[2]) \simeq \text{DHom}(\tau M_1, M_0[\ell]), \\ 0 \neq \text{Hom}(M_1, M_1) \simeq \text{DHom}(M_1[-1], \tau M_1). \end{cases}$$

Thus $\ell_T(\tau M_1) \geq \ell + 1$. □

3. TILTING MUTATIONS

The proofs of Theorem 1 and that of Theorem 2 use inductions based on tilting mutation. The inductive step produces a new tilting object T' from the given tilting object T , such that $\text{End}(T')^{\text{op}}$ has strong global dimension smaller than that of $\text{End}(T)^{\text{op}}$. This section therefore checks that tilting mutation permits a convenient use of the lower bounds presented in Section 2. It proceeds as follows.

- 3.1 checks that T' is indeed tilting.
- 3.2 compares the lengths of a given object $X \in \mathcal{T}$ with respect to T and T' . This leads to a comparison of the strong global dimensions of $\text{End}(T)^{\text{op}}$ and $\text{End}(T')^{\text{op}}$.
- 3.3 locates the indecomposable direct summands of T' in terms of convex subcategories of \mathcal{T} defined by indecomposable direct summands of T . This serves to 3.4.
- In view of applying the results of Section 2 to both T and T' , 3.4 compares the subcategories in which T and T' start.

3.1. Setting for the section. Let $T \in \mathcal{T}$ be tilting. Consider a direct sum decomposition $T = T_1 \oplus T_2$ such that $\text{Hom}(T_2, T_1) = 0$. Let $M \rightarrow T_2$ be a minimal right $\text{add } T_1$ -approximation. It fits into a triangle

$$(\Delta) \quad T'_2 \rightarrow M \rightarrow T_2 \rightarrow T'_2[1].$$

Let $T' = T_1 \oplus T'_2$. The following result is fundamental in this work. It is an application of [1, Theorem 2.31 and Theorem 2.32 (b)] since $\text{add}(T_1) = \mu^-(\text{add } T, \text{add } T_1)$ (with the notation introduced therein).

Proposition. *Under the previous setting, T' is a tilting object in \mathcal{T} .*

Proof. Since the point of view and notation in [1] are slightly different from the ones in this text a proof is given below for the convenience of the reader. Because of Δ , the smallest triangulated subcategory of \mathcal{T} containing T' and stable under direct summands is \mathcal{T} . Hence it suffices to prove that $\text{Hom}(T', T'[i]) = 0$ for every $i \neq 0$. Almost every argument below uses that T is tilting so this will be implicit. Let $i \in \mathbb{Z}$.

First $\text{Hom}(T_1, T_1[i]) = 0$ if $i \neq 0$ because $T_1 \in \text{add}(T)$.

Next there is an exact sequence obtained by applying $\text{Hom}(T_1, -)$ to Δ

$$\text{Hom}(T_1, M[i-1]) \rightarrow \text{Hom}(T_1, T_2[i-1]) \rightarrow \text{Hom}(T_1, T_2'[i]) \rightarrow \text{Hom}(T_1, M[i]).$$

Since $M \rightarrow T_2$ is an $\text{add } T_1$ -approximation it follows that $\text{Hom}(T_1, T_2'[i]) = 0$ if $i \neq 0$.

Next there is an exact sequence obtained by applying $\text{Hom}(-, T_1[i])$ to Δ

$$\text{Hom}(M, T_1[i]) \rightarrow \text{Hom}(T_2', T_1[i]) \rightarrow \text{Hom}(T_2, T_1[i+1]).$$

Since $\text{Hom}(T_2, T_1) = 0$ it follows that $\text{Hom}(T_2', T_1[i]) = 0$ if $i \neq 0$.

Next there is an exact sequence obtained by applying $\text{Hom}(T_2, -)$ to Δ

$$\underbrace{\text{Hom}(T_2, T_2[i])}_{=0 \text{ if } i \neq 0} \rightarrow \text{Hom}(T_2, T_2'[i+1]) \rightarrow \underbrace{\text{Hom}(T_2, M[i+1])}_{=0}$$

where the rightmost term is zero if $i = -1$ by assumption on the decomposition $T = T_1 \oplus T_2$. Hence $\text{Hom}(T_2, T_2'[i+1]) = 0$ if $i \neq 0$. Therefore the exact sequence obtained by applying $\text{Hom}(-, T_2')$ to (Δ)

$$\underbrace{\text{Hom}(M, T_2'[i])}_{=0 \text{ if } i \neq 0 \text{ (see above)}} \rightarrow \text{Hom}(T_2', T_2'[i]) \rightarrow \underbrace{\text{Hom}(T_2, T_2'[i+1])}_{=0 \text{ if } i \neq 0}$$

entails that $\text{Hom}(T_2', T_2'[i]) = 0$ if $i \neq 0$. All these considerations prove that T' is tilting. \square

3.2. Behaviour of the strong global dimension under tilting mutations. It is natural to ask for a relationship between $\ell_T(X)$ and $\ell_{T'}(X)$. The following table gives an answer assuming (up to suspension) that $\ell_{T'}(X) = 0$, and denoting $\ell_{T'}(X)$ by ℓ .

	$\text{Hom}(T_2, X[1]) \neq 0$	$\text{Hom}(T_2, X[1]) = 0$
$\text{Hom}(X, T_1[\ell]) \neq 0$	$\ell_T^-(X) = -1$, $\ell_T^+(X) = \ell$, and $\ell_T(X) = \ell + 1$	$\ell_T^-(X) = 0$, $\ell_T^+(X) = \ell$, and $\ell_T(X) = \ell$
$\text{Hom}(X, T_1[\ell]) = 0$	$\ell_T^-(X) = -1$, $\ell_T^+(X) = \ell - 1$, and $\ell_T(X) = \ell$	$\ell_T^-(X) = 0$, $\ell_T^+(X) = \ell - 1$, and $\ell_T(X) = \ell - 1$

Indeed, examining the long exact sequence obtained upon applying $\text{Hom}(-, X)$ to Δ yields that

- $\text{Hom}(T_1[i], X) = \text{Hom}(M[i], X) = 0$ if $i < 0$ and $\text{Hom}(T_2[i], X) = 0$ if $i < -1$,
- $\ell_T^-(X) \geq -1$, with an equality if and only if $\text{Hom}(T_2[-1], X) \neq 0$,
- if $\text{Hom}(T_2[-1], X) = 0$, then $\text{Hom}(T_1, X) \neq 0$, and hence $\ell_T^-(X) = 0$;

Using dual considerations yields that

- $\text{Hom}(X, T_1[i]) = \text{Hom}(X, M[i]) = \text{Hom}(X, T_2[i]) = 0$ if $i > \ell$,
- $\ell_T^+(X) \leq \ell$, with an equality if and only if $\text{Hom}(X, T_1[\ell]) \neq 0$,
- if $\text{Hom}(X, T_1[\ell]) = 0$, then $\text{Hom}(X, T_2[\ell-1]) \neq 0$, and hence $\ell_T^+(X) = \ell - 1$.

Using that table, the following is immediate (keeping the setting of 3.1).

Proposition. $|\text{s.gl.dim. End}(T)^{\text{op}} - \text{s.gl.dim. End}(T')^{\text{op}}| \leq 1$.

3.3. Indecomposable direct summands and tilting mutation. In view of comparing the subcategories of \mathcal{T} in which T starts or ends to the corresponding ones for T' , it is useful to locate the indecomposable direct summands of T' with respect to those of T . This is done in the following result.

Lemma. *Let $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ be a triangle in \mathcal{T} . The following conditions are equivalent.*

- (i) *X is an indecomposable direct summand of T_2 and $N \rightarrow X$ is a right minimal $\text{add } T_1$ -approximation.*
- (ii) *X' is an indecomposable direct summand of T'_2 and $X' \rightarrow N$ is a left minimal $\text{add } T_1$ -approximation.*

Moreover when these conditions are satisfied the object X' lies in the full and convex subcategory of \mathcal{T} generated by $X[-1]$ and the indecomposable direct summands of N .

Proof. Assume (i). Let $X \rightarrow T_2$ and $T_2 \rightarrow X$ be morphisms such that the composite morphism $X \rightarrow T_2 \rightarrow X$ is identity. Since $N \rightarrow X$ and $M \rightarrow T_2$ are $\text{add } T_1$ -approximations, there are commutative diagrams whose rows are triangles

$$\begin{array}{ccccccc} X' & \longrightarrow & N & \longrightarrow & X & \longrightarrow & X'[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ T'_2 & \longrightarrow & M & \longrightarrow & T_2 & \longrightarrow & T'_2[1] \end{array} \quad \text{and} \quad \begin{array}{ccccccc} T'_2 & \longrightarrow & M & \longrightarrow & T_2 & \longrightarrow & T'_2[1] \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ X' & \longrightarrow & N & \longrightarrow & X & \longrightarrow & X'[1] \end{array}.$$

Since $N \rightarrow X$ is right minimal and since the composite morphism $X \rightarrow T_2 \rightarrow X$ is identity it follows that the composite morphism $N \rightarrow M \rightarrow N$ is an isomorphism, and hence so is the composite morphism $X' \rightarrow T'_2 \rightarrow X'$. This proves that X' is a direct summand of T'_2 .

In order to prove that X' is indecomposable, let $e \in \text{End}(X')$ be an idempotent. Since $\text{Hom}(X[-1], N) = 0$ (T is tilting), there exist $f \in \text{End}(N)$ and $g \in \text{End}(X)$ making the following diagram commute

$$\begin{array}{ccccccc} X' & \longrightarrow & N & \longrightarrow & X & \longrightarrow & X'[1] \\ e \downarrow & & f \downarrow & & g \downarrow & & e[1] \downarrow \\ X' & \longrightarrow & N & \longrightarrow & X & \longrightarrow & X'[1] \end{array}.$$

If g is invertible then so is f because $N \rightarrow X$ is a right minimal $\text{add}(T_1)$ -approximation; therefore e is invertible, and hence $e = 1_X$. If g is non-invertible then there exists $n \geq 1$ such that $g^n = 0$ because X is indecomposable; therefore the previous argument applies to $(1 - e^n = 1 - e, 1 - f^n, 1 - g^n = 1)$ instead of to the triple (e, f, g) ; it entails that $e = 0$. This proves that X' is indecomposable.

The functor $\text{Hom}(-, T_1)$ applies to the triangle $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ and gives an exact sequence

$$\text{Hom}(N, T_1) \rightarrow \text{Hom}(X', T_1) \rightarrow \text{Hom}(X[-1], T_1)$$

where the rightmost term is zero because T is tilting. Thus $X' \rightarrow N$ is a left $\text{add } T_1$ -approximation.

Finally let $u \in \text{End}(N)$ be such that the composite morphism $X' \rightarrow N \xrightarrow{u} N$ equals $X' \rightarrow N$. Therefore there exists $v \in \text{End}(X)$ such that the following diagram commutes for every $n \geq 1$

$$\begin{array}{ccccccc} X' & \longrightarrow & N & \longrightarrow & X & \longrightarrow & X'[1] \\ \parallel & & u^n \downarrow & & v^n \downarrow & & \parallel \\ X' & \longrightarrow & N & \longrightarrow & X & \longrightarrow & X'[1] \end{array}.$$

Since $N \in \text{add } T_1$ and $X \in \text{add } T_2$, the objects N and $X \oplus X'$ are not isomorphic. Therefore, the triangle $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ does not split, and hence $X \rightarrow X'[1]$ is nonzero. Therefore, $v^n \neq 0$ for every $n \geq 1$. Since X is indecomposable, it follows that $v: X \rightarrow X$ is an isomorphism, and hence so is u . Thus,

$X' \rightarrow N$ is left minimal. Therefore (i) \Rightarrow (ii). The proof of the converse implication is obtained using dual considerations.

Assume that (i) and (ii) hold true. Let \mathcal{C} be the full and convex subcategory of \mathcal{T} generated by $X[-1]$ and the indecomposable direct summands of N . Note that $X[-1] \rightarrow X'$ is nonzero as observed earlier. If $N = 0$ then $X' = X[-1]$, and hence $X' \in \mathcal{C}$. If $N \neq 0$ then for every indecomposable direct summand Z of N and for every retraction $N \rightarrow Z$ the composite morphism $X' \rightarrow N \rightarrow Z$ is nonzero ([17, Lemma 1.2, part (ii)]). These morphisms together with $X[-1] \rightarrow X'$ show that $X' \in \mathcal{C}$ \square

3.4. Starting (or ending) subcategories and tilting mutation. In view of Section 2, consider subcategories $\mathcal{A}, \mathcal{B} \subseteq \mathcal{T}$ such that T starts in $\mathcal{A} \vee \mathcal{B}$ and ends in $\mathcal{B}[\ell]$ or in $\mathcal{A}[\ell]$ (for some ℓ). A description of the subcategories in which T' starts and ends is made 3.4.1. And 3.4.2 and 3.4.3 concentrate on the situations where T starts in a one-parameter family of pairwise orthogonal tubes and ends in the suspension of it, or, more generally, in the ℓ -th suspension of it, respectively.

3.4.1. Let $\mathcal{A}, \mathcal{B} \subseteq \mathcal{T}$ be full, additive and convex subcategories stable under taking direct summands and such that

- (a) $\mathcal{T} = \bigvee_{i \in \mathbb{Z}} (\mathcal{A}[i] \vee \mathcal{B}[i])$,
- (b) $\text{Hom}(\mathcal{A}, \mathcal{A}[i]) = 0$ and $\text{Hom}(\mathcal{B}, \mathcal{B}[i]) = 0$ if $i \neq 0, 1$,
- (c) $\text{Hom}(\mathcal{A}, \mathcal{B}[i]) = 0$ if $i \neq 0$ and $\text{Hom}(\mathcal{B}, \mathcal{A}[i]) = 0$ if $i \neq 1$.

Typical examples used later on are as follows.

- (1) $\mathcal{A} = 0$ and $\mathcal{B} = \mathcal{H}$ (where $\mathcal{H} \subseteq \mathcal{T}$ is a hereditary abelian generating subcategory).
- (2) $\mathcal{A} = \mathcal{H}_+$ and $\mathcal{B} = \mathcal{H}_0$ (assuming additionally that \mathcal{H} arises from a weighted projective line).
- (3) $\mathcal{A} = \mathcal{H}_0[-1]$ and $\mathcal{B} = \mathcal{H}_+$ under the same additional assumption.
- (4) $\text{ind } \mathcal{A}$ consists of the transjective Auslander-Reiten component of \mathcal{T} intersecting \mathcal{H} , and $\text{ind } \mathcal{B} = (\text{ind } \mathcal{H} \setminus \text{ind } \mathcal{A})$ (assuming additionally that \mathcal{H} arises from a hereditary algebra of wild representation type).

Lemma. *Under the above setup let $\ell \geq 1$ be an integer. Assume that T starts in $\mathcal{A} \vee \mathcal{B}$ and ends in $\mathcal{B}[\ell]$. Let $T = T_1 \oplus T_2$ be the direct sum decomposition such that $T_1 \in \mathcal{A} \vee \mathcal{B} \vee \dots \vee \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$ and $T_2 \in \mathcal{A}[\ell] \vee \mathcal{B}[\ell]$. Then T' starts in $\mathcal{A} \vee \mathcal{B}$ and ends in $\mathcal{B}[\ell - 1]$, and T and T' have the same indecomposable direct summands in $\mathcal{A} \vee \mathcal{B}$ when $l \geq 2$.*

Proof. Since $T' = T_1 \oplus T'_2$ it suffices to prove that $T'_2 \in \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$ and that T'_2 has at least one indecomposable direct summand in $\mathcal{B}[\ell - 1]$.

Let X' be an indecomposable direct summand of T'_2 . Let $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ be a triangle such that $X' \rightarrow N$ is a left minimal $\text{add } T_1$ -approximation. Then $N \rightarrow X$ is a right minimal $\text{add } T_1$ -approximation, X is an indecomposable direct summand of T_2 and X' lies in the full and convex subcategory of \mathcal{T} generated by $X[-1]$ and the indecomposable direct summands of N (3.3). On the one hand $X[-1] \in \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$ because $X \in \text{add } T_2$ and $T_2 \in \mathcal{A}[\ell] \vee \mathcal{B}[\ell]$. On the other hand $N \in \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$ because $N \rightarrow X$ is a right minimal $\text{add } T_1$ -approximation and because of the assumptions made on \mathcal{A} and \mathcal{B} . Therefore X' lies in the full and convex subcategory generated by $\mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$ which already is full and convex. Thus $X' \in \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$. This proves that $T'_2 \in \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$.

Let X be an indecomposable direct summand of T_2 such that $X \in \mathcal{B}[\ell]$. Let $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ be a triangle such that $N \rightarrow X$ is a right minimal $\text{add } T_1$ -approximation. Then X' is an indecomposable direct summand of T'_2 lying in the full and convex subcategory of \mathcal{T} generated by $X[-1]$ and the indecomposable direct summands of N (3.3). Repeating the arguments used earlier to prove that $T'_2 \in \mathcal{A}[\ell - 1] \vee \mathcal{B}[\ell - 1]$ and using that $X \in \mathcal{B}[\ell]$ entails that $N \in \mathcal{B}[\ell - 1]$ and hence $X' \in \mathcal{B}[\ell - 1]$ (by assumption, $\text{add}(N) \cap \mathcal{A}[\ell] = \emptyset$). This proves that T'_2 has at least one indecomposable direct summand in $\mathcal{B}[\ell - 1]$. \square

3.4.2. The situation where there exists a hereditary abelian generating subcategory $\mathcal{H} \subseteq \mathcal{T}$ arising from a weighted projective line and such that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[\ell]$ for some integer ℓ needs careful consideration. Indeed the lower bound that is relevant to this situation is 2.2, part (2). It requires a tube in \mathcal{H}_0 containing an indecomposable direct summand of T and such that its ℓ -th suspension also contains an indecomposable direct summand of T . This crucial fact is proved in 4.2.2 and 4.2.3. As a preparation, the following result (when $\ell = 1$) and the next one (when $\ell \geq 2$) explain how these requirements are preserved under tilting mutation.

Lemma. *Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory arising from a weighted projective line. Let $\mathcal{U} \subseteq \mathcal{H}_0$ be a tube. Assume that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[1]$, and that T has at least one indecomposable direct summand in \mathcal{U} and at least one indecomposable direct summand in $\mathcal{H}_0[1] \setminus \mathcal{U}[1]$. Let $T = T_1 \oplus T_2$ be the direct sum decomposition such that $T_2 \in \text{add}\mathcal{U}[1]$ and T_1 has no indecomposable direct summand in $\mathcal{U}[1]$. Then*

- (1) T' starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[1]$,
- (2) for every tube $\mathcal{V} \subseteq \mathcal{H}_0$ there exists an indecomposable direct summand of T' in \mathcal{V} if and only if there exists an indecomposable direct summand of T in \mathcal{V} ,
- (3) for every tube $\mathcal{V} \subseteq \mathcal{H}_0$ there exists an indecomposable direct summand of T' in $\mathcal{V}[1]$ if and only if $\mathcal{V} \neq \mathcal{U}$ and there exists an indecomposable direct summand of T in $\mathcal{V}[1]$.

Proof. Let X' be an indecomposable direct summand of T'_2 . It is sufficient to prove that either $X' \in \mathcal{U}$ or else $X' \in \mathcal{H}_+[1]$. Let $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ be a triangle such that $X' \rightarrow N$ is a left minimal $\text{add}T_1$ -approximation. Then X is an indecomposable direct summand of T_2 , the morphism $N \rightarrow X$ is a right minimal $\text{add}T_1$ -approximation and X' lies in the full and convex subcategory of \mathcal{T} generated by $X[-1]$ and the indecomposable direct summands of N (3.3). In particular $X[-1] \in \mathcal{U}$ because $T_2 \in \text{add}\mathcal{U}[1]$. Therefore in order to prove that $X' \in \mathcal{U}$ or $X' \in \mathcal{H}_+[1]$ it is sufficient to prove that $N \in \mathcal{U} \vee \mathcal{H}_+[1]$. Let Z be an indecomposable direct summand of N . Then $\text{Hom}(Z, X) \neq 0$ because $N \rightarrow X$ is right minimal. Moreover $X \in \mathcal{U}[1]$, $Z \in \text{add}T_1$, the indecomposable direct summands of T_1 lie either in \mathcal{H}_0 , or in $\mathcal{H}_+[1]$ or in $\mathcal{H}_0[1] \setminus \mathcal{U}[1]$ and, finally, the tubes in \mathcal{H}_0 are orthogonal. This implies that $Z \in \mathcal{U}$ or $Z \in \mathcal{H}_+[1]$. \square

3.4.3.

Lemma. *Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory arising from a weighted projective line. Let $\ell \geq 2$ be an integer. Assume that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[\ell]$. Let $T = T_1 \oplus T_2$ be the direct sum decomposition such that $T_1 \in \mathcal{H}_0 \vee \mathcal{H}_+[1] \vee \cdots \vee \mathcal{H}_0[\ell - 2] \vee \mathcal{H}_+[\ell - 1] \vee \mathcal{H}_0[\ell - 1]$ and $T_2 \in \mathcal{H}_+[\ell] \vee \mathcal{H}_0[\ell]$. Then*

- (1) T' starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[\ell - 1]$,
- (2) T and T' have the same indecomposable direct summands in \mathcal{H}_0 ,
- (3) for every tube $\mathcal{U} \subseteq \mathcal{H}_0$, if T has an indecomposable direct summand in $\mathcal{U}[\ell]$ then T' has an indecomposable direct summand in $\mathcal{U}[\ell - 1]$.

Proof. (1) and (2) follow from 3.4.1 applied with $\mathcal{A} = \mathcal{H}_+$ and $\mathcal{B} = \mathcal{H}_0$.

(3) Let $\mathcal{U} \subseteq \mathcal{H}_0$ be a tube. Let X be an indecomposable direct summand of T lying in $\mathcal{U}[\ell]$. In particular X is an indecomposable direct summand of T_2 . Let $X' \rightarrow N \rightarrow X \rightarrow X'[1]$ be a triangle such that $N \rightarrow X$ is a right minimal $\text{add}T_1$ -approximation. It follows from 3.3 that X' is an indecomposable direct summand of T'_2 and that X' lies in the full and convex subcategory of \mathcal{T} generated by $X[-1]$ and the indecomposable direct summands of N . In view of proving (3) it is therefore sufficient to prove that $N \in \text{add}\mathcal{U}[\ell - 1]$. By the choice made for the decomposition $T = T_1 \oplus T_2$ and since $X \in \mathcal{H}_0[\ell]$ the indecomposable direct summands of T_1 from which there exists a nonzero morphism to X all lie in $\mathcal{H}_0[\ell - 1]$. This forces $N \in \mathcal{H}_0[\ell - 1]$ because $N \rightarrow X$ is a right minimal $\text{add}T_1$ -approximation. Moreover

since $X \in \mathcal{U}[\ell]$ and since \mathcal{H}_0 consists of pairwise orthogonal tubes, the indecomposable objects in $\mathcal{H}_0[\ell-1]$ from which there exists a nonzero morphism to X all lie in $\mathcal{U}[\ell-1]$. Therefore $N \in \text{add}\mathcal{U}[\ell-1]$. \square

4. INDECOMPOSABLE DIRECT SUMMANDS OF TILTING OBJECTS IN THE AUSLANDER-REITEN QUIVER

Let $T \in \mathcal{T}$ be a tilting object. This section aims at giving important information on the position of certain indecomposable direct summands of T in view of determining $\text{s.gl.dim. End}(T)^{\text{op}}$. Recall that any Auslander-Reiten component of \mathcal{T} has one of the following shapes: transjective Auslander-Reiten component, tube or $\mathbb{Z}A_\infty$. This section studies two particular situations, each of which is studied in a separate subsection: when T starts in a transjective component, and when T starts in a one-parameter family of pairwise orthogonal tubes. In each one of these situations a particular hereditary abelian generating subcategory of \mathcal{T} appears to be determined by T . This will be crucial to prove Theorem 1 and Theorem 2.

4.1. When T starts in a transjective component.

Proposition. *Let $T \in \mathcal{T}$ be a tilting object and $\Gamma \subseteq \mathcal{T}$ be a transjective Auslander-Reiten component. Assume that T starts in $\text{add}\Gamma$. Then there exists a slice Σ in Γ such that every source of Σ is an indecomposable direct summand of T , and for every indecomposable direct summand Y of T lying in Γ there exists a path in Γ with source in Σ and target Y .*

Proof. Since T starts in $\text{add}\Gamma$ there exist indecomposable summands S_1, \dots, S_n of T lying in Γ such that $\text{Hom}(\oplus_{i=1}^n S_i, X) \neq 0$ for every indecomposable direct summand X of T lying in Γ , and such that $\text{Hom}(S_i, S_j) = 0$ if $i \neq j$. Let Σ be the full subquiver of Γ the vertices of which are those $X \in \Gamma$ such that X is the successor in Γ of at least one of S_1, \dots, S_n , and such that any path in Γ from any of S_1, \dots, S_n to X is sectional.

By definition, Σ is a convex subquiver of Γ intersecting each τ -orbit at most once. Since Γ is a transjective Auslander-Reiten component there exists $n \in \mathbb{Z}$ such that $\tau^n X$ is a successor in Γ of one of the vertices in Σ , and $\tau^{n+1} X$ is the successor in Γ of none of the vertices in Σ . Consider any path in Γ

$$(\gamma) \quad S_i \rightarrow L_1 \rightarrow L_2 \rightarrow \cdots \rightarrow L_r = \tau^n X$$

from one of S_1, \dots, S_n to $\tau^n X$. If (γ) were not sectional there would exist some hook

$$L_{t-1} \rightarrow L_t \rightarrow L_{t+1} = \tau^{-1} L_{t-1},$$

and hence a path in Γ

$$S_i \rightarrow L_1 \rightarrow L_2 \rightarrow \cdots \rightarrow L_{t-1} = \tau L_{t+1} \rightarrow \tau L_{t+2} \rightarrow \cdots \rightarrow \tau L_r = \tau^{n+1} X$$

which would contradict the definition of n . The path (γ) is therefore sectional. This proves that Σ is a slice in Γ fitting the requirements of the proposition. \square


4.2. When T starts in a one-parameter family of pairwise orthogonal tubes. Like when T starts in a transjective component of \mathcal{T} , there are relevant hereditary abelian generating subcategories associated with T when it starts in a one-parameter family of pairwise orthogonal tubes. They arise from weighted projective lines and there are two cases to distinguish according to whether T ends in (a suitable suspension of) the subcategory of torsion objects or of torsion-free objects, respectively. The former case is dealt with in 4.2.2 and 4.2.3. The latter case is dealt with in 4.2.4. In both cases it appears that T cannot end in a maximal convex family of $\mathbb{Z}A_\infty$ components (4.2.1).

4.2.1. When T starts in a one-parameter family of pairwise orthogonal tubes it cannot end in a maximal convex family of $\mathbb{Z}A_\infty$ components.

Lemma. Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory equivalent to the category of coherent sheaves over a weighted projective line. Assume that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_+[\ell]$ for some integer ℓ . Then the weighted projective line has nonnegative Euler characteristic, equivalently \mathcal{H}_+ does not consist of $\mathbb{Z}A_\infty$ components.

Proof. Note that it is necessary that $\ell \geq 1$ because T starts in \mathcal{H}_0 . Thus $T \in \mathcal{H}_0 \vee \mathcal{H}_+[1] \vee \mathcal{H}_0[1] \vee \cdots \vee \mathcal{H}_+[\ell]$. Assume by contradiction that \mathcal{H}_+ consists of $\mathbb{Z}A_\infty$ components. A contradiction is obtained by induction on $\ell \geq 1$.

Assume that $\ell = 1$. In particular $T \in \mathcal{H}_0 \vee \mathcal{H}_+[1]$ and T has at least an indecomposable direct summand in \mathcal{H}_0 and in $\mathcal{H}_+[1]$ respectively. Then $\text{End}(T)^{\text{op}}$ is not quasi-tilted for, otherwise, there would exist a hereditary abelian generating subcategory $\mathcal{H}' \subseteq \mathcal{T}$ such that $T \in \mathcal{H}'$; since $\text{End}(T)^{\text{op}}$ is a connected algebra there would therefore exist a tube $\mathcal{U} \subseteq \mathcal{H}_0$ and a $\mathbb{Z}A_\infty$ component $\mathcal{V} \subseteq \mathcal{H}_+[1]$ such that $\mathcal{U}, \mathcal{V} \subseteq \mathcal{H}'$ and $\text{Hom}(\mathcal{U}, \mathcal{V}) \neq 0$, which is impossible. Then, it follows from [17, Proposition 3.3] and 1.2 that $\text{s.gl.dim. End}(T)^{\text{op}} = 3$. The picture below shows the subcategories of \mathcal{T} containing indecomposable direct summands of T (\circ) and $T[3]$ (\bullet).



Following in (1) in A.2, there is no indecomposable $X \in \mathcal{T}$ such that both $\text{Hom}(T, X)$ and $\text{Hom}(X, T[3])$ vanish. This contradicts $\text{s.gl.dim. End}(T)^{\text{op}} = 3$.

Now assume that $\ell \geq 2$. Let $T = T_1 \oplus T_2$ be the direct sum decomposition such that $T_1 \in \mathcal{H}_0 \vee \mathcal{H}_+[1]$ and $T_2 \in \mathcal{H}_0[1] \vee \mathcal{H}_+[2] \vee \cdots \vee \mathcal{H}_+[\ell]$. Let $T_1 \rightarrow M \rightarrow T_1' \rightarrow T[1]$ be a triangle where $T_1 \rightarrow M$ is a minimal left add T_2 -approximation. The dual versions of 3.1 and 3.4.1 show that $T_1' \oplus T_2$ is tilting, lies in $\mathcal{H}_0[1] \vee \mathcal{H}_+[2] \vee \cdots \vee \mathcal{H}_+[\ell]$ and has indecomposable direct summands in $\mathcal{H}_0[1]$ and in $\mathcal{H}_+[\ell]$ respectively. This is impossible by the induction hypothesis. \square

4.2.2. When $\mathcal{H} \subseteq \mathcal{T}$ is a hereditary abelian generating subcategory arising from a weighted projective line and such that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[1]$, the following lemma gives information on the indecomposable direct summands of T lying in \mathcal{H}_0 or in $\mathcal{H}_0[1]$.


Lemma. Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory arising from a weighted projective line. Assume that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[1]$. Then

- (1) there is no hereditary abelian generating subcategory of \mathcal{T} which contains T ,
- (2) $\text{s.gl.dim. End}(T)^{\text{op}} = 3$,
- (3) there exists a tube $\mathcal{U} \subseteq \mathcal{H}_0$ such that
 - (a) every indecomposable direct summand of T lying in \mathcal{H}_0 lies in \mathcal{U} ,
 - (b) every indecomposable direct summand of T lying in $\mathcal{H}_0[1]$ lies in $\mathcal{U}[1]$.

Proof. (1) Proceed by absurd and assume that there exists a hereditary abelian generating subcategory $\mathcal{H}' \subseteq \mathcal{T}$ such that $T \in \mathcal{H}'$. There are two cases to distinguish according to whether \mathcal{H}' is equivalent to a module category or not.

Assume first that \mathcal{H}' is equivalent to $\text{mod } H$ for some finite-dimensional hereditary algebra H . Since $\mathcal{T} \simeq \mathcal{D}^b(\mathcal{H}) \simeq \mathcal{D}^b(\mathcal{H}')$ it follows that H is of tame representation type. Since moreover T has indecomposable direct summands in \mathcal{H}_0 and $T \in \mathcal{H}' = \text{mod } H$ it follows that $\text{ind } \mathcal{H}_+$ consists of the transjective

Auslander-Reiten component of \mathcal{T} containing the indecomposable projective H -modules, and \mathcal{H}_0 consists of direct sums of indecomposable regular H -modules.



Therefore $T \in \mathcal{H}'$ whereas it ends in $\mathcal{H}_0[1] \subseteq \mathcal{H}'[1]$. This is absurd.

Assume next that \mathcal{H}' arises from a weighted projective line. Again there are two cases to distinguish according to whether \mathcal{H}'_+ consists of tubes or not.

If \mathcal{H}'_+ consists of tubes then \mathcal{H}' arises from a weighted projective line with vanishing Euler characteristic. Let $\mathcal{U} \subseteq \mathcal{H}_0$ be a tube containing an indecomposable direct summand of T . Then $\mathcal{U} \subseteq \mathcal{H}'$ because $T \in \mathcal{H}'$. Therefore there exists $q \in \mathbb{Q} \cup \{\infty\}$ such that $\mathcal{U} \subseteq \mathcal{H}'^{(q)}$. In other words $\mathcal{U} \subseteq (\mathcal{H}'\langle q \rangle)_0$ (see A.1). Applying A.4 to \mathcal{H} and $\mathcal{H}'\langle q \rangle$ entails that $\mathcal{H} = \mathcal{H}'\langle q \rangle$, and hence $\mathcal{H}_0 = \mathcal{H}'^{(q)}$. Consequently $T \in \mathcal{H}'$ whereas T has at least one indecomposable direct summand in $\mathcal{H}_0[1] = \mathcal{H}'^{(q)}[1] \subseteq \mathcal{H}'[1]$. This is absurd.

There only remains to treat the case where \mathcal{H}_+ does not consist of tubes, and hence contains no tube. Since \mathcal{U} is a tube containing an indecomposable direct summand of T and since $T \in \mathcal{H}'$ it follows that $\mathcal{U} \subseteq \mathcal{H}'_0$. Once again, applying A.4 to \mathcal{H} and \mathcal{H}' entails that $\mathcal{H} = \mathcal{H}'$. As observed previously this leads to a contradiction since T ends in $\mathcal{H}_0[1]$ and $T \in \mathcal{H}'$.

(2) It follows from 1.2 that $\text{s.gl.dim. End}(T)^{\text{op}} \leq 3$. Moreover (1) implies that $\text{End}(T)^{\text{op}}$ is not quasi-tilted, and hence $\text{s.gl.dim. End}(T)^{\text{op}} \geq 3$ ([17, Proposition 3.3]).

(3) It is necessary to prove first that there exists a tube $\mathcal{U} \subseteq \mathcal{H}_0$ such that both \mathcal{U} and $\mathcal{U}[1]$ contain an indecomposable direct summand of T . Since $\text{s.gl.dim. End}(T)^{\text{op}} = 3$, there exists an indecomposable $X \in \mathcal{T}$ such that $\text{Hom}(T, X) \neq 0$ and $\text{Hom}(X, T[3]) \neq 0$. Since $T \in \mathcal{H}_0 \vee \mathcal{H}_+[1] \vee \mathcal{H}_0[1]$, it follows that $X \in \mathcal{H}_0[2]$ and there exists indecomposable direct summands Y, Z of T such that $Z \in \mathcal{H}_0[1]$, $\text{Hom}(Z, X) \neq 0$, $Y \in \mathcal{H}_0$ and $\text{Hom}(X, Y[3]) \neq 0$ (see A.2 and picture below where the other possible positions of the indecomposable direct summands of T or $T[3]$ are marked with \circ or \bullet , respectively)

$\mathcal{H}_0[0]$	$\mathcal{H}_+[1]$	$\mathcal{H}_0[1]$	$\mathcal{H}_+[2]$	$\mathcal{H}_0[2]$	$\mathcal{H}_+[3]$	$\mathcal{H}_0[3]$	$\mathcal{H}_+[4]$	$\mathcal{H}_0[4]$
\circ	\circ	Z		X		$Y[3]$	\bullet	\bullet

Since \mathcal{H}_0 consists of pairwise orthogonal tubes, there exists a tube $\mathcal{U} \subseteq \mathcal{H}_0$ such that $Z \in \mathcal{U}[1]$, $X \in \mathcal{U}[2]$ and $Y[3] \in \mathcal{U}[3]$. In particular $Y \in \mathcal{U}$ and $Z \in \mathcal{U}[1]$. This proves the claim. In other words if $\mathcal{E}(T)$ denotes the set of those tubes $\mathcal{U} \subseteq \mathcal{H}_0$ such that each one of \mathcal{U} and $\mathcal{U}[1]$ contains an indecomposable direct summand of T then $\mathcal{E}(T) \neq \emptyset$.

Next it useful to prove that $\mathcal{E}(T)$ consists of a single tube which contains all indecomposable direct summands of T lying in \mathcal{H}_0 . Applying 3.4.2 to T and repeating the application for every tube lying in $\mathcal{E}(T) \setminus \{\mathcal{U}\}$ eventually yields a tilting object $S \in \mathcal{T}$ such that

- S starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[1]$,
- $\mathcal{E}(S)$ consists of a single tube \mathcal{U} ,
- every indecomposable direct summand of S lying in $\mathcal{H}_0[1]$ lies in $\mathcal{U}[1]$,

- for every tube $\mathcal{V} \subseteq \mathcal{H}_0$ there exists an indecomposable direct summand of T lying in \mathcal{V} if and only if the same holds true for S .

It is not possible for S to have any indecomposable direct summand lying in $\mathcal{H}_0 \setminus \mathcal{U}$ for, otherwise, the dual version of 3.4.2 could apply to S and $\mathcal{U}[1]$ and yield a tilting object $S' \in \mathcal{T}$ starting in \mathcal{H}_0 , ending in $\mathcal{H}_0[1]$ and such that $\mathcal{E}(S') = \emptyset$. Therefore every indecomposable direct summand of S lying in \mathcal{H}_0 (and hence every indecomposable direct summand of T lying in \mathcal{H}_0) lies in \mathcal{U} . In particular $\mathcal{E}(T) = \{\mathcal{U}\}$.

Finally it is not possible for T to have any indecomposable direct summand in $\mathcal{H}_0[1] \setminus \mathcal{U}[1]$ for, otherwise, 3.4.2 could apply to T and \mathcal{U} , and yield a tilting object $T' \in \mathcal{T}$ starting in \mathcal{H}_0 , ending in $\mathcal{H}_0[1]$ and such that $\mathcal{E}(T') = \emptyset$. This proves (3). \square

4.2.3. The previous result extends as follows when T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[\ell]$ for a hereditary abelian generating subcategory $\mathcal{H} \subseteq \mathcal{T}$ arising from a weighted projective line and for $\ell \geq 1$.

Proposition. *Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory arising from a weighted projective line. Assume that T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[\ell]$ for some integer $\ell \geq 0$. Then there exists a tube $\mathcal{U} \subseteq \mathcal{H}_0$ such that*

- (a) \mathcal{U} contains every indecomposable direct summand of T lying in \mathcal{H}_0 ,
- (b) $\mathcal{U}[\ell]$ contains every indecomposable direct summand of T lying in $\mathcal{H}_0[\ell]$.

In particular, if $\ell = 0$ then \mathcal{U} contains every indecomposable direct summand of T .

Proof. When $\ell = 0$ the hypotheses entail that $T \in \mathcal{H}_0$. The conclusion then follows from the fact that \mathcal{H}_0 consists of pairwise orthogonal tubes.

The general case proceeds by induction on $\ell \geq 1$. The case $\ell = 1$ is dealt with using 4.2.2 so assume that $\ell \geq 2$. Let $T' \in \mathcal{T}$ be the tilting object introduced in 3.4.3. Therefore the induction hypothesis applies to T' . Let $\mathcal{U} \subseteq \mathcal{H}_0$ be the tube such that \mathcal{U} (or $\mathcal{U}[\ell]$) contains every indecomposable direct summand of T' lying in \mathcal{H}_0 (or in $\mathcal{H}_0[\ell - 1]$, respectively). Then, it follows from 3.4.3, parts (2) and (3), and from the fact that T ends in $\mathcal{H}_0[\ell]$ that the conclusion of the proposition holds true for T \square

4.2.4. When T starts in a one-parameter family of pairwise orthogonal tubes but does not end in a suitable suspension of that family (unlike 4.2.3), then there still exists a relevant hereditary abelian generating subcategory associated with T as explained in the following result.

Proposition. *If T starts in a one-parameter family of pairwise orthogonal tubes and does not end in a transjective component, then there exists a hereditary abelian generating subcategory $\mathcal{H} \subseteq \mathcal{T}$ arising from a weighted projective line and there exists an integer $\ell \geq 0$ such that*

- (a) T starts in a one-parameter family of pairwise orthogonal tubes contained in \mathcal{H} , and
- (b) T ends in $\mathcal{H}_0[\ell]$.

Proof. It follows from 4.2.1 that T ends in one-parameter family of pairwise orthogonal tubes. Moreover A.5 shows that there exists a hereditary abelian generating subcategory $\mathcal{H}' \subseteq \mathcal{T}$ arising from a weighted projective line and such that \mathcal{H}'_0 is that family. Let $\ell \geq 0$ be the integer such that T starts in $\mathcal{H}'[-\ell]$. Let $\mathcal{H} = \mathcal{H}'[-\ell]$. Then \mathcal{H} fits the conclusion of the proposition. \square

5. THE STRONG GLOBAL DIMENSION THROUGH AUSLANDER-REITEN THEORY

Let $T \in \mathcal{T}$ be a tilting object. The objective of this section is to determine $\text{s.gl.dim. End}(T)^{\text{op}}$ in terms of the position of the indecomposable direct summands of T in the Auslander-Reiten quiver of T . Recall that T may start either in a transjective component, or in a one-parameter family of pairwise orthogonal tubes, or in a maximal convex family of $\mathbb{Z}A_\infty$ components. The three following subsections therefore treat each one of these cases separately. The situation where T ends in a transjective component is dual

to that where T starts in a transjective component. Thus when assuming that T does not start in a transjective component it may be assumed also that T does not end in a transjective component.

5.1. When T starts in a transjective component.

Proposition. *Let $T \in \mathcal{T}$ be a tilting object. Assume that T starts in a transjective component Γ . Let Σ be the slice introduced in 4.1.*

- (1) *Let $\mathcal{H} = \{X \in \mathcal{T} \mid (\forall S \in \Sigma) (\forall i \neq 0) \operatorname{Hom}(S, X[i]) = 0\}$. Then \mathcal{H} is a hereditary abelian generating subcategory. Moreover there exists an integer $\ell \geq 0$ such that $T \in \bigvee_{i=0}^{\ell} \mathcal{H}[i]$ and such that T has an indecomposable summand in \mathcal{H} and in $\mathcal{H}[\ell]$;*
- (2) *If $\operatorname{End}(T)^{\operatorname{op}}$ is not a hereditary algebra then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} = \ell + 2$.*

Proof. (1) The first assertion follows from the fact that Σ is a slice in Γ . In particular $\mathcal{T} = \bigvee_{i \in \mathbb{Z}} \mathcal{H}[i]$. The second assertion follows from the following facts: the indecomposable projectives in \mathcal{H} are, up to isomorphism, the objects in Σ ; and the sources of Σ are all summands of T .

(2) If $\ell = 0$ then $\operatorname{End}(T)^{\operatorname{op}}$ is quasi-tilted and not hereditary. Thus $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} = 2$ ([17, Prop. 3.3]). If $\ell \geq 1$ then the conclusion follows from 1.2 and 2.1. \square

5.2. When T starts and ends in a one-parameter family of pairwise orthogonal tubes.

Proposition. *Assume that T starts in a one-parameter family of pairwise orthogonal tubes and does not end in a transjective component. Let \mathcal{H}, ℓ be like in 4.2.4. Then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} = \ell + 2$.*

Proof. Assume first that T starts in \mathcal{H}_0 . Let $\mathcal{U} \subseteq \mathcal{H}_0$ be the tube obtained upon applying 4.2.3 to T . Since T starts in \mathcal{H}_0 and ends in $\mathcal{H}_0[\ell]$ it follows that $T \in \mathcal{H} \vee \mathcal{H}[1] \vee \cdots \vee \mathcal{H}[\ell]$. Therefore $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \leq \ell + 2$ (1.2). On the other hand it follows from 2.2 (part (2)) that $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$.

If T does not start in \mathcal{H}_0 then it starts in \mathcal{H}_+ . The arguments used in the previous case lead to the same conclusion provided that 2.2, part (3), is used instead of 2.2, part (2). \square

5.3. When T starts in a maximal convex family of $\mathbb{Z}A_{\infty}$ components.

Proposition. *Assume that T starts in a maximal convex family of $\mathbb{Z}A_{\infty}$ components and does not end in a transjective component. Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory such that T starts in \mathcal{H} . Let $\ell \geq 0$ be such that T ends in $\mathcal{H}[\ell]$. Then $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} = \ell + 2$.*

Proof. First, $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \leq \ell + 2$ due to 1.2. Let $M_0 \in \mathcal{H}$ be an indecomposable direct summand of T lying in an Auslander-Reiten component of shape $\mathbb{Z}A_{\infty}$. There are two cases to distinguish according to whether or not there exists an indecomposable direct summand M_1 of T lying in an Auslander-Reiten component of shape $\mathbb{Z}A_{\infty}$ contained in $\mathcal{H}[\ell]$. Assume first that this is indeed the case. According to 2.2 (part (4)) it follows that $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$, and hence $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} = \ell + 2$.

Otherwise, it is necessary that \mathcal{H} arises from a weighted projective line and that T ends in $\mathcal{H}_0[\ell]$. Let M_1 be an indecomposable direct summand of T lying in $\mathcal{H}_0[\ell]$. According to 2.2 (part (3)) it follows that $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} \geq \ell + 2$, and hence $\operatorname{s.gl.dim.} \operatorname{End}(T)^{\operatorname{op}} = \ell + 2$. \square

6. PROOFS OF THE MAIN THEOREMS

It is worth noticing that if a statement holds true for tilting objects starting in a transjective component then so does its dual statement for tilting objects ending in a transjective component. Hence all the possible situations (up to dualising) for a tilting object T are covered by the three following cases:

- (a) T starts in a transjective component, or
- (b) T starts in a one-parameter family of pairwise orthogonal tubes and does not end in a transjective component, or

- (c) T starts in a maximal convex family of $\mathbb{Z}A_\infty$ components and does not end in a transjective component.

6.1. Proof of Theorem 1.

Proof. Assertion (1) follows from 1.2. Assertion (2) follows from 5.1, 5.2 and 5.3. □

6.2. Proof of Theorem 2.

Proof. Only the last assertion needs a proof (see 3.2). Proceed by induction on $d = \text{s.gl.dim. End}(T)^{\text{op}}$ (≥ 2). If $d = 2$ then $\text{End}(T)^{\text{op}}$ is quasi-tilted, and hence there is nothing to prove. Assume that $d > 2$. Clearly, it suffices to show that there exists a tilting object $T' \in \mathcal{T}$ obtained from T by a tilting mutation and such that $\text{s.gl.dim. End}(T')^{\text{op}} = d - 1$. Let $\mathcal{H} \subseteq \mathcal{T}$ be a hereditary abelian category and $\ell \geq 0$ be an integer like in 5.1, 4.2.4 or 5.3 according to whether (a), (b) or (c) (as stated at the beginning of the section) holds true for T .

In either case T starts in \mathcal{H} and ends in $\mathcal{H}[\ell]$. It follows from 5.1, 5.2 and 5.3 that $d = \ell + 2$. Let $T = T_1 \oplus T_2$ be the direct sum decomposition such that $T_1 \in \bigvee_{i=0}^{\ell-1} \mathcal{H}[i]$ and $T_2 \in \mathcal{H}[\ell]$. Let $T'_2 \rightarrow M \rightarrow T_2 \rightarrow T'_2[1]$ be the triangle such that $M \rightarrow T_2$ is a right minimal add T_1 -approximation. Let $T' = T_1 \oplus T'_2$. Clearly $\text{Hom}(T_2, T_1) = 0$. Thus T' is a tilting object (3.1).

Applying 3.4.1 to $\mathcal{A} = \{0\}$ and $\mathcal{B} = \mathcal{H}$ shows that T' starts in \mathcal{H} and ends in $\mathcal{H}[\ell - 1]$. In particular if $\ell = 1$, that is $d = 3$, then $T' \in \mathcal{H}$; therefore $\text{s.gl.dim. End}(T')^{\text{op}} = 2 = d - 1$. From now on assume that $\ell \geq 2$. Therefore T' starts in \mathcal{H} and ends in $\mathcal{H}[\ell - 1]$, and T and T' have the same indecomposable direct summands lying in \mathcal{H} (3.4.1, with $\mathcal{A} = \{0\}$ and $\mathcal{B} = \mathcal{H}$). The rest of the proof distinguishes three cases according to situations (a), (b) and (c) listed earlier.

(a) Because of the conditions satisfied by T and T' , the triple $(\Sigma', \mathcal{H}', \ell')$ arising from 5.1 applied to T' is such that $\Sigma' = \Sigma$, $\mathcal{H}' = \mathcal{H}$, $\ell' = \ell - 1$, and $\text{s.gl.dim. End}(T')^{\text{op}} = d - 1$.

(b) Note that \mathcal{H} arises from a weighted projective line and, by assumption, T ends in $\mathcal{H}_0[\ell]$ (4.2.4). Applying 3.4.1 to $\mathcal{A} = \mathcal{H}_+$ and $\mathcal{B} = \mathcal{H}_0$ shows that T' starts in a one-parameter family of pairwise orthogonal tubes contained in \mathcal{H} and ends in $\mathcal{H}_0[\ell - 1]$. Hence, $\text{s.gl.dim. End}(T')^{\text{op}} = \ell + 1 = d - 1$ (5.2).

(c) Note that \mathcal{H} arises either from a weighted projective line with negative Euler characteristic, or else from a hereditary algebra of wild representation type. Apply 3.4.1 in situation (2) (or, in situation (4)) to the former case (or, to the latter case, respectively). Since $\ell \geq 2$ this shows that T' starts in a maximal convex family of $\mathbb{Z}A_\infty$ components contained in \mathcal{H} , that it ends in $\mathcal{H}[\ell - 1]$ and that it does not end in a transjective component. Therefore $\text{s.gl.dim. End}(T')^{\text{op}} = \ell + 1 = d - 1$ (5.3). □

APPENDIX A. MORPHISMS IN BOUNDED DERIVED CATEGORIES OF WEIGHTED PROJECTIVE LINES

This section collects some known results on hereditary abelian categories arising from weighted projective lines, and which are used in the proof of the main theorems in this text. Also it proves some useful technical facts on morphism spaces in the corresponding bounded derived categories.

Recall that like everywhere else in this text, by "hereditary abelian generating subcategory of \mathcal{T} " is meant a full subcategory $\mathcal{H}' \subseteq \mathcal{T}$ which is hereditary abelian and such that the embedding $\mathcal{H}' \hookrightarrow \mathcal{T}$ extends to a triangle equivalence $\mathcal{D}^b(\mathcal{H}') \simeq \mathcal{T}$.

A.1. **Reminder on $\text{coh}(\mathbb{X})$.** Let \mathbb{X} be a weighted projective line in the sense of Geigle-Lenzing [10]. Let $\mathcal{H} = \text{coh}(\mathbb{X})$ the category of coherent sheaves. This section collects some essential properties of \mathcal{H} used in this text (see [2, 12, 21, 23] and references therein for details). The full subcategory of \mathcal{H} formed by objects of finite length in \mathcal{H} (i.e. torsion sheaves on \mathbb{X}) is denoted by \mathcal{H}_0 . The full subcategory of \mathcal{H} of vector bundles (or, torsion-free sheaves) is denoted by \mathcal{H}_+ . Note that

$$(1) \quad \text{Hom}(\mathcal{H}_0, \mathcal{H}_+) = 0 \text{ and } \mathcal{H}_+ = \{X \in \mathcal{H} \mid \text{Hom}(\mathcal{H}_0, X) = 0\}.$$

To each nonzero vector bundle E is associated its rank $\text{rk}(E) \in \mathbb{N} \setminus \{0\}$. *Line bundles* are rank one vector bundles. Given any vector bundle E , its rank is r if and only if there exists a filtration $0 = E_0 \subset E_1 \subset \cdots \subset E_r = E$ in \mathcal{H} such that E_i/E_{i-1} is a line bundle for every i .

The category \mathcal{H} is abelian, hereditary, and Krull-Schmidt; it has Serre duality and Auslander-Reiten sequences. The main specificities of the Auslander-Reiten structure of \mathcal{H} used in this text depend on the Euler characteristic $\chi(\mathbb{X})$ ([2, Sect. 10] and [12, Sect. 4]).

- The indecomposable objects in \mathcal{H}_0 form a disjoint union of tubes in the Auslander-Reiten quiver of \mathcal{H} . This union is parametrised by \mathbb{X} .
- If $\chi(\mathbb{X}) > 0$ then the indecomposable objects in \mathcal{H}_+ form a single Auslander-Reiten component of shape $\mathbb{Z}\Delta$ where Δ is a graph of extended Dynkin type.
- If $\chi(\mathbb{X}) = 0$ then \mathcal{H}_+ decomposes as $\mathcal{H}_+ = \bigvee_{q \in \mathbb{Q}} \mathcal{H}^{(q)}$ where each $\mathcal{H}^{(q)}$ is a full subcategory of \mathcal{H} isomorphic to \mathcal{H}_0 . The subcategory \mathcal{H}_0 is also denoted by $\mathcal{H}^{(\infty)}$, and $\text{Hom}(\mathcal{H}^{(p)}, \mathcal{H}^{(q)}) = 0$ if $q < p \leq \infty$; in particular each $\mathcal{H}^{(p)}$ is a disjoint union of orthogonal tubes. For every $q \in \mathbb{Q}$, the category $\left(\bigvee_{r \in \mathbb{Q} \cup \{\infty\}, q < r} \mathcal{H}^{(r)}[-1] \vee \bigvee_{s \in \mathbb{Q}, s \leq q} \mathcal{H}^{(s)}\right)$ is denoted by $\mathcal{H}\langle q \rangle$. Recall that this is a hereditary abelian generating subcategory whose subcategory of finite length objects is $\mathcal{H}^{(q)}$. Also, $\mathcal{H}\langle \infty \rangle$ denotes \mathcal{H} .
- If $\chi(\mathbb{X}) < 0$ then the indecomposable objects in \mathcal{H}_+ form a disjoint union of Auslander-Reiten components of shape $\mathbb{Z}A_\infty$ in the Auslander-Reiten quiver of \mathcal{H} .

The following two properties on morphism spaces in \mathcal{H} play a fundamental role in this text:

- Let $L \in \mathcal{H}_+$ be a line bundle and let $\mathcal{U} \subseteq \mathcal{H}_0$ be a tube, then there exists a unique quasi-simple $S \in \mathcal{U}$ such that $\text{Hom}(L, S)$ is nonzero (and, moreover, is one dimensional).
- In the wild type case, given nonzero vector bundles $E, F \in \mathcal{H}_+$ it exists an integer n_0 such that $\text{Hom}(E, \tau^n F) \neq 0$ for every $n \geq n_0$.

A.2. **Paths in $\mathcal{D}^b(\text{coh}(\mathbb{X}))$.** Let \mathbb{X} be a weighted projective line. Let $\mathcal{H} = \text{coh}(\mathbb{X})$. Let $\mathcal{T} = \mathcal{D}^b(\mathcal{H})$. The following result is useful to investigate s.gl.dim. of endomorphism algebras of tilting objects in \mathcal{T} .

Lemma. (1) *Let $E \in \mathcal{H}_+$ be indecomposable. Let $\mathcal{U} \subseteq \mathcal{H}_0$ be a tube. Then there exists $S \in \mathcal{U}$ quasi-simple such that $\text{Hom}(E, S) \neq 0$ and $\text{Ext}^1(S, \tau E) \neq 0$. In particular $\text{Hom}(E, \mathcal{U}) \neq 0$ and $\text{Ext}^1(\mathcal{U}, E) \neq 0$. Moreover, if $S = X_0 \rightarrow \cdots \rightarrow X_n \rightarrow \cdots$ is the unique infinite sectional path in the Auslander-Reiten quiver of \mathcal{U} then $\text{Hom}(E, X_n) \neq 0$ and $\text{Ext}^1(X_n, \tau E) \neq 0$ for every $n \geq 0$.*

(2) *Let $\mathcal{U}, \mathcal{V} \subseteq \mathcal{H}_0$ be tubes. Let j be a positive integer. Then there exist quasi-simples $S \in \mathcal{U}$ and $S' \in \mathcal{V}$ together with indecomposable vector bundles $E, E' \in \mathcal{H}_+$ and a path of nonzero morphisms in $\text{ind } \mathcal{T}$*

$$S \rightarrow E[1] \rightarrow \cdots \rightarrow E'[j] \rightarrow S'[j].$$

Proof. (1) Using the filtration of E there exists a line bundle $L \in \mathcal{H}_+$ and an epimorphism $E \rightarrow L$ in \mathcal{H} . Then there exists a quasi-simple $S \in \mathcal{U}$ such that $\text{Hom}(L, S) \neq 0$. Taking a composite morphism $E \rightarrow L \rightarrow S$ shows that $\text{Hom}(E, S) \neq 0$. Thus $\text{Ext}^1(S, \tau E) \neq 0$ because of Serre duality.

Let $E \rightarrow S$ be a nonzero morphism. For every $n \in \mathbb{N}$ let $X_n \rightarrow X_{n+1}$ be an irreducible morphism. This is a monomorphism in \mathcal{H} because \mathcal{U} is a tube. This and the path $E \rightarrow S \rightarrow X_1 \rightarrow \cdots \rightarrow X_n$ show that $\text{Hom}(E, X_n) \neq 0$. Serre duality entails $\text{Ext}^1(X_n, \tau E) \neq 0$.

(2) The proof is an induction on j . Assume that $j = 1$. Let $L \in \mathcal{H}_+$ be any line bundle. Then there exist quasi-simples $S \in \mathcal{U}, S' \in \mathcal{V}$ such that $\text{Hom}(L, S') \neq 0$ and $\text{Ext}^1(S, L) \neq 0$. Whence the path $S \rightarrow L[1] \rightarrow S'[1]$. Assume that $j > 1$ and that (2) holds true for $j - 1$. By induction hypothesis and because of the case $j = 1$, there exist paths $S \rightsquigarrow S''[1]$ and $S''[1] \rightsquigarrow S'[j]$ in $\text{ind } \mathcal{T}$ for some quasi-simples $S, S'' \in \mathcal{U}$ and $S' \in \mathcal{V}$. Whence a path $S \rightsquigarrow S'[j]$. \square

A.3. Morphisms between Auslander-Reiten components in \mathcal{T} . In order to understand better the hereditary abelian generating subcategories of \mathcal{T} which contain indecomposable summands of a given tilting object in \mathcal{T} , it is useful to know whether or not there exists a nonzero morphism between two given Auslander-Reiten components in \mathcal{T} . Recall that every Auslander-Reiten component in \mathcal{H} is stable and $\mathcal{T} = \bigvee_{i \in \mathbb{Z}} (\mathcal{H}_+[i] \vee \mathcal{H}_0[i])$. Therefore any given Auslander-Reiten component of \mathcal{T} equals $\mathcal{U}[i]$ where $i \in \mathbb{Z}$ and \mathcal{U} is either a tube in \mathcal{H}_0 or else consists of objects in \mathcal{H}_+ . Note that if $\mathcal{U}, \mathcal{U}' \subseteq \mathcal{H}_0$ are distinct tubes then $\text{Hom}(\mathcal{U}, \mathcal{U}'[i]) = 0$ for every $i \in \mathbb{Z}$ because \mathcal{U} and \mathcal{U}' are orthogonal, because \mathcal{H} is hereditary and because of Serre duality. These considerations together with A.2, yield the following proposition where (2), (3) and (4) follow from (1).

Proposition. (1) Let $\mathcal{U} \subseteq \mathcal{H}_0$ and $\mathcal{V} \subseteq \mathcal{H}_+$ be Auslander-Reiten components. Let $i \in \mathbb{Z}$ then

- $\text{Hom}(\mathcal{U}, \mathcal{U}[i]) \neq 0 \Leftrightarrow i = 0$ or $i = 1$,
- $\text{Hom}(\mathcal{U}, \mathcal{V}[i]) \neq 0 \Leftrightarrow i = 1$,
- $\text{Hom}(\mathcal{V}, \mathcal{U}[i]) \neq 0 \Leftrightarrow i = 0$,
- $\text{Hom}(\mathcal{V}, \mathcal{V}[i]) \neq 0 \Leftrightarrow i = 0$ or $i = 1$.

(2) Both \mathcal{H}_0 and \mathcal{H}_+ are convex in \mathcal{T} .

(3) Let $\mathcal{U} \subseteq \mathcal{H}_0$ be a tube. Let \mathcal{V} be an Auslander-Reiten component of \mathcal{T} distinct from $\mathcal{U}[i]$ for every $i \in \mathbb{Z}$. Then: $\mathcal{V} \subseteq \bigvee_{\ell \in \mathbb{Z}} \mathcal{H}_0[\ell] \Leftrightarrow (\forall i \in \mathbb{Z}) \text{Hom}(\mathcal{U}, \mathcal{V}[i]) = 0 \Leftrightarrow (\forall i \in \mathbb{Z}) \text{Hom}(\mathcal{V}, \mathcal{U}[i]) = 0$.

(4) Let \mathcal{V} be an Auslander-Reiten component of \mathcal{T} then: $\mathcal{V} \subseteq \mathcal{H}_+ \Leftrightarrow (\forall i \neq 0) \text{Hom}(\mathcal{V}, \mathcal{H}_0[i]) = 0$.

A.4. A sufficient criterion for two hereditary abelian generating subcategories to coincide.

The proofs of the main theorems use the following general fact on tilting objects in \mathcal{T} : Let $T \in \mathcal{T}$ be a tilting object such that $T \in \mathcal{H}_0 \vee \mathcal{H}_+[1] \vee \mathcal{H}_0[1]$ and T has indecomposable summands both in \mathcal{H}_0 and $\mathcal{H}_0[1]$, then there is no hereditary abelian generating subcategory $\mathcal{H}' \subseteq \mathcal{T}$ such that $T \in \mathcal{H}'$ (4.2.2). The proof of this fact is based on the following result which gives a sufficient condition for two hereditary abelian generating subcategories $\mathcal{H}, \mathcal{H}' \subseteq \mathcal{T}$ to coincide.

Proposition. Let $\mathcal{H}' \subseteq \mathcal{T}$ be a hereditary abelian generating subcategory having no nonzero projective object. Assume that there exists a tube $\mathcal{U} \subseteq \mathcal{H}_0$ such that $\mathcal{U} \subseteq \mathcal{H}'_0$. Then $\mathcal{H} = \mathcal{H}'$.

Proof. Note that since \mathcal{H}' is not a module category it is the category of coherent sheaves over some weighted projective line, and since $\mathcal{D}^b(\mathcal{H}') \simeq \mathcal{D}^b(\mathcal{H})$ it follows that \mathcal{H}' and \mathcal{H} have isomorphic Auslander-Reiten quivers [12, Sect. 4]. Hence \mathcal{H} and \mathcal{H}' play symmetric roles in the proposition.

It is useful to prove first that $\mathcal{H}_0 = \mathcal{H}'_0$. Let $\mathcal{V} \subseteq \mathcal{H}_0$ be a tube distinct from \mathcal{U} . Hence $\mathcal{V} \neq \mathcal{U}[j]$ for every $j \in \mathbb{Z}$. Applying A.3 (part (3)) to $\mathcal{H}, \mathcal{U}, \mathcal{V}$, yields $\text{Hom}(\mathcal{V}, \mathcal{U}[j]) = 0$ for every $j \in \mathbb{Z}$. Then applying the same result to $\mathcal{H}', \mathcal{U}, \mathcal{V}$ entails that there exists $j \in \mathbb{Z}$ such that $\mathcal{V} \in \mathcal{H}'_0[j]$. By absurd assume that $j > 0$. Using A.2 (part (2)) applied to $\mathcal{H}', \mathcal{U}, \mathcal{V}[-j]$ gives a path $S \rightarrow E[1] \rightsquigarrow S'$ in $\text{ind } \mathcal{T}$ such that $S \in \mathcal{U}, S' \in \mathcal{V}$ are quasi-simple and $E \in \mathcal{H}'_+$. But $S, S' \in \mathcal{H}_0$ and \mathcal{H}_0 is convex in \mathcal{T} . Therefore $E[1] \in \mathcal{H}_0$. Now because $\text{ind } \mathcal{H}_0$ is a disjoint union of pairwise orthogonal tubes and since $S \rightarrow E[1]$ is a nonzero morphism with $S \in \mathcal{U} \subseteq \mathcal{H}_0$ and $E[1] \in \mathcal{H}_0$, it follows that $E[1] \in \mathcal{U}$, and hence $E[1] \in \mathcal{H}'_0$ (recall that $\mathcal{U} \subseteq \mathcal{H}'_0$). This contradicts $E \in \mathcal{H}'_+$. Therefore $j \leq 0$. Dually, the same arguments show that

$j \geq 0$. Thus $j = 0$, and therefore $\mathcal{V} \subseteq \mathcal{H}'_0$. This proves that $\mathcal{H}_0 \subseteq \mathcal{H}'_0$. And because of the symmetry between \mathcal{H} and \mathcal{H}' it follows that $\mathcal{H}_0 = \mathcal{H}'_0$.

There only remains to prove that $\mathcal{H}_+ = \mathcal{H}'_+$. Let \mathcal{V} be an Auslander-Reiten component of \mathcal{T} such that $\mathcal{V} \subseteq \mathcal{H}_+$. Using A.3 (part (4)) applied to $\mathcal{H}, \mathcal{U}, \mathcal{V}$ it follows that $\text{Hom}(\mathcal{V}, \mathcal{H}_0[i]) = 0$ for $i \neq 0$. Since $\mathcal{H}_0 = \mathcal{H}'_0$ the same result applied to $\mathcal{H}', \mathcal{U}, \mathcal{V}$ shows that $\mathcal{V} \subseteq \mathcal{H}'_+$. Therefore $\mathcal{H}_+ \subseteq \mathcal{H}'_+$, and hence $\mathcal{H}_+ = \mathcal{H}'_+$ by symmetry between \mathcal{H} and \mathcal{H}' . \square

A.5. Description of one-parameter families of pairwise orthogonal tubes.

Proposition. (1) *The subcategory $\mathcal{H}_0 \subseteq \mathcal{T}$ is a one-parameter family of tubes.*

(2) *Let $\mathcal{T}_0 \subseteq \mathcal{T}$ be a one-parameter family of tubes, then there exists a unique hereditary abelian generating subcategory $\mathcal{H}' \subseteq \mathcal{T}$ which is not a module category and such that $\mathcal{T}_0 = \mathcal{H}'_0$.*

Proof. (1) \mathcal{H}_0 is a disjoint union of pairwise orthogonal tubes (A.1) and it is convex in \mathcal{T} (A.3). Following A.2 (part (2)), if a full convex subcategory intersects \mathcal{H}_0 and $\bigvee_{j \neq 0} \mathcal{H}_0[j]$ nontrivially, then it intersects $\bigvee_{j \in \mathbb{Z}} \mathcal{H}_+[j]$ nontrivially. Therefore \mathcal{H}_0 is maximal as a family of pairwise orthogonal tubes.

(2) The uniqueness follows from A.4. Let $\mathcal{U} \subseteq \mathcal{T}_0$ be a tube. Let $\ell \in \mathbb{Z}$ be such that $\mathcal{U} \subseteq \mathcal{H}[\ell]$. If $\chi(\mathbb{X}) \neq 0$, then $\mathcal{U} \subseteq (\mathcal{H}[\ell])_0 = \mathcal{H}_0[\ell]$ (see A.1), and hence $\mathcal{U} \subseteq \mathcal{H}'_0$ where \mathcal{H}' denotes $\mathcal{H}[\ell]$. If $\chi(\mathbb{X}) = 0$, then there exists $q \in \mathbb{Q} \cup \{\infty\}$ such that $\mathcal{U} \subseteq (\mathcal{H}[\ell])^{(q)}$, and hence $\mathcal{U} \subseteq \mathcal{H}'_0$ where \mathcal{H}' denotes $\mathcal{H}[\ell]\langle q \rangle$.

Let $\mathcal{V} \subseteq \mathcal{T}_0$ be a tube distinct from \mathcal{U} . By assumption on \mathcal{T}_0 , there exists $t \in \mathbb{Z}$ such that $\mathcal{V} \subseteq \mathcal{H}'_0[t]$ (A.3, part (3)). Should t be nonzero, \mathcal{T}_0 would intersect $\bigvee_{\ell \in \mathbb{Z}} \mathcal{H}'_+[\ell]$ because it is convex in \mathcal{T} (A.2, part (2)). This is impossible because the Auslander-Reiten components of \mathcal{T}_0 are pairwise orthogonal. Thus, $t = 0$, and hence $\mathcal{V} \subseteq \mathcal{H}'_0$. Therefore, $\mathcal{T}_0 \subseteq \mathcal{H}'_0$, and hence $\mathcal{T}_0 = \mathcal{H}'_0$. \square

ACKNOWLEDGEMENTS

The work presented in this text was done while the second named author was associated professor at Université Blaise Pascal and during visits to the first and third named authors. He would like to thank the members of the Department of Mathematics at Université Blaise Pascal as well as the first and third named authors for their warm hospitality.

REFERENCES

- [1] Takuma Aihara and Osamu Iyama. Silting mutation in triangulated categories. *J. Lond. Math. Soc. (2)*, 85(3):633–668, 2012.
- [2] Lidia Angeleri Hügel, Dieter Happel, and Henning Krause, editors. *Handbook of tilting theory*, volume 332 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2007.
- [3] Lidia Angeleri Hügel, Steffen Koenig, and Qunhua Liu. Jordan-Hölder theorems for derived module categories of piecewise hereditary algebras. *J. Algebra*, 352:361–381, 2012.
- [4] Ibrahim Assem, Thomas Brüstle, and Ralf Schiffler. Cluster-tilted algebras as trivial extensions. *Bull. Lond. Math. Soc.*, 40(1):151–162, 2008.
- [5] Ibrahim Assem, Daniel Simson, and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 1*, volume 65 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2006. Techniques of representation theory.
- [6] Ibrahim Assem and Andrzej Skowroński. Algebras with cycle-finite derived categories. *Math. Ann.*, 280(3):441–463, 1988.
- [7] Ibrahim Assem and Andrzej Skowroński. On tame repetitive algebras. *Fund. Math.*, 142(1):59–84, 1993.
- [8] Aslak Bakke Buan, Robert Marsh, Markus Reineke, Idun Reiten, and Gordana Todorov. Tilting theory and cluster combinatorics. *Adv. Math.*, 204(2):572–618, 2006.
- [9] Piotr Dowbor and Andrzej Skowroński. On Galois coverings of tame algebras. *Arch. Math. (Basel)*, 44(6):522–529, 1985.
- [10] Werner Geigle and Helmut Lenzing. A class of weighted projective curves arising in representation theory of finite-dimensional algebras. In *Singularities, representation of algebras, and vector bundles (Lambrecht, 1985)*, volume 1273 of *Lecture Notes in Math.*, pages 265–297. Springer, Berlin, 1987.

- [11] Dieter Happel. *Triangulated categories in the representation theory of finite-dimensional algebras*, volume 119 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1988.
- [12] Dieter Happel. Quasitilted algebras. In *Algebras and modules, I (Trondheim, 1996)*, volume 23 of *CMS Conf. Proc.*, pages 55–82. Amer. Math. Soc., Providence, RI, 1998.
- [13] Dieter Happel. A characterization of hereditary categories with tilting object. *Invent. Math.*, 144(2):381–398, 2001.
- [14] Dieter Happel, Idun Reiten, and Sverre O. Smalø. Tilting in abelian categories and quasitilted algebras. *Mem. Amer. Math. Soc.*, 120(575):viii+ 88, 1996.
- [15] Dieter Happel, Jeremy Rickard, and Aidan Schofield. Piecewise hereditary algebras. *Bull. London Math. Soc.*, 20(1):23–28, 1988.
- [16] Dieter Happel and Claus Michael Ringel. Tilted algebras. *Trans. Amer. Math. Soc.*, 274(2):399–443, 1982.
- [17] Dieter Happel and Dan Zacharia. A homological characterization of piecewise hereditary algebras. *Math. Z.*, 260(1):177–185, 2008.
- [18] Dieter Happel and Dan Zacharia. Homological properties of piecewise hereditary algebras. *J. Algebra*, 323(4):1139–1154, 2010.
- [19] Otto Kerner, Andrzej Skowroński, Kunio Yamagata, and Dan Zacharia. Finiteness of the strong global dimension of radical square zero algebras. *Cent. Eur. J. Math.*, 2(1):103–111 (electronic), 2004.
- [20] Dirk Kussin, Helmut Lenzing, and Hagen Meltzer. Triangle singularities, ADE-chains, and weighted projective lines. *Adv. Math.*, 237:194–251, 2013.
- [21] Helmut Lenzing. Hereditary Noetherian categories with a tilting complex. *Proc. Amer. Math. Soc.*, 125(7):1893–1901, 1997.
- [22] Helmut Lenzing and José Antonio de la Peña. Wild canonical algebras. *Math. Z.*, 224(3):403–425, 1997.
- [23] Helmut Lenzing and Idun Reiten. Hereditary Noetherian categories of positive Euler characteristic. *Math. Z.*, 254(1):133–171, 2006.
- [24] Helmut Lenzing and Andrzej Skowroński. Quasi-tilted algebras of canonical type. *Colloq. Math.*, 71(2):161–181, 1996.
- [25] Daniel Simson and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 2*, volume 71 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2007. Tubes and concealed algebras of Euclidean type.
- [26] Daniel Simson and Andrzej Skowroński. *Elements of the representation theory of associative algebras. Vol. 3*, volume 72 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2007. Representation-infinite tilted algebras.
- [27] Andrzej Skowroński. On algebras with finite strong global dimension. *Bull. Polish Acad. Sci. Math.*, 35(9-10):539–547, 1987.
- [28] Andrzej Skowroński and Kunio Yamagata. Selfinjective algebras of quasitilted type. In *Trends in representation theory of algebras and related topics*, EMS Ser. Congr. Rep., pages 639–708. Eur. Math. Soc., Zürich, 2008.

(Edson Ribeiro Alves) CENTRO POLITÉCNICO, DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DO PARANÁ, CP019081, JARDIM DAS AMERICAS, CURITIBA-PR, 81531-990, BRAZIL
E-mail address: rolo1rolo@gmail.com

(Patrick Le Meur) LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ BLAISE PASCAL & CNRS, COMPLEXE SCIENTIFIQUE LES CÉZEAUX, BP 80026, 63171 AUBIÈRE CEDEX, FRANCE
Current address: Université Paris Diderot, Sorbonne Paris Cité, Institut de Mathématiques de Jussieu-Paris Rive Gauche, UMR 7586, CNRS, Sorbonne Universités, UPMC Univ. Paris 06, F-75013, Paris, France
E-mail address: patrick.lemeur@imj-prg.fr

(Eduardo N. Marcos) IME-USP (DEPARTAMENTO DE MATEMÁTICA), RUA MATÃO 1010 CID. UNIV., SÃO PAULO, 055080-090, BRAZIL
E-mail address: enmarcos@ime.usp.br